Title: LrhA promotes CRISPR-Cas immunity by promoting reciprocal interplay between interference and primed adaptation in *Escherichia coli*

Running Title: LrhA coordinates CRISPR adaptation and interference in *E. coli*

Mengdie Fang [1,2], Na Li [1], Mingyue Fei [1], Yanan Lu [1], Mingjing Yu [1], Dongchang Sun [1*]

[1] College of Biotechnology and Bioengineering, Zhejiang University of Technology, Hangzhou, Zhejiang, 310014, China

[2] School of Laboratory Medicine and Bioengineering, Hangzhou Medical College, Hangzhou, Zhejiang, 310013, China

[*] Corresponding authors: sundch@zjut.edu.cn

Tel/Fax: +86 0571 88320057

ABSTRACT

The CRISPR-based defense system protects prokaryotes against invading plasmids and viruses by memorizing their nucleotide-sequence traits (adaptation) and destructing the nucleic acids (interference). Previous studies have indicated that CRISPR-Cas adaptive immunity is the outcome of the combined effect of adaptation and interference, but the environmental and physiological cues that coordinate these two processes are unclear. Here, we identify the LysR-type transcriptional regulator LrhA as an important CRISPR-Cas regulator, which promotes the clearance of CRISPR-targeted plasmid more efficiently at 37 ℃ than at 30 ℃ in *Escherichia coli*, by increasing the acquisition of new spacers via interference-driven adaptation. LrhA directly binds to the promoter of the *cas* operon to activate the transcription of *cas* genes required for adaptation and interference, thereby enhancing the CRISPR-Cas adaptive immunity. Altogether, our results demonstrate that LrhA plays a pivotal role in coordinating CRISPR adaptation and interference at the body temperature in a gut-dwelling bacterium.

INTRODUCTION

Prokaryotes have evolved diverse defense systems against plasmids and viruses. Among them, the CRISPR-Cas adaptive immune system provides bacteria and archaea with acquired and inheritable immunity against previously encountered invading nucleic acids[1-6]. The CRISPR-Cas system acts in three distinct stages. During the first contact with a foreign genetic element, short DNA fragments from this element are incorporated into the CRISPR array, which contains the genetic memory of previous infecting elements[7-9]. The incorporation of a new spacer derived from the foreign DNA together with an additional repeat leads to the expansion of the size of the CRISPR array. This process is termed adaptation[10-13]. Triggered by environmental and physiological cues, Cas proteins are expressed, accompanied by the transcription of the CRISPR array, and subsequent processing of the premature CRISPR RNA (pre-crRNA) into mature crRNAs[14,15]. In the final stage termed interference, crRNAs guide Cas proteins to the target sequence, where the Cas3 nuclease is recruited for cleaving the previously encountered foreign DNA[16,17]. Native CRISPR-Cas systems have been engineered for broad applications such as genome editing tools[18-21] and antimicrobials for combating multi-drug resistant (MDR) bacteria [22-27]. Given that delivering engineered CRISPR-Cas systems as antimicrobials in the form of DNA could be challenged by endogenous CRISPR-Cas systems, it is critical to understand how the endogenous CRISPR-Cas systems are activated.

CRISPR-Cas adaptive immunity represents a powerful weapon of defense in prokaryotes. On the other hand, invasive nucleic acids often rapidly evolve resistance to CRISPR immunity by mutating the CRISPR-targeted region. To fight back, a process named 'primed adaptation' is provoked to rapidly incorporate new spacers, derived from DNA fragments produced by interference, for upgrading the CRISPR-Cas adaptive immunity[28-30]. Primed adaptation requires all components of the CRISPR-Cas system for assembly of the interference and the adaptation complexes[28,31]. Nevertheless, a high level of CRISPR-Cas adaptive immunity increases the burden of producing Cas proteins and crRNAs, as well as the risk of

suicide from autoimmunity[32-35]. To finely tune the CRISPR immunity to a moderate

level, interference and adaptation are often inversely correlated: an intermediate level

of interference allows invasive nucleic acids to persist longer, leading to higher levels

of adaptation (interference-driven adaptation), while a strong interference leads to a

relatively low level of adaptation [36-38]. Although the working mechanisms of CRISPR

adaptation and interference have been intensively investigated[7,10,12,13,31,39-46], little is

known about environmental and physiological cues that adjust the CRISPR-Cas

adaptive immunity by leveraging both CRISPR interference and adaptation[14,15].

The type I CRISPR-Cas system, comprised of the multi-subunit RNA-guided

surveillance complex, the Cas3 nuclease, and the Cas1-Cas2 integrase, is most

prevalent in prokaryotes[47]. Regulation studies have focused on type I CRISPR-Cas

systems, especially the type I-E CRISPR-Cas system of *Escherichia coli*, in which

transcription of *cas* genes are positively regulated by the LysR type transcriptional

regulator (LTTR) LeuO[48] and the histone-like protein StpA[49,50], and negatively

controlled by the carbon catabolite regulator CRP[51] and the histone-like protein

H-NS[52]. Because the nuclease activity of Cas3, a limiting factor of CRISPR-Cas

interference, is low at the body temperature (i.e., 37 ℃)[53,54], previous studies have

been predominantly performed at the environmental temperature (i.e., 30 ℃) in *E.*

coli[50,52,55]. However, it remains unclear how the CRISPR-Cas system is activated at

the physiological temperature (37 ℃) in *E. coli*.

To study the regulation of the CRISPR-Cas adaptive immunity in *E. coli* at

physiological temperature, we screened for regulators of the CRISPR-Cas system at

37 ℃ by combining DNA pull-down and mass spectrometry, and identified the LTTR

protein LrhA as an important regulator. By promoting a positive feedback circuit

between interference and adaptation, LrhA augmented the clearance of

CRISPR-targeted plasmid at 37 ℃, via direct binding to the promoter of the *cas*

operon to stimulate the transcription of *cas* genes required for both adaptation and

interference. Our work uncovers a novel mechanism by which a bacterium

coordinates interference and adaptation to achieve optimal immunity in response to

temperature, providing a foundation for curbing the endogenous CRISPR-Cas system

when using CRISPR-Cas antimicrobials in different scenarios.

RESULTS

Screening for proteins binding to the promoter of *cas* genes

Transcription of all *cas* genes except for *cas3* starts from the promoter P_{cas} upstream

of the *cas* operon[52] (Fig. 1a). Both H-NS and LeuO regulates the expression of the *cas*

operon by binding to specific sites within P_{cas} [48,52] (Fig. 1b, Supplementary Fig. S1).

To screen for regulators that potentially affect the transcription of *cas* genes at 37 ℃,

we performed a DNA pull-down assay using a DNA probe containing P_{cas} as a bait

(Fig. 1c). Because H-NS binds to high-affinity sites within P_{cas} and spreads along the

adjacent DNA[56], potentially masking the binding sites of other regulators, we

screened for proteins binding to P_{cas} in *hns*-deletion mutants of two *E. coli* strains

(MC4100 and BW25113). Mass spectrometry analysis showed that more than one

hundred proteins in each strain were isolated with the biotinylated P_{cas}

(Supplementary Fig. S2a). Among them, 49 proteins were detected in both strains

(Supplementary Fig. S2b, Supplementary Table S1). According to the bioinformatic

analysis of proteins with high coverage of matched peptides, 18 proteins were

classified as transcriptional factors (TFs), and 2 proteins (NadR and GlpR) were

annotated as multifunctional proteins (Supplementary Table S2). Two previously

documented regulators of the CRISPR-Cas system (i.e. StpA[49,50] and SlyA[57,58]) were

identified in the protein mixture bound to P_{cas}, indicating that the screening approach

was effective. Remarkably, the LTTR LrhA was detected in protein mixtures bound to

P_{cas} in both strains (Fig. 1d, Supplementary Table S2), with high coverage of peptide

sequences (Fig. 1e, Supplementary Fig. S3). Subsequent pull-down assay confirmed

that purified LrhA with a 6×His-tag bound P_{cas}, indicating that LrhA is a potential

regulator of the CRISPR-Cas system in *E. coli* (Fig. 1f).

To check whether P_{cas}-binding TFs regulate transcription of the *cas* operon, their

effects on the activity of P_{cas} were evaluated with a previously described fluorescence

reporter system[50] (Fig. 1g upper panel). To avoid the effect of the known strong

regulators H-NS and LeuO, the 12 predicted TFs, and LeuO (as the positive control),

were separately over-expressed in an Δhns $\Delta leuO$ mutant. SlyA promoted the activity

of P_{cas} by 2.4-fold (Fig. 1g), in agreement with the previous report [59]. Eight TFs

contain the HTH-type DNA binding domain, which is also present in LeuO (Fig. 1h).

The LTTR ArgP, known to act either as transcriptional activators or repressors

depending on its binding with two effectors lysine and arginine[60], reduced the activity

of P_{cas} by 70%, while LrhA increased the activity of P_{cas} by more than 2-fold (Fig. 1g).

Although YdcI and AaeR also belong to LTTRs and bind to P_{cas}, they showed

minimal (if any) effect on the activity of P_{cas} (Fig. 1g). Overexpression of the glycerol

metabolic regulator GlpR also resulted in strong stimulation of the transcription of

P_{cas}, increasing their transcripts by more than 2 folds (Fig. 1g). Thus, we narrowed

down to the regulators LrhA, ArgP, GlpR and SlyA below.

Activation of clearance of the CRISPR-targeted plasmid by LrhA is affected by

temperature

We investigated the function of LrhA, ArgP, GlpR and SlyA on CRISPR interference

by examining clearance of the CRISPR-targeted plasmid (pT) and the

CRISPR-non-targeted plasmid (pNT) in the Δhns $\Delta leuO$ mutant at 37 ℃ (Fig. 2a). As

shown in Fig. 2b, overexpression of LrhA promoted the clearance rates of the pT but

not the pNT after 24h growth, while overexpression of SlyA or ArgP did not affect

the clearance rates of both plasmids. Overexpression of GlpR reduced cell viability to

a similar level regardless of the harboring plasmids, but showed no effect on CRISPR

interference (Fig. 2b). Therefore, we focused on LrhA in the following investigation.

To confirm the impact of LrhA on the CRISPR system, a $lrhA^{-}$ strain was generated

by in-frame deletion of $lrhA$ in the Δhns $\Delta leuO$ mutant, and the clearance of the pT

was determined in this mutant. The $lrhA^{-}$ strain formed swarm rings with increased

diameter compared with the $lrhA^{+}$ strain (Supplementary Fig. S4), in agreement with

the previous report that LrhA represses transcription of the motility master regulator

operon $flhDC$[61]. Activation of CRISPR interference by LrhA was examined with the

pT and the pNT, through a documented natural transformation method[50,62-64] and

plasmid loss in transformants. Although the ratios of transformation efficiencies

(pT/pNT) were similar in $lrhA^+$ and $lrhA^-$ mutants (Supplementary Fig. S5a), stability

of the pT varied in transformants which were grown in test tubes with antibiotic-free

LB medium at 37℃ and at 30 ℃. The clearance rate of the pT in the $lrhA^+$ strain was

decreased faster than in the $lrhA^-$ strain (Supplementary Fig. S5b). After 36 h

incubation, the clearance rate of the pT was more than 12-fold lower at 37 ℃ while

only 5-fold at 30 ℃ in the $lrhA^-$ strain than in the $lrhA^+$ strain (Fig. 2c).

Complementing $lrhA$ on genome ($lrhA^c$) partially regained the CRISPR interference at

both 37 ℃ and 30 ℃ (Fig. 2c-d). These results indicate that LrhA activates CRISPR

interference and the impact of LrhA on CRISPR interference is

temperature-dependent, stronger at 37 ℃ than at 30 ℃.

Promotion of the primed adaptation by LrhA is affected by temperature

In contrast to the naïve adaptation which requires only the Cas1-Cas2 integrase, all

components of the CRISPR-Cas system are involved in the acquisition of DNA

spacers from CRISPR-targeted DNA during the primed adaptation[10,11,13,65]. To

investigate the role of LrhA in adaptation, we monitored CRISPR array expansion in

response to the pT and the pNT in $lrhA^+$ and $lrhA^-$ strains (Fig. 2e). Compared with

that in the $lrhA^+$ strain, fewer expansion events were observed in the $lrhA^-$ mutant

carrying the CRISPR targeted plasmid after 1.5 or 3 days of incubation at 30 ℃ and

37 ℃ (Fig. 2f, Supplementary Fig. S6a). CRISPR array expansion was not observed

in the $lrhA^+$ and $lrhA^-$ strains carrying the pNT, or in the strains lacking the *cas*

operon (Fig. 2f, Supplementary Fig. S6b), indicating that LrhA promotes the primed

but not the naïve adaptation.

Consistent with a stronger impact of LrhA on CRISPR interference at higher

temperature, the expansion band of the CRISPR array from the $lrhA^+$ strain was more

evident at 37 ℃ than at 30 ℃ (Fig. 2e, Supplementary Fig. S6a), suggesting that

acquisition of new spaces occurred more frequently when LrhA was expressed at the

body temperature. Complementing $lrhA$ on the genome of the $lrhA^-$ mutant ($lrhA^c$)

restored adaptation at 37 ℃, but the restoration effect was weaker at 30 ℃

(Supplementary Fig. S6a). Sequencing analysis of the expansion bands revealed that

all newly formed spacers (20/20) were derived from the CRISPR targeted plasmid

(Fig. 2g), with a strong preference for an AAG PAM (16/20) during adaptation, and a

pronounced bias in the DNA strand (15/20) matching the orientation of the

CRISPR-targeting units (CR1) (Supplementary Table S3), in agreement with typical

traits of primed CRISPR adaptation[28,30].

Cas1-2 is essential to activation of CRISPR interference by LrhA

LrhA promoted CRISPR-Cas interference and adaptation over a period of days more

efficiently at 37 ℃ than at 30 ℃ (Fig. 2). Because acquisition of new spacers allows

bacteria to mount more efficient CRISPR interference[28,30,66], we hypothesized that

high CRISPR-Cas interference and adaptation were coupled and reinforced by LrhA

at 37 ℃. To test this, we deleted both *cas1* and *cas2* (Δ*cas1-2*) in *lrhA*[+] and *lrhA*[−]

strains, and examined the CRISPR interference in the two strains through the plasmid

loss assay. *cas1-cas2* deletion abolished the CRISPR array expansion at both 37 ℃

and 30 ℃ (Fig. 2j), consistent with their critical role in adaption. Strikingly, the

clearance rates of the pT in the *lrhA*[+] mutant with *cas1-cas2* deletion were similarly

low at 37 ℃ and 30 ℃ (Fig. 2h-i), indicating that the temperature effect on

LrhA-promoted interference disappeared when adaptation was abolished. In addition,

cas1-cas2 deletion reduced the effect on LrhA in CRISPR interference activity at both

37 ℃ and 30 ℃ (Fig. 2h-i), indicating that LrhA contributes to the positive feedback

of interference and adaptation process (Fig. 2k).

LrhA activates transcription of the *cas* operon

To understand how LrhA controls the CRISPR-Cas adaptive immunity, its role in

regulating the *cas* operon was evaluated with a previously documented reporter

system. We detected only a subtle effect of chromosomally expressed LrhA on the

activity of P$_{cas}$. Although *lrhA* inactivation decreased the activity of P$_{cas}$ by 31% and

37% in the Δ*hns* Δ*leuO* mutant grown at 30 ℃ and 37 ℃, respectively,

complementing *lrhA* fully restored the activity of P$_{cas}$ at 37 ℃, but only partially

restored it at 30 ℃ (Fig. 3a). A more pronounced effect of temperature on the activity

of P$_{cas}$ was detected when LrhA was overexpressed. Expressing LrhA from an

arabinose-inducible promoter on a multi-copy plasmid elevated transcription of the

cas operon by 4.75 folds in the Δ*hns* Δ*leuO* mutant after 36 h of incubation at 37 ℃

(Fig. 3b middle panel). However, no significant effect of expressing LrhA on

transcription of the *cas* operon was observed when the Δ*hns* Δ*leuO* mutant was grown

at 30 ℃ (Fig. 3b left panel), indicating that the impact of LrhA on the transcription of

the *cas* operon was temperature-dependent. Interestingly, deletion of *lrhA* did not

affect the activity of P$_{cas}$ in the Δ*hns* mutant grown at 30 ℃ and 37 ℃ (Fig. 3c). Also,

over-expressing LrhA in the Δ*hns* mutant only slightly increased transcription of the

cas operon after incubation for 36 h at 37 ℃, but not at 30 ℃ (Supplementary Fig.

S7a). In the wild-type strain, overexpressing LrhA did not affect the activity of P$_{cas}$ at

either 30 ℃ or 37 ℃ (Supplementary Fig. S7b), revealing that the effect of LrhA on

the expression of the *cas* operon was masked by H-NS and LeuO, and implicating the

presence of a hierarchical regulation of the *cas* operon.

LrhA activates the *cas* operon by directly binding to the promoter P$_{cas}$

The LrhA-binding site was reported to partially overlap with a LeuO-binding site on

the promoter of *leuO*[67]. Previous work has identified two LeuO binding sites (LBS I

and LBS II) within P$_{cas}$ (Supplementary Fig. S1). The observation that the presence of

LeuO interfered with the regulation of P$_{cas}$ by LrhA prompted us to consider whether

LrhA and LeuO compete for the same binding sites on P$_{cas}$. To check this, DNA

probes lacking LBS I and/or LBS II were generated (named 'P$_{cas}$ del LBS I', 'P$_{cas}$ del

LBS II', and 'P$_{cas}$ del LBS I&II') and EMSA was employed to determine the effect of

LBSI and LBS II on the efficiency of LrhA binding to P$_{cas}$. As shown in Fig. 4a, 700

nM of His-LrhA was sufficient to retard the mobility of the full-length P$_{cas}$ probe (P$_{cas}$

FL). However, 2800 nM of His-LrhA were required to completely retard the

migration of the P$_{cas}$ del LBS I and the P$_{cas}$ del LBS II probes, respectively (Fig. 4a).

Particularly, in the absence of both LBSI and LBSII (P$_{cas}$ del LBS I&II), more than

5600 nM of His-LrhA was necessary to saturate the binding of the probe. These

observations support that LrhA binds to both LBS I and LBS II. To confirm LBS I

and LBS II as binding sites of LrhA, we performed an EMSA assay with shortened

probes (I ~ IV). Sequences of probes I and II are identical to that of LBS I and LBS II

respectively, while sequences of probes III and IV are identical to regions of P$_{cas}$ with

high and low affinity to LrhA respectively (Fig. 4b), as predicted by the SELEX

database[68]. We observed that LrhA bound probes I and II with high affinity, the probe III with lower affinity, and the probe IV with the lowest affinity (Fig. 4b), confirming that LrhA binds to both LBS I and LBS II.

The results above indicate that both LrhA and LeuO bind to LBS I and LBS II on P_{cas} to regulate the expression of the *cas* operon (Fig. 4a and b). To know whether LBS I and/or LBS II are essential to activate P_{cas} by LrhA and LeuO *in vivo*, we generated P_{cas} containing disrupted LBS I and/or LBS II (Fig. 4c), and compared *in vivo* activities of disrupted P_{cas} with that of P_{cas} FL in response to expression of LrhA and LeuO. Over-expressing LrhA and LeuO increased the transcriptional activity of P_{cas} FL by more than 3- and 8-fold, respectively (Fig. 4c). In contrast, the activity of P_{cas} del LBS I in the Δhns $\Delta leuO$ mutant expressing LrhA or LeuO was decreased to the background level or 3.88-fold higher than the background level, respectively (Fig. 4c). This observation indicates that LBS I is indispensable for activation of P_{cas} by LrhA, and essential for full stimulation of P_{cas} by LeuO. Interestingly, the activity of P_{cas} del LBS II was also remarkably reduced in the Δhns $\Delta leuO$ mutant expressing either LrhA or LeuO (Fig. 4c), revealing that LBS II was required for activation of P_{cas} by both LrhA and LeuO. As expected, when both LBS I and LBS II were deleted, the expression of LrhA or LeuO no longer stimulated the expression of the impaired P_{cas} (Fig. 4c). Taken together, these data clearly show that both LBS I and LBS II are essential for activation of P_{cas} by LrhA and LeuO *in vivo*.

DISCUSSION

In this study, we have addressed the important question of how the CRISPR-Cas adaptive immunity is regulated in bacteria. Our work demonstrates that the LTTR LrhA activates CRISPR-Cas adaptive immunity more efficiently at 37 °C than at 30 °C by promoting reciprocal interplay between interference and adaptation in *E. coli*. Knowing how the endogenous CRISPR-Cas system is regulated under the physiological condition would not only unravel its ecological and evolutionary role, but also be a pre-requisite for devising strategies to steer the CRISPR-Cas system towards MDR bacteria.

LrhA triggers a positive feedback circuit between interference and primed adaptation

Early reports showed that primed adaptation was triggered by suboptimal Cascade-protospacer interactions that could lead to weak interference[28,69,70]. However, later studies reported that optimal Cascade-protospacer interactions also led to both interference and primed adaptation, which was fueled by interference[31,66,71,72]. It appears that the level of CRISPR immunity is determined by coordination of interference and adaptation. In this study, we detected remarkably higher interference and adaptation in the *lrhA$^+$* strain than that in the *lrhA$^-$* strain. Adaptation occurred more frequently in the *lrhA$^+$* *E. coli* strains harboring an intact *cas* operon and the CRISPR-targeted plasmid (Fig. 2, and Supplementary Fig. S6), revealing that LrhA can promote interference-driven spacer acquisition. Given that inactivation of *cas1* and *cas2* remarkably reduced LrhA-stimulated CRISPR interference (Fig. 2h), we propose that LrhA promotes the positive feedback circuit between CRISPR interference and adaptation: cleavage of the CRISPR-targeted plasmid produces more short DNA fragments for accelerating primed adaptation, which in turn enhances clearance of the CRISPR-targeted plasmid (Fig. 5). Analysis of the acquired spacers revealed an unequal distribution, indicating that certain sequences are preferentially integrated as new spacers. We also noticed that most spacers were originated from the same strands with a high preference for AAG PAM sequences (Fig. 2g and Supplementary Table S3), matching traits of the primed adaptation process[28,30]. Compared with other regions on the CRISPR-targeted plasmid, protospacers were mapped to encoding genes (i.e., *bla* and *dsRED*) and the replication origin with a remarkably higher frequency (Fig. 2g), suggesting that unwinding of double-stranded DNA during replication and transcription may facilitate the adaptation process.

Temperature affects regulation of the CRISPR-Cas system by LrhA

Previous work demonstrated that temperature regulates the CRISPR-Cas activity in *Pseudomonas aeruginosa* by controlling growth rate[73]. Our study supports that temperature affects the CRISPR-Cas adaptive immunity through controlling the activity of P$_{cas}$, since LrhA promoted a higher rate of clearance of CRISPR-targeted

plasmid and stronger transcription of the *cas* operon at 37 ℃ than 30 ℃ in *E. coli* (Fig.

2 to 3). Considering that the amount of LrhA was not reduced in *E. coli* grown at

30 ℃ with regard to that at 37 ℃ (Supplementary Fig. S8), it seems that temperature

does not function by affecting the expression or stability of LrhA. Because the

deletion of *cas1* and *cas2* abolished the effect of temperature on regulation of

CRISPR interference, we conclude that temperature affected the CRISPR-Cas

adaptive immunity by influencing regulation of the positive feedback circuit between

interference and primed adaptation by LrhA. Since more frequent adaptation in the

lrhA[+] strain at 37 ℃ than at 30 ℃ (Fig. 2f) could be associated with weaker nuclease

activity of Cas3 at 37 ℃ [66], LrhA may increase CRISPR interference with weak Cas3

by increasing the number of spacers matching sites on the targeted plasmid at 37 ℃.

Whereas, CRISPR interference with strong Cas3 at 30 ℃ could be unfavorable to

adaptation, restricting accumulation of spacers in the CRISPR array by LrhA.

Accordingly, we propose that, at the body temperature, LrhA may better coordinate

interference and adaptation to achieve maximal immunity, which is determined by the

combined effect of interference and adaptation, to reduce the risk of invading nucleic

acids.

Regulation of the CRISPR-Cas system by LTTRs

Both *in vivo* and *in vitro* evidence reveals that pleiotropic LTTRs LrhA and LeuO

activate the *cas* operon by binding to identical sites on P$_{cas}$. In contrast, the LTTR

ArgP was found to repress P$_{cas}$, showing diverse regulation of the CRISPR-Cas

system by LTTRs. In *Salmonella enterica*, transcription of the CRISPR-Cas system

was oppositely regulated by two LTTRs (i.e., LeuO and LRP) [74]. Considering the

widespread distribution of the type I CRISPR-Cas system (Supplementary Fig. S11),

it is possible that LTTR family proteins could extensively regulate CRISPR-Cas

systems in different bacteria. Although LeuO is normally considered as an antagonist

of H-NS, and activates gene transcription by relieving transcriptional suppression by

H-NS, we found that both LrhA and LeuO promoted the activity of P$_{cas}$ in the Δ*hns*

Δ*leuO* mutants (Fig. 3b, 4c), indicating that they both regulated the *cas* operon

independent of H-NS. Nevertheless, exact roles of LrhA and LeuO on activating P*cas* remain unclear, awaiting further biochemical and structural analysis.

Despite similarities between LrhA and LeuO, they behaved remarkably differently in regulating transcription of the *cas* operon and other genes as well. Most strikingly, temperature showed a more pronounced impact on the regulation of the CRISPR-Cas system by LrhA than that by LeuO (Fig. 3a and b). Although LeuO strongly stimulated the *cas* operon at both body and environmental temperatures (Fig. 3b), it stimulated weaker interference than LrhA did at the body temperature (Fig. 2b). Strong stimulation of interference by LrhA could be the outcome of a reinforced cycle of interference-driven adaptation and adaptation-promoted interference (Fig. 5). In light of disrupting either of the two binding sites on P*cas* abolished regulation by LrhA while regulation of LeuO was abolished only when both of the two binding sites were destroyed (Fig. 4c), the two regulators must interact with P*cas* in different ways. Apart from regulation of the CRISPR-Cas immunity, LrhA and LeuO showed remarkable differences in controlling other aspects of bacterial physiology. Although inactivating *lrhA* increased the swarming mobility by promoting transcription of the *flhDC* operon in the presence or absence of LeuO, disrupting *leuO* failed to affect bacterial swarming in both *lrhA*$^+$ and *lrhA*$^-$ strains (Supplementary Fig. S4). Phylogenetic analysis revealed divergent evolution of LrhA and LeuO, which could belong to distinct subfamilies and separately evolve in bacteria (Supplementary Fig. S9 and S10). Moreover, regulation of the CRISPR-Cas system by LrhA or LeuO could be evolved in response to different environmental cues, considering that expression of LeuO can be activated by the heterodimer BglJ-RcsB[75], Cyanase regulator CynR[67], and antibiotic responses regulator YdcI[67], while expression of LrhA is stimulated by acid resistance-related gene *hdeD* [76].

METHODS

1. *Bacterial strains, plasmids, primers, growth conditions, and media*

Bacterial strains, plasmids, and primers used in this study are listed in Supplementary Table S4, Table S5 and Table S6. *E. coli* gene-deletion mutants were constructed by

using a λ-RED recombination system expressed by the temperature-sensitive plasmid

pKD46[77]. Corresponding mutants were examined through PCR. Recombinant

plasmids expressing LrhA and other 10 TFs were constructed with a Transfer PCR

method[78]. Plasmids were isolated and purified with a plasmid isolation kit according

to the manufacturer's protocol (Axygen Biotech Co., Ltd.). *E. coli* cultures were

grown at 37 ℃ in LB broth containing 1% (wt/vol) tryptone, 0.5% (wt/vol) yeast

extract, and 1% (wt/vol) NaCl or on LB agar plates containing 1.5% (wt/vol) agar.

When evaluating the GFP/OD$_{600}$ level or the interference/adaptation effect, cells were

grown in M9 minimal medium containing 61 mM K$_2$HPO$_4$·3H$_2$O, 38.21 mM KH$_2$PO$_4$,

2.5 mM MgSO$_4$, 15.14 mM (NH$_4$)$_2$SO$_4$, 0.1% (wt/vol) tryptone, supplemented with

1% (wt/vol) glucose as the carbon source. When required, media were supplemented

with ampicillin (100 μg mL^{-1}), kanamycin (50 μg mL^{-1}), or chloramphenicol (25 μg

mL^{-1}). Arabinose was added for inducing gene expression in liquid cultures where

indicated. Cell growth was measured in a spectrophotometer (MD SpectraMax iD5 or

Biotek Synergy 2). All experiments were repeated independently at least three times

unless otherwise indicated.

2. *pull-down assay*

To prepare samples for mass spectrometric analysis of proteins binding to P$_{cas}$, 8 pmol

biotinylated P$_{cas}$ DNA fragment was coated with 100 μL streptavidin beads

(Invitrogen Dynabeads™ M-270) in the binding buffer A (10 mM Tris-HCl pH 7.5, 1

mM EDTA, 2 M NaCl) for 1 h at the room temperature. Biocytin, a derivative of

biotin, was used as the negative control. From a 5 mL culture of *E. coli hns*

deletion-mutant grown overnight at 37 ℃, proteins were extracted by sonication in

the binding buffer B (10 mM Tris-HCl pH 7.5, 1 mM EDTA, 50 mM NaCl, 0.4 mM

DTT and 10% glycerol), supplemented with protease inhibitor cocktail (Roche Co.,

Ltd.). The crude lysates were centrifuged at 15000×*g* for 10 minutes. The supernatant

was collected, pre-cleared by empty streptavidin beads, and then incubated with 50

μL P$_{cas}$ DNA coated beads overnight at 4 °C under constant rotation (20 rpm). Beads

were washed 5 times with the binding buffer B, followed by washing 3 times with the

binding buffer B supplemented with 1 mg salmon sperm DNA, and finally washing 5

times with the binding buffer B to remove non-specific proteins. DNA binding

proteins were eluted in SDS sample loading buffer and loaded onto 10% SDS-PAGE,

followed by the mass-spectrometric analysis of the gel bands as described below.

3. *Mass spectrometry analysis*

After 37℃ overnight in-gel digestion of proteins binding to P_{cas} with trypsin, peptides

were loaded onto a home-made reversed-phase analytical column (15-cm length, 75

μm i.d.), and analyzed with a gradient from 5% to 34% solvent B (0.1% formic acid

in 80% acetonitrile) over 40 min, 34% to 38% solvent B for 5 min, 38% to 90%

solvent B for 10 min, and then holding at 90% solvent B for 10 min, at a constant

flow rate of 300 nL/min on an EASY-nLC 1000 UPLC system. Peptides were

subjected to NSI source followed by tandem mass spectrometry (MS/MS) in Thermo

Scientific Orbitrap Fusion coupled online to the UPLC. Survey full-scan MS spectra

(m/z 350-1800) were acquired in the Orbitrap with resolution r of 70000 at m/z 400.

Mass spectrometry and analysis of the data were conducted by Micrometer Biotech

Company.

4. *Quantification of gene transcription*

Plasmid pGLO-P_{cas}-*gfp* was used for monitoring the transcription of the *cas* operon,

by measuring the intensity of GFP fluorescence and cell optical density (OD_{600}) on a

microplate reader (MD, SpectraMax iD5), with a method that has previously been

described[50]. For Fig. 3a-c, *E. coli* BW25113 cells expressing pGLO-P_{cas} -*gfp* were

grown in M9 minimum medium at 30 ℃ or 37 ℃. For Fig. 4c, Plasmids pGLO-P_{cas}

-*gfp*, pGLO-P_{cas} del LBS I-gfp, pGLO-P_{cas} del LBS II-gfp, pGLO-P_{cas} del LBS

I&II-*gfp*, were transformed into the Δ*hns* Δ*leuO* mutant cells separately, for

measuring GFP/OD_{600} of cells grown in M9 minimum medium at 37 ℃.

5. *Purification of LrhA Protein*

Full-length LrhA ORF with N terminal His-tag was cloned into the pET28A vector, to

generate pET28a-His-*lrhA*, which was transformed into *E. coli* BL21 (DE3). The

overnight culture was inoculated into 100 mL LB medium at a ratio of 1:100. When

the culture was grown to an OD_{600} of 0.4, expression of LrhA was induced with 0.5

mM IPTG (Beyotime, ST098) for 2 h. Cells were centrifuged at 8000 rpm for 5 min at

4 °C. The cell pellet was resuspended in PBS containing protease inhibitor and

sonicated, and the lysate was centrifuged at 12000×g for 5 min at 4 °C. His-LrhA

protein was purified by Ni-NTA Resin (ThermoFisher, 88222) in a gravity-flow

column, and the bound protein was eluted with PBS containing 250 mM imidazole.

The eluted protein was analyzed by SDS-PAGE, and dialyzed at 4°C in PBS with a

10 kDa dialysis tube (Millipore, UFC801096). Protein concentrations were

determined by the Bradford assay (Beyotime, P0006C).

6. Electrophoretic mobility shift assay (EMSA)

EMSA was performed by using the LightShift Chemiluminescent EMSA kit

(ThermoFisher, 20148) according to the manual guide. P$_{cas}$ (20 fmol) probes were

mixed with purified LrhA protein of different concentrations in the presence of

binding buffer (10 mM Tris-HCl pH 7.5, 50 mM KCl, 1 mM DTT, 2.5% glycerol, 1

mM EDTA). A Ctrl probe generated by PCR from the *lacZα* gene with primers P63

and P64 were used as control. The mixture was incubated at room temperature for 20

minutes, then separated by electrophoresis in 5% native polyacrylamide gels in

0.5×TBE buffer. Gels were visualized by SYBR Gold staining (ThermoFisher,

S11494). For Fig. 4b, bands were transferred to PVDF membrane using the semi-dry

transfer apparatus (Genscript) and visualized by Chemiluminescent Nucleic Acid

Detection Module (ThermoFisher, 89880). Images were collected with ChemiDoc™

XRS$^+$ (BioRad).

7. Plasmid loss assay

The *E. coli* strains harboring the CRISPR-targeted plasmid pT (pDsRED-CR1) or the

CRISPR-non-targeted plasmid pNT (pDsRED) were grown overnight in LB culture

supplemented with ampicillin (100 μg mL^{-1}). The overnight grown culture was

washed and inoculated at a ratio of 2% into antibiotic-free LB culture, followed by

incubation at 30 ℃ or 37 ℃ with shaking. At intervals, the culture was serially

diluted and spread on the LB plate supplemented with or without ampicillin (100 μg

mL^{-1}). The number of cells maintained the plasmid was evaluated by comparing

colonies on the LB plate supplemented with or without ampicillin.

8. Spacer acquisition assay

To monitor spacer acquisition, *E. coli* Δ*hns* Δ*leuO* and the Δ*hns* Δ*leuO* Δ*lrhA* strains

carrying pDsRED-CR1 or pDsRED were grown overnight in LB media supplemented

with ampicillin (100 μg mL^{-1}), then diluted to 1:50 times with antibiotic-free M9

media and growth was continued for 36 h at 30 ℃ or 37 ℃ with shaking. To monitor

CRISPR array expansion, cells from 200 μL of the culture were collected by

precipitation, washed thrice, and then resuspended in distilled water. These cells were

used as the template for PCR amplification of the CRISPR array I by using the primer

pair P65 (Adaptation-F) and P66 (Adaptation-R) as shown in Fig. 2e. All PCR

products were separated on 2% agarose gels to identify parental and expanded arrays

(parental array + 61bp). For sequencing newly acquired spacers, the expanded array

bands were excised, and corresponding DNA fragments were purified using a PCR

Clean-up kit (MN 740609.50), and ligated to T vectors (Takara, 6013).

Availability of data and materials

The raw data generated during the current study are available from the corresponding

authors on request. Source data are provided with this paper. Uncropped gels and

Western blots can be found in Supplementary Information. The mass spectrometry

proteomics data have been deposited to the ProteomeXchange Consortium

(http://proteomecentral.proteomexchange.org) via the iProX partner repository[79] with

the dataset identifier PXD036879.

COMPETING INTERESTS

476 We declare that we have no financial and personal relationships with other people or

477 organizations that can inappropriately influence our work, there is no professional or

478 other personal interest of any nature or kind in any product, service and/or company

479 that could be construed as influencing the position presented in, or the review of, the

480 manuscript entitled.

ACKNOWLEDGEMENT

This work was supported by the National Natural Science Foundation of China (grants 31670084 and 31100071), the Key Research and Development Program of Zhejiang Province (grant 2020C02031), and the Open Foundation of State Key Laboratory Cultivation Base of Hazard Factors and Risk Control of Quality and Safety of Agricultural Products (grant 2021DG700024-KF202105). We thank Prof. Shishen Du at Wuhan University and Dr. Yanmei Zhang at Hangzhou Medical College for scientific advice on this manuscript. We also thank Prof. Zhiyong Sun at Zhejiang University of Technology, Prof. Pengjun Zhang at Hangzhou Normal University, Ms. Minjia Shen at Université Paris-Saclay, Ms. Xia Yao at Zhejiang University of Technology, and all members of the Sun laboratory for critically reading the manuscript and helpful discussions.

    FIGURE LEGENDS

Figure 1. Screen for new regulators modulating transcription of *cas* genes.

(**a**) Schematic illustration of the CRISPR/Cas locus in *E. coli*. Repeats (diamonds) in

the CRISPR array are interspersed with 12 unique spacer sequences (rectangles). The

promoter of the *cas* operon (P_{cas}) is indicated by the black arrow. (**b**) The P_{cas} DNA

probe for pull-down assay contains a transcription start site (red letter), LeuO binding

site I and II (LBS I/II; purple bars), and H-NS binding site (HBS; yellow bar). (**c**)

Design of the pull-down proteomic experiment. A biotinylated probe (biotin-P_{cas} DNA)

was immobilized onto streptavidin beads, followed by incubation with extracts from *E.*

coli MC4100 Δhns or BW25113 Δhns before SDS-PAGE electrophoresis. (**d**)

SDS-PAGE gel electrophoresis of nontargeted biocytin (negative control, NT) or

biotin-P_{cas} DNA-binding proteins. NT or biotin-P_{cas} samples were digested in gel with

trypsin, and corresponding peptides were analyzed by mass spectrometry. An arrow

denotes the band corresponding to LrhA (34.593 kDa). (**e**) Peptides identified by mass

spectrometry (pink) were mapped to the amino acid sequence of LrhA. (**f**) Pull-down

assay of physical interaction between LrhA and P_{cas}. The biotinylated probe

(biotin-P_{cas} DNA) was immobilized onto streptavidin beads and incubated with

His-LrhA from cell extracts before (lane 3, input) and after (lane 2, P_{cas}) pull-down,

followed by western blot assay with antibody against His-tag. Nontargeted biocytin

(NT) was used as a negative control (lane 1). Arrow indicates the band of His-LrhA,

and the asterisk indicates a non-specific band. (**g**) Effects of 12 predicted TFs and

LeuO on the P_{cas} activity in *E. coli* BW25113 Δhns $\Delta leuO$. The cells were grown in

M9 medium supplemented with 10 mM arabinose, 100 μg/ml ampicillin and 25 μg/ml

chloramphenicol at 37 ℃ The P_{cas} activity was evaluated by GFP/OD$_{600}$, with the

relative expression level of Δhns $\Delta leuO$ carrying empty vector pSU19 (Vector) as 1

arbitrary unit. All bars represent the mean, and error bars denote standard deviation.

Statistical significance was determined using one-way ANOVA with Dunnett's

multiple comparisons test. (**h**) Phylogenetic analysis of 9 HTH-type P_{cas}-binding TFs

from 5-8 different bacteria species.

Figure 2. LrhA activates both CRISPR-Cas interference and adaptation.

(**a**) Schematic illustration for evaluating the CRISPR-Cas interference activity by

plasmid loss assay. (**b**) Plasmid loss assay were performed in *E. coli* strains BW25113

Δ*hns* Δ*leuO* expressing GlpR, LeuO, SlyA, LrhA or ArgP. Cells carrying the pT

(CRISPR-targeted plasmid) or the pNT (CRISPR-non-targeted plasmid) were grown

in test tubes containing antibiotic-free M9 medium at 37 ℃ for 24 h, then serially

diluted for drop-plating onto LB agar plates supplemented with or without 100 μg

mL^{-1} ampicillin. (**c** and **d**) *E. coli* strains BW25113 Δ*hns* Δ*leuO* (*lrhA*$^+$), Δ*hns* Δ*leuO*

Δ*lrhA* (*lrhA*$^-$) or genomic *lrhA* complementation (*lrhA*^{c}) carrying the pT or the pNT

were grown in antibiotic-free M9 medium at 37 ℃ or 30 ℃. After 36 h growth,

plasmid loss was evaluated by drop-plating at 37 ℃ (**c**) or 30 ℃ (**d**). (**e**) Schematics

of the spacer acquisition assay in (**f** and **j**). During the adaptation process, a new

spacer (a yellow rectangle) can be inserted into the leader end of the CRISPR array.

PCR was performed to amplify the leader-spacer region using the primers matching

the leader and the second spacer, which were indicated as flagged arrows. (**f**)

Spacer-acquisition assay by PCR amplification of Δ*hns*Δ*leuO* (*lrhA*$^+$), Δ*hns* Δ*leuO*

Δ*lrhA* (*lrhA*$^-$) cells carrying the pT or the pNT plasmids, which were grown at 37 ℃

or 30 ℃ for 36 h. PCR products were electrophoresed on a 2% agarose gel and

imaged. Parental (P) and expanded bands (E) are indicated on the right while DNA

marker (M) positions are displayed on the left. Gels are representative of two

experiments yielding similar results. (**g**) Mapping of newly formed spacers on the pT

plasmid DNA. The DNA of the expanded array was sequenced by ligating purified

PCR products into T vectors. The position and orientation of the sequenced spacers

are marked on the pT plasmid map (blue arrows pointing clockwise and red arrows

pointing counterclockwise). Protospacers are numbered according to the strain from

which the spacers were sequenced, as listed in Supplementary Table 3. CR1 indicates

4 spacers targeted by the type I-E CRISPR-Cas system of *E. coli*. (**h-i**) Plasmid loss

assay of *E. coli* strains BW25113 Δ*hns* Δ*leuO* (*lrhA*$^+$), Δ*hns* Δ*leuO* Δ*lrhA* (*lrhA*$^-$),

Δ*hns* Δ*leuO* Δ(*cas1-cas2*) (*lrhA⁺Δcas1-2*) and Δ*hns* Δ*leuO* Δ*lrhA* Δ(*cas1-cas2*) (*lrhA*

⁻Δ*cas1-2*) carrying the pT or the pNT, which were grown at 37 ℃ or 30 ℃ for 36 h.

(**j**) Spacer-acquisition assay by PCR amplification of Δ*hns*Δ*leuO* (*lrhA⁺*), Δ*hns* Δ*leuO*

Δ*lrhA* (*lrhA* ⁻), Δ*hns* Δ*leuO* Δ(*cas1-cas2*) (*lrhA⁺Δcas1-2*) and Δ*hns* Δ*leuO* Δ*lrhA*

Δ(*cas1-cas2*) (*lrhA* ⁻Δ*cas1-2*) cells carrying the pT, which were grown at 37 ℃ or

30 ℃ for 36 h. (**k**, upper panel) LrhA promotes the positive feedback circuit between

CRISPR interference and adaptation, which requires *cas1-2*. (**k**, lower panel) In

absence of cas1-2, LrhA showed a weak effect on interference since the adaptaion is

abolished.

Figure 3. LrhA promotes transcription of the *cas* operon.

(**a**) Effects of genomic *lrhA* on the P_{cas} activity in Δ*hns* Δ*leuO* grown in M9 medium

at 30 ℃ (left panel) or 37 ℃ (middle panel), with the relative expression level of

Δ*hns* Δ*leuO* of 24 h at 30℃ or 37℃ as 1 arbitrary unit. (**b**) Effects of LrhA or LeuO

expressed from the p15A derived plasmid pSU19 on the P_{cas} activity, indicated by

GFP/OD$_{600}$, were evaluated in BW25113 Δ*hns* Δ*leuO* mutants, which were grown in

M9 medium supplemented with 20 mM arabinose, 100 μg/ml ampicillin and 25 μg/ml

chloramphenicol at 30 ℃ (left) or 37 ℃ (middle). The relative expression level of *E.*

coli carrying empty vector pSU19 (Vector) was set as 1 arbitrary unit. (**c**) Effects of

genomic *lrhA* on the P_{cas} activity in Δ*hns* grown in M9 medium at 30 ℃ (left) or

37 ℃ (middle), with the relative expression level of Δ*hns* Δ*leuO* of 24 h incubation at

30 ℃ or 37 ℃ as 1 arbitrary unit. Mean values of at least three biological replicates

are shown as bars, and error bars indicate standard deviations. Data are shown as

means ± standard deviations (*n* = 4). Statistical significance was determined using

one-way ANOVA with Dunnett's multiple comparisons test (a, b) or an unpaired

two-tailed Student's t-test (c).

Figure 4. Identification of LrhA binding sites on the promoter of the *cas* operon.

(**a**, upper panel) Schematic illustration of the P_{cas} region for truncation analysis. LBS I

and LBS II indicate two DNA-binding sites of LeuO located upstream and

downstream of the transcriptional start site (TSS, black arrow) on P_{cas} (purple bars).

HBS indicates the DNA-binding site of the H-NS (yellow bar). (**a**, lower panel)

electrophoretic mobility shift (EMSA) assays of full-length or truncated P_{cas} DNA

fragments by LrhA. The P_{cas} probes including full-length P_{cas} (P_{cas} FL), P_{cas} with LBS

I deleted (P_{cas} del LBS I), P_{cas} with LBS II deleted (P_{cas} del LBS II) and P_{cas} with both

LBS I and LBS II deleted (P_{cas} del LBSI/II) were generated by PCR with primer P61

and P62. 20 fmol P_{cas} probes were separately incubated with a serial concentration of

purified His-LrhA, before EMSA assays with 5% native polyacrylamide gels (solid

arrow). A 175 bp *lacZα* fragment that was generated by PCR with primers P63 and

P64 was used as control (dashed arrow). (**b**, upper panel) Schematic of probes

matching the P_{cas} region for EMSA assay. DNA fragments "I" and "II" match LBS I

and LBS II respectively. DNA fragments "III" and "IV" match a region in between

LBS I and LBS II, and a region partially match downstream of LBS I respectively. (**b**,

lower panel) EMSA of DNA fragments "I" to "IV" by LrhA. Probes "I", "II", "III"

and "IV" were separately incubated with a serial concentration of purified His-LrhA

before EMSA assays with 5% native polyacrylamide gels. (**c**, upper panel) Schematic

illustration of DNA binding sites required for activating P_{cas} by LrhA and LeuO.

Transcriptional activities of full-length P_{cas}, P_{cas} del LBS I, P_{cas} del LBS II and P_{cas} del

LBS I/II were compared by using the P_{cas}-*gfp* reporter system. (**c**, lower panel) Effects

of disruption of LBS I and LBS II on activation of P_{cas} by LrhA. The P_{cas} activity was

evaluated by GFP/OD$_{600}$, with the relative expression level of Δhns $\Delta leuO$ carrying

the empty vector as 1 arbitrary unit. The data are shown as means with standard

deviations from three independent experiments. Statistical significance was

determined using one-way ANOVA with Dunnett's multiple comparisons test.

Figure 5. Schematic diagram of regulation of CRISPR-Cas immunity by LrhA in

E. coli. LrhA binds to P_{cas} and activates transcription of the *cas* operon, increasing the

amount of the surveillance complex Cascade and the integrase Cas1-Cas2. LrhA not

only augments interference of CRISPR-targeted plasmid (pT), which is mediated by

Cascade and the nuclease Cas3, but also promotes interference-driven spacer

acquisition which is mediated by Cas1-Cas2, together with Cascade and Cas3.

REFERENCES

1   Makarova, K. S. *et al.* Evolutionary classification of CRISPR-Cas systems: a burst of class 2
and   derived   variants.   *Nat.   Rev.   Microbiol.*   **18**,   67-83   (2020).
https://doi.org:10.1038/s41579-019-0299-x

2   Mohanraju, P. *et al.* Diverse evolutionary roots and mechanistic variations of the CRISPR-Cas
systems. *Science.* **353**, aad5147 (2016). https://doi.org:10.1126/science.aad5147

3   Hille, F. *et al.* The biology of CRISPR-Cas: backward and forward. *Cell.* **172**, 1239-1259
(2018). https://doi.org:10.1016/j.cell.2017.11.032

4   Koonin, E. V., Makarova, K. S. & Zhang, F. Diversity, classification and evolution of
CRISPR-Cas   systems.   *Curr.   Opin.   Microbiol.*   **37**,   67-78   (2017).
https://doi.org:10.1016/j.mib.2017.05.008

5   Garneau, J. E. *et al.* The CRISPR/Cas bacterial immune system cleaves bacteriophage and
plasmid DNA. *Nature* **468**, 67-71 (2010). https://doi.org:10.1038/nature09523

6   Marraffini, L. A. & Sontheimer, E. J. CRISPR interference limits horizontal gene transfer in
staphylococci   by   targeting   DNA.   *Science.*   **322**,   1843-1845   (2008).
https://doi.org:10.1126/science.1165771

7   Barrangou, R. *et al.* CRISPR provides acquired resistance against viruses in prokaryotes.
Science. **315**, 1709-1712 (2007). https://doi.org:10.1126/science.1138140

8   Horvath, P. *et al.* Diversity, activity, and evolution of CRISPR loci in *Streptococcus*
thermophilus. *J Bacteriol.* **190**, 1401-1412 (2008). https://doi.org:10.1128/jb.01415-07

9   Deveau, H. *et al.* Phage response to CRISPR-encoded resistance in *Streptococcus*
thermophilus. *J Bacteriol.* **190**, 1390-1400 (2008). https://doi.org:10.1128/jb.01412-07

10 Amitai, G. & Sorek, R. CRISPR-Cas adaptation: insights into the mechanism of action. *Nat.*
Rev. Microbiol. **14**, 67-76 (2016). https://doi.org:10.1038/nrmicro.2015.14

11 Lee, H. & Sashital, D. G. Creating memories: molecular mechanisms of CRISPR adaptation.
Trends Biochem Sci. **47**, 464-476 (2022). https://doi.org:10.1016/j.tibs.2022.02.004

12 Levy, A. *et al.* CRISPR adaptation biases explain preference for acquisition of foreign DNA.
Nature **520**, 505-510 (2015). https://doi.org:10.1038/nature14302

13 Sternberg, S. H., Richter, H., Charpentier, E. & Qimron, U. Adaptation in CRISPR-Cas
systems. *Mol. Cell* **61**, 797-808 (2016). https://doi.org:10.1016/j.molcel.2016.01.030

14 Patterson, A. G., Yevstigneyeva, M. S. & Fineran, P. C. Regulation of CRISPR-Cas adaptive
immune   systems.   *Curr.   Opin.   Microbiol.*   **37**,   1-7   (2017).
https://doi.org:10.1016/j.mib.2017.02.004

15 Shivram, H., Cress, B. F., Knott, G. J. & Doudna, J. A. Controlling and enhancing CRISPR
systems. *Nat. Chem. Biol.* **17**, 10-19 (2021). https://doi.org:10.1038/s41589-020-00700-7

16 Pacesa, M. *et al.* R-loop formation and conformational activation mechanisms of Cas9. *Nature*
609, 191-196 (2022). https://doi.org:10.1038/s41586-022-05114-0

17 Xiao, Y. *et al.* Structure basis for directional R-loop formation and substrate handover
mechanisms   in   type   I   CRISPR-Cas   system.   *Cell.*   **170**,   48-60   (2017).
https://doi.org:10.1016/j.cell.2017.06.012

18 Selle, K. & Barrangou, R. Harnessing CRISPR-Cas systems for bacterial genome editing.
Trends Microbiol. **23**, 225-232 (2015). https://doi.org:10.1016/j.tim.2015.01.008

19 Baumann, K. Genome editing: CRISPR-Cas becoming more human. *Nat. Rev. Mol. Cell. Biol.*

18, 591 (2017). https://doi.org:10.1038/nrm.2017.84

20   Stella, S., Alcón, P. & Montoya, G. Class 2 CRISPR-Cas RNA-guided endonucleases: Swiss
Army knives of genome editing. *Nat. Struct. Mol. Biol.* **24**, 882-892 (2017).
https://doi.org:10.1038/nsmb.3486

21   Wang, J. Y., Pausch, P. & Doudna, J. A. Structural biology of CRISPR-Cas immunity and
genome editing enzymes. *Nat. Rev. Microbiol.* **20**, 641-656 (2022).
https://doi.org:10.1038/s41579-022-00739-4

22   Bikard D, E. C., Jiang W, et al. Exploiting CRISPR-Cas nucleases to produce
sequence-specific antimicrobials. *Nat. Biotechnol.* **32**, 1146-1150 (2014).
https://doi.org:10.1038/nbt.3043

23   Citorik RJ, M. M., Lu TK. Sequence-specific antimicrobials using efficiently delivered
RNA-guided nucleases. *Nat. Biotechnol.* **32**, 1141-1145 (2014).
https://doi.org:10.1038/nbt.3011

24   Kiga K, T. X., Ibarra-Chavez R, et al. Development of CRISPR-Cas13a-based antimicrobials
capable of sequence-specific killing of target bacteria. *Nat. Commun.* **11**, 2934 (2020).
https://doi.org:10.1038/s41467-020-16731-6

25   Selle K, F. J., Tuson H, et al. In vivo targeting of clostridioides difficile using phage-delivered
CRISPR-Cas3 antimicrobials. *mBio* **11**, e00019-00020 (2020). https://doi.org:10.1128/mBio

26   Rosenblum D, G. A., Kedmi R, et al. CRISPR-Cas9 genome editing using targeted lipid
nanoparticles for cancer therapy. *Sci. Adv.* **6**, eabc9450 (2020).

27   Neil, K. *et al.* High-efficiency delivery of CRISPR-Cas9 by engineered probiotics enables
precise microbiome editing. *Mol. Syst. Biol.* **17**, e10335 (2021).
https://doi.org:10.15252/msb.202110335

28   Datsenko, K. A. *et al.* Molecular memory of prior infections activates the CRISPR/Cas
adaptive bacterial immunity system. *Nat. Commun.* **3**, 945 (2012).
https://doi.org:10.1038/ncomms1937

29   Fineran, P. C. & Charpentier, E. Memory of viral infections by CRISPR-Cas adaptive immune
systems: acquisition of new information. *Virology.* **434**, 202-209 (2012).
https://doi.org:10.1016/j.virol.2012.10.003

30   Swarts, D. C., Mosterd, C., van Passel, M. W. & Brouns, S. J. CRISPR interference directs
strand specific spacer acquisition. *PloS ONE.* **7**, e35888 (2012).
https://doi.org:10.1371/journal.pone.0035888

31   Semenova, E. *et al.* Highly efficient primed spacer acquisition from targets destroyed by the
Escherichia coli type I-E CRISPR-Cas interfering complex. *Proc. Natl Acad. Sci. USA.* **113**,
7626-7631 (2016). https://doi.org:10.1073/pnas.1602639113

32   Vale, P. F. *et al.* Costs of CRISPR-Cas-mediated resistance in *Streptococcus thermophilus*.
Proc Biol Sci **282**, 20151270 (2015). https://doi.org:10.1098/rspb.2015.1270

33   Jiang, W. *et al.* Dealing with the evolutionary downside of CRISPR immunity: bacteria and
beneficial plasmids. *PLoS Genet.* **9**, e1003844 (2013).
https://doi.org:10.1371/journal.pgen.1003844

34   Stern, A., Keren, L., Wurtzel, O., Amitai, G. & Sorek, R. Self-targeting by CRISPR: gene
regulation or autoimmunity? *Trends Genet.* **26**, 335-340 (2010).
https://doi.org:10.1016/j.tig.2010.05.008

35   Vercoe, R. B. *et al.* Cytotoxic chromosomal targeting by CRISPR/Cas systems can reshape

bacterial genomes and expel or remodel pathogenicity islands. *PLoS Genet.* **9**, e1003454
(2013). https://doi.org:10.1371/journal.pgen.1003454

36 Musharova, O. *et al.* Systematic analysis of Type I-E *Escherichia coli* CRISPR-Cas PAM
sequences ability to promote interference and primed adaptation. *Mol. Microbiol.* **111**,
1558-1570 (2019). https://doi.org:10.1111/mmi.14237

37 Blosser, T. R. *et al.* Two distinct DNA binding modes guide dual roles of a CRISPR-Cas
protein complex. *Mol. Cell* **58**, 60-70 (2015). https://doi.org:10.1016/j.molcel.2015.01.028

38 Redding, S. *et al.* Surveillance and processing of foreign DNA by the *Escherichia coli*
CRISPR-Cas system. *Cell* **163**, 854-865 (2015). https://doi.org:10.1016/j.cell.2015.10.003

39 Redding, S. *et al.* Surveillance and processing of foreign DNA by the *Escherichia coli*
CRISPR-Cas system. *Cell.* **163**, 854-865 (2015). https://doi.org:10.1016/j.cell.2015.10.003

40 Xiao, Y., Luo, M., Dolan, A. E., Liao, M. & Ke, A. Structure basis for RNA-guided DNA
degradation by Cascade and Cas3. *Science.* **361**, eaat0839 (2018).
https://doi.org:10.1126/science.aat0839

41 van Erp, P. B. *et al.* Mechanism of CRISPR-RNA guided recognition of DNA targets in
Escherichia coli. *Nucleic Acids Res.* **43**, 8381-8391 (2015).
https://doi.org:10.1093/nar/gkv793

42 Nunez, J. K., Lee, A. S., Engelman, A. & Doudna, J. A. Integrase-mediated spacer acquisition
during CRISPR-Cas adaptive immunity. *Nature* **519**, 193-198 (2015).
https://doi.org:10.1038/nature14237

43 Jackson, R. N. *et al.* Crystal structure of the CRISPR RNA-guided surveillance complex from
Escherichia coli. *Science.* **345**, 1473-1479 (2014). https://doi.org:10.1126/science.1256328

44 Sinkunas, T. *et al.* Cas3 is a single-stranded DNA nuclease and ATP-dependent helicase in the
CRISPR/Cas immune system. *EMBO J.* **30**, 1335-1342 (2011).
https://doi.org:10.1038/emboj.2011.41

45 Semenova, E. *et al.* Interference by clustered regularly interspaced short palindromic repeat
(CRISPR) RNA is governed by a seed sequence. *Proc. Natl Acad. Sci. USA.* **108**,
10098-10103 (2011). https://doi.org:10.1073/pnas.1104144108

46 .   (!!! INVALID CITATION !!!).

47 Makarova, K. S., Zhang, F. & Koonin, E. V. SnapShot: Class 1 CRISPR-Cas Systems. *Cell.*
168, 946-946 (2017). https://doi.org:10.1016/j.cell.2017.02.018

48 Westra, E. R. *et al.* H-NS-mediated repression of CRISPR-based immunity in *Escherichia coli*
K12 can be relieved by the transcription activator LeuO. *Mol. Microbiol.* **77**, 1380-1393
(2010). https://doi.org:10.1111/j.1365-2958.2010.07315.x

49 Mitic, D., Radovcic, M., Markulin, D. & Ivancic-Bace, I. StpA represses CRISPR-Cas
immunity in H-NS deficient *Escherichia coli*. *Biochimie* **174**, 136-143 (2020).
https://doi.org:10.1016/j.biochi.2020.04.020

50 Sun, D. *et al.* Histone-like nucleoid-structuring protein (H-NS) paralogue StpA activates the
Type I-E CRISPR-Cas system againstnatural transformation in *Escherichia coli*. *Appl.*
Environ. Microbiol. **86**, e00731-00720 (2020). https://doi.org:10.1128/aem.00731-20

51 Yang, C. D., Chen, Y. H., Huang, H. Y., Huang, H. D. & Tseng, C. P. CRP represses the
CRISPR/Cas system in *Escherichia coli*: evidence that endogenous CRISPR spacers impede
phage P1 replication. *Mol. Microbiol.* **92**, 1072-1091 (2014).
https://doi.org:10.1111/mmi.12614

52 Pul, U. *et al.* Identification and characterization of *E. coli* CRISPR-*cas* promoters and their
silencing   by   H-NS.   *Mol.   Microbiol.*   **75**,   1495-1512   (2010).
https://doi.org:10.1111/j.1365-2958.2010.07073.x

53 Yosef, I., Goren, M. G., Kiro, R., Edgar, R. & Qimron, U. High-temperature protein G is
essential for activity of the *Escherichia coli* clustered regularly interspaced short palindromic
repeats (CRISPR)/Cas system. *Proc. Natl Acad. Sci. USA.* **108**, 20136-20141 (2011).
https://doi.org:10.1073/pnas.1113519108

54 Majsec, K., Bolt, E. L. & Ivančić-Baće, I. Cas3 is a limiting factor for CRISPR-Cas immunity
in   *Escherichia   coli*   cells   lacking   H-NS.   *BMC   microbiology*   **16**,   28   (2016).
https://doi.org:10.1186/s12866-016-0643-5

55 Patterson, A. G. *et al.* Quorum sensing controls adaptive immunity through the regulation of
multiple   CRISPR-Cas   systems.   *Mol.   Cell*   **64**,   1102-1108   (2016).
https://doi.org:10.1016/j.molcel.2016.11.012

56 Fang, F. C. & Rimsky, S. New insights into transcriptional regulation by H-NS. *Curr. Opin.*
Microbiol. **11**, 113-120 (2008). https://doi.org:10.1016/j.mib.2008.02.011

57 Corbett, D., Bennett, H. J., Askar, H., Green, J. & Roberts, I. S. SlyA and H-NS regulate
transcription of the *Escherichia coli* K5 capsule gene cluster, and expression of *slyA* in
Escherichia coli is temperature-dependent, positively autoregulated, and independent of H-NS.
J Biol Chem **282**, 33326-33335 (2007). https://doi.org:10.1074/jbc.M703465200

58 Lithgow, J. K., Haider, F., Roberts, I. S. & Green, J. Alternate SlyA and H-NS nucleoprotein
complexes control *hlyE* expression in *Escherichia coli* K-12. *Mol. Microbiol.* **66**, 685-698
(2007). https://doi.org:10.1111/j.1365-2958.2007.05950.x

59 Curran, T. D. *et al.* Identification of new members of the *Escherichia coli* K-12 MG1655
SlyA   regulon.   *Microbiology   (Reading)*   **163**,   400-409   (2017).
https://doi.org:10.1099/mic.0.000423

60 Nguyen Le Minh, P. *et al.* Differential protein-DNA contacts for activation and repression by
ArgP, a LysR-type (LTTR) transcriptional regulator in *Escherichia coli*. *Microbiol Res* **206**,
141-158 (2018). https://doi.org:10.1016/j.micres.2017.10.009

61 Lehnen, D. *et al.* LrhA as a new transcriptional key regulator of flagella, motility and
chemotaxis   genes   in   *Escherichia   coli*.   *Mol.   Microbiol.*   **45**,   521-532   (2002).
https://doi.org:10.1046/j.1365-2958.2002.03032.x

62 Sun, D. *et al. Escherichia coli* is naturally transformable in a novel transformation system.
FEMS Microbiol Lett **265**, 249-255 (2006). https://doi.org:10.1111/j.1574-6968.2006.00503.x

63 Sun, D. *et al.* Transforming DNA uptake gene orthologs do not mediate spontaneous plasmid
transformation   in   *Escherichia   coli*.   *J   Bacteriol*   **191**,   713-719   (2009).
https://doi.org:10.1128/jb.01130-08

64 Zhang, Y., Shi, C., Yu, J., Ren, J. & Sun, D. RpoS regulates a novel type of plasmid DNA
transfer   in   *Escherichia   coli*.   *PLoS   One*   **7**,   e33514   (2012).
https://doi.org:10.1371/journal.pone.0033514

65 Sasnauskas, G. & Siksnys, V. CRISPR adaptation from a structural perspective. *Curr. Opin.*
Microbiol. **65**, 17-25 (2020). https://doi.org:10.1016/j.sbi.2020.05.015

66 Kunne, T. *et al.* Cas3-derived target DNA degradation fragments fuel primed CRISPR
adaptation. *Mol. Cell* **63**, 852-864 (2016). https://doi.org:10.1016/j.molcel.2016.07.011

67 Breddermann, H. & Schnetz, K. Activation of *leuO* by LrhA in *Escherichia coli*. *Mol.*

Microbiol. **104**, 664-676 (2017). https://doi.org:10.1111/mmi.13656

68   Ishihama, A., Shimada, T. & Yamazaki, Y. Transcription profile of *Escherichia coli*: genomic
SELEX search for regulatory targets of transcription factors. *Nucleic Acids Res.* **44**,
2058-2074 (2016). https://doi.org:10.1093/nar/gkw051

69   Fineran, P. C. *et al.* Degenerate target sites mediate rapid primed CRISPR adaptation. *Proc.*
Natl Acad. Sci. USA. **111**, E1629-1638 (2014). https://doi.org:10.1073/pnas.1400071111

70   Xue, C. *et al.* CRISPR interference and priming varies with individual spacer sequences.
Nucleic Acids Res. **43**, 10831-10847 (2015). https://doi.org:10.1093/nar/gkv1259

71   Staals, R. H. *et al.* Interference-driven spacer acquisition is dominant over naive and primed
adaptation in a native CRISPR-Cas system. *Nat. Commun.* **7**, 12853 (2016).
https://doi.org:10.1038/ncomms12853

72   Cooper, L. A., Stringer, A. M. & Wade, J. T. Determining the Specificity of Cascade Binding,
Interference, and Primed Adaptation In Vivo in the *Escherichia coli* Type I-E CRISPR-Cas
System. *mBio* **9** (2018). https://doi.org:10.1128/mBio.02100-17

73   Høyland-Kroghsbo, N. M., Muñoz, K. A. & Bassler, B. L. Temperature, by controlling
growth rate, regulates CRISPR-Cas activity in *Pseudomonas aeruginosa*. *mBio* **9**,
e02184-02118 (2018). https://doi.org:10.1128/mBio.02184-18

74   Medina-Aparicio, L. *et al.* The CRISPR/Cas immune system is an operon regulated by LeuO,
H-NS, and leucine-responsive regulatory protein in *Salmonella enterica* serovar Typhi. *J*
Bacteriol **193**, 2396-2407 (2011). https://doi.org:10.1128/JB.01480-10

75   Stratmann, T., Pul, Ü., Wurm, R., Wagner, R. & Schnetz, K. RcsB-BglJ activates the
Escherichia coli leuO gene, encoding an H-NS antagonist and pleiotropic regulator of
virulence determinants. *Mol. Microbiol.* **83**, 1109-1123 (2012).
https://doi.org:10.1111/j.1365-2958.2012.07993.x

76   Yamanaka, Y., Aizawa, S. I. & Yamamoto, K. The *hdeD* Gene Represses the Expression of
Flagellum Biosynthesis via LrhA in *Escherichia coli* K-12. *J Bacteriol.* **204**, e0042021 (2022).
https://doi.org:10.1128/jb.00420-21

77   Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in *Escherichia*
coli K-12 using PCR products. *Proc. Natl Acad. Sci. USA.* **97**, 6640-6645 (2000).
https://doi.org:10.1073/pnas.120163297

78   Erijman, A., Dantes, A., Bernheim, R., Shifman, J. M. & Peleg, Y. Transfer-PCR (TPCR): a
highway for DNA cloning and protein engineering. *J Struct Biol.* **175**, 171-177 (2011).
https://doi.org:10.1016/j.jsb.2011.04.005

79   Ma, J. *et al.* iProX: an integrated proteome resource. *Nucleic Acids Res.* **47**, D1211-d1217
(2019). https://doi.org:10.1093/nar/gky869

a

cas3 P*cas* *cse1* *cas7* *cas6* *cas2* Leader n=12

cse2 *cas5* *cas1*

b

250 nt LBS II HBS TAACTTGCAG ATTTGAAATG CATGCATTAT TGTCTTTAA**A** CAATTCAACA +1 LBS I

59 nt 20 nt 60 nt

c

Streptavidin beads + Biotin-P*cas* DNA → Pull down

d

MW(kDa) Input NT biotin-P*cas* Input NT biotin-P*cas*

100
70
50
40
35 ◄LrhA
25
20
15

BW25113Δ*hns* MC4100Δ*hns*

e

■ Peptides identified by LC-MS/MS Coverage

MC4100 55.4%

N-term C-term

BW25113 33.0%

f

Pull down

MW(kDa) NT P*cas* Input

35 *
30 ◄His-LrhA
25
20

Anti-His Ab

g

P*cas*-*gfp* reporter

P*cas* *gfp*

ArgP
HexR
OmpR
NadR
YfhH
DeoR
McbR
AaeR
Vector
YdcI
SlyA P = 0.0020
LrhA P < 0.0001
GlpR P < 0.0001
LeuO P < 0.0001

0   2   4   6   8

Relative P*cas* activity

h

AaeR
YdcI
LrhA
LeuO
ArgP
AllR
YfhH
HexR
BluR

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5