Multiomic ALS signatures highlight sex differences and molecular subclusters and identify the MAPK pathway as therapeutic target

Lucas Caldi Gomes[1*], Sonja Hänzelmann[2,3,4*], Fabian Hausmann[2,3], Robin Khatri[2,3], Sergio Oller[2,3], Mojan Parvaz[1], Laura Tzeplaeff[1], Laura Pasetto[5], Marie Gebelin[6], Melanie Ebbing[2,3], Constantin Holzapfel[2,3], Stefano Fabrizio Columbro[5], Serena Scozzari[5], Johanna Knöferle[1], Isabell Cordts[1], Antonia F. Demleitner[1], Marcus Deschauer[1], Claudia Dufke[7], Marc Sturm[7], Qihui Zhou[8,13], Pavol Zelina[9], Emma Sudria-Lopez[9], Tobias B. Haack[7,10], Sebastian Streb[11], Magdalena Kuzma-Kozakiewicz[12], Dieter Edbauer[8,13], R. Jeroen Pasterkamp[9], Endre Laczko[11], Hubert Rehrauer[11], Ralph Schlapbach[11], Christine Carapito[6], Valentina Bonetto[5], Stefan Bonn[2,3§], Paul Lingor[1,8,13§,#]

Technical University of Munich, School of Medicine, rechts der Isar Hospital, Clinical Department of Neurology, Munich, Germany.
Institute of Medical Systems Biology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany.
Center for Biomedical AI, University Medical Center Hamburg-Eppendorf, Hamburg, Germany.
Department of Medicine, University Medical Center Hamburg-Eppendorf, Hamburg, Germany.
Research Center for ALS, Istituto di Ricerche Farmacologiche Mario Negri IRCCS, Milan, Italy
Laboratoire de Spectrométrie de Masse Bio-Organique, Université de Strasbourg, Infrastructure Nationale de Protéomique, Strasbourg, France.
Institute of Medical Genetics and Applied Genomics, University of Tübingen, Tübingen, Germany
German Center for Neurodegenerative Diseases (DZNE), München, Germany.
Department of Translational Neuroscience, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands.
Center for Rare Diseases, University of Tübingen, Tübingen, Germany.
Functional Genomics Center Zürich, ETH Zürich and University of Zürich, Zürich, Switzerland.
Department of Neurology, Medical University of Warsaw, Warsaw, Poland.
Munich Cluster for Systems Neurology (SyNergy), Munich, Germany.

*,§ These authors contributed equally.

#Corresponding Author: Paul Lingor; email: paul.lingor@tum.de; Clinical Department of Neurology, School of Medicine, rechts der Isar Hospital, Technical University of Munich, Ismaninger Straße 22, 81675 Munich, Germany, Tel.: +498941408257

42   **Abstract**
43
Amyotrophic lateral sclerosis (ALS) is the most common motor neuron disease and lacks
effective disease-modifying treatments. Here, we performed a multiomic analysis of the
prefrontal cortex of 51 patients with sporadic ALS and 50 control subjects, as well as four
transgenic mouse models of C9orf72-, SOD1-, TDP-43-, and FUS-ALS to characterize early
and sex-specific disease mechanisms in ALS. Integrated analyses of transcriptomes,
(phospho)proteomes, and miRNAomes revealed more pronounced changes in males. We
identified transcriptome-based human ALS subclusters driven by the immune response,
ECM, mitochondrial respiration, and RNA metabolism. The molecular signatures of human
subclusters were reflected in specific mouse models. Individual and integrative multiomics
analysis highlighted the mitogen-activated protein kinase pathway as an early disease-
relevant mechanism. Its modulation by trametinib in vitro and in vivo validated that mitogen-
activated protein kinase kinase 2 is a promising therapeutic target with beneficial effects in
females.

Introduction

Amyotrophic lateral sclerosis (ALS) is the most frequent motor neuron disease, is more common in males, and leads to paralysis and death within a few years of symptom onset on average[1]. While most cases are sporadic (sALS) with no family history, approximately 10% of cases have a genetic cause (fALS). The most common genetic variants include *C9orf72*, *SOD1*, *TARDBP*, and *FUS,* with around 10% of patients with sALS carrying disease-causing mutations[2]. The etiology of sALS remains unclear, and effective disease-modifying treatments for the disease are currently unavailable[3]. Enhancing our understanding of early disease mechanisms could help identify the diagnostic and prognostic biomarkers and uncover more efficient therapeutic drug targets. Although direct analysis of affected nervous system tissues remains the gold standard in neuropathology, patient samples are only available postmortem and in limited quantities. These limitations increase the risk of primarily describing end-stages of the disease and obscuring mechanisms occurring in earlier phases, which are potentially more suitable drug targets. As in Alzheimer's or Parkinson's disease, ALS is believed to spread over time—from the motor cortex to other cortical brain areas[4–6]. Numerous studies have analyzed the motor cortex in ALS[7–9]. However, as this area is the most severely affected by ALS, it primarily reflects the final stages of the disease[6]. In contrast, the prefrontal cortex (PFC) in Brodmann area 6 typically exhibits only intermediate TDP-43 pathology at the time of death[6], suggesting that analyzing this area could provide insights into earlier disease-mediated alterations.

Previous investigations of ALS brain tissue have mainly focused on individual molecular subsets, such as transcripts[7,9,10], miRNAs[11], and proteins[12], suggesting that ALS is a complex and heterogeneous disease. Recent studies have identified potentially distinct ALS populations, which were stratified into different subclusters based on transcriptomics and gene set enrichment analyses (GSEA)[9,13].

In this study, we deciphered early ALS disease mechanisms by profiling the transcriptomes, miRNAomes, and (phospho)proteomes in the PFCs of patients with ALS and four ALS mouse models. Our findings highlighted significant sex differences and demonstrated that ALS is not a homogeneous disease but comprises different molecular subtypes that correlate with individual transgenic mouse models of the disease. Multiomic data integration identified several known ALS disease mechanisms, but also less-prioritized pathways, such as the MAPK pathway. Validation studies in vitro and in vivo underscored MEK2 as a potential target for early therapeutic interventions.

Results

Cohort and data description

We conducted an in-depth multiomic characterization of the human postmortem PFC, (Brodmann area 6) from 51 patients with neuropathologically confirmed ALS and 50 control (CTR) patients, to describe early molecular changes in sALS (Fig. 1a, Table 1, Supplementary Table 1). On average, 19.641 transcripts, 736 miRNAs (mature miRNAs and hairpin precursors), and 2.344 proteins were detected per sample (Supplementary Fig. 1 a-c). A C9orf72 repeat expansion was detected in one patient with ALS, and another individual carried a pathogenic variant of *NEK1*(c.3107C>G, p.Ser1036Ter)[14] (Supplementary Fig. 2, Supplementary Table 2). For clarity and accuracy, we refer to all cases simply as "ALS". Four transgenic mouse models were analyzed to identify parallels with human ALS. PFCs from presymptomatic or early disease stages were collected from C9orf72-, SOD1-, TDP-43-, or FUS-ALS, aiming for an equal distribution of wild-type and transgenic littermates, as well as sexes (n = 20 per model, survival and sampling time points depicted in Supplementary Fig. 3). In mice, an average of 17.020 transcripts, 842 miRNAs (mature miRNAs and hairpins), 2.568 proteins, and 6.755 phosphosites were detected. The overall sample quality was consistently high for all omics analyses (Supplementary Fig. 4).

Transcriptomic analysis reveals sex-specific alterations in ALS-affected PFC tissue
Principal component analysis was performed to evaluate the impact of disease, sex, and sample origin on the transcriptome, revealing a moderate separation by condition (silhouette score: 0.11 [ALS]; -0.03 [CTR]) but a robust separation by sex (silhouette score: 0.29 [male], 0.15 [female]) (Fig. 1b). Consequently, we analyzed differentially expressed genes (DEGs) separately for males and females. Notably, the number of DEGs was significantly higher in males (n=73) than in females (n=2), which was confirmed by bootstrapping down sampling analysis (Fig. 1c, Supplementary Table 3 and 4). Similar to transcriptional changes, also differential alternative splicing (DAS) analysis revealed marked sex differences in the abundance of different splicing events. We observed more DAS events in male human samples (Fig. 1d). DAS was observed in males for *CLTB*, *TPRN*, *NRN1*, and *CAMK2N1* and in females for *TPRN* and the gene encoding TMEM170A-CFDP1*,* a novel readthrough protein (AC009163.5). TPRN, a stereocilium-associated protein previously described only in non-syndromic deafness, may play additional roles in the pathogenesis of ALS (Supplementary Table 5 and 6). Overrepresentation analysis using GO showed enrichment for *negative regulation of ERK1 and ERK2 cascade* and *negative regulation of MAPK cascade* (p.adj < 0.1) for male samples (Supplementary Table 5).
Enrichment analyses (KEGG) with transcriptomics results for males showed a significant enrichment for several *synapse-related* pathways (*retrograde endocannabinoid signaling, synaptic vesicle cycle, long-term potentiation, glutamatergic synapse*), as well as pathways related to *immune response*, *extracellular matrix (ECM)*, and diverse *protein processing* and *protein metabolism* terms (which are important components of the KEGG pathway for neurodegenerative diseases [hsa05022]) (Fig. 1e, Supplementary Figure 5, Supplementary Table 7). For females, the most significant results were related to *ribosomal function* and *oxidative phosphorylation,* as well as *mitochondria-related* terms. Further significantly enriched categories included *neurodegenerative disorders* and *cell adhesion molecules*. *Oxidative stress* was inferred from the enrichment of *oxidative phosphorylation* pathways in both males and females, and the *MAPK pathway* was frequently enriched for weighted gene co-expression network analysis (WGCNA[15]) (details in Supplementary Tables 7 and 8 and in the Methods section).

Stratification of human ALS into four molecular subclusters based on transcriptomic data
Based on frequently enriched terms for transcriptomics results (Supplementary Table 7), we conducted hierarchical clustering analyses for pathways of interest (Fig. 1f). This analysis revealed four distinct clusters labeled as C1–C4. The regulation of *immune response* served as a dichotomizing factor, distinguishing patients with ALS into C1 and C2 vs. C3 and C4. At the second level of arborization, *the ECM* played a primary role (C1 vs. C2), along with *synaptic function* and *protein folding* (C3 vs. C4, Fig. 1f). These clusters are reminiscent, but not identical to previously proposed subtypes[9], where C1 and C2 align with ALS-Ox (oxidative stress) and showing less resemblance to ALS-TE (elevated transposable element expression), while C3 and C4 correspond to the ALS-Glia (glial dysfunction) subtype (Supplementary Fig. 6).
Finally, to characterize the clusters using similarly regulated RNA networks, we performed WGCNA[15], resulting in 20 modules (Fig. 1g, Supplementary Table 8). The turquoise module was enriched for *mitochondrial respiration* and upregulated in C1 and C2, and driven by neuronal alterations, suggesting increased oxidative respiration in PFC neurons. The yellow module, enriched for *synaptic function,* exhibited a similar regulation in C1 and C2 (Fig. 1h). In contrast, the tan and lightcyan modules, enriched for *immune response* and *RNA splicing,*

158 were upregulated in C3 and C4 (Fig. 1g). These findings indicate that molecular subclusters
159 and sex-specific differences drive heterogeneity in the PFC of patients with ALS.
160
Male-predominant deregulation of miRNA expression
To explore the role of miRNA-mediated regulation in ALS[16], we analyzed small RNAs,
confirming previously identified sex differences. Male ALS patients exhibited more significant
downregulation in both mature miRNAs and miRNA hairpins compared to females (Fig.
2a,b). Specifically, males showed 17 mature DE miRNAs (15 downregulated, 2 upregulated)
while females exhibited 9 DE miRNAs (4 downregulated, 5 upregulated) (p < 0.1, fold
change > 1.5, Supplementary Table 9). Further analysis of miRNA hairpins highlighted sex-
dependent differences in early miRNA biogenesis. Notably, males demonstrated a more
pronounced dysregulation with 82 DE hairpins (71 downregulated, 11 upregulated), while
females displayed 13 DE hairpins (11 downregulated, 1 upregulated). This suggests
potential defects in miRNA biogenesis as an early disease mechanism.
To assess the significance of miRNA expression changes in our human disease clusters, we
conducted a cluster-specific DE analysis (Fig. 2c, Supplementary Table 10). Only miRNAs
with a minimum fold change of 0.1 and significance in at least two clusters were considered,
ensuring the identification of robust expression alterations. Clusters 1, 2, and 4 exhibited
distinctive miRNA profiles, while such changes were not observed in Cluster 3. In more
detail, miRNA miR-4472, modulating E cadherin and vimentin via *RGMA*, presented the
overall highest significant fold change in cluster 2. miR-181c-3p, a miRNA proposed as a
novel circulating biomarker for ALS[17], was downregulated in both clusters 1 and 2. In cluster
4 the miR-340-5p, which is neuronal injury-related, presented the highest fold change[18].
Overall, miRNA dysregulation appeared particularly important for clusters 2 and 4, while less
pronounced effects were seen in clusters 1 and 3.
183
Human proteomic proteomic analysis confirms stronger deregulation in males
We also assessed the proteomic signatures of early-stage ALS using mass spectrometry in
the same samples. Here, we detected 379 and 251 differentially expressed proteins (DEPs)
in males and females, respectively, with annexin A2 (ANXA2) being the only protein
downregulated in both sexes (*p* < 0.1). Interestingly, we identified several
neurodegeneration-related proteins, such as MATR3, SPART, and SCNA (involved in the
genetic forms of ALS, spastic paraplegia, and Parkinson's disease, respectively)[19–21] (Fig.
2d, Supplementary Table 11). The projection of transcriptomic clusters onto proteomic data
did not reproduce the sub clustering, likely because of the much smaller number of mapped
entities (Supplementary Fig. 7). Functional enrichment and unsupervised clustering identified
relevant pathways in both sexes, such as *synaptic function*, *immune response*, and
ECM/cytoskeleton (Fig. 2e,f). In contrast, *transmembrane transport, lipid metabolism*,
development, *catalytic activity* and *ERK1/2 signaling* were enriched in females, whereas *cell
metabolism* and *tyrosine kinase-related pathways* were enriched in males (Fig. 2e,f,
Supplementary Table 12).
199
Integration of multiomic data reveals sex-specific molecular networks of ALS
After using various omics modalities to identify the molecular pathways associated with early
ALS, we employed a biologically motivated approach that focused on identifying valid
interaction triplets involving miRNAs, transcripts, and proteins in male and female samples.
The triplet miR-769-3p–*ANXA2*–ANXA2 was particularly intriguing (Supplementary Fig. 8,
Supplementary Table 13), since ANXA2 is involved in angiogenesis and autophagy[22], which
are mechanisms known to be altered in ALS[23,24].
Next, we integrated transcriptomic, small RNA, and proteomic data using Multi-Omics Factor
Analysis (MOFA)[25]. Because sex was an important differentiating factor (Fig. 2g), MOFA was

performed by sex (Fig. 2h, Supplementary Fig. 9, Supplementary Table 14). In males, factor
1, mainly driven by hairpin miRNAs, explained 23.7% of the variance. Downregulation of
hsa-miR-7851, -1285-1, -5096, and a cluster of -1273 isoforms strongly contributed to its
weight (Supplementary Fig. 10). Transcriptome-based factor 3 correlated best with disease
condition and was driven by genes responsible for *vesicular function* (*RAB3C, NSF*), *cell*
survival (*BCL2, BHLHB9*), and *RNA metabolism* (*SNORA73B, RN7SL2*) (Fig. 2h).
Proteome-dominated factor 4 contains ZO2 and CD44, which are involved in myelination
and blood–central nervous system barrier (BCNSB) formation, respectively. Finally, factor 7,
which also showed a strong correlation with the disease condition, was dominated by
neurofilament heavy, medium, and light polypeptides (NfH, NfM, and NfL, respectively) as
well as proteins involved in Ca^{2+}-binding (HPCL4) and ECM formation (PGCA)
(Supplementary Table 14).
In females, factors 1–3 explained 42.6% of the variance, but factors 10 and 12 correlated
best with the disease condition (Fig. 2h). Synaptic genes, such as *RAB3C*, *NAPB*, and
SNAP25, contribute to factor 1 and are upregulated in ALS. The hsa-miR-1285-1, miR-5096,
and miR-1273 clusters also contributed to factor 2. Interestingly, MAPK1 plays a central role
in the miR-1273 target network (Supplementary Figure 11). Factor 3 (similar to male factor
4) included the oligodendrocyte and myelin markers CD44 and ZO2. Factor 10 contained the
antiproteases SERPINA1 and SERPINA3[26] as well as chitinases CHI3L1 and CHI3L2, which
are known biomarkers for ALS[27]. Factor 12 showed negative correlation with NFH and
MAPK–ERK1/2 regulator PEA15. MOFA analysis underlined the mechanisms identified in
the individual omics analyses, such as the downregulation of miRNA clusters (particularly in
males, Supplementary Fig. 10), ECM components, and oligodendrocyte and myelin markers.
This unbiased data integration strategy highlighted known ALS biomarkers (neurofilaments
and chitinases) and the MAPK pathway, especially in females, as important molecular hubs
(Supplementary Fig. 10-12, Supplementary Table 14).
235
Murine models of ALS partially recapitulate human ALS
We used four mouse models representing the most frequent disease-causing mutations,
C9orf72, SOD1, TDP-43, and FUS and performed sex-specific analyses for male and female
animals. Strikingly, the C9orf72 model exhibited the most pronounced transcriptomic
changes, mirroring the rapid disease progression observed in this model (Fig. 3a,
Supplementary Fig. 3). In each mouse model, we found differentially regulated pathways
that were also identified in the human cohort: The C9orf72 model prominently featured
alterations in *immune* and *inflammatory response* pathways, the SOD1 model in the *ERK1/2*
cascade, *development*, and *response to oxidative stress,* whereas *transcription* and
endopeptidase activity were altered in the TDP-43 model. In contrast, the FUS model
exhibited the least transcriptional deregulation without enrichment of pathways (Fig 3b,
Supplementary Table 15).
Further, transcriptome-based analyses were performed to explore DAS in our mouse models
(Supplementary Fig. 13, Supplementary Table 6). It is well documented that TARDBP/TDP-
43 and FUS regulate alternative splicing and transcript usage in hundreds of genes[28,29].
TDP-43 also represses cryptic exon splicing events in *STMN2*[30] and *UNC13A*[31]. We were
not able to detect any significant cryptic splicing events for these two genes in our data.
However, it is important to note that such cryptic exon inclusion events are specific to
neurons with TDP-43 pathology[32]. They are also not conserved in mice[31] and are rarely
detectable in bulk RNA-Seq at our sequencing depth[33]. Despite the absence of notable
cryptic splicing events, our analysis revealed differential splicing in important genes such as
FLNB, CPLANE1 (involved in ciliogenesis and migration), and *ATP1B1* (a membrane-bound
Na^+/K^+-ATPase) in multiple mouse models (Supplementary Table 6). DAS analyses provided
additional insights into sex-specific variations for the mouse cohorts (Supplementary Fig. 13)

showing enrichment in the terms *mitochondria* and *myelin sheath* (C9orf72), *GTPase activity*
and *myelin sheath* (SOD1), *DNA binding* and *heat shock protein binding* (TDP-43), and
protein binding and *ribosome* (FUS) (Supplementary Table 5).
To assess if the observed alterations in gene expression and protein abundance reflect
changes in the cellular composition of the PFC, we estimated the cell-type fractions for the
subclusters of human and mouse ALS models using deep learning-based deconvolution[19]
from the gene expression data (Fig. 3c, Supplementary Fig. 14). Interestingly, the human
subclusters showed changes in the cell fractions that were reflected in the individual mouse
models. The SOD1 model and human C1 and C2 clusters showed a decrease in glial and
endothelial cells and a relative increase in excitatory neurons (Fig. 3d). The C9orf72 model
and human C3 were characterized by a strong increase in glial and endothelial cells and a
decrease in excitatory neurons, respectively, suggesting strong neuroinflammation and
neuronal loss. The FUS and TDP-43 models showed intermediate levels of glial and
neuronal cells, also assessed by cell type enrichment in WGCNA modules (Fig. 3d,
Supplementary Fig. 14-15). Thus, the observed transcriptional changes are partially driven
by cell composition. Interestingly, our data suggest that the neurovascular unit in ALS[20] may
be affected differently in different subclusters of patients with ALS. We then correlated the
deconvolved fractions of human and mouse with each other. Overall, our transcriptome
analyses revealed correlations between human clusters and mouse models: C1 and C2
showed the best correlation with the SOD1 model (C1: 0.11 and C2: 0.42), whereas C3
correlated best with the C9orf72 model (0.31) and, to a lesser extent, TDP-43 (0.23) and
FUS (0.14) models. Finally, C4 showed a weak correlation with the FUS model (0.14).
Overall, correlation analyses show remarkable similarities between human clusters and
mouse models (Fig. 3d).
To further correlate molecular alterations in mouse models with our human data, we
investigated the proteome in PFC samples, where again sex-specific differences were
reproduced (Supplementary Table 11, Supplementary Fig. 16). Similar to transcriptomic
data, the strongest changes were observed in the C9orf72 model. Sequestosome 1/p62, a
product of the ALS-causing SQSTM1[34], showed the strongest upregulation in C9orf72
males, indicating reduced autophagic flux in a dipeptide accumulation model[35]
(Supplementary Fig. 16). SOD1 mice showed one upregulated DEP, exportin-1 (XPO1), a
major regulator of nuclear RNA export. XPO1 was also among the significantly regulated
proteins in the C9orf72 model (Supplementary Table 11). Unsupervised clustering for
enrichment analysis of proteomic mouse data using REVIGO revealed multiple pathway
clusters that were also identified in the human data. For example, most models presented
clusters or individual terms related to *synaptic function*. Other enriched pathways include
cytoskeleton and morphogenesis (for C9orf72 females, SOD1 and FUS males);
mitochondria/cell respiration/cell metabolism (present for C9orf72 females, TDP43 females,
FUS males and SOD1 for both sexes). A cluster for the *MAPK cascade* was captured for
SOD1 animals (males), which also exhibited enrichment for terms related to *endocytosis and*
regulation of cellular transport (Supplementary Fig. 17).
To further compare the representation of human ALS-related proteomic changes in murine
models, we performed a comparative clustering analysis of Gene Ontology (GO) term in
semantic space[36]. Human proteomics results showed prominent clustering for *differentiation*
and *development* (females: groups 3, 4, 7, and 11; males: group 8), *synapse* (females:
groups 16 and 17; males: group 2, 10, 15)*, and *immune/defense response* (females: group
12; males: groups 3, 4, 6) (Fig. 3e). In contrast, the C9orf72 model showed clustering for
RNA processing, ribosome, translation, *ATP synthesis*, *development*, *cell adhesion*,
transport, and *synapse*. The SOD1 model showed strong clustering for *ATP synthesis,*

mitochondrial respiration, *translation*, and *vesicle-mediated transport*. Thus, enrichment
results underscored the pathways previously identified in the RNA sequencing data.
In addition, multiple mouse omics datasets, including phospho-proteomics, were also
integrated to visualize valid interacting partners based on their expression levels in so-called
quadruplets and identified a coherent regulation of XPO1, GFAP, SQSTM1, and ATXN2L
(Supplementary Fig. 18, Supplementary Table 13). Several different phosphosites of
SQSTM1 were also found significantly upregulated in the phosphoproteomic analysis of
C9orf72 mice (Supplementary Fig. 18).
317
Alterations of the MAPK/ERK pathway are identified across multiple omics layers in
mouse models and human patients
Finally, we wanted to use the multiomic data generated from humans and mice to identify
signaling pathways that could represent novel therapeutic targets for ALS. Although multiple
molecular pathways identified in our analysis merit therapeutic validation, we decided to
focus on the MAPK/ERK pathway, which was altered consistently across several data types,
integration methods, and different subclusters. For example, *ERK1/2 signaling* was enriched
in human females, whereas *cell metabolism* and *tyrosine kinase* pathways were enriched in
males (Fig. 2e-f, Supplementary Table 12). Integrative analyses performed with MOFA also
identified different interactors of the MAPK/ERK pathway: PEA15 and isoforms of miR-1273
were important contributors for the weight of different MOFA factors for both males and
females, and directly regulate MAPK signaling (Supplementary Fig. 10-12, Supplementary
Table 9 and 14). Enrichment analyses with transcriptomics data showed disruptions in the
ERK1/2 cascade for the SOD1 mouse model and DAS results for human males showed
enrichment for terms related to the ERK1/ERK2 cascade and the MAPK cascade
(Supplementary Table 15). To explore alterations of the MAPK pathway in more depth, we
analyzed mouse and human transcriptomics with WGCNA, showing the contribution of
MAPK signaling in multiple modules. While deregulations in the MAPK cascade are
observed for all mouse models and the human cohort (both for significant levels and
absolute counts), the C9orf72 model presented the most striking changes (Fig. 4a). Next,
based on the DAS analyses for human and mouse cohorts, we calculated enrichment
analyses across all species and sexes and grouped the top resulting pathways in five major
groups (Fig. 4b). Among them, MAPK clearly shows a sex specific expression in SOD1
males - corroborating the enrichment results from transcriptomics analyses for this model,
while human males showed enrichment for more general terms related to kinase-activity. We
also carried out a miRNA expression analysis, which identified miRNA-451a consistently
deregulated across mouse models and human samples with ALS (Fig. 4c, Supplementary
Fig. 19). Target prediction analyses revealed that miRNA-451a interacts with multiple
members of the MAPK pathway (Fig. 4d), including MAPK1, AKT1, and BCL2, which are
known for their roles in the regulation of cell growth, survival, and apoptosis[37]. In addition,
centrality measurements for targets of this miRNA showed MAPK1 as the top hit, followed by
AKT1, IKBKB, BCL2 and MYC. This indicates a pivotal role of miR-451a in the context of
MAPK signaling (Fig. 4e).
We next investigated the role of the MAPK pathway in human subclusters C1-C4.
Interestingly, the classical MAPK pathway was activated in C1 and C2 and downregulated in
C3. Meanwhile, C1 and C2 showed downregulation of the JNK and p38 MAPK pathways,
which were activated in C3 (Supplementary Fig. 20). In addition to individual omics, also
MOFA-results from the ALS-associated factor 12 pointed towards a prominent role of the
MAPK pathway, where multiple of its members contributed to the PPI network (Fig. 4f).
Overall, many of the analyses pointed towards a significant role of the MAPK pathway in
ALS, specifically ERK1/2, across organisms and data types.
359

Validation of MEK2 inhibition by trametinib in vitro and in vivo

Currently, licensed drugs target *glutamatergic synapse function* (riluzole), *oxidative stress* (edaravone), *mitochondrial function* (tauroursodeoxycholic acid/phenylbutyrate), or SOD1 itself (tofersen). Other mechanisms identified in this analysis, such as *immune response*, *ECM/BCNSB function*, or the *MAPK/ERK* pathway are not yet targeted by licensed drugs. We concentrated on mitogen-activated protein kinase kinase 2 (MAP2K2 or MEK2), as it appeared to be upregulated in the human PFC and multiple mouse models. Furthermore, MEK2 can be modulated by the clinically approved inhibitor, trametinib[38]. A scheme for its mechanism of action and the expected effects in the context of the MAPK pathway is depicted in Fig. 4g.

First, we used primary cortical neuronal cultures from P0–P1 C57BL/6J mice that were treated with glutamate as an in vitro model of excitotoxicity in ALS[39]. Treatment with 5 mM glutamate (6 hours) did not affect total MEK2 or ERK1/2 protein expression, but significantly increased pErk1/2 levels. Trametinib attenuates glutamate-induced phosphorylation of Erk1/2. We observed a significant increase of pMEK2 activation in response to trametinib treatment. This could be explained by inhibition of the feedback circuit: MEK2 inhibition induces a decrease of pERK1/2 (Fig. 5), which also counteracts the feedback loop, resulting in the activation of the pathway and the accumulation of activated pMEK2[40,41]. Furthermore, glutamate treatment also increased cell death (caspase-positive neurons) and reduced the average neurite length, both of which were counteracted by trametinib (Fig. 5a–d, Supplementary Figs. 21 and 22). Our data suggest that trametinib attenuates MEK2 activity and reduces Erk1/2 phosphorylation resulting in decreased cell death and increased neurite outgrowth under excitotoxic stress.

To validate the importance of the MAPK pathway in vivo, we selected the SOD1 mouse model, which showed the strongest similarity with the largest human ALS subcluster (C1 and C2) (Fig. 3c-d). In females, we observed pMEK2 levels substantially increased with disease progression, whereas in males, pMEK2 levels returned to control levels after week 14 (Fig. 6a). Furthermore, in an initial study, we treated SOD1 mice with trametinib for seven weeks, starting from week 9 (presymptomatic stage), and observed a reduction in ERK1/2 phosphorylation compared with that in vehicle-treated female and male mice (Fig. 6b). Trametinib significantly reduced the autophagy receptor p62 expression in the spinal cord of female but not male mice, correlating with its previously described neuroprotective role of increasing autophagy via transcription factor EB activation[42] (Fig. 6c). p62 also co-localizes with ubiquitin and mutant SOD1 in protein aggregates[43]. Accordingly, we detected a significant reduction in detergent-insoluble SOD1 and ubiquitin in trametinib-treated females but not in males (Fig. 6d-e). Finally, we investigated the effect of trametinib on neurodegeneration. NfL plasma concentration[44] was significantly reduced after trametinib-treatment in female SOD1 mice (Fig. 6f). We also observed strong MEK2 phosphorylation in the motor neurons of the spinal cord, the main tissue involved in the pathology of this animal model[45] (Fig. 6g-h). Phospho-proteomic analyses supported the regulation of the MAPK pathway by trametinib in females, demonstrating in addition a reduced phosphorylation of its upstream regulator CAMK2A (Supplementary Fig. 23).

We thus focused on female mice to assess the effect of trametinib on the onset and progression of the disease. In a follow-up study, a second group of female SOD1 mice was treated with the same schedule up to survival. Body weight monitoring and motor tests (grip strength, extension reflex) were performed in comparison with vehicle-treated mice. Although no effect was observed in body weight loss and behavioral analysis (data not shown), there was a significant delay in the age of paralysis (Fig. 6i), onset of the disease, and a significantly improved survival (Fig. 6j). We also observed that a significant effect on NfL plasma concentration was maintained along the progression of the disease (Fig. 6k).

410 Thus, trametinib markedly affects the clearance of protein aggregates, leading to
411 neuroprotection in female SOD1 mice, suggesting that MEK2 is a promising therapeutic
412 target for ALS, particularly in females.
413
Discussion
In this study, we conducted individual and combined analyses of multiple omics datasets to
comprehensively understand the molecular architecture of ALS in the PFC, an area affected
in the later stages by TDP-43 pathology and that has the potential to reveal early disease
mechanisms[6]. We observed that significant sex-specific differences are often more
pronounced in males, which is aligned with the higher prevalence of ALS in males[1]. Previous
studies have also identified sex-specific differences in the blood of patients with ALS[46,47] as
well as therapeutic responses in ALS mouse models[48,49]. However, current therapeutic
options, clinical trials, and Food and Drug Administration guidelines for clinical trial design[40]
do not consider patient sex in differential therapy. Our data, especially the sex-specific
differences in MAPK signaling and therapeutic tractability, argue for a stronger consideration
of sex as a covariate in ALS clinical trials.
Phenotypic heterogeneity is clearly recognized in ALS[50], as reflected by stratification in onset
or disease progression rate, and previous transcriptomic analyses have suggested
molecular subtypes[13]. We identified four clusters in our human ALS cohort based on the
transcriptome, partially mirroring previously identified ALS-Ox, ALS-Glia, and ALS-
Transposable elements subtypes[9]. While the ALS-Glia and ALS-Ox clusters correlated with
clusters C1 and C2 and C3 and C4, respectively, the ALS-TE cluster was only marginally
represented in our data. As we analyzed the PFC, an area wherein TDP-43 aggregation is
observed later than that in the motor cortex, our data reinforce that ALS-TE is driven by
TDP-43 dysfunction[9]. Because alterations in the PFC may represent earlier changes in ALS,
clustering differences between the two studies may also reflect the evolution of clusters in
different stages of the disease. Molecular subtypes of ALS may evolve with time and could
be subject to changes during the disease course. Exploring liquid biomarkers could further
aid in patient stratification and personalized clinical trials.
Furthermore, we suggest that the molecular phenotype in the four analyzed mouse models
can be approximated to these subtypes, making these models potential surrogates for the
molecular subgroups in humans. Notably, the oldest and most frequently used model, the
SOD1.G93A mouse, correlated best with the largest clusters C1 and C2. Although not
representative of all human ALS cases, our findings indicate that this model represents an
important subgroup of the disease based on the identified dysregulated pathways.
Integrated analysis of multiomic data sets has identified several dysregulated pathways
relevant for ALS, including mitochondrial respiration/oxidative stress, transcriptional
regulation/splicing, and protein misfolding/degradation[3]. Additionally, previously less
prioritized pathways—including the dysregulation of the ECM and BCNSB or the MAPK
pathway—were identified. The deregulation of pathways that are only distantly related
suggests the need for combinatorial drug therapies to address ALS mechanistically. It is
important to highlight that we found evidence of female-predominant deregulation of the
MAPK pathway in integrative analyses, but it did not contribute to the separation of clusters
C1–C4. Therefore, the MAPK pathway may be an interesting therapeutic target for all human
ALS subgroups. MAPKs are fundamental signal transducers that are involved in cell
proliferation, differentiation, survival, and death, and as such not specific for ALS[51].
Extracellular and intracellular signals are integrated by MAPK, and the overactivation of
MAPK signaling, for example, via the abnormal phosphorylation of ERK1/2, has been
reported in human and ALS mouse models[52]. A Ser/Thr kinase belonging to the MAPK
superfamily—the MAPK/MAK/MRK [MOK] kinase—has also been linked to impaired

microglial function and inflammatory insults associated with ALS pathophysiology[53].
Increased phosphorylation of ERK1/2 was also observed in our glutamate toxicity model,
and pMEK2 expression was increased in the SOD1 mouse model, reproducing the aberrant
activation of this pathway that was restored by the MEK2-inhibitor trametinib. However, this
effect was most pronounced in females, suggesting a sex-specific efficacy. Notably, our
experimental approach did not target neurons exclusively. Nevertheless, our primary
cultures mainly contained neurons (∼95%), and all staining procedures for both in vivo and
in vitro experiments focused on neurons. While other cell types may be involved, the
evidence presented here supports the significance of the MEK2–MAPK pathway in neurons.
Indeed, trametinib affected the onset and the survival of the disease in the SOD1 mice, but
not the rate of disease progression, in which non neuronal supporting cells are thought to
have a role[54]. Although the MAPK pathway may represent a less specific answer of the cell
in response to stressors, its involvement in ALS makes it a promising target, which could add
another step in a combination therapy. Currently, one phase I/II clinical trial is examining the
safety, tolerability, and efficacy of trametinib in patients with ALS (clinicaltrials.gov
identifier:NCT04326283). Our data support the independent evaluation of male and female
patients and suggest that the ECM, immune response, and RNA processing machinery are
potential targets for therapeutic intervention that need further exploration.
Despite the limitations of our study (e.g., the retrospective view permitted by postmortem
tissue analysis and restricted balancing of sex and disease stage due to the limited number
of well-characterized postmortem brain samples), we aimed to provide valuable insights into
the molecular architecture of ALS. While analyzing over 100 brain samples helped identify
molecular subclusters, a larger sample size may provide a more detailed view of the
subgroups. Nevertheless, analyzing the PFC tissue allowed us to study the earlier stages of
the disease, offering a unique perspective compared with those of previous studies. Our
robust multiomic, computational, and integrative approaches identified subclusters that are
reminiscent of previously identified subtypes, yet show clear differences. Additionally, we
emphasize the role of splicing and transcript usage in early disease regulation, involving
several ALS-related genes. Overall, our study contributes towards unraveling the complexity
of ALS and lays the groundwork for further research in the field.
Our study also compared human brain tissue with samples from transgenic mouse models of
ALS. Although clear correlations were found between mouse models and molecular
subtypes of sALS, the four analyzed models represent specific scenarios. Other models for
each of the four studied genes could potentially yield different results[55]. Furthermore, we did
not analyze DNA methylation status or other post-translational modifications (such as
glycation, methylation, or acetylation) that could provide further insights into ALS-specific
dysregulation, as much as a single cell-based analysis could yield an additional layer of
detail.
In this thorough integration of human tissue- and animal model-derived multiomic data, we
emphasized sex-specific differences in ALS pathology, identified molecular clusters, and
highlighted their importance for clinical trials and the development of novel therapeutic
strategies. The validation of omics results relies on multiple systems, as each mouse model
system only reproduces parts of the human pathology. Our data suggest the need for the
validation of additional molecules and pathways and justify the further exploration of the
MAPK pathway as therapeutic ALS treatment targets.
505
506 **Methods**
507 **Human postmortem PFC samples**
508 Human PFC samples were provided by four different brain banks: The Netherlands Brain
509 Bank (NBB), London Neurodegenerative Diseases Brain Bank (London NDBB), Imperial

College London - Multiple Sclerosis and Parkinson's Tissue Bank (ICL MS&PD TB), and the Oxford Brain Bank (OBB). In total, 51 ALS and 50 CTR samples (without any signs of neurodegeneration) were included. Frozen tissues were shipped on dry ice to the rechts der Isar Hospital Department of Neurology at the Technical University of Munich and stored at -80 °C (Supplementary Table 1). Ethical approval for the use of human tissues was obtained from the Ethics Committees of the University Medical Center Göttingen (2/8/18 AN) and the Technical University Munich (145/19 S-SR). For sampling, PFC samples were transferred to a cryostat chamber at -20 °C and punched with a 20-G Quincke Spinal Needle (Becton Dickinson, Franklin Lakes, NJ, USA). Subsequently, ~20 mg tissue was collected in RNAse-/DNAse-free tubes and stored at -80 °C until further use.

ALS animal models

Four transgenic mouse models representing the most frequent ALS-causing genes were used for the multiomic studies. Animal handling and experiments in this study were performed in accordance with the applicable animal welfare laws and approved by the respective regulatory organs of the involved research centers. The animals were kept in standard cages in a pathogen-free facility under a 12-hour light/dark cycle. The mice were provided with unrestricted access to food and water. B6;129S6-Gt(ROSA)26Sortm1(TARDBP*M337V/Ypet)Tlbt/J mice[56] (here simply referred to as TDP-43 mice) were provided by the Department of Translational Neuroscience of the University Medical Center Utrecht. This model was generated by inserting an 80 kb genomic fragment carrying the human TDP-43 locus (including a patient-derived M337V mutation). TDP-43 transgenic and control wild-type mice were euthanized at the age of 26 weeks (presymptomatic stage) for biomaterial collection. B6SJL-Tg(SOD1*G93A)1Gur/J mice[45] (here referred to as SOD1 mice) were provided by the Laboratory of Translational Biomarkers, Mario Negri Institute for Pharmacological Research, Milan. High-copy number B6 congenic Tg(SOD1*G93A)1Gur/J SOD1*G93A male mice from The Jackson Laboratory (Bar Harbor, ME, USA) were bred with C57BL/6 female mice to obtain non-transgenic and mutant transgenic G93A*SOD1-expressing mice. SOD1 transgenic and control animals were sacrificed 14 weeks after birth (presymptomatic stage). (Poly)GA-NES/C9orf72(R26(CAG-lsl-175GA)-29×Nes-Cre) mice[47] (here referred to as C9orf72 mice) were provided by the German Center for Neurodegenerative Diseases in Munich. To generate transgenic animals, plasmids were electroporated to enable the conditional expression of dipeptide-repeat proteins (DPRs). DPRs were created by inserting 175 GFP-(GA) genes encoded by non-repeating alternate codons downstream of a floxed stop cassette in the pEX CAG stop-bpA vector. Electroporation was performed on murine recombination-mediated cassette exchange embryonic stem cells at the Rosa26 safe harbor locus. Mouse lines with germline transmission, known as GAstop, were obtained and subsequently backcrossed to a C57BL/6J background until a confirmed purity of over 98% was achieved (using SNP genotyping). C9orf72 transgenic and control animals were sacrificed 4.5 weeks after birth (early symptomatic stage). Tg (Prnp-FUS)WT3Cshw/J mice[57] (hereafter referred to as FUS mice) were provided by the Laboratory of Translational Biomarkers, Mario Negri Institute for Pharmacological Research, and were sacrificed four weeks after birth. For each model, ten transgenic and ten non-transgenic mice were selected and balanced for sex.

Preparation of PFC from ALS mouse models

The mice were perfused with 50 mL of ice-cold phosphate-buffered saline (PBS) prior to microdissection. The head was removed by cutting the base of the skull followed by skin removal. The skull was removed using small incisions followed by microdissection of the PFC in both hemispheres. The olfactory bulb and cerebellum were removed by cutting the

560 cerebellar peduncle starting from the olfactory bulb and continuing along the
561 interhemispheric fissure using tweezers with fine tips. The cortex was removed from the rest
562 of the brain. Incisions were made in the middle of the cortex to excise the PFC
563 (Supplementary Fig. 24). Freshly prepared PFCs were collected in nuclease-free tubes and
564 stored at -80 °C until RNA and protein isolation experiments were performed.
565
RNA and DNA isolation from human and mouse tissue samples
Total RNA was isolated from human and animal PFC samples using TRIzol Reagent
(Sigma-Aldrich, Taufkirchen, Germany). To ensure proper handling of the RNA, all related
experiments were conducted in an RNA workstation fume hood. In brief, 500 µL TRIzol was
added to each sample, and the tissues were homogenized using a plastic homogenizer.
Subsequently, 50 µL of 1-bromo-3-chloro-propane (Sigma-Aldrich) was added. The reaction
tubes were inverted for 10–15 s and incubated at room temperature (22–25 °C) for three
minutes. The resulting lysates were centrifuged at 12 000 × g for 15 min at 4 °C, leading to
phase separation. The RNA-containing aqueous phase was carefully collected and
transferred to a fresh nuclease-free tube. To precipitate the RNA, 250 µL of 2-propanol
(AppliChem, Darmstadt, Germany) and 2 µL of GlycoBlue Coprecipitant (15 mg/mL)
(Thermo Fisher Scientific, Waltham, MA, USA) were added to each sample. The mixture
was thoroughly vortexed and incubated overnight at -20 °C. Samples were centrifuged at 12
000 × g for 30 min at 4 °C the following day. The supernatant was discarded, and the RNA
pellets were washed three times with 75% ice-cold ethanol (AppliChem). After air-drying the
pellets for a few minutes under a fume hood, they were reconstituted with 15–20 µL
nuclease-free water (Sigma-Aldrich). The reconstituted RNA was completely dissolved by
incubation at 55 °C for two minutes in a thermoshaker. Following RNA isolation, DNase
treatment was performed to eliminate DNA contamination. For this purpose, 5 µL 10×
DNAse I Incubation Buffer (Life Technologies, Carlsbad, CA, United States), 5 µL DNase I (2
U/µL), and 0.5 µL RNaseOUT (40 U/µL) were added to each sample. Nuclease-free water
was added to a final volume of 50 µL, and the samples were then incubated at 37 °C for 20
min. Finally, the RNA samples were purified and concentrated using an RNA Clean &
Concentrator-5 Kit (Zymo Research, Irvine, CA, USA) following the manufacturer's
instructions. A QIAamp DNA Mini Kit (Qiagen, Hilden, Germany) was used for DNA isolation
from human midbrain samples according to the manufacturer's instructions. After RNA and
DNA isolation, the concentration and purity of nucleic acids were assessed using a
NanoDrop One spectrophotometer (Thermo Fisher Scientific). RNA integrity was evaluated
using an Agilent 6000 NanoKit on a 2100 Bioanalyzer (Agilent, Santa Clara, CA, USA).
595
DNA sequencing experiments and C9orf72 repeat expansion analysis
Prior to the DNA sequencing experiments, the quality of the isolated DNA was determined
using a 4200 TapeStation System (Agilent). The AmpliSeq protocol (Illumina, San Diego,
CA, USA) was used in subsequent steps. We used a target panel of 566 amplicons covering
30 ALS-related genes: *TARDBP, DCTN1, ALS2, ERBB4, TUBA4A, CHMP2B, NEK1,*
MATR3, SQSTM1, FIG4, C9orf72, SIGMAR1, VCP, GLE1, SETX, OPTN, HNRNPA1,
KIF5A, TBK1, ANG, SPG11, CCNF, FUS, PFN1, MAPT, VAPB, SOD1, CHCHD10, NEFH,
and *UBQLN2.* Using this panel, DNA samples (50–100 ng) were amplified over 14 PCR
cycles. The amplicons were digested, and AmpliSeq combinatorial dual indices were ligated
for multiplexing. The quality and quantity of the enriched libraries were validated using a
4200 TapeStation System (Agilent). The average fragment size of the amplified product was
480 bp. The libraries were standardized to a concentration of 9 nM in Tris–Cl (10 mM, pH
8.5) supplemented with 0.1% Tween 20. Cluster generation and sequencing were performed
using the MiSeq platform (Illumina) following the standard protocol. The loading

concentration was 9 pM, and 15% phiX was added. The sequencing configuration was a
250-bp paired-end sequencing. C9orf72 repeat expansion analysis was performed using the
AmplideX PCR/CE C9orf72 Kit (Asuragen, Austin, TX, USA). Briefly, DNA samples (n = 51,
40 ng each) were amplified using the three-primer GGGCC-repeat-primed configuration
(combining flanking primers and a GGGCC-repeat-specific primer). This configuration allows
the sizing of GGGCC alleles of up to 145 repeats and simultaneous detection of expanded
GGGCC alleles with > 145 repeats. The PCR conditions consisted of initial denaturation (five
minutes, 98 °C), 37 cycles (35 s, 97 °C; 35 s, 62 °C; and 3 min, 72 °C), and final elongation
(10 min, 72 °C). Capillary electrophoresis was carried out on an ABI 3730 DNA Analyzer
(Applied Biosystems, Waltham, MA. USA) using a ROX 1000 Size Ladder (Asuragen),
followed by analysis with GeneMapper 4.0 software (Applied Biosystems) and conversion of
peak size to GGGGCC repeat length via the calibration curve method according to the
manufacturer's instructions.
623

Preparation of RNA libraries for sequencing experiments
The mRNA and small RNA sequencing experiments were conducted at the Functional
Genomics Center, Zürich. For mRNA sequencing, two different library preparation kits were
used: the TruSeq Stranded mRNA Kit (Illumina; short read sequencing) and the SMARTer
Stranded Total RNA-Seq Kit v2 Pico Input Mammalian (Takara Bio USA, San Jose, CA,
USA; short read sequencing). In the TruSeq protocol, total RNA samples (100–1000 ng)
were subjected to poly-A enrichment and reverse transcription to generate double-stranded
cDNA. The cDNA was fragmented, end-repaired, adenylated, and ligated to TruSeq
adapters containing unique dual indices for multiplexing. Fragments with adapters at both
ends were selectively enriched using PCR amplification, resulting in a smear with an
average fragment size of ∼260 bp. The libraries were subsequently normalized to 10 nM in
Tris–Cl (10 mM, pH 8.5) containing 0.1% Tween 20. For the SMARTer Stranded Total RNA-
Seq Kit protocol, total RNA samples (0.25–10 ng) were reverse-transcribed using random
priming into double-stranded cDNA in the presence of a template switch oligo (TSO). The
generated cDNA fragments contained sequences derived from random priming oligos and
the TSO. PCR amplification with primers binding to these sequences incorporated full-length
Illumina adapters, including a multiplexing index. The ribosomal cDNA was cleaved using
ZapR in the presence of mammalian R-Probes. The remaining fragments were enriched
through a second round of PCR amplification using primers designed to match the Illumina
adapters, resulting in a smear with an average fragment size of ∼360 bp. The libraries were
normalized to 5 nM in Tris–Cl (10 mM, pH 8.5) containing 0.1% Tween 20. To validate the
quality and quantity of the isolated RNA and enriched libraries, an Agilent Fragment
Analyzer was used for the TruSeq kit, while the 4200 TapeStation System (Agilent) was
used for the SMARTer Kit.
648

RNA sequencing
Sequencing was performed using the Illumina platforms NovaSeq 6000 (for transcriptomics)
and HiSeq 2500 (for small RNA sequencing) according to standard protocols. Small RNA
sequencing was performed using RealSeq-AC miRNA (SomaGenics, Santa Cruz, CA, USA)
(short read sequencing). All samples were quantified and quality was controlled using an
Agilent Fragment Analyzer. Briefly, RNA samples (1 ng–1 ug) were adaptor-ligated,
circularized, and reverse-transcribed into cDNA. The cDNA samples were amplified using
PCR, which also incorporated sample barcodes. The library product, a peak with a fragment
size of ∼149 bp, was normalized to 10nM in Tris–Cl (2 mM, pH 8.5) containing 0.1% Tween
20. The quality and quantity of the enriched libraries were validated using an Agilent
Fragment Analyzer. Transcriptomics data were processed using the Nextflow Core RNASeq

pipeline version 3.0[58]. The data were demultiplexed with bcl2fastq, and the fastq files
underwent several quality checks, including FastQC[59]. Salmon[60] was used for pseudo
alignment and quantitation with a Salmon index built using GRCm39 with annotations from
GENCODE vM26 for mouse data and GRCh38 with annotations from GENCODE v37 for
human data. Count matrices from Salmon were used for the downstream analyses. The
count matrices were filtered, retaining genes with at least ten counts in 50% of the samples
for any condition or sex. We used the clusterProfiler R package[61] and GO biological
processes and molecular functions for GSEA, filtering terms by size between 10 and 500
genes and correcting for multiple testing (Benjamini–Hochberg correction). For the DE
analysis we used a cut-off of adjusted p-value 0.05 and threshold of log2 fold change (±1.5).
670
Global proteomics of mouse and human PFC tissue samples
Tissues from both human PFCs and four different mouse models were prepared. The
tissues were ground with a biomasher using 350 µL MeOH:H_2O (4:1). Protein pellets were
resuspended in 200 µL Laemmli buffer (10% SDS, 1 M Tris pH 6.8, and glycerol) and then
centrifuged at 16.600 x g at 4 °C for five minutes. The protein concentration was determined
using a DC assay (Bio-Rad, Hercules, CA, USA) following the manufacturer's instructions.
Each sample, containing 100 µg protein lysate, was heated at 95 °C for five minutes and
loaded onto a 5% acrylamide SDS–PAGE stacking gel (prepared in-house). Prior to
overnight digestion at 37 °C using modified porcine trypsin (Mass Spectrometry Grade,
Promega, Madison, WI, USA) at an enzyme–protein ratio of 1:80, the gel bands were
reduced and alkylated. The resulting peptides were extracted using 60% acetonitrile (ACN)
followed by 100% ACN. The peptides were resuspended in 30 µL H_2O, 2% ACN, and 0.1%
fluoroacetic acid. iRT peptides (Biognosys, Schlieren, Switzerland) were added to each
sample as an internal quality control (QC). NanoLC–MS/MS analyses were conducted using
a nanoAcquity UltraPerformance LC (UPLC) system (Waters, Milford, MA, USA) coupled to
a Q-Exactive Plus Mass Spectrometer (Thermo Fisher Scientific). The solvent system
consisted of 0.1% formic acid (FA) in water (solvent A) and 0.1% FA in ACN (solvent B).
Samples equivalent to 800 ng protein were loaded onto a Symmetry C18 precolumn (20 mm
× 180 µm with 5 µm diameter particles, Waters) over three minutes at a flow rate of 5 µL/min
with 99% solvent A and 1% solvent B. Peptides were separated on an ACQUITY UPLC
BEH130 C18 column (250 mm × 75 µm with 1.7 µm diameter particles) at a flow rate of 400
nL/min. The following gradient of solvent B was used: 1–8% over 2 min, 8–35% over 77 min,
35–90% over 1 min, 90% for 5 min, and 90–1% over 2 min. Samples from each cohort were
injected in a random order. The system was operated in data-dependent acquisition (DDA)
mode, automatically switching between MS (mass range 300–1800 *m/z* with R = 70 000,
automatic gain control [AGC] fixed at 3.10^6 ions, and the maximum injection time set to 50
ms) and MS/MS (mass range 200–2000 *m/z* with R = 17 500, AGC fixed at 1.10^5, and the
maximum injection time set to 100 ms) modes. The ten most abundant precursor ions were
selected on each MS spectrum for further isolation and higher energy collisional dissociation,
excluding monocharged and unassigned ions. The dynamic exclusion time was set to 60 s.
A sample pool was injected as an external QC every six samples for the human cohort and
every five samples for the mouse cohort. MaxQuant[53] version 1.6.14 was used to process
raw data. The Andromeda[54] search engine was used to assign peaks with trypsin/P
specificity against a protein sequence database generated in-house containing all human
(20 421 entries as of August 24, 2020) or mouse entries (17 134 entries for SOD1 and TDP-
43 models as of March 27, 2020, and 17 061 entries for C9orf72 and FUS models as of
September 29, 2020) extracted from UniProtKB/Swiss-Prot. Methionine oxidation/acetylation
of the protein N-termini was used as a variable modification, whereas cysteine
carbamidomethylation was used as a fixed modification. The "match between runs" option

was enabled to facilitate protein quantification. To control false discoveries, a maximum false
discovery rate of 1% was implemented (at both peptide and protein levels), employing a
decoy strategy. Intensities were extracted from the Proteingroup.txt file for statistical
analyses. The MaxQuant protein vs. sample table was used for downstream analyses,
including label-free quantitation intensities. Only Swiss-Prot proteins were retained, whereas
TrEMBL proteins were removed for higher reliability. After filtering out low abundant proteins,
i.e., proteins that were detected in less than 50% of the samples in any combination of
condition and sex, and imputing missing values using the missForest[62] algorithm, the
intensities were log2-transformed and used for principal component exploration, constructing
heat maps, and differential abundance analysis. The limma package[63] was used for linear
modeling, and p-values were using the Benjamini–Hochberg correction. The protein names
were mapped to the corresponding genes and used to search for enriched biological
processes and molecular functions using the criteria described in the transcriptomics data
processing, with a $p < 0.01$ threshold for functional annotation analyses.
724
Phosphoproteomics on mouse PFC brain tissue samples
Starting with the protein extracts from the global proteomics experiments, protease inhibitors
(Sigma-Aldrich, P8340) and phosphatase inhibitors (final concentration in Na_3VO_4 = 1 mM)
were added to samples. Protein concentration was determined using the RC DC Protein
Assay (Bio-Rad) according to the manufacturer's instructions. Proteins for each sample (250
µg) were reduced and alkylated prior to an in-house optimized single-pot, solid-phase-
enhanced sample preparation protocol (adapted from Hughes et al., Nat Protoc, 2019).
Briefly, beads A (Sera-Mag SpeedBeads, Fisher Scientific, Schwerte, Germany,
45152105050250) and B (Sera-Mag SpeedBeads, Fisher Scientific, 65152105050250) were
combined (1:1) and, after three washing steps with H_2O, were added to the samples (bead–
protein ratio of 10:1 for each type of bead, meaning a 20:1 ratio for the combination of
beads). After inducing protein binding to the beads with 100% ACN for 18 min, the bead and
protein mixtures were washed twice with 80% ethyl alcohol and once with 100% ACN before
being resuspended in 95 µL NH_4HCO_3 prior to overnight on-bead digestion (enzyme–protein
ratio of 1:20) at 1 000 rpm at 37 °C using modified porcine trypsin/Lys-C Mix (Mass
Spectrometry Grade, Promega). The digestion was stopped using trifluoroacetic acid (TFA)
(final pH < 2). The recovered peptides were resuspended in 170 µL 80% ACN and 0.1%
TFA, and MS PhosphoMix I Light (Sigma-Aldrich) was added to each sample (peptide [µg]–
mix [fmol] ratio of 1:6). Phosphopeptide enrichment was performed on 5 µL phase Fe(III)–
NTA cartridges on an AssayMAP Bravo platform (Agilent) following the immobilized metal
affinity chromatography protocol. Briefly, cartridges were washed and primed with 50% ACN,
0.1% TFA and then equilibrated with 80% ACN, 0.1% TFA. Subsequently, 100 µL samples
were loaded onto the phase at a rate of 2 µL/min and then washed with 80% ACN, 0.1%
TFA before being eluted in 20 µL 1% NH_4OH at 5 µL/min. After enrichment, FA and MS
PhosphoMix I Heavy (Sigma-Aldrich) (peptide [µg]–mix [fmol] ratio of 1:6) were added to
each sample. Dried phosphopeptides were resuspended in 40 µL H_2O, 2% ACN, 0.1% FA.
Sample preparation steps for the C9orf72 and FUS mouse models were identical to those
previously described for SOD1 and TDP-43, except that proteins were extracted from new
tissue samples immediately before the phosphoproteomics experiment.
NanoLC–MS/MS analyses were conducted using a nanoAcquity UPLC system (Waters)
coupled with a Q-Exactive HF-X Mass Spectrometer (Thermo Scientific) equipped with a
Nanospray Flex ion source. The solvent system consisted of 0.1% FA in water (solvent A)
and 0.1% FA in ACN (solvent B). Samples were loaded onto an ACQUITY UPLC Peptide
BEH C18 Column (250 mm x 75 µm, 1.7 µm diameter particles) over three minutes at a flow
rate of 5 µL/min with 99% solvent A and 1% solvent B. Phosphopeptides were separated on

an ACQUITY UPLC M-Class Symmetry C18 Trap Column (20 mm x 180 µm, 5 µm diameter particles; Waters) at a flow rate of 400 nL/min using a gradient of solvent B as follows: 1–2% over 2 min, 2–35% over 77 min, and finally 35–90% over 1 min. The samples in each group were injected in random order. The instrument was operated in DDA mode with automatic switching between the MS and MS/MS modes. MS scans were performed in the mass range of 375–1500 m/z with R = 120 000, an AGC fixed at 3.106 ions, and a maximum injection time of 60 ms. MS/MS scans were conducted in the mass range of 200–2000 m/z with R = 15 000, an AGC fixed at 1.105 ions, and a maximum injection time of 60 ms. In each MS spectrum, the ten most abundant ions were selected for further isolation and higher-energy collisional dissociation, excluding monocharged and unassigned ions. A dynamic exclusion time of 40 s was used to prevent the reanalysis of previously selected ions. The obtained raw phosphoproteomics data were processed using MaxQuant version 1.6.14. Peaks were assigned using the Andromeda search engine with trypsin/P specificity against an in-house-generated protein sequence database containing mouse entries from UniProtKB/Swiss-Prot (17 061 entries as of September 29, 2020). The peptides were required to have a minimum length of seven amino acids, and a maximum of one missed cleavage was allowed. These modifications include methionine oxidation, acetylation of protein N-termini, and phosphorylation of serine, threonine, and tyrosine. Cysteine carbamidomethylation was used as a fixed modification. Protein quantification utilized the "match between runs" option. A maximum false discovery rate of 1% (at both peptide and protein levels) was implemented using a decoy strategy. The intensities were extracted from the Phospho (STY).txt file and processed using the Perseus platform[57] version 2.0.7.0, during which contaminants and reversed proteins, as well as proteins with negative scores, were removed. Using the "expand sites table" option, the intensities of the different phosphopeptides involved in one phosphosite were summed, and phosphosites with a localization probability below 75% were removed. The Perseus output table was used for statistical analyses.

Only proteins that were detected in more than 50% of the mouse samples of any combination of sex and condition were retained for each dataset. The intensities were log2-transformed, quantile normalization was applied, and missing values were imputed to the lowest quartile. The remaining data processing for phosphoproteomics followed the procedure described above for proteomics data processing. For the DE analysis we used a cut-off of p-value 0.1.

Differential expression analysis

To compare murine and human samples, transcript counts were normalized using DESeq2 size factor estimation. Subtype-specific differential expression of transcripts was determined using a 1.5-fold change cut-off and an adjusted p-value <0.05 (unless stated otherwise). Two different models were designed to analyze sex-specific differences: The cohort was divided into male and female cohorts. For each cohort we carried out differential expression between case and control samples using the RNA-seq batch as covariate for the human samples.

Transcriptomics data has been processed using the NextFlow Core RNASeq pipeline, version 3.0 described[58]. The data has been demultiplexed with bcl2fastq, and the fastq files have undergone several quality checks including FastQC[59]. FastQC: A Quality Control Tool for High Throughput Sequence Data. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and fastq screen[64]. Salmon[60] was used for pseudo alignment and quantitation, with a salmon index built using GRCm39 with annotations from GENCODE vM26 for the mice data and GRCh38 with annotations from GENCODE v37 for the human data. Count matrices from Salmon were used in downstream analyses.

The Principal Component Analysis and heatmaps used the count matrices from Salmon,
after filtering and normalization using a variance stabilizing transformation blind to the
experimental design. The count matrices were filtered, keeping genes with at least ten
counts in 50% of the samples of any condition and sex.

Following differential expression analysis by DESeq2[65], we searched for relevant biological
processes and molecular functions using gene set enrichment analysis on their Gene
Ontology terms, using the clusterProfiler R package[66], filtering terms by size between 10 and
500 genes, and adjusting p-values for multiple testing with the Benjamini-Hochberg
correction. To assess the heterogeneity of the samples silhouette scores were calculated on
the first two principal components and averaged across conditions.

820
Small RNA sequencing data-processing and miRNA target prediction
Small RNA data were processed using the Nextflow Core smRNASeq pipeline version 1.0[58].
Reads were trimmed and aligned against miRBase version 22.1 using Bowtie1[67], both for
mature miRNAs and hairpins. miRNAs with at least ten counts in 50% of the samples of any
condition and sex were retained, and the rest were filtered out. Unnormalized count matrices
were used for subsequent DESeq2[65] differential expression analysis after stratifying by sex
and for p-values using the Benjamini–Hochberg correction. Mature miRNAs were mapped to
their corresponding genes using miRDB version 6.0[68], excluding matches with scores < 60
or > 800 targets, as recommended. For each miRNA present in the miRNA expression
matrices, we obtained experimentally validated targets from miRTarBase[69] 8.0 and predicted
targets from miRDB miRBase version 22[70]. miRTarBase[69] provides the most extensive
curated database of validated miRNA–target interactions (MTI) collected from literature
using natural language processing to select functional miRNA studies. Additionally, the
miRDB database includes MTIs predicted by MirTarget, which uses a support vector
machine (SVM) to analyze thousands of high-throughput sequencing experiments[63]. Each
experiment is assigned a probability score, which serves as the SVM's output. A higher
probability score indicates a higher likelihood of accurate target prediction. Therefore, we set
a threshold of 0.6 on the output probabilities to select only highly likely MTIs. Finally, we
joined the miRNA–target pairs from both sources for further analysis. For the DE analysis we
used a cut-off of p-value 0.1 and threshold of log2 fold change (±1.5).

841
Cell-type deconvolution analyses
Single-cell reference-based cell-type deconvolution of RNA-Seq in mouse models was
performed using Scaden[71]. We used healthy adult scRNA-Seq datasets for mouse[64] and
human[65]. Scaden uses a fully connected deep neural network ensemble trained on pseudo-
bulks simulated from reference scRNA-Seq data. Before deconvolution, we filtered the
scRNA-seq data using Scanpy[72] to maintain at least 200 genes expressed per cell and at
least five cells expressing one gene. For Scaden, counts per million (CPM) of simulated
pseudo-bulks and transcripts per million (TPMs) of the data to be deconvolved were used. In
this study, CPM was used for scRNA-Seq instead of TPMs because scRNA-Seq consists of
unique molecular identifier counts and does not include gene-length bias. We used a
variance cutoff of 0.01, and the mean squared error was calculated for each batch as loss.

853
Differential alternative splicing analysis
The splicing tool SUPPA2[73] version 2.3 was used to analyze DAS for seven alternative
splicing events: exon skipping, mutually exclusive exons, intron retention, alternative 3'-
splice site, alternative 5'-splice site, alternative first exon, and alternative last exon. SUPPA2
was used with multipleFieldSelection() to select the TPM values of the transcripts, followed
by GenerateEvents with the parameters -f ioe -e SE SS MX RI FL and annotation files

GENCODE v37 for human data and GENCODE vM26 for mouse data. The inclusion values
(PSI) were calculated using psiPerEvent, and the differences in PSI (ΔPSI) between the
mutant and control conditions were determined using diffSplice with parameters -m empirical
-l 0.05 -gc –save tpm_events to detect anomalies in the splicing landscape. For the DE
analysis we used a cut-off of p-value 0.1.
865

Multi-omics factor analysis
We used MOFA2[25] version 1.4.0 to integrate data from multiple omics levels for the human
cohort, including transcriptomics, miRNA, and proteomics. The MOFA model was trained on
the data, and downstream analyses were performed. Each omics type was preprocessed in
its own manner, and the default training parameters were used. The MOFA models were
initialized with 15 initial factors, and convergence was reached when the evidence lower
bound (ELBO) value did not change with more than a deltaELBO value of 1e-4%.
873

Enrichment analyses
Gene ontology, pathway enrichment analyses, and protein interaction networks
GSEA[74] was performed using gseGO and gseKEGG from the clusterProfiler R package, with
biological processes and molecular functions chosen as background databases for GO
enrichment. The p-value cutoff was set at 0.05. Differential expression was presented using
volcano plots generated with the Enhanced Volcano plot package in R, and over-
representation analysis was performed on genes that showed at least one significant DAS
event using the clusterProfiler function enrichGO, with a p-value cutoff of 0.1 and Benjamini–
Hochberg correction for multiple hypothesis testing. Protein–protein interaction networks
were created using the STRING protein interaction network database version 11.0 using
standard settings. We summarized the enriched GO terms by clustering pathways based on
their descriptions, which is called clustering in semantic space using GO-Figure[36]. In
addition, for proteomics results, we performed additional clustering analyses using
REVIGO[75]. This tool allows GO term clustering by hierarchy, considering the following such
as enrichment p-adjusted values, semantic similarity and term proximity.
889

Weighted Gene Correlation Network Analysis (WGCNA)
To conduct a weighted gene co-expression analysis, we employed the WGCNA package[15].
Pairwise Pearson correlations were calculated to establish signed regulatory networks within
the WGCNA. By constructing an adjacency matrix, we applied a soft-thresholding technique
to approximate a scale-free topological network. Eigengenes or eigenproteins were
calculated as the first principal components of each module. This resulted in the
development of several modules. We merged similar modules based on hierarchical
clustering (SOD1: 0.4, C9orf72: 0.4, FUS: 0.5, TDP-43: 0.4 for RNA-Seq and 0.25 for
proteomics). We calculated the relationships between WGCNA modules and traits. Sex was
also included in the traits resulting in four traits: male ALS, female ALS, male CTR, and
female CTR. First, we filtered modules based on the significance of the module–gene
relationship (p < 0.05) and then selected modules that were highly correlated with either
male or female ALS. The correlation cutoffs differed between the mouse models (SOD1: 0.5,
C9orf72: 0.3, FUS: 0.3, and TDP-43: 0.5 for RNA-Seq and SOD1: 0, C9orf72: 0, FUS: 0, and
TDP-43: 0 for proteomics). Using WGCNA, we also analyzed co-expression networks in
mouse models. The minimal module size was set to 30 with a merge height of 0.4–0.5 and a
correlation threshold of 0.3–0.5.
907

Pathway selection for clustering analyses with transcriptomics datasets

The selection of the pathways used for clustering analyses of human samples was based on
the enrichment analyses performed with the transcriptomics dataset. We selected the top
hits from gseGO and gseKEGG analyses [from the clusterProfiler R package], as well as a
co-expression network analysis (weighted gene co-expression analysis - (WGCNA) analysis,
analyzing the top 30 most highly enriched terms for each dataset, and then grouped these
terms by more general umbrella terms, to select ten pathways that summarized the enriched
terms best: mitochondria/respiration; synapse; MAPK cascade; oxidative stress;
nucleocytoplasmic transport; protein folding/metabolism; lipid metabolism; RNA splicing;
extracellular matrix and activation of immune response. In more detail, for overlapping
KEGG pathway and GO results, a predominant enrichment for extracellular matrix (ECM)
and immune response pathways and synapse-related terms was observed for both males
and females. The convergence of these themes across sexes suggests common biological
themes relevant for the disease and justified their selection for the clustering analyses. In
addition to neurodegenerative disease pathways driven by DE genes involved in protein
metabolism and oxidative stress mechanisms (Supplementary Fig. 5), female specific
results showed enrichment for aerobic respiration (GO) and ribosome (KEGG), while protein
metabolism was captured for male results (KEGG). Oxidative stress was also inferred from
the enrichment of oxidative phosphorylation pathways in both males and females (KEGG).
WGCNA results revealed key associations with all of the selected themes, especially for
terms related to mitochondria/respiration and protein metabolism, synapse,
nucleocytoplasmic transport, lipid metabolism; immune response and RNA splicing. WGCNA
results were also very frequently enriched for MAPK-related terms (Supplementary Table 8),
justifying its selection for the clustering analyses.
932
933
Identification of transcriptome-based subclusters in human ALS patients
We used decoupleR (v1.1.0) [76] to aggregate scores from gene sets derived from GO terms
of interest. Gene lists with the components of GO terms of interest were extracted using the
AmiGO2 database (https://amigo.geneontology.org/). For the unsupervised per sample
enrichment analyses, we used decoupleR with the consensus from the mlm, ulm, wsum
functions[76] and calculated activity scores. These are numerical values associated with
genes and represent gene expression levels.
941
Mouse primary cortical cultures
Primary cortical neuronal cell cultures were generated from neonatal mouse C57BL/6J pups
aged postnatal day 0–1 (P0–P1) in accordance with ethical guidelines for animal
experimentation at local and international levels. Animal care strictly followed official
governmental protocols, with utmost effort to minimize the number of animals utilized and to
mitigate any potential suffering or distress. Pups were decapitated, and the brains were
collected in dissection media containing 10x Hanks balanced salt solution and $NaHCO_3$. The
cortex was dissected, the meninges were removed, and small pieces of the cortex were
collected in a Falcon tube. Tissues were trypsinized at 37 °C in a water bath for 12 min and
treated with 200 µL DNAse I (10 mg/mL). The tissue strips were gently triturated (until the
tissue fragmented) in fetal bovine serum using a fire-polished Pasteur pipette. The mixture
was subsequently centrifuged at 800 × *g* for four minutes, and the resulting cell pellet was
resuspended and cultured in a neurobasal medium supplemented with B27 and antibiotics
(0.06 µg/mL penicillin and 0.1 µg/mL streptomycin). Cells were seeded at a density of 3 ×
10^5 cells per well in 24-well plates. Prior to cell seeding, the coverslips were acid-washed,
rinsed several times with water, sterilized with ethanol and UV light, and placed in a well
plate. The plates were then coated with poly-L-ornithine (0.05 mg/mL) overnight and laminin

(10 µg/mL) for 2 h in an incubator before use. The cells were maintained at 37 °C in a
humidified incubator under 5% CO_2 for seven days prior to the experiments. The cell culture
medium was replaced every three days during this period. To induce glutamate
excitotoxicity, L-glutamic acid (Tocris, UK) was dissolved in 50 mM NaOH, and a stock
solution of 50 mM was prepared prior to use. An appropriate concentration of glutamate was
prepared in the maintenance medium (neurobasal medium supplemented with B27 and
antibiotics). The cells were exposed to 5 mM glutamate by exchanging 1:3 medium at seven
days in vitro. After six hours of incubation, glutamate was thoroughly removed by washing,
and the cells were fixed for immunocytochemistry or lysed for protein extraction.
968

Immunocytochemistry and microscopy for in vitro experiments

Cells were cultured on coverslips following previously described methods[70] and
immunostained at seven days in vitro according to standard techniques. Cells were fixed in
4% paraformaldehyde in PBS at room temperature for 10 min. To quench the free aldehyde
groups, the cells were treated with 50 mM NH_4Cl for 15 min and washed with PBS. For the
permeabilization of the cell membrane, PBS with 0.25% Triton X-100 was added for 10 min
at room temperature. Non-specific binding sites were blocked by applying 10% goat serum
in PBS for at least 20 min. Dilutions of primary antibodies were prepared in blocking solution
to a final volume of 180 µL per 18-mm coverslip, and cells were incubated for 90 min at 37
°C with shaking. The following primary antibodies were used: mouse anti-MAP2 (Invitrogen,
Waltham, MA, USA; #MA5-12826, RRID:AB_10976831) (1:500) and rabbit anti-cleaved
caspase 3 (Cell Signaling Technology, Danvers, MA, USA; #9661, RRID: AB_2341188)
(1:250). The cells were washed three times for five minutes each with PBS before applying
the secondary antibodies. The cells were then incubated with secondary antibodies for 30
min and washed, respectively Alexa Fluor 488 (Invitrogen, Waltham, MA, USA #A-11034,
RRID: AB_2576217) (1:250) and goat Anti-rabbit Cy3 (Jackson ImmunoResearch, Ely,
United Kingdom, #111-165-144, RRID: AB_2338006) (1:250). For double staining, a second
primary antibody was added, and the same steps were repeated. Coverslips were mounted
on slides using a mounting medium supplemented with DAPI. Images were captured using
an inverted fluorescence microscope (Zeiss, Jena, Germany) with a 63× oil objective and
analyzed using ImageJ software. Fifteen random images from each coverslip were analyzed
for cell death by counting the number of cleaved caspase-3 positive cells. Neurite lengths
were measured using a simple neurite tracing plugin in ImageJ. Statistical analyses were
conducted using GraphPad Prism version 9.4.1 (GraphPad, San Diego, CA, USA). Outliers
were identified and removed using the Grubbs test ($α = 0.1$). Comparisons were done using
one-way analysis of variance (ANOVA), and data were plotted as the mean ± standard error
of the mean (SEM) of at least five independent experiments. Differences were considered
statistically significant at $p < 0.05$.
997

Protein extraction and western blotting

For protein analysis, cells were washed once with 1× PBS, and after adding RIPA lysis
buffer, protease inhibitor cocktail (1:25), and phosphatase inhibitor (1:20), cells were
incubated on ice for five minutes. The cells were scraped off with a cell scraper, transferred
into 1-mL reaction tubes, and homogenized by passage through a U-100 insulin syringe
several times. Protein concentration was determined using a Pierce BCA Protein Assay Kit
(Thermo Fisher Scientific) following the manufacturer's instructions. One microliter of protein
sample was used for the assay. Colorimetric reactions were analyzed using an Infinite M200
PRO ELISA plate reader (Tecan, Männedorf, Switzerland). Twenty grams of each sample
were loaded onto a gel (NuPAGE 4 to 12%, Bis-Tris) (Invitrogen, Waltham, MA, USA).
NuPAGE LDS sample buffer (1:4) and sample-reducing buffer (1:10) were added to lysed

protein before loading onto the gels, incubated with shaking at 75 °C for 13 min, and
centrifuged at 12 000 × *g* at 4 °C. The proteins were subjected to gel electrophoresis at 200
V. Subsequently, the proteins were transferred onto nitrocellulose membranes using an
iBlot2 gel transfer device (Thermo Fisher Scientific, Waltham, MA, USA). The membranes
were then blocked with 5% nonfat milk in PBST at room temperature for 30 min. For
antibody incubation, primary antibodies (diluted in blocking buffer) were added to the
membranes and allowed to incubate overnight at 4 °C with rotation. Rabbit anti Lamin B1
(Proteintech, Planegg-Martinsried, Germany, #12987-1-AP, RRID: AB_2136290) (1:2000),
rabbit anti p44/42 mitogen-activated protein kinase (anti-Erk1/2) (Cell Signaling Technology,
Danvers, MA, USA, #9102, RRID: AB_330744) (1:1000), rabbit anti MEK2 (Cell Signaling
Technology, Danvers, MA, USA, #9125, RRID: AB_2140644) (1:1000), rabbit anti Phospho-
p44/42 MAPK(ERK1/2) (Cell Signaling Technology, Danvers, MA, USA, #9101, RRID:
AB_331646) (1:1000) and rabbit anti Phospho-MEK1/2 (Ser217/221) (Cell Signaling
Technology, Danvers, MA, USA, #9121, RRID: AB_331648) (1:1000). After four washes with
PBST (five minutes each), the membranes were incubated with peroxidase-conjugated goat
anti-rabbit secondary antibodies (Vector Laboratories, Newark, CA, USA, #PI-1000, RRID:
AB_2336198) (diluted 1:10 000 in blocking buffer) at room temperature for one hour. Finally,
the membranes were washed thoroughly with PBST to remove the unbound antibodies.
Blots were incubated with enhanced chemiluminescence reagent and were imaged using a
Molecular Imager ChemiDoc (Bio-Rad, Hercules, CA, USA) imaging system. Band signal
intensities were quantified using ImageJ software and normalized to housekeeping proteins
and controls. Statistical analyses were conducted using GraphPad Prism version 9.4.1.
Outliers were identified and removed using the Grubbs test (α = 0.1). Comparisons were
done using one-way ANOVA, and data were plotted as the mean ± SEM of at least three
independent experiments. Differences were considered statistically significant at p < 0.05.
1034

Tissue protein extraction for immunoblot analysis
Spinal cords were homogenized in five volumes (w/v) of 1% boiling SDS[71]. Protein
homogenates were sonicated, boiled for 10 min, and centrifuged at 13 500 × *g* for five
minutes. Supernatants were analyzed by dot blotting. For detergent-insoluble protein
extraction, mouse tissues were homogenized in 10 volumes (w/v) of buffer (15 mM Tris–HCl
pH 7.6, 1 mM dithiothreitol, 0.25 M sucrose, 1 mM $MgCl_2$, 2.5 mM EDTA, 1 mM EGTA, 0.25
M Na_3VO_4, 2 mM $Na_4P_2O_7$, 25 mM NaF, 5 µM MG132) and a protease inhibitor cocktail
(Roche, Basel, Switzerland), as previously described[72]. Briefly, the samples were centrifuged
at 10 000 × *g*, and the pellet was suspended in an ice-cold homogenization buffer containing
2% Triton X-100 and 150 mM KCl. The samples were then centrifuged at 10 000 × *g* to
obtain the Triton-insoluble (insoluble) fraction.
1046

Immunohistochemistry
Mice were anesthetized and underwent transcardial perfusion using 50 mL PBS followed by
100 mL of 4% paraformaldehyde (in PBS). The spinal cord was promptly dissected,
postfixed for three hours, and subsequently transferred to a solution of 20% sucrose in PBS
overnight. Once sunk, the spinal cord was transferred to a 30% sucrose solution until it
reached the desired consistency. Next, the spinal cord was frozen in n-pentane at 45 °C and
stored at ∼-80 °C. Prior to freezing, the spinal cord was divided into cervical, thoracic, and
lumbar segments and embedded in Tissue-Tek O.C.T. compound (Sakura Finetek,
Torrance, CA, USA). Coronal sections (30 µm thick) were sliced from the lumbar spinal cord
and subjected to immunohistochemistry. Rabbit monoclonal anti-pMEK (Ser221) (pMEK2)
antibody (1:50, Cell Signaling, Danvers, MA, USA, RRID: AB_490903) was used. In brief,
slices were incubated with blocking solutions (0.2% Triton X-100 and 2% normal goat

serum) at room temperature for one hour. Subsequently, the slices were incubated overnight at 4 °C with primary antibodies. The sections were then incubated with biotinylated secondary antibodies (1:200) for one hour at room temperature, followed by immunostaining using an avidin–biotin kit and diaminobenzidine. To facilitate visualization, sections were counterstained with 0.5% cresyl violet. Stained sections were captured at 20× and 40× magnification using an Olympus BX-61 Virtual Stage microscope (Olympus Life Sciences, Tokyo, Japan), ensuring complete stitching of the entire section with a pixel size of 0.346 μm. Images were acquired using 6-μm-thick stacks (step size of 2 μm). The different focal planes were merged into a single stack using mean intensity projection to maintain consistent focus across the sample. Finally, the acquired signals were analyzed for each slice using ImageJ and OlyVIA software (Olympus Life Sciences, Tokyo, Japan).

Immunoblotting for in vivo experiments

Protein levels were quantified using a BCA Protein Assay Kit (Pierce Biotechnology, Waltham, MA, USA) and analyzed by western blotting and dot blotting following previously described methods[73]. Membranes were blocked with a solution of 3% (w/v) BSA (Sigma-Aldrich, St. Louis, MO, USA) and 0.1% (v/v) Tween 20 in Tris-buffered saline (pH 7.5). Subsequently, the membranes were incubated with primary antibodies followed by peroxidase-conjugated secondary antibodies (GE HealthCare, Chicago, IL, USA). The antibodies utilized were rabbit monoclonal anti-pMEK2 (1:2000, Cell Signaling Technology; Danvers, MA, USA, RRID: AB_490903), rabbit monoclonal anti-phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (1:2000, Cell Signaling Technology, Danvers, MA, USA,; RRID: AB_2315112), rabbit polyclonal anti-human SOD1 (1:1000, StressMarq Biosciences, Victoria, Canada; RRID: AB_2704217), rabbit polyclonal anti-ubiquitin (1:1000, Abcam, Cambridge, UK; RRID: AB_306069), and mouse monoclonal anti-SQSTM1/p62 (p62) (1:500, Abcam, Cambridge, United Kingdom). Peroxidase-conjugated secondary antibodies (goat anti-mouse and anti-rabbit) (GE HealthCare, Chicago, IL, USA) were used at dilutions of 1:20 000 and 1:10 000, respectively. The blots were developed using the Luminata Forte Western Chemiluminescent HRP Substrate (MilliporeSigma, Burlington, MA, USA) and visualized on a ChemiDoc Imaging System (Bio-Rad, Hercules, CA, USA). Densitometry analysis was performed using Image Lab software version 6.0 (Bio-Rad, Hercules, CA, USA). Protein immunoreactivity was normalized to Ponceau Red staining (Honeywell Fluka, Charlotte, NC, USA) for accurate quantification.

Preclinical study in SOD1^{G93A} mice

SOD1^{G93A} female and male mice received 3 mg/kg dose of trametinib or vehicle (PBS) through intranasal delivery, twice per week. The treatment of SOD1^{G93A} female and male mice started from 9 weeks of age until 16 weeks of age, when they were sacrificed for biochemical analysis. The effect of trametinib on the progression of the disease in SOD1^{G93A} female mice was assessed twice a week from 9 weeks of age by measuring body weight, hind limb extension reflex and grip strength. The extension reflex was quantified using the following 3-point score system: 3, hind limbs extending to form an angle of 120 degrees; 2.5, hind limbs extending to < 90 degrees with decreased reflex in a hind limb; 2.0, as 2.5 with decreased reflex in both hind limbs; 1.5, loss of reflex with marked flexion in a hind limb; 1, as 1.5 with marked flexion in both hind limbs; 0.5, loss of reflex with hind limbs and paws held close to the body, but still able to walk; 0, as 0.5 but unable to walk[77]. The grip strength test was performed by placing mice on a horizontal metallic grid which was then gently inverted. The latency to fall of each mouse was recorded. The test ended after 90 s. In the case of failure, the measurement was repeated three times and the best performance of the session was considered for the statistical analysis[78]. Age of paralysis was defined as the age

at which the mice lost reflex with hind limbs and paws held close to the body (0.5 score at extension reflex test). Disease onset was retrospectively determined as the average age at which the mouse exhibits the first failure, at two consecutive time points, from both maximum weight and maximum performance in extension reflex score and grip strength test. The animals were euthanized when they showed rigid paralysis in the hind limb and a 0.5 score in hind limb extension reflex. Data of behavioral tests and body weight loss were evaluated by two-way ANOVA for repeated measures. Disease onset and survival length were evaluated by log-rank Mantel-Cox test. The animal experimental protocols used in this study (Prot. no. 9F5F5.143 and 9F5F5.250) were prospectively reviewed and approved by the Mario Negri Institutional Animal Care and Use Committee and the Italian Ministry of Health (*Direzione Generale della Sanità Animale e dei Farmaci Veterinari, Ufficio 6).* This study complied with international and local animal welfare standards.

Plasma neurofilament light chain quantification
Mouse plasma samples were collected in K2-EDTA BD Microtainer blood collection tubes and centrifuged at 5 000 × *g* for five minutes to obtain the plasma. The plasma NfL was quantified using a Simoa NF-light Advantage (SR-X) Kit (#103400) on a Quanterix SR-X platform (Quanterix, Boston, MA, USA). All reagents used were from a single lot, and measurements were performed according to the manufacturer's protocol.

Statistical analyses of in vivo experiments
Statistical analyses of in vivo experimental data were performed using GraphPad Prism 7.0 and statsmodels Python package (v0.13.0). Student's *t*-tests or one-way ANOVAs, followed by pairwise TukeyHSD post hoc tests, were used to analyze the differences between the experimental groups for each variable. Statistical significance was set at $p < 0.05$.

Data availability
Raw RNA-Seq and processed gene expression data were deposited to the National Center for Biotechnology Information Gene Expression Omnibus database (GSE234246). Encrypted raw RNA-Seq data for the human cohort were deposited to the European Genome Phenome Archive (registered study: EGAS00001007318). Proteomics and phosphoproteomics datasets have been deposited in the ProteomeXchange Consortium database with the identifiers PXD043300 and PXD043297, respectively.

Code availability
Given the size and complexity of the datasets, data processing, and data analysis methods, we integrated all analyses and raw data into a Data Version Control pipeline[79].
The code for the computational analyses is available at: https://github.com/imsb-uke/MAXOMOD_Pipeline.

References

1.Mehta, P. *et al.* Prevalence of amyotrophic lateral sclerosis in the United States using established and novel methodologies, 2017. *Amyotroph. Lateral Scler. Front. Degener.* 24, 108–116 (2023).

2. Suzuki, N., Nishiyama, A., Warita, H. & Aoki, M. Genetics of amyotrophic lateral sclerosis: seeking therapeutic targets in the era of gene therapy. *J. Hum. Genet.* 68, 131–152 (2023).

3. Mead, R. J., Shan, N., Reiser, H. J., Marshall, F. & Shaw, P. J. Amyotrophic lateral sclerosis: a neurodegenerative disorder poised for successful therapeutic translation. *Nat. Rev. Drug Discov.* 22, 185–212 (2023).

4. Braak, H. & Braak, E. Neuropathological stageing of Alzheimer-related changes. *Acta*
Neuropathol. (Berl.) 82, 239–259 (1991).
5. Braak, H. *et al.* Staging of brain pathology related to sporadic Parkinson's disease.
Neurobiol. Aging 24, 197–211 (2003).
6. Brettschneider, J. *et al.* Stages of pTDP-43 pathology in amyotrophic lateral sclerosis.
Ann. Neurol. 74, 20–38 (2013).
7. Aronica, E. *et al.* Molecular classification of amyotrophic lateral sclerosis by unsupervised
clustering of gene expression in motor cortex. *Neurobiol. Dis.* 74, 359–376 (2015).
8. Morello, G. *et al.* Integrative multi-omic analysis identifies new drivers and pathways in
molecularly distinct subtypes of ALS. *Sci. Rep.* 9, 9968 (2019).
9. Tam, O. H. *et al.* Postmortem Cortex Samples Identify Distinct Molecular Subtypes of
ALS: Retrotransposon Activation, Oxidative Stress, and Activated Glia. *Cell Rep.* 29, 1164-
1177.e5 (2019).
10. Prudencio, M. *et al.* Distinct brain transcriptome profiles in C9orf72-associated and
sporadic ALS. *Nat. Neurosci.*18, 1175–1182 (2015).
11. Figueroa-Romero, C. *et al.* Expression of microRNAs in human post-mortem
amyotrophic lateral sclerosis spinal cords provides insight into disease mechanisms. *Mol.*
Cell. Neurosci. 71, 34–45 (2016).
12. Umoh, M. E. *et al.* A proteomic network approach across the ALS-FTD disease spectrum
resolves clinical phenotypes and genetic vulnerability in human brain. *EMBO Mol. Med.* 10,
48–62 (2018).
13. Eshima, J. *et al.* Molecular subtypes of ALS are associated with differences in patient
prognosis. *Nat. Commun.*14, 95 (2023).
14. Kenna, K. P. *et al.* NEK1 variants confer susceptibility to amyotrophic lateral sclerosis.
Nat. Genet. 48, 1037–1042 (2016).
15. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics* 9, 559 (2008).
16. Rinchetti, P., Rizzuti, M., Faravelli, I. & Corti, S. MicroRNA Metabolism and
Dysregulation in Amyotrophic Lateral Sclerosis. *Mol. Neurobiol.* 55, 2617–2630 (2018).
17. Magen, I. *et al.* Circulating miR-181 is a prognostic biomarker for amyotrophic lateral
sclerosis. *Nat. Neurosci.* 24, 1534–1541 (2021).
18. Song, N., Song, R. & Ma, P. MiR-340-5p alleviates neuroinflammation and neuronal
injury via suppressing STING in subarachnoid hemorrhage. *Brain Behav.* 12, e2687 (2022).
19. Chia, R., Chiò, A. & Traynor, B. J. Novel genes associated with amyotrophic lateral
sclerosis: diagnostic and clinical implications. *Lancet Neurol.* 17, 94–102 (2018).
20. Ciccarelli, F. D. *et al.* The identification of a conserved domain in both spartin and
spastin, mutated in hereditary spastic paraplegia. *Genomics* 81, 437–441 (2003).
21. Han, W. *et al.* Alpha-synuclein (SNCA) polymorphisms and susceptibility to Parkinson's
disease: a meta-analysis. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet. Off. Publ. Int.*
Soc. Psychiatr. Genet. 168B, 123–134 (2015).
22. Xi, Y., Ju, R. & Wang, Y. Roles of Annexin A protein family in autophagy regulation and
therapy. *Biomed. Pharmacother. Biomedecine Pharmacother.* 130, 110591 (2020).
23. Chua, J. P., De Calbiac, H., Kabashi, E. & Barmada, S. J. Autophagy and ALS:
mechanistic insights and therapeutic implications. *Autophagy* 18, 254–282 (2022).
24. Månberg, A. *et al.* Altered perivascular fibroblast activity precedes ALS disease onset.
Nat. Med. 27, 640–646 (2021).
25. Argelaguet, R. *et al.* Multi-Omics Factor Analysis-a framework for unsupervised
integration of multi-omics data sets. *Mol. Syst. Biol.* 14, e8124 (2018).
26. Kenigsbuch, M. *et al.* A shared disease-associated oligodendrocyte signature among
multiple CNS pathologies. *Nat. Neurosci.* 25, 876–886 (2022).

27. Costa, J. *et al.* Cerebrospinal Fluid Chitinases as Biomarkers for Amyotrophic Lateral
Sclerosis. *Diagn. Basel Switz.* 11, 1210 (2021).
28. Buratti, E. & Baralle, F. E. The multiple roles of TDP-43 in pre-mRNA processing and
gene expression regulation. *RNA Biol.* 7, 420–429 (2010).
29. Polymenidou, M. *et al.* Misregulated RNA processing in amyotrophic lateral sclerosis.
Brain Res. 1462, 3–15 (2012).
30. Melamed, Z. *et al.* Premature polyadenylation-mediated loss of stathmin-2 is a hallmark
of TDP-43-dependent neurodegeneration. *Nat. Neurosci.* 22, 180–190 (2019).
31. Ma, X. R. *et al.* TDP-43 represses cryptic exon inclusion in the FTD–ALS gene UNC13A.
Nature 603, 124–130 (2022).
32. Ling, J. P., Pletnikova, O., Troncoso, J. C. & Wong, P. C. TDP-43 repression of
nonconserved cryptic exons is compromised in ALS-FTD. *Science* 349, 650–655 (2015).
33. Hansen, K. D., Brenner, S. E. & Dudoit, S. Biases in Illumina transcriptome sequencing
caused by random hexamer priming. *Nucleic Acids Res.* 38, e131 (2010).
34. Hirano, M. *et al.* Mutations in the gene encoding p62 in Japanese patients with
amyotrophic lateral sclerosis. *Neurology* 80, 458–463 (2013).
35. Beckers, J., Tharkeshwar, A. K. & Van Damme, P. C9orf72 ALS-FTD: recent evidence
for dysregulation of the autophagy-lysosome pathway at multiple levels. *Autophagy* 17,
3306–3322 (2021).
36. Reijnders, M. J. M. F. & Waterhouse, R. M. Summary Visualizations of Gene Ontology
Terms With GO-Figure! *Front. Bioinforma.* 1, 638255 (2021).
37. Franke, T. F., Hornik, C. P., Segev, L., Shostak, G. A. & Sugimoto, C. PI3K/Akt and
apoptosis: size matters. *Oncogene* 22, 8983–8998 (2003).
38. Yoshida, T. *et al.* Identification and characterization of a novel chemotype MEK inhibitor
able to alter the phosphorylation state of MEK1/2. *Oncotarget* 3, 1533–1545 (2012).
39. Dong, X., Wang, Y. & Qin, Z. Molecular mechanisms of excitotoxicity and their relevance
to pathogenesis of neurodegenerative diseases. *Acta Pharmacol. Sin.* 30, 379–387 (2009).
40. Pratilas, C. A. *et al.* (V600E)BRAF is associated with disabled feedback inhibition of
RAF-MEK signaling and elevated transcriptional output of the pathway. *Proc. Natl. Acad.*
Sci. U. S. A. 106, 4519–4524 (2009).
41. Lake, D., Corrêa, S. A. L. & Müller, J. Negative feedback regulation of the ERK1/2 MAPK
pathway. *Cell. Mol. Life Sci. CMLS* 73, 4397–4413 (2016).
42. Chun, Y. S. *et al.* MEK1/2 inhibition rescues neurodegeneration by TFEB-mediated
activation of autophagic lysosomal function in a model of Alzheimer's Disease. *Mol.*
Psychiatry 27, 4770–4780 (2022).
43. Gal, J., Ström, A.-L., Kilty, R., Zhang, F. & Zhu, H. p62 accumulates and enhances
aggregate formation in model systems of familial amyotrophic lateral sclerosis. *J. Biol.*
Chem. 282, 11068–11077 (2007).
44. Benatar, M. Lost in translation: treatment trials in the SOD1 mouse and in human ALS.
Neurobiol. Dis. 26, 1–13 (2007).
45. Gurney, M. E. *et al.* Motor neuron degeneration in mice that express a human Cu,Zn
superoxide dismutase mutation. *Science* 264, 1772–1775 (1994).
46. Santiago, J. A., Quinn, J. P. & Potashkin, J. A. Network Analysis Identifies Sex-Specific
Gene Expression Changes in Blood of Amyotrophic Lateral Sclerosis Patients. *Int. J. Mol.*
Sci. 22, 7150 (2021).
47. Murdock, B. J., Goutman, S. A., Boss, J., Kim, S. & Feldman, E. L. Amyotrophic Lateral
Sclerosis Survival Associates With Neutrophils in a Sex-specific Manner. *Neurol.*
Neuroimmunol. Neuroinflammation 8, e953 (2021).
48. Günther, R. *et al.* The rho kinase inhibitor Y-27632 improves motor performance in male
SOD1(G93A) mice. *Front. Neurosci.* 8, 304 (2014).

49.   Torres, P. *et al.* Gender-Specific Beneficial Effects of Docosahexaenoic Acid Dietary
Supplementation in G93A-SOD1 Amyotrophic Lateral Sclerosis Mice. *Neurother. J. Am. Soc.*
Exp. Neurother. 17, 269–281 (2020).
50.   Tahedl, M. *et al.* Propagation patterns in motor neuron diseases: Individual and
phenotype-associated disease-burden trajectories across the UMN-LMN spectrum of MNDs.
Neurobiol. Aging 109, 78–87 (2022).
51. Sahana, T. G. & Zhang, K. Mitogen-Activated Protein Kinase Pathway in Amyotrophic
Lateral Sclerosis. *Biomedicines* 9, 969 (2021).
52. Ayala, V. *et al.* Cell stress induces TDP-43 pathological changes associated with ERK1/2
dysfunction: implications in ALS. *Acta Neuropathol. (Berl.)* 122, 259–270 (2011).
53. Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass
spectrometry-based proteomics evidences. *Nucleic Acids Res.* 50, D543–D552 (2022).
54. Ilieva, H., Polymenidou, M. & Cleveland, D. W. Non-cell autonomous toxicity in
neurodegenerative disorders: ALS and beyond. *J. Cell Biol.* 187, 761–772 (2009).
55. Bonifacino, T. *et al.* Nearly 30 Years of Animal Models to Study Amyotrophic Lateral
Sclerosis: A Historical Overview and Future Perspectives. *Int. J. Mol. Sci.* 22, 12236 (2021).
56. Gordon, D. *et al.* Single-copy expression of an amyotrophic lateral sclerosis-linked TDP-
43 mutation (M337V) in BAC transgenic mice leads to altered stress granule dynamics and
progressive motor dysfunction. *Neurobiol. Dis.* 121, 148–162 (2019).
57. Mitchell, J. C. *et al.* Overexpression of human wild-type FUS causes progressive motor
neuron degeneration in an age- and dose-dependent fashion. *Acta Neuropathol. (Berl.)* 125,
273–288 (2013).
58. Ewels, P. A. *et al.* The nf-core framework for community-curated bioinformatics pipelines.
Nat. Biotechnol. 38, 276–278 (2020).
59. Andrews, S. FastQC: A Quality Control Tool for High Throughput Sequence Data.
(2010).
60. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast
and bias-aware quantification of transcript expression. *Nat. Methods* 14, 417–419 (2017).
61. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
Innov. Camb. Mass 2, 100141 (2021).
62. Stekhoven, D. J. & Bühlmann, P. MissForest--non-parametric missing value imputation
for mixed-type data. *Bioinforma. Oxf. Engl.* 28, 112–118 (2012).
63. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Res.* 43, e47 (2015).
64. Wingett, S. W. & Andrews, S. FastQ Screen: A tool for multi-genome mapping and
quality control. *F1000Research* 7, 1338 (2018).
65. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* 15, 550 (2014).
66. Wu, H., Kirita, Y., Donnelly, E. L. & Humphreys, B. D. Advantages of Single-Nucleus
over Single-Cell RNA Sequencing of Adult Kidney: Rare Cell Types and Novel Cell States
Revealed in Fibrosis. *J. Am. Soc. Nephrol. JASN* 30, 23–32 (2019).
67. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol.* 10, R25 (2009).
68. Chen, S., Lake, B. B. & Zhang, K. High-throughput sequencing of the transcriptome and
chromatin accessibility in the same cell. *Nat. Biotechnol.* 37, 1452–1457 (2019).
69. Huang, H.-Y. *et al.* miRTarBase update 2022: an informative resource for experimentally
validated miRNA-target interactions. *Nucleic Acids Res.* 50, D222–D230 (2022).
70. Chen, Y. & Wang, X. miRDB: an online database for prediction of functional microRNA
targets. *Nucleic Acids Res.* 48, D127–D131 (2020).
71. Menden, K. *et al.* Deep learning-based cell composition analysis from tissue expression

profiles. *Sci. Adv.* 6, eaba2619 (2020).

72. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression

data analysis. *Genome Biol.* 19, 15 (2018).

73. Trincado, J. L. *et al.* SUPPA2: fast, accurate, and uncertainty-aware differential splicing

analysis across multiple conditions. *Genome Biol.* 19, 40 (2018).

74. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for

interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* 102, 15545–

15550 (2005).

75. Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO summarizes and visualizes long

lists of gene ontology terms. *PloS One* 6, e21800 (2011).

76. Badia-i-Mompel, Pau, *et al.* decoupleR: ensemble of computational methods to infer

biological activities from omics data. *Bioinformatics Advances,* vbac016 (2022).

77. Pasetto, L. *et al.* Defective cyclophilin A induces TDP-43 proteinopathy: implications for

amyotrophic lateral sclerosis and frontotemporal dementia. *Brain J. Neurol.* 144, 3710–3726

(2021).

78. Vallarola, A. *et al.* RNS60 exerts therapeutic effects in the SOD1 ALS mouse model

through protective glia and peripheral nerve rescue. *J. Neuroinflammation* 15, 65 (2018).

79. Kuprieiev, R. *et al.* DVC: Data Version Control - Git for Data & Models. Zenodo

https://doi.org/10.5281/zenodo.7093084 (2022).

1328

Acknowledgements

We thank all the members of the Lingor, Bonn, Carapito, Schlapbach, Pasterkamp, and Bonetto laboratories for their feedback on the manuscript. This work was performed by the research consortium "Multi-omic analysis of axono-synaptic degeneration in motoneuron disease (MAXOMOD)" funded in the scope of the E-Rare Joint Transnational Call for Proposals 2018 "Transnational research projects on hypothesis-driven use of multi-omic integrated approaches for discovery of disease causes and/or functional validation in the context of rare diseases." Consortium members: S. B., J. P., V. B., C. C., R. S., and M. K., coordinated by P. L.

L. C. G., M. P., and P. L. were supported by the Bundesministerium für Bildung und Forschung (01GM1917A). P. L.; Q.Z. and D. E. were further supported by the Munich Cluster for Systems Neurology (SyNergy). S. H. was supported by SFB1192 B8; S. O. by SFB1286 SP02 and KFO296 P8; R. K. by FOR5068 P9; F. H. by the M3I excellence initiative and a UKE postdoctoral stipend; C. H. by KFO306 P11; and S. B. by SFB1286 SP02, SFB1192 PB8, and PC3. J. P. was supported by Stichting ALS Nederland (TOTALS). D. E. received funding from the European Union's Horizon Europe research and innovation program under grant agreement No 101057649 and the Association for Frontotemporal Degeneration (AFTD).

The funders had no role in the study design, data collection and analysis, decision to publish, or manuscript preparation.

Author contributions

P. L. and L. C. G. conceived the project, and S. H., F. H., and S. O. conceptualized the computational analysis. P. L., L. C. G., J. K., and A. D. designed the sample collection methodology, reviewed the sample and data quality, and coordinated the acquisition of human tissue samples from the brain banks as well as pathological, genetic, and clinical information. L. C. G, M. P., and Q. Z. processed mouse and human brain samples for multiomics experiments with the infrastructure provided by V. B., C. C., D. E., and P. L.

C. D., M. S., T. B. H., I. C., and M. D. analyzed the genetic data.

S. H., S. O., R. K., and F. H. were responsible for the computational data analysis, and conceived and planned the statistical analyses. L. C. G, S. H., F. H., R. K., P. L., L.T., M. P., and L. P. contributed to figure generation and assembly.

M. P, L. P., L. C. G., S. S., M. G. and S. C. conducted the validation experiments with conceptual inputs from P. L., C. C. and V. B.

L. C. G., S. H., S. B., and P. L. wrote the manuscript with input from all co-authors.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information - Supplementary Tables and Supplementary Figures are provided as appendices.

Correspondence and requests for materials should be addressed to Paul Lingor.

 Figure legends

Fig. 1 Identification of transcriptomic subclusters and sex-specific differences in
human ALS patients
a. Overview of the sample processing workflow. Prefrontal cortex samples were prepared for
multiomics experiments from the human cohort (51 ALS/50 CTR samples), as well as from
four selected ALS mouse models (C9orf72, FUS, SOD1, and TDP-43; 10 TG/10 CTR
animals per group). RNA, DNA, and protein lysates were prepared for mRNA and small RNA
sequencing, targeted DNA sequencing, and proteomics/phosphoproteomic (ph-proteomic)
experiments (for animal tissues). **b.** Principal component analysis (PCA) on the 500 most
variable genes of the human samples. Blue and orange indicate the sex, and the condition is
indicated in pink (ALS) or gray (CTR). **c.** Volcano plot of deregulated proteins in human
samples. x-axis: log2 fold change; y-axis: -log10 *p*-value for each protein. Blue and orange
circles indicate significant differential changes: left side, decrease (low ALS); right side,
increase (high ALS). **d.** Differential alternative splicing (DAS) analysis. The plot displays the
results for human male and female samples for various splice events, i.e., alternative exon
(AE), skipped exon (SE), alternative 5'-splice site (A5), alternative 3'-splice site (A3),
alternative last exon (AL), and retained intron (RI) events. Each event is represented by a
separate bar, the height of which represents the fraction of significant events in ALS vs.
CTR. Blue bars represent events with significant differential splicing in males, whereas
orange bars represent events with significant differential splicing in females. **e.** Sex-specific
enrichment analysis reveals crucial pathways in neurodegeneration and ALS pathology. In
males, significant enrichment is observed in "synapse-related" pathways, pathways related
to immune system activation, ECM, and various neurodegenerative diseases. Females show
a distinct pattern with significant results in "ribosomal function" and the "oxidative
phosphorylation" pathway. Pathways associated with neurodegenerative disorders, immune
system activation and cell adhesion molecules are also evident. **f.** Hierarchical clustering
analysis of enriched pathways and literature insights reveals four distinct clusters (C1–C4) in
ALS patients. Immune response regulation dichotomizes patients into C1/C2 vs. C3/C4,
while ECM, synaptic function, and protein folding further differentiate C1 vs. C2 and C3 vs.
C4. **g.** Heatmap showing modules from weighted gene co-expression network analysis
(WGCNA) associated with the clusters through similarly regulated RNA networks. **h**.
Pathway enrichment per WGCNA cluster (top hits). The turquoise module, upregulated in C1
and C2, especially in males, is enriched for mitochondrial respiration, suggesting increased
oxidative activity in PFC neurons. The yellow module, associated with synaptic function,
exhibits a similar regulation in C1 and C2, while the tan and lightcyan modules, enriched for
immune response and RNA splicing, respectively, are upregulated in C3 and C4.

1411

1412

Fig. 2 Validation of sex-specific dysregulation in ALS patients in microRNAomic,
proteomic and multiomic analyses
a-b. Volcano plot of microRNA analyses of human samples, separated by sex for mature (**a**)
and hairpin (**b**) microRNAs. The x-axis shows the log2 fold change in ALS vs CTR, whereas
the y-axis shows the -log10 p-value. Orange and blue dots represent DEGs in females and
males, respectively. miRNA-mediated regulation in ALS reveals a notable sex-dependent
pattern. Male patients with ALS exhibit a more pronounced downregulation of mature and
hairpin miRNAs compared to their female counterparts. **c.** Heatmap showing miRNA
expression changes for the identified ALS clusters. Considering log2 fold change ≥ 0.1, and
miRNA candidates that are significant in at least two conditions, clusters 1, 2, and 4 emerge

with distinct miRNA profiles. Cluster 3, in contrast, displays no differentially expressed (DE) miRNAs under these parameters, suggesting a relatively stable miRNA landscape within this particular cluster. The scale (left-hand side) shows the range of log2 fold change values. **d.** Volcano plot of differentially expressed proteins (DEP) of human samples. The x-axis shows the log2 fold change ALS and CTR, whereas the y-axis shows the -log10 p-values . Orange and blue dots represent DEGs in females and males, respectively. 379 DEPs in males and 251 in females. ANXA2 emerges as the sole protein downregulated in both sexes (p < 0.1). **e-f.** Functional enrichment and unsupervised clustering using REVIGO, a tool used for summarizing and visualizing Gene Ontology (GO) terms. Important pathway clusters are revealed in both sexes, including synaptic function, immune response, and ECM/cytoskeleton. Females (**e**) exhibited enrichment in transmembrane transport, lipid metabolism, development, catalytic activity, and ERK1/2 signaling, while males (**f**) showed enrichment in cell metabolism and tyrosine kinase-related pathways. The size of the circles represents the number of genes in the GO-BP terms, and the color of each node represents the enrichment FDR values. **g.** Uniform manifold approximation and projection (UMAP) of MOFA factor analysis demonstrates that sex emerges as a robust differentiating factor when integrating transcriptomic, small RNA, and proteomic data. **h.** MOFA correlation analysis displays which factors are dominated by which omic layer. Components of the representative factors 3 (males) and 12 (females) are displayed in feature weight plots (bottom part). These factors are the ones that better correlate with disease condition for each sex.

Fig. 3 ALS mouse models partially resemble human ALS subclusters
a. Volcano plots of DE genes for male and female mice across ALS models (C9orf72, SOD1, TDP-43, and FUS). The x-axis illustrates the log2 fold change between ALS and control (CTR), while the y-axis displays the -log10 adjusted p-values. Orange and blue dots denote differentially expressed proteins (DEPs) in females and males, respectively. **b.** Analysis of enriched Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways using ClusterProfiler in male and female samples. The dot plot displays significantly enriched pathways in males and females, represented by circles colored by their corresponding adjusted p-values (-log10 transformed) on the x-axis and gene count on the y-axis. The size of each circle corresponds to the number of genes annotated in the KEGG gene set. Distinct fingerprints of pathway deregulation were observed in each ALS model. The C9orf72 model prominently featured alterations in immune/inflammatory response pathways, the SOD1 model showed disruptions in the ERK1/2 cascade, development, and response to oxidative stress, while the TDP-43 model exhibited changes in transcription and endopeptidase activity. The FUS model exhibited no significant pathway enrichment. **c.** Heatmap showing the differences (ALS vs. ctrl) in the estimated cell-type fractions for human subclusters and mouse ALS models. The SOD1 model and human C1/C2 groups showed a decrease in glial and endothelial cells and a relative increase in excitatory neurons. In contrast, the C9orf72 model and human C3 cluster exhibited a strong increase in glial and endothelial cells and a decrease in excitatory neurons, suggesting neuroinflammation and neuronal loss. The FUS and TDP-43 models displayed intermediate levels of glial and neuronal cells, indicating that observed transcriptional changes are partially driven by cell composition. **d.** Correlation analysis of deconvolved cell-type fraction changes (as shown in d) between human and mouse ALS models. Subgroups C1 and C2 demonstrated the strongest correlation with the SOD1 model, while C3 exhibited the best correlation with the C9orf72 model, with additional correlations with the TDP-43 and FUS models. A weak correlation was noted between C4 and the FUS model. **e.** GO-Figure comparative analysis of enrichment results for proteomics results showed clustering for differentiation and

development in human females (groups 3, 4, 7, 11) and in human males (groups 8), as well as synapse (females: groups 16 and 17; males: groups 2, 10, 15) and immune/defense response (females: group 12; males: groups 3, 4, 6). C9orf72 exhibited clustering for RNA processing, ribosome, translation, ATP synthesis, development, cell adhesion, transport, and synapse. The SOD1 model showed strong clustering for ATP synthesis, mitochondrial respiration, translation, and vesicle-mediated transport. Enrichment results underscored pathways previously identified in RNA sequencing data.

Fig. 4 Consistent dysregulation of the MAPK pathway in human ALS patients and mouse models

a. The occurrence and importance of MAPK pathways and other kinase pathways are shown across all mouse models highlighting distinct activities within the co-expression modules identified by WGCNA. The upper panel shows the significance (-log10 p-value) and the lower panel shows absolute counts. **b.** Enrichment results for DAS genes. Enriched terms include translation, myelin sheath (C9orf72), GTPase activity, myelin sheath (SOD1), translation, heat shock protein binding (TDP-43), and purine metabolism, basal plasma membrane (FUS). Regulation of kinase activity is particularly enriched in DAS of male patients with ALS. **c.** Differential expression analyses for mature miRNAs for humans and ALS mouse models. Mice exhibited pronounced differential expression (DE) of miRNAs, with the C9orf72 model showing the most significant changes. Among the top regulated miRNAs, miR-451a is consistently deregulated across all mouse models and human patients with ALS. While C9ORF72, FUS and TDP43 male mice demonstrate a trend for an increase of miRNA-451a levels, the opposite trend is observed in the SOD1 male mouse model (log2 fold change = -2.34, padj = 0.1685) and a significant decrease is visible in human males with ALS (log2 fold change = -0.91, padj = 0.0002). miR-451a also shows significant decreases in the female C9ORF72 mouse model (log2 fold change = -1.82, padj = 0.0363) and trends towards downregulation are observed across all other female mouse models (SOD1: log2 fold change = -2.34; TDP43: log2 fold change=-0.55) and female humans PFC (log2 fold change = -0.08). **d.** Targets of miR-451a, such as MAPK1, AKT1, BCL2, Il6R, IKBKB, MIF, MMP2, and MMP9, are involved in cell growth, survival, apoptosis, inflammatory signaling, and ECM remodeling. **e.** Target centrality measurement for miR-451a reveals MAPK1 as the top hit for target centrality measures based on the network shown in panel (d). **f.** Protein-protein interaction (PPI) network of the genes in MOFA factor 12 in females. MAPK1 is an important molecular hub and interacts with PEA15, PRRT2, MEK2, DUSP6, HSPA4 and HSP90AA1. Further proteins of interest are highlighted (bold/dark purple). **g.** Scheme for the mechanism of action and the expected effects of trametinib in the context of the MAPK pathway. This pathway is dependent on a series of phosphorylation steps that ultimately lead to MEK and Erk activation through the Ras/RAF/PI3K cascade, exerting pivotal roles for cell proliferation and survival. The physiological activation of ERK is subject to negative feedback regulation both downstream and upstream of MEK, and this regulation involves the expression of DUSPs, as well as the direct phosphorylation and inhibition of proteins such as RAF. Trametinib binds to the activation loop of MEK, disrupting the Raf-dependent phosphorylation of MEK. This results in reduced expression of P-MEK and P-Erk. The pharmacological inhibition of MEK by trametinib swiftly eliminates this feedback mechanism, inducing the phosphorylation of MEK (panel created with BioRender.com).

Fig. 5 Effects of the MEK2-inhibitor trametinib on the MAPK pathway in vitro

a-b. Effects of different concentrations of trametinib on apoptosis in glutamate (5 mM)- and non-glutamate-treated cells analyzed by immunostaining. Representative photomicrographs for two of the analyzed conditions (control [vehicle] and 200 nM trametinib). Scale bar: 40 μm. **c-d.** Quantification plots showing the effects of treatment with trametinib on cell survival (caspase 3 staining) (**c**) and neurite outgrowth (**d**) for glutamate- or vehicle-treated (control) cells, for all analyzed conditions (control [vehicle]; 2 nM trametinib; 20 nM trametinib; 200 nM trametinib). Data are presented as the mean□±□ standard error of the mean (SEM) of at least five different cultures and were tested using a one-way analysis of variance. **e-f.** Western blot analysis and quantification of trametinib effects on pErk1/2 with and without glutamate treatment. Data represent the mean□±□SEM of at least three different cultures and were tested by one-way analysis of variance, $p < 0.05$.

Fig. 6 Modulation of the MAPK pathway member MEK2 by trametinib attenuates ALS pathology in vivo.

a. Western blot analysis of pMEK2 in the lumbar spinal cord of SOD1 and non-transgenic mice at 9, 14, and 19 weeks of age. Data are mean□±□SEM (n = 6–4 in each experimental group) and are expressed as relative immunoreactivity (RI). *$p < 0.05$, pairwise Tukey's test after one-way analysis of variance. **b-f.** Results of the study in female and male SOD1 mice treated or not with trametinib from 9 to 16 weeks of age. Dot blot analysis of pERK1/2 (**b**), p62 (**c**), insoluble SOD1 (**d**), and ubiquitin (**e**) in the spinal cord of SOD1 female and male mice treated with trametinib or vehicle. Data are mean□±□SEM (n = 3/5 in each experimental group) and are expressed as relative immunoreactivity (RI). *$p < 0.05$, Student's *t*-test. Plasma neurofilament light chain levels were analyzed in female and male (**f**) SOD1 mice treated or not with trametinib. Data are mean ± SEM (n□= 3/5 in each experimental group). *$p < 0.05$, Student's *t*-test. **g-h.** Diffuse pMEK2 immunostaining in the lumbar spinal cord of non-transgenic (**g**) and SOD1 mice (**h**) at 19 weeks of age. In the ventral horns, pMEK2 staining was mainly observed in motor neurons. Scale bar: 50 μm. Tg: transgenic mice; Ntg: non-transgenic mice. **i-k.** Results of the preclinical study in female SOD1 mice treated or not with trametinib from 9 weeks of age up to survival. **i.** Paralysis is assessed as the age at which the mice lost reflex with hind limbs and paws held close to the body. The mean age was significantly different in the trametinib-treated (150.2 ± 8 days) with respect to the vehicle-treated (143.5 ± 6.5 days) mice. Data are mean ± SEM (n□= 8/10 in each experimental group). *$p < 0.05$, Student's *t*-test. **j.** Kaplan–Meier curve for disease onset and survival of SOD1 female mice treated with vehicle (n = 8) or trametinib (n = 10). Log–rank Mantel–Cox test for disease onset, $p = 0.0177$ (vehicle 103.4 ± 3.5 days and trametinib 108.7 ± 5 days). Log–rank Mantel–Cox test for survival, $p = 0.0405$ (vehicle 147 ± 6.5 days and trametinib 153.7 ± 8 days). **k.** Plasma NfL were analyzed in SOD1 female mice treated or not with trametinib, at 16 and 19 weeks of age. Data are mean ± SEM (n□= 3/6 in each experimental group). *$p < 0.05$, Benjamini-Hochberg corrected p-values after one-way analysis of variance.

Table 1: Summary of the demographics of the human cohort
Continuous variables are presented as medians and ranges. Categorical variables were
compared using Fisher's exact test (for single groups) or the chi-square test (for multiple
groups); continuous variables were compared using the Mann–Whitney test. The p-values
shown are uncorrected for multiple testing; p-values less than 0.05 are indicated in bold.
NBB: The Netherlands Brain Bank; OBB: Oxford Brain Bank; ICL MS&PD TB: Imperial
College London - Multiple Sclerosis and Parkinson's Tissue Bank; London NDBB: London
Neurodegenerative Diseases Brain Bank.

	Control	ALS	p-value
Donors (number)	50	51	
Age at death (years)	75 (43–94)	67 (44–83)	**0.0032**
Sex female	56.0%	31.4%	**0.0163**
Postmortem interval (hours)	22 (5–95)	23 (2.5–98)	0.4324
Unknown	8.0%	5.9%	
Onset			
Bulbar	-	17.6%	
Spinal	-	37.3%	
Upper limb	-	31.6%	
Lower limb	-	63.1%	
Limb unknown	-	5.3%	
Thoracic	-	2.0%	
Unknown	-	43.1%	
Disease duration (years)	-	3 (1–28)	
Unknown	-	39.2%	
Brain bank origin			**< 0.0001**
NBB	18.0%	17.6%	
Oxford BB	20.0%	27.5%	
ICL MS & PD TB	38.0%	0.0%	
London NDBB	24.0%	54.9%	

1573 ALS, amyotrophic lateral sclerosis.

a 0mM glutamate

control
MAP2   caspase

200nM trametinib
MAP2   caspase

merge

b 5mM glutamate

control
MAP2   caspase

200nM trametinib
MAP2   caspase

merge

c

0mM glut   5mM glut

caspase 3 pos. cells (relative to CTR)

P < 0.0001

P = 0.0237
P = 0.0027

trametinib concentration (nM)
0 2 20 200   0 2 20 200

d

0mM glut   5mM glut

Neurite lenght/cell (relative to CTR)

p = 0.0051

p = 0.0003
p < 0.0001

trametinib concentration (nM)
0 2 20 200   0 2 20 200

e

0 mM Glut   5 mM Glut

Lamin b

Erk1/2

p-Erk1/2

0 2 20 200   0 2 20 200

trametinib concentration (nM)

f

0 mM glut   5 mM glut

p-Erk1/2 / Erk1/2 band intensity ratio

p = 0.0139

p = 0.0042
p = 0.0103

trametinib concentration (nM)
0 2 20 200   0 2 20 200

a pMEK2 (RI)

females · males

b pERK1/2 (RI)

females · males

c p62 (RI)

females · males

d insoluble SOD1 (RI)

females · males

e insoluble ubiquitin (RI)

females · males

f NfL (pg/ml)

females · males

● trametinib-treated females   ● trametinib-treated males   ● vehicle-treated

g Ntg

pMEK2
Nissl

h SOD1-G93A Tg

pMEK2
Nissl

i age of paralysis (days)

vehicle   trametinib

j disease onset   survival

onset / survival (%)

age (days)

k NfL (pg/ml)

veh. tram.   veh. tram.

16 weeks   19 weeks

● trametinib-treated   ● vehicle-treated