Multiomic ALS signatures highlight sex differences and molecular subclusters and identify the MAPK pathway as therapeutic target

Lucas Caldi Gomes[1*], Sonja Hänzelmann[2,3,4*], Sergio Oller[2,3], Mojan Parvaz[1], Fabian Hausmann[2,3], Robin Khatri[2,3], Melanie Ebbing[2,3], Constantin Holzapfel[2,3], Laura Pasetto[5], Stefano Fabrizio Columbro[5], Serena Scozzari[5], Marie Gebelin[6], Johanna Knöferle[1], Isabell Cordts[1], Antonia F. Demleitner[1], Laura Tzeplaeff[1], Marcus Deschauer[1], Claudia Dufke[7], Marc Sturm[7], Qihui Zhou[8], Pavol Zelina[9], Emma Sudria-Lopez[9], Tobias B. Haack[7,10], Sebastian Streb[11], Magdalena Kuzma-Kozakiewicz[12], Dieter Edbauer[8,13], R. Jeroen Pasterkamp[9], Endre Laczko[11], Hubert Rehrauer[11], Ralph Schlapbach[11], Christine Carapito[6], Valentina Bonetto[5], Stefan Bonn[2,3§], Paul Lingor[1,8,13§,#]

Technical University of Munich, School of Medicine, rechts der Isar Hospital, Clinical Department of Neurology, Munich, Germany.
Institute of Medical Systems Biology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany.
Center for Biomedical AI, University Medical Center Hamburg-Eppendorf, Hamburg, Germany.
Department of Medicine, University Medical Center Hamburg-Eppendorf, Hamburg, Germany.
Research Center for ALS, Mario Negri Institute for Pharmacological Research, Milan, Italy
Laboratoire de Spectrométrie de Masse Bio-Organique, Université de Strasbourg, Infrastructure Nationale de Protéomique, Strasbourg, France.
Institute of Medical Genetics and Applied Genomics, University of Tübingen, Tübingen, Germany
German Center for Neurodegenerative Diseases (DZNE), München, Germany.
Department of Translational Neuroscience, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands.
Center for Rare Diseases, University of Tübingen, Tübingen, Germany.
Functional Genomics Center Zürich, ETH Zürich and University of Zürich, Zürich, Switzerland.
Department of Neurology, Medical University of Warsaw, Warsaw, Poland.
Munich Cluster for Systems Neurology (SyNergy), Munich, Germany.

*,§ These authors contributed equally.

#Corresponding Author: Paul Lingor; email: paul.lingor@tum.de; Clinical Department of Neurology, School of Medicine, rechts der Isar Hospital, Technical University of Munich, Ismaninger Straße 22, 81675 Munich, Germany, Tel.: +498941408257

42
Abstract
44

Amyotrophic lateral sclerosis (ALS) is the most common motor neuron disease and lacks effective disease-modifying treatments. Here, we performed a multiomic analysis of the prefrontal cortex of 51 patients with sporadic ALS and 50 control subjects, as well as four transgenic mouse models of C9orf72-, SOD1-, TDP-43-, and FUS-ALS to characterize early and sex-specific disease mechanisms in ALS. Integrated analyses of transcriptomes, (phospho)proteomes, and miRNAomes revealed more pronounced changes in male patients. We identified transcriptome-based human ALS subclusters driven by the immune response, extracellular matrix, mitochondrial respiration, and RNA metabolism. The molecular signatures of human subclusters were reflected in specific mouse models. Individual and integrative multiomics analyses highlighted the mitogen-activated protein kinase pathway as an early disease-relevant mechanism. Its modulation by trametinib in vitro and in vivo validated that mitogen-activated protein kinase kinase 2 is a promising therapeutic target with beneficial effects in female patients.

Main, Introduction

Amyotrophic lateral sclerosis (ALS) is the most common motor neuron disease, largely prevalent in males, and causes paralysis and death within a few years of symptom onset[1]. Most cases are sporadic (sALS), i.e., without a family history, but 10% of cases have a genetic cause (fALS). The most common genetic variants include *C9orf72*, *SOD1*, *TARDBP*, and *FUS* and approximately 10% of patients with sALS carry disease-causing mutations[2]. The etiology of sALS remains unclear, and there are no effective disease-modifying treatments for the disease[3]. An improved understanding of early disease mechanisms could help identify the diagnostic and prognostic biomarkers, thereby uncovering more efficient therapeutic drug targets. Although direct analysis of affected nervous system tissues remains the gold standard in neuropathology, patient samples are only available postmortem and in limited quantities. This increases the risk of describing the end stages of the disease and obscuring mechanisms that occur in earlier phases, which are potentially more favorable drug targets. As in Alzheimer's or Parkinson's disease, ALS is believed to spread over time—from the motor cortex to other cortical brain areas[4–6]. Numerous studies have analyzed the motor cortex in ALS[7–9]. However, as this area is the most severely affected by ALS, it primarily reflects the final stages of the disease[6]. In contrast, the prefrontal cortex (PFC) in Brodmann area 6 typically exhibits only intermediate TDP-43 pathology at the time of death[6], suggesting that analyzing this area could provide insights into earlier disease-mediated alterations.

Previous investigations of ALS brain tissue have primarily focused on individual molecular subsets, including transcripts[7,9,10], miRNAs[11], and proteins[12], suggesting that ALS is a complex and heterogeneous disease. Recent studies have identified potentially distinct ALS populations, which were stratified into different subclusters based on transcriptomics and gene set enrichment analyses (GSEA)[9,13].

In this study, we deciphered early ALS disease mechanisms by profiling the transcriptomes, miRNAomes, and (phospho)proteomes in the PFCs of patients with ALS and four ALS mouse models. We identified strong sex differences and demonstrated that ALS is not a homogeneous disease but consists of different molecular subtypes that are correlated in individual transgenic mouse models of the disease. Multiomic data integration identified several known ALS patho-mechanisms and pathways that have been less prioritized in ALS research to date—particularly the Mitogen-activated protein kinase (MAPK) pathway, which is important for modulating cell survival and differentiation. In vitro and in vivo validation studies underscored MAPK as a putative target for early therapeutic interventions.

Results

Cohort composition and depth of analysis

A deep multiomic characterization of the human postmortem PFC (Brodmann area 6) from 51 patients with neuropathologically confirmed ALS and 50 control (CTR) patients was performed to describe early molecular changes in sALS (Fig. 1a, Table 1, Supplementary Table 1). On average, 19.641 transcripts, 736 miRNAs (mature miRNAs and hairpin precursors), and 2.344 proteins were detected per sample (Supplementary Fig. 1). C9orf72 repeat expansion was detected in one patient with ALS, and another individual carried a pathogenic variant of *NEK1* (c.3107C>G, p.Ser1036Ter)[14] (Supplementary Fig. 2, Supplementary Table 2). For clarity and accuracy, we refer to all cases simply as "ALS". Four transgenic mouse models were analyzed to identify parallels with human ALS. PFCs from presymptomatic or early disease stages were collected from C9orf72-, SOD1-, TDP-43-, or FUS-ALS, with an equal distribution of wild-type and transgenic littermates and sex (n = 20 per model, Fig. 2, Supplementary Fig. 3). In mice, an average of 17.020 transcripts, 842 miRNAs (mature miRNAs and hairpins), 2.568 proteins, and 6.755 phosphosites were detected. Overall, sample quality was high (Supplementary Fig. 4).

Transcriptomic stratification of human ALS into four molecular subclusters

Principal component analysis was performed to assess the effects of disease, sex, and sample origin on the transcriptome, revealing a moderate separation by condition (silhouette score: 0.11 [ALS]; -0.03 [CTR]) but a strong separation by sex (Supplementary Fig. 5). Therefore, we analyzed differentially expressed genes (DEGs) separately for males and females. The number of DEGs was significantly higher in males (72) than in females (2) (Fig. 1b, Supplementary Table 3). Overrepresentation analysis revealed the upregulation of genes involved in *retinoic acid* and *lipid metabolism* in females and downregulation of *extracellular matrix (ECM), collagen,* and *vasculature* in males (Supplementary Fig. 6). Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis showed suppressed *complement/cytokine* and *ECM-receptor signaling* and activation of *oxidative phosphorylation* and *glutamatergic synapse* in males, whereas decreased *complement/coagulation* and activated *ribosomal function* and *oxidative phosphorylation* were found in females (Fig. 1d). Hierarchical clustering of the enriched pathways revealed four distinct clusters: C1–C4. Regulation of *immune response* dichotomized patients with ALS best (C1 and C2 vs. C3 and C4), whereas second-level arborization was mainly driven by *ECM* (C1 vs. C2), *synaptic function*, *RNA splicing,* and *protein folding* (C3 vs. C4, Fig. 1c, Supplementary Table 4). These clusters are reminiscent of previously proposed subtypes[9], where C1 and C2 show similarities to ALS-Oxidative stress (ALS-Ox) (and less to ALS-Retrotransposon reactivated), and C3 and C4 correspond to the ALS-Glial dysfunction (ALS-Glia) subtype (Supplementary Fig. 7). To characterize the clusters using similarly regulated RNA networks, we performed weighted gene co-expression network analysis (WGCNA[15]), resulting in 20 modules (M) (Fig. 1e, Supplementary Fig. 8, Supplementary Table 5). The Mturquoise module was enriched for *mitochondrial respiration* and upregulated in C1 and C2, particularly in males, and driven by neuronal alterations, suggesting increased oxidative respiration in PFC neurons. The Myellow module was enriched for *synaptic function* and showed a similar regulation in C1 an C2. By contrast, the Mtan and Mlightcyan modules were enriched for *immune response,* and *RNA splicing* was upregulated in C3 and C4 (Fig. 1e,f). Molecular subclusters and sex-specific differences drive heterogeneity in the PFCs of patients with ALS.

Finally, transcriptome-based analyses were performed to explore differential alternative splicing (DAS) in the analyzed cohort. TARDBP/TDP-43 and FUS regulate alternative splicing and transcript usage in hundreds of genes[16,17]. TDP-43 also represses cryptic exon splicing

events in *UNC13A*[18]. However, these events are specific to neurons with TDP-43 pathology
and are rarely detectable in bulk RNA-Seq at our sequencing depth. The sequences are also
not conserved in mice. We identified significant DAS events that occurred more frequently in
male patients with ALS than in females (Fig. 3a). DAS was observed in males for *CLTB*, *TPRN*,
NRN1, and *CAMK2N1* and in females for *TPRN* and the gene encoding TMEM170A-CFDP1,
a novel readthrough protein (AC009163.5) (Supplementary Table 6). Genes involved in the
regulation of *kinase activity* were particularly enriched in the DAS of male patients with ALS
(Supplementary Table 7). Early disease mechanisms appear to be influenced by splicing
events, particularly in males. TPRN, a stereocilium-associated protein previously described
only in non-syndromic deafness, may play additional roles in the pathogenesis of ALS (Fig.
3d).
155
Transcriptomes of murine ALS models reflect human ALS subclusters
To explore sex differences in mice, we performed separate differential analyses for male and
female mice (Fig. 2, Supplementary Fig. 9). The strongest transcriptomic changes were
observed in the C9orf72 mice, reflecting the rapid disease progression observed in this model.
Each mouse model was predominantly characterized by the deregulation of specific pathways,
that is, *immune/inflammatory response* in C9orf72 mice, *ERK1/2 cascade, development* and
response to H_2O_2 in SOD1, and *transcription* and *endopeptidase activity* in TDP-43. No
enrichment was found in the FUS model (Fig 2a–e, Supplementary Table 8). DAS analyses
were also conducted in mice, which showed more balanced results between the sexes (Fig.
3b, Supplementary Fig. 10). We identified *FINB, CPLANE1* (involved in ciliogenesis and
migration), and *ATP1B1* (a membrane-bound Na$^+$/K$^+$-ATPase) as differentially spliced in male
and female models (Fig. 3c), showing enrichment in the terms *translation* and *myelin sheath*
(C9orf72), *GTPase activity* and *myelin sheath* (SOD1), *translation* and *heat shock protein*
binding (TDP-43), and *purine metabolism* and *basal plasma membrane* (FUS)
(Supplementary Table 7).
Next, to assess to which degree the observed alterations in gene expression reflected
changes in the cellular composition of the PFC, we estimated the cell-type fractions for the
subclusters of human and mouse ALS models using deep learning-based deconvolution[19].
Interestingly, the human subclusters showed changes in the cell fractions that were reflected
in the individual mouse models. The SOD1 model and human C1 and C2 groups showed a
decrease in glial and endothelial cells and a relative increase in excitatory neurons. The
C9orf72 model and human C3 were characterized by a strong increase in glial and endothelial
cells and a decrease in excitatory neurons, respectively, suggesting strong neuroinflammation
and neuronal loss. The FUS and TDP-43 models showed intermediate levels of glial and
neuronal cells (Fig. 2c, Supplementary Fig. 11). Transcriptional changes are partially driven
by cell composition. Interestingly, our data suggest that the neurovascular unit in ALS[20] may
be affected differently in different subgroups of patients with ALS. Overall, our transcriptomic
analyses revealed remarkable similarities between human clusters and mouse models: C1
and C2 showed the best correlation with the SOD1 model, whereas C3 correlated best with
the C9orf72 model and, to a lesser extent, TDP-43 and FUS models. Finally, C4 showed a
weak correlation with the FUS model (Fig. 2d). While we observed transcriptional and cell
composition differences between mouse models and human ALS clusters, we found
consistent changes in *MAPK signaling* in the TDP-43, SOD1, and C9orf72 models, as well as
human ALS samples (Supplementary Fig.12). Moreover, analysis of the subclusters showed
that the classical MAPK pathway was activated in C1 and C2, whereas it was downregulated
in C3. Meanwhile, C1 and C2 showed downregulation of the JNK and p38 MAPK pathways,
which were activated in C3 (Fig. 2e). Thus, mouse models of ALS reflect the molecular

subgroups of human ALS, which are partially driven by cell fraction changes. The deregulation
of MAPK pathways is a common theme in both human ALS and mouse transcriptomes.
195
Downregulation of hairpin and mature miRNA in males characterize early ALS
To address the role of miRNA-mediated regulation in ALS mechanisms[21], we sequenced small
RNAs. Male patients with ALS showed a stronger downregulation of mature miRNAs and
miRNA hairpins than female patients (Fig. 3e,f). The higher number of differentially expressed
(DE) hairpins compared with the number of mature miRNAs indicates defects in early miRNA
biogenesis as a potential early disease mechanism. In males, we observed 17 mature DE
miRNAs (15 downregulated, 2 upregulated) and in females, there were 9 DE miRNAs (4
downregulated, 5 upregulated) ($p < 0.1$). For miRNA hairpins, females presented 24 DE
hairpins (12 downregulated, 12 upregulated), whereas males presented 99 DE hairpins (86
downregulated, 13 upregulated). The top hits common to males and females included the
miRNA hairpins let-7a-3, miR-26a-1, let-7f-1 (all downregulated), and miR-1227 (upregulated)
(Fig. 3e–h). Enrichment analysis of the targets of the DE miRNAs revealed pathways involved
in *cell growth*, *focal adhesion, integrin binding,* and *kinase activity regulation,* indicating that
the ECM/neurovascular unit is an important target of miRNA regulation (Supplementary Fig.
13). Mouse models showed the most pronounced DE of miRNAs in the C9orf72 model (targets
annotated for *cell survival/PI3K-Akt, mTOR, MAPK signaling*) followed by the SOD1 model
(*cancer* and *lipid metabolism*) (Supplementary Fig. 14, Supplementary Table 9). Among the
top five regulated miRNAs, miR-451a was found to be significantly deregulated across all
mouse models and human patients with ALS (Fig. 3g,h). Targets of hsa-miR-451a, such as
MAPK1, *AKT1,* and *BCL2*, are known for their roles in the regulation of cell growth, survival,
and apoptosis, whereas other targets, such as *Il6R*, *IKBKB*, and *MIF,* are involved in
inflammatory signaling or ECM remodeling, such as *MMP2* and *MMP9* (Fig. 3i, Supplementary
Fig. 15). In addition to the pronounced sex-specific regulation of miRNAs, consistent miRNA
changes related to the regulation of the MAPK pathway, ECM, and inflammatory signaling
were identified and matched the findings of the transcriptomic dataset.
221
Human proteomic landscape most closely correlates with the TDP-43 model
Subsequently, we assessed the proteomic signatures of early-stage ALS using mass
spectrometry. In human samples, we detected 49 (30) differentially expressed proteins (DEPs)
in males (females), with annexin A2 (ANXA2) being the only protein downregulated in both
sexes ($p < 0.1$). Interestingly, we identified several neurodegeneration-related proteins, such
as MATR3, SPART, and SCNA (involved in the genetic forms of ALS, spastic paraplegia, and
Parkinson's disease, respectively) (Fig. 4a, Supplementary Table 10). The projection of
transcriptomic clusters onto proteomic data did not reproduce subclustering, likely because of
the much smaller number of mapped entities (Supplementary Fig. 16). Functional enrichment
and unsupervised clustering identified relevant pathways in both sexes, such as *synaptic*
function, *immune response*, and *ECM/cytoskeleton*. In contrast, *transmembrane transport,*
lipid metabolism, *development*, and *catalytic activity* were enriched in female mice, whereas
cell metabolism was enriched in male mice (Fig. 4b,c, Supplementary Table 11). In mice,
proteomic analyses revealed the strongest changes in the C9orf72 model, where
sequestosome 1/p62, a product of the ALS-causing *SQSTM1*[22], showed the strongest
upregulation, indicating reduced autophagic flux in a dipeptide accumulation model[23]
(Supplementary Figs. 17, 18). pSQSTM1 was also significantly upregulated in the
phosphoproteomic analysis (Supplementary Fig. 19). Male SOD1 mice showed one
upregulated DEP—exportin-1 (XPO1)—a major regulator of nuclear RNA export. This
corresponded to increased *XPO1* expression in this model, which was also observed in TDP-
43 females, FUS males, and both sexes in the C9orf72 model (Supplementary Table 3). To

compare Gene Ontology (GO) term enrichment in human and animal proteomics, we used
clustering in semantic space[24]. Human proteomics results showed clustering for *differentiation*
and *development* (females: clusters 3, 4, 7, and 11; males: cluster 8), *synapse,* and
immune/defense response (Fig. 4e,f). In contrast, the C9orf72 model showed clustering for
RNA processing, ribosome, translation, *ATP synthesis*, *development*, *cell adhesion*, *transport*,
and *synapse*. Remarkably, the SOD1 model showed strong clustering for *ATP synthesis,*
mitochondrial respiration, *translation*, and *vesicle-mediated transport*. Thus, enrichment
results underscored the pathways previously identified in the RNA sequencing data. Overall,
the human proteome showed the strongest similarity with the TDP-43 model (Fig. 4d). Several
mouse models (especially male models) showed increased expression of XPO1—a target in
an ALS clinical trial[25] (Supplementary Fig. 20).
254
Integration of multiomic data reveals sex-specific molecular networks of ALS
After using various omics modalities to identify the molecular pathways associated with early
ALS, we employed a biologically motivated approach that focused on identifying valid
interaction triplets involving miRNAs, transcripts, and proteins. Triplet miR-769-3p–*ANXA2*–
ANXA2 was particularly intriguing because ANXA2 was the only common differentially
expressed protein in both sexes (Supplementary Fig. 21, Supplementary Table 12). Valid
quadruplets in mouse models, including phosphoproteomic data, highlighted GFAP, SQSTM1,
ATXN2L, and XPO1 as salient target proteins (Supplementary Fig. 22, Supplementary Table
12).
Next, we integrated transcriptomic, small RNA, and proteomic data using Multi-Omics Factor
Analysis (MOFA)[26]. Because sex was an important differentiating factor (Fig. 4g), MOFA was
performed by sex (Fig. 4h, Supplementary Fig. 23). In males, factor 1, mainly driven by hairpin
small RNA, explained 23.7% of the variance. Downregulation of hsa-miR-7851, -1285-1, -
5096, and a cluster of -1273 isoforms strongly contributed to its weight (Supplementary Fig.
24). MAPK1 plays a central role in the miR-1273 targets (Supplementary Fig. 25).
Transcriptome-based factor 3 correlated best with disease condition and was driven by genes
responsible for *vesicular function* (*RAB3C, NSF*), *oxidative phosphorylation* (*ND1, ND2*), *cell*
survival (*BCL2, BHLHB9*), and *RNA metabolism* (*SNORA73B, RN7SL2*). This factor also
contained miR-3648-2, which recruits amyloid precursor protein intracellular domains and
regulates survival and neurogenesis[27]. Proteome-dominated factor 4 contains ZO2 and CD44,
which are involved in myelination and blood–central nervous system barrier (BCNSB)
formation, respectively. Finally, factor 7, which also showed a strong correlation with the
disease condition, was dominated by neurofilament heavy, medium, and light polypeptides
(NfH, NfM, and NfL, respectively) as well as proteins involved in Ca^{2+}-binding (HPCL4) and
ECM formation (PGCA) (Supplementary Fig. 24). In females, factors 1–3 explained 42.6% of
the variance, but factors 10 and 12 correlated best with the disease condition (Fig. 4h).
Synaptic genes, such as *RAB3C*, *NAPB*, and *SNAP25*, contributed to factor 1 and are
upregulated in ALS. The hsa-miR-1285-1, miR-5096, and miR-1273 clusters also contributed
to factor 2, similar to factor 1, in males. Factors 3 and 7 (similar to male factor 4) included the
oligodendrocyte and myelin markers CD44 and ZO2 (Fig. 4i). Factor 10 contained the
antiproteases SERPINA1 and SERPINA3[28] as well as chitinases CHI3L1 and CHI3L2, which
are known biomarkers for ALS[29] (Supplementary Fig. 26). Factor 12 downregulated the NFH
and MAPK–ERK1/2 regulator PEA15 (Fig. 4j). MOFA analysis underlined the mechanisms
identified in the individual omics analyses, such as the downregulation of miRNA clusters
(particularly in males, Supplementary Fig. 27), ECM components, and oligodendrocyte and
myelin markers. This unbiased data integration strategy highlighted known ALS biomarkers
(neurofilaments and chitinases) and the MAPK pathway, especially in females, as important
molecular hubs (Fig. 4k).

MAPK pathway emerges as therapeutic target based on multiomic data integration

Single-omics analyses identified several molecular signaling pathways as pharmacological targets that could be more appropriate for different subclusters or sexes of patients with ALS. Currently, licensed drugs target *glutamatergic synapse function* (riluzole), *oxidative stress* (edaravone), *mitochondrial function* (tauroursodeoxycholic acid/phenylbutyrate), or SOD1 itself (tofersen). The major mechanisms revealed in our analysis, such as *immune response* and *ECM/BCNSB function*, have not yet been targeted by licensed drugs. Although multiple molecular pathways identified in our analysis merit therapeutic validation, we decided to focus on the MAPK pathway, which was altered consistently in humans, albeit more pronounced in females and mouse models, across several data types, integration methods, and different subclusters. More specifically, we concentrated on mitogen-activated protein kinase kinase 2 (MAP2K2 or MEK2), as it appeared to be upregulated in the human PFC and multiple mouse models and is central to the female-specific ALS-associated factor 12 (Fig. 4k, Supplementary Fig. 12). Furthermore, MEK2 can be modulated by the clinically approved inhibitor, trametinib[30].

Validation of MEK2 inhibition by trametinib in vitro and in vivo

First, we used primary cortical neuronal cultures from P0–P1 C57BL/6J mice that were treated with glutamate as an in vitro model of excitotoxicity in ALS[31]. Treatment with 5 mM glutamate (six hours) did not affect MEK2 protein expression, but significantly increased pErk1/2 levels. Trametinib attenuates glutamate-induced phosphorylation of Erk1/2. Glutamate treatment also increased cell death (caspase-positive neurons) and reduced the average neurite length, both of which were counteracted by trametinib (Fig. 5a–f, Supplementary Fig. 28). Our data suggest that trametinib attenuates MEK2 activity and reduces Erk1/2 phosphorylation resulting in decreased cell death and increased neurite outgrowth under excitotoxic stress.

To validate the importance of the MAPK pathway in vivo, we selected the SOD1 mouse model, which showed the strongest similarity with the largest human ALS subcluster (C1 and C2) (Fig. 2h). We observed strong MEK2 phosphorylation in the motor neurons of the spinal cord, the main tissue involved in the pathology of this animal model[32] (Fig. 5g,h). In females, pMEK2 levels substantially increased with disease progression, whereas in males, the levels returned to control levels after week 14 (Fig. 5i). We treated SOD1 mice with trametinib for 7 weeks, starting from week 9 (presymptomatic stage), and observed a reduction in ERK1/2 phosphorylation compared with that in vehicle-treated female and male mice (Fig. 5j). Trametinib significantly reduced the autophagy receptor p62 expression in the spinal cord of female but not male mice, correlating with its previously described neuroprotective role of increasing autophagy via transcription factor EB activation[33] (Fig. 5k). p62 also co-localizes with ubiquitin and mutant SOD1 in protein aggregates[34]. Accordingly, we detected a significant reduction in detergent-insoluble SOD1 and ubiquitin in trametinib-treated females but not in males (Fig. 5l,m). Finally, we investigated the effect of trametinib on neurodegeneration. NfL plasma concentration[35] was significantly reduced after trametinib-treatment in female SOD1 mice (Fig. 5n). The striking sex-specific differences in the trametinib treatment response correlated with increased MEK2 phosphorylation during disease progression in female but not male SOD1 mice. Thus, trametinib markedly affects the clearance of protein aggregates, leading to neuroprotection in female SOD1 mice, suggesting that MEK2 is a promising therapeutic target for ALS, particularly in females.

Discussion

In this study, we conducted individual and combined analyses of multiple omics datasets to comprehensively understand the molecular architecture of ALS in the PFC, an area affected in the later stages by TDP-43 pathology and that has the potential to reveal early disease mechanisms[6]. We observed that significant sex-specific differences are often more pronounced in males, which is aligned with the higher prevalence of ALS in males[1]. Previous studies have also identified sex-specific differences in the blood of patients with ALS[36,37] as well as therapeutic responses in ALS mouse models[38,39]. However, current therapeutic options, clinical trials, and Food and Drug Administration guidelines for clinical trial design[40] do not consider patient sex in differential therapy. Our data, especially the sex-specific differences in MAPK signaling and therapeutic tractability, argue for a stronger consideration of sex as a covariate in ALS clinical trials.

Phenotypic heterogeneity is clearly recognized in ALS[41], as reflected by stratification in onset or disease progression rate, and previous transcriptomic analyses have suggested molecular subtypes[13]. We identified four clusters in our human ALS cohort based on the transcriptome, partially mirroring previously identified ALS-Ox, ALS-Glia, and ALS-Transposable elements subtypes[9]. While the ALS-Glia and ALS-Ox clusters correlated with clusters C1 and C2 and C3 and C4, respectively, the ALS-TE cluster was only marginally represented in our data. As we analyzed the PFC, an area wherein TDP-43 aggregation is observed later than that in the motor cortex, our data reinforce that ALS-TE is driven by TDP-43 dysfunction[9]. Moreover, this suggests that molecular subtypes of ALS may evolve with time and are subject to changes during the disease course. Thus, exploring liquid biomarkers could aid in patient stratification and personalized clinical trials.

Furthermore, we suggest that the molecular phenotype in the four analyzed mouse models can be approximated to these subtypes, making these models potential surrogates for the molecular subgroups in humans. Notably, the oldest and most frequently used model, the SOD1.G93A mouse, correlated best with the largest clusters C1 and C2. Although not representative of all human ALS cases, our findings indicate that this model represents an important subgroup of the disease based on the identified dysregulated pathways.

Integrated analysis of multiomic datasets has identified several dysregulated pathways relevant for ALS, including mitochondrial respiration/oxidative stress, transcriptional regulation/splicing, and protein misfolding[3]. Additionally, previously less prioritized pathways—including the dysregulation of the ECM and BCNSB or the MAPK pathway—were identified. The deregulation of pathways that are only distantly related suggests the need for combinatorial drug therapies to address ALS mechanistically. Remarkably, we found evidence of female-predominant deregulation of the MAPK pathway in integrative analyses, but it did not contribute to the separation of clusters C1–C4. Therefore, the MAPK pathway may be an interesting therapeutic target for all human ALS subgroups. MAPKs are fundamental signal transducers that are involved in cell proliferation, differentiation, survival, and death[42]. Extracellular and intracellular signals are integrated by MAPK, and the overactivation of MAPK signaling, for example, via the abnormal phosphorylation of ERK1/2, has been reported in human and ALS mouse models[43]. A Ser/Thr kinase belonging to the MAPK superfamily—the MAPK/MAK/MRK [MOK] kinase—has also been linked to impaired microglial function and inflammatory insults associated with ALS pathophysiology[44]. Increased phosphorylation of ERK1/2 was also observed in our glutamate toxicity model, and pMEK2 expression was increased in the SOD1 mouse model, reproducing the aberrant activation of this pathway that was restored by the MEK2-inhibitor trametinib. However, this effect was most pronounced in females, suggesting a sex-specific efficacy. Notably, our experimental approach did not target neurons exclusively. Nevertheless, our primary cultures mainly contained neurons (~95%), and all staining procedures for both in vivo and in vitro experiments focused on neurons. While

other cell types may be involved, the evidence presented here supports the significance of the
MEK2–MAPK pathway in neurons.
Currently, one phase I/II clinical trial is examining the safety, tolerability, and efficacy of
trametinib in patients with ALS (clinicaltrials.gov identifier:NCT04326283). Our data support
the independent evaluation of male and female patients and suggest that the ECM, immune
response, and RNA processing machinery are potential targets for therapeutic intervention
that need further exploration.
Despite the limitations of our study (e.g., the retrospective view permitted by postmortem
tissue analysis and restricted balancing of sex and disease stage due to the limited number of
well-characterized postmortem brain samples), we aimed to provide valuable insights into the
molecular architecture of ALS. While analyzing over 100 brain samples helped identify
molecular subclusters, a larger sample size may provide a more detailed view of the
subgroups. Nevertheless, analyzing the PFC tissue allowed us to study the earlier stages of
the disease, offering a unique perspective compared with those of previous studies. Our robust
multiomic, computational, and integrative approaches identified subclusters that are
reminiscent of previously identified subtypes, yet show clear differences. Additionally, we
emphasize the role of splicing and transcript usage in early disease regulation, involving
several ALS-related genes. Overall, our study contributes towards unraveling the complexity
of ALS and lays the groundwork for further research in the field.
Our study also compared human brain tissue with samples from transgenic mouse models of
ALS. Although clear correlations were found between mouse models and molecular subtypes
of sALS, the four analyzed models represent specific scenarios. Other models for each of the
four studied genes could potentially yield different results[45]. Furthermore, we did not analyze
DNA methylation status or other post-translational modifications (such as glycation,
methylation, or acetylation) that could provide further insights into ALS-specific dysregulation,
as much as a single cell-based analysis could yield an additional layer of detail.
In this comprehensive integration of human tissue- and animal model-derived multiomic data,
we emphasized sex-specific differences in ALS pathology, identified molecular clusters, and
highlighted their importance for clinical trials and the development of novel therapeutic
strategies. The validation of omics results relies on multiple systems, as each mouse model
system only reproduces parts of the human pathology. Our data suggest the need for the
validation of additional molecules and pathways and justify the further exploration of the MAPK
pathway as therapeutic ALS treatment targets.
424
425

Methods

Human postmortem PFC samples

Human PFC samples were provided by four different brain banks: The Netherlands Brain Bank, King's College London Brain Bank (London Neurodegenerative Diseases Brain Bank), Parkinson's UK Brain Bank, and the Oxford Brain Bank. In total, 51 ALS and 50 CTR samples (without any signs of neurodegeneration) were included. Frozen tissues were shipped on dry ice to the rechts der Isar Hospital Department of Neurology at the Technical University of Munich and stored at -80 °C (Supplementary Table 1). Ethical approval for the use of human tissues was obtained from the Ethics Committees of the University Medical Center Göttingen (2/8/18 AN) and the Technical University Munich (145/19 S-SR). For sampling, PFC samples were transferred to a cryostat chamber at -20 °C and punched with a 20-G Quincke Spinal Needle (Becton Dickinson, Franklin Lakes, NJ, USA). Subsequently, ~20 mg tissue was collected in RNAse-/DNAse-free tubes and stored at -80 °C until further use.

ALS animal models

Four transgenic mouse models representing the most frequent ALS-causing genes were used for the multiomic studies. Animal handling and experiments in this study were performed in accordance with the applicable animal welfare laws and approved by the respective regulatory organs of the involved research centers. The animals were kept in standard cages in a pathogen-free facility under a 12-hour light/dark cycle. The mice were provided with unrestricted access to food and water. B6;129S6-Gt(ROSA)26Sortm1(TARDBP*M337V/Ypet)Tlbt/J mice[46] (here simply referred to as TDP-43 mice) were provided by the Department of Translational Neuroscience of the University Medical Center Utrecht. This model was generated by inserting an 80 kb genomic fragment carrying the human TDP-43 locus (including a patient-derived M337V mutation). TDP-43 transgenic and control wild-type mice were euthanized at the age of 26 weeks (presymptomatic stage) for biomaterial collection. B6SJL-Tg(SOD1*G93A)1Gur/J mice[32] (here referred to as SOD1 mice) were provided by the Laboratory of Translational Biomarkers, Mario Negri Institute for Pharmacological Research, Milan. High-copy number B6 congenic Tg(SOD1*G93A)1Gur/J SOD1*G93A male mice from The Jackson Laboratory (Bar Harbor, ME, USA) were bred with C57BL/6 female mice to obtain non-transgenic and mutant transgenic G93A*SOD1-expressing mice. SOD1 transgenic and control animals were sacrificed 14 weeks after birth (presymptomatic stage). (Poly)GA-NES/C9orf72(R26(CAG-Isl-175GA)-29 × Nes-Cre) mice[47] (here referred to as C9orf72 mice) were provided by the German Center for Neurodegenerative Diseases in Munich. To generate transgenic animals, plasmids were electroporated to enable the conditional expression of dipeptide-repeat proteins (DPRs). DPRs were created by inserting 175 GFP-(GA) genes encoded by non-repeating alternate codons downstream of a floxed stop cassette in the pEX CAG stop-bpA vector. Electroporation was performed on murine recombination-mediated cassette exchange embryonic stem cells at the Rosa26 safe harbor locus. Mouse lines with germline transmission, known as GAstop, were obtained and subsequently backcrossed to a C57BL/6N background until a confirmed purity of over 98% was achieved (using SNP genotyping). C9orf72 transgenic and control animals were sacrificed 4.5 weeks after birth (early symptomatic stage). Tg (Prnp-FUS)WT3Cshw/J mice[48] (hereafter referred to as FUS mice) were provided by the Laboratory of Translational Biomarkers, Mario Negri Institute for Pharmacological Research, and were sacrificed four weeks after birth. For each model, ten transgenic and ten non-transgenic mice were selected and balanced for sex.

Preparation of PFC from ALS mouse models

The mice were perfused with 50 mL of ice-cold phosphate-buffered saline (PBS) prior to
microdissection. The head was removed by cutting the base of the skull followed by skin
removal. The skull was removed using small incisions followed by microdissection of the PFC
in both hemispheres. The olfactory bulb and cerebellum were removed by cutting the
cerebellar peduncle starting from the olfactory bulb and continuing along the interhemispheric
fissure using tweezers with fine tips. The cortex was removed from the rest of the brain.
Incisions were made in the middle of the cortex to excise the PFC (Supplementary Fig. 29).
Freshly prepared PFCs were collected in nuclease-free tubes and stored at -80 °C until RNA
and protein isolation experiments were performed.
484
RNA and DNA isolation from human and mouse tissue samples
Total RNA was isolated from human and animal PFC samples using TRIzol Reagent (Sigma-
Aldrich, Taufkirchen, Germany). To ensure proper handling of the RNA, all related
experiments were conducted in an RNA workstation fume hood. In brief, 500 µL TRIzol was
added to each sample, and the tissues were homogenized using a plastic homogenizer.
Subsequently, 50 µL of 1-bromo-3-chloro-propane (Sigma-Aldrich) was added. The reaction
tubes were inverted for 10–15 s and incubated at room temperature (22–25°C) for three
minutes. The resulting lysates were centrifuged at 12 000 × g for 15 min at 4 °C, leading to
phase separation. The RNA-containing aqueous phase was carefully collected and transferred
to a fresh nuclease-free tube. To precipitate the RNA, 250 µL of 2-propanol (AppliChem,
Darmstadt, Germany) and 2 µL of GlycoBlue Coprecipitant (15 mg/mL) (Thermo Fisher
Scientific, Waltham, MA, USA) were added to each sample. The mixture was thoroughly
vortexed and incubated overnight at -20 °C. Samples were centrifuged at 12 000 × g for 30
min at 4 °C the following day. The supernatant was discarded, and the RNA pellets were
washed three times with 75% ice-cold ethanol (AppliChem). After air-drying the pellets for a
few minutes under a fume hood, they were reconstituted with 15–20 µL nuclease-free water
(Sigma-Aldrich). The reconstituted RNA was completely dissolved by incubation at 55 °C for
two minutes in a thermoshaker. Following RNA isolation, DNase treatment was performed to
eliminate DNA contamination. For this purpose, 5 µL 10× DNAse I Incubation Buffer (Life
Technologies, Carlsbad, CA, United States), 5 µL DNase I (2 U/µL), and 0.5 µL RNaseOUT
(40 U/µL) were added to each sample. Nuclease-free water was added to a final volume of 50
µL, and the samples were then incubated at 37 °C for 20 min. Finally, the RNA samples were
purified and concentrated using an RNA Clean & Concentrator-5 Kit (Zymo Research, Irvine,
CA, USA) following the manufacturer's instructions. A QIAamp DNA Mini Kit (Qiagen, Hilden,
Germany) was used for DNA isolation from human midbrain samples according to the
manufacturer's instructions. After RNA and DNA isolation, the concentration and purity of
nucleic acids were assessed using a NanoDrop One spectrophotometer (Thermo Fisher
Scientific). RNA integrity was evaluated using an Agilent 6000 NanoKit on a 2100 Bioanalyzer
(Agilent, Santa Clara, CA, USA).
514
DNA sequencing experiments and C9orf72 repeat expansion analysis
Prior to the DNA sequencing experiments, the quality of the isolated DNA was determined
using a 4200 TapeStation System (Agilent). The AmpliSeq protocol (Illumina, San Diego, CA,
USA) was used in subsequent steps. We used a target panel of 566 amplicons covering 30
ALS-related genes: *TARDBP, DCTN1, ALS2, ERBB4, TUBA4A, CHMP2B, NEK1, MATR3,*
SQSTM1, FIG4, C9orf72, SIGMAR1, VCP, GLE1, SETX, OPTN, HNRNPA1, KIF5A, TBK1,
ANG, SPG11, CCNF, FUS, PFN1, MAPT, VAPB, SOD1, CHCHD10, NEFH, and *UBQLN2.*
Using this panel, DNA samples (50–100 ng) were amplified over 14 PCR cycles. The
amplicons were digested, and AmpliSeq combinatorial dual indices were ligated for
multiplexing. The quality and quantity of the enriched libraries were validated using a 4200

TapeStation System (Agilent). The average fragment size of the amplified product was 480
bp. The libraries were standardized to a concentration of 9 nM in Tris–Cl (10 mM, pH 8.5)
supplemented with 0.1% Tween 20. Cluster generation and sequencing were performed using
the MiSeq platform (Illumina) following the standard protocol. The loading concentration was
9 pM, and 15% phiX was added. The sequencing configuration was a 250-bp paired-end
sequencing. C9orf72 repeat expansion analysis was performed using the AmplideX PCR/CE
C9orf72 Kit (Asuragen, Austin, TX, USA). Briefly, DNA samples (n = 51, 40 ng each) were
amplified using the three-primer GGGCC-repeat-primed configuration (combining flanking
primers and a GGGCC-repeat-specific primer). This configuration allows the sizing of GGGCC
alleles of up to 145 repeats and simultaneous detection of expanded GGGCC alleles with >
145 repeats. The PCR conditions consisted of initial denaturation (five minutes, 98 °C), 37
cycles (35 s, 97 °C; 35 s, 62 °C; and 3 min, 72 °C), and final elongation (10 min, 72 °C).
Capillary electrophoresis was carried out on an ABI 3730 DNA Analyzer (Applied Biosystems,
Waltham, MA. USA) using a ROX 1000 Size Ladder (Asuragen), followed by analysis with
GeneMapper 4.0 software (Applied Biosystems) and conversion of peak size to GGGGCC
repeat length via the calibration curve method according to the manufacturer's instructions.
541

Preparation of RNA libraries for sequencing experiments
The mRNA and small RNA sequencing experiments were conducted at the Functional
Genomics Center, Zürich. For mRNA sequencing, two different library preparation kits were
used: the TruSeq Stranded mRNA Kit (Illumina; short read sequencing) and the SMARTer
Stranded Total RNA-Seq Kit v2 Pico Input Mammalian (Takara Bio USA, San Jose, CA, USA;
short read sequencing). In the TruSeq protocol, total RNA samples (100–1000 ng) were
subjected to poly-A enrichment and reverse transcription to generate double-stranded cDNA.
The cDNA was fragmented, end-repaired, adenylated, and ligated to TruSeq adapters
containing unique dual indices for multiplexing. Fragments with adapters at both ends were
selectively enriched using PCR amplification, resulting in a smear with an average fragment
size of ~260 bp. The libraries were subsequently normalized to 10 nM in Tris–Cl (10 mM, pH
8.5) containing 0.1% Tween 20. For the SMARTer Stranded Total RNA-Seq Kit protocol, total
RNA samples (0.25–10 ng) were reverse-transcribed using random priming into double-
stranded cDNA in the presence of a template switch oligo (TSO). The generated cDNA
fragments contained sequences derived from random priming oligos and the TSO. PCR
amplification with primers binding to these sequences incorporated full-length Illumina
adapters, including a multiplexing index. The ribosomal cDNA was cleaved using ZapR in the
presence of mammalian R-Probes. The remaining fragments were enriched through a second
round of PCR amplification using primers designed to match the Illumina adapters, resulting
in a smear with an average fragment size of ~360 bp. The libraries were normalized to 5 nM
in Tris–Cl (10 mM, pH 8.5) containing 0.1% Tween 20. To validate the quality and quantity of
the isolated RNA and enriched libraries, an Agilent Fragment Analyzer was used for the
TruSeq kit, while the 4200 TapeStation System (Agilent) was used for the SMARTer Kit.
565

RNA sequencing
Sequencing was performed using the Illumina platforms NovaSeq 6000 (for transcriptomics)
and HiSeq 2500 (for small RNA sequencing) according to standard protocols. Small RNA
sequencing was performed using RealSeq-AC miRNA (SomaGenics, Santa Cruz, CA, USA)
(short read sequencing). All samples were quantified and quality was controlled using an
Agilent Fragment Analyzer. Briefly, RNA samples (1 ng–1 ug) were adaptor-ligated,
circularized, and reverse-transcribed into cDNA. The cDNA samples were amplified using
PCR, which also incorporated sample barcodes. The library product, a peak with a fragment
size of ~149 bp, was normalized to 10nM in Tris–Cl (2 mM, pH 8.5) containing 0.1% Tween

20. The quality and quantity of the enriched libraries were validated using an Agilent Fragment
Analyzer. Transcriptomics data were processed using the Nextflow Core RNASeq pipeline
version 3.0[49]. The data were demultiplexed with bcl2fastq, and the fastq files underwent
several quality checks, including FastQC[50]. Salmon[51] was used for pseudoalignment and
quantitation with a Salmon index built using GRCm39 with annotations from GENCODE vM26
for mouse data and GRCh38 with annotations from GENCODE v37 for human data. Count
matrices from Salmon were used for the downstream analyses (Supplementary Table 13).
The count matrices were filtered, retaining genes with at least ten counts in 50% of the
samples for any condition or sex. We used the clusterProfiler R package[52] and GO biological
processes and molecular functions for GSEA, filtering terms by size between 10 and 500
genes and correcting for multiple testing (Benjamini–Hochberg correction).
586
Global proteomics of mouse and human PFC tissue samples
Tissues from both human PFCs and four different mouse models were prepared. The tissues
were ground with a biomasher using 350 µL MeOH:H_2O (4:1). Protein pellets were
resuspended in 200 µL Laemmli buffer (10% SDS, 1 M Tris pH 6.8, and glycerol) and then
centrifuged at 16.600 x g at 4 °C for five minutes. The protein concentration was determined
using a DC assay (Bio-Rad, Hercules, CA, USA) following the manufacturer's instructions.
Each sample, containing 100 µg protein lysate, was heated at 95 °C for five minutes and
loaded onto a 5% acrylamide SDS–PAGE stacking gel (prepared in-house). Prior to overnight
digestion at 37 °C using modified porcine trypsin (Mass Spectrometry Grade, Promega,
Madison, WI, USA) at an enzyme–protein ratio of 1:80, the gel bands were reduced and
alkylated. The resulting peptides were extracted using 60% acetonitrile (ACN) followed by
100% ACN. The peptides were resuspended in 30 µL H_2O, 2% ACN, and 0.1% fluoroacetic
acid. iRT peptides (Biognosys, Schlieren, Switzerland) were added to each sample as an
internal quality control (QC). NanoLC–MS/MS analyses were conducted using a nanoAcquity
UltraPerformance LC (UPLC) system (Waters, Milford, MA, USA) coupled to a Q-Exactive
Plus Mass Spectrometer (Thermo Fisher Scientific). The solvent system consisted of 0.1%
formic acid (FA) in water (solvent A) and 0.1% FA in ACN (solvent B). Samples equivalent to
800 ng protein were loaded onto a Symmetry C18 precolumn (20 mm × 180 µm with 5 µm
diameter particles, Waters) over three minutes at a flow rate of 5 µL/min with 99% solvent A
and 1% solvent B. Peptides were separated on an ACQUITY UPLC BEH130 C18 column (250
mm × 75 µm with 1.7 µm diameter particles) at a flow rate of 400 nL/min. The following gradient
of solvent B was used: 1–8% over 2 min, 8–35% over 77 min, 35–90% over 1 min, 90% for 5
min, and 90–1% over 2 min. Samples from each cohort were injected in a random order. The
system was operated in data-dependent acquisition (DDA) mode, automatically switching
between MS (mass range 300–1800 *m/z* with R = 70 000, automatic gain control [AGC] fixed
at 3.10^6 ions, and the maximum injection time set to 50 ms) and MS/MS (mass range 200–
2000 *m/z* with R = 17 500, AGC fixed at 1.10^5, and the maximum injection time set to 100 ms)
modes. The ten most abundant precursor ions were selected on each MS spectrum for further
isolation and higher energy collisional dissociation, excluding monocharged and unassigned
ions. The dynamic exclusion time was set to 60 s. A sample pool was injected as an external
QC every six samples for the human cohort and every five samples for the mouse cohort.
MaxQuant[53] version 1.6.14 was used to process raw data. The Andromeda[54] search engine
was used to assign peaks with trypsin/P specificity against a protein sequence database
generated in-house containing all human (20 421 entries as of August 24, 2020) or mouse
entries (17 134 entries for SOD1 and TDP-43 models as of March 27, 2020, and 17 061 entries
for C9orf72 and FUS models as of September 29, 2020) extracted from UniProtKB/Swiss-
Prot. Methionine oxidation/acetylation of the protein N-termini was used as a variable
modification, whereas cysteine carbamidomethylation was used as a fixed modification. The

"match between runs" option was enabled to facilitate protein quantification. To control false
discoveries, a maximum false discovery rate of 1% was implemented (at both peptide and
protein levels), employing a decoy strategy. Intensities were extracted from the
Proteingroup.txt file for statistical analyses. The MaxQuant protein vs. sample table was used
for downstream analyses, including label-free quantitation intensities. Only Swiss-Prot
proteins were retained, whereas TrEMBL proteins were removed for higher reliability. After
filtering out low abundant proteins, i.e., proteins that were detected in less than 50% of the
samples in any combination of condition and sex, and imputing missing values using the
missForest[55] algorithm, the intensities were log2-transformed and used for principal
component exploration, constructing heat maps, and differential abundance analysis. The
limma package[56] was used for linear modeling, and p-values were adjusted using the
Benjamini–Hochberg correction. The protein names were mapped to the corresponding genes
and used to search for enriched biological processes and molecular functions using the criteria
described in the transcriptomics data processing, with a $p < 0.01$ threshold for functional
annotation analyses.
640
Phosphoproteomics on mouse PFC brain tissue samples
Starting with the protein extracts from the global proteomics experiments, protease inhibitors
(Sigma-Aldrich, P8340) and phosphatase inhibitors (final concentration in Na_3VO_4 = 1 mM)
were added to samples. Protein concentration was determined using the RC DC Protein Assay
(Bio-Rad) according to the manufacturer's instructions. Proteins for each sample (250 µg)
were reduced and alkylated prior to an in-house optimized single-pot, solid-phase-enhanced
sample preparation protocol (adapted from Hughes et al., Nat Protoc, 2019). Briefly, beads A
(Sera-Mag SpeedBeads, Fisher Scientific, Schwerte, Germany, 45152105050250) and B
(Sera-Mag SpeedBeads, Fisher Scientific, 65152105050250) were combined (1:1) and, after
three washing steps with H_2O, were added to the samples (bead–protein ratio of 10:1 for each
type of bead, meaning a 20:1 ratio for the combination of beads). After inducing protein binding
to the beads with 100% ACN for 18 min, the bead and protein mixtures were washed twice
with 80% ethyl alcohol and once with 100% ACN before being resuspended in 95 µL NH_4HCO_3
prior to overnight on-bead digestion (enzyme–protein ratio of 1:20) at 1 000 rpm at 37 °C using
modified porcine trypsin/Lys-C Mix (Mass Spectrometry Grade, Promega). The digestion was
stopped using trifluoroacetic acid (TFA) (final pH < 2). The recovered peptides were
resuspended in 170 µL 80% ACN and 0.1% TFA, and MS PhosphoMix I Light (Sigma-Aldrich)
was added to each sample (peptide [µg]–mix [fmol] ratio of 1:6). Phosphopeptide enrichment
was performed on 5 µL phase Fe(III)–NTA cartridges on an AssayMAP Bravo platform
(Agilent) following the immobilized metal affinity chromatography protocol. Briefly, cartridges
were washed and primed with 50% ACN, 0.1% TFA and then equilibrated with 80% ACN,
0.1% TFA. Subsequently, 100-µL samples were loaded onto the phase at a rate of 2 µL/min
and then washed with 80% ACN, 0.1% TFA before being eluted in 20 µL 1% NH_4OH at 5
µL/min. After enrichment, FA and MS PhosphoMix I Heavy (Sigma-Aldrich) (peptide [µg]–mix
[fmol] ratio of 1:6) were added to each sample. Dried phosphopeptides were resuspended in
40 µL H_2O, 2% ACN, 0.1% FA. Sample preparation steps for the C9orf72 and FUS mouse
models were identical to those previously described for SOD1 and TDP-43, except that
proteins were extracted from new tissue samples immediately before the phosphoproteomics
experiment.
NanoLC–MS/MS analyses were conducted using a nanoAcquity UPLC system (Waters)
coupled with a Q-Exactive HF-X Mass Spectrometer (Thermo Scientific) equipped with a
Nanospray Flex ion source. The solvent system consisted of 0.1% FA in water (solvent A) and
0.1% FA in ACN (solvent B). Samples were loaded onto an ACQUITY UPLC Peptide BEH
C18 Column (250 mm x 75 µm, 1.7 µm diameter particles) over three minutes at a flow rate

of 5 µL/min with 99% solvent A and 1% solvent B. Phosphopeptides were separated on an
ACQUITY UPLC M-Class Symmetry C18 Trap Column (20 mm x 180 µm, 5 µm diameter
particles; Waters) at a flow rate of 400 nL/min using a gradient of solvent B as follows: 1–2%
over 2 min, 2–35% over 77 min, and finally 35–90% over 1 min. The samples in each group
were injected in random order. The instrument was operated in DDA mode with automatic
switching between the MS and MS/MS modes. MS scans were performed in the mass range
of 375–1500 m/z with R = 120 000, an AGC fixed at 3.106 ions, and a maximum injection time
of 60 ms. MS/MS scans were conducted in the mass range of 200–2000 m/z with R = 15 000,
an AGC fixed at 1.105 ions, and a maximum injection time of 60 ms. In each MS spectrum,
the ten most abundant ions were selected for further isolation and higher-energy collisional
dissociation, excluding monocharged and unassigned ions. A dynamic exclusion time of 40 s
was used to prevent the reanalysis of previously selected ions. The obtained raw
phosphoproteomics data were processed using MaxQuant version 1.6.14. Peaks were
assigned using the Andromeda search engine with trypsin/P specificity against an in-house-
generated protein sequence database containing mouse entries from UniProtKB/Swiss-Prot
(17 061 entries as of September 29, 2020). The peptides were required to have a minimum
length of seven amino acids, and a maximum of one missed cleavage was allowed. These
modifications include methionine oxidation, acetylation of protein N-termini, and
phosphorylation of serine, threonine, and tyrosine. Cysteine carbamidomethylation was used
as a fixed modification. Protein quantification utilized the "match between runs" option. A
maximum false discovery rate of 1% (at both peptide and protein levels) was implemented
using a decoy strategy. The intensities were extracted from the Phospho (STY).txt file and
processed using the Perseus platform[57] version 2.0.7.0, during which contaminants and
reversed proteins, as well as proteins with negative scores, were removed. Using the "expand
sites table" option, the intensities of the different phosphopeptides involved in one phosphosite
were summed, and phosphosites with a localization probability below 75% were removed. The
Perseus output table was used for statistical analyses.
Only proteins that were detected in more than 50% of the mouse samples of any combination
of sex and condition were retained for each dataset. The intensities were log2-transformed,
quantile normalization was applied, and missing values were imputed to the lowest quartile.
The remaining data processing for phosphoproteomics followed the procedure described
above for proteomics data processing.
707
Small RNA sequencing data-processing and miRNA target prediction
Small RNA data were processed using the Nextflow Core smRNASeq pipeline version 1.0[49].
Reads were trimmed and aligned against miRBase[58] version 22.1 using Bowtie1[59], both for
mature miRNAs and hairpins. miRNAs with at least ten counts in 50% of the samples of any
condition and sex were retained, and the rest were filtered out. Unnormalized count matrices
were used for subsequent DESeq2[60] differential expression analysis after stratifying by sex
and adjusting for p-values using the Benjamini–Hochberg correction. Mature miRNAs were
mapped to their corresponding genes using miRDB version 6.0[61], excluding matches with
scores < 60 or > 800 targets, as recommended. For each miRNA present in the miRNA
expression matrices, we obtained experimentally validated targets from miRTarBase[62] 8.0 and
predicted targets from miRDB miRBase version 22[61]. miRTarBase[62] provides the most
extensive curated database of validated miRNA–target interactions (MTI) collected from
literature using natural language processing to select functional miRNA studies. Additionally,
the miRDB database includes MTIs predicted by MirTarget, which uses a support vector
machine (SVM) to analyze thousands of high-throughput sequencing experiments[63]. Each
experiment is assigned a probability score, which serves as the SVM's output. A higher
probability score indicates a higher likelihood of accurate target prediction. Therefore, we set

a threshold of 0.6 on the output probabilities to select only highly likely MTIs. Finally, we joined
the miRNA–target pairs from both sources for further analysis.

Cell-type deconvolution analyses

Single-cell reference-based cell-type deconvolution of RNA-Seq in mouse models was
performed using Scaden[19]. We used healthy adult scRNA-Seq datasets for mouse[64] and
human[65]. Scaden uses a fully connected deep neural network ensemble trained on pseudo-
bulks simulated from reference scRNA-Seq data. Before deconvolution, we filtered the
scRNA-seq data using Scanpy[66] to maintain at least 200 genes expressed per cell and at least
five cells expressing one gene. For Scaden, counts per million (CPM) of simulated pseudo-
bulks and transcripts per million (TPMs) of the data to be deconvolved were used. In this study,
CPM was used for scRNA-Seq instead of TPMs because scRNA-Seq consists of unique
molecular identifier counts and does not include gene-length bias. We used a variance cutoff
of 0.01, and the mean squared error was calculated for each batch as loss.

Differential Alternative Splicing analysis

The splicing tool SUPPA2 version 2.3 was used to analyze DAS for seven alternative splicing
events: exon skipping, mutually exclusive exons, intron retention, alternative 3'-splice site,
alternative 5'-splice site, alternative first exon, and alternative last exon. SUPPA2 was used
with multipleFieldSelection() to select the TPM values of the transcripts, followed by
GenerateEvents with the parameters -f ioe -e SE SS MX RI FL and annotation files GENCODE
v37 for human data and GENCODE vM26 for mouse data. The inclusion values
(PSupplementary) were calculated using psiPerEvent, and the differences in PSupplementary
(ΔPSupplementary) between the mutant and control conditions were determined using
diffSplice with parameters -m empirical -l 0.05 -gc –save tpm_events to detect anomalies in
the splicing landscape.

Multi-Omics Factor Analysis

We used MOFA 2 version 1.4.0 to integrate data from multiple omics levels for the human
cohort, including transcriptomics, miRNA, and proteomics. The MOFA model was trained on
the data, and downstream analyses were performed. Each omics type was preprocessed in
its own manner, and the default training parameters were used. The MOFA models were
initialized with 15 initial factors, and convergence was reached when the evidence lower bound
(ELBO) value did not change with more than a deltaELBO value of 1e-4%.

Gene ontology, pathway enrichment analyses, and protein interaction networks

GSEA was performed using gseGO and gseKEGG from the clusterProfiler R package, with
biological processes and molecular functions chosen as background databases for GO
enrichment. The *p*-value cutoff was set at 0.05. Differential expression was presented using
volcano plots generated with the Enhanced Volcano plot package in R, and over-
representation analysis was performed on genes that showed at least one significant DAS
event using the clusterProfiler function enrichGO, with a *p*-value cutoff of 0.1 and Benjamini–
Hochberg correction for multiple hypothesis testing. Protein–protein interaction networks were
created using the STRING protein interaction network database version 11.0[67,68] using
standard settings.

Weighted Correlation Network Analysis

To conduct a weighted gene co-expression analysis, we employed the WGCNA package[15].
Pairwise Pearson correlations were calculated to establish signed regulatory networks within
the WGCNA. By constructing an adjacency matrix, we applied a soft-thresholding technique

to approximate a scale-free topological network. Eigengenes or eigenproteins were calculated
as the first principal components of each module. This resulted in the development of several
modules. We merged similar modules based on hierarchical clustering (SOD1: 0.4, C9orf72:
0.4, FUS: 0.5, TDP-43: 0.4 for RNA-Seq and 0.25 for proteomics). We calculated the
relationships between WGCNA modules and traits. Sex was also included in the traits resulting
in four traits: male ALS, female ALS, male CTR, and female CTR. First, we filtered modules
based on the significance of the module–gene relationship ($p < 0.05$) and then selected
modules that were highly correlated with either male or female ALS. The correlation cutoffs
differed between the mouse models (SOD1: 0.5, C9orf72: 0.3, FUS: 0.3, and TDP-43: 0.5 for
RNA-Seq and SOD1: 0, C9orf72: 0, FUS: 0, and TDP-43: 0 for proteomics). Using WGCNA,
we also analyzed co-expression networks in mouse models. The minimal module size was set
to 30 with a merge height of 0.4–0.5 and a correlation threshold of 0.3–0.5 (Supplementary
Table 14).
788
Reproducible data pipeline
Given the size and complexity of the datasets, data processing, and data analysis methods,
we integrated all analyses and raw data into a Data Version Control pipeline[69]. The pipeline
uses both Nextflow and Docker images to guarantee a reproducible execution environment.
This enabled us to keep all results consistent because any change or exploration of a data-
processing method automatically triggered the recalculation of all dependent analyses and
integrations.
796
Mice primary cortical cultures
Primary cortical neuronal cell cultures were generated from neonatal mouse C57BL/6J pups
aged postnatal day 0–1 (P0–P1) in accordance with ethical guidelines for animal
experimentation at local and international levels. Animal care strictly followed official
governmental protocols, with utmost effort to minimize the number of animals utilized and to
mitigate any potential suffering or distress. Pups were decapitated, and the brains were
collected in dissection media containing 10× Hanks balanced salt solution and $NaHCO_3$. The
cortex was dissected, the meninges were removed, and small pieces of the cortex were
collected in a Falcon tube. Tissues were trypsinized at 37 °C in a water bath for 12 min and
treated with 200 µL DNAse I (10 mg/mL). The tissue strips were gently triturated (until the
tissue fragmented) in fetal bovine serum using a fire-polished Pasteur pipette. The mixture
was subsequently centrifuged at 800 × g for four minutes, and the resulting cell pellet was
resuspended and cultured in a neurobasal medium supplemented with B27 and antibiotics
(0.06 µg/mL penicillin and 0.1 µg/mL streptomycin). Cells were seeded at a density of 3 × 10^5
cells per well in 24-well plates. Prior to cell seeding, the coverslips were acid-washed, rinsed
several times with water, sterilized with ethanol and UV light, and placed in a well plate. The
plates were then coated with poly-L-ornithine (0.05 mg/mL) overnight and laminin (10 µg/mL)
for 2 h in an incubator before use. The cells were maintained at 37 °C in a humidified incubator
under 5% CO_2 for seven days prior to the experiments. The cell culture medium was replaced
every three days during this period. To induce glutamate excitotoxicity, L-glutamic acid (Tocris,
UK) was dissolved in 50 mM NaOH, and a stock solution of 50 mM was prepared prior to use.
An appropriate concentration of glutamate was prepared in the maintenance medium
(neurobasal medium supplemented with B27 and antibiotics). The cells were exposed to 5
mM glutamate by exchanging 1:3 medium at seven days in vitro. After six hours of incubation,
glutamate was thoroughly removed by washing, and the cells were fixed for
immunocytochemistry or lysed for protein extraction.
823
Immunocytochemistry and microscopy for in vitro experiments

Cells were cultured on coverslips following previously described methods[70] and
immunostained at seven days in vitro according to standard techniques. Cells were fixed in
4% paraformaldehyde in PBS at room temperature for 10 min. To quench the free aldehyde
groups, the cells were treated with 50 mM NH_4Cl for 15 min and washed with PBS. For the
permeabilization of the cell membrane, PBS with 0.25% Triton X-100 was added for 10 min at
room temperature. Non-specific binding sites were blocked by applying 10% goat serum in
PBS for at least 20 min. Dilutions of primary antibodies were prepared in blocking solution to
a final volume of 180 µL per 18-mm coverslip, and cells were incubated for 90 min at 37 °C
with shaking. The following primary antibodies were used: mouse anti-MAP2 (Invitrogen,
Waltham, MA, USA; #MA5-12826) (1:500) and rabbit anti-cleaved caspase 3 (Cell Signaling
Technology, Danvers, MA, USA; #9661) (1:250). The cells were washed three times for five
minutes each with PBS before applying the secondary antibodies. The cells were then
incubated with secondary antibodies for 30 min and washed. For double staining, a second
primary antibody was added, and the same steps were repeated. Coverslips were mounted
on slides using a mounting medium supplemented with DAPI. Images were captured using an
inverted fluorescence microscope (Zeiss, Jena, Germany) with a 63× oil objective and
analyzed using ImageJ software. Fifteen random images from each coverslip were analyzed
for cell death by counting the number of cleaved caspase-3 positive cells. Neurite lengths were
measured using a simple neurite tracing plugin in ImageJ. Statistical analyses were conducted
using GraphPad Prism version 9.4.1 (GraphPad, San Diego, CA, USA). Outliers were
identified and removed using the Grubbs test (α = 0.1). Comparisons were done using one-
way analysis of variance (ANOVA), and data were plotted as the mean ± standard error of the
mean (SEM) of at least five independent experiments. Differences were considered
statistically significant at $p < 0.05$.
849
Protein extraction and western blotting
For protein analysis, cells were washed once with 1× PBS, and after adding RIPA lysis buffer,
protease inhibitor cocktail (1:25), and phosphatase inhibitor (1:20), cells were incubated on
ice for five minutes. The cells were scraped off with a cell scraper, transferred into 1-mL
reaction tubes, and homogenized by passage through a U-100 insulin syringe several times.
Protein concentration was determined using a Pierce BCA Protein Assay Kit (Thermo Fisher
Scientific) following the manufacturer's instructions. One microliter of protein sample was used
for the assay. Colorimetric reactions were analyzed using an Infinite M200 PRO ELISA plate
reader (Tecan, Männedorf, Switzerland). Twenty grams of each sample were loaded onto a
gel (NuPAGE 4 to 12%, Bis-Tris, Invitrogen). NuPAGE LDS sample buffer (1:4) and sample-
reducing buffer (1:10) were added to lysed protein before loading onto the gels, incubated with
shaking at 75 °C for 13 min, and centrifuged at 12 000 × g at 4 °C. The proteins were subjected
to gel electrophoresis at 200 V. Subsequently, the proteins were transferred onto nitrocellulose
membranes using an iBlot2 gel transfer device (Thermo Fisher Scientific). The membranes
were then blocked with 5% nonfat milk in PBST at room temperature for 30 min. For antibody
incubation, primary antibodies (diluted in blocking buffer) were added to the membranes and
allowed to incubate overnight at 4 °C with rotation. After four washes with PBST (five minutes
each), the membranes were incubated with highly sensitive horseradish peroxidase-labeled
secondary antibodies (diluted 1:10 000 in blocking buffer) at room temperature for one hour.
Finally, the membranes were washed thoroughly with PBST to remove the unbound
antibodies. Blots were incubated with enhanced chemiluminescence reagent and were
imaged using a Molecular Imager ChemiDoc (Bio-Rad) imaging system. Band signal
intensities were quantified using ImageJ software and normalized to housekeeping proteins
and controls. Statistical analyses were conducted using GraphPad Prism version 9.4.1.
Outliers were identified and removed using the Grubbs test (α = 0.1). Comparisons were done

using one-way ANOVA, and data were plotted as the mean ± SEM of at least three
independent experiments. Differences were considered statistically significant at $p < 0.05$.
877

Tissue protein extraction for immunoblot analysis
Spinal cords were homogenized in five volumes (w/v) of 1% boiling SDS[71]. Protein
homogenates were sonicated, boiled for 10 min, and centrifuged at 13 500 × g for five minutes.
Supernatants were analyzed by dot blotting. For detergent-insoluble protein extraction, mouse
tissues were homogenized in 10 volumes (w/v) of buffer (15 mM Tris–HCl pH 7.6, 1 mM
dithiothreitol, 0.25 M sucrose, 1 mM $MgCl_2$, 2.5 mM EDTA, 1 mM EGTA, 0.25 M Na_3VO_4, 2
mM $Na_4P_2O_7$, 25 mM NaF, 5 µM MG132) and a protease inhibitor cocktail (Roche, Basel,
Switzerland), as previously described[72]. Briefly, the samples were centrifuged at 10 000 × g,
and the pellet was suspended in an ice-cold homogenization buffer containing 2% Triton X-
100 and 150 mM KCl. The samples were then centrifuged at 10 000 × g to obtain the Triton-
insoluble (insoluble) fraction.
889

Immunohistochemistry
Mice were anesthetized and underwent transcardial perfusion using 50 mL PBS followed by
100 mL of 4% paraformaldehyde (in PBS). The spinal cord was promptly dissected, postfixed
for three hours, and subsequently transferred to a solution of 20% sucrose in PBS overnight.
Once sank, the spinal cord was transferred to a 30% sucrose solution until it reached the
desired consistency. Next, the spinal cord was frozen in n-pentane at 45 °C and stored at ~-
80 °C. Prior to freezing, the spinal cord was divided into cervical, thoracic, and lumbar
segments and embedded in Tissue-Tek O.C.T. compound (Sakura Finetek, Torrance, CA,
USA). Coronal sections (30-µm thick) were sliced from the lumbar spinal cord and subjected
to immunohistochemistry. Rabbit monoclonal anti-pMEK (Ser221) (pMEK2) antibody (1:50,
Cell Signaling; RRID: AB_490903) was used. In brief, slices were incubated with blocking
solutions (0.2% Triton X-100 and 2% normal goat serum) at room temperature for one hour.
Subsequently, the slices were incubated overnight at 4 °C with primary antibodies. The
sections were then incubated with biotinylated secondary antibodies (1:200) for one hour at
room temperature, followed by immunostaining using an avidin–biotin kit and
diaminobenzidine. To facilitate visualization, sections were counterstained with 0.5% cresyl
violet. Stained sections were captured at 20× and 40× magnification using an Olympus BX-61
Virtual Stage microscope (Olympus Life Sciences, Tokyo, Japan), ensuring complete stitching
of the entire section with a pixel size of 0.346 µm. Images were acquired using 6-µm-thick
stacks (step size of 2 µm). The different focal planes were merged into a single stack using
mean intensity projection to maintain consistent focus across the sample. Finally, the acquired
signals were analyzed for each slice using ImageJ and OlyVIA software (Olympus Life
Sciences).

Immunoblotting for in vivo experiments
Protein levels were quantified using a BCA Protein Assay Kit (Pierce Biotechnology, Waltham,
MA, USA) and analyzed by western blotting and dot blotting following previously described
methods[73]. Membranes were blocked with a solution of 3% (w/v) BSA (Sigma-Aldrich) and
0.1% (v/v) Tween 20 in Tris-buffered saline (pH 7.5). Subsequently, the membranes were
incubated with primary antibodies followed by peroxidase-conjugated secondary antibodies
(GE HealthCare, Chicago, IL, USA). The antibodies utilized were rabbit monoclonal anti-
pMEK2 (1:2000, Cell Signaling Technology; RRID: AB_490903), rabbit monoclonal anti-
phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (1:2000, Cell Signaling Technology; RRID:
AB_2315112), rabbit polyclonal anti-human SOD1 (1:1000, StressMarq Biosciences, Victoria,
Canada; RRID: AB_2704217), rabbit polyclonal anti-ubiquitin (1:1000, Abcam, Cambridge,

UK; RRID: AB_306069), and mouse monoclonal anti-SQSTM1/p62 (p62) (1:500, Abcam).
Peroxidase-conjugated secondary antibodies (goat anti-mouse and anti-rabbit) (GE
HealthCare) were used at dilutions of 1:20 000 and 1:10 000, respectively. The blots were
developed using the Luminata Forte Western Chemiluminescent HRP Substrate
(MilliporeSigma, Burlington, MA, USA) and visualized on a ChemiDoc Imaging System (Bio-
Rad). Densitometry analysis was performed using Image Lab software version 6.0 (Bio-Rad).
Protein immunoreactivity was normalized to Ponceau Red staining (Honeywell Fluka,
Charlotte, NC, USA) for accurate quantification.
932
Preclinical study in SOD1^{G93A} mice
Starting at the presymptomatic stage (nine weeks of age), SOD1^{G93A} female and male mice
received a 3 mg/kg dose of trametinib (n = 10) or vehicle (PBS) (n = 10) via intranasal delivery.
The drug was administered twice per week for seven weeks (from 9 to 16 weeks of age). At
16 weeks of age, the mice were sacrificed, and spinal cords and plasma were collected for
subsequent biochemical analysis. The animal experimental protocols used in this study (Prot.
no. 9F5F5.143 and 9F5F5.250) were prospectively reviewed and approved by the Mario Negri
Institutional Animal Care and Use Committee and the Italian Ministry of Health (*Direzione
Generale della Sanità Animale e dei Farmaci Veterinari, Ufficio 6).* This study complied with
international and local animal welfare standards.
943
Plasma neurofilament light chain quantification
Mouse plasma samples were collected in K2-EDTA BD Microtainer blood collection tubes and
centrifuged at 5 000 × *g* for five minutes to obtain the plasma. The plasma NfL was quantified
using a Simoa NF-light Advantage (SR-X) Kit (#103400) on a Quanterix SR-X platform
(Quanterix, Boston, MA, USA). All reagents used were from a single lot, and measurements
were performed according to the manufacturer's protocol.
950
Statistical analyses of in vivo experiments
Statistical analyses of in vivo experimental data were performed using GraphPad Prism 7.0.
Student's *t*-tests or one-way ANOVAs, followed by posthoc tests, were used to analyze the
differences between the experimental groups for each variable. Statistical significance was
set at *p* < 0.05.
Data availability
Raw RNA-Seq and processed gene expression data were deposited to the National Center
for Biotechnology Information Gene Expression Omnibus database (GSE234246). Encrypted
raw RNA-Seq data for the human cohort were deposited to the European Genome Phenome
Archive (registered study: EGAS00001007318). Proteomics and phosphoproteomics datasets
have been deposited in the ProteomeXchange Consortium database with the identifiers
PXD043300 and PXD043297, respectively.
963
Code availability
For RNA-Seq, we used the Nextflow pipeline for alignment and DESeq2 for differential
analysis: https://www.nextflow.io/,
https://www.bioconductor.org/packages//2.10/bioc/html/DESeq.html
Alternative splicing was performed using SUPPA, DRIMSeq, and DEXSeq
(https://github.com/comprna/SUPPA,
https://bioconductor.org/packages/release/bioc/html/DRIMSeq.html,
https://bioconductor.org/packages/release/bioc/html/DEXSeq.html

Enrichment was performed with:

https://bioconductor.org/packages/release/bioc/html/clusterProfiler.html, http://revigo.irb.hr/

Gene–gene interaction network analysis was done with:

https://cran.r-project.org/web/packages/WGCNA/index.html

Deconvolution was performed with Scaden:

https://scaden.readthedocs.io/en/latest/

References

1. Mehta, P. *et al.* Prevalence of amyotrophic lateral sclerosis in the United States using established and novel methodologies, 2017. *Amyotroph. Lateral Scler. Front. Degener.* **24**, 108–116 (2023).
2. Suzuki, N., Nishiyama, A., Warita, H. & Aoki, M. Genetics of amyotrophic lateral sclerosis: seeking therapeutic targets in the era of gene therapy. *J. Hum. Genet.* **68**, 131–152 (2023).
3. Mead, R. J., Shan, N., Reiser, H. J., Marshall, F. & Shaw, P. J. Amyotrophic lateral sclerosis: a neurodegenerative disorder poised for successful therapeutic translation. *Nat. Rev. Drug Discov.* **22**, 185–212 (2023).
4. Braak, H. & Braak, E. Neuropathological stageing of Alzheimer-related changes. *Acta Neuropathol. (Berl.)* **82**, 239–259 (1991).
5. Braak, H. *et al.* Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol. Aging* **24**, 197–211 (2003).
6. Brettschneider, J. *et al.* Stages of pTDP-43 pathology in amyotrophic lateral sclerosis. *Ann. Neurol.* **74**, 20–38 (2013).
7. Aronica, E. *et al.* Molecular classification of amyotrophic lateral sclerosis by unsupervised clustering of gene expression in motor cortex. *Neurobiol. Dis.* **74**, 359–376 (2015).
8. Morello, G. *et al.* Integrative multi-omic analysis identifies new drivers and pathways in molecularly distinct subtypes of ALS. *Sci. Rep.* **9**, 9968 (2019).
9. Tam, O. H. *et al.* Postmortem Cortex Samples Identify Distinct Molecular Subtypes of ALS: Retrotransposon Activation, Oxidative Stress, and Activated Glia. *Cell Rep.* **29**, 1164-1177.e5 (2019).
10. Prudencio, M. *et al.* Distinct brain transcriptome profiles in C9orf72-associated and sporadic ALS. *Nat. Neurosci.* **18**, 1175–1182 (2015).
11. Figueroa-Romero, C. *et al.* Expression of microRNAs in human post-mortem amyotrophic lateral sclerosis spinal cords provides insight into disease mechanisms. *Mol. Cell. Neurosci.* **71**, 34–45 (2016).
12. Umoh, M. E. *et al.* A proteomic network approach across the ALS-FTD disease spectrum resolves clinical phenotypes and genetic vulnerability in human brain. *EMBO Mol. Med.* **10**, 48–62 (2018).
13. Eshima, J. *et al.* Molecular subtypes of ALS are associated with differences in patient prognosis. *Nat. Commun.* **14**, 95 (2023).
14. Kenna, K. P. *et al.* NEK1 variants confer susceptibility to amyotrophic lateral sclerosis. *Nat. Genet.* **48**, 1037–1042 (2016).
15. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* **9**, 559 (2008).
16. Buratti, E. & Baralle, F. E. The multiple roles of TDP-43 in pre-mRNA processing and gene expression regulation. *RNA Biol.* **7**, 420–429 (2010).
17. Polymenidou, M. *et al.* Misregulated RNA processing in amyotrophic lateral sclerosis. *Brain Res.* **1462**, 3–15 (2012).
18. Ma, X. R. *et al.* TDP-43 represses cryptic exon inclusion in the FTD-ALS gene UNC13A. *Nature* **603**, 124–130 (2022).
19. Menden, K. *et al.* Deep learning-based cell composition analysis from tissue expression profiles. *Sci. Adv.* **6**, eaba2619 (2020).
20. Månberg, A. *et al.* Altered perivascular fibroblast activity precedes ALS disease onset. *Nat. Med.* **27**, 640–646 (2021).
21. Rinchetti, P., Rizzuti, M., Faravelli, I. & Corti, S. MicroRNA Metabolism and Dysregulation in Amyotrophic Lateral Sclerosis. *Mol. Neurobiol.* **55**, 2617–2630 (2018).

22. Hirano, M. *et al.* Mutations in the gene encoding p62 in Japanese patients with amyotrophic lateral sclerosis. *Neurology* **80**, 458–463 (2013).

23. Beckers, J., Tharkeshwar, A. K. & Van Damme, P. C9orf72 ALS-FTD: recent evidence for dysregulation of the autophagy-lysosome pathway at multiple levels. *Autophagy* **17**, 3306–3322 (2021).

24. Reijnders, M. J. M. F. & Waterhouse, R. M. Summary Visualizations of Gene Ontology Terms With GO-Figure! *Front. Bioinforma.* **1**, 638255 (2021).

25. Biogen. *A Phase 1, Double-Blind, Placebo-Controlled, Single-Ascending-Dose Study to Evaluate the Safety, Tolerability, Pharmacokinetics, and Pharmacodynamics of BIIB100 Administered Orally to Adult Participants With Amyotrophic Lateral Sclerosis.* https://clinicaltrials.gov/ct2/show/NCT03945279 (2023).

26. Argelaguet, R. *et al.* Multi-Omics Factor Analysis-a framework for unsupervised integration of multi-omics data sets. *Mol. Syst. Biol.* **14**, e8124 (2018).

27. Shu, R. *et al.* APP intracellular domain acts as a transcriptional regulator of miR-663 suppressing neuronal differentiation. *Cell Death Dis.* **6**, e1651 (2015).

28. Kenigsbuch, M. *et al.* A shared disease-associated oligodendrocyte signature among multiple CNS pathologies. *Nat. Neurosci.* **25**, 876–886 (2022).

29. Costa, J. *et al.* Cerebrospinal Fluid Chitinases as Biomarkers for Amyotrophic Lateral Sclerosis. *Diagn. Basel Switz.* **11**, 1210 (2021).

30. Yoshida, T. *et al.* Identification and characterization of a novel chemotype MEK inhibitor able to alter the phosphorylation state of MEK1/2. *Oncotarget* **3**, 1533–1545 (2012).

31. Dong, X., Wang, Y. & Qin, Z. Molecular mechanisms of excitotoxicity and their relevance to pathogenesis of neurodegenerative diseases. *Acta Pharmacol. Sin.* **30**, 379–387 (2009).

32. Gurney, M. E. *et al.* Motor neuron degeneration in mice that express a human Cu,Zn superoxide dismutase mutation. *Science* **264**, 1772–1775 (1994).

33. Chun, Y. S. *et al.* MEK1/2 inhibition rescues neurodegeneration by TFEB-mediated activation of autophagic lysosomal function in a model of Alzheimer's Disease. *Mol. Psychiatry* **27**, 4770–4780 (2022).

34. Gal, J., Ström, A.-L., Kilty, R., Zhang, F. & Zhu, H. p62 accumulates and enhances aggregate formation in model systems of familial amyotrophic lateral sclerosis. *J. Biol. Chem.* **282**, 11068–11077 (2007).

35. Benatar, M., Wuu, J. & Turner, M. R. Neurofilament light chain in drug development for amyotrophic lateral sclerosis: a critical appraisal. *Brain J. Neurol.* awac394 (2022) doi:10.1093/brain/awac394.

36. Santiago, J. A., Quinn, J. P. & Potashkin, J. A. Network Analysis Identifies Sex-Specific Gene Expression Changes in Blood of Amyotrophic Lateral Sclerosis Patients. *Int. J. Mol. Sci.* **22**, 7150 (2021).

37. Murdock, B. J., Goutman, S. A., Boss, J., Kim, S. & Feldman, E. L. Amyotrophic Lateral Sclerosis Survival Associates With Neutrophils in a Sex-specific Manner. *Neurol. Neuroimmunol. Neuroinflammation* **8**, e953 (2021).

38. Günther, R. *et al.* The rho kinase inhibitor Y-27632 improves motor performance in male SOD1(G93A) mice. *Front. Neurosci.* **8**, 304 (2014).

39. Torres, P. *et al.* Gender-Specific Beneficial Effects of Docosahexaenoic Acid Dietary Supplementation in G93A-SOD1 Amyotrophic Lateral Sclerosis Mice. *Neurother. J. Am. Soc. Exp. Neurother.* **17**, 269–281 (2020).

40. Food and Drug Administration; U.S. Department of Health and Human Services. Amyotrophic Lateral Sclerosis: Developing Drugs for Treatment Guidance for Industry. (2019).

41. Tahedl, M. *et al.* Propagation patterns in motor neuron diseases: Individual and
phenotype-associated disease-burden trajectories across the UMN-LMN spectrum of
MNDs. *Neurobiol. Aging* **109**, 78–87 (2022).

42. Sahana, T. G. & Zhang, K. Mitogen-Activated Protein Kinase Pathway in Amyotrophic
Lateral Sclerosis. *Biomedicines* **9**, 969 (2021).

43. Ayala, V. *et al.* Cell stress induces TDP-43 pathological changes associated with ERK1/2
dysfunction: implications in ALS. *Acta Neuropathol. (Berl.)* **122**, 259–270 (2011).

44. Pérez-Cabello, J. A. *et al.* MAPK/MAK/MRK overlapping kinase (MOK) controls microglial
inflammatory/type-I IFN responses via Brd4 and is involved in ALS. *Proc. Natl. Acad. Sci.*
120, e2302143120 (2023).

45. Bonifacino, T. *et al.* Nearly 30 Years of Animal Models to Study Amyotrophic Lateral
Sclerosis: A Historical Overview and Future Perspectives. *Int. J. Mol. Sci.* **22**, 12236
(2021).

46. Gordon, D. *et al.* Single-copy expression of an amyotrophic lateral sclerosis-linked TDP-
43 mutation (M337V) in BAC transgenic mice leads to altered stress granule dynamics
and progressive motor dysfunction. *Neurobiol. Dis.* **121**, 148–162 (2019).

47. LaClair, K. D. *et al.* Congenic expression of poly-GA but not poly-PR in mice triggers
selective neuron loss and interferon responses found in C9orf72 ALS. *Acta Neuropathol.*
(Berl.) **140**, 121–142 (2020).

48. Mitchell, J. C. *et al.* Overexpression of human wild-type FUS causes progressive motor
neuron degeneration in an age- and dose-dependent fashion. *Acta Neuropathol. (Berl.)*
125, 273–288 (2013).

49. Ewels, P. A. *et al.* The nf-core framework for community-curated bioinformatics pipelines.
Nat. Biotechnol. **38**, 276–278 (2020).

50. Andrews, S. FastQC: A Quality Control Tool for High Throughput Sequence Data. (2010).

51. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and
bias-aware quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).

52. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
Innov. Camb. Mass **2**, 100141 (2021).

53. Cox, J. & Mann, M. MaxQuant enables high peptide identification rates, individualized
p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat. Biotechnol.*
26, 1367–1372 (2008).

54. Cox, J. *et al.* Andromeda: A Peptide Search Engine Integrated into the MaxQuant
Environment. *J. Proteome Res.* **10**, 1794–1805 (2011).

55. Stekhoven, D. J. & Bühlmann, P. MissForest--non-parametric missing value imputation for
mixed-type data. *Bioinforma. Oxf. Engl.* **28**, 112–118 (2012).

56. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Res.* **43**, e47–e47 (2015).

57. Tyanova, S. *et al.* The Perseus computational platform for comprehensive analysis of
(prote)omics data. *Nat. Methods* **13**, 731–740 (2016).

58. Griffiths-Jones, S., Grocock, R. J., van Dongen, S., Bateman, A. & Enright, A. J. miRBase:
microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res.* **34**, D140-144
(2006).

59. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25 (2009).

60. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

61. Chen, Y. & Wang, X. miRDB: an online database for prediction of functional microRNA
targets. *Nucleic Acids Res.* **48**, D127–D131 (2020).

62. Huang, H.-Y. *et al.* miRTarBase update 2022: an informative resource for experimentally
validated miRNA–target interactions. *Nucleic Acids Res.* **50**, D222–D230 (2022).
63. Liu, W. & Wang, X. Prediction of functional microRNA targets by integrative modeling of
microRNA binding and target expression data. *Genome Biol.* **20**, 18 (2019).
64. Bhattacherjee, A. *et al.* Cell type-specific transcriptional programs in mouse prefrontal
cortex during adolescence and addiction. *Nat. Commun.* **10**, 4169 (2019).
65. Mathys, H. *et al.* Single-cell transcriptomic analysis of Alzheimer's disease. *Nature* **570**,
332–337 (2019).
66. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression
data analysis. *Genome Biol.* **19**, 15 (2018).
67. Szklarczyk, D. *et al.* STRING v11: protein-protein association networks with increased
coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic*
Acids Res. **47**, D607–D613 (2019).
68. Szklarczyk, D. *et al.* The STRING database in 2021: customizable protein–protein
networks, and functional characterization of user-uploaded gene/measurement sets.
Nucleic Acids Res. **49**, D605–D612 (2021).
69. Kuprieiev, R. *et al.* DVC: Data Version Control - Git for Data & Models. (2022)
doi:10.5281/zenodo.7083235.
70. Zhang, J.-N. *et al.* Calpain-mediated cleavage of collapsin response mediator protein-2
drives acute axonal degeneration. *Sci. Rep.* **6**, 37050 (2016).
71. Filareti, M. *et al.* Decreased Levels of Foldase and Chaperone Proteins Are Associated
with an Early-Onset Amyotrophic Lateral Sclerosis. *Front. Mol. Neurosci.* **10**, 99 (2017).
72. Pasetto, L. *et al.* Defective cyclophilin A induces TDP-43 proteinopathy: implications for
amyotrophic lateral sclerosis and frontotemporal dementia. *Brain J. Neurol.* **144**, 3710–
3726 (2021).
73. Luotti, S. *et al.* Diagnostic and prognostic values of PBMC proteins in amyotrophic lateral
sclerosis. *Neurobiol. Dis.* **139**, 104815 (2020).

1154
1155
1156
1157

Acknowledgements
We thank all the members of the Lingor, Bonn, Carapito, Schlapbach, Pasterkamp, and
Bonnetto laboratories for their feedback on the manuscript. This work was performed by the
research consortium "Multi-omic analysis of axono-synaptic degeneration in motoneuron
disease (MAXOMOD)" funded in the scope of the E-Rare Joint Transnational Call for
Proposals 2018 "Transnational research projects on hypothesis-driven use of multi-omic
integrated approaches for discovery of disease causes and/or functional validation in the
context of rare diseases." Consortium members: S. B., J. P., V. B., C. C., R. S., and M. K.,
coordinated by P. L.
L. C. G., M. P., and P. L. were supported by the Bundesministerium für Bildung und Forschung
(01GM1917A). P. L. and D. E. were further supported by the Munich Cluster for Systems
Neurology (SyNergy). S. H. was supported by SFB1192 B8; S. O. by SFB1286 SP02 and
KFO296 P8; R. K. by FOR5068 P9; F. H. by the M3I excellence initiative and a UKE
postdoctoral stipend; C. H. by KFO306 P11; and S. B. by SFB1286 SP02, SFB1192 PB8, and
PC3. J. P. was supported by Stichting ALS Nederland (TOTALS). D. E. received funding from
the European Union's Horizon Europe research and innovation program under grant
agreement No 101057649 and the Association for Frontotemporal Degeneration (AFTD).
The funders had no role in the study design, data collection and analysis, decision to publish,
or manuscript preparation.

1177
Author contributions
P. L. and L. C. G. conceived the project, and S. H., S. O., and S. B. conceptualized the
computational analysis. P. L., L. C. G., J. K., and A. D. designed the sample collection
methodology, reviewed the sample and data quality, and coordinated the acquisition of human
tissue samples from the brain banks as well as pathological, genetic, and clinical information.
L. C. G, M. P., and Q. Z. processed mouse and human brain samples for multiomics
experiments with the infrastructure provided by V. B., C. C., D. E., and P. L.
C. D., M. S., T. B. H., I. C., and M. D. analyzed the genetic data.
S. H., S. O., R. K., and F. H. were responsible for the computational data analysis, and
conceived and planned the statistical analyses. L. C. G, S. H., F. H., R. K., P. L., M. P., and L.
P. contributed to figure generation and assembly.
M. P, L. P., L. C. G., S. S., and S. C. conducted the validation experiments with conceptual
inputs from P. L. and V. B.
L. C. G., S. H., S. B., and P. L. wrote the manuscript with input from all co-authors.
1192
Competing interests
1194   The authors declare no competing interests.
1195
Additional information
1197   Supplementary information
1198
1199   Correspondence and requests for materials should be addressed to Paul Lingor.
1200

Figure Legends

1202

Fig. 1 Study overview and the transcriptomic landscape of brains affected by amyotrophic lateral sclerosis

a. Overview of the sample processing workflow. Prefrontal cortex samples were prepared for multiomics experiments from the human cohort (51 ALS/50 CTR samples), as well as from four selected ALS mouse models (C9orf72, FUS, SOD1, and TDP-43; 10 TG/10 CTR animals per group). RNA, DNA, and protein lysates were prepared for mRNA and small RNA sequencing, targeted DNA sequencing, and proteomics/phosphoproteomic (ph-proteomic) experiments. The latter was performed exclusively on animal tissues. **b.** Volcano plot of deregulated genes in human samples. x-axis: log2(fold change); y-axis: -log10(adjusted *p*-values) for each gene. Blue and orange circles indicate significant differential expression: left side, decrease (low ALS); right side, increase (high ALS). **c.** Heat map of the activity scores (calculated by Decoupler) of each pathway throughout the ALS human cohort. Pathways were selected based on their increased occurrence across different enrichment analyses. Pathways are indicated on the y-axis and ALS samples on the x-axis (top) along with metadata, sex, and age at death. Only selected pathways are shown. A full heat map of all chosen pathways is shown in Supplementary Figure 16. **d.** Dot plot analysis of enriched Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways using ClusterProfiler in male and female samples. The dot plot displays significantly enriched pathways in males and females, represented by circles colored by their corresponding adjusted *p*-values (-log10 transformed) on the x-axis and gene count on the y-axis. The size of each circle corresponds to the number of genes annotated in the KEGG gene set. Similar pathways were also identified. Pathway clusters for human males were composed of the following terms: retrograde endocannabinoid signaling, synaptic vesicle cycle, long-term potentiation, glutamatergic synapse (*synapse related terms*), *Staphylococcus aureus*

infection, malaria, complement and coagulation cascades, and cytokine–cytokine receptor interaction (for *immune system activation/infectious diseases*). Pathway clusters for human females were composed of the following terms: Parkinson's disease, Huntington's disease, spinocerebellar ataxia, pathway of neurodegeneration-multiple diseases, and amyotrophic lateral sclerosis (for *neurodegenerative/neurological diseases*); complement and coagulation cascades, viral protein interaction with cytokine and cytokine receptor, *Staphylococcus aureus* infection, Epstein–Barr virus infection, Influenza A cytokine–cytokine receptor interaction (for *immune system activation/infectious diseases*). **e.** Heat map showing Pearson's correlation between the weighted gene co-expression network analysis (WGCNA) modules and each sample group. Each cluster of the ALS samples and controls is shown on the y-axis (substratified by sex). Representative WGCNA modules were selected for the identified ALS clusters (turquoise (1) and yellow (2) for clusters C1 and C2, and tan (3) and light cyan (4) for clusters C3 and C4) and explored further by enrichment analyses.
f. Enrichment results (over-representation analysis) for each selected WGCNA module (top five terms based on significance).

Fig. 2 Overview of the results for the selected ALS mouse models, correlation with human results, and evidence for mitogen-activated protein kinase deregulation in ALS subclusters
a. Volcano plot of deregulated genes in the four transgenic mouse models, represented by dots colored according to sex. The x-axis shows the log2(fold change) between transgenic and wild-type mice, whereas the y-axis shows the -log10(adjusted *p*-values). Orange and blue dots represent DEGs in females and males, respectively.
b. Dot plot analysis of enriched gene ontology (GO) biological processes (BP) using Genetonics in male and female samples for ALS mouse models. The dot plot displays enriched pathways in males and females, represented by circles colored by their corresponding adjusted *p*-values on the x-axis and gene count on the y-axis. The size of each circle corresponds to the number of genes annotated in the GO BP gene sets.
c. Heat map showing changes in cell type proportions across ALS mouse models and human clusters. The indicated changes are the relative differences between the median cell type composition of the ALS cluster/TG and the median cell type composition in the corresponding CTR samples. Comparisons were grouped according to their similarity using hierarchical clustering with Euclidean distance.
d. Heat map showing the pairwise Pearson correlation of relative changes in median cell type proportions in ALS mouse models and human clusters. Comparisons were grouped according to similarity using hierarchical clustering, as depicted by the dendrograms.
e. Summarized visualization of the MAPK pathway (Kyoto Encyclopedia of Genes and Genomes: hsa04010) colored by log2(fold change) of transcriptomics clusters C1 (orange) and C3 (purple) compared with healthy controls.

Fig. 3 Differential alternative splicing and miRNAomic profiling
a,b. Differential alternative splicing (DAS) analysis. The plot displays the results for human male and female samples for various splice events, i.e., alternative exon (AE), skipped exon (SE), alternative 5'-splice site (A5), alternative 3'-splice site (A3), alternative last exon (AL), and retained intron (RI) events. Each event is represented by a separate bar, the height of which represents the fraction of significant events in ALS vs. CTR. Blue bars represent events with significant differential splicing in males, whereas orange bars represent events with significant differential splicing in females.
c. Venn diagram showing the overlap of DAS genes in four transgenic mouse models (TDP-43, FUS, C9orf72, and SOD1) for the ALS vs. CTR comparison. Each circle represents the

number of genes with significantly differential splicing in each model. Overlapping regions between circles represent the number of genes with significantly differential splicing in more than one model.

d. Enrichment of selected themes for DAS genes in four transgenic mouse models and humans, male and female samples, and percentage of genes in each functional term. The plot displays two sets of information: the enrichment of DAS genes in different samples and the percentage of genes in each functional term. The y-axis represents the percentage of genes in each term, whereas the x-axis represents different functional terms. The bars are colored according to sex, mouse models, and human samples.

e,f. Volcano plots showing differential expression of mature miRNAs (e) and hairpin miRNAs (f) in male and female human samples. The plots display the log2(fold change) (x-axis) and the -log10(adjusted p-value) (y-axis) for each miRNA between male and female samples. The miRNAs are represented as dots in the plot, and the color of the dots indicates the significance of differential expression (blue for males and orange for females).

g,h. The heat maps display the log2(fold change) of the top miRNAs in female and male samples from the mouse and human datasets. The miRNAs were selected based on the ordered absolute log2(fold change) per dataset. Red indicates miRNA upregulation, whereas blue indicates miRNA downregulation. Statistical significance of the differential expression of miRNAs was inferred using DESeq2 and is indicated by asterisks. **p < 0.01 and *p < 0.05 represents statistical significance at a high and moderate level, respectively. The top five miRNAs significantly differentially expressed in at least two mouse models are shown.

i. Protein–protein interaction network of the target genes of miRNA451a. MAPK1, AKT1, IKBKB, MYC, BCL2, and IL6 are important molecular hubs.

Fig. 4 Proteomics overview and multiomics factor integration

a. Volcano plot of deregulated proteins in human samples. x-axis: log2(fold change); y-axis: -log10(adjusted p-value) for each protein. Blue and orange circles indicate significant differential changes: left side, decrease (low ALS); right side, increase (high ALS).

b,c. Revigo-based summary of proteomic gene set enrichment results for human samples (left and right females). The plot summarizes the functional similarity of the proteins by reducing the redundant gene ontology (GO) biological processes (BP) terms and clustering the remaining non-redundant terms. The plot is divided into several areas, each of which represents a cluster of similar GO BP terms. The size of the circles represents the number of genes in the GO BP terms, and the color represents significance.

d,e. Plots displaying the results of GO enrichment analysis of the proteins identified in the proteomic data of the four mouse models (TDP-43, FUS, C9orf7, and SOD1) separated by sex. GO terms were summarized by clustering pathways in semantic space based on their descriptions using the GO-Figure clustering tool.

f. Pearson correlation of log2(fold change) of proteins found to be significantly deregulated (p < 0.1) in any mouse model or human sample. Comparisons were grouped according to similarity using hierarchical clustering, as depicted by the dendrograms.

g. Multi-Omics Factor Analysis (MOFA) was performed to integrate and visualize transcriptomic, proteomic, and mature and hairpin miRNA data in male and female samples. Uniform Manifold Approximation and Projection (UMAP) representation showing the distribution of male (blue) and female (orange) human samples.

h. Variance decomposition plot showing the contribution of each factor in the MOFA to the overall variance of the integrated dataset separated by sex. The plot displays the variance explained by transcriptomics, proteomics, and mature and hairpin miRNAs in human male and female samples. The y-axis represents the different omics, and the x-axis represents the

different factors in the analysis (left: male, right: female). Blue indicates the proportion of
explained variance, and red indicates Pearson's correlation with the condition.
i. MOFA weight plot for human male miRNA data (factor 3) displayed the contribution of each
gene to the MOFA model. Each line represents a gene, and the x-axis shows the MOFA weight
of that gene, reflecting its importance to the factor.
j. MOFA weight plot for proteomic human female data (factor 12) displays the contribution of
each protein to the MOFA model. Each line represents a protein, and the x-axis shows the
MOFA weight of that protein, reflecting its importance to the factor.
k. Protein–protein interaction network of genes in factor 12 females. MAPK1 is an important
molecular hub protein that interacts with PEA15, PRRT2, MEK2, DUSP6, HSPA4, and
HSP90AA1. Proteins of interest are highlighted (bold and dark purple).
1337
Fig. 5 Validation experiments
a,b. Effects of different concentrations of trametinib on apoptosis in glutamate (5 mM)- and
non-glutamate-treated cells analyzed by immunostaining. Representative photomicrographs
for two of the analyzed conditions (control [vehicle] and 200 nM trametinib). Scale bar: 40 μm.
c,d. Quantification plots showing the effects of treatment with trametinib on cell survival
(caspase 3 staining) (**c**) and neurite outgrowth (**d**) for glutamate- or vehicle-treated (control)
cells, for all analyzed conditions (control [vehicle]; 2 nM trametinib; 20 nM trametinib; 200 nM
trametinib). Data are presented as the mean ± standard error of the mean (SEM) of at least
five different cultures and were tested using one-way analysis of variance.
e,f. Western blot analysis and quantification of trametinib effects on pErk1/2 with and without
glutamate treatment. Data represent the mean ± SEM of at least three different cultures and
were tested by one-way analysis of variance, $p < 0.05$.
g,h. Diffuse pMEK2 immunostaining in the lumbar spinal cord of non-transgenic (**g**) and
SOD1^{G93A} mice (**h**) at 19 weeks of age. In the ventral horns, pMEK2 staining was mainly
observed in motor neurons. Scale bar: 50 μm.
i. Western blot analysis of pMEK2 in the lumbar spinal cord of SOD^{1G93A} and non-transgenic
mice at 9, 14, and 19 weeks of age. Data are mean ± SEM (n = 6–4 in each experimental
group) and are expressed as relative immunoreactivity (RI). *$p < 0.05$, pairwise Tukey's test
after one-way analysis of variance. Dot blot analysis of pERK1/2 (**j**), p62 (**k**), insoluble SOD1
(**l**), and ubiquitin (**m**) in the spinal cord of SOD1^{G93A} female and male mice treated with
trametinib or vehicle, at 16 weeks of age. Data represent mean ± SEM (n = 3/5 in each
experimental group) and are expressed as relative immunoreactivity (RI). *$p < 0.05$, Student's
t-test. Plasma neurofilament light chain levels were analyzed in female and male (**n**) SOD1^{G93A}
mice treated with or without trametinib at 16 weeks of age. Data are presented as the mean ±
SEM (n = 3/5 in each experimental group). *$p < 0.05$, Student's *t*-test.
1363

Table 1: Summary of the demographics of the human cohort
Continuous variables are presented as medians and ranges. Categorical variables were
compared using Fisher's exact test (for single groups) or the chi-square test (for multiple
groups); continuous variables were compared using the Mann–Whitney test. The p-values
shown are uncorrected for multiple testing; p-values less than 0.05 are indicated in bold.
Netherlands BB, Netherlands Brain Bank; Oxford BB, Oxford Brain Bank; PD and MS TB,
Parkinson's and Multiple Sclerosis UK Tissue/Brain Banks; London ND BB, King's College
London Brain Bank.

	Control	ALS	*p*-value
Donors	50	51	
Age at death, years	75 (43–94)	67 (44–83)	**0.0032**
Sex female	56.0%	31.4%	**0.0163**
Postmortem interval, h	22 (5–95)	23 (2.5–98)	0.4324
Unknown	8.0%	5.9%	
Onset			
Bulbar	-	17.6%	
Spinal	-	37.3%	
Upper limb	-	31.6%	
Lower limb	-	63.1%	
Limb unknown	-	5.3%	
Thoracic	-	2.0%	
Unknown	-	43.1%	
Disease duration, years	-	3 (1–28)	
Unknown	-	39.2%	
Brain bank origin			**< 0.0001**
Netherlands BB	18.0%	17.6%	
Oxford BB	20.0%	27.5%	
ICL MS & PD TB	38.0%	0.0%	
London ND BB	24.0%	54.9%	

1372 ALS, amyotrophic lateral sclerosis.

a

−Log₁₀ *P* (y-axis), log2FC (x-axis)

↑132
↑211
↓119
↓168

significant for:
males (blue)
females (orange)

Labels: TGM2, CO6A3, ANXA2, STAT3, LEG3, ANXA2, TAGL, TAGL, CO6A3, DOC2A, SYUA

b human females

cytoskeleton / extracellular matrix
catalytic activity
development
synapse
transmembrane transport
ERK1/ERK2 signaling
cytokine / inflammatory response
lipid metabolism

c human males

synapse
metabolism
immune response
cytoskeleton / extracellular matrix

d female

C9orf72 (blue)
FUS (orange)
human (green)
SOD1 (red)

1. ATP synthesis
2. rRNA processing
3. generation of neurons
4. neuron development
5. mitochondrial ATP synth.
6. neuron projection dev.
7. neuron differentiation
8. homeostatic process
9. cell-cell adhesion
10. reg. of transport
11. nervous system dev.
12. defense response
13. brain development
14. ribosome biogenesis
15. nervous system process
16. reg.of membrane potential
17. synaptic vesicle exocytosis
18. reg.of protein localization
19. cell projection
20. aerobic respiration

e male

1. translation
2. neurotransmitter secretion
3. B cell activation
4. defense response
5. ATP synthesis
6. humoral immune response
7. cellular detoxification
8. nervous system dev.
9. pmembrane invagination
10. postsynaptic membrane
11. response to toxin
12. cellular respiration
13. vesicle-med.transport
14. protein transport
15. synaptic plasticity
16. electron transport
17. reg.endocytosis
18. ribosomal biogenesis
19. vesicle localization
20. metabolic process

f

FUS fem.
SOD1 fem.
SOD1 male
C1
C2
C3
C4
TDP43 fem.
TDP43 male
FUS male
C9orf72 fem.
C9orf72 male

(columns): FUS fem., SOD1 fem., SOD1 male, C1, C2, C3, C4, TDP43 fem., TDP43 male, FUS male, C9orf72 fem., C9orf72 male

g

human males (blue)
human females (orange)

UMAP2 (y-axis), UMAP1 (x-axis)

h

human males

Condition *r²*
proteomics
miRNAhairpin
miRNAmature
transcriptomics

factors: 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15

human females

Condition *r²*
proteomics
miRNAhairpin
miRNAmature
transcriptomics

factors: 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15

correlation
males: .05 .10 .15
females: .02 .04 .06 .08 .10
factors: 10 20 30

i male - factor 3

RAB3C −
MIR3648-2 +
BHLHB9 −
TP53INP2 +
SNORA73B +
RN7SL2 +
BCL2L2-PABPN1 −
GOLGA6L17P +
PAQR6 +
OPRM1 −

0.0 0.3 ... 0.9

j female - factor 12

NFH +
STX1A −
PRIO −
PEA15 −
NT5D1 +
IPYR2 +
PRRT2 −
KCC4 −
NAC2 −
ANK1 +

0.0 0.3 ... 0.9

k

synaptic compartment (pink)
MAPK pathway (dark purple)
other proteins from factor 12 and closely interacting proteins (light)

Labels: STX1A, SNAP23, VAMP8, CPLX1, STX3, VAMP3, SYT1, SNAP25, VAMP2, STX4, STX11, PEA15, DUSP6, PRRT2, STXBP1, NEFH, MAPK1, MAP2K2, HSP90AA1, ITGAM, ANK1, OBSCN, PRNP, ICAM1, STIP1, HSPA4

a 0mM glutamate

control — 200nM trametinib

MAP2 caspase / MAP2 caspase

merge / merge

b 5mM glutamate

control — 200nM trametinib

MAP2 caspase / MAP2 caspase

merge / merge

c

0mM glutamate | 5mM glutamate

P = 0.0237
P < 0.0001
P = 0.0027

caspase 3 pos. cells (relative to CTR)

0 2 20 200 | 0 2 20 200

trametinib concentration (nM)

d

0mM glutamate | 5mM glutamate

p = 0.0051
p = 0.0003
p < 0.0001

Neurite lenght/cell (relative to CTR)

0 2 20 200 | 0 2 20 200

trametinib concentration (nM)

e

0mM glutamate
phospho ERK1/2
lamin b
0 2 20 200
trametinib concentration (nM)

5mM glutamate
phospho ERK1/2
lamin b
0 2 20 200
trametinib concentration (nM)

f

0mM glutamate | 5mM glutamate

p = 0.0071
p = 0.0011
p < 0.0001

phospho-Erk1/2 / lamin b band intensity ratio

0 2 20 200 | 0 2 20 200

trametinib concentration (nM)

g

pMEK2
Nissl

h

i pMEK2 (RI)

females | males

***
**
*
**

Ntg 9 weeks 14 weeks 19 weeks | Ntg 9 weeks 14 weeks 19 weeks

j pERK1/2 (RI)

females | males

vehicle trametinib | vehicle trametinib

k p62 (RI)

females | males

*

vehicle trametinib | vehicle trametinib

l insoluble SOD1 (RI)

females | males

*

vehicle trametinib | vehicle trametinib

m insoluble ubiquitin (RI)

females | males

*

vehicle trametinib | vehicle trametinib

n NfL (pg/ml)

females | males

*

vehicle trametinib | vehicle trametinib