1 # Different coexistence patterns between apex carnivores and mesocarnivores
2 # based on temporal, spatial, and dietary niche partitioning analysis in Qilian
3 # Mountain National Park, China
4

5 Wei Cong[a], Jia Li[b], Charlotte Hacker[c], Ye Li[a], Yu Zhang[a], Lixiao Jin[a], Yi Zhang[a], Diqiang Li[a], Yadong
6 Xue[a]*, Yuguang Zhang[a]*
7
8 a. Ecology and Nature Conservation Institute, Chinese Academy of Forestry, Key Laboratory of
9 Biodiversity Conservation, State Forestry and Grassland Administration, Beijing 100091, China
10 b. Institute of Ecological Conservation and Restoration, Chinese Academy of Forestry, Beijing, 100091,
11 China
12 c. Department of Biological Sciences, Duquesne University, Pittsburgh, PA, 15282, USA
13
Wei Cong, 779237048@qq.com
Jia Li, lijia2530@126.com
Charlotte Hacker, charlotte.hacker00@gmail.com
Ye Li, liye19821012@sina.com
Yu Zhang: yuzhang420@126.com
Lixiao Jin: 2507476267@qq.com
Yi Zhang: 527808577@qq.com
Diqiang Li: lidq@caf.ac.cn
*Correspondence:
Yadong Xue, xueyadong334@163.com;
Yuguang Zhang, yugzhang@sina.com.cn
25

Abstract

Carnivores play key roles in maintaining ecosystem structure and function as well as ecological processes.

Understanding how sympatric species coexist in natural ecosystems is a central research topic in

community ecology and biodiversity conservation. In this study, we explored intra- and interspecific

niche partitioning along spatial, temporal, and dietary niche partitioning between apex carnivores (wolf

Canis lupus, snow leopard *Panthera uncia*, Eurasian lynx *Lynx lynx*) and mesocarnivores (Pallas's cat

Otocolobus manul, red fox *Vulpes vulpes*, Tibetan fox *Vulpes ferrilata*) in Qilian Mountain National Park,

China using camera trapping data and DNA metabarcoding sequencing data. Our study showed that apex

carnivore species had more overlap temporally (coefficients of interspecific overlap ranging from 0.661

to 0.900) or trophically (Pianka's index ranging from 0.458 to 0.892), mesocarnivore species had high

dietary overlap with each other (Pianka's index ranging from 0.945 to 0.997), and apex carnivore and

mesocarnivore species had high temporal overlap (coefficients of interspecific overlap ranging from

0.497 to 0.855). Large dietary overlap was observed between wolf and snow leopard (Pianka's

index=0.892) and Pallas's cat and Tibetan fox (Pianka's index=0.997), suggesting the potential for

increased resource competition for these species pairs. We concluded that spatial niche partitioning is

likely to key driver in facilitating the coexistence of apex carnivore species, while spatial and temporal

niche partitioning likely facilitate the coexistence of mesocarnivore species, and spatial and dietary niche

partitioning facilitate the coexistence between apex and mesocarnivore species. Our findings consider

partitioning across temporal, spatial and dietary dimensions while examining diverse coexistence patterns

of carnivore species in Qilian Mountain National Park, China. These findings will contribute

substantially to current understanding of carnivore guilds and effective conservation management in

fragile alpine ecosystems.

Keywords: Species coexistence mechanism, niche partitioning, species occupancy model, camera

trapping, DNA metabarcoding technology

Introduction

Carnivores play key roles in maintaining ecosystem structure and function, as well as ecological processes (Ripple et al., 2014). Apex carnivores are often classically specialized hunters occupying top trophic positions that work to suppress the number of herbivores and mesocarnivores through predation, competition and trophic cascades (Ripple et al., 2014; Newsome et al., 2017; Ritchie and Johnson, 2009). Understanding how sympatric carnivore species coexist in natural ecosystems is a central focus in community ecology and biodiversity conservation (Chu et al., 2017). Human activities and climate change are causing large-scale habitat loss and fragmentation, posing significant challenges to carnivore interspecies relationships (Li et al., 2021; Manlick and Pauli, 2020; Ripple et al., 2014). Such relationships are becoming increasingly intricate, leading to greater research efforts aimed at determining the strategies that facilitate coexistence despite intense competition (Kuijper et al., 2016; Smith et al., 2017).

The competitive exclusion principle dictates that species with similar ecological requirements are unable to successfully coexist (Hardin, 1960; Gause, 1934). Thus, carnivores within a guild occupy different ecological niches based on a combination of three niche dimensions, i.e. spatial, temporal, and trophic (Schoener, 1974). Spatially, carnivore species within the same geographic area exhibit distinct distributions that minimize overlap in resource use and competition. For example, carnivores can partition habitats based on habitat feature preferences and availability of prey (De Satgé et al., 2017; Garrote and Pérez De Ayala, 2019; Gołdyn et al., 2003; Strampelli et al., 2023). Temporally, differences in seasonal or daily activity patterns among sympatric carnivores can reduce competitive interactions and facilitate coexistence. For example, carnivores can exhibit temporal segregation in their foraging behaviors, such as diurnal versus nocturnal activity, to avoid direct competition (Finnegan et al., 2021; Nasanbat et al., 2021; Searle et al., 2021). Trophically, carnivore species can diversify their diets to exploit different prey species or sizes, thereby reducing competition for food resources. For example, carnivores can exhibit dietary specialization to optimize their foraging efficiency and minimize competitive pressures (Steinmetz et al., 2021). Currently, research on the niche partitioning of sympatric carnivores primarily focuses on one or two dimensions, with little attention paid to all three (Alexander et al., 2016; Karanth et al., 2017; Li et al., 2022a; Santos et al., 2019; Shao et al., 2021; Strampelli et al., 2023; Tsunoda et al., 2020; Vilella et al., 2020). This limitation restricts the multi-scale understanding of coexistence mechanisms among carnivores.

In recent years, camera trapping and DNA metabarcoding technology has been widely used in wildlife monitoring and research. Camera trapping enables the monitoring of elusive species in remote areas (Boitani, 2016; Palencia et al., 2022), and has become a helpful tool for accumulating large amounts of time-recorded data that can provide detailed information surrounding animal behavior and activity patterns, which is widely used to investigate species interspecies relationships (Frey et al., 2017). For example, Li et al. (2019) used camera trap data to conclude that temporal segregation is a key mechanism for promoting the coexistence of tigers (*Panthera tigris*) and leopards (*P. pardus*). DNA metabarcoding provides a noninvasive molecular tool that is more accurate than traditional dietary analysis methods, such as microhistology (Deiner et al., 2017; Newmaster et al., 2013). DNA metabarcording technology has been applied to many studies and has offered profound insight into the ecology, conservation and biological monitoring of rare and endangered species (Deagle et al., 2019; Kartzinel et al., 2015). For example, Shao et al. (2021) found that dietary niche partitioning promoted the

coexistence of species in the mountains of southwestern China based on DNA metabarcoding. The

combination of camera trapping and DNA metabarcoding can result in robust data for exploring the

mechanisms of coexistence surrounding carnivore guilds.

The Qilian Mountains constitute a biodiversity hotspot with one of the richest carnivore assemblages

in the world (Di Minin et al., 2016). To understand the competition and coexistence among different

carnivore species in this area, we explored the habitat use, activity patterns, and prey item composition of

sympatric carnivore species comprised of apex carnivores and mesocarnivores across Qilian Mountain

National Park using camera trap data and DNA metabarcoding data. Based on theories surrounding

resource partitioning and niche differentiation, as well as studies on interactions and coexistence among

carnivorous species (Haswell et al., 2018; Linnell and Strand, 2000), we hypothesized that differentiation

along one or more niche axes is beneficial for the coexistence of the carnivorous guild in the Qilian

Mountains. We expected that spatial niche differentiation promotes the coexistence of large carnivores in

the Qilian Mountain region, as they are more likely than small carnivores to spatially avoid interspecific

competition (Davis et al., 2018). Mesocarnivores may coexist either spatially or temporally due to

increased interspecific competition for similar prey (Di Bitetti et al., 2010; Donadio and Buskirk, 2006).

Dietary niche differentiation may be a significant factor for promoting coexistence between large and

mesocarnivore species due to differences in body size (Gómez-Ortiz et al., 2015; Lanszki et al., 2019).

Material and methods

Study sites

The Qilian Mountains laterally span Gansu and Qinghai Provinces in China, located on the

northeastern edge of the Qinghai-Tibetan Plateau (Figure 1). Qilian Mountain National Park covers an

area of approximately 52,000 km^2, with an average elevation of over 3000 m. The area is an alpine

ecosystem with a typical plateau continental climate. The average annual temperature is below -4℃ and

the average annual rainfall is about 400 mm, with habitats mainly consisting of deserts, grassland,

meadows, and wetland (Zheng, 2011). Wildlife present include the wolf (*Canis lupus*), snow leopard (*P.*

uncia), Eurasian lynx (*Lynx lynx*), red fox (*Vulpes vulpes*), Tibetan fox (*V. ferrilata*), Tibetan brown bear

(*Ursus arctos*), Chinese mountain cat (*Felis bieti*), wild yak (*Bos mutus*), blue sheep (*Pseudois nayaur*),

alpine musk-deer (*Moschus chrysogaster*), Tibetan antelope (*Pantholops hodgsonii*), Himalayan marmot

(*Marmota himalayana*), woolly hare *(Lepus oiostolus)*, and plateau pika (*Ochotona curzoniae*), among

others (Ma et al., 2021; Xue et al., 2019).

Camera-trap monitoring and non-invasive sampling

The study area was subdivided into sample squares of 25 km^2 (5×5 km) as a geographical reference

for placing camera survey sites and collecting scat samples (Xue et al., 2019). Species occurrence was

recorded using camera-trap monitoring (Ltl-6210; Shenzhen Ltl Acorn Electronics Co. Ltd). Two camera

traps were placed in each square with a distance of at least 1 km between them. However, due to

limitations of terrain, landform, road accessibility and other factors, the number of camera trap in some

squares was adjusted in accordance to field conditions. Camera traps were set in areas believed to be

important to and heavily used by wildlife, such as the bottoms of cliffs, sides of boulders, valleys and

ridges along movement corridors. Taking into account the fact that mammalian communities are sensitive

to seasonality, we used camera traps to monitor animals with an extensive survey effort from December

2016 to February 2022, covering the activity of animal species in different seasons, which can reflect the

overall distribution of carnivores. We placed a total of 280 infrared cameras at the study site, set them to be active for 4 to 6 months, and considered possible relocation to another position based on animal detection in an effort to improve estimates of the occupancy and detection rates for both common and rare species (Figure 1) (Kays et al., 2020). The camera trap was set to record the time and date on a 24 hr clock when triggered, and to record a 15s video and 1 photo with an interval of 2 minutes between any two consecutive triggers. The sum of camera trap effective days was defined by the total amount of trapping effort during the sampling period, which was calculated from the time the camera was placed in operation to the time the last video or photograph was taken. We visited each camera 2 to 3 times a year to download photos and check batteries.

Non-invasive sampling of scats believed to be of carnivore origin were collected within camera trapping areas. A small portion of scat (approximately 1/3) was broken off and stored in a 15 ml centrifuge tubes with silica desiccant covered by clean filter paper to separate the desiccant from the scat (Janecka et al., 2008). Gloves were replaced between sampling to avoid cross-contamination. Sampling place, date, and sample number were labeled on the tube; GPS coordinates, elevation, and nearby landscape features were recorded on the sample collection sheet (Hacker et al., 2021). A total 480 scat samples were collected from April 2019 to June 2021 (Figure 1).

Data Analysis

Spatial analysis

To investigate the spatial distribution of carnivores, as well as the influence of environmental factors on the site occupancy of species in the study area, we performed single-season, single-species occupancy models to estimate carnivores' occupancy (ψ) and detection (*Pr*) probability (Li et al., 2022b; MacKenzie, 2018; Moreno-Sosa et al., 2022). To ensure capture independence, only photo or video records at intervals of 30 min were was included in the data analysis (Li et al., 2020). We created a matrix recording whether each carnivore species was detected (1) or not (0) across several 30-day intervals (that is 0-30, 31-60, 61-90, 91-120, 121-150, >150 days) for each camera location. Based on the previous studies of habitat use of carnivores (Greenspan and Giordano, 2021; Alexander et al., 2016; Gorczynski et al., 2022), we selected terrain, vegetation, biological factors and disturbance to construct the model. Terrain is a fundamental element of wildlife habitat and closely linked to other environmental factors (Chen et al., 2024). Terrain variables include elevation (ele) and roughness index (rix). Vegetation variables include normalized difference vegetation index (ndvi), and provide information on the level of habitat concealment. Biological variables include prey abundance (the number of independent photos of their preferred prey based on dietary analysis in this study, wolf and snow leopard: artiodactyla including livestock; Eurasian lynx and Pallas's cat: lagomorpha; red fox and Tibetan fox: lagomorpha and rodentia) and reflect habitat preference and distribution patterns of carnivores. Disturbance variables include distance to roads (disrd) and human disturbances (hdis, the number of independent photos of herdsman and livestock) and can provide insight into the habitat selection and behavior patterns of carnivores. In addition, we used elevation, human disturbance and prey as covariates that affect detection rate. Road data were obtained from Open Street Map (OSM, https://www.openstreetmap.org). Others environmental data were obtained from the Resource and Environment Science and Data Center (https://www.resdc.cn). We fitted all possible combinations of covariates and used Akaike's information criterion (AIC) to rank candidate models, and selected ΔAIC≤2 model as the optimal model. If more than one optimal model

resulted, then covariate estimates were obtained by using the equal-weight average.

Carnivore co-occurrence was evaluated using the Sørensen similarity index (value = 0, indicating

maximum segregation and value = 1, indicating maximum co-occurrence) based on binary presence

absence data within the 5 km × 5 km camera trap grid (Torretta et al., 2021; Sorensen, 1948). Spatial

analyses were performed using ArcGIS 10.8 (ESRI Inc.), the "vegan" packages (Oksanen et al., 2019)

and "unmarked" package (Fiske and Chandler, 2011) for R studio (version 1.1.463).

Temporal analysis

Estimates of the coefficient of overlap (Δ) for activity patterns were estimated using the

non-parametric kernel density method and applying time data obtained by the camera traps. Because the

smallest sample had more than 50 records, we used the Δ_4 estimator for pairwise comparisons between

carnivore activity patterns, and used a smooth bootstrap scheme to generate 1000 resamples with 95%

confidence intervals to test the reliability of the overlap value (Ridout and Linkie, 2009). Activity pattern

analyses were performed using the "overlap" R package. Values of the Δ_4 estimates were calculated

relative to 1000 simulated null models of randomized overlap data using the 'compareCkern' function in

the "activity" R package to test for differences in daily activity patterns (Ridout and Linkie, 2009;

Rowcliffe et al., 2014).

Species identification and dietary analysis

Host species and diet were identified using metabarcoding. DNA was extracted using the QIAamp

Fast DNA Stool Mini Kit (QIAGEN, Germany) following standard protocols and the MT-RNR1 (12S)

and COX1 (cytochrome c oxidase subunit I) gene segments amplified using 12SV5-F/R primer and

COX1 primers, respectively (Hacker et al., 2021; Riaz et al., 2011). PCR reaction conditions followed the

methods described in Hacker et al. (Hacker et al., 2021). The resulting library was sequenced on an

Illumina NovaSeq platform and 250 bp paired-end reads were generated (Guangdong Magigene

Biotechnology Co., Ltd. Guangzhou, China).

We used CLC Genomics Workbench v12.0 (QIAGEN, Denmark) to determine the host species as

well as the prey consumed by each carnivore by mapping sequence reads to reference sequences of

possible prey downloaded from GenBank and BOLD (Barcode of Life Data Systems) with representative

haplotypes compiled into one .fasta file. Raw reads were required to have at least 98% similarity across at

least 90% of the reference sequence for mapping (Hacker et al., 2021). Species and prey identification

were made based on the reference taxa with the highest number of reads mapped and the fewest

mismatches. Samples in which species could not be identified were analyzed to ensure the reference file

was not incomplete by using the *de nova* assembly tool in CLC, then blasting the resulting contig

sequence with the nucleotide databases in NCBI (https://blast.ncbi.nlm.nih.gov/Blast.cgi). As an

additional precaution, the geographic range of the determined host and prey species was researched using

the IUCN Red List (https://www.iucnredlist.org/) to ensure that it overlapped with the study site. For

complete methods on data parameters and methods used, see Hacker et al. (Hacker et al., 2021).

Dietary data were summarized by the frequency of occurrence of prey species in scats observed. The

"bipartite" R package was used to construct food web networks (Dormann, 2011). Dietary diversity for

each carnivore host species was assessed by calculating richness and Shannon's Index (Shannon and

Weaver, 1949). Interspecific dietary niche overlap was evaluated using Pianka's Index (O_{jk}) (value = 0,

no dietary overlap and value = 1, complete dietary overlap) and 95% confidence intervals were obtained

by bootstrapping with 1000 resamples via the "spaa" R package (Zhang, 2016). Dietary similarity

between any two given carnivore species was assessed by calculating the inversed value of Jaccard's Index

based on binary presence absence data of prey.

Results

Sympatric carnivore identification

Of the 480 scat samples sequenced, those which had no sequencing data, were inconclusive,

consisted of non-target species, or host species with low sample sizes (1 Asian badger (*Meles leucurus*)

and 4 upland buzzard (*Buteo hemilasius*)) were removed (Figure S1). The remaining 404 scat samples

were composed of three apex carnivores (49 wolf, 147 snow leopard, 19 Eurasian lynx) and three

mesocarnivores (63 Pallas's cat, 87 red fox, 39 Tibetan fox).

Spatial distribution difference and overlap

A total of 322 camera trap sites were surveyed after relocating infrared cameras that did not capture

any target carnivore species. A total of 3 cameras were considered to have failed due to loss. We analyzed

data from 319 camera sites and obtained 14,316 independent detections during a total effort of 37,192

effective camera trap days. We recorded wolf in 26 sites, snow leopard in 109 sites, Eurasian lynx in 36

sites, red fox in 92 sites, and Tibetan fox in 34 sites. However, the camera detection rates of Pallas's cat

were too low to analyze species occupancy and daily activity patterns.

According to the findings derived from single-season, single-species occupancy models, the snow

leopard demonstrated a notably higher probability of occupancy overall compared to other carnivore

species, estimated at 0.437 (Table 1). Conversely, the Eurasian lynx exhibited a lower occupancy

probability, estimated at 0.161. Further analysis revealed that the occupancy probabilities of the wolf and

Eurasian lynx declined with increasing Normalized Difference Vegetation Index (NDVI) (Table 2, Figure

2). Additionally, wolf occupancy probability displayed a negative relationship with roughness index and a

positive relationship with prey availability. Snow leopard occupancy probabilities exhibited a negative

relationship with distance to roads and NDVI. In contrast, both red fox and Tibetan fox demonstrated a

positive relationship with distance to roads. Moreover, red fox occupancy probability increased with

higher human disturbance and greater prey availability.

The detection probabilities of wolf, snow leopard, red fox, and Tibetan fox exhibited an increase

with elevation (Table 2). Moreover, there was a positive relationship between the detection probability of

Tibetan fox and prey availability. The detection probabilities of snow leopard and Eurasian lynx declined

as human disturbance increased.

The Sørensen similarity index (S) ranged from 0.1 to 0.5 (Table 3). Compared with other

combinations of apex-mesocarnivore pairs, snow leopard and red fox (S=0.477) had relatively high

spatial overlap, while Eurasian lynx and Tibetan fox (S=0.198) had the lowest spatial overlap. Moreover,

spatial overlap of apex versus apex carnivores and mesocarnivores versus mesocarnivores was relatively

low.

Daily activity patterns and differences

A total of 1444 independent records were obtained for five carnivore species, consisting of 79

records of wolf, 458 records of snow leopard, 126 records of Eurasian lynx, 421 records of red fox, and

359 records of Tibetan fox. Among apex carnivores (Table 3, Figure 3), the daily activity was similar

between snow leopard and Eurasian lynx and their diel activity overlap was close to 1 (Δ_4=0.900,

P=0.285), their daily activity peak was at 21:00 hr and dawn. However, the wolf had a significantly

different daily activity pattern with snow leopard (Δ_4=0.676, P<0.001) and Eurasian lynx (Δ_4=0.661,

P<0.001), and its daily activity peaked happened around 9:00 and 18:00 hr. Tibetan fox and red fox had

different activity patterns and peaks (P<0.001). The activity peak for red fox activity peaked at 3:00 and

21:00 hr, while the Tibetan fox had a prolonged active bout between noon and dusk. Temporal activity

patterns between apex carnivores and mesocarnivores were significantly different, except for wolf and

Tibetan fox (Δ_4=855, P=0.118).

Dietary composition, diversity and similarity

A total of 26 unique prey species were identified from 9 taxonomic orders (Figure 4, Table S1).

Artiodactyla and lagomorpha were the most frequently detected in the diets of apex carnivores and

mesocarnivores, accounting for 32.81% and 70.18% of prey counts, respectively. Blue sheep made up

26.50% of prey counts in apex carnivore diet, while plateau pika made up 67.11% of prey counts in

mesocarnivore diet. Livestock were present in 17.98% of apex carnivore diet counts and were present in

4.82% mesocarnivore diet counts.

The dietary niche overlap among all carnivore species can be found in Table 3. Wolf and snow

leopard had the highest dietary niche overlap value among apex carnivores (O_{jk}=0.892). The value of

Pianka's index was generally low between apex carnivores and mesocarnivores, except wolf and red fox

(O_{jk}=0.811). In contrast, observed dietary overlap was greatest among the mesocarnivores, especially

Pallas's cat and Tibetan fox, with a value of 0.997.

Red fox had the greatest richness of prey with a value of 16, while Pallas's cat and Tibetan fox had

the lowest diversity of prey with richness value of 6 (Table S2). Dietary similarity was assessed using

inversed Jaccard's Distances, Table S3. Diets were most similar between wolf and snow leopard with a

value of 0.588 and least similar between Pallas's cat and Tibetan fox with a value of 0.200. All other pairs

fell between the values of 0.2 and 0.5.

Discussion

Our study addresses, for the first time, the coexistence patterns of carnivore species present in Qilian

Mountain National Park across multiple dimensions of niche partitioning. This work substantially

contributes to current understanding of carnivore guilds and offers helpful information for biodiversity

conservation at a regional scale. Moreover, our study provides important insights into the potentional

mechanisms of niche partitioning among sympatric carnivores, particularly intra- and interspecific

relations between apex carnivores and mesocarnivores. Specifically, we found that the overall trend of

spatial overlap across carnivores is relatively low, that apex carnivores overlap in time and diet, that

mesocarnivores showed a high degree of dietary overlap, and that there was substantial similarity in diel

activity patterns between apex carnivores and mesocarnivores. These results indicate that carnivores with

similar ecological traits foster co-occurrence by adjusting their daily activity patterns and using differing

food resources to minimize competitive interactions.

We found dietary and temporal overlap among apex carnivores, showing that spatial partitioning is

responsible for their successful coexistence in this area, consistent with our hypothesis. Wolf and snow

leopard had the highest dietary overlap and prey similarity between apex carnivore pairs in this study,

showing that the avoidance of space and time plays an important role in their coexistence. Recent

evidence proves that habitat preference facilitates the coexistence of wolf and snow leopard (Shrotriya et al., 2022). Their hunting strategies may be impacted by their habitat selection. Solitary snow leopards are more suitable for hiding in habitats with features that favor ambush predators, while wolves hunt in packs (Shrotriya et al., 2022). It is clear that wild ungulates (e.g., blue sheep) constituted the primary proportion of wolf and snow leopard diet, followed by small mammals such as plateau pika, Himalayan marmot (*Marmota himalayana*), and woolly hare (*Lepus oiostolus*). In addition, livestock consumption also contributed to the high degree of overlap in their diets (Wang et al., 2014). This supports the optimal foraging theory, in which large predators preferentially select food resources that provide maximum benefit (Brown et al., 1999), but also showed that greater competition for resources is likely to occur between wolf and snow leopard due to their use of the same prey species in this area. This may be especially true in times of habitat stress when resources are poor.

Snow leopard and Eurasian lynx had the highest temporal overlap between apex carnivore pairs in our case, showing that spatial and dietary partitioning facilitate their coexistence. Eurasian lynx is considered an opportunistic predator, and its prey varies among different regions with its primary dietary resource being ungulates and small mammals. For example, Eurasian lynx showed a strong preference for brown hare *L. europaeus* in Turkey, edible dormice *Glis glis* in Slovenia and Croatia, and chamois *Rupicapra rupicapra* or roe deer *Capreolus capreolus* in Switzerland (Mengüllüoğlu et al., 2018; Krofel et al., 2011; Molinari-Jobin et al., 2007). Varied prey selection may be related to sex, age, population density and season (Mengüllüoğlu et al., 2018; Odden et al., 2006). Our results show that woolly hare make up the majority of the Eurasian lynx diet, followed by blue sheep. Woolly hare is mainly distributed in shrubland, meadow, desert and wetland, while blue sheep tend to choose highly sheltered areas, close to bare rocks and cliffs as habitat. Therefore, prey preferences among snow leopard and Eurasian lynx also contributes to spatial avoidance.

Mesocarnivores had substantial overlap in diet, underscoring food resources as a primary competitive factor that necessitates spatial and temporal partitioning for successful coexistence. This finding aligns with previous research indicating that mesocarnivores use temporal and spatial segregation to reduce competition and the probability of antagonistic interspecific encounters (Ferreiro-Arias et al., 2021; Li et al., 2022a). Differences in habitat preference may lead to spatial niche partitioning among mesocarnivores (Wang et al., 2022). In addition, species can adjust temporal periods of behavior to respond to environmental change, competition, and predation risk (Gallo et al., 2022; Finnegan et al., 2021b; van der Vinne et al., 2019). Pallas's cat hunts during crepuscular and diurnal periods and inhabit meadow habitat with greater prey abundance (Anile et al., 2021; Greco et al., 2022; Ross et al., 2019). In contrast, red fox is primarily nocturnal and occupies diverse habitats depending on prey abundance (Gołdyn et al., 2003; Pandolfi et al., 1997; Reshamwala et al., 2022). Tibetan fox is a diurnal hunter of the Tibetan plateau, preferring shrub meadow, meadow steppe and alpine meadow steppe (Gong and Hu, 2003). It is worth noting the substantial overlap in diet between Pallas's cat and Tibetan fox. The dietary overlap between the two was extremely high, with a Pianka's value close to 1. Dietary analyses showed that pika contributed to more than 85% of their collective diets, with 90% of Pallas's cat diet being pika. Pika may be an optimal prey item in the area because of size and year-round activity (Ross, 2009). Previous studies have shown that the Pallas's cat and Tibetan fox are specialist predators of pikas (Harris et al., 2014; Ross, 2009). However, specialization on pika is facultative in that Pallas's cat and Tibetan fox can select other prey items when pika availability is low (Harris et al., 2014; Ross, 2009). This was

observed in our study, even though dietary diversity was low.

Apex carnivores and mesocarnivores exhibited considerable overlap in time overall, showing that

spatial and dietary partitioning may play a large role in facilitating their coexistence. As confirmed by

previous research, kit foxes (*V. macrotis*) successfully coexisted with dominant carnivores by a

combination of spatial avoidance and use of alternative resources (Lonsinger et al., 2017). Differences in

body mass may play a crucial role in minimizing dietary overlap, effectively reducing interspecific

competition between apex and mesocarnivores. Of exception in our study, however, was wolf and red fox,

who exhibited more dietary overlap, showing that temporal and spatial avoidance may promote their

coexistence. As canid generalist-opportunist species, the wolf and red fox consumed similar prey, albeit

the red fox may have obtained livestock and ungulate species via scavenging or by preying on very

young individuals (Hacker et al., 2022). In addition, the wolf and red fox had different peak activity times,

suggesting temporal segregation as a potentially strong driver of coexistence. Our occupancy model also

revealed that mesocarnivores prefer habitats that coincide with human disturbance, possible contributing

to spatial niche differentiation with other apex carnivore species like the wolf. Our findings echo other

recent research which supports the idea that red foxes can coexist with wolves by exploiting a broader

ecological niche (Shrotriya et al., 2022).

Our study highlights the effectiveness of combining camera trapping with DNA metabarcoding for

detecting and identifying both cryptic and rare species within a sympatric carnivore guild. This integrated

approach allowed us to capture a more comprehensive view of species presence and interactions

compared to traditional visual surveys. whereas, it is important to acknowledge the challenges associated

with this technique, including the high costs of equipment and the need for specialized training and

computational resources to manage and analyze the large volumes of sequence data. Despite these

challenges, the benefits of this combined method in improving biodiversity assessments and

understanding species coexistence outweigh the drawbacks. However, several restrictions remain for this

research. The first limitation involves differences in samples sizes. Although the scat samples of Tibetan

fox were relatively low, the accuracy of DNA metabarcoding in informing species presence in diet

ensures that data are informative and thus still important for species conservation management decisions

(Hacker et al., 2022). Second, the methodology of foraging (e.g., predation or scavenging) and the

condition of the prey item (e.g., age or size) cannot be identified in dietary studies (Hacker et al., 2022).

Pika is a prime component of diet among mesocarnivores, especially in the diet of the Pallas's cat and the

Tibetan fox. We surmise that the simultaneous dependence on pika led to partial overlap in spatial and

activity patterns, resulting in increased potential competitive interactions. Due to the lack of spatial and

temporal analysis of Pallas's cat in our study, further monitoring is needed to develop a comprehensive

conservation plan. Despite these limitations, our study provides a foundation from which future studies

interested in niche partitioning among carnivores along spatial, temporal, and dietary dimensions, can be

modeled.

In summary, our study has shown that the coexistence of carnivore species in the landscapes of

Qilian Mountain National Park can be facilitated along three niche axes, with spatial segregation being

the most pronounced of the three. Apex carnivore species tended to overlap temporally or trophically,

mesocarnivore species had high dietary overlap with each other, and apex carnivore and mesocarnivore

species displayed similarity in temporal use. Pika, blue sheep and livestock were found to make up a

large proportion of carnivore diet. Resource competition between wolf and snow leopard and the

interspecific competition between Pallas's cat and Tibetan fox were strong in this area. Based on the findings presented above, we recommend targeted efforts for enhanced protection and management in several key areas. First, greater efforts are needed to protect habitat, including the establishment of habitat corridors and optimizing grassland fence layouts to safeguard migration routes. Second, resource competition should be carefully monitored between snow leopards and wolves, as well as between Pallas's cat and Tibetan foxes. More attention is needed for pika at our study site considering the role they play in the conservation of Pallas's cat and Tibetan fox populations. Previous poisoning campaigns targeting pikas were initiated out of concern for grassland degradation (Smith and Foggin, 1999). Recent research underscores the importance of pika population health and habitat for the distribution of Pallas's cats (Greenspan and Giordano, 2021). It's crucial to note that poisoning campaigns targeting small mammals may be incredibly dangerous for mesocarnivores feeding on them due to secondary poisoning, necessitating caution in such conservation strategies. Further, vigilance regarding the preponderance of smaller prey in predator diets is vital, as this may indicate severe loss of larger prey, which will increase the risk of interference competition (Steinmetz et al., 2021). Third, efforts should focus on restoring vulnerable wild prey populations, strengthening grazing area management, and supporting livelihoods of herders to mitigate livestock predation and consequent human-wildlife conflicts. Regular patrols of protected areas and core habitats are essential, along with educational outreach to conservation staff and herders. Our study corroborates and complements the findings of prior studies on these species and their coexistence mechanisms, offering insights crucial for wildlife conservation in the region.

Acknowledgements

We would like to thank Mr. Jiong He, Yayue Gao, Duifang Ma, Liji Wu, Dazhi Hu and other colleagues of Qilianshan National Park for their generous assistance in the field surveys. We thank Dr. Charlotte Hacker for editing the English text of this manuscript. This work was supported by the Welfare Project of the National Scientific Research Institution (No. CAFYBB2019ZE003 and CAFYBB2018ZD001) and the National Natural Science Foundation of China (No. 32201430 and No. 32101409).

Author contributions

Wei Cong, Conceptualization, Formal analysis, Investigation, Methodology, Software, Validation, Visualization, Writing-original draft, Writing-review and editing; Jia Li, Conceptualization, Funding acquisition, Investigation, Methodology, Resources, Software, Writing-review and editing; Charlotte Hacker, Investigation, Methodology, Writing-review and editing; Ye Li, Investigation; Yu Zhang, Investigation, Funding acquisition; Lixiao Jin, Investigation; Yi Zhang, Investigation; Diqiang Li, Funding acquisition, Methodology, Resources, Supervision, Writing-review & editing; Yadong Xue, Investigation, Resources, Writing-review and editing; Yuguang Zhang, Conceptualization, Funding acquisition, Investigation, Methodology, Resources, Supervision, Writing-review and editing.

Declaration of interests

The authors declare no competing interests.

References

Alexander JS, Shi K, Tallents LA, Riordan P. 2016. On the high trail: examining determinants of site use by the Endangered snow leopard Panthera uncia in Qilianshan, China. *Oryx* **50**:231–238. doi:10.1017/S0030605315001027

Anile S, Augugliaro C, Munkhtsog B, Dartora F, Vendramin A, Bombieri G, Nielsen CK. 2021. Density

424   and activity patterns of Pallas's cats, Otocolobus manul, in central Mongolia. *Wildl Res* **48**:264.
425   doi:10.1071/WR20061

426 Boitani L. 2016. Camera trapping for wildlife research. Pelagic Publishing Ltd.

427 Brown JS, Laundré JW, Gurung M. 1999. The Ecology of Fear: Optimal Foraging, Game Theory, and
428   Trophic Interactions. *Journal of Mammalogy* **80**:385–399. doi:10.2307/1383287

429 Chen X, Tian T, Pan H, Jin Y, Zhang X, Xiang Y, Yang B, Zhang L. 2024. Coexistence mechanisms of
430   small carnivores in a near-pristine area within the mountains of Southwest China. *Global*
431   *Ecology and Conservation* **49**:e02777. doi:10.1016/j.gecco.2023.e02777

432 Chu C, Wang Y, Liu Y, Jiang L, He F. 2017. Advances in species coexistence theory. *Biodiversity Science*
433   **25**:345–354. doi:10.17520/biods.2017034

434 Davis CL, Rich LN, Farris ZJ, Kelly MJ, Di Bitetti MS, Blanco YD, Albanesi S, Farhadinia MS,
435   Gholikhani N, Hamel S, Harmsen BJ, Wultsch C, Kane MD, Martins Q, Murphy AJ, Steenweg R,
436   Sunarto S, Taktehrani A, Thapa K, Tucker JM, Whittington J, Widodo FA, Yoccoz NG, Miller
437   DAW. 2018. Ecological correlates of the spatial co-occurrence of sympatric mammalian
438   carnivores worldwide. *Ecol Lett* **21**:1401–1412. doi:10.1111/ele.13124

439 De Satgé J, Teichman K, Cristescu B. 2017. Competition and coexistence in a small carnivore guild.
440   *Oecologia* **184**:873–884. doi:10.1007/s00442-017-3916-2

441 Deagle BE, Thomas AC, McInnes JC, Clarke LJ, Vesterinen EJ, Clare EL, Kartzinel TR, Eveson JP. 2019.
442   Counting with DNA in metabarcoding studies: How should we convert sequence reads to dietary
443   data? *Mol Ecol* **28**:391–406. doi:10.1111/mec.14734

444 Deiner K, Bik HM, Mächler E, Seymour M, Lacoursière-Roussel A, Altermatt F, Creer S, Bista I, Lodge
445   DM, de Vere N, Pfrender ME, Bernatchez L. 2017. Environmental DNA metabarcoding:
446   Transforming how we survey animal and plant communities. *Mol Ecol* **26**:5872–5895.
447   doi:10.1111/mec.14350

448 Di Bitetti MS, De Angelo CD, Di Blanco YE, Paviolo A. 2010. Niche partitioning and species
449   coexistence in a Neotropical felid assemblage. *Acta Oecologica* **36**:403–412.
450   doi:10.1016/j.actao.2010.04.001

451 Di Minin E, Slotow R, Hunter LTB, Montesino Pouzols F, Toivonen T, Verburg PH, Leader-Williams N,
452   Petracca L, Moilanen A. 2016. Global priorities for national carnivore conservation under land
453   use change. *Sci Rep* **6**:23814. doi:10.1038/srep23814

454 Donadio E, Buskirk SW. 2006. Diet, Morphology, and Interspecific Killing in Carnivora. *The American*
455   *Naturalist* **167**:524–536. doi:10.1086/501033

456 Dormann CF. 2011. How to be a specialist? Quantifying specialisation in pollination networks. *Network*
457   *Biology* **1**:1–20.

458 Ferreiro-Arias I, Isla J, Jordano P, Benítez-López A. 2021. Fine-scale coexistence between Mediterranean
459   mesocarnivores is mediated by spatial, temporal, and trophic resource partitioning. *Ecol Evol*
460   **11**:15520–15533. doi:10.1002/ece3.8077

461 Finnegan SP, Gantchoff MG, Hill JE, Silveira L, Tôrres NM, Jácomo AT, Uzal A. 2021a. "When the
462   felid's away, the mesocarnivores play": seasonal temporal segregation in a neotropical carnivore
463   guild. *Mamm Biol* **101**:631–638. doi:10.1007/s42991-021-00110-9

464 Finnegan SP, Gantchoff MG, Hill JE, Silveira L, Tôrres NM, Jácomo AT, Uzal A. 2021b. "When the
465   felid's away, the mesocarnivores play": seasonal temporal segregation in a neotropical carnivore

guild. *Mamm Biol* **101**:631–638. doi:10.1007/s42991-021-00110-9

Fiske I, Chandler R. 2011. unmarked: An R Package for Fitting Hierarchical Models of Wildlife
Occurrence and Abundance. *Journal of Statistical Software* **43**:1–23.

Frey S, Fisher J, Burton C, Volpe J. 2017. Investigating animal activity patterns and temporal niche
partitioning using camera trap data: Challenges and Opportunities. *Remote Sensing in Ecology*
and Conservation **3**. doi:10.1002/rse2.60

Gallo T, Fidino M, Gerber B, Ahlers A, Angstmann J, Amaya M, Concilio A, Drake D, Gay D, Lehrer E,
Murray M, Ryan T, St. Clair C, Salsbury C, Sander H, Stankowich T, Williamson J, Belaire J,
Simon K, Magle S. 2022. Mammals adjust diel activity across gradients of urbanization. *eLife* **11**.
doi:10.7554/eLife.74756

Garrote G, Pérez De Ayala R. 2019. Spatial segregation between Iberian lynx and other carnivores. *Anim*
Biodiv Conserv 347–354. doi:10.32800/abc.2019.42.0347

Gause GF. 1934. The Struggle for Existence. Williams and Wilkins, Baltimore.

Gołdyn B, Hromada M, Surmacki A, Tryjanowski P. 2003. Habitat use and diet of the red fox Vulpes
vulpes in an agricultural landscape in Poland. *Zeitschrift für Jagdwissenschafte* **49**:191–200.
doi:10.1007/BF02189737

Gómez-Ortiz Y, Monroy-Vilchis O, Mendoza-Martínez GD. 2015. Feeding interactions in an assemblage
of terrestrial carnivores in central Mexico. *Zoological Studies* **54**:16.
doi:10.1186/s40555-014-0102-7

Gong M, Hu J. 2003. The Summer Microhabitat Selection of Tibetan Fox in the Northwest Plateau of
Sichuan. *Acta Theriologica Sinica* **23**:266–269.

Gorczynski D, Hsieh C, Ahumada J, Akampurira E, Andrianarisoa MH, Espinosa S, Johnson S,
Kayijamahe C, Lima MGM, Mugerwa B, Rovero F, Salvador J, Santos F, Sheil D, Uzabaho E,
Beaudrot L. 2022. Human density modulates spatial associations among tropical forest terrestrial
mammal species. *Global Change Biology* **28**:7205–7216. doi:10.1111/gcb.16434

Greco I, Oberosler V, Monti IE, Augugliaro C, Barashkova A, Rovero F. 2022. Spatio‐temporal
occurrence and sensitivity to livestock husbandry of Pallas's cat in the Mongolian Altai. *J Wildl*
Manag **86**:e22150. doi:10.1002/jwmg.22150

Greenspan E, Giordano AJ. 2021. A rangewide distribution model for the Pallas's cat (Otocolobus manul):
identifying potential new survey regions for an understudied small cat. *Mammalia* **85**:574–587.
doi:10.1515/mammalia-2020-0094

Hacker C, Cong W, Xue Y, Li J, Zhang Yu, Wu L, Ji Y, Dai Y, Li Y, Jin L, Li D, Janecka J, Zhang
Yuguang. 2022. Dietary diversity and niche-partitioning of carnivores across the
Qinghai-Tibetan Plateau of China using DNA metabarcoding. *Journal of Mammalogy*
103:1005–1018. doi:https://doi.org/10.1093/jmammal/gyac044

Hacker C, Jevit M, Hussain S, Muhammad G, Munkhtsog Bariushaa, Munkhtsog Bayaraa, Zhang Y, Li D,
Liu Y, Farrington J, Balbakova F, Alamanov A, Kurmanaliev O, Buyanaa C, Bayandonoi G,
Ochirjav M, Liang X, Bian X, Weckworth B, Janecka J. 2021. Regional comparison of snow
leopard (panthera uncia) diet using dna metabarcoding. *Biodiversity and Conservation*
30:797–817. doi:10.1007/s10531-021-02118-6

Hardin G. 1960. The Competitive Exclusion Principle. *Science* **131**:1292–1297.

Harris R, Jiake Z, Ji Y, Zhang K, Chunyan Y, Yu D. 2014. Evidence that the Tibetan fox is an obligate

predator of the plateau pika: Conservation implications. *Journal of Mammalogy* **95**:0–0.
doi:10.1644/14-MAMM-A-021

Haswell P, Jones K, Kusak J, Hayward M. 2018. Fear, foraging and olfaction: how mesopredators avoid
costly interactions with apex predators. *Oecologia* **187**:573–583.
doi:10.1007/s00442-018-4133-3

Janecka J, Jackson R, Yuquang Z, Li D, Munkhtsog B, Buckley-Beason V, Murphy W. 2008. Population
monitoring of snow leopards using noninvasive collection of scat samples: A pilot study. *Animal*
Conservation **11**:401–411. doi:10.1111/j.1469-1795.2008.00195.x

Karanth KU, Srivathsa A, Vasudev D, Puri M, Parameshwaran R, Kumar NS. 2017. Spatio-temporal
interactions facilitate large carnivore sympatry across a resource gradient. *Proc R Soc B*
284:20161860. doi:10.1098/rspb.2016.1860

Kartzinel TR, Chen PA, Coverdale TC, Erickson DL, Kress WJ, Kuzmina ML, Rubenstein DI, Wang W,
Pringle RM. 2015. DNA metabarcoding illuminates dietary niche partitioning by African large
herbivores. *Proc Natl Acad Sci U S A* **112**:8019–8024. doi:10.1073/pnas.1503283112

Kays R, Arbogast BS, Baker☐Whatton M, Beirne C, Boone HM, Bowler M, Burneo SF, Cove MV, Ding
P, Espinosa S, Gonçalves ALS, Hansen CP, Jansen PA, Kolowski JM, Knowles TW, Lima MGM,
Millspaugh J, McShea WJ, Pacifici K, Parsons AW, Pease BS, Rovero F, Santos F, Schuttler SG,
Sheil D, Si X, Snider M, Spironello WR. 2020. An empirical evaluation of camera trap study
design: How many, how long and when? *Methods Ecol Evol* **11**:700–713.
doi:10.1111/2041-210X.13370

Krofel M, Huber D, Kos I. 2011. Diet of Eurasian lynx Lynx lynx in the northern Dinaric Mountains
(Slovenia and Croatia). *Acta Theriologica* **56**:315–322. doi:10.1007/s13364-011-0032-2

Kuijper DPJ, Sahlén E, Elmhagen B, Chamaillé-Jammes S, Sand H, Lone K, Cromsigt JPGM. 2016.
Paws without claws? Ecological effects of large carnivores in anthropogenic landscapes. *Proc R*
Soc B **283**:20161625. doi:10.1098/rspb.2016.1625

Lanszki J, Heltai M, Kövér G, Zalewski A. 2019. Non-linear relationship between body size of terrestrial
carnivores and their trophic niche breadth and overlap. *Basic and Applied Ecology* **38**:36–46.
doi:10.1016/j.baae.2019.06.004

Li J, Li D, Hacker C, Dong W, Wu B, Xue Y. 2022a. Spatial co-occurrence and temporal activity patterns
of sympatric mesocarnivores guild in Qinling Mountains. *Global Ecology and Conservation*
36:e02129. doi:10.1016/j.gecco.2022.e02129

Li J, Xue Y, Hacker CE, Zhang Yu, Li Y, Cong W, Jin L, Li G, Wu B, Li D, Zhang Yuguang. 2021.
Projected impacts of climate change on snow leopard habitat in Qinghai Province, China. *Ecol*
Evol **11**:17202–17218. doi:10.1002/ece3.8358

Li J, Xue Y, Liao M, Dong W, Wu B, Li D. 2022b. Temporal and Spatial Activity Patterns of Sympatric
Wild Ungulates in Qinling Mountains, China. *Animals* **12**:1666. doi:10.3390/ani12131666

Li Z, Wang B, Smith J, Feng R, Feng L, Mou P, Ge J. 2019. Coexistence of two sympatric flagship
carnivores in the human-dominated forest landscapes of Northeast Asia. *Landscape Ecology*
34:291–305. doi:10.1007/s10980-018-0759-0

Linnell JDC, Strand O. 2000. Interference interactions, co☐existence and conservation of mammalian
carnivores. *Diversity and Distributions* **6**:169–176. doi:10.1046/j.1472-4642.2000.00069.x

Lonsinger RC, Gese EM, Bailey LL, Waits LP. 2017. The roles of habitat and intraguild predation by

coyotes on the spatial dynamics of kit foxes. *Ecosphere* **8**:e01749.
doi:https://doi.org/10.1002/ecs2.1749

Ma D, Sun Z, Hu D, An B, Chen L, Zhang D, Dong K, Zhang L. 2021. Camera-trapping survey of the
mammal diversity in the Qilian Mountains National Nature Reserve, Gansu Province. *ACTA*
Theriologica Sinica **41**:90–98. doi:10.16829/j.slxb.150418

MacKenzie DI. 2018. Occupancy estimation and modeling: inferring patterns and dynamics of species
occurrence, Second edition. ed. London, United Kingdom: Academic Press, an imprint of
Elsevier.

Manlick PJ, Pauli JN. 2020. Human disturbance increases trophic niche overlap in terrestrial carnivore
communities. *Proc Natl Acad Sci U S A* **117**:26842–26848. doi:10.1073/pnas.2012774117

Mengüllüoğlu D, Ambarli H, Berger A, Hofer H. 2018. Foraging ecology of Eurasian lynx populations in
southwest Asia: Conservation implications for a diet specialist. *Ecology and Evolution*
8:9451–9463. doi:10.1002/ece3.4439

Molinari-Jobin A, Zimmermann F, Ryser A, Breitenmoser C, Capt S, Breitenmoser U, Molinari P, Haller
H, Eyholzer R. 2007. Variation in diet, prey selectivity and home-range size of Eurasian lynx
Lynx lynx in Switzerland. *Wildlife Biology* **13**:393–405.
doi:10.2981/0909-6396(2007)13[393:VIDPSA]2.0.CO;2

Moreno-Sosa AM, Yacelga M, Craighead KA, Kramer-Schadt S, Abrams JF. 2022. Can prey occupancy
act as a surrogate for mesopredator occupancy? A case study of ocelot (Leopardus pardalis).
Mamm Biol **102**:163–175. doi:10.1007/s42991-022-00232-8

Nasanbat B, Ceacero F, Ravchig S. 2021. A small neighborhood well-organized: seasonal and daily
activity patterns of the community of large and mid-sized mammals around waterholes in the
Gobi Desert, Mongolia. *Front Zool* **18**:25. doi:10.1186/s12983-021-00412-1

Newmaster SG, Thompson ID, Steeves RAD, Rodgers AR, Fazekas AJ, Maloles JR, McMullin RT,
Fryxell JM. 2013. Examination of two new technologies to assess the diet of woodland caribou:
video recorders attached to collars and DNA barcoding. *Can J For Res* **43**:897–900.
doi:10.1139/cjfr-2013-0108

Newsome TM, Greenville AC, Ćirović D, Dickman CR, Johnson CN, Krofel M, Letnic M, Ripple WJ,
Ritchie EG, Stoyanov S, Wirsing AJ. 2017. Top predators constrain mesopredator distributions.
Nat Commun **8**:15469. doi:10.1038/ncomms15469

Odden J, Linnell J, Andersen R. 2006. Diet of Eurasian lynx, Lynx lynx, in the boreal forest of
southeastern Norway: The relative importance of livestock and hares at low roe deer density.
European Journal of Wildlife Research **52**:237–244. doi:10.1007/s10344-006-0052-4

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB,
Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2019. vegan: Community Ecology
Package.

Palencia P, Vicente J, Soriguer RC, Acevedo P. 2022. Towards a best☐practices guide for camera trapping:
assessing differences among camera trap models and settings under field conditions. *Journal of*
Zoology **316**:197–208. doi:10.1111/jzo.12945

Pandolfi M, Forconi P, Montecchiari L. 1997. Spatial behaviour of the red fox (Vulpes vulpes) in a rural
area of central Italy. *Italian Journal of Zoology* **64**:351–358. doi:10.1080/11250009709356222

Reshamwala H, Raina P, Hussain Z, Khan S, Dirzo R, Habib B. 2022. On the move: spatial ecology and

habitat use of red fox in the Trans-Himalayan cold desert. *PeerJ* **10**:e13967.
doi:10.7717/peerj.13967

Riaz T, Shehzad W, Viari A, Pompanon F, Taberlet P, Coissac É. 2011. EcoPrimers: Inference of new
DNA barcode markers from whole genome sequence analysis. *Nucleic acids research* **39**:e145.
doi:10.1093/nar/gkr732

Ridout M, Linkie M. 2009. Estimating overlap of daily activity patterns from camera trap data. *Journal of*
Agricultural, Biological, and Environmental Statistics **14**:322–337.
doi:10.1198/jabes.2009.08038

Ripple WJ, Estes JA, Beschta RL, Wilmers CC, Ritchie EG, Hebblewhite M, Berger J, Elmhagen B,
Letnic M, Nelson MP, Schmitz OJ, Smith DW, Wallach AD, Wirsing AJ. 2014. Status and
ecological effects of the world's largest carnivores. *Science* **343**:1241484.
doi:10.1126/science.1241484

Ritchie EG, Johnson CN. 2009. Predator interactions, mesopredator release and biodiversity conservation.
Ecol Lett **12**:982–998. doi:10.1111/j.1461-0248.2009.01347.x

Ross S. 2009. Providing an ecological basis for the conservation of the Pallas's cat (Otocolobus manul).
PhD thesis, University of Bristol, UK. doi:10.13140/RG.2.1.4338.1521

Ross S, Barashkova A, Kirilyuk V, Naidenko S. 2019. The behaviour and ecology of the manul.

Rowcliffe M, Kays R, Kranstauber B, Carbone C, Jansen P. 2014. Quantifying levels of animal activity
using camera-trap data. *Methods in Ecology and Evolution* **5**. doi:10.1111/2041-210X.12278

Santos F, Carbone C, Wearn O, Rowcliffe M, Espinosa S, Lima M, Ahumada J, Gonçalves A, Trevelin L,
Alvarez-Loayza P, Spironello W, Jansen P, Juen L, Peres C. 2019. Prey availability and temporal
partitioning modulate felid coexistence in Neotropical forests. *PLoS ONE* **14**.
doi:10.1371/journal.pone.0213671

Schoener TW. 1974. Resource Partitioning in Ecological Communities. *Science*.

Searle CE, Smit JB, Cusack JJ, Strampelli P, Grau A, Mkuburo L, Macdonald DW, Loveridge AJ,
Dickman AJ. 2021. Temporal partitioning and spatiotemporal avoidance among large carnivores
in a human-impacted African landscape. *PLoS ONE* **16**:e0256876.
doi:10.1371/journal.pone.0256876

Shannon CE, Weaver W. 1949. The Mathematical Theory of Communication. *Philosophical Review* **60**.

Shao X, Lu Q, Xiong M, Bu H, Shi X, Wang D, Zhao J, Li S, Yao M. 2021. Prey partitioning and
livestock consumption in the world's richest large carnivore assemblage. *Curr Biol*
31:4887-4897.e5. doi:10.1016/j.cub.2021.08.067

Shrotriya S, Reshamwala H, Lyngdoh S, Jhala Y, Habib B. 2022. Feeding Patterns of Three Widespread
Carnivores—The Wolf, Snow Leopard, and Red Fox—in the Trans-Himalayan Landscape of
India. *Frontiers in Ecology and Evolution* **10**:815996. doi:10.3389/fevo.2022.815996

Smith A, Foggin M. 1999. The plateau pika (Ochotona curzoniae) is a keystone species for biodiversity
on the Tibetan plateau. *Animal Conservation* **2**:235–240. doi:10.1017/S1367943099000566

Smith JA, Suraci JP, Clinchy M, Crawford A, Roberts D, Zanette LY, Wilmers CC. 2017. Fear of the
human 'super predator' reduces feeding time in large carnivores. *Proc R Soc B* **284**:20170433.
doi:10.1098/rspb.2017.0433

Sorensen TJ. 1948. A method of establishing groups of equal amplitude in plant sociology based on
similarity of species content and its application to analyses of the vegetation on Danish commons.

634     Kobenhavn,: Ejnar Minksgaard,.

635 Steinmetz R, Seuaturien N, Intanajitjuy P, Inrueang P, Prempree K. 2021. The effects of prey depletion on

636     dietary niches of sympatric apex predators in Southeast Asia. *Integr Zool* **16**:19–32.

637     doi:10.1111/1749-4877.12461

638 Strampelli P, Henschel P, Searle CE, Macdonald DW, Dickman AJ. 2023. Spatial co-occurrence patterns

639     of sympatric large carnivores in a multi-use African system. *PLoS ONE* **18**:e0280420.

640     doi:10.1371/journal.pone.0280420

641 Torretta E, Riboldi L, Costa E, Delfoco C, Frignani E, Alberto M. 2021. Niche partitioning between

642     sympatric wild canids: the case of the golden jackal (Canis aureus) and the red fox (Vulpes

643     vulpes) in north-eastern Italy. *BMC Ecology and Evolution* **21**:129.

644     doi:10.1186/s12862-021-01860-3

645 Tsunoda H, Newman C, Peeva S, Raichev E, Buesching CD, Kaneko Y. 2020. Spatio-temporal

646     partitioning facilitates mesocarnivore sympatry in the Stara Planina Mountains, Bulgaria.

647     *Zoology (Jena)* **141**:125801. doi:10.1016/j.zool.2020.125801

648 van der Vinne V, Tachinardi P, Riede SJ, Akkerman J, Scheepe J, Daan S, Hut RA. 2019. Maximising

649     survival by shifting the daily timing of activity. *Ecol Lett* **22**:2097–2102. doi:10.1111/ele.13404

650 Vilella M, Ferrandiz-Rovira M, Sayol F. 2020. Coexistence of predators in time: Effects of season and

651     prey availability on species activity within a Mediterranean carnivore guild. *Ecol Evol*

652     **10**:11408–11422. doi:https://doi.org/10.1002/ece3.6778

653 Wang D, Sai Q, Wang Z, Zhao H, Lian X. 2022. Spatiotemporal overlap among sympatric Pallas's cat

654     (Otocolobus manul), Tibetan fox (Vulpes ferrilata) and red fox (V. vulpes) in the source region of

655     the Yangtze River. *Biodiv Sci* **30**:21365. doi:10.17520/biods.2021365

656 Wang J, Laguardia A, Damerell PJ, Riordan P, Shi K. 2014. Dietary overlap of snow leopard and other

657     carnivores in the Pamirs of Northwestern China. *Chin Sci Bull* **59**:3162–3168.

658     doi:10.1007/s11434-014-0370-y

659 Xue Y, Li J, Hu Y, Ma L, Qian W, Yan P, Yang M, Chen D, Wu B, Li D. 2019. Camera-trapping survey of

660     the mammals and birds in the Qilian Mountain National Park (Qinghai area), China. *ACTA*

661     *Theriologica Sinica* **39**:466–475. doi:10.16829/j.slxb.150307

662 Zhang J. 2016. spaa: SPecies Association Analysis.

663 Zheng J. 2011. The study of Qinghai nature reserve. Qinghai People's Publishing House.

666 **Figure 1** Locations of camera trap stations and scat collection sites in this study.

Figure 2 The relationship between carnivore species occupancy probability and covariates by the optimal

models (ΔAIC≤2). The solid line represents the fitted polynomial regression and the gray area represents

95% confidence intervals.

673
674 **Figure 3** Daily activity patterns of carnivore species. Shades of gray indicate temporal areas of overlap.
675

676
Figure 4 The food web of carnivore species (SL-snow leopard, EL-Eurasian lynx, PC-Pallas's cat,
RF-red Fox, TF-Tibetan Fox). The heights of the left bars represent the frequency of occurrence of the
taxonomic order of prey species, the middle bars represent the number of samples for each carnivore, and
the heights of the right bars represent the frequency of occurrence of prey species in scats. The colors of
prey match the taxonomic orders. The connecting line widths represent the prey frequency of occurrence
in the diet of each carnivore species.
683

684   **Table 1** Summary of occupancy rate and detection probability of different species for the optimal models

685   (ΔAIC≤2).

Species	Models	Number of parameters	AIC	ΔAIC	AIC $_{Wt}$	ψ	Pr
	Psi (rix+ndvi+prey); *P* (ele+prey)	7	282.14	0.000	0.077	0.180	0.153
	Psi (rix+ndvi+prey); *P* (ele+hdis)	7	282.45	0.302	0.066	0.183	0.145
	Psi (rix+ndvi+prey); *P* (ele)	6	283.67	1.525	0.036	0.217	0.111
Wolf	*Psi* (rix+ndvi+prey); *P* (ele+hdis+prey)	8	284.10	1.959	0.029	0.179	0.154
	Psi (ele+rix+ndvi+prey); *P* (ele+prey)	8	284.12	1.972	0.029	0.168	0.165
	Psi (rix+ndvi+disrd+prey); *P* (ele+prey)	8	284.14	2.000	0.028	0.178	0.156
	Model average					**0.184**	**0.147**
	Psi (disrd+ndvi); *P* (hdis+prey)	6	970.69	0.000	0.035	0.436	0.419
	Psi (ele+disrd); *P* (hdis+prey)	6	970.81	0.113	0.033	0.435	0.421
	Psi (.); *P* (hdis+prey)	4	970.99	0.292	0.030	0.437	0.420
	Psi (disrd+ndvi); *P* (ele+hdis+prey)	7	971.12	0.429	0.028	0.440	0.413
	Psi (ele+disrd+ndvi); *P* (hdis+prey)	7	971.13	0.439	0.028	0.433	0.420
	Psi (.); *P* (ele+hdis+prey)	5	971.21	0.511	0.027	0.442	0.413
	Psi (disrd); *P* (ele+hdis+prey)	6	971.22	0.525	0.027	0.444	0.410
Snow leopard	*Psi* (disrd); *P* (hdis+prey)	5	971.52	0.825	0.023	0.437	0.420
	Psi (ele); *P* (hdis+prey)	5	971.62	0.928	0.022	0.435	0.421
	Psi (ndvi); *P* (hdis+prey)	5	971.64	0.947	0.022	0.437	0.420
	Psi (ele+disrd); *P* (ele+hdis+prey)	7	971.92	1.222	0.019	0.436	0.416
	Psi (ndvi); *P* (ele+hdis+prey)	6	972.42	1.726	0.015	0.440	0.414
	Psi (ele+disrd+ndvi); *P* (ele+hdis+prey)	8	972.45	1.757	0.014	0.435	0.416
	Psi (disrd+ndvi+rix); *P* (hdis+prey)	7	972.67	1.974	0.013	0.436	0.419
	Model average					**0.437**	**0.417**
	Psi (ndvi); *P* (hdis)	4	371.32	0.000	0.025	0.163	0.321
	Psi (rix+ndvi); *P* (hdis)	5	371.40	0.076	0.024	0.161	0.322
	Psi (ndvi); *P* (hdis+prey)	5	371.64	0.325	0.021	0.170	0.296
	Psi (rix+ndvi); *P* (hdis+prey)	6	371.86	0.539	0.019	0.167	0.300
	Psi (ndvi+hdis); *P* (prey)	5	371.94	0.623	0.019	0.163	0.315
	Psi (ndvi+disrd); *P* (hdis)	5	372.02	0.702	0.018	0.163	0.321
	Psi (rix+ndvi+hdis); *P* (prey)	6	372.23	0.907	0.016	0.161	0.318
Eurasian lynx	*Psi* (ndvi+hdis); *P* (.)	4	372.24	0.916	0.016	0.156	0.343
	Psi (rix+ndvi+hdis); *P* (.)	5	372.36	1.044	0.015	0.154	0.345
	Psi (rix+ndvi+disrd); *P* (hdis)	6	372.39	1.072	0.015	0.161	0.321
	Psi (ndvi+disrd); *P* (hdis+prey)	6	372.64	1.318	0.013	0.168	0.300
	Psi (ndvi+prey); *P* (hdis)	5	372.67	1.352	0.013	0.163	0.320
	Psi (rix+ndvi+prey); *P* (hdis)	6	372.99	1.672	0.011	0.160	0.323
	Psi (ndvi+disrd+hdis); *P* (.)	5	373.06	1.744	0.011	0.156	0.344

Psi (ndvi+hdis); *P* (hdis)	5	373.09	1.775	0.010	0.156	0.325
Psi (ndvi+disrd+hdis); *P* (prey)	6	373.11	1.792	0.010	0.162	0.319
Psi (rix+ndvi+disrd); *P* (hdis+prey)	7	373.12	1.796	0.010	0.165	0.303
Psi (rix+ndvi+hdis); *P* (hdis)	6	373.23	1.913	0.010	0.154	0.326
Psi (ele+ndvi); *P* (hdis)	5	373.32	1.999	0.009	0.164	0.319
Model average					**0.161**	**0.320**
Psi (disrd+hdis+prey); *P* (.)	5	894.44	0.000	0.039	0.369	0.391
Psi (disrd+hdis+prey); *P* (ele)	6	894.59	0.152	0.037	0.370	0.387
Psi (disrd+hdis+ndvi+prey); *P* (.)	6	895.30	0.864	0.026	0.370	0.391
Psi (disrd+hdis+ndvi+prey); *P* (ele)	7	895.37	0.934	0.025	0.370	0.387
Red fox *Psi* (disrd+hdis+prey+rix); *P* (.)	6	895.99	1.544	0.018	0.369	0.392
Psi (disrd+hdis+prey+rix); *P* (ele)	7	896.16	1.717	0.017	0.370	0.388
Psi (disrd+hdis+prey); *P* (prey)	6	896.29	1.850	0.016	0.368	0.388
Psi (disrd+hdis+prey); *P* (hdis)	6	896.37	1.932	0.015	0.368	0.394
Model average					**0.369**	**0.390**
Psi (ele+disrd+prey+rix); *P* (ele+prey)	8	326.43	0.000	0.066	0.147	0.236
Psi (disrd+prey+rix); *P* (ele+prey+hdis)	8	327.14	0.714	0.046	0.277	0.135
Psi (ele+disrd+hdis+prey+rix); *P* (ele+prey)	9	327.20	0.775	0.045	0.151	0.230
Psi (ele+disrd+rix); *P* (ele+prey)	7	327.29	0.866	0.043	0.158	0.219
Tibetan fox *Psi* (ele+disrd+prey+rix); *P* (ele+prey+hdis)	9	327.53	1.100	0.038	0.153	0.222
Psi (ele+disrd+prey); *P* (ele+prey)	7	327.69	1.263	0.035	0.149	0.234
Psi (disrd+hdis+prey+rix); *P* (ele+prey)	8	328.01	1.585	0.030	0.260	0.142
Psi (ele+disrd); *P* (ele+prey)	6	328.42	1.986	0.024	0.159	0.219
Model average					**0.182**	**0.205**

686 Abbreviations: Disrd-distance to roads, Ele-elevation, NDVI-normalized difference vegetation index, Rix- roughness index,

687 hdis-human disturbance.

689 **Table 2** Covariates influencing carnivore occupancy rate and detection probability based on the optimal

690 models (ΔAIC≤2).

Species	Model component	Covariates	Estimate (β)	SE	Z	P
Wolf	Occupancy	Intercept	-2.031	0.497	4.083	<0.001***
		Ndvi	-0.843	0.287	2.935	0.003**
		Prey	1.318	0.534	2.470	0.014*
		Rix	-1.166	0.570	2.046	0.041*
	Detection	Intercept	-1.920	0.457	4.198	<0.001***
		Ele	0.618	0.266	2.320	0.020*
Snow leopard	Occupancy	Intercept	-0.256	0.143	1.783	0.075
		Disrd	-0.163	0.172	0.953	0.341
		Ndvi	-0.095	0.145	0.653	0.514
	Detection	Intercept	-0.451	0.124	3.628	<0.001***
		Hdis	-1.830	0.520	3.519	<0.001***
		Ele	0.661	0.247	2.678	0.007**
Eurasian lynx	Occupancy	Intercept	-1.879	0.316	5.941	<0.001***
		Ndvi	-0.583	0.201	2.899	0.004**
	Detection	Intercept	-0.999	0.375	2.665	0.008**
		Hdis	-1.785	1.690	1.056	0.291
Red fox	Occupancy	Intercept	-0.441	0.177	2.488	0.013*
		Disrd	0.331	0.178	1.866	0.062
		Hdis	1.333	0.611	2.184	0.029*
		Prey	0.665	0.319	2.083	0.037*
	Detection	Intercept	-0.451	0.121	3.729	<0.001***
		Ele	0.057	0.093	0.606	0.545
Tibetan fox	Occupancy	Intercept	-2.087	0.643	3.247	<0.001***
		Disrd	0.955	0.321	2.973	0.003**
		Ele	0.768	0.547	1.403	0.161
	Detection	Intercept	-1.679	0.685	2.453	0.014*
		Ele	0.882	0.417	2.112	0.035*
		Prey	0.369	0.167	2.216	0.027*

691 Abbreviations: Disrd-distance to roads, Ele-elevation, NDVI-normalized difference vegetation index, Rix- roughness index,

692 hdis-human disturbance.

693 The different superscript letters represent significance, *** p< 0.001, **0.001<p<0.01, * 0.01 < p< 0.05.

694 **Table 3** Spatial overlap (Sørensen's index), diel activity overlap (Δ) and dietary overlap (Pianka's index),
695 as well as confidence intervals for carnivore species.

	Sørensen's index	Δ	Pianka's index
Wolf – Snow leopard	0.277	0.676 (0.562-0.756)	0.892 (0.804-0.982)
Wolf – Eurasian lynx	0.272	0.661 (0.541-0.759)	0.585 (0.141-0.881)
Snow leopard – Eurasian lynx	0.305	0.900 (0.854-0.992)	0.458 (0.160-0.886)
Wolf – Pallas's cat	—	—	0.658 (0.053-0.950)
Wolf – Red fox	0.365	0.497 (0.359-0.563)	0.811 (0.497-0.962)
Wolf – Tibetan fox	0.350	0.855 (0.777-0.937)	0.689 (0.456-0.967)
Snow leopard – Pallas's cat	—	—	0.354 (0.092-0.827)
Snow leopard – Red fox	0.477	0.814 (0.756-0.863)	0.586 (0.568-0.941)
Snow leopard – Tibetan fox	0.292	0.711 (0.629-0.744)	0.390 (0.299-0.905)
Eurasian lynx – Pallas's cat	—	—	0.376 (0-0.921)
Eurasian lynx – Red fox	0.205	0.800 (0.728-0.878)	0.536 (0.442-0.910)
Eurasian lynx – Tibetan fox	0.198	0.695 (0.601-0.756)	0.385 (0.078-0.919)
Pallas's cat – Red fox	—	—	0.945 (0.369-0.988)
Pallas's cat – Tibetan fox	—	—	0.997 (0-1)
Red fox – Tibetan fox	0.349	0.576 (0.467-0.580)	0.949 (0.279-0.988)