1 **H2-O deficiency promotes regulatory T cell differentiation and CD4 hyperactivity**
2

3 Robin A. Welsh[1], Nianbin Song[1], Chan-su Park[1,2], J. David Peske[1], Scheherazade Sadegh-
4 Nasseri[1,4]*
5

6 [1]Department of Pathology, Johns Hopkins School of Medicine, Baltimore, MD, USA
7 [2]Current Address: Department of Pharmaceutics, College of Pharmacy, Chungbuk National
8 University, 1 Chungdae-ro, Seowon-gu, Cheongju 28644, Korea
9 [4]Senior Author
10 *Corresponding Author: ssadegh@jhmi.edu

11
12
13

14   **Abstract**
15
Regulatory T cells (Treg) are crucial immune modulators, yet the exact mechanism of thymic
Treg development remains controversial. Here, we present the first direct evidence for H-2O, a
peptide editing molecular chaperon for MHC class II antigen processing, on selection of thymic
Tregs. We demonstrate that lack of H2-O in the thymic medulla promotes thymic Treg
development and leads to an increased peripheral Treg frequency. Single-cell RNA-sequencing
(scRNA-seq) analysis of splenic CD4 T cells revealed an enrichment of both effector-like Tregs
and activated CD4 T cells in the absence of H2-O. H2-O has been shown to increase the
presentation of high-affinity class II epitopes, thus leading to proper negative selection in the
medullary thymus. In spite of this notion, our data support two concepts; a) lack of expression
of H2-O in thymic medulla creates an environment permissive to Treg development and, b) that
loss of H2-O drives increased basal auto-stimulation of CD4 T cells. These findings can help in
better understanding of autoimmunity and design of therapeutics for treatment of
autoimmune diseases.
29
30
31

Introduction

T cells are key players in humoral immune responses. Upon infection with a pathogen,
CD4 T cells utilize their T cell receptor (TCR) to survey peptides presented by professional
antigen presenting cells (APCs) on class II molecules (pMHCII) for recognition. Identification of a
cognate pMHCII complex by the TCR leads to CD4 T cell activation and ultimately clearance of
the foreign pathogen. Faulty activation, however, can lead to deleterious inflammation causing
possible autoimmune diseases and cancer development. Therefore, multiple regulatory
processes exist to ensure proper T cell activation.
Regulation of T cell activation begins during thymic development where immature
thymocytes are screened for self-reactivity. Broadly divided into positive and negative
selection, this process ensures that CD4 T cells expressing high avidity self-reactive TCRs are
either eliminated (1), or become CD4 regulatory T cells (Tregs) (2). Critical to the screening
process for identifying auto-reactive T cells is the Class II antigen processing machinery
expressed by medullary thymic epithelium cells (mTECs) and thymic antigen presenting cells
(APCs). Two chaperone proteins, H2-M (murine; human, HLA-DM), and H2-O (murine; human,
HLA-DO) are major components of the MHC II processing pathway. While H2-M is expressed in
all APCs, H2-O is expressed mainly in thymic medulla and B cells (3). H2-M plays a critical role in
MHC class II antigen processing by dissociating the Class II Invariant Chain peptide (CLIP) from
the newly synthesized MHC II. Dissociation of CLIP promotes a peptide-receptive MHC II
conformation that can readily scan for the best MHC II groove fitting epitopes of denatured
antigens. However, a peptide-receptive MHC II conformation is highly transient (4, 5, 6, 7). We
have proposed that H2-O binds to H2-M (8) and works cooperatively with H2-M by stabilizing
the peptide-receptive MHC II conformation in order to optimize epitope selection (9). Together,
these two molecules ensure that the best MHC II groove fitting epitopes are presented to
cognate CD4 T cells.
While the exact mechanism of Treg selection remains to be fully understood (10), two
critical requirements have been identified as necessary for successful thymic Treg development.
First, thymic Treg development requires relatively strong TCR signaling in the thymic medulla
(11), and second, Treg development relies on signaling by common γ chain (γC) cytokines,
mainly IL-2, for driving Foxp3 expression (12). Considering that strong TCR signaling during
negative selection normally leads to CD4 T cell deletion (1), when investigating Treg selection,
both the nature and density of medulla presented self-peptides must be considered. If epitopes
are in high abundance and more ubiquitously expressed, then cognate CD4 T cells will undergo
clonal deletion. However, if epitopes are in lower abundance and have a sparser expression,
leading to discontinuous TCR stimulation, then cognate CD4 T cells might undergo Treg
selection (10, 13, 14). This model of Treg selection relies on the level of TCR signaling that
medulla localized CD4 single-positive (SP) T cells receive. Indeed, loss of H2-O function has been
shown to alter both the repertoire and density of peptides presented by naïve peripheral B cells
(15, 16). Thus, loss of H2-O could affect whether CD4 T cells are signaled for deletion, or
survival during thymic deletion. Should a lower density of self pMHCII be presented by thymic
epithelium, an increased number of self-reactive CD4 T cells might escape deletion leading to

an increased frequency of auto-reactive T cell clones in the periphery. Or, alternatively,
presentation of a lower density of self pMHCII could promote selection of regulatory T cells.
Here, we demonstrate that loss of H2-O generates a more stimulatory *in vivo*
environment impacting both the thymic development and peripheral activation of regulatory T
cells.
80
81   **Results**
Loss of H2-O increases the activation state of auditing medulla CD4 T cells
Previously, we demonstrated that loss of H2-O expression correlated with both an
increased B cell presentation of low-affinity MHC II peptides, and an increased frequency of a
MOG_{35-55} specific, self-reactive CD4 T cell (15). Because of H2-O expression in the medullary
thymus we speculated that H2-O deficiency might lead to presentation of lower densities of
high-affinity self-peptides in the medulla, thereby causing an altered clonal deletion process.
Medulla CD4 T cells can be divided into two groups based upon expression of the pro-apoptotic
protein Caspase-3 (17); "Auditing" CD4 T cells (CCR7+ CD4+ Caspase-3_{neg}) that are still
undergoing self-auditing, and "Clonally Deleted" CD4 T cells (CCR7+ CD4+ Caspase-3_{pos}) that
have been selected for thymic deletion.
We found that loss of H2-O significantly increased expression of the activation marker
CD69 on auditing, but not CD4 T cells selected for clonal deletion (**Fig 1a**). CD69 in combination
with MHC-I levels defines 3 medullary maturation stages: semi-mature (CD69+ MHC-I -), mature
1 (CD69+ MHC-I +), and mature 2 (CD69- MHC-I -) (17). Subdivision of auditing H2-O-KO CD4 T
cells identified a significant increase in the Mature 1 stage (**Fig 1b, middle**). Conversely, both
the semi-mature and mature 2 stages were decreased in H2-O KO mice (**Fig 1b left/right**). H2-O
deficiency, however, did not appear to alter the rate of CD4 T cells undergoing clonal deletion
(**Fig 1c**). Importantly, no differences were observed in thymocytes undergoing positive selection
(**sFig1**). These data suggest that loss of H2-O drives a more stimulatory thymic medulla
environment, but with similar levels of clonal deletion. It is therefore likely that the increased
peripheral frequencies of MOG-specific CD4 T cells previously identified is due to increased
peripheral expansion of the MOG-reactive clone, not a general altered clonal deletion.
104
H2-O KO thymi have increased regulatory T cell development
The increase in the Mature 1 (M1) state was noteworthy as Treg development has been
shown to begin in the M1 stage (17). With increased levels of peripheral Treg previously
identified in H2-O KO mice (15), we wondered if H2-O deficiency was affecting Treg selection. In
fact, one model of thymic Treg selection centers around the concept of antigen density (13, 18).
Within this "mosaic" model, sporadic MHC-TCR interactions with sparsely presented self-
epitopes leads to Treg development. Having previously shown that peripheral loss of H2-O
leads to increased presentation of lower affinity peptides (15), we postulated that altering the
level of self-epitopes present in the medulla could alter Treg selection. In support of this idea,
we found an increased frequency of CD25+Foxp3+ T cells in the H2-O KO CD4 SP population (**Fig
2a**). Furthermore, H2-O KO thymic Tregs (tTregs) expressed higher levels of CD25, CD5 and the
orphan nuclear receptor Nur77 (**Fig 2b**). As Nur77 has been associated with the level of TCR
engagement (19), increased Nur77 signal was particularly interesting since it indicates that
absence of H2-O leads to increased TCR engagement suggesting an increased self-reactivity. We

also found that H2-O KO tTregs were enriched in the M1 maturation stage (**Fig 2c**). This could
mean two things; (1) loss of H2-O leads to increased presentation of specific Treg epitopes
thereby increasing the frequency of Treg selection, or (2) lower antigen density allows an
increased proportion of self-reactive CD4 T cells to engage their cognate ligands at suboptimal
levels for clonal deletion but optimal ranges for Treg selection. Cumulatively, evaluation of H2-
O KO thymi showed that only CD4 T cells undergoing the auditing process were affected by H2-
O loss.
126
127
Loss of H2-O correlates with increased peripheral CD4 T cell activation
In light of the observation that lack of H2-O KO did not alter clonal deletion frequencies,
we next evaluated whether peripheral loss of H2-O increased CD4 T cells activation. We found
that H2-O KO spleens had a dramatic increase in the proportion of CD3+ CD4 T cells (**Fig 3a**).
Critically, unimmunized H2-O KO spleens had increased levels of CD4 T cells co-expressing the
key activation markers CD44+ and CD69+ (**Fig 3b**). We further assessed the levels of "non-
activated" (CCR7+ CD62L+) versus "activated" (CCR7- CD62L-) CD4 T cells (20), and found an
overall decreased percentage of CCR7 expressing CD4 T cells in H2-O KO spleens (**Fig 3c**) and
lower frequencies of non-activated CD4 T cells in H2-O KO mice (**Fig 3d**). Importantly, this
correlated with an increase in percentage of activated CD4 T cells (**Fig 3d**). Collectively, these
phenotypic analyses support the idea that loss of H2-O leads to increased basal levels of
activated CD4 T cells in unimmunized H2-O KO mice.
As discussed above, H2-O KO thymi promoted Tregs selection. Consistent with these
observations and our previously published data (15, 21), H2-O KO spleens had an increased
frequency of CD25+ Foxp3+ Tregs (**Fig 4a**). Furthermore, H2-O KO Tregs showed decreased
levels of CD62L expression indicating these cells are likely more activated and circulating
through the periphery (**Fig 4b**). Evaluation of the intracellular levels of Nur77 expression again
revealed an increase, suggesting a higher TCR stimulation (**Figure 4c)** further validating the
observations that loss of H2-O is driving a more activated state in both conventional CD4 T cells
and Treg cells.
148
Single cell RNA-Sequencing of H2-O KO splenic CD4 T cells confirms increased activation
Based upon the strong FACS data above exhibiting increased numbers of Tregs, and
more activated CD4 T cells, we attempted single-cell RNA-sequencing (scRNA-seq) to gain a
more holistic unbiased characterization. CD3+ CD4+ NK1.1- CD19- cells were sorted from
spleens of 3 unimmunized H2-O WT and 3 H2-O KO mice and subjected to 10x Genomics
scRNA-seq analyses. Principle Component Analysis (PCA) of the normalized scRNA-seq data
identified 11 distinct CD4 T cell clusters (**Fig 5a & sFig 2**). Separation of the clusters based upon
H2-O WT or H2-O KO genotypes identified a dramatic increase in cluster 0, cluster 2 and cluster
3 in the H2-O KO samples. Conversely, clusters 1 and 4 were significantly increased in H2-O WT
samples (**Fig 5b**).
Comparing the genes from Clusters 0, 1 and 5 to the published CD4 T cell datasets using
the "My Geneset" function on the Immunological Genome Project (www.imgen.org) suggested
that upregulated genes (Log$_2$Avg Fold-change (FC) $\geq$ 0.5) were present in both "naïve" and
"activated" CD4 T cell datasets. Based upon the high expression of CCR7 however, these

clusters were labeled as "non-activated". Clusters 3, 6, 7, and 11 aligned with mainly an
"activated" CD4 T phenotype. Cluster 3 was found to have increased expression of the
activation marker CD44, while Cluster 11 expressed high Ki67. Clusters 2 and 4 contained the
known Treg genes Foxp3 and Il2ra (CD25) (**Fig 5a & Sup Table 1**). Clusters 8, 9, and 10 were
small populations of Macrophages and NKT cell (**Sup Table 1**). Condensing the cluster analysis
further supported our initial FACS data that unimmunized H2-O-KO mice have a decreased
frequency of "Non-activated" CD4 T cells (68.90% H2-O KO vs 74.17% H2-O WT with a
coinciding increase in "Activated" (14.14% H2-O KO vs 8.53% H2-O WT) CD4 T cells (**Fig 5b**). To
further test whether the higher activated state in H2-O KO T cells were due to the H2-O KO CD4
T cells themselves, or the environment, we performed an   adoptive transfer experiment of dye
labeled naïve (CD4+ CD44- CD25-) Thy1.1+ CD4 T cells from H2-O WT into either unimmunized
Thy1.2+ H2-O WT, or Thy1.2+ H2-O KO hosts. As shown in **Fig 5d**, H2-O KO hosts induced
significantly more *in vivo* proliferation of the WT cells than the H2-O WT hosts. These data
additionally confirms that loss of H2-O creates a more activating CD4 T cell environment by
presenting a wider range of self-epitopes (15).
178
H2-O KO Tregs have a more activated phenotype
Analyses of the two Treg clusters revealed that the frequency of Clusters 2 and 4 were
roughly equal in H2-O WT samples, whereas H2-O KO samples had an overrepresentation of
Cluster 2 (**Fig 6a, left**). Both cluster 2 and 4 expressed the classic Treg genes Foxp3, IL2ra
[CD25], CTLA4, and Pdcd1 [PD-1] (**Fig 6a, right**; **Sup Table 1**). Also identified was the
transcription factor Ikzf2 [Helios], a marker of stable Treg lineage commitment in inflammatory
conditions (22) and debated marker of thymic Tregs (23, 24, 25, 26). Supporting our scRNA-Seq
analysis, Helios protein expression was found in both H2-O WT and H2-O KO CD4 Tregs (**Fig 6b**).
The data also showed significant Helios expression in H2-O KO Foxp3- CD4 T cells supporting
that loss of H2-O increases CD4 T cell activation. Interestingly, comparison of the cluster 2
genetic signature with a published Treg data set indicated that these cells are likely effector-like
Tregs. Furthermore, breaking down Cluster 2 by genotype showed that H2-O KO Tregs had a
larger fold change in expression of the core effector genes signature (27) (**Fig 6c**). We
hypothesize that this activated Treg state is likely driven by the increased basal CD4 T cell
activation. Expansion of this Treg population could certainly be a mechanism controlling
spontaneous autoimmunity in H2-O KO mice.
In summary, both scRNA-seq and FACs analyses showed that H2-O KO mice have an
increased frequency of both conventional and regulatory CD4 T cells in the spleens of
unimmunized mice. Loss of thymic H2-O expression correlated with an increased tTreg
population that have received increased TCR stimulation which suggests increased
autoreactivity. Furthermore, increased activation of conventional splenic CD4 T cells is likely a
driving factor for the increased effector-like Treg status that was identified by scRNA-seq
analyses of H2-O-KO CD4 cells.
202
203   **Discussion**
In this study, we have refined our understanding of how loss of H2-O alters CD4 T cell
activity *in vivo*. Thymic analysis of medulla localized CD4 T cells pointed to a novel role for H2-O

on CD4 T cells undergoing self-auditing. Importantly, lack of H2-O expression led to an
increased frequency of both thymic and peripheral Tregs.
Tregs play a fundamental role in maintenance of homeostasis, hence a better
understanding of the mechanism of their development is highly desirable and the subject of
numerous studies (13, 18). Our studies here are the first attempt in demonstrating that H2-O is
a critical player in Treg development. While the exact mechanism by which thymic Treg cells
are selected remains somewhat uncertain, recent studies support the idea that intermittent
TCR signaling along with cytokine signaling drive Treg development (14). These findings and
others (28, 29, 30, 31, 32) support a more avidity based model of Treg selection, in which
alterations to the density of self-ligands present in the medulla will have differential effects on
CD4 T cell selection. Intermittent TCR signaling as the driving force in Treg selection is very
relevant to our understanding of how HLA-DO contributes to epitope selection during antigen
processing. HLA-DO in complex with HLA-DM leads to better refinement of the epitopes from
denatured antigens, promoting selection of the best fitting epitopes in the groove of MHC II
molecules (9, 33). As such, when DO is present, higher affinity peptides are more likely to be
presented. Conversely, in the absence of DO a larger percent of lower affinity epitopes are
selected. Indeed, we have recently reported that peptides eluted from H2-O KO mice
expressing either murine I-A^b (15) or human HLA-DR1 (Welsh et al, unpublished data) were of a
lower general affinity. Accordingly, presentation of a larger portion of lower affinity peptides in
the thymic medulla of H2-O KO mice is expected. Intermittent TCR signaling is typically
generated by either pMHC II that are less stable (34, 35, 36, 37), or TCR/pMHC of lower affinity
(38). Our findings here support the former. We associate increased Treg development as a
consequence of the presentation of lower affinity peptides to self-auditing CD4 T cells.
Due to the continuous egress of mature CD4 T cells from the thymus, we also analyzed
splenic CD4 T cells from unimmunized H2-O WT and H2-O KO mice. H2-O KO splenic CD4 T cells
in unimmunized mice revealed an increased frequency of CD44hiCD69+ antigen experienced
CD4 T cells, which suggests an increased level of antigen-specific TCR signaling in H2-O KO mice.
As previously implied (39), a simple explanation for this activated state is incomplete thymic
deletion in H2-O KO mice. While we were able to show that H2-O KO mice failed to delete
specific CD4 clonal populations (15), detection of differences at the global levels did not show
significant changes between the two genotypes. Nonetheless, we suggest that presentations of
different arrays of self-antigens as well as their lower densities on thymic medulla leads to a
less than optimal thymic deletion of self-reactive CD4 T cells and their routing to the periphery.
Similar to activated CD4 T cells in periphery, H2-O KO Treg cells also had a more effector-like
phenotype, indicating enhanced Treg activation in unimmunized mice (27, 40).
In conclusion, our studies add a new dimension to our understanding of the role of H2-O
in both CD4 T cell selection and activation. For the first time, we report on thymic negative
selection in H2-O KO mice and demonstrate that loss of H2-O enhances thymic selection of
regulatory T cells. Once in the periphery, an increased proportion of H2-O KO Tregs appear to
be activated in a TCR dependent manner. These effector-like Tregs will likely help control
increased basal CD4 T cell activation. However, the exact mechanism by which increased Treg
cells are selected in the H2-O KO thymus remains to be determined. While we propose that
alterations in medulla pMHCII-TCR avidity interactions could lead to enhanced Treg selection,

increased presentation of self-antigens in a more tissue restricted antigenic (TRA) manner could
also be possible (13).
251
252
Methods
Mice
Male and female C57BL/6J (H2-O WT, stock no: 000664), Female B6.PL-Thy1a/CyJ (stock
no: 000406) were purchased from Jackson Laboratories and bred in house. Generation of the
H2-O knock-out mice has been previously described (41) and mice bred in house. All mouse
procedures were approved by the Johns Hopkins University Animal Care and Use Committee
and were following relevant ethical regulations.
260
Antibodies/reagents
Flow Cytometry: anti-CD3e (17A2), anti-CD4 (RM4-5), anti-CD5 (53-7.3), anti-CD8α (53-6.7),
CD11c (N418), anti-CD19 (ID3), anti-CD25 (PC61), anti-CD44 (IM7), anti-CD45R/B220 (RA3-6B2),
anti-CD62L (MEL-14), anti-CD69 (H1.2F3), anti-CD197/CCR7 (4B12), anti-NK1.1 (PK136), anti-
TCRβ (H57-597), anti-TCRγδ (GL3), anti-Cleaved Caspase 3 (D3E9) Cell signaling (Danvers, MA);
Foxp3 (150D), anti-Helios (22F6), anti-Nur77 (12.14), Fixable viability dye eFluor 780
(eBioscience).
268
269 **FACs Sorting**: anti-CD3e, anti-CD4, anti-CD19, anti-NK1.1, Propidium iodide. Briefly, 30,000 live
270 CD4 T cells [Gating: CD3+, CD4, CD19-, NK1.1-, PI-] were FACs sorted from the spleens of 3
271 unimmunized 10 week-old H2-O WT and 3 unimmunized 10 week-old unimmunized H2-O KO
272 female mice.
273
Cell staining
For cleaved caspase 3 staining (17) homogenized mice thymocyte cells were stained
with anti-CCR7/CD197 at a final dilution of 1:50 for 30 min at 37˚C prior to additional surface
stains. Following surface staining, cells were fixed with Cytofix/Cytoperm (BD Biosciences) for
20 min at 4˚C. Cells were then washed with Perm/Wash buffer (BD Biosciences) twice. Cells
were stained with anti–cleaved caspase 3 at a 1:50 dilution at 23˚C for 30 min.
For transcription factor staining, cells were incubated with surface antibody at 4˚C for 20
min, permeabilized at 4˚C for 30 min using a Foxp3/Transcription factor buffer set (Invitrogen,
ThermoFisher Scientific), and then stained with anti–Foxp3 and/or anti-Helios at 23˚C for 30
min.
284
Adoptive Cell Transfer
Naïve CD4 T cells were isolated from the pooled spleens and LNs of 4 week old Thy1.1
expressing C57BL/6J mice using the EasySep Mouse Naive CD4 T cell Isolation Kit (StemCell
Technologies). Isolated cells were stained with eFluor 450 viability dye (eBiosciences) according
to manufactures directions. $3x10^6$ dye labeled Thy1.1 cells were IV injected into 10 week old
H2-O WT or H2-O KO hosts. Transferred cells were recovered 7 days post-transfer.
291

scRNA-Sequencing

Library & sequencing

The samples were prepared using the 10x Genomics Chromium Next GEM Single Cell 5' Library and Gel Bead Kit v1.1, Chromium Next GEM Chip and Dual Index Kit TN Set A. They were run on the Illumina NovaSeq6000 with a run configuration of 28bp x 10bp x 10bp x 91bp.

Analysis

Analyses of T-cell scRNA-seq were performed with the package Seurat (42), as follows. Data was filtered to remove cells with low gene count (<200), large number of UMIs (>12,000) and high (>5%) fraction of mitochondrial reads. Expression levels of genes were log-normalized, and the most variable 2000 genes were selected for linear dimensionality reduction with Principal Component Analysis (PCA). The first 15 principal components were then used to performed unsupervised clustering using the Seurat SNN clustering package, with a resolution of 0.2. To identify cell types, potential markers for each cluster were calculated as the set of genes significantly differentially expressed in each cluster compared to all others, using the function FindAllMarkers, and by searching existing literature and marker databases. Lastly, differentially expressed genes for each cell type between conditions were determined using the function FindMarkers with the default function (bimod).

Defining clusters – Genes with an average fold-change (avgFC) >0.5 and an adjusted p-value <0.05 were run against CD4 T cell data sets available on the Immunological Genome Project (https://www.immgen.org/) using the "My Geneset" data browser function.

Statistical testing

GraphPad Prism was used for all statistical analyses. A standard Student T-test was used for estimation of statistical significance. Data is shown as mean ± SEM. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Acknowledgments

Supported by grants from NIAID, R01AI063764, R21AI101987, and R01AI120634, to SS-N. FACs analysis was performed in the BD Immunology and Flow Cytometry Laboratory at the Johns Hopkins School of Public Health.

Bibliography

1.   Murphy K, Weaver C. Janeway's Immunobiology. New York & London: Garland Science;
2016 2016.

2.   Nishizuka Y, Sakakura T. Thymus and reproduction: sex-linked dysgenesia of the gonad
after neonatal thymectomy in mice. Science. 1969;166(3906):753-5.

3.   Karlsson L, Surh CD, Sprent J, Peterson PA. A novel class II MHC molecule with unusual
tissue distribution. Nature. 1991;351(6326):485-8.

4.   Narayan K, Su KW, Chou CL, Khoruzhenko S, Sadegh-Nasseri S. HLA-DM mediates
peptide exchange by interacting transiently and repeatedly with HLA-DR1. Mol Immunol.
2009;46(15):3157-62.

5.   Chou CL, Mirshahidi S, Su KW, Kim A, Narayan K, Khoruzhenko S, et al. Short peptide
sequences mimic HLA-DM functions. Mol Immunol. 2008;45(7):1935-43.

6.   Natarajan SK, Assadi M, Sadegh-Nasseri S. Stable peptide binding to MHC class II
molecule is rapid and is determined by a receptive conformation shaped by prior association
with low affinity peptides. J Immunol. 1999;162(7):4030-6.

7.   Sadegh-Nasseri S. A step-by-step overview of the dynamic process of epitope selection
by major histocompatibility complex class II for presentation to helper T cells. F1000Res.
2016;5.

8.   Alfonso C, Williams GS, Han JO, Westberg JA, Winqvist O, Karlsson L. Analysis of H2-O
influence on antigen presentation by B cells. J Immunol. 2003;171(5):2331-7.

9.   Poluektov YO, Kim A, Hartman IZ, Sadegh-Nasseri S. HLA-DO as the optimizer of epitope
selection for MHC class II antigen presentation. PloS One. 2013;8(8):e71228.

10.   Savage PA, Klawon DEJ, Miller CH. Regulatory T Cell Development. Annu Rev Immunol.
2020;38:421-53.

11.   Hsieh CS, Lee HM, Lio CW. Selection of regulatory T cells in the thymus. Nat Rev
Immunol. 2012;12(3):157-67.

12.   Tai X, Erman B, Alag A, Mu J, Kimura M, Katz G, et al. Foxp3 transcription factor is
proapoptotic and lethal to developing regulatory T cells unless counterbalanced by cytokine
survival signals. Immunity. 2013;38(6):1116-28.

13.   Klein L, Robey EA, Hsieh CS. Central CD4(+) T cell tolerance: deletion versus regulatory T
cell differentiation. Nat Rev Immunol. 2019;19(1):7-18.

14.   Tai X, Indart A, Rojano M, Guo J, Apenes N, Kadakia T, et al. How autoreactive
thymocytes differentiate into regulatory versus effector CD4(+) T cells after avoiding clonal
deletion. Nat Immunol. 2023;24(4):637-51.

15.   Welsh RA, Song N, Foss CA, Boronina T, Cole RN, Sadegh-Nasseri S. Lack of the MHC
class II chaperone H2-O causes susceptibility to autoimmune diseases. PLoS Biol.
2020;18(2):e3000590.

16.   Nanaware PP, Jurewicz MM, Leszyk J, Shaffer SA, Stern LJ. HLA-DO modulates the
diversity of the MHC-II self-peptidome. Mol Cell Proteomics. 2018.

17.   Breed ER, Watanabe M, Hogquist KA. Measuring Thymic Clonal Deletion at the
Population Level. J Immunol. 2019;202(11):3226-33.

18.   Li MO, Rudensky AY. T cell receptor signalling in the control of regulatory T cell
differentiation and function. Nat Rev Immunol. 2016;16(4):220-33.

19.   Moran AE, Holzapfel KL, Xing Y, Cunningham NR, Maltzman JS, Punt J, et al. T cell
receptor signal strength in Treg and iNKT cell development demonstrated by a novel
fluorescent reporter mouse. J Exp Med. 2011;208(6):1279-89.

20.   Golubovskaya V, Wu L. Different Subsets of T Cells, Memory, Effector Functions, and
CAR-T Immunotherapy. Cancers (Basel). 2016;8(3).

21.   Welsh R, Sadegh-nasseri S. The love and hate relationship of HLA-DM/DO in the
selection of immunodominant epitopes. Curr Opin Immunol. 2020;64:117–23.

22.   Lam AJ, Uday P, Gillies JK, Levings MK. Helios is a marker, not a driver, of human Treg
stability. Eur J Immunol. 2022;52(1):75-84.

23.   Thornton AM, Korty PE, Tran DQ, Wohlfert EA, Murray PE, Belkaid Y, et al. Expression of
Helios, an Ikaros transcription factor family member, differentiates thymic-derived from
peripherally induced Foxp3+ T regulatory cells. J Immunol. 2010;184(7):3433-41.

24.   Thornton AM, Lu J, Korty PE, Kim YC, Martens C, Sun PD, et al. Helios(+) and Helios(-)
Treg subpopulations are phenotypically and functionally distinct and express dissimilar TCR
repertoires. Eur J Immunol. 2019;49(3):398-412.

25.   Serre K, Benezech C, Desanti G, Bobat S, Toellner KM, Bird R, et al. Helios is associated
with CD4 T cells differentiating to T helper 2 and follicular helper T cells in vivo independently of
Foxp3 expression. PLoS One. 2011;6(6):e20731.

26.   Akimova T, Beier UH, Wang L, Levine MH, Hancock WW. Helios expression is a marker of
T cell activation and proliferation. PLoS One. 2011;6(8):e24226.

27.   Miragaia RJ, Gomes T, Chomka A, Jardine L, Riedel A, Hegazy AN, et al. Single-Cell
Transcriptomics of Regulatory T Cells Reveals Trajectories of Tissue Adaptation. Immunity.
2019;50(2):493-504 e7.

28.   Simons DM, Picca CC, Oh S, Perng OA, Aitken M, Erikson J, et al. How specificity for self-
peptides shapes the development and function of regulatory T cells. J Leukoc Biol.
2010;88(6):1099-107.

29.   Relland LM, Mishra MK, Haribhai D, Edwards B, Ziegelbauer J, Williams CB. Affinity-
based selection of regulatory T cells occurs independent of agonist-mediated induction of
Foxp3 expression. J Immunol. 2009;182(3):1341-50.

30.   Picca CC, Oh S, Panarey L, Aitken M, Basehoar A, Caton AJ. Thymocyte deletion can bias
Treg formation toward low-abundance self-peptide. Eur J Immunol. 2009;39(12):3301-6.

31. Feuerer M, Jiang W, Holler PD, Satpathy A, Campbell C, Bogue M, et al. Enhanced thymic
selection of FoxP3+ regulatory T cells in the NOD mouse model of autoimmune diabetes. Proc
Natl Acad Sci U S A. 2007;104(46):18181-6.

32. Atibalentja DF, Murphy KM, Unanue ER. Functional Redundancy between Thymic
CD8{alpha}+ and Sirp{alpha}+ Conventional Dendritic Cells in Presentation of Blood-Derived
Lysozyme by MHC Class II Proteins. J Immunol. 2011;186(3):1421-31.

33. Poluektov YO, Kim A, Sadegh-Nasseri S. HLA-DO and Its Role in MHC Class II Antigen
Presentation. Frontiers in immunology. 2013;4:260.

34. Mirshahidi S, Ferris LC, Sadegh-Nasseri S. The magnitude of TCR engagement is a critical
predictor of T cell anergy or activation. J Immunol. 2004;172(9):5346-55.

35. Sadegh-Nasseri S, Dalai SK, Korb Ferris LC, Mirshahidi S. Suboptimal engagement of the
T-cell receptor by a variety of peptide-MHC ligands triggers T-cell anergy. Immunology.
2010;129(1):1-7.

36. Baumgartner CK, Ferrante A, Nagaoka M, Gorski J, Malherbe LP. Peptide-MHC class II
complex stability governs CD4 T cell clonal selection. J Immunol. 2010;184(2):573-81.

37. Smyth LA, Williams O, Huby RDJ, Norton T, Acuto O, Ley SC, et al. Altered peptide
ligands induce quantitatively but not qualitatively different intracellular signals in primary
thymocytes. PNAS. 1998;95(14):8193-8.

38. Lucas B, Stefanova I, Yasutomo K, Dautigny N, Germain RN. Divergent changes in the
sensitivity of maturing T cells to structurally related ligands underlies formation of a useful T
cell repertoire. Immunity. 1999;10(3):367-76.

39. Gu Y, Jensen PE, Chen X. Immunodeficiency and autoimmunity in H2-O-deficient mice.
Journal of immunology. 2013;190(1):126-37.

40. Shevyrev D, Tereshchenko V. Treg Heterogeneity, Function, and Homeostasis. Front
Immunol. 2019;10:3100.

41. Liljedahl M, Winqvist O, Surh CD, Wong P, Ngo K, Teyton L, et al. Altered antigen
presentation in mice lacking H2-O. Immunity. 1998;8(2):233-43.

42. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al.
Comprehensive Integration of Single-Cell Data. Cell. 2019;177(7):1888-902 e21.

 Figure Legends

Figure 1: Loss of H2-O function supports increased thymic Regulatory T cell development

A. Left: representative gating showing how "Auditing" vs "Clonal Deleted" CCR7+CD4+ T cells

are defined. Middle: Frequency of the activation marker CD69 in either "auditing" medulla

CD4+ T cells or (Right) "clonally deleted" medulla CD4+ T cells.

B. Subdivision of auditing CD4 T cells into three maturation stages: Semi-Mature (Left), Mature

1 (Middle), and Mature 2 (Right). Expression has been normalized to the average H2-O WT

levels within each experiment to allow for comparison across experiments. Data from 5

replicate experiments.

C. Frequency of medulla specific (CCR7+) CD4 T cells selected for clonal deleted (Caspase-3+)

ns = not significant

* = <0.05

** = <0.001

***= <0.0001

**** = <0.00001

Statistics: simple student T-test

Figure 2: DO-KO thymi have an increased proportion of medulla CD4 T cells at the Mature 1

stage

A. The percentage of Foxp3+ CD25+ cells within the CD4 single-positive thymus population in

H2-O WT (white) or H2-O KO (red) cells. Data from 3 replicate experiments

B. Geometric mean fluorescence intensity (gMFI) of CD25 (left), CD5 (right) and Nur77

(Bottom) expressed by Foxp3+ CD4+ T cells in the thymus of H2-O WT (white) or H2-O KO (red)

mice.

C. Subdivision of Treg cells into three maturation stages: Semi-Mature (Left), Mature 1

(Middle), and Mature 2 (Right). Expression has been normalized to the average H2-O WT levels

within each experiment to allow for comparison across experiments. Data from 2 replicate

experiments.

Figure 3: **H2-O KO mice have an increased activated peripheral CD4 T cell population**

A. Percentage of CD4+ T cells within the splenic CD3+ population of unimmunized 6-10-week-

old H2-O WT (white) and H2-O KO (red) mice. Representative of >4 independent experiments, N

= 16 mice per group.

B. Percentage of CD4+ T cells expressing the lymphoid tissue homing receptor CCR7

C. CCR7 and CD62L expression levels in unimmunized splenic CD4 T cells

D. Unimmunized levels of CD44 and CD69 expressed by splenic CD4 T cells. Increased CD69

expression, a marker of recent activation showed increased levels on H2-O KO CD4 T cells

* = <0.05

** = <0.001

*** = <0.0001

**** = <0.00001

Figure 4: **Peripheral H2-O KO Tregs have increased TCR stimulation**
A. Frequency of Foxp3+CD25+ cells in splenic CD4+ T cell population of unimmunized mice
B. H2-O KO peripheral Tregs express decreased levels of naïve (CD44+ CD62L+) expressing cells
C. Normalized Nur77 gMFI levels in Foxp3+CD25+ Treg cells in H2-O WT (white) and H2-O KO
(red) cells. To account for experimental variation the average Nur77 gMFI level in H2-O WT
samples was calculated. gMFI levels in both H2-O WT and H2-O KO samples were then divided
by the calculated H2-O WT average. An increased Nur77 ratio indicates increased Nur77 gMFI
levels. Summary of 3 repeat experiments.
484
Figure 5: **Loss of H2-O function causes increased basal CD4 cell activation**
A. scRNA-seq clustering of CD4 T cells after Seurat analysis. Data represents the average of of 3
biological replicates per genotype.
B. Breakdown of clusters in H2-O WT (Left) or H2-O KO (Right) samples. Clusters are grouped
based upon, (1) known CD4 T cell subset markers and (2) gene comparison to published CD4 T
cell data sets available on the Immunological Genome Project (www.immgen.org). Identified
CD4 Cell phenotypes were: Non-activated, Activated, and Regulatory T cells. "Other" refers to a
minor macrophage and NKT cell contamination from the sorting process.
C. Distribution of the for the Non-activated and Activated phenotypes across the H2-O WT and
H2-O KO biological replicates (N = 3 mice per genotype).
D. In vivo proliferation of adoptively transferred naïve Thy1.1+ CD4 T cells (Thy1.1+ CD3+ CD4+
CD44- CD25-) after 7 days in either Thy1.2 H2-O WT (white) or Thy1.2 H2-O KO (red) hosts.
Pooled data from 2 independent experiments (N= 6 H2-O WT, 5 H2-O KO)
498
Figure 6: **The H2-O KO Treg population has a more effector like phenotype**
A. (Left) distribution of the two Treg populations identified by scRNA-seq clustering in either
H2-O WT (left) or H2-O KO (right) samples. (Right) Expression of key Treg phenotypic markers:
Foxp3 (Top) and CD25 [Il2ra] (Bottom)
B. Subdivision of the splenic CD4+ T cell population by Foxp3 and Helios expression
C. Comparison of cluster 2 upregulated genes (Log$_2$Avg FC >0.5) in H2-O WT and H2-KO samples
to a published (Miragaia *et al*.) splenic effector Treg genetic profile
506
Supplemental Figure 1
A. Representative plots showing CD4, CD8 and Double Positive (DP) percentages in H2-O WT
(Top) and H2-O KO (Bottom)
B. Summary plots of the single-positive CD4 (Top) and CD8 (Bottom) percentages from 5 repeat
experiments.
C. (Left) representative flow plots showing the "Signaled" vs "Non-signaled" thymocytes in H2-
O WT (Top) or H2-O KO (Bottom).
(Right) Summary plots of >5 repeat experiments.
D. Frequency of signaled CCR7+ medulla CD4 T cells in H2-O WT (white) and H2-O KO (red) mice
516
517 **Supplemental Figure 2**:
518 Top: breakdown of H2-O KO (right) and H2-O WT (Left) Seurat clustering
519 Bottom: Seurat clustering for each individual biological replicate

     Figure 1

522 **Figure 2**

A.

Treg (%)

B.

C.

Semi-Mature

Mature 1

Mature 2

524 **Figure 3**

 Figure 4

 Figure 5

Figure 6

 Supplemental Figure 1

A.

B.

C.

D.

534   **Supplemental Figure 2**