Agent-based model demonstrates the impact of nonlinear, complex interactions between cytokines on muscle regeneration

Megan Haase[1], Tien Comlekoglu[1], Alexa Petrucciani[2], Shayn M. Peirce[1], Silvia S. Blemker[1].

[1]University of Virginia, Charlottesville, VA.

[2]Purdue University, West Lafayette, IN

* Correspondence:
Silvia S. Blemker
ssblemker@virginia.edu

Abstract

Muscle regeneration is a complex process due to dynamic and multiscale biochemical and cellular interactions, making it difficult to determine optimal treatments for muscle injury using experimental approaches alone. To understand the degree to which individual cellular behaviors impact endogenous mechanisms of muscle recovery, we developed an agent-based model (ABM) using the Cellular Potts framework to simulate the dynamic microenvironment of a cross-section of murine skeletal muscle tissue. We referenced more than 100 published studies to define over 100 parameters and rules that dictate the behavior of muscle fibers, satellite stem cells (SSC), fibroblasts, neutrophils, macrophages, microvessels, and lymphatic vessels, as well as their interactions with each other and the microenvironment. We utilized parameter density estimation to calibrate the model to temporal biological datasets describing cross-sectional area (CSA) recovery, SSC, and fibroblast cell counts at multiple time points following injury. The calibrated model was validated by comparison of other model outputs (macrophage, neutrophil, and capillaries counts) to experimental observations. Predictions for eight model perturbations that varied cell or cytokine input conditions were compared to published experimental studies to validate model predictive capabilities. We used Latin hypercube sampling and partial rank correlation coefficient to identify *in silico* perturbations of cytokine diffusion coefficients and decay rates to enhance CSA recovery. This analysis suggests that combined alterations of specific cytokine decay and diffusion parameters result in greater fibroblast and SSC proliferation compared to individual perturbations with a 13% increase in CSA recovery compared to unaltered regeneration at 28 days. These results enable guided development of therapeutic strategies that similarly alter muscle physiology (i.e. converting ECM-bound cytokines into freely diffusible forms as studied in cancer therapeutics or delivery of exogenous cytokines) during regeneration to enhance muscle recovery after injury.

Introduction

Skeletal muscle injuries account for more than 30% of all injuries and are one of the most common complaints in orthopedics[1,2,3]. The standard treatment for muscle injuries is limited mostly to rest, ice, compression, elevation, anti-inflammatory drugs, and immobilization[1]. These treatments lack a firm scientific basis and have varied outcomes, some resulting in incomplete functional recovery, formation of scar tissue, and high injury recurrence rates[4,5]. Our fundamental understanding of the individual cellular and subcellular behaviors of muscle cells has advanced and made it clear that interactions between cells and their microenvironment is critical for healthy regeneration. These interactions are dynamic, involve feedback mechanisms, and lead to complex emergent phenomena; therefore, there are numerous possible interventions that could enhance muscle regeneration.

Muscle regeneration requires an abundance of cells and cytokines to interact in a highly coordinated mechanism involving five interrelated cascading phases including degeneration, inflammation, regeneration, remodeling, and functional recovery[6]. Following an acute muscle injury, there is a time-dependent recruitment of neutrophils, monocytes, and macrophages to remove necrotic tissue and release factors that regulate fibroblast behavior and SSC activation, proliferation, and division[7]. Following initial inflammatory response, fibroblasts and SSCs activate and proliferate with the macrophages shifting from their pro to anti-inflammatory phenotype. In healthy muscle, this process would be followed by remodeling of the muscle where the fibroblasts apoptose and SSCs differentiate and fuse to repair the myofibers[8]. Each cell involved in this process secretes cytokines that help regulate cell recruitment and chemotaxes to modulate the dynamics of the recovery. It has also been shown that the molecular events implicated in angiogenesis occur at early stages of muscle regeneration to restore microvascular networks that are crucial for successful muscle recovery[9].

There are numerous cytokines involved in muscle regeneration, many of which have been

individually studied to examine their influence on muscle regeneration[10]. These cytokines play

key roles in dictating cell behaviors and are major drivers of the regeneration cascade[11]. The

dynamics of these cytokines control many aspects of the microenvironment and altering their

properties to optimize treatments has been proposed in a variety of settings[12]. Testing alterations

in cytokine dynamics experimentally has proven to be complex and expensive due to difficulties

in cytokine identification and quantification as well as confounding factors due to pleiotropic

activities of cytokines and interactions with soluble receptors[13]. These challenges make it difficult

to holistically test different diffusion and decay properties for numerous cytokines[14]. However, if

we could better understand the synergistic effects of alteration in cytokines, we could design a

more effective therapy for treating muscle injury.

There are over a million possible combinations of cytokine alterations, making it unrealistic to

study all combinations with experiments alone. For this reason, an *in silico* approach is needed

to fully explore the possible treatment landscape for muscle injury. Over the last several years,

Agent Based Models (ABM) of muscle regeneration have developed to study muscle

regeneration in a variety of applications[8,15–19]. These models were foundational for exploring the

role of SSCs in a variety of muscle milieus[8,15,17] and for demonstrating how ABMs can be used

to simulate therapeutic interventions[16]. However, previous models employed simplistic, non-

spatial representations of cytokine behaviors and properties, which limited their ability to

recapitulate cytokine alterations such as injection of TGF-β[19]. Furthermore, these prior models

did not include microvessel adaptations and dynamic ECM properties which are crucial for

understanding the altered microenvironmental state following muscle injury. These critical

limitations must be addressed in order for ABMs of muscle regeneration to provide meaningful

insights into treatments for muscle injury.

The goals of this work were to: 1) develop an ABM of muscle regeneration that includes cellular

and cytokine spatial dynamics as well as the microvascular environment, 2) calibrate the model

to capture cell behaviors from published experimental studies, 3) validate model outcomes by

comparison with multiple published experimental studies, 4) conduct *in silico* experiments to

predict how altering cytokine dynamics impacts muscle regeneration. For model calibration we

implemented an iterative and robust parameter density estimation protocol to refine the

parameter space and calibrate to temporal biological datasets[20]. Partial rank correlation

coefficient (PRCC) was used to guide *in silico* experiments by identifying parameters and

timepoints that were most critical for ideal regeneration metrics.

Methods

Agent-Based Model Development Overview

ABMs represent the behaviors and interactions of autonomous agents, such as cells, which are

governed by literature-derived rules[15,21,22]. Agent-based modeling provides an excellent platform

for studying complex cellular dynamics because they reveal how the interactions between

individual cellular behaviors lead to emergent behaviors in the whole system.

We implemented the ABM in CompuCell3D (version 4.3.1), a Python-based modeling

software[23]. The ABM's code is available for download (https://simtk.org/projects/muscle_regen).

To build the model, we extended upon about 40 rules developed in previous agent based

models of muscle regeneration[8,15,16] in combination with a deep literature search referencing

over 100 published studies to define approximately 100 total rules that dictate the behavior of

fiber cells, SSC, fibroblasts, neutrophils, and macrophages, as well as their interactions with the

microenvironment, including microvasculature remodeling and cytokine diffusion and secretion

(Fig. 1). For a rule to be incorporated into the model, there had to be an established

understanding within the literature supporting the behavior (i.e. multiple studies reporting similar

findings or supported by other reputable publications). When available, we used experimental

data to define the parameters associated with the model rules. There were 52 parameters that

could not be related to known physiological measurement; therefore, these parameters were

calibrated using parameter density estimation which will be described below in *Model*

Calibration. Following calibration of model parameters, separate model outputs were validated

by comparison with experimental data, and various model perturbations were conducted and

compared to literature results. This process allowed us to have confidence in the predictive

capabilities of the model so that we could simulate and predict the sensitivity of muscle

regeneration to changes in cytokines.

Cellular-Potts Modeling Framework

Prior work to construct computational models to represent muscle recovery have used ordinary

differential equations[24] or agent-based modeling software, such as software such as Netlogo[16]

or Repast[19]. While these models have yielded great insights into skeletal muscle damage and

recovery processes, they have limited capacity to represent the spatial diffusion of cytokines

accurately and explicitly throughout the skeletal muscle. The Cellular-Potts model framework[23]

(CPM, also known as the Glazier-Graner-Hogeweg model), proved an ideal choice because it

represents diffusive species in a physically accurate manner while also allowing for logic-based

representation of cellular behavior characteristic of agent-based modeling.

ABM Design

The ABM spatially represents a two-dimensional male murine skeletal muscle fascicle cross-

section of approximately 50 muscle fibers (Fig. 2). The ABM depicts the microenvironment of

the cross-section as well as the spatial migration of cells and diffusion of various cytokines. The

ABM simulates the emergent phenomenon of muscle tissue from an acute injury over the

course of 28 days. The spatial agents in the model include muscle fibers, necrotic muscle

tissues, extracellular matrix (ECM), capillaries, lymphatic vessels, quiescent and activated

fibroblasts, myofibroblasts, quiescent and activated SSCs, myoblasts, myocytes, immature

myotubes, neutrophils, monocytes, resident macrophages, pro-inflammatory macrophages

(M1), and anti-inflammatory macrophages (M2). In addition, the ABM includes seven diffusing

factors, such as hepatocyte growth factor (HGF), monocyte chemoattractant protein-1 (MCP-1),

matrix metalloproteinase-9 (MMP-9), transforming growth factor beta (TGF-β), tumor necrosis

factor-alpha (TNF-α), vascular endothelial growth factor A (VEGF-A), and interleukin 10 (IL-10).

A review of the literature led us to determine that these factors and cytokine isoforms were most

critical for representing the behaviors of each cell during the regeneration cascade[25,26].

The muscle cross-section geometry was created by importing a histology image stained with

laminin α2 into a custom MATLAB script that masked the histology image to distinguish between

the fibers and ECM. The mask was imported into an initialization CC3D script that defined the

muscle fibers, ECM, and microvasculature to specific cell types and generated a PIFF file that

was imported into the ABM as the starting cross-section. The injury is simulated by

stochastically selecting a region within the cross-section to replace the fiber elements with

necrotic elements, where the percentage of CSA damage is an input parameter. When a

threshold of fiber elements within a muscle fiber becomes damaged the entire muscle fiber turns

necrotic and requires clearance. If the damage is below the threshold, only the region of

necrosis must be removed and the SSCs can fuse to the remaining fiber. During model

initialization, the injury criteria can be altered to simulate various degrees of myotoxin injury by

changing the percent of necrotic tissue following injury.

Each Monte Carlo step (mcs) represents a 15-minute timestep, and the model simulations were

run until 28 days post-injury. We used the CC3D feature Flip2DimRatio to control the number of

times the Cellular-Potts algorithm runs per mcs. The cell velocity is limited by how many times

the Cellular-Potts algorithm is run, so we increased the Flip2DimRatio until we could obtain cell

speeds that were consistent with speeds derived from literature sources (Table 7). At each mcs,

the agent behaviors are governed by rules that were derived from experimental data found in

the literature. The behaviors of each agent are based on environmental conditions, such as

nearby cells and cytokine gradients, as well as probability-based rules. As an example, a

capillary located near a damaged fiber has a probability of becoming non-perfused and then

senses the amount of VEGF-A and MMP-9 at its location to decide if the levels are adequate to

induce angiogenesis (Table 6). Model outputs include CSA recovery (sum of total healthy fiber

elements normalized by the initial CSA), capillary and collagen density, cell counts, relative

cytokine abundance, and spatial coordinates of cells and cytokines.

Overview of Agent Behaviors

Simulated behaviors (Fig. 2) of the neutrophils and macrophages include cytokine-dependent

recruitment, chemotaxis, phagocytosis of damaged fibers (neutrophils, monocytes, and M1

macrophages), phagocytosis of apoptotic neutrophils (monocytes and M1 macrophages),

secretion and uptake of cytokines, and apoptosis. The SSC and fibroblast agent behaviors also

include cytokine-dependent recruitment, chemotaxis, secretion and uptake of cytokines, and

apoptosis, in addition to quiescence, activation, division, and differentiation. The biological

intricacy of some cell types, such as SSCs which have a more complex cell cycle and are

regulated by dynamic interplay of intrinsic factors and an array of microenvironmental stimuli,

led to the necessity for adding more rules that govern their behaviors[27]. The SSCs have 33

parameters dictating the 17 agent rules (Table 3), fibroblasts have 27 parameters for 11 agent

rules (Table 4), macrophages have 31 parameters for 15 agent rules (Table 2) , neutrophils

have 18 parameters for 7 agent rules (Table 1), fibers have 18 parameters for 4 agent rules

(Table 5), and microvessels have 22 parameters for 6 agent rules (Table 6).   At each mcs,

cytokines are secreted by agents if certain conditions were met. For cell recruitment, the levels

of recruiting cytokines for each agent are checked, and if the concentration is high enough to

signal cell recruitment (Supplemental Table 1), a new agent is added to the field at the location

of the highest concentration. The agents also undergo chemotaxis by sensing the surrounding

cytokine gradients and move towards higher concentrations of cytokines, binding and removing

that cytokine as they move along it to simulate physical binding of the cytokine to the receptor.

Agents that are in a quiescent state require a certain threshold level of cytokines to become

activated and cannot chemotax, secrete, divide, or differentiate until this threshold is reached.

Our model assumes each unique cell type secretes the same concentration of cytokines per

timestep for all relevant cytokines to drive model agent decisions. Each computational timestep

represents 15 minutes of real-world time. We assume that this is of sufficient resolution to

accurately reproduce immune cell agent behaviors during regeneration.

Neutrophil Agents

Neutrophils are recruited through capillaries to sites of necrotic tissue (Table 1). Neutrophils

move to areas of necrotic tissue with high concentrations of HGF by chemotaxing along the

HGF gradient to reach areas of necrosis[28,29]. Neutrophils phagocytose necrotic tissue and

facilitate remodeling into ECM with low-collagen density. During phagocytosis, neutrophils

secrete MMP-9, MCP-1, and TNF-α[30,28,31,32]. Individual neutrophil agents apoptose after

phagocytosing two necrotic cells (based on calibration) or 12.5 hours after their recruitment[33].

Macrophage Agents

Resident macrophages are distributed randomly throughout the tissue at a ratio of 1 macrophage

per 5 myofibers at model initialization and secrete MCP-1[34] (Table 2). Resident macrophages

chemotax along MCP-1 and HGF chemical gradients and secrete MMP-9, TNF-α, and MCP-1

during simulation[35–40]. After tissue injury, monocytes are recruited through healthy capillary

microvasculature and chemotax along MCP-1, VEGF-A, TGF-β[41,42–44]. Monocytes infiltrate into

the tissue if the MCP-1 concentration is above a specified threshold at a capillary site. Resident

macrophages, monocytes, and the M1 macrophages differentiated from monocytes may

phagocytose areas of necrotic tissue and apoptotic neutrophil agents[45–47]. During phagocytosis,

these agents secrete MMP-9, HGF, TGF-β, and IL-10[48–52].

Monocytes transition to M1 polarized macrophages when the monocyte agent experiences a large

enough TNF-α concentration or if enough time has passed that a predefined transition time

threshold is met. Each monocyte agent at creation has a defined transition time sampled from a

gaussian distribution with mean and standard deviation (SD) set to reproduce literature-defined

populations of M1 macrophages over time[51,53].

M1 macrophages may transition to M2 macrophages if the M1 macrophage agent experiences

an IL-10 concentration that exceeds a threshold value or if the M1 macrophage has phagocytosed

enough to meet a calibrated threshold value (as discussed in *Model Calibration*)[51,54,55]. Following

the transition to the anti-inflammatory phenotype, the M2 macrophages can proliferate, secrete

TGF-β and IL-10, and chemotax along an MCP-1 gradient[38,56,57].

SSC Agents

The model is initialized with 1 quiescent SSC per every 4 fibers and upon injury[58]. Additional

SSCs are recruited based on the amount of HGF, MMP-9, and TGF-β[59–62] (Table 3). For SSC

activation there has to be enough HGF at the location of the quiescent SSC to induce activation[62–

65]. The SSCs also chemotax up the MMP-9 gradient, removing some of the MMP-9 as they move

along it. Activated SSCs can also undergo symmetric or asymmetric division and differentiation

given that the required cytokine signaling is met locally. Activated SSCs differentiated into

myoblasts and myoblasts differentiate into myocytes[66–68]. Myocytes can fuse to other myocytes

to form new myotubes or fuse to fibers as long as the fiber is not fusion incompetent (i.e., fully

necrotic)[27,69–71]. Maturation of myotubes is required for fusion of additional myocytes to the new

fiber[70,72,73]. If the damage signal is not sustained, activated SSCs return to quiescence. If there is

enough TGF-β to induce apoptosis and not enough VEGF-A or macrophages nearby to block it,

the SSC undergoes cell death and leaves the simulation[42,74–76].

Fibroblast Agents

For model initialization, fibroblasts are randomly placed within the ECM at a population size that

is proportional to the number of fibers[77] (Table 4). Fibroblasts are activated based on the

concentration of TGF-β around the fibroblast[78,79]. Fibroblasts include an additional expression in

their effective energy function that directs their migration towards areas of low-density collagen

ECM [80]. Specifically, fibroblasts can form spring-like links to drag them towards areas of low-

density ECM which are implemented with the relation $\lambda_{ij}(l_{ij} - L_{ij})^2$ where λ_{ij} denotes a

hookean spring constant of a link between cells *i* and *j, l* represents the current distance

between the centers of mass between the two cells, and *L* is the target length of the spring-like

link. In addition to the cytokines secreted by fibroblasts (Table 4), collagen is secreted at low-

density collagen ECM[81–86]. Fibroblasts divide when they are near dividing SSCs and can

differentiate into myofibroblasts with extended exposure to TGF-β[77,87,88]. The myofibroblasts can

secrete more collagen regardless of the ECM density[89]. Fibroblasts can undergo apoptosis if

there are adequate levels of TNF-α at the site of the cell but it can be blocked if there is

sufficient TGF-β[90].

ECM Agents

ECM elements surround the fiber elements and are assigned a collagen density parameter which

varies based on the amount of necrotic tissue removed and the extent of fibroblast/myofibroblast

collagen secretion. When necrotic elements are removed, the phagocytosing inflammatory cells

secrete MMP-9s which degrade some of the collagen within that section of the ECM, thereby

causing that element to have a lower collagen density[28]. The collagen density of the ECM alters

the diffusivity of the secreted factors, and fiber placement is dependent on the collagen density

(discussed below). The fibroblasts help rebuild the ECM by secreting collagen on low collagen

density ECM elements[81]. Myofibroblasts can secrete collagen on any ECM element and if

prolonged results in high density collagen elements, representing a fibrotic state.

Fiber and Necrotic Agents

Upon model initialization, a portion of the muscle fiber agents are converted to necrotic fibers

based on the user prescribed injury. Fibers that reach a damaged threshold became fully necrotic

whereas those surrounding the area of necrosis were regenerating but not fully apoptotic cells.

Healthy fiber elements secrete VEGF-A, and necrotic elements secrete HGF and TGF-β[64,91,92]

(Table 5). Phagocytosing agents chemotax along those gradients to clear the necrosis, but before

a new fiber can be deposited, the collagen has to be restored so that there is a scaffold to hold

the fiber in place[34]. Fully necrotic fibers are fusion incompetent and require myocyte-to-myocyte

fusion to form a new myofiber and require maturation before additional myocyte fusion[70,72,73].

Damaged fibers are regenerated by myocytes fusion to the healthy fiber edge[93].

Capillary and Lymphatic Agents

The muscle fascicle environment includes approximately 4 capillaries per fiber and 1 lymphatic

vessel[94,95] (Table 6). The model defines perfused capillaries as capillary agents that can transport

neutrophils and monocytes into the system proportional to the concentration of recruiting

cytokines[29,41]. The lymphatic vessel agents act to drain the nearby cytokines and cells which

removes them from the microenvironment[96]. Capillaries that are neighboring areas of necrosis

become non-perfused and therefore are unable to transport cells into the microenvironment until

regenerated[97]. Angiogenesis can occur as long as there is enough VEGF-A present at the non-

perfused capillary[98]. Similar to published studies, there is an increase in the capillary-to-myofiber

ratio during muscle regeneration, which is due to the formation of new capillary sprouts modulated

in part by local MMP-9 and VEGF-A levels[97,99,100].

Binding, Diffusivity, and Collagen Density

For many of the agent behaviors described above there are associated binding events that play

a role in regulation of the cytokine fields. Any cytokine dependent behavior is coupled with removal

of a portion of that cytokine once the behavior is initiated. For example, upon SSC activation the

amount of HGF required to activate is taken up by the SSC and removed from the cytokine field

to simulate the binding event that led to SSC activation. Similar binding events were modeled for

SSC and fibroblast division and differentiation, macrophage transitions, cell apoptosis, and

chemotaxis along a cytokine gradient.

To estimate the diffusivity of the various cytokines within the ECM equation 5 was used to account

for the combined effects of collagen and glycosaminoglycans (GAGs) which are known to hinder

diffusivity (Table 7).[101]. The expression includes the radius of the cytokine (r_s), the radius of the

fiber (r_f), the volume fraction (ϕ), D and D_∞ are the diffusivities of the cytokines in the polymer

solution and in free solution, respectively.

$$D = D_\infty \left(-\phi^{\frac{1}{2}} \frac{r_s}{r_f}\right)_{collagen} \times \exp\left(-\phi^{\frac{1}{2}} \frac{r_s}{r_f}\right)_{GAG} \tag{5}$$

Throughout the model simulation, the diffusivity is recalculated with the updated collagen volume

fraction, as the collagen density changes throughout the microenvironment. This allows the

changes in collagen density within the ECM to be reflected in the diffusion rate of each of the

cytokines in the model.

Model Calibration

Known parameters were fixed to literature values, and uncertain parameters were calibrated by

comparing simulation outcomes to published experimental data. Calibration data included

published findings from injury models that have synchronous regeneration after tissue necrosis

(i.e., cardiotoxin, notexin, and barium chloride)[99]. Specifically, data that were used to calibrate

the model included time-varying CSA[102], SSC counts[77] , and fibroblast counts[77]. These metrics

were used for calibration because of their key roles in the regeneration of muscle and the

complex interplay between these outputs. Cell count data were normalized by the number of

cells on the day of the experimental peak to allow for comparison between experiments and

simulations. For CSA, experimental fold-change from pre-injury was compared with fold-change

in model-simulated CSA. Model cell counts were normalized by the number of cells at the peak

time point in the experimental data. SSC and fibroblast counts were normalized to day 5.

Neutrophil counts were normalized to day 1. Total macrophage, M1, and M2 counts were

normalized to day 3. The capillaries were normalized to fiber area, as done in the experimental

data.

Initial ranges for the 52 unknown parameters were determined by literature review or by running

the model to test possible upper and lower thresholds for parameters (Supplemental Table 1).

To narrow the parameter ranges beyond those initial ranges, we used a recently published

calibration protocol, CaliPro, which utilizes parameter density estimation to refine parameter

space and calibrate to temporal biological datasets[20]. CaliPro was selected as the calibration

method because it is model-agnostic which allows it to handle the complexities of stochastic

models such as ABMs, selects viable parameter ranges in the setting of a very high-dimensional

parameter space, and circumvents the need for a cost function, a challenge when there are

many objectives, as in our case. Briefly, Latin hypercube sampling (LHS) was used to generate

600 samples which were run in triplicate. These runs were then evaluated against a set of pass

criteria, and the density functions of the passing runs and failing runs were calculated

(Supplemental Table 2). Parameter ranges were narrowed by alternative density subtraction

(ADS), where the new ranges were determined by the smallest and largest parameter values

where the density of passing is higher than the density of failing. The sensitivity of the model

outputs to the parameters was examined using LHS in combination with partial rank correlation

coefficient (PRCC)[103]. LHS/PRCC methods have been used for various differential equation

models and ABMs[104]. PRCC was computed using MATLAB to determine the correlation

between ABM parameters (i.e. cytokine threshold for activation) and the ABM output (i.e.

fibroblast cell count). Correlations with a p-value less than 0.05 were assumed to be statistically

significant. This helped refine initial parameter bounds as well as make model adjustments

based on the parameter dynamics elucidated from PRCC. This process of sampling parameter

ranges, evaluating the model, and narrowing parameter ranges was repeated in an iterative

fashion while updating pass criteria until a parameter set was identified that consistently met the

strictest criteria (Supplemental Fig. 1). The final passing criteria were set to be within 1 SD for

CSA recovery and 2.5 SD for SSC and fibroblast count. These criteria were selected so that the

model followed experimental trends and accounted for both model stochasticity and

experimental variability in datasets that had narrower SDs for certain timepoints. Following 8

iterations of narrowing the parameter space with CaliPro, we reached a set that had fewer

passing runs than the previous iteration. We then returned to the runs from the prior iteration

and set the bounds such that all 3 runs from the parameter set fell within the final passing

criteria. The final parameter set was run 100 times to verify that the variation from the stochastic

nature of the rules did not cause output that was inconsistent with experimental trends.

Model Validation

We compared model outputs (M1, M2, and total macrophage counts[99,105], neutrophil counts[106],

and capillary counts[102]) that were kept separate from the calibration criteria with published

experimental data to verify that these outputs followed trends from the experimental data

without requiring extra model tuning. In addition, we also altered various model input conditions

(cell input conditions, cytokine dynamics, and microvessel dynamics) to simulate an array of

model perturbations (Table 8) which allowed comparison of a set of model outputs with separate

published experiments. For example, we simulated an IL-10 knockout condition by adjusting the

diffusion and decay parameters so that the concentration of IL-10 throughout the simulation was

zero. One hundred replicates of each model perturbation were performed, and perturbation

outputs were compared with control simulation outputs via a two-sample t-test with a

significance level of 0.05. We were then able to compare how the model outputs aligned with

published experimental findings to determine if the model could capture the altered regeneration

dynamics.

Sensitivity Analysis

A sensitivity analysis was performed using LHS-PRCC to examine the impact of cytokine-

related parameters on model outputs of interest. Diffusion coefficients and decay rates for the

seven cytokines (HGF, TGF-β, MMP-9, TNF-α, VEGF-A, IL-10, MCP-1) were sampled across a

range from 0.1 to 10 times the calibrated value while holding the other parameters constant.

Three hundred samples were generated, and these parameter sets were simulated in triplicate.

PRCCs were calculated with $\alpha=0.05$ and a Bonferroni correction for the number of tests every

10 ticks/hours for CSA and cell counts for SSCs, fibroblasts, non-perfused capillaries,

myoblasts, myocytes, neutrophils, M1 macrophages, and M2 macrophages.

In Silico Experiments

To gain insight into the recovery response with altered angiogenesis, we simulated different

levels of VEGF-A injections and conditions of hindered angiogenesis in which damaged

capillaries were unable to reperfuse following injury (n = 100 for each simulation condition).

Simulations were also conducted to examine correlations between cytokines and their impact on

various cell behaviors and regeneration outcomes. Next, a sensitivity analysis was performed to

understand how alterations in cytokines influence key metrics of regeneration. LHS-PRCC was

used to quantify the impact of cytokine-related parameters (i.e. diffusion rates and decay

coefficients) on outputs of interest (CSA, SSC, fibroblasts, non-perfused capillaries, myoblasts,

myocytes, neutrophils, M1, and M2). A single time point for each output is summarized in Table

9, and these were chosen at the timepoint when PRCC values were peaking, with complete

results available in Supplementary Figure 2.

This sensitivity analysis was then used to guide *in silico* experiments based on which cytokine

parameters promoted favorable regeneration outcomes (i.e. improved recovery, fewer non-

perfused capillaries, increased SSCs). Following individual cytokine parameter alterations, we

combined the cytokine alterations based on beneficial outcomes from the initial *in silico*

experiments to determine if the benefits would be cumulative.

Results

ABM outputs align with calibration and validation data

Following parameter density-based calibration, the unknown parameters were narrowed into a

final calibration parameter set (Supplemental table 1). The simulations captured SSC and

fibroblast cellular behaviors, as well as CSA outcomes, that aligned with experimental studies

(Fig. 3A-C). The model data were consistent with the experimental trends, and the 95%

confidence interval was within the SD for all calibration data time points except for SSCs at day

3 (Fig. 3B). Macrophage (total, M1, and M2), neutrophil, and capillary counts, which were not

used for model calibration, were found to also be consistent with experimental trends and

allowed us to independently validate model outputs (Fig. 3D-H).

ABM perturbations are consistent with published experiments

The model reproduced published findings for a variety of altered conditions that lead to both

improved and diminished muscle regeneration (Fig. 4). For example, injections of VEGF-A led

to faster CSA recovery, more damaged tissue clearance, and a concentration dependent dose

response but with cell depletion there was an overall decrease in all markers of

regeneration[74,107,108]. When simulating hindered angiogenesis conditions, the model aligned

with experimental studies showing detriments in CSA recovery, increased neutrophil and

macrophage cells, and elevated ECM collagen density, indicating progression of fibrosis within

the microenvironment[109].

Analysis of ABM perturbations leads to new insights regarding cytokine and cell

dynamics

The model allowed for new insights into the dynamics of muscle regeneration by providing

additional timepoints and metrics to evaluate the response to exogenous delivery of VEGF-A

and hindered angiogenesis. VEGF-A levels remained elevated compared to control simulations

following the injection at day 5 post injury (Fig. 5A). CSA recovery had the highest increase at

28 days post injury with the high (10^3) relative concentration delivered) VEGF-A injection

followed by the extra high (2×10^3) relative concentration delivered) injection (Fig. 5B). The

medium (750 relative concentration delivered) and low (500 relative concentration delivered)

VEGF-A injections had higher CSA recovery 15 days post injury but were not significantly

different from the control at day 28. All VEGF-A injections had a higher capillary count and were

proportional to the level of VEGF-A injection (Fig. 5C). The impact of VEGF-A injection on peak

SSC and fibroblast counts was dependent on dosage amount, with the extra high VEGF-A

injection resulting in the largest peaks (Fig. 5E&F). Cytokine concentration trends were similar

for all injections, but most peak levels were dosage dependent (Fig. 5G-L). In contrast, HGF

levels were elevated from day 5 to day 28 with hindered angiogenesis, as were TGF-β and IL-

10 (Fig. 5I & L). MCP-1 concentration had a lower overall peak level with elevated levels from

days 21 to 28 (Fig. 5H). Hindered angiogenesis had lower CSA recovery throughout the

simulation and did not achieve unaltered regeneration levels (Fig. 5B).

Cytokine knockout perturbations revealed crosstalk and temporal interplay between cytokines

(Fig. 6). For example, with MCP-1 KO there was an overall increase in cytokine levels for all

other cytokines within the microenvironment except for VEGF-A at 12 hours post injury (Fig.

6A). By 7 days post injury TNF-α, TGF-β, IL-10, and MMP-9 had decreased from unaltered

regeneration day 7 levels but VEGF-A and HGF were elevated. With TNF-α KO there was a

decrease in TGF-β at early timepoints but a strong increase by day 28 (Fig. 6B). Following IL-10

KO there was an increase in TNF-α that peaked at 7 days post injury (Fig. 6C). HGF was

slightly decreased throughout and TGF-β was strongly decreased by day 7. MMP-9 was

decreased at 12 hours and 28 days post injury but heavily increased at day 7.

Cytokine dynamic analysis leads to new model perturbations that predict improved

regeneration

LHS-PRCC of cytokine decay and diffusion parameters elucidated temporal relationships

between cytokine parameters and key regeneration metrics, such as positive correlations

between CSA and TGF-β and MMP-9 decay (Table 9). Of all cytokine parameters, the model

outputs were most sensitive to HGF decay, with all outputs expect M1 cell count being

significantly impacted. PRCC plots showed that TGF-β and MMP decay were positively

correlated and HGF decay was negatively correlated with CSA recovery, with higher

significance at timepoints after 12 days (Supplemental Fig 2). Correlation plots for various

cytokine concentrations and regeneration metrics showed trends in cytokine dependent cell

behaviors such as the TNF-α concentration that led to heightened fibroblast cell counts and

thresholds above that diminish fibroblast response (Supplemental Fig 3). These PRCC trends

guided cytokine parameter perturbations to include lower HGF and VEGF-A decay, higher TGF-

β, MMP-9, and MCP-1 decay, and higher MCP-1 diffusion because each of the cytokine

modifications indicated some form of enhanced regeneration outcome metrics (Supplemental

Table 3). All these perturbations except MCP-1 decay show increased CSA, increased healthy

capillaries, and increased SSCs (Fig. 8). Finally, a combination of all changes except for MCP-1

decay was simulated. The combined cytokine alteration resulted in the highest CSA recovery

(Fig. 8A), as well as increased M1 macrophage counts (Fig. 8B), decreased M2 macrophage

counts (Fig. 8C), increased fibroblasts (Fig. 8D) and SSCs cell counts (Fig. 8E). Capillaries

regenerated faster in the combined perturbation than at unaltered (Fig. 8F). The combined

cytokine perturbation predicted a 13% improvement in CSA recovery compared to the unaltered

regeneration amount at 28. The combined cytokine perturbation also had higher peaks in SSC

and fibroblast counts than any of the singular cytokine perturbations, indicating the additive

effects of altering the cytokine dynamics in combination.

Discussion

We developed a novel ABM that recapitulates muscle regeneration and, unique from prior work,

includes spatial interactions between cytokines and the microvasculature based on relevant

literature[8,15,16]. Model predictions aligned with experimental data under various altered inputs.

Through *in silico* experiments, we gained new insight into how the combination of key cytokine

dynamic alterations could increase SSC cells and enhance CSA recovery. The ability for altered

cytokine concentrations to change regeneration outcomes is consistent with studies that have

found enhanced muscle recovery with delivery of platelet-rich plasma (PRP) which contain

VEGF and TGF- β[110]. These model perturbations allow development of hypotheses for new

experiments and potential therapeutic interventions such as delivery of cytokines to enhance

muscle recovery.

ABM provides biological insight on nonlinear effects of cytokine levels

The ABM offers valuable insights into the muscle regeneration dynamics under various altered

conditions, elucidating the complex interplay of cytokines, angiogenesis, and cell behaviors.

Systematic simulations reveal critical thresholds, nonlinear effects, and synergistic cytokine

combinations impacting regeneration. Perturbations varying VEGF-A injection doses shows

increased CSA recovery up to a threshold (high VEGF-A injection simulation), beyond which

further improvements in CSA recovery cease. Relationships between cytokines and cellular

outputs exhibit nonlinear effects, as seen with the limited impact of elevated HGF on CSA

recovery beyond a threshold and the non-monotonic relationship between TNF-α and fibroblast

counts (Supplemental Fig. 3). Further analysis revealed that specific combinations of cytokine

perturbations could enhance regeneration beyond singular cytokine interventions. For example,

a combined intervention of: 1) decreasing HGF and VEGF-A decay, 2) increasing TGF-β and

MMP-9 decay, and 3) increasing MCP-1 diffusion enhanced muscle regeneration. Prior studies

have shown that individually, HGF[111], VEGF-A[74], and MCP-1[112] stimulate muscle regeneration

whereas reduced TGF-β[113] and MMP-9[114] activity improve recovery. The model suggests that

combined alterations have a stronger regenerative effect than individual cytokine changes,

enhancing muscle recovery through distinct mechanisms—increasing healthy capillaries, SSC

counts, and reducing inflammatory cells.

Cytokine modifications intended to enhance muscle recovery can have clinical relevance and

have been studied in various settings. For example, synthetic biomaterials coated with IL-4 have

been implanted as a cytokine delivery vehicle and were successful in increasing M2 cells within

the muscle[115]. Cytokine antagonist have been successful at promoting muscle regeneration,

seen in prior work with anti-IL-6[116]. Studies have also shown that activation of plasmin is able to

induce the release of ECM bound VEGF, increasing angiogenesis[14,117]. Due to the complex

network of cytokines, studies that deliver simple modulation of one or two cytokines typically

have an insufficient response to generate appreciable improvements. This suggests that using a

combination of biologic and synthetic biomaterials to modulate multiple cytokines is necessary,

which aligns with our findings[115]. Multiple cytokines have been modulated through the use of

PRP which contains VEGF-A and an array of other cytokines, but PRP has had mixed success

in a clinical setting[118]. Our model could test and optimize combinations of cytokines, guiding

future experiments and treatments.

Advancements from prior muscle regeneration models

Previous studies have employed computational models to investigate muscle regeneration

across diverse contexts, such as Duchenne muscular dystrophy and volumetric muscle loss[8,15].

Earlier muscle regeneration ABMs from our group have been used to test the effects of priming

muscle with inflammatory cells prior to injury[16]. While these models laid the foundation for

simulating muscle adaptations, they were constrained by limited diffusion capabilities and an

absence of critical features related to microvessel growth and remodeling throughout the

regeneration process. Similarly, other ABMs from our group have examined altered

microenvironments, but their omission of spatial cytokine diffusion hindered comprehensive

representation of cell behaviors pivotal to regeneration[8,15]. Recently, new ABMs have been

published that focus on cerebral palsy and the impact of injury type on eccentric contraction-

induced damage[17,18]. These models incorporate the effects of some cytokines but have

simplified spatial behaviors that do not incorporate cytokine-specific diffusion and decay or

dynamic changes in cytokine diffusion as a function of varied ECM properties. In addition, these

models were not robustly calibrated or validated against experimental data and lack

microvasculature components. Our model predictions are consistent with these prior models,

though the added biological complexity our model led several new important insights. For

example, in the hindered angiogenesis simulation there was a decrease in SSCs which was

paired with poor CSA recovery similar to how lower SSC counts resulted in lower CSA recovery

in perturbations in both healthy and DMD simulations[15]. Our model provides additional

understanding about the corresponding spatial cytokine changes that ultimately result in

modulation of SSC dynamics within the microenvironment. The additional model advancements

incorporated address prior muscle regeneration modeling gaps in understanding of how

angiogenesis alters recovery outcomes as well as the response of complex spatial cell and

cytokine dynamics.

The computational model introduced in this paper provides three key advancements from

previous models of muscle regenerations[8,15–18]. First is the incorporation of microvasculature,

second is the spatial interactions of cytokines, and third is the use of advanced calibration and

sensitivity analysis techniques. The addition of microvessel growth and remodeling dynamics

empowers investigations into how interventions impact angiogenesis during regeneration,

thereby influencing muscle recovery outcomes. By considering the intricate relationship

between microvessels and regeneration, our model opens avenues for evaluating the effects of

interventions on the broader recovery process. Secondly, understanding how cytokines

influence cell behaviors at different times during regeneration is crucial for determining optimal

treatment targets and dosing. While cytokine dynamics can be altered experimentally, doing so

is expensive and time consuming[12,14] so exploring many combinations of alterations would be

practically infeasible. Our model incorporates decay and diffusion dynamics of a subset of

cytokines to allow testing of far more alterations in cytokines than would be reasonable to

conduct experimentally. Lastly, we leveraged the CaliPro technique for parameter density

estimation-based calibration and LHS-PRCC to gain biological insight by analyzing how altered

microenvironmental parameters could benefit regeneration outcomes. This approach of

implementing parameter identification to guide model perturbations demonstrates the

capabilities of the model as a novel tool for generating new hypotheses and identifying

mechanisms to target for enhanced regeneration outcomes.

Limitations and future work

There are some important limitations of this study that should be discussed. First, the model

does not include all cytokines that are known to influence muscle regeneration, such as IGF,

FGF, PDGF, and additional isoforms of IL. These cytokines likely have redundant functions,

given the model effectively captures relevant cell behaviors using the included cytokines.

Second, while this model does not currently consider hypertrophy during regeneration, which

restricts CSA recovery from surpassing 100%, the cell dynamics it portrays remain consistent

with those observed in studies that lead to hypertrophy following injury. By integrating datasets

from diverse sources, we capture the intricate dynamics of muscle regeneration, while

acknowledging inherent limitations arising from variations in sample sizes and experimental

techniques across sources. Lastly, the current model was calibrated to male mice data despite

known sex difference in skeletal muscle, regeneration mechanisms, and the timeline of

recovery[119–121]. Data on female muscle regeneration is fairly limited because most muscle injury

studies only use male mice or do not distinguish between sexes, making it difficult to incorporate

sex differences into the model[122].

This paper describes a significant advancement in modeling the complex process of muscle

regeneration. Future efforts will extend the use of parameter density estimation to optimize the

selection, doses, and timing of injections of exogenously delivered cytokines. Additionally, we

aim to explore diverse muscle injury types and their varying recovery responses, addressing

challenges in comparing different acute injury techniques found in the literature. This study

underscores the significance of cellular and cytokine spatial dynamics in muscle regeneration.

Further inclusion of additional factors and hormones would provide a more holistic

understanding of the system and how treatments may be altered based on microenvironmental

conditions, providing a unique framework for the study of personalized muscle injury treatment.

586   **Acknowledgements**

587   The authors acknowledge NIH Grant #R21AR080415, Wu-Tsai Foundation Agility Project
588   Funding, and the National Science Foundation Graduate Research Fellowship for financially
589   supporting this research and to James Glazier and T.J. Sego for providing technical support
590   with the CC3D ABM platform.

Figure 1. Flowchart of ABM rules. The model starts with initialization of the geometry and the prescribed injury. This is followed by recruitment of cells based on relative cytokine amounts within the microenvironment. The inflammatory cells, SSCs, and fibroblasts follow their literature defined rules and probability-based decision tree to govern their behaviors. The boxes represent the behavior that the agent completes during that timestep given the appropriate conditions and the circles represent the uptake that occurs as a result of the simulated binding with microenvironmental factors for certain cell behaviors. ABM, agent-based model; SSC, satellite stem cell; ECM, extracellular matrix; TGF-β, transforming growth factor β; HGF, hepatocyte growth factor; TNF-α , tumor necrosis factor α; VEGF-A, vascular endothelial growth factor A; MMP-9, matrix metalloproteinase-9; MCP-1, monocyte chemoattractant protein-1; IL-10; interleukin 10.

Figure 2. Overview of ABM simulation of muscle regeneration following an acute injury. **A.** Simulated
cross-sections of a muscle fascicle that was initially defined by spatial geometry from a histology image.
Muscle injury was simulated by replacing a section of the healthy fibers with necrotic elements. In
response to the injury, a variety of factors are secreted in the microenvironment which impacts the
behavior of the cells. The colors correspond with those typically seen in H&E staining. **B.** ABM screen
captures show the spatial locations of the cells throughout the 28-day simulation. The agent colors were
matched to those typically seen in IHC-stained muscle sections.

Table 1. Neutrophil Agent Rules

Neutrophil Agent Behavior	Sources
Recruitment signal: necrosis	28
Neutrophils are brought to site of injury via capillaries	29
Phagocytose necrosis	30
Secretes MMP-9, MCP-1, TNF-α during phagocytosis	16,28,31,32
Undergoes apoptosis after phagocytosis or 12.5 hours	33
Migrates towards areas of high HGF	39
Migration speed ~7.5 µm/min	123,124

Table 2. Macrophage Agent Rules

Macrophage Agent Behavior	Sources
Initial count: 1 resident macrophages per 5 myofibers	34
Recruitment signal: MCP-1	38,42
Monocytes are brought to the site of injury via microvessels	41
Resident macrophages secrete MMP-9, MCP-1, and TNF-α and chemotax along MCP-1 and HGF	35–40
Monocytes chemotax along MCP-1, VEGF-A, and TGF-β	42–44,125

Monocyte migration speed ~4 μm/min	126
M1 macrophages secrete VEGF-A, MMP-9, and TNF-α and chemotax along MCP-1	127–130
Monocytes, resident and M1 macrophages phagocytose apoptotic neutrophils and necrosis	45–47
Monocytes and macrophages secrete MMP-9, HGF, TGF-β, and IL-10 during phagocytosis	16,48–52
Monocyte transitions to M1 occurs when TNF-α threshold is met or based on literature means and standard deviation properties	51,53
M1 transition into M2 is mediated by the amount of IL-10 and the amount the M1 has phagocytosed	16,51,54,55
M2 macrophages secrete TGF-β and IL-10 and chemotax along MCP-1	16,38,56,131
Macrophages can proliferate following the transition to the anti-inflammatory (M2) state	57
Macrophage migration speed ~0.62 μm/min	126
Macrophages apoptose in a Poisson distribution	132

619 **Table 3.** SSC Agent Rules

SSC Agent Behavior	Sources
Initial count: 1 SSC per 4 fibers	15,58
Recruitment signal: HGF + MMP-9 - TGF-β	15,59–62
Activation signal: HGF	15,62–65
Activated SSCs secrete MCP-1 and VEGF-A	42
Activated SSCs migrate towards areas of high MMP-9	60,133
Myoblasts migrate towards high TNF-α	134
Division signal: TNF-α + VEGF-A - TGF-β	15,61,135,136
Differentiation signal: 3*IL-10 - HGF – TNF-α - TGF-β	15,54,137–139
Activated SSCs differentiate to myoblasts, myoblasts to myocytes, and myocytes to myotubes/myofibers	66–68
Differentiated myocytes fuse at damaged fiber edge or fuse together to from new, immature myotubes	27,69–71
50% cell divisions are symmetric, 50% asymmetric	15,140,141
Division probability decreases with each cell division; 1st division 85%; 2nd 65%; 3rd 20%	15,142
VEGF-A and macrophages nearby can block apoptosis	42,74,75
TGF-β triggers apoptosis	76
Time to divide: 10 hrs	15,142,143
Migration speed ~0.94 μm/min	144
Return activated SSCs to quiescence without sustained HGF	62

621 **Table 4.** Fibroblast Agent Rules

Fibroblast Agent Behavior	Sources
Initial count: 1 fibroblast per every 2 fibers	15,77
Activation signal: TGF-β	145
Fibroblasts move to low collagen ECM	15,80
Fibroblasts secrete TNF-α, TGF-β, MMP-9, VEGF-A. Collagen is secreted at low density ECM	15,81–86

Fibroblast division signaled by SSC division	[15,77]
Division probability decreases with each cell division; 1st division 100%; 2nd 25%; 3rd 6%	[146]
Fibroblast differentiation to myofibroblasts with extended TGF-β exposure	[15,87,88]
Myofibroblasts secrete double the amount of collagen and secretion is not dependent on collagen density	[15,89]
Fibroblasts apoptose with sustained exposure to TNF-α	[15,90]
Fibroblast migration speed ~0.73 µm/min	[147]
Sufficient TGF-β can block fibroblast apoptosis	[15,90]

623 **Table 5.** Fiber Agent Rules

Fiber Agent Behavior	Sources
Damaged muscle fibers secrete HGF and TGF-β	[64,91]
Healthy fibers secrete VEGF-A	[92]
Fibers that are fully necrotic are fusion incompetent, but damaged fibers are fusion competent	[93]
Immature myotubes gain functional capacity as they fully mature overtime	[70,72,73]

625 **Table 6.** Microvasculature Rules

Microvessel Agent Behavior	Sources
Initial count: ~4 capillaries per fiber, 1 lymphatic vessel per fascicle	[94,95]
Capillaries near necrosis will become damaged and unable to perfuse	[97]
With sufficient VEGF-A damaged capillaries will undergo angiogenesis	[98]
MMP-9 is elevated during capillary growth	[100,148]
Increasing capillary to myofiber ratio during muscle regeneration from new sprouting capillaries at areas with enough MMP-9 and VEGF-A	[97,99,100]
Cells and cytokines near lymphatic vessel will be drained via the vessel and removed from microenvironment	[96]

627 **Table 7.** Model parameters of spatial mechanisms

Parameter	Value	Source/Justification
Volume parameters		
Target volume (V_{target}) neutrophil	12 µm^2	Chosen for an average cell diameter of 12 µm[149]
Target volume (V_{target}) SSC	10 µm^2	Chosen for an average cell diameter of 10 µm[150]
Target volume (V_{target}) macrophage	21 µm^2	Chosen for an average cell diameter of 21 µm[151]
Target volume (V_{target}) monocyte	8.5 µm^2	Chosen for an average cell diameter of 8.5 µm[152]
Target volume (V_{target}) fibroblast	15 µm^2	Chosen for an average cell diameter of 15 µm[153]
Volume multiplier (λ_{volume})	50	Volume constraint to maintain target[23]
Diffusion coefficients		

HGF	66.38 μm^2/s	Estimated diffusivity within the ECM accounting for baseline GAGs and collagen[101]
MMP-9	63.40 μm^2/s	
MCP-1	189.27 μm^2/s	
VEGF-A	112.10 μm^2/s	
TGF-β	90.33 μm^2/s	
TNF-α	138.95 μm^2/s	
IL-10	135.17 μm^2/s	
Chemotaxis parameters λ_c		
Neutrophils	750	Chosen for a cell velocity between 1-20 µm/min[123]
Macrophage	9.3	Chosen for a cell velocity around 0.62 µm/min[126]
Monocyte	75	Chosen for a cell velocity around 4 µm/min[126]
SSC	11.3	Chosen for a cell velocity around 0.94 µm/min[144]
Fibroblast	23	Chosen for a cell velocity around 0.73 µm/min[8]

629   **Table 8.** Model perturbation input conditions and corresponding published experimental results

Perturbation	Specific model conditions	Published outcomes
IL-10 knockout	Adjust diffusion and decay parameters so IL-10 is removed from the system	Attenuates shift to M2, hinders differentiation[52]
Neutrophil depletion	Lower neutrophil recruitment proportion	Abundant necrotic tissue 7 days post injury[107]
Macrophage depletion	Lower macrophage recruitment proportion	Decreased HGF, increased TGF-β and TNF-α, impaired regeneration[108]
MCP-1 Knockout	Adjust diffusion and decay parameters so MCP-1 is removed from the system	Increased necrosis at day 7, lower CSA at day 21, impaired phagocytosis[112]
Directed M2 polarization (Anti-inflammatory nanoparticles)	Require less phagocytosis and IL-10 for transition	Improved muscle histology and inflammatory resolution[154]
TNF-α Knockout	Adjust diffusion and decay parameters so TNF-α is removed from the system	Impaired recovery at days 5 and 12, increased inflammation[155]
Hindered angiogenesis	Increase VEGF-A and MMP-9 threshold required for angiogenesis	Delayed regeneration with toxin injury, and persistent immune cell infiltration with freeze injury[109]

| VEGF-A injection | Add VEGF-A at specified concentration (100 for low and 1000 relative concentration for high), radius (300 pixels), and timepoint (5 days post injury) | Lower injury area at day 20 post injury with injection 5 days after damage[74] |

Table 9. Summary of cytokine sensitivity analysis. Significance was determined with α=0.05, and a Bonferroni correction for the number of tests. + and - represent statistically significant positive and negative correlations, respectively.

	CSA	SSC	Fibroblasts	Non-perfused capillaries	Myoblasts	Myocytes	Neutrophils	M1	M2
Day	16.7	6.3	10.5	8.4	6.3	8.4	8.4	4.2	6.3
HGF decay	-	-	-	+	-	-	+		+
TGF-β decay	+	+	+	-	+				-
MMP-9 decay	+	+	+	-	+	+			
TNF-α decay			+						-
VEGF-A decay				+					
MCP-1 decay								+	+
MCP-1 diffusion		+		-				+	

Model Calibration

Model Validation

Figure 3. ABM calibration and validation. ABM parameters were calibrated so that model outputs for CSA recovery, SSC, and fibroblast counts were consistent with experimental data (A-C)[77,102]. Separate outputs from those used in calibration were compared to experimental data[99,102,105,106] to validate the ABM (D-H). Error bars represent experimental standard deviation, and model 95% confidence interval is indicated by the shaded region. Cell count data were normalized by number of cells on the day of the experimental peak to allow for comparison between experiments and simulations.

Figure 4. ABM perturbation outputs are compared to various literature experimental results. Each perturbation model output is compared to the available corresponding published result. The top triangles indicate the literature findings and the bottom triangles indicate the model outputs. Red triangles represent a decrease, blue represents an increase, and gray represents no significant change. Time points of comparison were based on which time points were available from published experimental data.

Figure 5. Dose dependent response with VEGF-A injection compared to hindered angiogenesis. VEGF-A concentration response to varied levels of VEGF injection (A). Hindered angiogenesis resulted in slower and overall decreased CSA recovery (B). Capillary count was dependent on VEGF-A injection level (C). Total macrophage count was similar between control and VEGF-A injections perturbations but macrophage count was higher in later time points in the hindered angiogenesis simulation (D). SSC peak varied with VEGF-A injection level and counts were prolonged in the hindered angiogenesis simulations (E). The fibroblast peak was lower for the hindered angiogenesis perturbation and highest with the extra high VEGF-A injection. In contrast to the other simulations, the fibroblast count was trending upwards at later time points in the hindered angiogenesis perturbation (F). HGF levels were consistent between control and VEGF-A injection perturbations but was significantly elevated in the hindered angiogenesis perturbation (G). MCP-1, TGF-β, and IL-10 concentrations were elevated a later stages of regeneration with hindered angiogenesis (H, I, L). TNF-α was elevated with the extra high VEGF-A injection and lower with hindered angiogenesis (J). MMP-9 concentration was lower at the simulation midpoint but elevated at late regeneration stages (K).

Figure 6. Heatmaps of changes in cytokine concentration at various timepoints throughout regeneration
following individual cytokine knockout (KO) demonstrating cross-talk between cytokines. With MCP-1 KO
there was an increase in all cytokines except VEGF-A at 12 hours post injury. Over the course of
regeneration there was continued increasing elevation of HGF, increases in VEGF-A, and TGF-β
decreased at day 7 followed by a strong increase by day 28 post injury (A). In the TNF-α KO simulations,
there was an early decrease in TGF-β that shifts to strong increases by day 28. MMP-9 increased
throughout the duration, HGF and IL-10 were decreased, VEGF-A lagged in the beginning but was
increased during mid to late timepoints (B). Following IL-10 KO there were increases in TNF-α, decreases
in HGF and TGF-β, and elevated MMP-9 at day 7 that decreased by day 28 (C).

Figure 7. Combined alterations of various cytokine dynamics enhance muscle regeneration outcomes.
All tested alterations except higher MCP-1 decay resulted in higher CSA recovery compared to the
control (A). M1 cell count was higher for all perturbations with the highest peaks with increased MCP-1
diffusion and the combined cytokine alteration perturbation (B). Higher MCP-1 decay resulted in the
largest M2 peak and higher MCP-1 diffusion, higher TGF-β decay, and the combined cytokine alteration
had a lower M2 peak than the control (C). Fibroblasts had the largest increase in cell count with the
higher TGF-β decay and the cytokine combination perturbations (D). All perturbations resulted in an
increased in SSC count with the largest increase resulting from the combined cytokine alteration (E). All
perturbations except the combined and higher MMP-9 decay resulted in increased capillaries as a result
of additional capillary sprouts (F).

Supplemental

Supplemental Figure 1. Overview of Calibration Methods. Latin hypercube sampling is used to generate 600 unique parameter sets given starting bounds, each of which was run in triplicate. The simulations were filtered given specified criteria (i.e. fitting within experimental bounds for CSA recovery) and then alternative density subtraction (ADS) was used to narrow in the parameter bounds. Partial rank squared correlation coefficient (PRCC) was used to gain insight into model sensitivity and adjust the bounds in case initial parameter bounds were too wide or too narrow. This method also allowed for model rule execution refinement to correct cases that interfere with the dynamics of other cell types.

Supplemental Table 1. Unknown model parameters calibrated using LHS to recapitulate published literature

Parameter	Description	Range Tested	Value
neutroRecruitmentProportion	Number of neutrophils generated at every recruitment as proportion of necrotic cells	(4, 12)	7.8322
secretionUptakeValNeu	Cytokine secreted by neutrophil per step during and after "eating necrosis" also max uptake value	(20, 80)	37.3001
necrosisThreshold	Neutrophils are recruited as long as the necrosis amount is greater than this parameter	(0.75, 0.98)	0.9202
macroRecruitmentProportion	Recruitment of macrophages is proportional to mean MCP-1	(0.1, 0.7)	0.4701

secretionUptakeValMac	Cytokine secreted by macrophages per step, also max uptake value	(0.3, 95)	48.2090
TNFconverter	Value of TNF-α to convert monocyte to M1	(15, 150)	109.9025
macRecruitThres	Required mean MCP-1 to signal recruitment of macrophage	(0.05, 4.5)	1.4289
macRecruitStop	Required mean TGF-β to signal stop of macrophage recruitment	(0.05, 0.4)	0.1406
IL10TransThreshold	Local IL-10 that can induce M1 to M2 transition	(15, 600)	78.5442
phagoThresholdforNeuM2	Phagocytosis threshold for neutrophil apoptosis or M1 to M2 transition	(0, 8)	2.0706
sscActivationThreshold	Level of HGF to activate SSC	(10, 100)	44.7091
recruitmentProportionSSC	Number of SSC recruited every time step is proportional to sum of HGF + MMP-9 – TGF-β means	(0.08, 1)	0.8213
quiescentProb	Probability of SSC apoptosis vs remaining quiescent	(1, 4)	3.3617
secretionUptakeValSSC	Cytokine secreted by SSC per step when activated also max uptake value	(0.5, 200)	44.4650
sscTGFApoptosisThreshold	Amount of TGF-β to induce apoptosis	(50, 190)	79.1563
sscApoptosisProb	Probability of SSC apoptosis with apoptosis signal	(0.2, 0.99)	0.7651
VEGFblockApop	Amount of VEGF-A required to block SSC apoptosis	(60, 190)	80.6055
SSCdivisionThreshold	TNF-α + VEGF-A – TGF-β has to be greater than this for SSC to divide	(50, 400)	126.6268
SSCdiffThreshold	IL-10 – HGF – TNF-α – TGF-β has to be greater than this for SSC diff	(5, 145)	57.9870
sscDivideProb	Probability of SSC to divide if signal is not present	(0.01, 0.6)	0.0972
sscDiffProb	Probability for SSC to differentiate without signal	(0.06, 1)	0.6136
fuseProb	Probability of SSC fusing even if the collagen amount is still low	(0.06, 1)	0.3
quiescentThreshold	SSC will return to quiescence if HGF is lower than this at their location	(0.2, 100)	36.3109

necrosisSecretionVal	Amount of HGF and TGF-β secreted by necrotic cells per step	(0.0005, 35)	21.9514
VEGFsecretionValFib	VEGF-A secreted by fibers per step	(0.02, 0.1)	0.0498
VEGF_MMPthreshold	Level of VEGF-A to trigger sprouting/elongation from non-perfused capillaries	(98, 800)	469.0449
capVEGFUptakeMax	Max uptake of VEGF-A by capillaries	(8, 32)	23.8174
MMPsecretionValCap	MMP-9 secreted as capillary is regenerating/sprouting	(4, 16)	8.2444
NumFibers	Number of fibers nearby must be higher than this to check for capillary distances	(1, 2)	1.3899
sproutProb	Probability of capillary sprouting behavior	(0.08, 0.32)	0.1566
secretionUptakeValFibro	Cytokine secretion from fibroblasts per mcs also max uptake value	(0.3, 77)	9.0436
collagenSecretionValFibro	Collagen secretion from fibroblasts per timestep when it is above ECM	(0.09, 0.24)	0.1388
distNearSSCdiv	If fibroblast is within this distance of a dividing SSC, it will divide	(5, 65)	35
TGFActivationThreshold	Level of TGF-β required to activate fibroblast	(20, 80)	40.3643
TGFMyoThreshold	Level of sustained TGF-β required to turn fibroblast into myofibroblast	(280, 650)	536.7525
fibroblastTNFApoptosisThreshold	Level of TNF-α to initiate fibroblast apoptosis	(100, 550)	310.5323
TGFBlockApoptosisThreshold	Level of TGF-β required to block TNF-initiated apoptosis	(160, 1000)	848.3706
fibroblastApoptosisProb	probability of fibroblast apoptosis if TNF-α and TGF-β conditions are met	(0.65,.99)	0.8423
IL10decay	Decay rate of IL-10	(0.00046, 0.00184)	0.0012
HGFdecay	Decay rate of HGF	(0.000135, 0.0003)	0.000228
TGFdecay	Decay rate of TGF	(0.0005, 0.05)	0.0242
TNFdecay	Decay rate of TNF	(0.0000682, 0.002260)	0.00076272
MCPdecay	Decay rate of MCP-1	(0.00596, 0.02384)	0.0124
MMPdecay	Decay rate of MMP-9	(0.0012, 0.1)	0.0112

VEGFdecay	Decay rate of VEGF-A	(0.000464, 0.005569)	0.0015
MinCapDist	Minimum distance between capillary and another capillary	(2.5, 40)	9.6060
capPlacementThreshold	Threshold for if a capillary will be placed/distance required between fiber and cap	(8, 45)	34.6346
M2prolifProb	Probability of M2 proliferation behavior occurring	(0.145305, 0.60)	0.2
M1HalfLife	Corresponds with rate of M1 apoptosis	(15, 55)	24.6360
M2HalfLife	Corresponds with rate of M1 apoptosis	(30, 70)	58.1243
reperfusionLag	Probability of capillary regaining perfusion	(0.001, 0.98)	0.1

709 **Supplemental Table 2.** Criteria utilized for CaliPro model calibration

Iteration	Criteria
1	End criteria: SSC, M1 macrophages, M2 macrophage, fibroblasts, and neutrophils must be below max threshold at the final timepoint (threshold held constant for all iterations).
2	SSC, CSA, and end criteria: within 4 times the experimental std for SSC and 1 times CSA std.
3	Fibroblast, SSC, CSA, replicate check, and end criteria: within 2.1 times the experimental std for fibroblasts, 2.5 for SSC, and 1 times CSA std. More than 1 replicate of the parameter set must meet the criteria.
4	Fibroblast, SSC, CSA, and end criteria: within 2.2 times the experimental std for fibroblasts and SSC and 1 times CSA std.
5	Fibroblast, SSC, CSA, and end criteria: within 1.5 times the experimental std for fibroblasts, 1.75 for SSC, and 1 times CSA std.
6	Fibroblast, SSC, CSA, and end criteria: within 1.75 times the experimental std for fibroblasts, 1.5 for SSC, and 1 times CSA std.
7	Fibroblast, SSC, CSA, replication, and end criteria: within 2.5 times the experimental std for fibroblasts and SSC, and 1 times CSA std. All replicates are required to pass for the parameter set to pass.

Supplemental Figure 2. PRCC plots for various model outputs over time to illustrate how the significance of cytokine decay and diffusion parameters varies at different points throughout regeneration. Black dots indicate statistically significant (P < 0.05) correlation for that timepoint. (A) CSA recovery had correlations with HGF, TGF-β, and MMP-9 decay. (B) SSC count was correlated with HGF, TGF-β, MMP-9 decay and MCP-1 and TND diffusion. (C) Fibroblast count was correlated with HGF, TGF-β, MMP-9, and TNF-α decay. (D) HGF, TGF-β, MMP-9, VGEF decay and MCP-1 diffusion were correlated with the number of non-perfused capillaries. (E) Myoblast cell count was correlated with HGF, TGF-β, MMP-9, and IL-10 decay. (F) Myocyte cell count was correlated with HGF, TGF-β, and MMP-9 decay and TNF-α diffusion. (G) HGF and MCP-1 decay as well as MCP-1 diffusion were correlated with neutrophil count. (H) M1 macrophage cell count was correlated with TGF-β, VEGF-A, IL-10, and MCP-1 decay and MCP-1 diffusion. (I) M2 macrophage count was correlated with HGF, TGF-β, MMP-9, TNF-α, VEGF-A, MCP-1 decay and MCP-1 diffusion.

Supplemental Figure 3. Cytokine concentrations are correlated with cell counts and recovery metrics at
various stages of regeneration. There is an optimal MCP-1 concentration that tends to result in higher M1
counts 1 day post injury (A). IL-10 concentration is positively correlated with M2 count 3 days post injury
(B). VEGF-A concentration is negatively correlated with the number of fragmented (non-perfused)
capillaries 5 days post injury (C). Higher TGF-β concentrations tends to result in lower SSC cell count 7
days post injury (D). Fibroblasts cell count is highest at an optimal TNF-α concentration with higher or
lower levels hindering cell count 14 days post injury (E). HGF concentration is positively correlated with
CSA recovery at day 28 post injury but there appears to be a threshold where high HGF is no longer
correlated with increased recovery (F).

Supplemental Table 3. Cytokine perturbations based on PRCC

Perturbation	Original cytokine parameter	Altered cytokine parameter
Decrease HGF decay	0.000228	0.0000776
Increase TGF-β decay	0.0242	0.241899
Increase MMP-9 decay	0.0112	0.101994
Decrease VEGF-A decay	0.0015	0.000249267
Increase MCP-1 diffusion	0.18627	1.76943
Increase MCP-1 decay	0.0124	0.118216

| Combination | Above alterations excluding increased MCP-1 decay |

Supplemental Figure 4. Non-perfused capillaries for each cytokine perturbation. The combined cytokine perturbation had the lowest number of non-perfused capillaries and all other perturbations resulted in less non-perfused capillaries compared to the control.

References

1. Quintero AJ, Wright VJ, Fu FH, Huard J. Stem Cells for the Treatment of Skeletal Muscle Injury Sports injury Stem cells Tissue engineering Fibrosis Regeneration Skeletal muscle. *Clin Sports Med*. 2009;28:1-11. doi:10.1016/j.csm.2008.08.009

2. Barroso GC, Thiele ES. MUSCLE INJURIES IN ATHLETES. *Rev Bras Ortop (English Ed*. 2011;46(4):354-358. doi:10.1016/s2255-4971(15)30245-7

3. Valle X. Clinical practice guide for muscular injuries: epidemiology, diagnosis, treatment and prevention. *Br J Sports Med*. 2011;45(2):e2-e2. doi:10.1136/bjsm.2010.081570.20

4. Järvinen TAH, Järvinen TLN, Kääriäinen M, et al. Muscle injuries: optimising recovery. *Best Pract Res Clin Rheumatol*. 2007;21(2):317-331. doi:10.1016/j.berh.2006.12.004

5. Huard J, Li Y, Fu FH. Muscle injuries and repair: Current trends in research. *J Bone Jt Surg - Ser A*. Published online 2002. doi:10.2106/00004623-200205000-00022

6. Forcina L, Cosentino M, Musarò A. Mechanisms Regulating Muscle Regeneration: Insights into the Interrelated and Time-Dependent Phases of Tissue Healing. *Cells*.

755   2020;9(5). doi:10.3390/cells9051297

7.   Howard EE, Pasiakos SM, Blesso CN, Fussell MA, Rodriguez NR. Divergent Roles of Inflammation in Skeletal Muscle Recovery From Injury. *Front Physiol.* 2020;11:87. doi:10.3389/FPHYS.2020.00087/BIBTEX

8.   Westman AM, Peirce SM, Christ GJ, Blemker SS. Agent-based model provides insight into the mechanisms behind failed regeneration following volumetric muscle loss injury. Merks RMH, ed. *PLOS Comput Biol.* 2021;17(5):e1008937. doi:10.1371/journal.pcbi.1008937

9.   Wagatsuma A. Endogenous expression of angiogenesis-related factors in response to muscle injury. *Mol Cell Biochem 2006 2981.* 2006;298(1):151-159. doi:10.1007/S11010-006-9361-X

10. Chen C-H, Chang C-H, Lee C-H. Usage of Growth Factors in Acute Muscle Injuries. In: *Sports Injuries.* ; 2015:2343-2351. doi:10.1007/978-3-642-36569-0_174

11. Husmann I, Soulet L, Gautron J, Martelly I, Barritault D. Growth factors in skeletal muscle regeneration. *Cytokine Growth Factor Rev.* 1996;7(3):249-258. doi:10.1016/S1359-6101(96)00029-9

12. Itoh Y. Proteolytic modulation of tumor microenvironment signals during cancer progression. *Front Oncol.* 2022;12. doi:10.3389/FONC.2022.935231

13. Ciano-Petersen NL, Muñiz-Castrillo S, Birzu C, et al. Cytokine dynamics and targeted immunotherapies in autoimmune encephalitis. *Brain Commun.* 2022;4(4):1-16. doi:10.1093/braincomms/fcac196

14. Ferrara N. Binding to the Extracellular Matrix and Proteolytic Processing: Two Key Mechanisms Regulating Vascular Endothelial Growth Factor Action. *Mol Biol Cell.* 2010;21(5):687. doi:10.1091/MBC.E09-07-0590

15. Virgilio KM, Martin KS, Peirce SM, Blemker SS. Agent-based model illustrates the role of the microenvironment in regeneration in healthy and mdx skeletal muscle. *J Appl Physiol.* 2018;125(5):1424-1439. doi:10.1152/japplphysiol.00379.2018

16. Martin KS, Virgilio KM, Peirce SM, Blemker SS. Computational modeling of muscle regeneration and adaptation to advance muscle tissue regeneration strategies. *Cells Tissues Organs.* 2016;202(3-4):250-266. doi:10.1159/000443635

17. Khuu S, Fernandez JW, Handsfield GG. A Coupled Mechanobiological Model of Muscle Regeneration In Cerebral Palsy. *Front Bioeng Biotechnol.* 2021;9. doi:10.3389/FBIOE.2021.689714/FULL

18. Khuu S, Fernandez JW, Handsfield GG. Delayed skeletal muscle repair following inflammatory damage in simulated agent-based models of muscle regeneration. Peirce SM, ed. *PLOS Comput Biol.* 2023;19(4):e1011042. doi:10.1371/JOURNAL.PCBI.1011042

19. Virgilio KM, Jones BK, Miller EY, et al. Computational Models Provide Insight into In Vivo Studies and Reveal the Complex Role of Fibrosis in mdx Muscle Regeneration. *Ann Biomed Eng.* Published online 2020. doi:10.1007/s10439-020-02566-1

20. Joslyn LR, Kirschner DE, Linderman JJ. CaliPro: A Calibration Protocol That Utilizes Parameter Density Estimation to Explore Parameter Space and Calibrate Complex

Biological Models. *Cell Mol Bioeng.* 2021;14(1):31-47. doi:10.1007/S12195-020-00650-
Z/FIGURES/7

21.   Martin KS, Blemker SS, Peirce SM. Agent-based computational model investigates
muscle-specific responses to disuse-induced atrophy. *J Appl Physiol.* 2015;118(10):1299-
1309. doi:10.1152/japplphysiol.01150.2014

22.   Gianlupi JF, Mapder T, Sego TJ, et al. Multiscale Model of Antiviral Timing, Potency, and
Heterogeneity Effects on an Epithelial Tissue Patch Infected by SARS-CoV-2. *Viruses
2022, Vol 14, Page 605.* 2022;14(3):605. doi:10.3390/V14030605

23.   Swat MH, Thomas GL, Belmonte JM, Shirinifard A, Hmeljak D, Glazier JA. Multi-Scale
Modeling of Tissues Using CompuCell3D. *Methods Cell Biol.* 2012;110:325.
doi:10.1016/B978-0-12-388403-9.00013-8

24.   Stephenson ER, Kojouharov H V. A mathematical model of skeletal muscle regeneration.
Math Methods Appl Sci. 2018;41(18):8589-8602. doi:10.1002/mma.4908

25.   Waldemer-Streyer RJ, Kim D, Chen J. Muscle cell-derived cytokines in skeletal muscle
regeneration. *FEBS J.* 2022;289(21):6463-6483. doi:10.1111/febs.16372

26.   Rucavado A, Escalante T, Teixeira CFP, Fernándes CM, Díaz C, Gutiérrez JM.
Increments in cytokines and matrix metalloproteinases in skeletal muscle after injection of
tissue-damaging toxins from the venom of the snake Bothrops asper. *Mediators Inflamm.*
2002;11(2):121-128. doi:10.1080/09629350220131980

27.   Yin H, Price F, Rudnicki MA. Satellite cells and the muscle stem cell niche. *Physiol Rev.*
2013;93(1):23-67. doi:10.1152/PHYSREV.00043.2011

28.   Madaro L, Bouché M. From innate to adaptive immune response in muscular dystrophies
and skeletal muscle regeneration: The role of lymphocytes. *Biomed Res Int.* 2014;2014.
doi:10.1155/2014/438675

29.   Wang X, Hossain M, Bogoslowski A, Kubes P, Irimia D. Chemotaxing neutrophils enter
alternate branches at capillary bifurcations. *Nat Commun 2020 111.* 2020;11(1):1-12.
doi:10.1038/s41467-020-15476-6

30.   Butterfield TA, Best TM, Merrick MA. The Dual Roles of Neutrophils and Macrophages in
Inflammation: A Critical Balance Between Tissue Damage and Repair. *J Athl Train.*
2006;41(4):457. doi:10.1016/s0162-0908(08)79217-1

31.   Wang J. Neutrophils in tissue injury and repair. *Cell Tissue Res.* 2018;371(3):531-539.
doi:10.1007/S00441-017-2785-7/FIGURES/1

32.   Soehnlein O, Zernecke A, Eriksson EE, et al. Neutrophil secretion products pave the way
for inflammatory monocytes. *Blood.* 2008;112(4):1461. doi:10.1182/BLOOD-2008-02-
139634

33.   Fox S, Leitch AE, Duffin R, Haslett C, Rossi AG. Neutrophil Apoptosis: Relevance to the
Innate Immune Response and Inflammatory Disease. *J Innate Immun.* 2010;2(3):216.
doi:10.1159/000284367

34.   Oishi Y, Manabe I. Macrophages in inflammation, repair and regeneration. *Int Immunol.*
2018;30(11):511-528. doi:10.1093/INTIMM/DXY054

35.   Elkington PT, Green JA, Friedland JS. Analysis of matrix metalloproteinase secretion by

838   macrophages. *Methods Mol Biol*. 2009;531:253-265. doi:10.1007/978-1-59745-396-7_16

839   36.   Chen GY, Nuñez G. Sterile inflammation: sensing and reacting to damage. *Nat Rev*
840   *Immunol 2010 1012*. 2010;10(12):826-837. doi:10.1038/nri2873

841   37.   Lacy P, Stow JL. Cytokine release from innate immune cells: association with diverse
842   membrane trafficking pathways. *Blood*. 2011;118(1):9-18. doi:10.1182/BLOOD-2010-08-
843   265892

844   38.   Vogel DYS, Heijnen PDAM, Breur M, et al. Macrophages migrate in an activation-
845   dependent manner to chemokines involved in neuroinflammation. *J Neuroinflammation*.
846   2014;11:23. doi:10.1186/1742-2094-11-23

847   39.   Molnarfi N, Benkhoucha M, Funakoshi H, Nakamura T, Lalive PH. Hepatocyte growth
848   factor: A regulator of inflammation and autoimmunity. *Autoimmun Rev*. 2015;14(4):293-
849   303. doi:10.1016/J.AUTREV.2014.11.013

850   40.   Furrer R, Handschin C. Optimized engagement of macrophages and satellite cells in
851   the repair and regeneration of exercised muscle. *Res Perspect Endocr Interact*.
852   2017;(9783319727899):57-66. doi:10.1007/978-3-319-72790-5_5/FIGURES/1

853   41.   Kratofil RM, Kubes P, Deniset JF. Monocyte conversion during inflammation and injury.
854   *Arterioscler Thromb Vasc Biol*. 2017;37(1):35-42. doi:10.1161/ATVBAHA.116.308198

855   42.   Chazaud B, Sonnet C, Lafuste P, et al. Satellite cells attract monocytes and use
856   macrophages as a support to escape apoptosis and enhance muscle growth. *J Cell Biol*.
857   2003;163(5):1133. doi:10.1083/JCB.200212046

858   43.   Owen JL, Mohamadzadeh M. Macrophages and chemokines as mediators of
859   angiogenesis. *Front Physiol*. 2013;4. doi:10.3389/FPHYS.2013.00159

860   44.   Reibman J, Meixler S, Lee TC, et al. Transforming growth factor beta 1, a potent
861   chemoattractant for human neutrophils, bypasses classic signal-transduction pathways.
862   *Proc Natl Acad Sci U S A*. 1991;88(15):6805. doi:10.1073/PNAS.88.15.6805

863   45.   Greenlee-Wacker MC. Clearance of apoptotic neutrophils and resolution of inflammation.
864   *Immunol Rev*. 2016;273(1):357. doi:10.1111/IMR.12453

865   46.   Watanabe S, Alexander M, Misharin A V., Budinger GRS. The role of macrophages in the
866   resolution of inflammation. *J Clin Invest*. 2019;129(7):2619. doi:10.1172/JCI124615

867   47.   Uribe-Querol E, Rosales C. Phagocytosis: Our Current Understanding of a Universal
868   Biological Process. *Front Immunol*. 2020;11:1066.
869   doi:10.3389/FIMMU.2020.01066/BIBTEX

870   48.   Yoon YS, Lee YJ, Choi YH, Park YM, Kang JL. Macrophages programmed by apoptotic
871   cells inhibit epithelial-mesenchymal transition in lung alveolar epithelial cells via PGE2,
872   PGD2, and HGF. *Sci Reports 2016 61*. 2016;6(1):1-18. doi:10.1038/srep20992

873   49.   D'Angelo F, Bernasconi E, Schäfer M, et al. Macrophages promote epithelial repair
874   through hepatocyte growth factor secretion. *Clin Exp Immunol*. 2013;174(1):60.
875   doi:10.1111/CEI.12157

876   50.   Popov Y, Sverdlov DY, Bhaskar KR, et al. Macrophage-mediated phagocytosis of
877   apoptotic cholangiocytes contributes to reversal of experimental biliary fibrosis. *Am J*
878   *Physiol - Gastrointest Liver Physiol*. 2010;298(3):G323. doi:10.1152/AJPGI.00394.2009

879   51.   Arnold L, Henry A, Poron F, et al. Inflammatory monocytes recruited after skeletal muscle
880   injury switch into antiinflammatory macrophages to support myogenesis. *J Exp Med.*
881   2007;204(5):1057-1069. doi:10.1084/jem.20070075

882   52.   Chung EY, Liu J, Homma Y, et al. Interleukin-10 Expression in Macrophages during
883   Phagocytosis of Apoptotic Cells Is Mediated by the TALE homeoproteins Pbx-1 and
884   Prep-1. *Immunity.* 2007;27(6):952. doi:10.1016/J.IMMUNI.2007.11.014

885   53.   Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev
886   Immunol 2008 812.* 2008;8(12):958-969. doi:10.1038/nri2448

887   54.   Saini J, McPhee JS, Al-Dabbagh S, Stewart CE, Al-Shanti N. Regenerative function of
888   immune system: Modulation of muscle stem cells. *Ageing Res Rev.* 2016;27:67-76.
889   doi:10.1016/J.ARR.2016.03.006

890   55.   Das A, Sinha M, Datta S, et al. Monocyte and Macrophage Plasticity in Tissue Repair and
891   Regeneration. *Am J Pathol.* 2015;185(10):2596. doi:10.1016/J.AJPATH.2015.06.001

892   56.   Arabpour M, Saghazadeh A, Rezaei N. Anti-inflammatory and M2 macrophage
893   polarization-promoting effect of mesenchymal stem cell-derived exosomes. *Int
894   Immunopharmacol.* 2021;97:107823. doi:10.1016/J.INTIMP.2021.107823

895   57.   Arnold L, Henry A, Poron F, et al. Inflammatory monocytes recruited after skeletal muscle
896   injury switch into antiinflammatory macrophages to support myogenesis. *J Exp Med.*
897   2007;204(5):1057-1069. doi:10.1084/JEM.20070075

898   58.   Reimann J, Irintchev A, Wernig A. Regenerative capacity and the number of satellite cells
899   in soleus muscles of normal and mdx mice. *Neuromuscul Disord.* 2000;10(4-5):276-282.
900   doi:10.1016/S0960-8966(99)00118-2

901   59.   Kawamura K, Takano K, Suetsugu S, et al. N-WASP and WAVE2 acting downstream of
902   phosphatidylinositol 3-kinase are required for myogenic cell migration induced by
903   hepatocyte growth factor. *J Biol Chem.* 2004;279(52):54862-54871.
904   doi:10.1074/JBC.M408057200

905   60.   Wang W, Pan H, Murray K, Jefferson BS, Li Y. Matrix metalloproteinase-1 promotes
906   muscle cell migration and differentiation. *Am J Pathol.* 2009;174(2):541-549.
907   doi:10.2353/AJPATH.2009.080509

908   61.   Allen RE, Boxhorn LK. Regulation of skeletal muscle satellite cell proliferation and
909   differentiation by transforming growth factor-beta, insulin-like growth factor I, and
910   fibroblast growth factor. *J Cell Physiol.* 1989;138(2):311-315.
911   doi:10.1002/JCP.1041380213

912   62.   González MN, de Mello W, Butler-Browne GS, et al. HGF potentiates extracellular matrix-
913   driven migration of human myoblasts: Involvement of matrix metalloproteinases and
914   MAPK/ERK pathway. *Skelet Muscle.* 2017;7(1):1-13. doi:10.1186/S13395-017-0138-
915   6/FIGURES/5

916   63.   Allen RE, Sheehan SM, Taylor RG, Kendall TL, Rice GM. Hepatocyte growth factor
917   activates quiescent skeletal muscle satellite cells in vitro. *J Cell Physiol.* 1995;165(2):307-
918   312. doi:10.1002/JCP.1041650211

919   64.   Miller KJ, Thaloor D, Matteson S, Pavlath GK. Hepatocyte growth factor affects satellite
920   cell activation and differentiation in regenerating skeletal muscle. *Am J Physiol Cell
921   Physiol.* 2000;278(1). doi:10.1152/AJPCELL.2000.278.1.C174

65. Tatsumi R, Anderson JE, Nevoret CJ, Halevy O, Allen RE. HGF/SF is present in normal
adult skeletal muscle and is capable of activating satellite cells. *Dev Biol*.
1998;194(1):114-128. doi:10.1006/DBIO.1997.8803

66. Cooper RN, Tajbakhsh S, Mouly V, Cossu G, Buckingham M, Butler-Browne GS. In vivo
satellite cell activation via Myf5 and MyoD in regenerating mouse skeletal muscle. *J Cell
Sci*. 1999;112 (Pt 17)(17):2895-2901. doi:10.1242/JCS.112.17.2895

67. Flamini V, Ghadiali RS, Antczak P, Rothwell A, Turnbull JE, Pisconti A. The Satellite Cell
Niche Regulates the Balance between Myoblast Differentiation and Self-Renewal via p53.
Stem Cell Reports. 2018;10(3):970-983. doi:10.1016/J.STEMCR.2018.01.007

68. Bentzinger CF, Wang YX, Rudnicki MA. Building Muscle: Molecular Regulation of
Myogenesis. *Cold Spring Harb Perspect Biol*. 2012;4(2):a008342.
doi:10.1101/CSHPERSPECT.A008342

69. Wang YX, Dumont NA, Rudnicki MA. Muscle stem cells at a glance. *J Cell Sci*.
2014;127(Pt 21):4543-4548. doi:10.1242/JCS.151209

70. Nguyen JH, Chung JD, Lynch GS, Ryall JG. The Microenvironment Is a Critical Regulator
of Muscle Stem Cell Activation and Proliferation. *Front Cell Dev Biol*. 2019;7:254.
doi:10.3389/FCELL.2019.00254/BIBTEX

71. Ruiz-Gómez M, Coutts N, Suster ML, Landgraf M, Bate M. myoblasts incompetent
encodes a zinc finger transcription factor required to specify fusion-competent myoblasts
in Drosophila. *Development*. 2002;129(1):133-141. doi:10.1242/DEV.129.1.133

72. Abmayr SM, Pavlath GK. Myoblast fusion: lessons from flies and mice. *Development*.
2012;139(4):641-656. doi:10.1242/DEV.068353

73. Isesele PO, Mazurak VC. Regulation of Skeletal Muscle Satellite Cell Differentiation by
Omega-3 Polyunsaturated Fatty Acids: A Critical Review. *Front Physiol*. 2021;12:750.
doi:10.3389/FPHYS.2021.682091/BIBTEX

74. Arsic N, Zacchigna S, Zentilin L, et al. Vascular endothelial growth factor stimulates
skeletal muscle regeneration in Vivo. *Mol Ther*. 2004;10(5):844-854.
doi:10.1016/J.YMTHE.2004.08.007

75. Sonnet C, Lafuste P, Arnold L, et al. Human macrophages rescue myoblasts and
myotubes from apoptosis through a set of adhesion molecular systems. *J Cell Sci*.
2006;119(Pt 12):2497-2507. doi:10.1242/JCS.02988

76. Cencetti F, Bernacchioni C, Tonelli F, Roberts E, Donati C, Bruni P. TGFβ1 evokes
myoblast apoptotic response via a novel signaling pathway involving S1P4 transactivation
upstream of Rho-kinase-2 activation. *FASEB J*. 2013;27(11):4532-4546.
doi:10.1096/FJ.13-228528

77. Murphy MM, Lawson JA, Mathew SJ, Hutcheson DA, Kardon G. Satellite cells,
connective tissue fibroblasts and their interactions are crucial for muscle regeneration.
Development. 2011;138(17):3625-3637. doi:10.1242/dev.064162

78. Beanes SR, Dang C, Soo C, Ting K. Skin repair and scar formation: the central role of
TGF-β. *Expert Rev Mol Med*. 2003;5(8):1-22. doi:10.1017/S1462399403005817

79. Chellini F, Tani A, Zecchi-Orlandini S, Sassoli C. Influence of platelet-rich and platelet-
poor plasma on endogenous mechanisms of skeletal muscle repair/regeneration. *Int J*

Mol Sci. 2019;20(3). doi:10.3390/ijms20030683

80.   Dickinson RB, Guido S, Tranquillo RT. Biased cell migration of fibroblasts exhibiting
contact guidance in oriented collagen gels. *Ann Biomed Eng.* 1994;22(4):342-356.
doi:10.1007/BF02368241

81.   Zou Y, Zhang RZ, Sabatelli P, Chu ML, Bönnemann CG. Muscle interstitial fibroblasts are
the main source of collagen VI synthesis in skeletal muscle: implications for congenital
muscular dystrophy types Ullrich and Bethlem. *J Neuropathol Exp Neurol.*
2008;67(2):144-154. doi:10.1097/NEN.0B013E3181634EF7

82.   Sanderson RD, Fitch JM, Linsenmayer TR, Mayne R. Fibroblasts promote the formation
of a continuous basal lamina during myogenesis in vitro. *J Cell Biol.* 1986;102(3):740.
doi:10.1083/JCB.102.3.740

83.   Yokoyama T, Sekiguchi K, Tanaka T, et al. Angiotensin II and mechanical stretch induce
production of tumor necrosis factor in cardiac fibroblasts. *Am J Physiol.* 1999;276(6).
doi:10.1152/AJPHEART.1999.276.6.H1968

84.   Skutek M, Van Griensven M, Zeichen J, Brauer N, Bosch U. Cyclic mechanical stretching
modulates secretion pattern of growth factors in human tendon fibroblasts. *Eur J Appl*
Physiol. 2001;86(1):48-52. doi:10.1007/S004210100502

85.   Lindner D, Zietsch C, Becher PM, et al. Differential expression of matrix metalloproteases
in human fibroblasts with different origins. *Biochem Res Int.* Published online 2012.
doi:10.1155/2012/875742

86.   Newman AC, Nakatsu MN, Chou W, Gershon PD, Hughes CCW. The requirement for
fibroblasts in angiogenesis: Fibroblast-derived matrix proteins are essential for
endothelial cell lumen formation. *Mol Biol Cell.* 2011;22(20):3791-3800.
doi:10.1091/MBC.E11-05-0393/ASSET/IMAGES/LARGE/3791FIG5.JPEG

87.   Desmouliere A, Geinoz A, Gabbiani F, Gabbiani G. Transforming growth factor-beta 1
induces alpha-smooth muscle actin expression in granulation tissue myofibroblasts and in
quiescent and growing cultured fibroblasts. *J Cell Biol.* 1993;122(1):103-111.
doi:10.1083/JCB.122.1.103

88.   Wipff PJ, Rifkin DB, Meister JJ, Hinz B. Myofibroblast contraction activates latent TGF-
beta1 from the extracellular matrix. *J Cell Biol.* 2007;179(6):1311-1323.
doi:10.1083/JCB.200704042

89.   Petrov V V., Fagard RH, Lijnen PJ. Stimulation of Collagen Production by Transforming
Growth Factor-β1 During Differentiation of Cardiac Fibroblasts to Myofibroblasts.
Hypertension. 2002;39(2 I):258-263. doi:10.1161/HY0202.103268

90.   Lemos DR, Babaeijandaghi F, Low M, et al. Nilotinib reduces muscle fibrosis in chronic
muscle injury by promoting TNF-mediated apoptosis of fibro/adipogenic progenitors. *Nat*
Med. 2015;21(7):786-794. doi:10.1038/NM.3869

91.   Kim J, Lee J. Role of transforming growth factor-β in muscle damage and regeneration:
focused on eccentric muscle contraction. *J Exerc Rehabil.* 2017;13(6):621.
doi:10.12965/JER.1735072.536

92.   Huey KA. Potential Roles of Vascular Endothelial Growth Factor during Skeletal Muscle
Hypertrophy. *Exerc Sport Sci Rev.* 2018;46(3):195-202.
doi:10.1249/JES.0000000000000152

93. Snijders T, Nederveen JP, McKay BR, et al. Satellite cells in human skeletal muscle
plasticity. *Front Physiol*. 2015;6(OCT):283. doi:10.3389/FPHYS.2015.00283/BIBTEX

94. Wickler SJ. Capillary supply of skeletal muscles from acclimatized white-footed mice
Peromyscus. *Am J Physiol*. 1981;241(5):357-361.
doi:10.1152/AJPREGU.1981.241.5.R357

95. Gehlert S, Theis C, Weber S, et al. Exercise-induced decline in the density of lyve-1-
positive lymphatic vessels in human skeletal muscle. *Lymphat Res Biol*. 2010;8(3):165-
173. doi:10.1089/LRB.2009.0035

96. Hampton HR, Chtanova T. Lymphatic Migration of Immune Cells. *Front Immunol*.
2019;10(MAY):1168. doi:10.3389/FIMMU.2019.01168

97. Jacobsen NL, Norton CE, Shaw RL, Cornelison D, Segal S, Segal SS. Myofiber injury
induces capillary disruption and regeneration of disorganized microvascular networks.
bioRxiv. Published online August 4, 2021:2021.08.02.454805.
doi:10.1101/2021.08.02.454805

98. Frey SP, Jansen H, Raschke MJ, Meffert RH, Ochman S. VEGF improves skeletal
muscle regeneration after acute trauma and reconstruction of the limb in a rabbit model.
Clin Orthop Relat Res. 2012;470(12):3607-3614. doi:10.1007/S11999-012-2456-7

99. Hardy D, Besnard A, Latil M, et al. Comparative Study of Injury Models for Studying
Muscle Regeneration in Mice. *PLoS One*. 2016;11(1):e0147198.
doi:10.1371/JOURNAL.PONE.0147198

100. Haas TL, Milkiewicz M, Davis SJ, et al. Matrix metalloproteinase activity is required for
activity-induced angiogenesis in rat skeletal muscle. *Am J Physiol Heart Circ Physiol*.
2000;279(4). doi:10.1152/AJPHEART.2000.279.4.H1540

101. Filion RJ, Popel AS. Intracoronary administration of FGF-2: A computational model of
myocardial deposition and retention. *Am J Physiol - Hear Circ Physiol*. 2005;288(1 57-
1):263-279. doi:10.1152/AJPHEART.00205.2004/SUPPL_FILE/TABLE

102. Ochoa O, Sun D, Reyes-Reyna SM, et al. Delayed angiogenesis and VEGF production in
CCR2-/- mice during impaired skeletal muscle regeneration. *Am J Physiol - Regul Integr*
Comp Physiol. 2007;293(2):651-661.
doi:10.1152/AJPREGU.00069.2007/ASSET/IMAGES/LARGE/ZH60080759050009.JPEG

103. Marino S, Hogue IB, Ray CJ, Kirschner DE. A methodology for performing global
uncertainty and sensitivity analysis in systems biology. *J Theor Biol*. 2008;254(1):178-
196. doi:10.1016/J.JTBI.2008.04.011

104. Segovia-Juarez JL, Ganguli S, Kirschner D. Identifying control mechanisms of granuloma
formation during M. tuberculosis infection using an agent-based model. *J Theor Biol*.
2004;231(3):357-376. doi:10.1016/J.JTBI.2004.06.031

105. Wang X, Zhao W, Ransohoff RM, Zhou L. Infiltrating macrophages are broadly activated
at the early stage to support acute skeletal muscle injury repair. *J Neuroimmunol*.
2018;317:55. doi:10.1016/J.JNEUROIM.2018.01.004

106. Nguyen MH, Cheng M, Koh TJ. Impaired Muscle Regeneration in Ob/ob and Db/db Mice.
Sci World J. 2011;11:1525. doi:10.1100/TSW.2011.137

107. Teixeira CFP, Zamunér SR, Zuliani JP, et al. Neutrophils do not contribute to local tissue

damage, but play a key role in skeletal muscle regeneration, in mice injected with
Bothrops asper snake venom. *Muscle Nerve*. 2003;28(4):449-459.
doi:10.1002/MUS.10453

108.   Liu X, Liu Y, Zhao L, Zeng Z, Xiao W, Chen P. Macrophage depletion impairs skeletal
muscle regeneration: The roles of regulatory factors for muscle regeneration. *Cell Biol Int*.
2017;41(3):228-238. doi:10.1002/CBIN.10705

109.   Hardy D, Fefeu M, Besnard A, et al. Defective angiogenesis in CXCL12 mutant mice
impairs skeletal muscle regeneration. *Skelet Muscle*. 2019;9(1):1-15.
doi:10.1186/S13395-019-0210-5/FIGURES/5

110.   Kunze KN, Hannon CP, Fialkoff JD, Frank RM, Cole BJ. Platelet-rich plasma for muscle
injuries: A systematic review of the basic science literature. *World J Orthop*.
2019;10(7):278-291. doi:10.5312/wjo.v10.i7.278

111.   Choi W, Lee J, Lee J, Lee SH, Kim S. Hepatocyte growth factor regulates macrophage
transition to the M2 phenotype and promotes murine skeletal muscle regeneration. *Front*
Physiol. 2019;10(JUL). doi:10.3389/fphys.2019.00914

112.   Lu H, Huang D, Ransohoff RM, Zhou L. Acute skeletal muscle injury: CCL2 expression
by both monocytes and injured muscle is required for repair. *FASEB J*.
2011;25(10):3344-3355. doi:10.1096/FJ.10-178939

113.   Girardi F, Taleb A, Ebrahimi M, et al. TGFβ signaling curbs cell fusion and muscle
regeneration. *Nat Commun*. 2021;12(1). doi:10.1038/s41467-020-20289-8

114.   Zimowska M, Olszynski KH, Swierczynska M, Streminska W, Ciemerych MA. Decrease
of MMP-9 activity improves soleus muscle regeneration. *Tissue Eng - Part A*.
2012;18(11-12):1183-1192. doi:10.1089/ten.tea.2011.0459

115.   Dziki JL, Velayutham M, Hussey GS, Turnquist HR. Cytokine networks in immune-
mediated muscle regeneration. *J Immunol Regen Med*. 2018;1:32-44.
doi:10.1016/j.regen.2018.03.001

116.   Fujita R, Kawano F, Ohira T, et al. Anti-interleukin-6 receptor antibody (MR16-1)
promotes muscle regeneration via modulation of gene expressions in infiltrated
macrophages. *Biochim Biophys Acta - Gen Subj*. 2014;1840(10):3170-3180.
doi:10.1016/j.bbagen.2014.01.014

117.   Ismail AA, Shaker BT, Bajou K. The plasminogen–activator plasmin system in
physiological and pathophysiological angiogenesis. *Int J Mol Sci*. 2022;23(1).
doi:10.3390/ijms23010337

118.   Alsousou J, Ali A, Willett K, Harrison P. The role of platelet-rich plasma in tissue
regeneration. *Platelets*. 2013;24(3):173-182. doi:10.3109/09537104.2012.684730

119.   Haizlip KM, Harrison BC, Leinwand LA. Sex-based differences in skeletal muscle kinetics
and fiber-type composition. *Physiology*. 2015;30(1):30-39.
doi:10.1152/physiol.00024.2014

120.   Knewtson KE, Ohl NR, Robinson JL. Estrogen Signaling Dictates Musculoskeletal Stem
Cell Behavior: Sex Differences in Tissue Repair. *Tissue Eng - Part B Rev*.
2022;28(4):789-812. doi:10.1089/ten.teb.2021.0094

121.   Liu, Siyu; Kostek, Matthew; Omstead K. Sex based differences in muscle regeneration.

Am J Physiol. 2023;38(S1).

122. Enns DL, Tiidus PM. The influence of estrogen on skeletal muscle: Sex matters. *Sport Med.* 2010;40(1):41-58. doi:10.2165/11319760-000000000-00000

123. Zhao W, Zhao H, Li M, Huang C. Microfluidic devices for neutrophil chemotaxis studies. *J Transl Med.* 2020;18(1):1-19. doi:10.1186/S12967-020-02335-7/FIGURES/5

124. Heit B, Liu L, Colarusso P, Puri KD, Kubes P. PI3K accelerates, but is not required for, neutrophil chemotaxis to fMLP. *J Cell Sci.* 2008;121(2):205-214. doi:10.1242/JCS.020412

125. Martin KS, Kegelman CD, Virgilio KM, et al. In Silico and In Vivo Experiments Reveal M-CSF Injections Accelerate Regeneration Following Muscle Laceration. *Ann Biomed Eng.* 45. doi:10.1007/s10439-016-1707-2

126. Bos E van den, Walbaum S, Horsthemke M, Bachg AC, Hanley PJ. Time-lapse Imaging of Mouse Macrophage Chemotaxis. *JoVE (Journal Vis Exp.* 2020;2020(158):e60750. doi:10.3791/60750

127. Corliss BA, Azimi MS, Munson JM, Peirce SM, Murfee WL. Macrophages: An Inflammatory Link between Angiogenesis and Lymphangiogenesis. *Microcirculation.* 2016;23(2):95. doi:10.1111/MICC.12259

128. Newby AC. Metalloproteinase expression in monocytes and macrophages and its relationship to atherosclerotic plaque instability. *Arterioscler Thromb Vasc Biol.* 2008;28(12):2108-2114. doi:10.1161/ATVBAHA.108.173898

129. Lu HL, Huang XY, Luo YF, Tan WP, Chen PF, Guo YB. Activation of M1 macrophages plays a critical role in the initiation of acute lung injury. *Biosci Rep.* 2018;38(2):20171555. doi:10.1042/BSR20171555

130. Cui K, Ardell CL, Podolnikova NP, Yakubenko VP. Distinct migratory properties of M1, M2, and resident macrophages are regulated by αdβ2and αmβ2integrin-mediated adhesion. *Front Immunol.* 2018;9(NOV):2650. doi:10.3389/FIMMU.2018.02650/BIBTEX

131. da Silva MD, Bobinski F, Sato KL, Kolker SJ, Sluka KA, Santos ARS. IL-10 Cytokine Released from M2 Macrophages Is Crucial for Analgesic and Anti-inflammatory Effects of Acupuncture in a Model of Inflammatory Muscle Pain. *Mol Neurobiol.* 2015;51(1):19. doi:10.1007/S12035-014-8790-X

132. Moncayo AR. Poisson Convergence and Family Trees. *Ann Probab.* 1975;3(6):1059-1061.

133. Xiaoping C, Yong L. Role of matrix metalloproteinases in skeletal muscle: migration, differentiation, regeneration and fibrosis. *Cell Adh Migr.* 2009;3(4). doi:10.4161/CAM.3.4.9338

134. Terrente Y, El Fahime E, Caron NJ, et al. Tumor necrosis factor-alpha (TNF-alpha) stimulates chemotactic response in mouse myogenic cells. *Cell Transplant.* 2003;12(1):91-100. doi:10.3727/000000003783985115

135. Bakkar N, Wang J, Ladner KJ, et al. IKK/NF-κB regulates skeletal myogenesis via a signaling switch to inhibit differentiation and promote mitochondrial biogenesis. *J Cell Biol.* 2008;180(4):787. doi:10.1083/JCB.200707179

136. Saclier M, Yacoub-Youssef H, Mackey AL, et al. Differentially activated macrophages

orchestrate myogenic precursor cell fate during human skeletal muscle regeneration. *Stem Cells*. 2013;31(2):384-396. doi:10.1002/STEM.1288

137. Perandini LA, Chimin P, Lutkemeyer D da S, Câmara NOS. Chronic inflammation in skeletal muscle impairs satellite cells function during regeneration: can physical exercise restore the satellite cell niche? *FEBS J*. 2018;285(11):1973-1984. doi:10.1111/FEBS.14417

138. Gal-Levi R, Leshem Y, Aoki S, Nakamura T, Halevy O. Hepatocyte growth factor plays a dual role in regulating skeletal muscle satellite cell proliferation and differentiation. *Biochim Biophys Acta - Mol Cell Res*. 1998;1402(1):39-51. doi:10.1016/S0167-4889(97)00124-9

139. Ten Broek RW, Grefte S, Von Den Hoff JW. Regulatory factors and cell populations involved in skeletal muscle regeneration. *J Cell Physiol*. 2010;224(1):7-16. doi:10.1002/JCP.22127

140. Kuang S, Kuroda K, Le Grand F, Rudnicki MA. Asymmetric self-renewal and commitment of satellite stem cells in muscle. *Cell*. 2007;129(5):999-1010. doi:10.1016/J.CELL.2007.03.044

141. Yennek S, Burute M, Théry M, Tajbakhsh S. Cell adhesion geometry regulates non-random DNA segregation and asymmetric cell fates in mouse skeletal muscle stem cells. *Cell Rep*. 2014;7(4):961-970. doi:10.1016/J.CELREP.2014.04.016

142. Siegel AL, Kuhlmann PK, Cornelison DDW. Muscle satellite cell proliferation and association: new insights from myofiber time-lapse imaging. *Skelet Muscle*. 2011;1(1). doi:10.1186/2044-5040-1-7

143. Rocheteau P, Gayraud-Morel B, Siegl-Cachedenier I, Blasco MA, Tajbakhsh S. A subpopulation of adult skeletal muscle stem cells retains all template DNA strands after cell division. *Cell*. 2012;148(1-2):112-125. doi:10.1016/J.CELL.2011.11.049

144. Otto A, Collins-Hooper H, Patel A, Dash PR, Patel K. Adult skeletal muscle stem cell migration is mediated by a blebbing/amoeboid mechanism. *Rejuvenation Res*. 2011;14(3):249-260. doi:10.1089/REJ.2010.1151/ASSET/IMAGES/LARGE/FIGURE6.JPEG

145. Gibb AA, Lazaropoulos MP, Elrod JW. Myofibroblasts and Fibrosis. *Circ Res*. 2020;127(3):427-447. doi:10.1161/CIRCRESAHA.120.316958

146. Alberts B, Johnson A, Lewis J, Raff M, Roberts K, Walter P. Extracellular Control of Cell Division, Cell Growth, and Apoptosis. Published online 2002. Accessed March 14, 2023. https://www.ncbi.nlm.nih.gov/books/NBK26877/

147. Cornwell KG, Downing BR, Pins GD. Characterizing fibroblast migration on discrete collagen threads for applications in tissue regeneration. *J Biomed Mater Res Part A*. 2004;71A(1):55-62. doi:10.1002/JBM.A.30132

148. Qutub A, Gabhann F, Karagiannis E, Vempati P, Popel A. Multiscale models of angiogenesis. In: *IEEE Engineering in Medicine and Biology Magazine*. Vol 28. NIH Public Access; 2009:14-31. doi:10.1109/MEMB.2009.931791

149. Tigner A, Ibrahim SA, Murray I. Histology, White Blood Cell. *StatPearls*. Published online November 19, 2021. Accessed May 16, 2022. https://www.ncbi.nlm.nih.gov/books/NBK563148/

1175   150.   Garcia SM, Tamaki S, Lee S, et al. Stem Cell Reports Resource High-Yield Purification,
1176   Preservation, and Serial Transplantation of Human Satellite Cells. Published online 2018.
1177   doi:10.1016/j.stemcr.2018.01.022

1178   151.   Krombach F, Münzing S, Allmeling AM, Gerlach JT, Behr J, Dörger M. Cell size of
1179   alveolar macrophages: an interspecies comparison. *Environ Health Perspect*.
1180   1997;105(Suppl 5):1261. doi:10.1289/EHP.97105S51261

1181   152.   Downey GP, Doherty DE, Schwab B, Elson EL, Henson PM, Worthen GS. Retention of
1182   leukocytes in capillaries: role of cell size and deformability.
1183   *https://doi.org/101152/jappl19906951767*. 1990;69(5):1767-1778.
1184   doi:10.1152/JAPPL.1990.69.5.1767

1185   153.   Freitas RJ. *Nanomedicine: Basic Capabilities* . Vol 1. Landes Bioscience; 1999.
1186   Accessed May 16, 2022. http://www.nanomedicine.com/NMI/8.5.1.htm

1187   154.   Raimondo TM, Mooney DJ. Functional muscle recovery with nanoparticle-directed M2
1188   macrophage polarization in mice. *Proc Natl Acad Sci U S A*. 2018;115(42):10648-10653.
1189   doi:10.1073/PNAS.1806908115/SUPPL_FILE/PNAS.1806908115.SAPP.PDF

1190   155.   Warren GL, Hulderman T, Jensen N, et al. Physiological role of tumor necrosis factor
1191   alpha in traumatic muscle injury. *FASEB J*. 2002;16(12):1630-1632. doi:10.1096/FJ.02-
1192   0187FJE

1193