Agent-based model demonstrates the impact of nonlinear, complex interactions between cytokines on muscle regeneration

Megan Haase[1], Tien Comlekoglu[1], Alexa Petrucciani[2], Shayn M. Peirce[1], Silvia S. Blemker[1].

[1]University of Virginia, Charlottesville, VA.

[2]Purdue University, West Lafayette, IN

* Correspondence:
Silvia S. Blemker
ssblemker@virginia.edu

Abstract

Muscle regeneration is a complex process due to dynamic and multiscale biochemical and cellular interactions, making it difficult to identify microenvironmental conditions that are beneficial to muscle recovery from injury using experimental approaches alone. To understand the degree to which individual cellular behaviors impact endogenous mechanisms of muscle recovery, we developed an agent-based model (ABM) using the Cellular Potts framework to simulate the dynamic microenvironment of a cross-section of murine skeletal muscle tissue. We referenced more than 100 published studies to define over 100 parameters and rules that dictate the behavior of muscle fibers, satellite stem cells (SSC), fibroblasts, neutrophils, macrophages, microvessels, and lymphatic vessels, as well as their interactions with each other and the microenvironment. We utilized parameter density estimation to calibrate the model to temporal biological datasets describing cross-sectional area (CSA) recovery, SSC, and fibroblast cell counts at multiple time points following injury. The calibrated model was validated by comparison of other model outputs (macrophage, neutrophil, and capillaries counts) to experimental observations. Predictions for eight model perturbations that varied cell or cytokine input conditions were compared to published experimental studies to validate model predictive capabilities. We used Latin hypercube sampling and partial rank correlation coefficient to identify *in silico* perturbations of cytokine diffusion coefficients and decay rates to enhance CSA recovery. This analysis suggests that combined alterations of specific cytokine decay and diffusion parameters result in greater fibroblast and SSC proliferation compared to individual perturbations with a 13% increase in CSA recovery compared to unaltered regeneration at 28 days. These results enable guided development of therapeutic strategies that similarly alter muscle physiology (i.e. converting ECM-bound cytokines into freely diffusible forms as studied in cancer therapeutics or delivery of exogenous cytokines) during regeneration to enhance muscle recovery after injury.

Introduction

Skeletal muscle injuries account for more than 30% of all injuries and are one of the most common complaints in orthopedics[1,2,3]. The standard treatment for muscle injuries is limited mostly to rest, ice, compression, elevation, anti-inflammatory drugs, and immobilization[1]. These treatments lack a firm scientific basis and have varied outcomes, some resulting in incomplete functional recovery, formation of scar tissue, and high injury recurrence rates[4,5]. Our fundamental understanding of the individual cellular and subcellular behaviors of muscle cells has advanced and made it clear that interactions between cells and their microenvironment is critical for healthy regeneration. These interactions are dynamic, involve feedback mechanisms, and lead to complex emergent phenomena; therefore, there are numerous possible interventions that could enhance muscle regeneration.

Muscle regeneration requires an abundance of cells and cytokines to interact in a highly coordinated mechanism involving five interrelated cascading phases including: degeneration, inflammation, regeneration, remodeling, and functional recovery[6]. Following an acute muscle injury, there is a time-dependent recruitment of neutrophils, monocytes, and macrophages to remove necrotic tissue and release factors that regulate fibroblast behavior and SSC activation, proliferation, and division[7]. Following initial inflammatory response, fibroblasts and SSCs activate and proliferate with the macrophages shifting from their pro to anti-inflammatory phenotype. In healthy muscle, this process would be followed by remodeling of the muscle where the fibroblasts apoptose and SSCs differentiate and fuse to repair the myofibers[8]. Each cell involved in this process secretes cytokines that help regulate cell recruitment and chemotaxes to modulate the dynamics of the recovery. It has also been shown that the molecular events implicated in angiogenesis occur at early stages of muscle regeneration to restore microvascular networks that are crucial for successful muscle recovery[9].

There are numerous cytokines involved in muscle regeneration, many of which have been

individually studied to examine their influence on muscle regeneration[10]. These cytokines play

key roles in dictating cell behaviors and are major drivers of the regeneration cascade[11]. The

dynamics of these cytokines control many aspects of the microenvironment and altering their

properties to optimize treatments has been proposed in a variety of settings[12]. Testing alterations

in cytokine dynamics experimentally has proven to be complex and expensive due to difficulties

in cytokine identification and quantification as well as confounding factors due to pleiotropic

activities of cytokines and interactions with soluble receptors[13]. These challenges make it difficult

to holistically test different diffusion and decay properties for numerous cytokines[14]. However, if

we could better understand the synergistic effects of alteration in cytokines, we could design a

more effective therapy for treating muscle injury.

There are over a million possible combinations of cytokine alterations, making it unrealistic to

study all combinations with experiments alone. For this reason, an *in silico* approach is needed

to fully explore the possible treatment landscape and make predictions on potential targets to

enhance muscle recovery. Over the last several years, Agent Based Models (ABM) of muscle

regeneration have developed to study muscle regeneration in a variety of applications[8,15–19].

These models were foundational for exploring the role of SSCs in a variety of muscle

milieus[8,15,17] and for demonstrating how ABMs can be used to simulate therapeutic

interventions[16]. However, previous models employed simplistic, non-spatial representations of

cytokine behaviors and properties, which limited their ability to recapitulate cytokine alterations

such as injection of TGF-β[19]. Furthermore, these prior models did not include microvessel

adaptations and dynamic ECM properties which are crucial for understanding the altered

microenvironmental state following muscle injury. These critical limitations must be addressed in

order for ABMs of muscle regeneration to provide meaningful insights into treatments for muscle

injury.

The goals of this work were to: 1) develop an ABM of muscle regeneration that includes cellular

and cytokine spatial dynamics as well as the microvascular environment, 2) calibrate the model

to capture cell behaviors from published experimental studies, 3) validate model outcomes by

comparison with multiple published experimental studies, 4) conduct *in silico* experiments to

predict how altering cytokine dynamics impacts muscle regeneration. For model calibration, we

implemented an iterative and robust parameter density estimation protocol to refine the

parameter space and calibrate to temporal biological datasets[20]. Partial rank correlation

coefficient (PRCC) was used to guide *in silico* experiments by identifying parameters and

timepoints that were most critical for ideal regeneration metrics.

Methods

Agent-Based Model Development Overview

ABMs represent the behaviors and interactions of autonomous agents, such as cells, which are

governed by literature-derived rules[15,21,22]. Agent-based modeling provides an excellent platform

for studying complex cellular dynamics because they reveal how the interactions between

individual cellular behaviors lead to emergent behaviors in the whole system.

We implemented the ABM in CompuCell3D (version 4.3.1), a Python-based modeling

software[23]. The ABM's code is available for download (https://zenodo.org/records/10403014).

To build the model, we extended upon about 40 rules developed in previous agent based

models of muscle regeneration[8,15,16] in combination with a deep literature search referencing

over 100 published studies to define approximately 100 total rules that dictate the behavior of

fiber cells, SSC, fibroblasts, neutrophils, and macrophages, as well as their interactions with the

microenvironment, including microvasculature remodeling and cytokine diffusion and secretion

(Fig. 1). For a rule to be incorporated into the model, there had to be an established

understanding within the literature supporting the behavior (i.e. multiple studies reporting similar

findings or supported by other reputable publications). When available, we used experimental

data to define the parameters associated with the model rules. There were 52 parameters that

could not be related to known physiological measurement; therefore, these parameters were

calibrated using parameter density estimation which will be described below in *Model*

Calibration. Following calibration of model parameters, separate model outputs were validated

by comparison with experimental data, and various model perturbations were conducted and

compared to literature results. This process allowed us to have confidence in the predictive

capabilities of the model so that we could simulate and predict the sensitivity of muscle

regeneration to changes in cytokines.

Cellular-Potts Modeling Framework

Prior work to construct computational models to represent muscle recovery have used ordinary

differential equations[24] or agent-based modeling software, such as software such as Netlogo[16]

or Repast[19]. While these models have yielded great insights into skeletal muscle damage and

recovery processes, they have limited capacity to represent the spatial diffusion of cytokines

accurately and explicitly throughout the skeletal muscle. The Cellular-Potts model framework[23]

(CPM, also known as the Glazier-Graner-Hogeweg model), proved an ideal choice because it

allows for logic-based representation of cellular behavior and interactions characteristic of

agent-based modeling (see Supplemental Text 1 for additional details on CPM).

ABM Design

The ABM spatially represents a two-dimensional male murine skeletal muscle fascicle cross-

section of approximately 50 muscle fibers (Fig. 2A). The ABM depicts the microenvironment of

the cross-section as well as the spatial migration of cells and diffusion of various cytokines. The

ABM simulates the emergent phenomenon of muscle tissue from an acute injury over the

course of 28 days. The spatial agents in the model include muscle fibers, necrotic muscle

tissues, extracellular matrix (ECM), capillaries, lymphatic vessels, quiescent and activated

fibroblasts, myofibroblasts, quiescent and activated SSCs, myoblasts, myocytes, immature

myotubes, neutrophils, monocytes, resident macrophages, pro-inflammatory macrophages

(M1), and anti-inflammatory macrophages (M2). In addition, the ABM includes seven diffusing

factors, such as hepatocyte growth factor (HGF), monocyte chemoattractant protein-1 (MCP-1),

matrix metalloproteinase-9 (MMP-9), transforming growth factor beta (TGF-β), tumor necrosis

factor-alpha (TNF-α), vascular endothelial growth factor A (VEGF-A), and interleukin 10 (IL-10).

A review of the literature led us to determine that these factors and cytokine isoforms were most

critical for representing the behaviors of each cell during the regeneration cascade[25,26].

The muscle cross-section geometry was created by importing a histology image stained with

laminin α2 into a custom MATLAB script that masked the histology image to distinguish between

the fibers and ECM. The mask was imported into an initialization CC3D script that defined the

muscle fibers, ECM, and microvasculature to specific cell types and generated a PIF file that

was imported into the ABM as the starting cross-section. The injury is simulated by

stochastically selecting a region within the cross-section to replace the fiber elements with

necrotic elements, where the percentage of CSA damage is an input parameter. When a

threshold of fiber elements within a muscle fiber becomes damaged the entire muscle fiber turns

necrotic and requires clearance. If the damage is below the threshold, only the region of

necrosis must be removed and the SSCs can fuse to the remaining fiber. During model

initialization, the injury criteria can be altered to simulate various degrees of myotoxin injury by

changing the percent of necrotic tissue following injury.

Each Monte Carlo step (mcs) represents a 15-minute timestep, and the model simulations were

run until 28 days post-injury. The cell velocity is limited by how many times the Cellular-Potts

algorithm is run, so we set 45 Cellular-Potts evaluations per mcs to ensure stability in migratory

agent behavior. The number of Cellular-Potts evaluations per mcs and the lambda chemotaxis

parameters were tuned in a simplified simulation of individual cells and their respective

chemotactic gradients so we could obtain cell speeds that were consistent with speeds derived

from literature sources (Table 7). At each mcs, the agent behaviors are governed by rules that

were derived from experimental data found in the literature. The behaviors of each agent are

based on environmental conditions, such as nearby cells and cytokine gradients, as well as

probability-based rules. As an example, a capillary located near a damaged fiber has a

probability of becoming non-perfused and then senses the amount of VEGF-A and MMP-9 at its

location to decide if the levels are adequate to induce angiogenesis (Table 6). Model outputs

include CSA recovery (sum of total healthy fiber elements normalized by the initial CSA),

capillary and collagen density, cell counts, relative cytokine abundance, and spatial coordinates

of cells and cytokines.

Overview of Agent Behaviors

Simulated behaviors (Fig. 2B) of the neutrophils and macrophages include cytokine-dependent

recruitment, chemotaxis, phagocytosis of damaged fibers (neutrophils, monocytes, and M1

macrophages), phagocytosis of apoptotic neutrophils (monocytes and M1 macrophages),

secretion and uptake of cytokines, and apoptosis. The SSC and fibroblast agent behaviors also

include cytokine-dependent recruitment, chemotaxis, secretion and uptake of cytokines, and

apoptosis, in addition to quiescence, activation, division, and differentiation. The biological

intricacy of some cell types, such as SSCs which have a more complex cell cycle and are

regulated by dynamic interplay of intrinsic factors and an array of microenvironmental stimuli,

led to the necessity for adding more rules that govern their behaviors[27]. The SSCs have 33

parameters dictating the 17 agent rules (Table 3), fibroblasts have 27 parameters for 11 agent

rules (Table 4), macrophages have 31 parameters for 15 agent rules (Table 2), neutrophils have

18 parameters for 7 agent rules (Table 1), fibers have 18 parameters for 4 agent rules (Table 5),

and microvessels have 22 parameters for 6 agent rules (Table 6). At each mcs, cytokines are

secreted by agents if certain conditions were met. For cell recruitment, the levels of recruiting

cytokines for each agent are checked, and if the concentration is high enough to signal cell

recruitment (Supplemental Table 1), a new agent is added to the field at the location of the

highest concentration. The agents also undergo chemotaxis by sensing the surrounding

cytokine gradients and move towards higher concentrations of cytokines, binding and removing

that cytokine as they move along it to simulate physical binding of the cytokine to the receptor.

Agents that are in a quiescent state require a certain threshold level of cytokines to become

activated and cannot chemotax, secrete, divide, or differentiate until this threshold is reached.

Our model assumes each unique cell type secretes the same concentration of cytokines per

timestep for all relevant cytokines to drive model agent decisions. Each computational timestep

represents 15 minutes of real-world time. We assume that this is of sufficient resolution to

accurately reproduce immune cell agent behaviors during regeneration.

Neutrophil Agents

Neutrophils are recruited through capillaries to sites of necrotic tissue (Table 1). Neutrophils

move to areas of necrotic tissue with high concentrations of HGF by chemotaxing along the

HGF gradient to reach areas of necrosis[28,29]. Neutrophils phagocytose necrotic tissue and

facilitate remodeling into ECM with low-collagen density. During phagocytosis, neutrophils

secrete MMP-9, MCP-1, and TNF-α[30,28,31,32]. Individual neutrophil agents apoptose after

phagocytosing two necrotic cells (based on calibration) or 12.5 hours after their recruitment[33].

Macrophage Agents

Resident macrophages are distributed randomly throughout the tissue at a ratio of 1 macrophage

per 5 myofibers at model initialization and secrete MCP-1[34] (Table 2). Resident macrophages

chemotax along MCP-1 and HGF chemical gradients and secrete MMP-9, TNF-α, and MCP-1

during simulation[35–40]. After tissue injury, monocytes are recruited through healthy capillary

microvasculature and chemotax along MCP-1, VEGF-A, TGF-β[41,42–44]. Monocytes infiltrate into

the tissue if the MCP-1 concentration is above a specified threshold at a capillary site. Resident

macrophages, monocytes, and the M1 macrophages differentiated from monocytes may

phagocytose areas of necrotic tissue and apoptotic neutrophil agents[45–47]. During phagocytosis,

these agents secrete MMP-9, HGF, TGF-β, and IL-10[48–52].

Monocytes transition to M1 polarized macrophages when the monocyte agent experiences a large

enough TNF-α concentration or if enough time has passed that a predefined transition time

threshold is met. Each monocyte agent at creation has a defined transition time sampled from a

gaussian distribution with mean and standard deviation (SD) set to reproduce literature-defined

populations of M1 macrophages over time[51,53].

M1 macrophages may transition to M2 macrophages if the M1 macrophage agent experiences

an IL-10 concentration that exceeds a threshold value or if the M1 macrophage has phagocytosed

enough to meet a calibrated threshold value (as discussed in *Model Calibration*)[51,54,55]. Following

the transition to the anti-inflammatory phenotype, the M2 macrophages can proliferate, secrete

TGF-β and IL-10, and chemotax along an MCP-1 gradient[38,51,56].

SSC Agents

The model is initialized with 1 quiescent SSC per every 4 fibers and upon injury[57]. Additional

SSCs are recruited based on the amount of HGF, MMP-9, and TGF-β[58–61] (Table 3). For SSC

activation there has to be enough HGF at the location of the quiescent SSC to induce activation[61–

64]. The SSCs also chemotax up the MMP-9 gradient, removing some of the MMP-9 as they move

along it. Activated SSCs can also undergo symmetric or asymmetric division and differentiation

given that the required cytokine signaling is met locally. Activated SSCs differentiated into

myoblasts and myoblasts differentiate into myocytes[65–67]. Myocytes can fuse to other myocytes

to form new myotubes or fuse to fibers as long as the fiber is not fusion incompetent (i.e., fully

necrotic)[27,68–70]. Maturation of myotubes is required for fusion of additional myocytes to the new

fiber[69,71,72]. If the damage signal is not sustained, activated SSCs return to quiescence. If there is

enough TGF-β to induce apoptosis and not enough VEGF-A or macrophages nearby to block it,

the SSC undergoes cell death and leaves the simulation[42,73–75].

Fibroblast Agents

For model initialization, fibroblasts are randomly placed within the ECM at a population size that

is proportional to the number of fibers[76] (Table 4). Fibroblasts are activated based on the

concentration of TGF-β around the fibroblast[77,78]. Fibroblasts include an additional expression in

their effective energy function that directs their migration towards areas of low-density collagen

ECM [79]. Specifically, fibroblasts can form spring-like links to drag them towards areas of low-

density ECM which are implemented with the relation $\lambda_{ij}(l_{ij} - L_{ij})^2$ where λ_{ij} denotes a

hookean spring constant of a link between cells *i* and *j, l* represents the current distance

between the centers of mass between the two cells (in our case, fibroblast and low collagen

ECM), and *L* is the target length of the spring-like link. In addition to the cytokines secreted by

fibroblasts (Table 4), collagen is secreted at low-density collagen ECM[80–85]. Fibroblasts divide

when they are near dividing SSCs and can differentiate into myofibroblasts with extended

exposure to TGF-β[76,86,87]. The myofibroblasts can secrete more collagen regardless of the ECM

density[88]. Fibroblasts can undergo apoptosis if there are adequate levels of TNF-α at the site of

the cell but it can be blocked if there is sufficient TGF-β[89].

ECM Agents

ECM elements surround the fiber elements and are assigned a collagen density parameter which

varies based on the amount of necrotic tissue removed and the extent of fibroblast/myofibroblast

collagen secretion. When necrotic elements are removed, the phagocytosing inflammatory cells

secrete MMP-9s which degrade some of the collagen within that section of the ECM, thereby

causing that element to have a lower collagen density[28]. The collagen density of the ECM alters

the diffusivity of the secreted factors, and fiber placement is dependent on the collagen density

(discussed below). The fibroblasts help rebuild the ECM by secreting collagen on low collagen

density ECM elements[80]. Myofibroblasts can secrete collagen on any ECM element and if

prolonged results in high density collagen elements, representing a fibrotic state.

Fiber and Necrotic Agents

Upon model initialization, a portion of the muscle fiber agents are converted to necrotic fibers

based on the user prescribed injury. Fibers that reach a damaged threshold became fully necrotic

whereas those surrounding the area of necrosis were damaged but not fully apoptotic cells.

Healthy fiber elements secrete VEGF-A, and necrotic elements secrete HGF and TGF-β[63,90,91]

(Table 5). Phagocytosing agents chemotax along those gradients to clear the necrosis, but before

a new fiber can be deposited, the collagen has to be restored so that there is a scaffold to hold

the fiber in place[34]. Fully necrotic fibers are fusion incompetent and require myocyte-to-myocyte

fusion to form a new myofiber and require maturation before additional myocyte fusion[69,71,72].

Damaged fibers are regenerated by myocytes fusion to the healthy fiber edge[92].

Capillary and Lymphatic Agents

The muscle fascicle environment includes approximately 4 capillaries per fiber and 1 lymphatic

vessel[93,94] (Table 6). The model defines perfused capillaries as capillary agents that can transport

neutrophils and monocytes into the system proportional to the concentration of recruiting

cytokines[29,41]. The neutrophils and monocytes are added to the simulation at the lattice sites

above capillaries (within the cell layer Fig. 2B) and chemotax along their respective gradients.

The recruitment of the neutrophils and monocytes are distributed among the healthy capillaries

with a higher affinity for capillaries at locations with higher concentrations of HGF and MCP-1,

respectively. Under physiologically reasonable chemotactic gradient conditions, the recruited

immune cells dispersed efficiently, with no aggregation. Capillaries that are neighboring areas of

necrosis become non-perfused and therefore are unable to transport cells into the

microenvironment until regenerated[95]. Angiogenesis can occur as long as there is enough VEGF-

A present at the non-perfused capillary[96]. Similar to published studies, there is an increase in the

capillary-to-myofiber ratio during muscle regeneration, which is due to the formation of new

capillary sprouts modulated in part by local MMP-9 and VEGF-A levels[95,97,98].

The lymphatic vessel uptakes cytokines at lattice locations corresponding to the lymphatic vessel

and will remove cells located in lattice sites neighboring those corresponding to the lymphatic

vessel[99]. In addition, we have included a rule in our ABM to encourage cells to migrate towards

the lymphatic vessel utilizing CompuCell3D External Potential Plugin[100]. The influence of this rule

is inversely proportional to the distance of the cells to the lymphatic vessel.

Binding, Diffusivity, and Collagen Density

For many of the agent behaviors described above, there are associated binding events that play

key roles in regulation of the cytokine fields. Any cytokine dependent behavior is coupled with

removal of a portion of that cytokine once the behavior is initiated. For example, upon SSC

activation the amount of HGF required to activate is taken up by the SSC and removed from the

cytokine field to simulate the ligand binding and endocytosis resulting from SSC activation. Similar

binding events were modeled for SSC and fibroblast division and differentiation, macrophage

transitions, cell apoptosis, and chemotaxis along a cytokine gradient.

Due to limited data availability quantifying the diffusion constants of the modeled cytokines in the

context of the tissue microenvironment (which includes diffusion-altering elements including

collagen and glycosaminoglycans (GAGs)), we applied a diffusivity estimation technique[101]. To

do so, previously developed methods (equation 1) were applied to account for the combined

effects of collagen and GAGs (Table 7)[101]. The expression includes the radius of the cytokine (r_s),

the radius of the fiber (r_f), the volume fraction (ϕ), D and D_∞ are the diffusivities of the cytokines

in the polymer solution and in free solution, respectively. This estimation technique allowed for

consistent conditions for cytokine diffusion calculations and fluctuations based on changes in

collagen density within the model.

$$D = D_\infty \left(-\phi^{\frac{1}{2}} \frac{r_s}{r_f} \right)_{collagen} \times \exp \left(-\phi^{\frac{1}{2}} \frac{r_s}{r_f} \right)_{GAG} \qquad\qquad (1)$$

Throughout the model simulation, the diffusivity is recalculated with the updated collagen volume

fraction, as the collagen density changes throughout the microenvironment. This allows the

changes in collagen density within the ECM to be reflected in the diffusion rate of each of the

cytokines in the model.

Model Calibration

Known parameters were fixed to literature values, and uncertain parameters were calibrated by

comparing simulation outcomes to published experimental data. Calibration data included

published findings from injury models that have synchronous regeneration after tissue necrosis

(i.e., cardiotoxin, notexin, and barium chloride)[97]. The metrics that were used to calibrate the

model included time-varying CSA[102], SSC counts[76], and fibroblast counts[76]. These metrics were

used for calibration because of their key roles in the regeneration of muscle and the complex

interplay between these outputs. Cell count data were normalized by the number of cells on the

day of the experimental peak to allow for comparison between experiments and simulations. For

CSA, the experimental and model outcomes were normalized using fold-change from pre-injury

to compare model-simulated with experimental CSA, as percent change from baseline is

commonly used experimentally[103,104]. Model cell counts were normalized by the number of cells

at the peak time point in the experimental data. SSC and fibroblast counts were normalized to

day 5. Neutrophil counts were normalized to day 1. Total macrophage, M1, and M2 counts were

normalized to day 3. The capillaries were normalized to fiber area, as done in the experimental

data.

Initial ranges for the 52 unknown parameters were determined by literature review or by running

the model to test possible upper and lower thresholds for parameters (Supplemental Table 1).

To narrow the parameter ranges beyond those initial ranges, we used a recently published

calibration protocol, CaliPro, which utilizes parameter density estimation to refine parameter

space and calibrate to temporal biological datasets[20]. CaliPro was selected as the calibration

method because it is model-agnostic which allows it to handle the complexities of stochastic

models such as ABMs, selects viable parameter ranges in the setting of a very high-dimensional

parameter space, and circumvents the need for a cost function, a challenge when there are

many objectives, as in our case. Briefly, Latin hypercube sampling (LHS) was used to generate

600 samples which were run in triplicate. These runs were then evaluated against a set of pass

criteria, and the density functions of the passing runs and failing runs were calculated

(Supplemental Table 2). Parameter ranges were narrowed by alternative density subtraction

(ADS), where the new ranges were determined by the smallest and largest parameter values

where the density of passing is higher than the density of failing. The sensitivity of the model

outputs to the parameters was examined using LHS in combination with partial rank correlation

coefficient (PRCC)[105]. LHS/PRCC methods have been used for various differential equation

models and ABMs[106]. PRCC was computed using MATLAB to determine the correlation

between ABM parameters (i.e. cytokine threshold for activation) and the ABM output (i.e.

fibroblast cell count). Correlations with a p-value less than 0.05 were assumed to be statistically

significant. This helped refine initial parameter bounds as well as make model adjustments

based on the parameter dynamics elucidated from PRCC. This process of sampling parameter

ranges, evaluating the model, and narrowing parameter ranges was repeated in an iterative

fashion while updating pass criteria until a parameter set was identified that consistently met the

strictest criteria (Supplemental Fig. 1). The final passing criteria were set to be within 1 SD of

the experimental data for CSA recovery and 2.5 SD for SSC and fibroblast count. These criteria

were selected so that the model followed experimental trends and accounted for both model

stochasticity and experimental variability in datasets that had narrower SDs for certain

timepoints. Early iterations had a wide parameter range to avoid missing portions of the realistic

parameter space. At first, narrowing the parameter space increased passing simulations, but

upon reaching the ideal parameter space, further narrowing eliminated viable parameters,

resulting in fewer passing runs. Following 8 iterations of narrowing the parameter space with

CaliPro, we reached a set of parameters that had fewer passing runs than the previous iteration.

We then returned to the runs from the prior iteration and set the bounds such that all 3 runs from

the parameter set fell within the final passing criteria. The final parameter set was run 100 times

to verify that the variation from the stochastic nature of the rules did not cause output that was

inconsistent with experimental trends.

Model Validation

We compared model outputs (M1, M2, and total macrophage counts[97,107], neutrophil counts[108],

and capillary counts[102]) that were kept separate from the calibration criteria with published

experimental data to verify that these outputs followed trends from the experimental data

without requiring extra model tuning. In addition, we also altered various model input conditions

(cell input conditions, cytokine dynamics, and microvessel dynamics) to simulate an array of

model perturbations (Table 8) which allowed comparison of a set of model outputs with separate

published experiments. For example, we simulated an IL-10 knockout condition by eliminating

IL-10 secretion and adjusting the diffusion and decay parameters so that the concentration of IL-

10 throughout the simulation was reduced, decreasing the behaviors driven by the cytokine as a

result of the KO condition. One hundred replicates of each model perturbation were performed,

and perturbation outputs were compared with control simulation outputs via a two-sample t-test

with a significance level of 0.05. We were then able to compare how the model outputs aligned

with published experimental findings to determine if the model could capture the altered

regeneration dynamics.

Sensitivity Analysis

A sensitivity analysis was performed using LHS-PRCC to examine the impact of cytokine-

related parameters on model outputs of interest. Diffusion coefficients and decay rates for the

seven cytokines (HGF, TGF-β, MMP-9, TNF-α, VEGF-A, IL-10, MCP-1) were sampled across a

range from 0.1 to 10 times the calibrated value while holding the other parameters constant.

Three hundred samples were generated, and these parameter sets were simulated in triplicate.

PRCCs were calculated with α=0.05 and a Bonferroni correction for the number of tests every

10 ticks/hours for CSA and cell counts for SSCs, fibroblasts, non-perfused capillaries,

myoblasts, myocytes, neutrophils, M1 macrophages, and M2 macrophages.

In Silico Experiments

To gain insight into the recovery response with altered angiogenesis, we simulated different

levels of VEGF-A injections to test how increases in VEGF-A impacted regeneration outcomes.

In addition, we simulated conditions of hindered angiogenesis in which damaged capillaries

were unable to reperfuse following injury (n = 100 for each simulation condition). Simulations

were also conducted to examine correlations between cytokines and their impact on various cell

behaviors and regeneration outcomes. Next, a sensitivity analysis was performed to understand

how alterations in cytokines influence key metrics of regeneration. LHS-PRCC was used to

quantify the impact of cytokine-related parameters (i.e. diffusion rates and decay coefficients) on

outputs of interest (CSA, SSC, fibroblasts, non-perfused capillaries, myoblasts, myocytes,

neutrophils, M1, and M2). A single time point for each output is summarized in Table 9, and

these were chosen at the timepoint when PRCC values were peaking, with complete results

available in Supplementary Figure 2.

This sensitivity analysis was then used to guide *in silico* experiments based on which cytokine

parameters promoted favorable regeneration outcomes (i.e. improved recovery, fewer non-

perfused capillaries, increased SSCs). Following individual cytokine parameter alterations, we

combined the cytokine alterations based on beneficial outcomes from the initial *in silico*

experiments to determine if the benefits would be cumulative.

Results

ABM outputs align with calibration and validation data

Following parameter density-based calibration, the unknown parameters were narrowed into a

final calibration parameter set (Supplemental table 1). The simulations captured SSC and

fibroblast cellular behaviors, as well as CSA outcomes, that aligned with experimental studies

(Fig. 3A-C). The model data were consistent with the experimental trends, and the 95%

confidence interval was within the SD for all calibration data time points except for SSCs at day

3 (Fig. 3B). Macrophage (total, M1, and M2), neutrophil, and capillary counts, which were not

used for model calibration, were found to also be consistent with experimental trends and

allowed us to independently validate model outputs (Fig. 3D-H).

ABM perturbations are consistent with published experiments

Overall, the model reproduced findings from multiple studies, replicating how altered conditions

lead to both improved and diminished muscle regeneration (Fig. 4). Injections of VEGF-A led to

faster CSA recovery, more damaged tissue clearance, and a concentration dependent dose

response, consistent with prior studies[73]. Cell depletion simulations predicted decrease in all

markers of regeneration, consistent with prior studies[73,109,110]. When simulating hindered

angiogenesis conditions, the model aligned with experimental studies showing detriments in

CSA recovery, increased neutrophil and macrophage cells, and elevated ECM collagen density,

indicating progression of fibrosis within the microenvironment[111]. There were a few cases in

which model predictions did not align with published studies. First, simulations of TNF-α KO

predicted increased CSA recovery, while experiments measured decreased recovery of CSA.

This difference is likely due to the fact that the model did not include cross regulation with

interferons which are upregulated with TNF-α KO[112]. Second, macrophage depletion simulations

predicted decreased TGF-β concentrations throughout the simulation while experiments

measured an initial decrease in concentration followed by increased concentrations at days 7

and 14. This difference may be due to the fact that macrophage depletion was experimentally

induced with clodronate-containing liposomes which could have reduced consistency of

depletion across the time course and other downstream impacts that were not represented by

decreasing macrophages in the model perturbation[110].

Analysis of ABM perturbations leads to new insights regarding cytokine and cell

dynamics

The model allowed for new insights into the dynamics of muscle regeneration by providing

additional timepoints and metrics to evaluate the response to exogenous delivery of VEGF-A

and hindered angiogenesis. VEGF-A levels remained elevated compared to control simulations

following the injection at day 5 post injury (Fig. 5A). CSA recovery had the highest increase at

28 days post injury with the high (10^3) relative concentration delivered) VEGF-A injection

followed by the extra high ($2x10^3$) relative concentration delivered) injection (Fig. 5B). The

medium (750 relative concentration delivered) and low (500 relative concentration delivered)

VEGF-A injections had higher CSA recovery 15 days post injury but were not significantly

different from the control at day 28. All VEGF-A injections had a higher capillary count and were

proportional to the level of VEGF-A injection (Fig. 5C). The impact of VEGF-A injection on peak

SSC and fibroblast counts was dependent on dosage amount, with the extra high VEGF-A

injection resulting in the largest peaks (Fig. 5E&F). Cytokine concentration trends were similar

for all injections, but most peak levels were dosage dependent (Fig. 5G-L). In contrast, HGF

levels were elevated from day 5 to day 28 with hindered angiogenesis, as were TGF-β and IL-

10 (Fig. 5I & L). MCP-1 concentration had a lower overall peak level with elevated levels from

days 21 to 28 (Fig. 5H). Hindered angiogenesis had lower CSA recovery throughout the

simulation and did not achieve unaltered regeneration levels (Fig. 5B).

Cytokine knockout perturbations revealed crosstalk and temporal interplay between cytokines

(Fig. 6). For example, with MCP-1 KO there was an overall increase in cytokine levels for all

other cytokines within the microenvironment except for VEGF-A at 12 hours post injury (Fig.

6A). By 7 days post injury TNF-α, TGF-β, IL-10, and MMP-9 had decreased from unaltered

regeneration day 7 levels but VEGF-A and HGF were elevated. With TNF-α KO there was a

decrease in TGF-β at early timepoints but a strong increase by day 28 (Fig. 6B). Following IL-10

KO there was an increase in TNF-α that peaked at 7 days post injury (Fig. 6C). HGF was

slightly decreased throughout and TGF-β was strongly decreased by day 7. MMP-9 was

decreased at 12 hours and 28 days post injury but heavily increased at day 7.

Cytokine dynamic analysis leads to new model perturbations that predict improved

regeneration

LHS-PRCC of cytokine decay and diffusion parameters elucidated temporal relationships

between cytokine parameters and key regeneration metrics, such as positive correlations

between CSA and TGF-β and MMP-9 decay (Table 9). Of all cytokine parameters, the model

outputs were most sensitive to HGF decay, with all outputs except M1 cell count being

significantly impacted. PRCC plots showed that TGF-β and MMP decay were positively

correlated and HGF decay was negatively correlated with CSA recovery, with higher

significance at timepoints after 12 days (Supplemental Fig. 2). Correlation plots for various

cytokine concentrations and regeneration metrics showed trends in cytokine dependent cell

behaviors such as the TNF-α concentration that led to heightened fibroblast cell counts as well

as the corresponding TNF-α concentration threshold that results in diminished fibroblast

response (Supplemental Fig 3). These PRCC trends guided cytokine parameter perturbations

to include lower HGF and VEGF-A decay, higher TGF-β, MMP-9, and MCP-1 decay, and higher

MCP-1 diffusion because each of the cytokine modifications indicated some form of enhanced

regeneration outcome metrics (Supplemental Table 3). All these perturbations except MCP-1

decay show increased CSA, increased healthy capillaries, and increased SSCs (Fig. 7). Finally,

a combination of all changes except for MCP-1 decay was simulated. The combined cytokine

alteration resulted in the highest CSA recovery (Fig. 7A), as well as increased M1 macrophage

counts (Fig. 7B), decreased M2 macrophage counts (Fig. 7C), increased fibroblasts (Fig. 7D)

and SSCs cell counts (Fig. 7E). Capillaries regenerated faster in the combined perturbation than

under unaltered conditions (Fig. 7F). It is likely that the combination of cytokines perturbed cell

dynamics in a manner that promoted regeneration in both the early and later phases. During

early regeneration, lower HGF decay, higher TGF decay, and MCP-1 diffusion contributed to

increased SSCs while lowered VEGF decay increased angiogenesis. During late regeneration,

lower HGF decay and higher MMP decay contributed to an increased anti-inflammatory state

and SSC differentiation. The combined cytokine perturbation predicted a 13% improvement in

CSA recovery compared to the unaltered regeneration amount at 28 days. The combined

cytokine perturbation also had higher peaks in SSC and fibroblast counts than any of the

singular cytokine perturbations, indicating the synergistic effects of altering the cytokine

dynamics in combination.

Discussion

We developed a novel ABM that recapitulates muscle regeneration and, unique from prior work,

includes spatial interactions between cytokines and the microvasculature based on relevant

literature[8,15,16]. The creation of the model provides a more controlled environment for studying

muscle regeneration, reducing error and variation commonly encountered with *in vivo*

experiments. Model predictions aligned with experimental data under various altered inputs.

Through *in silico* experiments, we gained new insight into how the combination of key cytokine

dynamic alterations could increase SSC cells and enhance CSA recovery. The ability for altered

cytokine concentrations to change regeneration outcomes is consistent with studies that have

found enhanced muscle recovery with delivery of platelet-rich plasma (PRP) which contain

VEGF and TGF- β[113]. These model perturbations allow development of hypotheses and can

provide the basis for future experiments and potential therapeutic interventions such as

plasminogen activators to alter cytokines dynamics to enhance muscle recovery.

ABM provides biological insight on nonlinear effects of cytokine levels

The ABM offers valuable insights into the muscle regeneration dynamics under various altered conditions, elucidating the complex interplay of cytokines, angiogenesis, and cell behaviors. Systematic simulations reveal critical thresholds, nonlinear effects, and synergistic cytokine combinations impacting regeneration. Perturbations varying VEGF-A injection doses showed increased CSA recovery up to a threshold (high VEGF-A injection simulation), beyond which further improvements in CSA recovery cease. Cytokine KO simulations revealed the complex nature of the relationship between cytokines; removal of one cytokine from the system has a cascading temporal impact. Relationships between cytokines and cellular outputs exhibit nonlinear effects, as seen with the limited impact of elevated HGF on CSA recovery beyond a threshold and the non-monotonic relationship between TNF-α and fibroblast counts (Supplemental Fig. 3). Further analysis revealed that specific combinations of cytokine perturbations could enhance regeneration beyond singular cytokine interventions. For example, a combined intervention of: 1) decreasing HGF and VEGF-A decay, 2) increasing TGF-β and MMP-9 decay, and 3) increasing MCP-1 diffusion enhanced muscle regeneration. Prior studies have shown that individually, increased HGF[114], VEGF-A[73], and MCP-1[115] stimulate muscle regeneration whereas reduced TGF-β[116] and MMP-9[117] stimulate muscle regeneration. The model suggests that combined alterations have a stronger regenerative effect than individual cytokine changes, enhancing muscle recovery through distinct mechanisms—increasing healthy capillaries, SSC counts, and reducing inflammatory cells.

Cytokine modifications intended to enhance muscle recovery can have clinical relevance and have been studied in various settings. For example, synthetic biomaterials coated with IL-4 have been implanted as a cytokine delivery vehicle and were successful in increasing M2 cells within the muscle[118]. Cytokine antagonist have been successful at promoting muscle regeneration, seen in prior work with anti-IL-6[119]. Studies have also shown that activation of plasmin is able to

induce the release of ECM bound VEGF, increasing angiogenesis[14,120]. Due to the complex

network of cytokines, studies that deliver simple modulation of one or two cytokines typically

have an insufficient response to generate appreciable improvements. This suggests that using a

combination of biologic and synthetic biomaterials to modulate multiple cytokines is necessary,

which aligns with our findings[118]. Multiple cytokines have been modulated through the use of

PRP which contains VEGF-A and an array of other cytokines, but PRP has had mixed success

in a clinical setting[121]. Our model has the capability to test and optimize various combinations of

cytokines, along with exploring different temporal schedules for delivering specific treatments.

For instance, it can predict whether modified combinations of cytokines prove beneficial at

specific timepoints, aiding in the development of optimal treatment compositions aligned with

the temporal dynamics of the regeneration cascade. These predictions provide novel concepts

for future experiments and potential interventions. For example, the predictions from the model

suggest that interventions that combine activation of plasmins for bound VEGF release[14,120] with

delivery of synthetic biomaterials coated with HGF[122], TGF-β antagonist[123], nuclear factor-kappa

B inhibitory peptide to inhibit MMP-9[124], and recombinant MCP-1 hydrogels[125] to alter diffusion

rate would result in improved regeneration outcomes.

Advancements from prior muscle regeneration models

Previous studies have employed computational models to investigate muscle regeneration

across diverse contexts, such as Duchenne muscular dystrophy and volumetric muscle loss[8,15].

Earlier muscle regeneration ABMs from our group have been used to test the effects of priming

muscle with inflammatory cells prior to injury[16]. While these models laid the foundation for

simulating muscle adaptations, they were constrained by limited diffusion capabilities and an

absence of critical features related to microvessel growth and remodeling throughout the

regeneration process. Similarly, other ABMs from our group have examined altered

microenvironments, but their omission of spatial cytokine diffusion hindered comprehensive

representation of cell behaviors pivotal to regeneration[8,15]. Recently, new ABMs have been

published that focus on cerebral palsy and the impact of injury type on eccentric contraction-

induced damage[17,18].

The model presented here provides advancements over prior models in three areas: 1) explicit

modeling of cytokine-specific diffusion and decay that depends on the ECM environment, 2)

addition of microvasculature, and 3) incorporation of a robust and rigorous calibration and

validation process. The addition of microvessel growth and remodeling dynamics empowers

investigations into how interventions impact angiogenesis during regeneration, thereby

influencing muscle recovery outcomes. By considering the intricate relationship between

microvessels and regeneration, our model opens avenues for evaluating the effects of

interventions on the broader recovery process. Secondly, understanding how cytokines

influence cell behaviors at different times during regeneration is crucial for determining optimal

treatment targets and dosing. While cytokine dynamics can be altered experimentally, doing so

is expensive and time consuming[12,14] so exploring many combinations of alterations would be

practically infeasible. Our model incorporates decay and diffusion dynamics of a subset of

cytokines to allow testing of far more alterations in cytokines than would be reasonable to

conduct experimentally. Lastly, we leveraged the CaliPro technique for parameter density

estimation-based calibration and LHS-PRCC to gain biological insight by analyzing how altered

microenvironmental parameters could benefit regeneration outcomes. This approach of

implementing parameter identification to guide model perturbations demonstrates the

capabilities of the model as a novel tool for generating new hypotheses and identifying

mechanisms to target for enhanced regeneration outcomes.

Our model predictions are generally consistent with these prior models, with added biological

complexity that has yielded several new important insights. For example, simulation of hindered

angiogenesis predicted a decrease in SSCs leading to poor CSA recovery, similar to how lower

SSC counts resulted in lower CSA recovery in perturbations in both healthy and DMD

simulations[15]. Our model provides additional understanding about the corresponding spatial

cytokine changes that ultimately result in modulation of SSC dynamics within the

microenvironment. The additional model advancements incorporated address prior muscle

regeneration modeling gaps in understanding of how angiogenesis alters recovery outcomes as

well as the response of complex spatial cell and cytokine dynamics.

Limitations and future work

There are some important limitations of this study that should be discussed. First, the model

does not include all cell types and cytokines that are known to influence muscle regeneration

and does not account for cytokine subtype or differences between endogenous and exogenous

cytokines. These cells and cytokines likely have redundant functions, given the model effectively

captures muscle regeneration using the included cells and cytokines. Second, the model does

not currently represent hypertrophy during regeneration, which restricts CSA recovery from

surpassing 100%; however, the cell dynamics it portrays remain consistent with those observed

in studies that lead to hypertrophy following injury. Third, we assume a 2D cross-section based

on similar ABMs that have explored the relations of 2D to 3D simulations. These studies found

that the diffusion accuracy is not greatly varied and that 2D is sufficient to predict the same

mechanisms seen in 3D simulations[126,127]. To determine the robustness of the 2D initial cross-

section, preliminary testing has shown that the initial spatial configuration can be altered and still

achieve similar results (Supplemental Fig. 5), but further examination is needed to determine

sensitivity to numerous configurations. Fourth, the calibration and validation dataset integrated

multiple datasets from diverse sources. We acknowledge inherent limitations arising from

variations in sample sizes and experimental techniques across sources. Fifth, it is also possible

that the calibrated parameters are unable to capture behaviors that were not exhibited within the

experimental datasets used in parameterization. While we tested ranges for each parameter

and settled on a single parameter set that best fit the calibration data, there may be additional

parameters sets that fit the calibration data but have varied levels of stochasticity and altered

reproducibility of replicate simulations. Lastly, the current model was calibrated to male mice

data despite known sex difference in skeletal muscle, regeneration mechanisms, and the

timeline of recovery[128–130]. Experimental measurements of female muscle regeneration are fairly

limited because most muscle injury studies only use male mice or do not distinguish between

sexes, making it difficult to incorporate sex differences into the model[131]. Experiments that

incorporate female mice and measure hormone levels are needed to accurately incorporate

rules to distinguish between the sex dependent dynamics of muscle regeneration.

This paper describes a significant advancement in modeling the complex process of muscle

regeneration. Future efforts will extend the use of parameter density estimation to optimize the

selection, doses, and timing of injections of exogenously delivered cytokines. Further refinement

of analysis methods could be pursued to disentangle specific underlying mechanisms of the

dynamic feedbacks that drive the observed model outputs. Predictions from model simulations

will also be used to inform future experiments by highlighting crucial time points to measure and

predicted effect sizes for power analysis. Additionally, we aim to explore diverse muscle injury

types and locations (i.e. injury relative to microvascular components) and their varying recovery

responses, addressing challenges in comparing different acute injury techniques found in the

literature. This study underscores the significance of cellular and cytokine spatial dynamics in

muscle regeneration. Further inclusion of additional factors and hormones would provide a more

holistic understanding of the system and how treatments may be altered based on

microenvironmental conditions, providing a unique framework for the study of personalized

muscle injury treatment.

Acknowledgements

The authors acknowledge NIH Grant #R21AR080415, Wu-Tsai Foundation Agility Project

Funding, and NSF GRFP GRANT #1842490 for financially supporting this research and to

James Glazier and T.J. Sego for providing technical support with the CC3D ABM platform.

Additional Resources

The ABM source code is publicly available at the following sites:

SimTK: https://simtk.org/docman/?group_id=2635

Zendo: https://zenodo.org/records/10403014

GitHub: https://github.com/mh2uk/ABM-of-Muscle-Regeneration-with-Microvascular-
Remodeling

Figure 1. Flowchart of ABM rules. The model starts with initialization of the geometry and the prescribed
injury. This is followed by recruitment of cells based on relative cytokine amounts within the
microenvironment. The inflammatory cells, SSCs, and fibroblasts follow their literature defined rules and
probability-based decision tree to govern their behaviors. The boxes represent the behavior that the agent
completes during that timestep given the appropriate conditions and the circles represent the uptake that
occurs as a result of the simulated binding with microenvironmental factors for certain cell behaviors.
ABM, agent-based model; SSC, satellite stem cell; ECM, extracellular matrix; TGF-β, transforming growth
factor β; HGF, hepatocyte growth factor; TNF-α , tumor necrosis factor α; VEGF-A, vascular endothelial
growth factor A; MMP-9, matrix metalloproteinase-9; MCP-1, monocyte chemoattractant protein-1; IL-10;
interleukin 10.

Figure 2. Overview of ABM simulation of muscle regeneration following an acute injury. **A.** Simulated cross-sections of a muscle fascicle that was initially defined by spatial geometry from a histology image. Muscle injury was simulated by replacing a section of the healthy fibers with necrotic elements. In response to the injury, a variety of factors are secreted in the microenvironment which impacts the behavior of the cells. The colors correspond with those typically seen in H&E staining. **B.** ABM screen captures show the spatial locations of the cells throughout the 28-day simulation. The agent colors were matched to those typically seen in IHC-stained muscle sections.

Table 1. Neutrophil Agent Rules

Neutrophil Agent Behavior	Sources
Recruitment signal: necrosis	28
Neutrophils are brought to site of injury via capillaries	29
Phagocytose necrosis	30
Secretes MMP-9, MCP-1, TNF-α during phagocytosis	16,28,31,32
Undergoes apoptosis after phagocytosis or 12.5 hours	33
Migrates towards areas of high HGF	39
Migration speed ~7.5 µm/min	132,133

Table 2. Macrophage Agent Rules

Macrophage Agent Behavior	Sources
Initial count: 1 resident macrophages per 5 myofibers	34
Recruitment signal: MCP-1	38,42
Monocytes are brought to the site of injury via microvessels	41
Resident macrophages secrete MMP-9, MCP-1, and TNF-α and chemotax along MCP-1 and HGF	35–40
Monocytes chemotax along MCP-1, VEGF-A, and TGF-β	42–44,134

Monocyte migration speed ~4 µm/min	[135]
M1 macrophages secrete VEGF-A, MMP-9, and TNF-α and chemotax along MCP-1	[136–139]
Monocytes, resident and M1 macrophages phagocytose apoptotic neutrophils and necrosis	[45–47]
Monocytes and macrophages secrete MMP-9, HGF, TGF-β, and IL-10 during phagocytosis	[16,48–52]
Monocyte transitions to M1 occurs when TNF-α threshold is met or based on literature means and standard deviation properties	[51,53]
M1 transition into M2 is mediated by the amount of IL-10 and the amount the M1 has phagocytosed	[16,51,54,55]
M2 macrophages secrete TGF-β and IL-10 and chemotax along MCP-1	[16,38,56,140]
Macrophages can proliferate following the transition to the anti-inflammatory (M2) state	[51]
Macrophage migration speed ~0.62 µm/min	[135]
Macrophages apoptose in a Poisson distribution	[141]

684 **Table 3.** SSC Agent Rules

SSC Agent Behavior	Sources
Initial count: 1 SSC per 4 fibers	[15,57]
Recruitment signal: HGF + MMP-9 - TGF-β	[15,58–61]
Activation signal: HGF	[15,61–64]
Activated SSCs secrete MCP-1 and VEGF-A	[42]
Activated SSCs migrate towards areas of high MMP-9	[59,142]
Myoblasts migrate towards high TNF-α	[143]
Division signal: TNF-α + VEGF-A - TGF-β	[15,60,144,145]
Differentiation signal: 3*IL-10 - HGF – TNF-α - TGF-β	[15,54,146–148]
Activated SSCs differentiate to myoblasts, myoblasts to myocytes, and myocytes to myotubes/myofibers	[65–67]
Differentiated myocytes fuse at damaged fiber edge or fuse together to from new, immature myotubes	[27,68–70]
50% cell divisions are symmetric, 50% asymmetric	[15,149,150]
Division probability decreases with each cell division; 1st division 85%; 2nd 65%; 3rd 20%	[15,151]
VEGF-A and macrophages nearby can block apoptosis	[42,73,74]
TGF-β triggers apoptosis	[75]
Time to divide: 10 hrs	[15,151,152]
Migration speed ~0.94 µm/min	[153]
Return activated SSCs to quiescence without sustained HGF	[61]

686 **Table 4.** Fibroblast Agent Rules

Fibroblast Agent Behavior	Sources
Initial count: 1 fibroblast per every 2 fibers	[15,76]
Activation signal: TGF-β	[154]
Fibroblasts move to low collagen ECM	[15,79]
Fibroblasts secrete TNF-α, TGF-β, MMP-9, VEGF-A. Collagen is secreted at low density ECM	[15,80–85]

Fibroblast division signaled by SSC division	15,76
Division probability decreases with each cell division; 1st division 100%; 2nd 25%; 3rd 6%	155
Fibroblast differentiation to myofibroblasts with extended TGF-β exposure	15,86,87
Myofibroblasts secrete double the amount of collagen and secretion is not dependent on collagen density	15,88
Fibroblasts apoptose with sustained exposure to TNF-α	15,89
Fibroblast migration speed ~0.73 µm/min	156
Sufficient TGF-β can block fibroblast apoptosis	15,89

688 **Table 5.** Fiber Agent Rules

Fiber Agent Behavior	Sources
Damaged muscle fibers secrete HGF and TGF-β	63,90
Healthy fibers secrete VEGF-A	91
Fibers that are fully necrotic are fusion incompetent, but damaged fibers are fusion competent	92
Immature myotubes gain functional capacity as they fully mature overtime	69,71,72

690 **Table 6.** Microvasculature Rules

Microvessel Agent Behavior	Sources
Initial count: ~4 capillaries per fiber, 1 lymphatic vessel per fascicle	93,94
Capillaries near necrosis will become damaged and unable to perfuse	95
With sufficient VEGF-A damaged capillaries will undergo angiogenesis	96
MMP-9 is elevated during capillary growth	98,157
Increasing capillary to myofiber ratio during muscle regeneration from new sprouting capillaries at areas with enough MMP-9 and VEGF-A	95,97,98
Cells and cytokines near lymphatic vessel will be drained via the vessel and removed from microenvironment	99

692 **Table 7.** Model parameters of spatial mechanisms

Parameter	Value	Source/Justification
Volume parameters		
Target volume (V_{target}) neutrophil	12	Chosen for an average cell diameter of 12 µm[158]
Target volume (V_{target}) SSC	10	Chosen for an average cell diameter of 10 µm[159]
Target volume (V_{target}) macrophage	21	Chosen for an average cell diameter of 21 µm[160]
Target volume (V_{target}) monocyte	8.5	Chosen for an average cell diameter of 8.5 µm[161]
Target volume (V_{target}) fibroblast	15	Chosen for an average cell diameter of 15 µm[162]
Volume multiplier (λ_{volume})	50	Volume constraint to maintain target[23]
Diffusion coefficients		

HGF	66.38 μm^2/s	Estimated diffusivity within the ECM accounting for baseline GAGs and collagen[101]
MMP-9	63.40 μm^2/s	
MCP-1	189.27 μm^2/s	
VEGF-A	112.10 μm^2/s	
TGF-β	90.33 μm^2/s	
TNF-α	138.95 μm^2/s	
IL-10	135.17 μm^2/s	
Chemotaxis parameters λ_c		
Neutrophils	750	Chosen for a cell velocity between 1-20 μm/min[132]
Macrophage	9.3	Chosen for a cell velocity around 0.62 μm/min[135]
Monocyte	75	Chosen for a cell velocity around 4 μm/min[135]
SSC	11.3	Chosen for a cell velocity around 0.94 μm/min[153]
Fibroblast	23	Chosen for a cell velocity around 0.73 μm/min[8]

694   **Table 8.** Model perturbation input conditions and corresponding published experimental results

Perturbation	Specific model conditions	Published outcomes
IL-10 knockout	Adjust diffusion and decay parameters so IL-10 is removed from the system	Attenuates shift to M2, disrupted SSC differentiation, slowed regeneration[163]
Neutrophil depletion	Lower neutrophil recruitment proportion	Abundant necrotic tissue 7 days post injury[109]
Macrophage depletion	Lower macrophage recruitment proportion	Decreased HGF, increased TGF-β and TNF-α, impaired regeneration[110]
MCP-1 Knockout	Adjust diffusion and decay parameters so MCP-1 is removed from the system	Increased necrosis at day 7, lower CSA at day 21, impaired phagocytosis[115]
Directed M2 polarization (Anti-inflammatory nanoparticles)	Require less phagocytosis and IL-10 for transition	Improved muscle histology and inflammatory resolution[164]
TNF-α Knockout	Adjust diffusion and decay parameters so TNF-α is removed from the system	Impaired recovery at days 5 and 12, increased inflammation[165]
Hindered angiogenesis	Increase VEGF-A and MMP-9 threshold required for angiogenesis	Delayed regeneration with toxin injury, and persistent immune cell infiltration with freeze injury[111]

VEGF-A injection	Add VEGF-A at specified concentration (100 for low and 1000 relative concentration for high), radius (300 pixels), and timepoint (5 days post injury)	Lower injury area at day 20 post injury with injection 5 days after damage[73]

Table 9. Summary of cytokine sensitivity analysis. Significance was determined with α=0.05, and a Bonferroni correction for the number of tests. + and - represent statistically significant positive and negative correlations, respectively.

	CSA	SSC	Fibroblasts	Non-perfused capillaries	Myoblasts	Myocytes	Neutrophils	M1	M2
Day	16.7	6.3	10.5	8.4	6.3	8.4	8.4	4.2	6.3
HGF decay	-	-	-	+	-	-	+		+
TGF-β decay	+	+	+	-	+				-
MMP-9 decay	+	+	+	-	+	+			
TNF-α decay			+						-
VEGF-A decay				+					
MCP-1 decay								+	+
MCP-1 diffusion		+		-				+	

Model Calibration

Model Validation

Figure 3. ABM calibration and validation. ABM parameters were calibrated so that model outputs for CSA recovery, SSC, and fibroblast counts were consistent with experimental data (A-C)[76,102]. Separate outputs from those used in calibration were compared to experimental data[97,102,107,108] to validate the ABM (D-H). Error bars represent experimental standard deviation, and model 95% confidence interval is indicated by the shaded region. Cell count data were normalized by number of cells on the day of the experimental peak to allow for comparison between experiments and simulations.

Figure 4. ABM perturbation outputs are compared to various literature experimental results. Each
perturbation model output is compared to the available corresponding published result. The top triangles
indicate the literature findings and the bottom triangles indicate the model outputs. Red triangles
represent a decrease, blue represents an increase, and gray represents no significant change. Time
points of comparison were based on which time points were available from published experimental data.
Refer to table 8 for model input conditions and supplemental table 6 for information on experimental
references.

Figure 5. Dose dependent response with VEGF-A injection compared to hindered angiogenesis. VEGF-A
concentration response to varied levels of VEGF injection (A). Hindered angiogenesis resulted in slower
and overall decreased CSA recovery (B). Capillary count was dependent on VEGF-A injection level (C).
Total macrophage count was similar between control and VEGF-A injections perturbations but
macrophage count was higher in later time points in the hindered angiogenesis simulation (D). SSC peak
varied with VEGF-A injection level and counts were prolonged in the hindered angiogenesis simulations
(E). The fibroblast peak was lower for the hindered angiogenesis perturbation and highest with the extra
high VEGF-A injection. In contrast to the other simulations, the fibroblast count was trending upwards at
later time points in the hindered angiogenesis perturbation (F). HGF levels were consistent between
control and VEGF-A injection perturbations but was significantly elevated in the hindered angiogenesis
perturbation (G). MCP-1, TGF-β, and IL-10 concentrations were elevated a later stages of regeneration
with hindered angiogenesis (H, I, L). TNF-α was elevated with the extra high VEGF-A injection and lower
with hindered angiogenesis (J). MMP-9 concentration was lower at the simulation midpoint but elevated
at late regeneration stages (K).

Figure 6. Heatmaps of changes in cytokine concentration at various timepoints throughout regeneration following individual cytokine knockout (KO) demonstrating cross-talk between cytokines. With MCP-1 KO there was an increase in all cytokines except VEGF-A at 12 hours post injury. Over the course of regeneration there was continued increasing elevation of HGF, increases in VEGF-A, and TGF-β decreased at day 7 followed by a strong increase by day 28 post injury (A). In the TNF-α KO simulations, there was an early decrease in TGF-β that shifts to strong increases by day 28. MMP-9 increased throughout the duration, HGF and IL-10 were decreased, VEGF-A lagged in the beginning but was increased during mid to late timepoints (B). Following IL-10 KO there were increases in TNF-α, decreases in HGF and TGF-β, and elevated MMP-9 at day 7 that decreased by day 28 (C).

Figure 7. Combined alterations of various cytokine dynamics enhance muscle regeneration outcomes.
All tested alterations except higher MCP-1 decay resulted in higher CSA recovery compared to the
control (A). M1 cell count was higher for all perturbations with the highest peaks with increased MCP-1
diffusion and the combined cytokine alteration perturbation (B). Higher MCP-1 decay resulted in the
largest M2 peak and higher MCP-1 diffusion, higher TGF-β decay, and the combined cytokine alteration
had a lower M2 peak than the control (C). Fibroblasts had the largest increase in cell count with the
higher TGF-β decay and the cytokine combination perturbations (D). All perturbations resulted in an
increased in SSC count with the largest increase resulting from the combined cytokine alteration (E). All
perturbations except the combined and higher MMP-9 decay resulted in increased capillaries as a result
of additional capillary sprouts (F).

Supplemental

Supplemental Figure 1. Overview of Calibration Methods. Latin hypercube sampling is used to generate 600 unique parameter sets given starting bounds, each of which was run in triplicate. The simulations were filtered given specified criteria (i.e. fitting within experimental bounds for CSA recovery) and then alternative density subtraction (ADS) was used to narrow in the parameter bounds. Partial rank squared correlation coefficient (PRCC) was used to gain insight into model sensitivity and adjust the bounds in case initial parameter bounds were too wide or too narrow. This method also allowed for model rule execution refinement to correct cases that interfere with the dynamics of other cell types.

Supplemental Table 1. Unknown model parameters calibrated using LHS to recapitulate published literature

Parameter	Description	Range Tested	Value
neutroRecruitmentProportion	Number of neutrophils generated at every recruitment as proportion of necrotic cells	(4, 12)	7.8322
secretionUptakeValNeu	Cytokine secreted by neutrophil per step during and after "eating necrosis" also max uptake value	(20, 80)	37.3001
necrosisThreshold	Neutrophils are recruited as long as the necrosis amount is greater than this parameter	(0.75, 0.98)	0.9202
macroRecruitmentProportion	Recruitment of macrophages is proportional to mean MCP-1	(0.1, 0.7)	0.4701

secretionUptakeValMac	Cytokine secreted by macrophages per step, also max uptake value	(0.3, 95)	48.2090
TNFconverter	Value of TNF-α to convert monocyte to M1	(15, 150)	109.9025
macRecruitThres	Required mean MCP-1 to signal recruitment of macrophage	(0.05, 4.5)	1.4289
macRecruitStop	Required mean TGF-β to signal stop of macrophage recruitment	(0.05, 0.4)	0.1406
IL10TransThreshold	Local IL-10 that can induce M1 to M2 transition	(15, 600)	78.5442
phagoThresholdforNeuM2	Phagocytosis threshold for neutrophil apoptosis or M1 to M2 transition	(0, 8)	2.0706
sscActivationThreshold	Level of HGF to activate SSC	(10, 100)	44.7091
recruitmentProportionSSC	Number of SSC recruited every time step is proportional to sum of HGF + MMP-9 – TGF-β means	(0.08, 1)	0.8213
quiescentProb	Probability of SSC apoptosis vs remaining quiescent	(1, 4)	3.3617
secretionUptakeValSSC	Cytokine secreted by SSC per step when activated also max uptake value	(0.5, 200)	44.4650
sscTGFApoptosisThreshold	Amount of TGF-β to induce apoptosis	(50, 190)	79.1563
sscApoptosisProb	Probability of SSC apoptosis with apoptosis signal	(0.2, 0.99)	0.7651
VEGFblockApop	Amount of VEGF-A required to block SSC apoptosis	(60, 190)	80.6055
SSCdivisionThreshold	TNF-α + VEGF-A – TGF-β has to be greater than this for SSC to divide	(50, 400)	126.6268
SSCdiffThreshold	IL-10 – HGF – TNF-α – TGF-β has to be greater than this for SSC diff	(5, 145)	57.9870
sscDivideProb	Probability of SSC to divide if signal is not present	(0.01, 0.6)	0.0972
sscDiffProb	Probability for SSC to differentiate without signal	(0.06, 1)	0.6136
fuseProb	Probability of SSC fusing even if the collagen amount is still low	(0.06, 1)	0.3
quiescentThreshold	SSC will return to quiescence if HGF is lower than this at their location	(0.2, 100)	36.3109

necrosisSecretionVal	Amount of HGF and TGF-β secreted by necrotic cells per step	(0.0005, 35)	21.9514
VEGFsecretionValFib	VEGF-A secreted by fibers per step	(0.02, 0.1)	0.0498
VEGF_MMPthreshold	Level of VEGF-A to trigger sprouting/elongation from non-perfused capillaries	(98, 800)	469.0449
capVEGFUptakeMax	Max uptake of VEGF-A by capillaries	(8, 32)	23.8174
MMPsecretionValCap	MMP-9 secreted as capillary is regenerating/sprouting	(4, 16)	8.2444
NumFibers	Number of fibers nearby must be higher than this to check for capillary distances	(1, 2)	1.3899
sproutProb	Probability of capillary sprouting behavior	(0.08, 0.32)	0.1566
secretionUptakeValFibro	Cytokine secretion from fibroblasts per mcs also max uptake value	(0.3, 77)	9.0436
collagenSecretionValFibro	Collagen secretion from fibroblasts per timestep when it is above ECM	(0.09, 0.24)	0.1388
distNearSSCdiv	If fibroblast is within this distance of a dividing SSC, it will divide	(5, 65)	35
TGFActivationThreshold	Level of TGF-β required to activate fibroblast	(20, 80)	40.3643
TGFMyoThreshold	Level of sustained TGF-β required to turn fibroblast into myofibroblast	(280, 650)	536.7525
fibroblastTNFApoptosisThreshold	Level of TNF-α to initiate fibroblast apoptosis	(100, 550)	310.5323
TGFBlockApoptosisThreshold	Level of TGF-β required to block TNF-initiated apoptosis	(160, 1000)	848.3706
fibroblastApoptosisProb	probability of fibroblast apoptosis if TNF-α and TGF-β conditions are met	(0.65,.99)	0.8423
IL10decay	Decay rate of IL-10	(0.00046, 0.00184)	0.0012
HGFdecay	Decay rate of HGF	(0.000135, 0.0003)	0.000228
TGFdecay	Decay rate of TGF	(0.0005, 0.05)	0.0242
TNFdecay	Decay rate of TNF	(0.0000682, 0.002260)	0.00076272
MCPdecay	Decay rate of MCP-1	(0.00596, 0.02384)	0.0124
MMPdecay	Decay rate of MMP-9	(0.0012, 0.1)	0.0112

VEGFdecay	Decay rate of VEGF-A	(0.000464, 0.005569)	0.0015
MinCapDist	Minimum distance between capillary and another capillary	(2.5, 40)	9.6060
capPlacementThreshold	Threshold for if a capillary will be placed/distance required between fiber and cap	(8, 45)	34.6346
M2prolifProb	Probability of M2 proliferation behavior occurring	(0.145305, 0.60)	0.2
M1HalfLife	Corresponds with rate of M1 apoptosis	(15, 55)	24.6360
M2HalfLife	Corresponds with rate of M2 apoptosis	(30, 70)	58.1243
reperfusionLag	Probability of capillary regaining perfusion	(0.001, 0.98)	0.1

776 **Supplemental Table 2.** Criteria utilized for CaliPro model calibration

Iteration	Criteria
1	End criteria: SSC, M1 macrophages, M2 macrophage, fibroblasts, and neutrophils must be below max threshold at the final timepoint (threshold held constant for all iterations).
2	SSC, CSA, and end criteria: within 4 times the experimental std for SSC and 1 times CSA std.
3	Fibroblast, SSC, CSA, replicate check, and end criteria: within 2.1 times the experimental std for fibroblasts, 2.5 for SSC, and 1 times CSA std. More than 1 replicate of the parameter set must meet the criteria.
4	Fibroblast, SSC, CSA, and end criteria: within 2.2 times the experimental std for fibroblasts and SSC and 1 times CSA std.
5	Fibroblast, SSC, CSA, and end criteria: within 1.5 times the experimental std for fibroblasts, 1.75 for SSC, and 1 times CSA std.
6	Fibroblast, SSC, CSA, and end criteria: within 1.75 times the experimental std for fibroblasts, 1.5 for SSC, and 1 times CSA std.
7	Fibroblast, SSC, CSA, replication, and end criteria: within 2.5 times the experimental std for fibroblasts and SSC, and 1 times CSA std. All replicates are required to pass for the parameter set to pass.

Supplemental Figure 2. PRCC plots for various model outputs over time to illustrate how the significance
of cytokine decay and diffusion parameters varies at different points throughout regeneration. Black dots
indicate statistically significant (P < 0.05) correlation for that timepoint. (A) CSA recovery had correlations
with HGF, TGF-β, and MMP-9 decay. (B) SSC count was correlated with HGF, TGF-β, MMP-9 decay and
MCP-1 and TND diffusion. (C) Fibroblast count was correlated with HGF, TGF-β, MMP-9, and TNF-α
decay. (D) HGF, TGF-β, MMP-9, VGEF decay and MCP-1 diffusion were correlated with the number of
non-perfused capillaries. (E) Myoblast cell count was correlated with HGF, TGF-β, MMP-9, and IL-10
decay. (F) Myocyte cell count was correlated with HGF, TGF-β, and MMP-9 decay and TNF-α diffusion.
(G) HGF and MCP-1 decay as well as MCP-1 diffusion were correlated with neutrophil count. (H) M1
macrophage cell count was correlated with TGF-β, VEGF-A, IL-10, and MCP-1 decay and MCP-1
diffusion. (I) M2 macrophage count was correlated with HGF, TGF-β, MMP-9, TNF-α, VEGF-A, MCP-1
decay and MCP-1 diffusion.

Supplemental Figure 3. Cytokine concentrations are correlated with cell counts and recovery metrics at
various stages of regeneration. There is an optimal MCP-1 concentration that tends to result in higher M1
counts 1 day post injury (A). IL-10 concentration is positively correlated with M2 count 3 days post injury
(B). VEGF-A concentration is negatively correlated with the number of fragmented (non-perfused)
capillaries 5 days post injury (C). Higher TGF-β concentrations tends to result in lower SSC cell count 7
days post injury (D). Fibroblasts cell count is highest at an optimal TNF-α concentration with higher or
lower levels hindering cell count 14 days post injury (E). HGF concentration is positively correlated with
CSA recovery at day 28 post injury but there appears to be a threshold where high HGF is no longer
correlated with increased recovery (F).

Supplemental Table 3. Cytokine perturbations based on PRCC

Perturbation	Original cytokine parameter	Altered cytokine parameter
Decrease HGF decay	0.000228	0.0000776
Increase TGF-β decay	0.0242	0.241899
Increase MMP-9 decay	0.0112	0.101994
Decrease VEGF-A decay	0.0015	0.000249267
Increase MCP-1 diffusion	0.18627	1.76943
Increase MCP-1 decay	0.0124	0.118216

Combination	Above alterations excluding increased MCP-1 decay

Supplemental Figure 4. Non-perfused capillaries for each cytokine perturbation. The combined cytokine
perturbation had the lowest number of non-perfused capillaries and all other perturbations resulted in less
non-perfused capillaries compared to the control.

Supplemental Figure 5. Overview of ABM simulation with different initial histology configuration.

Supplemental Text 1. Cellular-Potts

Cellular-Potts model implementation

In the CPM, individual cells are represented as a collection of pixels on a square lattice. These

computational representations of cells have properties of predefined volume, contact energy

with surrounding cells in their environment, and affinity or aversion to diffusing species that drive

chemotactic behaviors[23]. These properties are defined mathematically in the Effective Energy

function *H* which is evaluated in every computational timestep. Equation (1) represents the

effective energy function that imposes the physical properties of all cells in the agent-based

CPM.

$$\mathcal{H} = \sum J\left(\tau_{(\sigma(i)),\tau_{(\sigma(j))}}\right)\left(1 - \delta_{(\sigma(i)),(\sigma(j))}\right) + \lambda_{volume}\left(V_{cell} - V_{target}\right)^2 \quad (1)$$

The first term describes contact energy of each cell for neighboring cells in its environment

governed by *J* the contact coefficient, where *i,j* denote neighboring lattice sites, τ denotes cell

types, and σ denotes individual cells in the simulation. The delta function localizes contact energy

contributions to cell-cell interfaces. The second term represents a volume constraint scaled by

λ_{volume} on each cell. Motile cells in the simulation that exhibit chemotactic behavior towards a

target have additional terms to drive their movement described in (2).

$$\sum_c \lambda_c\left(\tau(\sigma(y_i, t), t)\right)c(y_i, t)1 + c_{CM}(\sigma(y_i, t), t) \qquad (2)$$

Which models logarithmic chemotaxis by cell type and chemokine field concentration influenced

by chemotaxis parameter λ_c, chemical field concentration *c,* and cell body center-of-mass

measurement c_{CM} from which chemotaxis behaviors are calculated. Diffusion and secretion

behavior of chemical species is governed by the general diffusion equation described in (3).

$$\frac{\partial c}{\partial t} = D\nabla^2 c + kc + S \qquad (3)$$

Where *k* defines the decay constant of diffusive species concentration *c*, for which *D* is the

diffusion constant. The *S* term is an additional term to represent secretion of diffusive species.

To apply these relations to recreate cell movement, the CPM randomly selects a pair of

neighboring pixels and evaluates whether one pixel may copy itself to the location of the other,

called a "copy attempt." The probability of acceptance or rejection of that pixel copy attempt is

governed described by the Boltzmann acceptance function described in (4).

$$\Pr\left(\sigma(y_i, t) \rightarrow \sigma(y_i', t)\right) = e^{-\max\left\{0, \frac{\Delta \mathcal{H}}{\mathcal{H}^*}\right\}}$$ (4)

This function describes the probability of the copy attempt $\sigma(y_i, t) \rightarrow \sigma(y_i', t)$ where pixel y_i' copies

to pixel y_i in the CPM algorithm[23]. The value $\mathcal{H}^*$ represents our *temperature* parameter in the

CPM, or the parameter that affects stochasticity of lattice site copy attempts. $\Delta \mathcal{H}$ is the change in

effective energy due to a lattice site copy attempt. The Cellular-Potts simulation discussed here

was implemented using CompuCell3D (CC3D), an open-source Python-based CPM platform[23].

Cellular-Potts model initialization:

The muscle cross-section geometry was initialized by importing a histology image of (murine

gastrocnemius stained with laminin α2) masked to distinguish between muscle fiber and ECM.

The mask was imported into an initialization CC3D script that defined the muscle fibers, ECM,

and microvasculature to specific Cellular-Potts lattice types and generated a Potts Initialization

File (PIF) that was imported into the ABM as the starting cross-section. This initial configuration

resulted in a square lattice of dimensions 321x417 pixels. The model consisted of 2 layers in the

z axis: a tissue environment layer consisting of muscle, ECM, capillaries, and lymphatic

channels, and an immune cell layer where immune cell agents migrated and interacted in the

"2.5D" model. CPM interaction neighbor order was 4 for typical adhesion energy terms, and 1

for all other CPM Hamiltonian terms. Cellular-Potts model parameters are documented in

Supplemental Table 4, adhesion terms are described in Supplemental Table 5. Non-zero

parameter values for adhesion energies between lattice types were calibrated visually to

prevent aggregation of migratory cell types.

Supplemental Table 4. CPM model parameters

Parameter	Value	Justification

Simulation lattice size	321x417x2	2 z-layers: Tissue microenvironment layer and migratory cell layer
Metropolis Algorithm Temperature	10	This is a default, numerically stable temperature for CPM models described in CompuCell3D
Neighbor Order	1	Interactions spanning 1 pixel distance for Cellular-Potts cell agents was sufficient for agent-based behaviors

859   **Supplemental Table 5.** CPM agent Adhesion parameters

CPM Lattice Interaction	Parameter Value
Medium - Medium	0
Medium - Fiber	0
Medium - ECM	0
Medium - SSC	0
Medium - Capillary	0
Medium - Neutrophil	0
Medium - Macrophage	0
Medium - Necrotic	0
Medium - Wall	0
Medium - Lymphatic	0
Fiber - Fiber	0
Fiber - ECM	0
Fiber - SSC	35
Fiber - Capillary	0
Fiber - Neutrophil	35
Fiber - Macrophage	20
Fiber - Necrotic	0
Fiber - Wall	0
Fiber - Lymphatic	0
ECM - ECM	0

ECM - SSC	30
ECM - Capillary	0
ECM - Neutrophil	30
ECM - Macrophage	20
ECM - Necrotic	0
ECM - Wall	0
ECM - Lymphatic	0
SSC - SSC	35
SSC - Capillary	30
SSC - Neutrophil	15
SSC - Macrophage	15
SSC - Necrotic	30
SSC - Wall	20
SSC - Lymphatic	10
Capillary - Capillary	0
Capillary - Neutrophil	35
Capillary - Macrophage	20
Capillary - Necrotic	0
Capillary - Wall	0
Capillary - Lymphatic	0
Neutrophil - Neutrophil	35
Neutrophil - Macrophage	15
Neutrophil - Necrotic	35
Neutrophil - Wall	20
Neutrophil - Lymphatic	10
Macrophage - Macrophage	15
Macrophage - Necrotic	20
Macrophage - Wall	20
Macrophage - Lymphatic	10
Necrotic - Necrotic	0

Necrotic - Wall	0
Necrotic - Lymphatic	0
Wall - Wall	0
Wall - Lymphatic	0
Lymphatic - Lymphatic	0
Fibroblast - Medium	0
Fibroblast - Fiber	20
Fibroblast - ECM	20
Fibroblast - SSC	15
Fibroblast - Capillary	20
Fibroblast - Neutrophil	15
Fibroblast - Macrophage	15
Fibroblast - Necrotic	20
Fibroblast - Wall	20
Fibroblast - Lymphatic	10
Fibroblast - Fibroblast	15
Neighbor Order	4

862 **Supplemental Table 6.** Experimental data description for model comparison

Experimental data description and relevant figure from literature	Model comparison
CTX injury to mouse hindlimb muscle and quantification of average cross-sectional area (μm^2) of individual muscle fibers for a given animal was determined after manually outlining individual muscle fibers in each of the digitally captured images (Fig. 3B)[102]	Muscle CSA recovery calibration (Fig. 3A)
BaCl$_2$ injury to mouse hindlimb and IF quantification of SSC (Pax7+) and fibroblasts (Tcf4+) cells (Fig 2DD & EE)[76]	SSC and fibroblast count calibration (Fig. 3B & C)
NTX injury to mouse hindlimb and macrophage quantification based on number of F4/80+ cells (Fig. 4E)[97]	Total macrophage count validation (Fig. 3D)
CTX injury to mouse hindlimb muscle and capillary density was calculated using biotinylated lectin and H&E (Fig. 6B)[102]	Capillaries/mm^2 validation (Fig. 3H)

BaCl$_2$ injury to mouse hindlimb and flow cytometry to assess CD45$^+$/F4/80$^+$/Ly6C^{hi} (M1) and CD45$^+$/F4/80$^+$/Ly6C^{lo} (M2) macrophages (Fig. 2A)[107]	M1 and M2 cell count validation (Fig. 3E & H)
CTX injury to mouse hindlimb muscle and IHC analysis was used to label neutrophils using Ly6G antibody (Fig. 3A)[108]	Neutrophil cell count validation (Fig. 3G)
Bothrops asper snake venom induced hindlimb injury to mice pretreated with either an antimouse granulocyte rat monoclonal immunoglobulin G (IgG) antibody or with isotype-matched control antibody. Inflammatory infiltrate was quantified from muscle tissue (Fig. 7A) and the extent of regeneration was quantitatively assessed by determining the percentage of the area comprised of regenerating muscle fibers in relation to the total area of muscle damage[109]	Neutrophil depletion perturbation (Fig. 4)
IL-4–conjugated gold nanoparticles were injected into murine skeletal muscle following hindlimb ischemia. M1 and M2 cells were quantified with flow cytometry (Fig. 6 A, C, and E) and muscle force was measured for quantification of regeneration (Fig. 4G)[164]	Anti-inflammatory nanoparticles perturbation (Fig. 4)
IL-10 null mice underwent hindlimb unloading and reloading to induce muscle damage. Muscle CSA (Fig. 6C), M1 and M2 cells (Fig. 3A-B), and SSC differentiation markers were quantified (Fig. 3C-F)[163]	IL-10 KO perturbation (Fig. 4)
Noetoxin or freeze injury to CXCL12Gagtm/Gagtm and control mouse hindlimb muscle with H&E for assessment of muscle regeneration (Fig. 3), transcriptome analysis to asses collagen organization and synthesis (Fig. 2C), and immune cell infiltration (Fig. 4S)[111]	Hindered angiogenesis perturbation (Fig. 4)
Following CTX injury to mouse hindlimb muscle and injection of intramuscular administration of rAAV vectors injury area was quantified for various doses of AAV-VEGF (Fig. 3D) and H&E was used to assess muscle recovery (Fig. 5)[73]	VEGF-A injection perturbation (Fig. 4)
CCL2-/- and control mice received BaCl$_2$ injection to induce acute injury. CSA quantification was used to assess muscle regeneration (Fig. 1I). Flow cytometry was used to quantify macrophage (Fig. 2A), and neutrophil (Fig. 2B) cells[115]	MCP-1 KO perturbation (Fig. 4)
CTX induced injury to control and TNF-α receptor double-knockout mice. Force was measured to quantify muscle recovery (Fig. 7), immunohistochemical staining of Mac-1 for quantifying immune cells (Fig. 6B), and western blot analysis was used to assess SSC differentiation (Fig. 2)[165]	TNF-α KO perturbation (Fig. 4)
Mice were treated with clodronate-containing or control liposomes and underwent a contusion injury. H&E was used for muscle regeneration assessment (Fig. 2), HGF (Fig. 4A) and TGF (Fig. 5A) mRNA were quantified[110]	Macrophage depletion perturbation (Fig. 4)

References

1. Quintero AJ, Wright VJ, Fu FH, Huard J. Stem Cells for the Treatment of Skeletal Muscle Injury Sports injury Stem cells Tissue engineering Fibrosis Regeneration Skeletal muscle. *Clin Sports Med.* 2009;28:1-11. doi:10.1016/j.csm.2008.08.009

2. Barroso GC, Thiele ES. MUSCLE INJURIES IN ATHLETES. *Rev Bras Ortop (English Ed.*

870   2011;46(4):354-358. doi:10.1016/s2255-4971(15)30245-7

3.   Valle X. Clinical practice guide for muscular injuries: epidemiology, diagnosis, treatment and prevention. *Br J Sports Med*. 2011;45(2):e2-e2. doi:10.1136/bjsm.2010.081570.20

4.   Järvinen TAH, Järvinen TLN, Kääriäinen M, et al. Muscle injuries: optimising recovery. *Best Pract Res Clin Rheumatol*. 2007;21(2):317-331. doi:10.1016/j.berh.2006.12.004

5.   Huard J, Li Y, Fu FH. Muscle injuries and repair: Current trends in research. *J Bone Jt Surg - Ser A*. Published online 2002. doi:10.2106/00004623-200205000-00022

6.   Forcina L, Cosentino M, Musarò A. Mechanisms Regulating Muscle Regeneration: Insights into the Interrelated and Time-Dependent Phases of Tissue Healing. *Cells*. 2020;9(5). doi:10.3390/cells9051297

7.   Howard EE, Pasiakos SM, Blesso CN, Fussell MA, Rodriguez NR. Divergent Roles of Inflammation in Skeletal Muscle Recovery From Injury. *Front Physiol*. 2020;11:87. doi:10.3389/fphys.2020.00087

8.   Westman AM, Peirce SM, Christ GJ, Blemker SS. Agent-based model provides insight into the mechanisms behind failed regeneration following volumetric muscle loss injury. Merks RMH, ed. *PLOS Comput Biol*. 2021;17(5):e1008937. doi:10.1371/journal.pcbi.1008937

9.   Wagatsuma A. Endogenous expression of angiogenesis-related factors in response to muscle injury. *Mol Cell Biochem 2006 2981*. 2006;298(1):151-159. doi:10.1007/S11010-006-9361-X

10.   Chen CH, Chang CH, Lee CH. Usage of Growth Factors in Acute Muscle Injuries. In: *Sports Injuries*. ; 2015:2343-2351. doi:10.1007/978-3-642-36569-0_174

11.   Husmann I, Soulet L, Gautron J, Martelly I, Barritault D. Growth factors in skeletal muscle regeneration. *Cytokine Growth Factor Rev*. 1996;7(3):249-258. doi:10.1016/S1359-6101(96)00029-9

12.   Itoh Y. Proteolytic modulation of tumor microenvironment signals during cancer progression. *Front Oncol*. 2022;12. doi:10.3389/FONC.2022.935231

13.   Ciano-Petersen NL, Muñiz-Castrillo S, Birzu C, et al. Cytokine dynamics and targeted immunotherapies in autoimmune encephalitis. *Brain Commun*. 2022;4(4):1-16. doi:10.1093/braincomms/fcac196

14.   Ferrara N. Binding to the Extracellular Matrix and Proteolytic Processing: Two Key Mechanisms Regulating Vascular Endothelial Growth Factor Action. *Mol Biol Cell*. 2010;21(5):687. doi:10.1091/MBC.E09-07-0590

15.   Virgilio KM, Martin KS, Peirce SM, Blemker SS. Agent-based model illustrates the role of the microenvironment in regeneration in healthy and mdx skeletal muscle. *J Appl Physiol*. 2018;125(5):1424-1439. doi:10.1152/japplphysiol.00379.2018

16.   Martin KS, Virgilio KM, Peirce SM, Blemker SS. Computational modeling of muscle regeneration and adaptation to advance muscle tissue regeneration strategies. *Cells Tissues Organs*. 2016;202(3-4):250-266. doi:10.1159/000443635

17.   Khuu S, Fernandez JW, Handsfield GG. A Coupled Mechanobiological Model of Muscle Regeneration In Cerebral Palsy. *Front Bioeng Biotechnol*. 2021;9.

911   doi:10.3389/fbioe.2021.689714

18. Khuu S, Fernandez JW, Handsfield GG. Delayed skeletal muscle repair following inflammatory damage in simulated agent-based models of muscle regeneration. Peirce SM, ed. *PLOS Comput Biol*. 2023;19(4):e1011042. doi:10.1371/JOURNAL.PCBI.1011042

19. Virgilio KM, Jones BK, Miller EY, et al. Computational Models Provide Insight into In Vivo Studies and Reveal the Complex Role of Fibrosis in mdx Muscle Regeneration. *Ann Biomed Eng*. Published online 2020. doi:10.1007/s10439-020-02566-1

20. Joslyn LR, Kirschner DE, Linderman JJ. CaliPro: A Calibration Protocol That Utilizes Parameter Density Estimation to Explore Parameter Space and Calibrate Complex Biological Models. *Cell Mol Bioeng*. 2021;14(1):31-47. doi:10.1007/s12195-020-00650-z

21. Martin KS, Blemker SS, Peirce SM. Agent-based computational model investigates muscle-specific responses to disuse-induced atrophy. *J Appl Physiol*. 2015;118(10):1299-1309. doi:10.1152/japplphysiol.01150.2014

22. Gianlupi JF, Mapder T, Sego TJ, et al. Multiscale Model of Antiviral Timing, Potency, and Heterogeneity Effects on an Epithelial Tissue Patch Infected by SARS-CoV-2. *Viruses 2022, Vol 14, Page 605*. 2022;14(3):605. doi:10.3390/V14030605

23. Swat MH, Thomas GL, Belmonte JM, Shirinifard A, Hmeljak D, Glazier JA. Multi-Scale Modeling of Tissues Using CompuCell3D. *Methods Cell Biol*. 2012;110:325. doi:10.1016/B978-0-12-388403-9.00013-8

24. Stephenson ER, Kojouharov H V. A mathematical model of skeletal muscle regeneration. *Math Methods Appl Sci*. 2018;41(18):8589-8602. doi:10.1002/mma.4908

25. Waldemer-Streyer RJ, Kim D, Chen J. Muscle cell-derived cytokines in skeletal muscle regeneration. *FEBS J*. 2022;289(21):6463-6483. doi:10.1111/febs.16372

26. Rucavado A, Escalante T, Teixeira CFP, Fernándes CM, Díaz C, Gutiérrez JM. Increments in cytokines and matrix metalloproteinases in skeletal muscle after injection of tissue-damaging toxins from the venom of the snake Bothrops asper. *Mediators Inflamm*. 2002;11(2):121-128. doi:10.1080/09629350220131980

27. Yin H, Price F, Rudnicki MA. Satellite cells and the muscle stem cell niche. *Physiol Rev*. 2013;93(1):23-67. doi:10.1152/PHYSREV.00043.2011

28. Madaro L, Bouché M. From innate to adaptive immune response in muscular dystrophies and skeletal muscle regeneration: The role of lymphocytes. *Biomed Res Int*. 2014;2014. doi:10.1155/2014/438675

29. Wang X, Hossain M, Bogoslowski A, Kubes P, Irimia D. Chemotaxing neutrophils enter alternate branches at capillary bifurcations. *Nat Commun 2020 111*. 2020;11(1):1-12. doi:10.1038/s41467-020-15476-6

30. Butterfield TA, Best TM, Merrick MA. The dual roles of neutrophils and macrophages in inflammation: A critical balance between tissue damage and repair. *J Athl Train*. 2006;41(4):457-465. doi:10.1016/s0162-0908(08)79217-1

31. Wang J. Neutrophils in tissue injury and repair. *Cell Tissue Res*. 2018;371(3):531-539. doi:10.1007/s00441-017-2785-7

32. Soehnlein O, Zernecke A, Eriksson EE, et al. Neutrophil secretion products pave the way
for inflammatory monocytes. *Blood.* 2008;112(4):1461. doi:10.1182/BLOOD-2008-02-
139634

33. Fox S, Leitch AE, Duffin R, Haslett C, Rossi AG. Neutrophil Apoptosis: Relevance to the
Innate Immune Response and Inflammatory Disease. *J Innate Immun.* 2010;2(3):216.
doi:10.1159/000284367

34. Oishi Y, Manabe I. Macrophages in inflammation, repair and regeneration. *Int Immunol.*
2018;30(11):511-528. doi:10.1093/INTIMM/DXY054

35. Elkington PT, Green JA, Friedland JS. Analysis of matrix metalloproteinase secretion by
macrophages. *Methods Mol Biol.* 2009;531:253-265. doi:10.1007/978-1-59745-396-7_16

36. Chen GY, Nuñez G. Sterile inflammation: sensing and reacting to damage. *Nat Rev*
Immunol 2010 1012. 2010;10(12):826-837. doi:10.1038/nri2873

37. Lacy P, Stow JL. Cytokine release from innate immune cells: association with diverse
membrane trafficking pathways. *Blood.* 2011;118(1):9-18. doi:10.1182/BLOOD-2010-08-
265892

38. Vogel DYS, Heijnen PDAM, Breur M, et al. Macrophages migrate in an activation-
dependent manner to chemokines involved in neuroinflammation. *J Neuroinflammation.*
2014;11:23. doi:10.1186/1742-2094-11-23

39. Molnarfi N, Benkhoucha M, Funakoshi H, Nakamura T, Lalive PH. Hepatocyte growth
factor: A regulator of inflammation and autoimmunity. *Autoimmun Rev.* 2015;14(4):293-
303. doi:10.1016/J.AUTREV.2014.11.013

40. Furrer R, Handschin C. Optimized engagement of macrophages and satellite cells in the
repair and regeneration of exercised muscle. In: *Research and Perspectives in Endocrine*
Interactions. Springer Verlag; 2017:57-66. doi:10.1007/978-3-319-72790-5_5

41. Kratofil RM, Kubes P, Deniset JF. Monocyte conversion during inflammation and injury.
Arterioscler Thromb Vasc Biol. 2017;37(1):35-42. doi:10.1161/ATVBAHA.116.308198

42. Chazaud B, Sonnet C, Lafuste P, et al. Satellite cells attract monocytes and use
macrophages as a support to escape apoptosis and enhance muscle growth. *J Cell Biol.*
2003;163(5):1133. doi:10.1083/JCB.200212046

43. Owen JL, Mohamadzadeh M. Macrophages and chemokines as mediators of
angiogenesis. *Front Physiol.* 2013;4. doi:10.3389/FPHYS.2013.00159

44. Reibman J, Meixler S, Lee TC, et al. Transforming growth factor beta 1, a potent
chemoattractant for human neutrophils, bypasses classic signal-transduction pathways.
Proc Natl Acad Sci U S A. 1991;88(15):6805. doi:10.1073/PNAS.88.15.6805

45. Greenlee-Wacker MC. Clearance of apoptotic neutrophils and resolution of inflammation.
Immunol Rev. 2016;273(1):357. doi:10.1111/IMR.12453

46. Watanabe S, Alexander M, Misharin A V., Budinger GRS. The role of macrophages in the
resolution of inflammation. *J Clin Invest.* 2019;129(7):2619. doi:10.1172/JCI124615

47. Uribe-Querol E, Rosales C. Phagocytosis: Our Current Understanding of a Universal
Biological Process. *Front Immunol.* 2020;11:1066. doi:10.3389/fimmu.2020.01066

48. Yoon YS, Lee YJ, Choi YH, Park YM, Kang JL. Macrophages programmed by apoptotic

cells inhibit epithelial-mesenchymal transition in lung alveolar epithelial cells via PGE2,
PGD2, and HGF. *Sci Reports 2016 61*. 2016;6(1):1-18. doi:10.1038/srep20992

49. D'Angelo F, Bernasconi E, Schäfer M, et al. Macrophages promote epithelial repair
through hepatocyte growth factor secretion. *Clin Exp Immunol*. 2013;174(1):60.
doi:10.1111/CEI.12157

50. Popov Y, Sverdlov DY, Bhaskar KR, et al. Macrophage-mediated phagocytosis of
apoptotic cholangiocytes contributes to reversal of experimental biliary fibrosis. *Am J
Physiol - Gastrointest Liver Physiol*. 2010;298(3):G323. doi:10.1152/AJPGI.00394.2009

51. Arnold L, Henry A, Poron F, et al. Inflammatory monocytes recruited after skeletal muscle
injury switch into antiinflammatory macrophages to support myogenesis. *J Exp Med*.
2007;204(5):1057-1069. doi:10.1084/jem.20070075

52. Chung EY, Liu J, Homma Y, et al. Interleukin-10 Expression in Macrophages during
Phagocytosis of Apoptotic Cells Is Mediated by the TALE homeoproteins Pbx-1 and
Prep-1. *Immunity*. 2007;27(6):952. doi:10.1016/J.IMMUNI.2007.11.014

53. Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev
Immunol 2008 812*. 2008;8(12):958-969. doi:10.1038/nri2448

54. Saini J, McPhee JS, Al-Dabbagh S, Stewart CE, Al-Shanti N. Regenerative function of
immune system: Modulation of muscle stem cells. *Ageing Res Rev*. 2016;27:67-76.
doi:10.1016/J.ARR.2016.03.006

55. Das A, Sinha M, Datta S, et al. Monocyte and Macrophage Plasticity in Tissue Repair and
Regeneration. *Am J Pathol*. 2015;185(10):2596. doi:10.1016/J.AJPATH.2015.06.001

56. Arabpour M, Saghazadeh A, Rezaei N. Anti-inflammatory and M2 macrophage
polarization-promoting effect of mesenchymal stem cell-derived exosomes. *Int
Immunopharmacol*. 2021;97:107823. doi:10.1016/J.INTIMP.2021.107823

57. Reimann J, Irintchev A, Wernig A. Regenerative capacity and the number of satellite cells
in soleus muscles of normal and mdx mice. *Neuromuscul Disord*. 2000;10(4-5):276-282.
doi:10.1016/S0960-8966(99)00118-2

58. Kawamura K, Takano K, Suetsugu S, et al. N-WASP and WAVE2 acting downstream of
phosphatidylinositol 3-kinase are required for myogenic cell migration induced by
hepatocyte growth factor. *J Biol Chem*. 2004;279(52):54862-54871.
doi:10.1074/JBC.M408057200

59. Wang W, Pan H, Murray K, Jefferson BS, Li Y. Matrix metalloproteinase-1 promotes
muscle cell migration and differentiation. *Am J Pathol*. 2009;174(2):541-549.
doi:10.2353/AJPATH.2009.080509

60. Allen RE, Boxhorn LK. Regulation of skeletal muscle satellite cell proliferation and
differentiation by transforming growth factor-beta, insulin-like growth factor I, and
fibroblast growth factor. *J Cell Physiol*. 1989;138(2):311-315.
doi:10.1002/JCP.1041380213

61. González MN, de Mello W, Butler-Browne GS, et al. HGF potentiates extracellular matrix-
driven migration of human myoblasts: Involvement of matrix metalloproteinases and
MAPK/ERK pathway. *Skelet Muscle*. 2017;7(1):1-13. doi:10.1186/s13395-017-0138-6

62. Allen RE, Sheehan SM, Taylor RG, Kendall TL, Rice GM. Hepatocyte growth factor

activates quiescent skeletal muscle satellite cells in vitro. *J Cell Physiol.* 1995;165(2):307-
312. doi:10.1002/JCP.1041650211

63.   Miller KJ, Thaloor D, Matteson S, Pavlath GK. Hepatocyte growth factor affects satellite
cell activation and differentiation in regenerating skeletal muscle. *Am J Physiol Cell*
Physiol. 2000;278(1). doi:10.1152/AJPCELL.2000.278.1.C174

64.   Tatsumi R, Anderson JE, Nevoret CJ, Halevy O, Allen RE. HGF/SF is present in normal
adult skeletal muscle and is capable of activating satellite cells. *Dev Biol.*
1998;194(1):114-128. doi:10.1006/DBIO.1997.8803

65.   Cooper RN, Tajbakhsh S, Mouly V, Cossu G, Buckingham M, Butler-Browne GS. In vivo
satellite cell activation via Myf5 and MyoD in regenerating mouse skeletal muscle. *J Cell*
Sci. 1999;112 (Pt 17)(17):2895-2901. doi:10.1242/JCS.112.17.2895

66.   Flamini V, Ghadiali RS, Antczak P, Rothwell A, Turnbull JE, Pisconti A. The Satellite Cell
Niche Regulates the Balance between Myoblast Differentiation and Self-Renewal via p53.
Stem Cell Reports. 2018;10(3):970-983. doi:10.1016/J.STEMCR.2018.01.007

67.   Bentzinger CF, Wang YX, Rudnicki MA. Building Muscle: Molecular Regulation of
Myogenesis. *Cold Spring Harb Perspect Biol.* 2012;4(2):a008342.
doi:10.1101/CSHPERSPECT.A008342

68.   Wang YX, Dumont NA, Rudnicki MA. Muscle stem cells at a glance. *J Cell Sci.*
2014;127(Pt 21):4543-4548. doi:10.1242/JCS.151209

69.   Nguyen JH, Chung JD, Lynch GS, Ryall JG. The Microenvironment Is a Critical Regulator
of Muscle Stem Cell Activation and Proliferation. *Front Cell Dev Biol.* 2019;7:254.
doi:10.3389/fcell.2019.00254

70.   Ruiz-Gómez M, Coutts N, Suster ML, Landgraf M, Bate M. myoblasts incompetent
encodes a zinc finger transcription factor required to specify fusion-competent myoblasts
in Drosophila. *Development.* 2002;129(1):133-141. doi:10.1242/DEV.129.1.133

71.   Abmayr SM, Pavlath GK. Myoblast fusion: lessons from flies and mice. *Development.*
2012;139(4):641-656. doi:10.1242/DEV.068353

72.   Isesele PO, Mazurak VC. Regulation of Skeletal Muscle Satellite Cell Differentiation by
Omega-3 Polyunsaturated Fatty Acids: A Critical Review. *Front Physiol.* 2021;12:750.
doi:10.3389/fphys.2021.682091

73.   Arsic N, Zacchigna S, Zentilin L, et al. Vascular endothelial growth factor stimulates
skeletal muscle regeneration in Vivo. *Mol Ther.* 2004;10(5):844-854.
doi:10.1016/J.YMTHE.2004.08.007

74.   Sonnet C, Lafuste P, Arnold L, et al. Human macrophages rescue myoblasts and
myotubes from apoptosis through a set of adhesion molecular systems. *J Cell Sci.*
2006;119(Pt 12):2497-2507. doi:10.1242/JCS.02988

75.   Cencetti F, Bernacchioni C, Tonelli F, Roberts E, Donati C, Bruni P. TGFβ1 evokes
myoblast apoptotic response via a novel signaling pathway involving S1P4 transactivation
upstream of Rho-kinase-2 activation. *FASEB J.* 2013;27(11):4532-4546.
doi:10.1096/FJ.13-228528

76.   Murphy MM, Lawson JA, Mathew SJ, Hutcheson DA, Kardon G. Satellite cells,
connective tissue fibroblasts and their interactions are crucial for muscle regeneration.

Development. 2011;138(17):3625-3637. doi:10.1242/dev.064162

77.   Beanes SR, Dang C, Soo C, Ting K. Skin repair and scar formation: the central role of
TGF-β. *Expert Rev Mol Med*. 2003;5(8):1-22. doi:10.1017/S1462399403005817

78.   Chellini F, Tani A, Zecchi-Orlandini S, Sassoli C. Influence of platelet-rich and platelet-
poor plasma on endogenous mechanisms of skeletal muscle repair/regeneration. *Int J*
Mol Sci. 2019;20(3). doi:10.3390/ijms20030683

79.   Dickinson RB, Guido S, Tranquillo RT. Biased cell migration of fibroblasts exhibiting
contact guidance in oriented collagen gels. *Ann Biomed Eng*. 1994;22(4):342-356.
doi:10.1007/BF02368241

80.   Zou Y, Zhang RZ, Sabatelli P, Chu ML, Bönnemann CG. Muscle interstitial fibroblasts are
the main source of collagen VI synthesis in skeletal muscle: implications for congenital
muscular dystrophy types Ullrich and Bethlem. *J Neuropathol Exp Neurol*.
2008;67(2):144-154. doi:10.1097/NEN.0B013E3181634EF7

81.   Sanderson RD, Fitch JM, Linsenmayer TR, Mayne R. Fibroblasts promote the formation
of a continuous basal lamina during myogenesis in vitro. *J Cell Biol*. 1986;102(3):740.
doi:10.1083/JCB.102.3.740

82.   Yokoyama T, Sekiguchi K, Tanaka T, et al. Angiotensin II and mechanical stretch induce
production of tumor necrosis factor in cardiac fibroblasts. *Am J Physiol*. 1999;276(6).
doi:10.1152/AJPHEART.1999.276.6.H1968

83.   Skutek M, Van Griensven M, Zeichen J, Brauer N, Bosch U. Cyclic mechanical stretching
modulates secretion pattern of growth factors in human tendon fibroblasts. *Eur J Appl*
Physiol. 2001;86(1):48-52. doi:10.1007/S004210100502

84.   Lindner D, Zietsch C, Becher PM, et al. Differential expression of matrix metalloproteases
in human fibroblasts with different origins. *Biochem Res Int*. Published online 2012.
doi:10.1155/2012/875742

85.   Newman AC, Nakatsu MN, Chou W, Gershon PD, Hughes CCW. The requirement for
fibroblasts in angiogenesis: Fibroblast-derived matrix proteins are essential for
endothelial cell lumen formation. *Mol Biol Cell*. 2011;22(20):3791-3800.
doi:10.1091/mbc.E11-05-0393

86.   Desmouliere A, Geinoz A, Gabbiani F, Gabbiani G. Transforming growth factor-beta 1
induces alpha-smooth muscle actin expression in granulation tissue myofibroblasts and in
quiescent and growing cultured fibroblasts. *J Cell Biol*. 1993;122(1):103-111.
doi:10.1083/JCB.122.1.103

87.   Wipff PJ, Rifkin DB, Meister JJ, Hinz B. Myofibroblast contraction activates latent TGF-
beta1 from the extracellular matrix. *J Cell Biol*. 2007;179(6):1311-1323.
doi:10.1083/JCB.200704042

88.   Petrov V V., Fagard RH, Lijnen PJ. Stimulation of Collagen Production by Transforming
Growth Factor-β1 During Differentiation of Cardiac Fibroblasts to Myofibroblasts.
Hypertension. 2002;39(2 I):258-263. doi:10.1161/HY0202.103268

89.   Lemos DR, Babaeijandaghi F, Low M, et al. Nilotinib reduces muscle fibrosis in chronic
muscle injury by promoting TNF-mediated apoptosis of fibro/adipogenic progenitors. *Nat*
Med. 2015;21(7):786-794. doi:10.1038/NM.3869

90. Kim J, Lee J. Role of transforming growth factor-β in muscle damage and regeneration:
focused on eccentric muscle contraction. *J Exerc Rehabil*. 2017;13(6):621.
doi:10.12965/JER.1735072.536

91. Huey KA. Potential Roles of Vascular Endothelial Growth Factor during Skeletal Muscle
Hypertrophy. *Exerc Sport Sci Rev*. 2018;46(3):195-202.
doi:10.1249/JES.0000000000000152

92. Snijders T, Nederveen JP, McKay BR, et al. Satellite cells in human skeletal muscle
plasticity. *Front Physiol*. 2015;6(OCT):283. doi:10.3389/fphys.2015.00283

93. Wickler SJ. Capillary supply of skeletal muscles from acclimatized white-footed mice
Peromyscus. *Am J Physiol*. 1981;241(5):357-361.
doi:10.1152/AJPREGU.1981.241.5.R357

94. Gehlert S, Theis C, Weber S, et al. Exercise-induced decline in the density of lyve-1-
positive lymphatic vessels in human skeletal muscle. *Lymphat Res Biol*. 2010;8(3):165-
173. doi:10.1089/LRB.2009.0035

95. Jacobsen NL, Norton CE, Shaw RL, Cornelison D, Segal S, Segal SS. Myofiber injury
induces capillary disruption and regeneration of disorganized microvascular networks.
bioRxiv. Published online August 4, 2021:2021.08.02.454805.
doi:10.1101/2021.08.02.454805

96. Frey SP, Jansen H, Raschke MJ, Meffert RH, Ochman S. VEGF improves skeletal
muscle regeneration after acute trauma and reconstruction of the limb in a rabbit model.
Clin Orthop Relat Res. 2012;470(12):3607-3614. doi:10.1007/S11999-012-2456-7

97. Hardy D, Besnard A, Latil M, et al. Comparative Study of Injury Models for Studying
Muscle Regeneration in Mice. *PLoS One*. 2016;11(1):e0147198.
doi:10.1371/JOURNAL.PONE.0147198

98. Haas TL, Milkiewicz M, Davis SJ, et al. Matrix metalloproteinase activity is required for
activity-induced angiogenesis in rat skeletal muscle. *Am J Physiol Heart Circ Physiol*.
2000;279(4). doi:10.1152/AJPHEART.2000.279.4.H1540

99. Hampton HR, Chtanova T. Lymphatic Migration of Immune Cells. *Front Immunol*.
2019;10(MAY):1168. doi:10.3389/FIMMU.2019.01168

100. ExternalPotential Plugin — CC3D Reference Manual 4.4.1 documentation. Accessed
January 4, 2024.
https://compucell3dreferencemanual.readthedocs.io/en/latest/external_potential_plugin.ht
ml

101. Filion RJ, Popel AS. Intracoronary administration of FGF-2: A computational model of
myocardial deposition and retention. *Am J Physiol - Hear Circ Physiol*. 2005;288(1 57-
1):263-279. doi:10.1152/ajpheart.00205.2004

102. Ochoa O, Sun D, Reyes-Reyna SM, et al. Delayed angiogenesis and VEGF production in
CCR2-/- mice during impaired skeletal muscle regeneration. *Am J Physiol - Regul Integr
Comp Physiol*. 2007;293(2):651-661. doi:10.1152/ajpregu.00069.2007

103. Pratt SJP, Shah SB, Ward CW, Kerr JP, Stains JP, Lovering RM. Recovery of altered
neuromuscular junction morphology and muscle function in mdx mice after injury. *Cell
Mol Life Sci*. 2014;72(1):153-164. doi:10.1007/s00018-014-1663-7

104. You JS, Barai P, Chen J. Sex differences in skeletal muscle size, function, and myosin
heavy chain isoform expression during post-injury regeneration in mice. *Physiol Rep*.
2023;11(16). doi:10.14814/PHY2.15791

105. Marino S, Hogue IB, Ray CJ, Kirschner DE. A methodology for performing global
uncertainty and sensitivity analysis in systems biology. *J Theor Biol*. 2008;254(1):178-
196. doi:10.1016/J.JTBI.2008.04.011

106. Segovia-Juarez JL, Ganguli S, Kirschner D. Identifying control mechanisms of granuloma
formation during M. tuberculosis infection using an agent-based model. *J Theor Biol*.
2004;231(3):357-376. doi:10.1016/J.JTBI.2004.06.031

107. Wang X, Zhao W, Ransohoff RM, Zhou L. Infiltrating macrophages are broadly activated
at the early stage to support acute skeletal muscle injury repair. *J Neuroimmunol*.
2018;317:55. doi:10.1016/J.JNEUROIM.2018.01.004

108. Nguyen MH, Cheng M, Koh TJ. Impaired Muscle Regeneration in Ob/ob and Db/db Mice.
Sci World J. 2011;11:1525. doi:10.1100/TSW.2011.137

109. Teixeira CFP, Zamunér SR, Zuliani JP, et al. Neutrophils do not contribute to local tissue
damage, but play a key role in skeletal muscle regeneration, in mice injected with
Bothrops asper snake venom. *Muscle Nerve*. 2003;28(4):449-459.
doi:10.1002/MUS.10453

110. Liu X, Liu Y, Zhao L, Zeng Z, Xiao W, Chen P. Macrophage depletion impairs skeletal
muscle regeneration: The roles of regulatory factors for muscle regeneration. *Cell Biol Int*.
2017;41(3):228-238. doi:10.1002/CBIN.10705

111. Hardy D, Fefeu M, Besnard A, et al. Defective angiogenesis in CXCL12 mutant mice
impairs skeletal muscle regeneration. *Skelet Muscle*. 2019;9(1):1-15.
doi:10.1186/s13395-019-0210-5

112. Cantaert T, Baeten D, Tak PP, van Baarsen LGM. Type I IFN and TNFα cross-regulation
in immune-mediated inflammatory disease: Basic concepts and clinical relevance.
Arthritis Res Ther. 2010;12(5):1-10. doi:10.1186/AR3150/TABLES/2

113. Kunze KN, Hannon CP, Fialkoff JD, Frank RM, Cole BJ. Platelet-rich plasma for muscle
injuries: A systematic review of the basic science literature. *World J Orthop*.
2019;10(7):278-291. doi:10.5312/wjo.v10.i7.278

114. Choi W, Lee J, Lee J, Lee SH, Kim S. Hepatocyte growth factor regulates macrophage
transition to the M2 phenotype and promotes murine skeletal muscle regeneration. *Front
Physiol*. 2019;10(JUL). doi:10.3389/fphys.2019.00914

115. Lu H, Huang D, Ransohoff RM, Zhou L. Acute skeletal muscle injury: CCL2 expression
by both monocytes and injured muscle is required for repair. *FASEB J*.
2011;25(10):3344-3355. doi:10.1096/FJ.10-178939

116. Girardi F, Taleb A, Ebrahimi M, et al. TGFβ signaling curbs cell fusion and muscle
regeneration. *Nat Commun*. 2021;12(1). doi:10.1038/s41467-020-20289-8

117. Zimowska M, Olszynski KH, Swierczynska M, Streminska W, Ciemerych MA. Decrease
of MMP-9 activity improves soleus muscle regeneration. *Tissue Eng - Part A*.
2012;18(11-12):1183-1192. doi:10.1089/ten.tea.2011.0459

118. Dziki JL, Velayutham M, Hussey GS, Turnquist HR. Cytokine networks in immune-

mediated muscle regeneration. *J Immunol Regen Med*. 2018;1:32-44. doi:10.1016/j.regen.2018.03.001

119. Fujita R, Kawano F, Ohira T, et al. Anti-interleukin-6 receptor antibody (MR16-1) promotes muscle regeneration via modulation of gene expressions in infiltrated macrophages. *Biochim Biophys Acta - Gen Subj*. 2014;1840(10):3170-3180. doi:10.1016/j.bbagen.2014.01.014

120. Ismail AA, Shaker BT, Bajou K. The plasminogen–activator plasmin system in physiological and pathophysiological angiogenesis. *Int J Mol Sci*. 2022;23(1). doi:10.3390/ijms23010337

121. Alsousou J, Ali A, Willett K, Harrison P. The role of platelet-rich plasma in tissue regeneration. *Platelets*. 2013;24(3):173-182. doi:10.3109/09537104.2012.684730

122. Van De Kamp J, Jahnen-Dechent W, Rath B, Knuechel R, Neuss S. Hepatocyte Growth Factor-Loaded Biomaterials for Mesenchymal Stem Cell Recruitment. *Stem Cells Int*. 2013;2013. doi:10.1155/2013/892065

123. Akhurst RJ. TGF-β antagonists: Why suppress a tumor suppressor? *J Clin Invest*. 2002;109(12):1533. doi:10.1172/JCI15970

124. Li H, Mittal A, Makonchuk DY, Bhatnagar S, Kumar A. Matrix metalloproteinase-9 inhibition ameliorates pathogenesis and improves skeletal muscle regeneration in muscular dystrophy. *Hum Mol Genet*. 2009;18(14):2584. doi:10.1093/HMG/DDP191

125. Lin CC, Boyer PD, Aimetti AA, Anseth KS. Regulating MCP-1 Diffusion in Affinity Hydrogels for Enhancing Immuno-isolation. *J Control Release*. 2010;142(3):384. doi:10.1016/J.JCONREL.2009.11.022

126. Marino S, Hult C, Wolberg P, Linderman JJ, Kirschner DE. The Role of Dimensionality in Understanding Granuloma Formation. *Comput (Basel, Switzerland)*. 2018;6(4). doi:10.3390/COMPUTATION6040058

127. Sego TJ, Kasacheuski U, Hauersperger D, Tovar A, Moldovan NI. A heuristic computational model of basic cellular processes and oxygenation during spheroid-dependent biofabrication. *Biofabrication*. 2017;9(2):024104. doi:10.1088/1758-5090/AA6ED4

128. Haizlip KM, Harrison BC, Leinwand LA. Sex-based differences in skeletal muscle kinetics and fiber-type composition. *Physiology*. 2015;30(1):30-39. doi:10.1152/physiol.00024.2014

129. Knewtson KE, Ohl NR, Robinson JL. Estrogen Signaling Dictates Musculoskeletal Stem Cell Behavior: Sex Differences in Tissue Repair. *Tissue Eng - Part B Rev*. 2022;28(4):789-812. doi:10.1089/ten.teb.2021.0094

130. Liu S, Kostek M, Omstead K. Sex based differences in muscle regeneration. *Physiology*. 2023;38(S1). doi:10.1152/physiol.2023.38.s1.5729941

131. Enns DL, Tiidus PM. The influence of estrogen on skeletal muscle: Sex matters. *Sport Med*. 2010;40(1):41-58. doi:10.2165/11319760-000000000-00000

132. Zhao W, Zhao H, Li M, Huang C. Microfluidic devices for neutrophil chemotaxis studies. *J Transl Med*. 2020;18(1):1-19. doi:10.1186/s12967-020-02335-7

133. Heit B, Liu L, Colarusso P, Puri KD, Kubes P. PI3K accelerates, but is not required for,
neutrophil chemotaxis to fMLP. *J Cell Sci*. 2008;121(2):205-214. doi:10.1242/JCS.020412

134. Martin KS, Kegelman CD, Virgilio KM, et al. In Silico and In Vivo Experiments Reveal M-
CSF Injections Accelerate Regeneration Following Muscle Laceration. *Ann Biomed Eng*.
45. doi:10.1007/s10439-016-1707-2

135. Bos E van den, Walbaum S, Horsthemke M, Bachg AC, Hanley PJ. Time-lapse Imaging
of Mouse Macrophage Chemotaxis. *JoVE (Journal Vis Exp*. 2020;2020(158):e60750.
doi:10.3791/60750

136. Corliss BA, Azimi MS, Munson JM, Peirce SM, Murfee WL. Macrophages: An
Inflammatory Link between Angiogenesis and Lymphangiogenesis. *Microcirculation*.
2016;23(2):95. doi:10.1111/MICC.12259

137. Newby AC. Metalloproteinase expression in monocytes and macrophages and its
relationship to atherosclerotic plaque instability. *Arterioscler Thromb Vasc Biol*.
2008;28(12):2108-2114. doi:10.1161/ATVBAHA.108.173898

138. Lu HL, Huang XY, Luo YF, Tan WP, Chen PF, Guo YB. Activation of M1 macrophages
plays a critical role in the initiation of acute lung injury. *Biosci Rep*. 2018;38(2):20171555.
doi:10.1042/BSR20171555

139. Cui K, Ardell CL, Podolnikova NP, Yakubenko VP. Distinct migratory properties of M1,
M2, and resident macrophages are regulated by $\alpha d\beta2$ and $\alpha m\beta2$ integrin-mediated
adhesion. *Front Immunol*. 2018;9(NOV):2650. doi:10.3389/fimmu.2018.02650

140. da Silva MD, Bobinski F, Sato KL, Kolker SJ, Sluka KA, Santos ARS. IL-10 Cytokine
Released from M2 Macrophages Is Crucial for Analgesic and Anti-inflammatory Effects of
Acupuncture in a Model of Inflammatory Muscle Pain. *Mol Neurobiol*. 2015;51(1):19.
doi:10.1007/S12035-014-8790-X

141. Moncayo AR. Poisson Convergence and Family Trees. *Ann Probab*. 2007;3(6).
doi:10.1214/aop/1176996235

142. Xiaoping C, Yong L. Role of matrix metalloproteinases in skeletal muscle: migration,
differentiation, regeneration and fibrosis. *Cell Adh Migr*. 2009;3(4).
doi:10.4161/CAM.3.4.9338

143. Terrente Y, El Fahime E, Caron NJ, et al. Tumor necrosis factor-alpha (TNF-alpha)
stimulates chemotactic response in mouse myogenic cells. *Cell Transplant*.
2003;12(1):91-100. doi:10.3727/000000003783985115

144. Bakkar N, Wang J, Ladner KJ, et al. IKK/NF-κB regulates skeletal myogenesis via a
signaling switch to inhibit differentiation and promote mitochondrial biogenesis. *J Cell
Biol*. 2008;180(4):787. doi:10.1083/JCB.200707179

145. Saclier M, Yacoub-Youssef H, Mackey AL, et al. Differentially activated macrophages
orchestrate myogenic precursor cell fate during human skeletal muscle regeneration.
Stem Cells. 2013;31(2):384-396. doi:10.1002/STEM.1288

146. Perandini LA, Chimin P, Lutkemeyer D da S, Câmara NOS. Chronic inflammation in
skeletal muscle impairs satellite cells function during regeneration: can physical exercise
restore the satellite cell niche? *FEBS J*. 2018;285(11):1973-1984.
doi:10.1111/FEBS.14417

147.   Gal-Levi R, Leshem Y, Aoki S, Nakamura T, Halevy O. Hepatocyte growth factor plays a
1287   dual role in regulating skeletal muscle satellite cell proliferation and differentiation.
1288   *Biochim Biophys Acta - Mol Cell Res*. 1998;1402(1):39-51. doi:10.1016/S0167-
1289   4889(97)00124-9

1290   148.   Ten Broek RW, Grefte S, Von Den Hoff JW. Regulatory factors and cell populations
1291   involved in skeletal muscle regeneration. *J Cell Physiol*. 2010;224(1):7-16.
1292   doi:10.1002/JCP.22127

1293   149.   Kuang S, Kuroda K, Le Grand F, Rudnicki MA. Asymmetric self-renewal and commitment
1294   of satellite stem cells in muscle. *Cell*. 2007;129(5):999-1010.
1295   doi:10.1016/J.CELL.2007.03.044

1296   150.   Yennek S, Burute M, Théry M, Tajbakhsh S. Cell adhesion geometry regulates non-
1297   random DNA segregation and asymmetric cell fates in mouse skeletal muscle stem cells.
1298   *Cell Rep*. 2014;7(4):961-970. doi:10.1016/J.CELREP.2014.04.016

1299   151.   Siegel AL, Kuhlmann PK, Cornelison DDW. Muscle satellite cell proliferation and
1300   association: new insights from myofiber time-lapse imaging. *Skelet Muscle*. 2011;1(1).
1301   doi:10.1186/2044-5040-1-7

1302   152.   Rocheteau P, Gayraud-Morel B, Siegl-Cachedenier I, Blasco MA, Tajbakhsh S. A
1303   subpopulation of adult skeletal muscle stem cells retains all template DNA strands after
1304   cell division. *Cell*. 2012;148(1-2):112-125. doi:10.1016/J.CELL.2011.11.049

1305   153.   Otto A, Collins-Hooper H, Patel A, Dash PR, Patel K. Adult skeletal muscle stem cell
1306   migration is mediated by a blebbing/amoeboid mechanism. *Rejuvenation Res*.
1307   2011;14(3):249-260. doi:10.1089/rej.2010.1151

1308   154.   Gibb AA, Lazaropoulos MP, Elrod JW. Myofibroblasts and Fibrosis. *Circ Res*.
1309   2020;127(3):427-447. doi:10.1161/CIRCRESAHA.120.316958

1310   155.   Alberts B, Johnson A, Lewis J, Raff M, Roberts K, Walter P. Extracellular Control of Cell
1311   Division, Cell Growth, and Apoptosis. Published online 2002. Accessed March 14, 2023.
1312   https://www.ncbi.nlm.nih.gov/books/NBK26877/

1313   156.   Cornwell KG, Downing BR, Pins GD. Characterizing fibroblast migration on discrete
1314   collagen threads for applications in tissue regeneration. *J Biomed Mater Res Part A*.
1315   2004;71A(1):55-62. doi:10.1002/JBM.A.30132

1316   157.   Qutub A, Gabhann F, Karagiannis E, Vempati P, Popel A. Multiscale models of
1317   angiogenesis. In: *IEEE Engineering in Medicine and Biology Magazine*. Vol 28. NIH
1318   Public Access; 2009:14-31. doi:10.1109/MEMB.2009.931791

1319   158.   Tigner A, Ibrahim SA, Murray I. Histology, White Blood Cell. *StatPearls*. Published online
1320   November 19, 2021. Accessed May 16, 2022.
1321   https://www.ncbi.nlm.nih.gov/books/NBK563148/

1322   159.   Garcia SM, Tamaki S, Lee S, et al. Stem Cell Reports Resource High-Yield Purification,
1323   Preservation, and Serial Transplantation of Human Satellite Cells. Published online 2018.
1324   doi:10.1016/j.stemcr.2018.01.022

1325   160.   Krombach F, Münzing S, Allmeling AM, Gerlach JT, Behr J, Dörger M. Cell size of
1326   alveolar macrophages: an interspecies comparison. *Environ Health Perspect*.
1327   1997;105(Suppl 5):1261. doi:10.1289/EHP.97105S51261

1328   161.   Downey GP, Doherty DE, Schwab B, Elson EL, Henson PM, Worthen GS. Retention of
1329   leukocytes in capillaries: role of cell size and deformability.
1330   *https://doi.org/101152/jappl19906951767*. 1990;69(5):1767-1778.
1331   doi:10.1152/JAPPL.1990.69.5.1767

1332   162.   Freitas RJ. *Nanomedicine: Basic Capabilities* . Vol 1. Landes Bioscience; 1999.
1333   Accessed May 16, 2022. http://www.nanomedicine.com/NMI/8.5.1.htm

1334   163.   Deng B, Wehling-Henricks M, Villalta SA, Wang Y, Tidball JG. IL-10 Triggers Changes in
1335   Macrophage Phenotype That Promote Muscle Growth and Regeneration. *J Immunol*.
1336   2012;189(7):3669-3680. doi:10.4049/jimmunol.1103180

1337   164.   Raimondo TM, Mooney DJ. Functional muscle recovery with nanoparticle-directed M2
1338   macrophage polarization in mice. *Proc Natl Acad Sci U S A*. 2018;115(42):10648-10653.
1339   doi:10.1073/pnas.1806908115

1340   165.   Chen SE, Gerken E, Zhang Y, et al. Role of TNF-α signaling in regeneration of
1341   cardiotoxin-injured muscle. *Am J Physiol Cell Physiol*. 2005;289(5):C1179.
1342   doi:10.1152/AJPCELL.00062.2005

1343