Neural geometrodynamics, complexity, and plasticity: a psychedelics perspective

G. Ruffini[1]*, E. Lopez-Sola[1,2], J. Vohryzek[2,3] and R. Sanchez-Todo[1,2]

[1] Brain Modeling Department, Neuroelectrics, Barcelona

[2] Computational Neuroscience Group, UPF, Barcelona

[3] Centre for Eudaimonia and Human Flourishing, Linacre College, University of Oxford

Dec 1st, 2023

Abstract

We explore the intersection of neural dynamics and the effects of psychedelics in light of distinct timescales in a framework integrating concepts from dynamics, complexity, and plasticity. We call this framework *neural geometrodynamics* for its parallels with general relativity's description of the interplay of spacetime and matter. Trajectories within the dynamical landscape of "fast time" dynamics are shaped by the structure of a differential equation and its connectivity parameters, which themselves evolve over "slow time" driven by state-dependent and state-independent plasticity mechanisms. Finally, the adjustment of plasticity processes (metaplasticity) takes place in an "ultraslow" time scale. Psychedelics flatten the neural landscape, leading to heightened entropy and complexity of neural dynamics, as observed in neuroimaging and modeling studies linking increases in complexity with a disruption of functional integration. We highlight the relationship between the complexity of fast neural dynamics and synaptic plasticity. Pathological, rigid, or "canalized" neural dynamics result in an ultrastable confined repertoire, allowing slower plastic changes to consolidate them further. However, under the influence of psychedelics, the destabilizing emergence of complex dynamics leads to a more fluid and adaptable neural state in a process that is amplified by the plasticity-enhancing effects of psychedelics. This shift manifests as an acute systemic increase of disorder and a possibly longer-lasting increase in complexity affecting both short-term dynamics and long-term plastic processes. Our framework offers a holistic perspective of the acute effects of these substances and their potential long-term impacts on neural structure and function.

---

*Corresponding author: giulio.ruffini@neuroelectrics.com

Contents

1   Introduction

> Spacetime tells matter how to move;
> matter tells spacetime how to curve.
>
> ———————————————
> *John Archibald Wheeler*, in
> *Gravitation* (1973)

In this paper, we explore new perspectives to interpret changes in the brain's landscape and connectivity, focusing on the subtle interplay between structural and dynamical aspects across timescales (fast, slow, and ultraslow). Our primary goal is to present a framework that enhances the understanding of the intricate relationships among brain dynamics, complexity, structure, and plasticity. This framework, which we call "neural geometrodynamics", draws on principles from non-linear dynamics and is further inspired by conceptual links to general relativity in physics.

In describing neural dynamics, we will employ the mathematical formalism of neural mass models (NMMs), although other computational neuroscience formulations are equally relevant [1, 2]. Neural mass models have been extensively utilized to model various brain activities, from localized brain functions to the coordinated activity observed in different brain regions. By employing mathematical formulations that include essential features like synaptic connectivity and neuronal excitability, NMMs enable the simulation and analysis of complex brain activities in various dynamic regimes [3]. NMMs are particularly useful because they provide a link between the mesoscopic physiological scale and macroscopic brain function, allowing for the connection of effects on neurons at the molecular level, such as those of psychedelics, with those of whole-brain connectivity [4, 5].

Psychedelics serve as an illustrative case of the framework, given the immediate and potentially lasting plastic changes these substances can provoke in the brain [6]. By altering neural dynamics and connectivity, psychedelics are thought to induce both transient and sustained shifts in cognition and perception [7]. Several studies underscore the role of psychedelics in inducing neuroplasticity with antidepressant effects, revealing mechanisms at molecular, synaptic, and dendritic levels [8, 9], and with significant potential for treating neuropsychiatric disorders [10, 11], although the duration and permanence of these effects remain to be fully understood.

Recent conceptual perspectives have enhanced our understanding of the brain's response to psychedelics, combining biological, dynamical systems, complexity science, and artificial intelligence viewpoints. The REBUS (RElaxed Beliefs Under pSychedelics)

framework [12], grounded in the Free Energy Principle and the entropic and anarchic brain models, offers a perspective on the effects of psychedelics on the brain whereby psychedelic action results in the collapse of brain functional hierarchies or, in other words, in the "flattening of the landscape" of brain's dynamics to allow the brain state to escape a deep local minimum. The term *annealing* is also used in this context in relation to physical annealing in metallurgy and simulated annealing in numerical optimization [13].

These changes are reflected in the complexity of neural dynamics, which can be evaluated using various techniques, including Ising modeling, showing that the effects of LSD increase the complexity of brain dynamics (e.g., increased Ising temperature, Lempel-Ziv, and the Block Decomposition Method complexity) by loosening interhemispheric connectivity—especially homotopic links [14], corroborating earlier modeling studies suggesting the important of long-range connections in phase transitions [15].

It has been argued that the observed expansion of the repertoire of functional patterns elicited by hallucinogenic substances can be understood as being driven by an enhancement of brain entropy [16]. In turn, this may take place while the brain is moving to a more disordered state, with this being related to the relaxation of high-level cognitive priors [12, 17]. This may in turn lead to a favorable context for conducting psychotherapy [12, 18]. Studies on functional neuroimaging regarding psilocybin and LSD effects have shown initial evidence of the mechanistic alterations on brain dynamics at the network level, with the majority of the findings suggesting a relative weakening of usual functional configurations giving place to an increase of brain entropy, global functional integration, and more flexible brain dynamics [16, 19–27].

In [14], we showed that psychedelics push brain dynamics towards disorder and away from criticality in the context of Ising modeling. This aligns with the REBUS framework and the Free-Energy Principle (FEP), which links the vividness of experience to the entropy of brain activity. The primary action of psychedelics starts with the stimulation of the 5-HT$_{2A}$ serotonin receptor, leading to increased neuronal excitability and altered cortical activity. This disruption affects high-level neural functions and precision-weighting of internal models, allowing lower-level models to emerge more prominently in experience. This mechanism leads to increased Shannon complexity of brain activity, as supported by measures like LZ complexity. Extending this, we incorporated algorithmic information theory-oriented methods like BDM, finding increased signal complexity and system temperature under psychedelics. This supports the idea that a wakeful brain, in tracking structured world data, exhibits dimensionality reduction and criticality. Psychedelics,

however, elevate these dynamics into higher dimensions, as evidenced by the shift from a critical to a supercritical state in the LSD-induced state. These results are also consistent with the algorithmic information theory of consciousness (KT) [17, 28], where the psychedelic shift towards disorder is associated with a breakdown of world-modeling / world-tracking system in the brain.

A related perspective for this paper is the CANAL framework [11] for describing the pathological plasticity of "being stuck in a rut" in certain mood disorders and the potential therapeutic role of psychedelics through the concept of metaplasticity. In contrast to psychedelics, these changes are reflected in neural dynamics with brain signatures of excessively rigid and highly ordered functional states [29]. The CANAL framework can be further extended by establishing connections with Deep Neural Networks [30] to introduce a distinction between two distinct pathological phenomena — one related to fast brain dynamics and their slow and ultraslow counterparts. These distinctions will be naturally integrated into the presented framework.

While the discussion is centered on the effects of psychedelics, the proposed framework readily extends to other phenomena related to plasticity, including pathological plasticity in mood disorders [31] and interventions like transcranial brain stimulation (tES) [32].

In what follows, we formalize the notions of brain dynamics, plasticity, and their associated timescales and subsequently use them to study the impact of psychedelics on the brain. In the last section, we draw connections between the proposed framework and concepts from general relativity in physics. We hope these parallels will illuminate the complex relationship between the structure and function of brain dynamics.

2   Dynamics across timescales

The state of a system can be defined by a set of coordinates in phase space: a multidimensional manifold in which each dimension corresponds to one of the variables. For a single particle moving in one dimension, the phase space is two-dimensional, with one axis representing its position and the other representing its momentum. For example, Figure 1 illustrates the phase space of a pendulum with friction. In phase space, and perhaps after some transient period, the possible trajectories of the states of the system lie in a reduced or invariant manifold, which we may refer to as the "geometry" or latent "structure" of the phase space. In convergent phase space dynamics, these are called stable manifolds, including attractors or repellors. Together, the structure of

the phase space with its invariant properties can be referred to as the dynamical landscape, where the depth or shallowness of the "valleys" can, in some cases, be interpreted as the stability of the state in that location given some stochastic forcing. For example, in mechanics, the landscape can be labeled by potential energy isolines, e.g., in a physical system such as in the pendulum example in Figure 1 (bottom right), or their generalization, Lyapunov functions [37].

Fast time: neural dynamics

Here, we discuss the first equation in neural geometrodynamics in the context of neural mass models, but the ideas are applicable more extensively in computational neuroscience. The standard equation we use in neural mass modeling is a multidimensional ODE of the form

$$\dot{x} = f\big(x; w, \eta(t)\big) \tag{1}$$

with $x \in \mathbb{R}^n$ and where w denotes connectivity parameters[1] and where, as usual, a dot over a variable denotes its time derivative. This equation governs dynamics at short time scales (seconds or less) when connectivity parameters w are assumed to be constant.

We may think of this equation describing phenomena in **fast time** scales as providing the "structure" for the dynamics of neuronal population state. The fast timescale is set by synaptic transmission (milliseconds) and by ephaptic coupling (electromagnetic waves) [39–42] in a nanosecond or subnanosecond scale [41].

Equation 1 characterizes the **dynamical landscape**, which is established through the geometric structure of the phase space, where trajectories are shaped by the given set

---

[1]In the REBUS model [38], from the Free Energy perspective, w would correspond to the weights or precision assigned to priors/beliefs; from the Entropic Brain perspective, w would correspond to the weights of the effective connectivity between neuronal populations on the macroscopic scale.

Figure 1: Dynamics of a pendulum with friction: time series, phase space, and energy landscape. Attractors in phase space are valleys in the "energy" landscape (or more generally, defined by the level sets of a Lyapunov function).

of ordinary differential equations. Such landscape is analogous to the Neural Activation Landscape proposed in [30]. More specifically, we talk about the landscape as defined by the manifold generated by the motion of trajectories with coordinates $x \in \mathbb{R}^n$. Typically, trajectories lie in a reduced manifold of dimensionality lower than $\mathbb{R}^n$. The fact that such a reduced space exists means that it can be generated by coordinates in a reduced latent space.

As we can see, the landscape is determined by the functional form of $f(x; w, \eta(t))$ and by the parameters w. The external input term $\eta(t)$ makes the equations non-autonomous (an autonomous ODE does not explicitly depend on time). This term can refer to external forces providing random kicks to the trajectory or to a more steady and purposeful forcing from unspecified internal systems or external inputs.

Slow time: connectodynamics

The landscape, like that on planet Earth, may appear to be static, but in reality, it is not fixed. It also flows in **slow time**. We thus consider changes in connectivity in the system, that is, now $w = w(t)$. We call the potential for such changes **plasticity of the system**. The general form of this equation is $\dot{w} = g(x, w; \theta)$, with θ standing for a set

of parameters controlling plasticity.

To be more concrete, we can think of two types of process: one that modifies the connectivity parameters independently of the system's state (ψ) and another that is a function of the state (e.g., Hebbian plasticity [43], h). We express this by writing

$$\dot{w} = \psi(w; \gamma) + h(x, w; \alpha) \tag{2}$$

This decomposition separates out *state-dependent* (via the term $h(x, w; \alpha)$) and *state-independent plasticity* (with $\psi(w; \gamma)$) processes. The set of parameters θ is similarly decomposed as $\theta = (\alpha, \gamma)$: we separate out the plasticity-controlling parameters in order to differentiate the *state-dependent (α)* and *state-independent (γ)* plasticity control parameters (e.g., Hebbian vs. drug-enhanced structural plasticity [11]).

State dependence implies that system temperature, phase transitions, and critical phenomena are relevant for the study of the dynamics of plasticity. In particular, within the scope of slower "slow time" (taking place over many hours), we include *homeostatic plasticity* [44–47], which may itself target desired complexity states as a homeostatic goal [48, 49].

In summary, the functions h and ψ with parameters α and γ regulate *connectodynamics*, defining to where and how fast the effective connectivity will change in a state-dependent or state-independent way.

These connectodynamics differential equations define a new dynamical landscape, which we can call the *plasticity landscape* (analogous to the Synaptic Weight Landscape in [30]). The state w in this plasticity landscape will determine the shape of the neural dynamics landscape.

Ultraslow time: metaplasticity

Next, we allow the plasticity parameters to be dynamic, i.e.,

$$\dot{\theta} = \xi(x, w, \theta) \tag{3}$$

This equation is again state-dependent, allowing the system to respond to changes in the neural dynamics, including critical phenomena and complexity. Plasticity dynamics reflect changes in plasticity, such as the changes regulating state-dependent (Hebbian) plasticity (changes in α) during neurodevelopment, and state-independent plasticity, such as the ones induced by psychedelics in the acute or post-acute phases (changes in ψ). In the latter case of plasticity, there are numerous candidates for these plastic processes,

such as heterosynaptic plasticity [50], critical-period plasticity [51] and metaplasticity [52]. They all reflect a common physiological principle well described by Abraham et al. in the definition of metaplasticity:

> **Metaplasticity** *[...] is manifested as* **a change in the ability to induce subsequent synaptic plasticity**, *such as long-term potentiation or depression. Thus, metaplasticity is a higher-order form of synaptic plasticity [52].*

Thus, metaplasticity and its counterparts are terms used in neuroscience to refer to the plasticity of synaptic plasticity. That is, the idea that the ability of synapses to strengthen or weaken in response to increases or decreases in their activity (which is called synaptic plasticity) can be modulated based on the history of the synaptic activity.

There are various mechanisms through which metaplasticity can occur, such as changes in the amount of neurotransmitters that are released, alterations in the types or numbers of receptors on the post-synaptic neuron, changes in the internal signaling pathways within neurons, and more. Metaplasticity has important implications for the learning and memory of an organism, as it can regulate the ability of synaptic connections to change and adapt over time. In extreme scenarios, such regulation can lead to two opposite outcomes in which metaplasticity can either become overly exuberant, reflecting the notion of catastrophic forgetting in artificial neural networks, or impoverished, reflecting general plasticity loss [30].

Metaplasticity may also play a role in certain neurological and psychiatric conditions. For example, reduced metaplasticity could underlie conditions such as major depressive disorder, obsessive-compulsive disorder, anxiety, or substance abuse on one hand, and overly malleable metaplasticity could define bipolar and personality disorders on the other [30, 53].

We call the set of equations 1,2 and 3 — somewhat whimsically — the equations for *neural geometrodynamics* in reference to the equations of general relativity in physics. We recall that general relativity provides equations defining the dynamics of spacetime geometry (via the "metric") coupled with matter (see Section 4) [54].

3 Dynamics under psychedelics

Psychedelics like psilocybin and LSD act as agonists or partial agonists for serotonin 5-hydroxytryptamine 2A (5-HT$_{2A}$) receptors, specifically targeting Layer V cortical pyra-

Figure 2: Geometrodynamics of the acute and post-acute plastic effects of psychedelics. The **acute** plastic effects can be represented by rapid state-independent changes in connectivity parameters, i.e., the term $\psi(w;\gamma)$ in Eq. 2. This will result in the flattening or de-weighting of the dynamical landscape. Such flattening allows for the exploration of a wider range of states, eventually creating new minima through state-dependent plasticity, represented by the term $h(x,w;\alpha)$ in Eq. 2. As the psychedelic action fades out, the landscape gradually transitions towards its initial state but with lasting changes due to the creation of new attractors during the acute state. The **post-acute** plastic effects can be described as a "window of plasticity" characterized by a period of enhanced plasticity. These transitions are brought about by time-dependent changes of the parameters γ and α, each controlling the behavior of state-independent and state-dependent plasticity, respectively. In this post-acute phase, the landscape is more malleable to internal and external influences.

midal neurons [11, 16, 38, 55, 56]. Recognized for their potent and immediate impact on the brain, these drugs cause a swift reconfiguration of neural dynamics. As we explain, this immediate effect is represented in our model by alterations in the state-independent connectivity parameters (w).

5-HT$_{2A}$ receptors are strongly expressed in the pyramidal neurons of the cortex along a steep anteroposterior gradient [57]. When psychedelics bind to them, they can lead to a gradual increase of the excitability of these pyramidal neurons — depolarizing them and making them more susceptible to excitatory inputs such as those associated with glutamate receptors [57] — much as the gain knob in an amplifier. This increased excitability and susceptibility to inputs can lead to changes in the firing patterns of these neurons and alterations in the overall neural circuit activity.

How are these effects represented in Equations 1 and 2? If we include the neuromodula-

tory nodes in our model — the dorsal raphe and median raphe nuclei in the brainstem are the source of most serotonergic neurons projecting throughout the brain [57] —, the modulation of serotonin receptors can be represented by changes in neuromodulatory connectivity (the subset of w parameters in the model connecting the raphe nuclei to other nodes). Alternatively, if neuromodulatory nodes are not explicitly included in the model, for the purposes at hand, we can think of the changes in the excitability of the nodes affected by neuromodulatory inputs as leading to changes in their effective connectivity to other nodes.

This abrupt shift induced by psychedelics can be thought of as a transformation of the phase space's geometry, allowing the neural state to explore new trajectories. This process manifests in an increase of complexity and disorder, which can be measured using various tools in different modalities (e.g., EEG or fMRI BOLD with measures such as entropy, fractal dimension, algorithmic complexity, etc. [14, 58–60]). Psychedelic-induced changes in connectivity correspond to a flattening of the dynamical landscape [12] or a destabilization of it [30]. Effective connectivity, as inferred using Ising modeling, is seen to decrease in the acute phase of LSD (especially in interhemispheric homotopic links), with a subsequent increase in algorithmic complexity of BOLD signals measured using fMRI imaging [14]. The alteration of effective connective results in an immediate and state-independent remodeling of the dynamical landscape during the acute phase of psychedelics, which is represented by the term $\psi(w; \gamma)$ in the connectodynamics equation (Eq. 2)[2].

The instantaneous modification of the landscape is, however, ephemeral, gradually fading as the acute effects of the psychedelics wear off. The system returns to near its original geometrical configuration but with lasting influences brought about by the plastic changes resulting from the exploration of new trajectories in the acute phase. These residual changes are captured by the state-dependent plasticity term, $h(x, w; \alpha)$, which reflects changes in connectivity due to Hebbian plasticity that arise from the co-activation of neurons during the psychedelic acute stage.

In the literature, there is an increasing body of evidence suggesting a post-acute phase following psychedelic exposure characterized by a period of enhanced plasticity [11, 12, 61–63]. This phase can be interpreted as an extended window of malleability of the landscape, which could have profound implications for learning and therapy. Such window of

---

[2]In the REBUS model and the Entropic Brain perspective [38], the weights of the effective connectivity during the psychedelic-induced state are "flattened" or "de-weighted", representing a more symmetrical and non-hierarchical connectivity profile.

plasticity has been related to increased neurogenesis and upregulation of Brain-Derived Neurotropic Factor (BDNF) in humans and mice [8]. The activity-dependent release of BDNF plays a crucial role in selectively strengthening active synapses while weakening inactive ones, a critical process for Hebbian-type plasticity. Intriguingly, recent studies with mice have found psychedelic-induced changes in plasticity and antidepressant-like behavior dependent on the increase of endogenous BDNF and TrkB binding (the receptor of BDNF), but independent from the activation of 5-HT$_{2A}$ [9, 10].

In terms of our model, these two pathways correspond to changes of connectivity through Equation 2 due to a temporary modulation of the parameters γ and α (i.e., metaplasticity, see Equation 3) upregulating state-independent and state-dependent plasticity processes, respectively. The strong acute-phase increase of state-independent plasticity (ψ) would be directly associated with the activation of serotonergic receptors, as discussed above, with a possible gradual decrease during the post-acute phase (solid white line in Figure 2). The sustained increase of state-dependent plasticity (h) in the post-acute phase (dashed black line in Figure 2) would be linked to dendritic growth, neurogenesis, upregulation of BDNF, and other related changes. This means that in the post-acute period, the landscape would be more responsive to state changes (itself influenced by external factors), offering a potential mechanism for the long-lasting changes reported after psychedelic experiences. Such external influences are modeled by the external input term $\eta(t)$ in the state equation (Eq. 1) and can represent environmental/sensory inputs, psychotherapy, or neuromodulatory brain stimulation techniques such as transcranial electrical current stimulation (tES).

Dynamics of psychedelics and psychopathology

Recently, psychedelic medicine has emerged as a promising direction for treating mental disorders such as depression or addiction [64]. The nuanced interaction between the brain's neurophysiology and the emergent brain activity underlies the pathophysiology of mood disorders, often resulting in a persistent and maladaptive rigidity in cognitive and emotional processes [65]. Such changes to the brain's neurophysiology can be explained through the CANAL framework whereby pathological plasticity, often caused by a traumatic event, asserts itself and dominates brain activity, driving the brain state to be "stuck in a rut" [11], i.e., a deepening minimum in the dynamical landscape (see Figure 3).

The interplay between external inputs, neural (fast time), and connectivity (slow time) dynamics can drive the system into a joint canalized, stable state of lower complexity.

Under the influence of psychedelics, more diverse and complex dynamics destabilize the plasticity equilibrium point, leading to a more fluid and adaptable neural state in a process that is amplified by the plasticity-enhancing effects of psychedelics. This shift manifests as an acute systemic increase of disorder and possibly a longer-lasting increase in complexity (Ising temperature, LZ complexity, etc.) that affects both short-term dynamics and long-term plastic processes.

The CANAL framework offers insight into the neural mechanisms underlying the persistence of various brain disorders. In particular, psychedelics may mediate their effects by altering the balance between stability and plasticity in neural networks through metaplasticity and thus act as potential therapeutic treatments. By acting on the serotonergic receptors, they trigger a cascade of neurochemical events, subsequently facilitating the reorganization of entrenched neural patterns. As discussed above, this alteration of the neural network during the acute phase (connectodynamics) can be interpreted as a rapid deformation or flattening of the landscape that allows the trapped state to escape and access more adaptive cognitive and emotional patterns. The rapid increase in complexity (a change in the dynamics) is in itself a likely driver of metaplasticity. The acute phase is believed to be followed by an extended window of malleability of the landscape, otherwise known as a "window of plasticity", where treatments such as psychotherapy and transcranial electrical stimulation can further alter the pathological rigidity characteristic of various brain disorders (see Figure 3).

4 Neural geometrodynamics and general relativity

A parallel can be drawn between neural geometrodynamics and Einstein's equations of general relativity — the original geometrodynamics. Both frameworks involve the dynamical interaction between structure and resulting activity, each influencing and being influenced by the other. The Einstein field equations, including the cosmological constant Λ, are

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu}. \tag{4}$$

Here, $g_{\mu\nu}$ is the metric tensor, $R_{\mu\nu} = R_{\mu\nu}[g_{\mu\nu}]$ is the Ricci curvature tensor and a function of $g_{\mu\nu}$, $R[g_{\mu\nu}]$ is the Ricci scalar (or curvature scalar) and a function of $g_{\mu\nu}$, $T_{\mu\nu}$ is the stress-energy tensor[3], a function of the mass and energy distribution (all the indices refer

---

[3]The stress-energy tensor (also called the energy-momentum tensor) is a central concept in general relativity. It encapsulates the distribution and flow of energy and momentum in spacetime, and its components include energy density, momentum density, and stress (pressure and shear stress) within a given region.

Figure 3: Psychedelics and psychopathology: a dynamical systems perspective. From left to right, we provide three views of the transition from health to canalization following a traumatic event and back to a healthy state following the acute effects and post-acute effects of psychedelics and psychotherapy. The top row provides the *neural network (NN) and effective connectivity (EC) view.* Circles represent nodes in the network and edge connectivity between them, with edge thickness representing the connectivity strength between nodes. The middle row provides *the landscape view*, with three schematic minima and colors depicting the valence of each corresponding state (positive, neutral, or negative). The bottom row represents the *transition probabilities across states* and how they change across the different phases. Due to traumatic events, excessive canalization may result in a pathological landscape reflected as a deepening of a negative valence minimum where the state may be trapped. During the acute psychedelic state, the landscape is deformed, enabling the state to escape. Moreover, plasticity is enhanced during the acute and post-acute phases, benefiting interventions such as psychotherapy or brain stimulation (i.e., changes in effective connectivity). Not shown is the possibility that a deeper transformation of the landscape may take place during the acute phase (see the discussion on the wormhole analogy in Section 4).

337 to spacetime dimensions), G is the gravitational constant, c is the speed of light, and Λ
338 is the cosmological constant. These equations, which can be seen as an equation of the
339 spacetime metric $g_{\mu\nu}$ if the stress-energy tensor $T_{\mu\nu}$ is given, describe the fundamental
340 interaction of gravitation as a result of spacetime being curved by matter and energy.
341 Specifically, they equate local spacetime curvature (on the left-hand side) with the local

energy and momentum within that spacetime (on the right-hand side).

To complete these equations, the *geodesic equation* portrays how particles (matter) move in this curved spacetime, encapsulated by the notion that particles follow the straightest possible paths (geodesics) in curved spacetime,

$$\frac{d^2 x^\mu}{d\tau^2} + \Gamma^\mu_{\rho\sigma} \frac{dx^\rho}{d\tau} \frac{dx^\sigma}{d\tau} = 0 \tag{5}$$

where x^μ are the coordinates of the particle, τ is the proper time along the particle's path, and $\Gamma^\mu_{\rho\sigma}[g_{\mu\nu}]$ are the Christoffel symbols, which are a function of $g_{\mu\nu}$ and encode the *connection* (a mathematical object that describes how vectors change as they are parallel transported along curves in spacetime). The stress-energy tensor $T_{\mu\nu}$ can be computed from the state of the particles, closing the system of equations. For example, for N particles, it is given by $T^{\mu\nu} = \sum_i m_i u_i^\mu u_i^\nu \delta(\mathbf{x} - \mathbf{x}_i)$, where m_i and u_i are the mass and velocity of the ith particle. More generally, the stress-energy tensor represents the state of matter and energy, which corresponds to x in our neural model. The metric $g_{\mu\nu}$, which specifies the geometry of spacetime, is akin to the connectivity w — which shapes the structure of the space where fast dynamics occur.

In the context of neural mass models, the state equation, $\dot{x} = f(x; w)$, is analogous to the geodesic equation – "the state of the system evolves according to the landscape geometry specified by the parameters w". On the other hand, the connectodynamics equation, $\dot{w} = h(x, w; \theta)$ (with θ standing for plasticity parameters), is analogous to Einstein's field equations — the parameters w, which describe the "structure" of the space where dynamics take place, evolve according to the current state of the system x and its 'readiness' for plasticity (parametrized by θ).

This comparison is largely metaphorical: the mutual influence between particles and spacetime in general relative is akin to the state of the neural system and its underlying connectivity parameters. In both cases, dynamics and structure are intertwined (see Figure 4).

Metaplasticity and variable constants in cosmology

In our neural mass model framework, the concept of metaplasticity is introduced as dynamic variations in the plasticity control constants, namely θ in the connectodynamics equation. This set of constants can be represented as evolving over time as a function of the state of the system or other relevant variables,

$$\dot{\theta} = \xi(x, w, \theta; \lambda) \tag{6}$$

Figure 4: General Relativity and Neural Geometrodynamics. Left: Equations for general relativity (the original geometrodynamics), coupling dynamics of matter with the dynamics of spacetime. Right: Equations for neural geometrodynamics, coupling neural state, and connectivity. Only fast and slow time equations are shown (ultraslow time endows with dynamics the "constants" appearing in these equations).

In this equation, ξ defines the evolution of the plasticity control constants with parameters λ.

Analogously, in the realm of general relativity and cosmology, it has been speculated that the fundamental constants, such as the speed of light c, the gravitational constant G, or the cosmological constant Λ, may in fact be dynamic. Although not part of the mainstream cosmological model, theories proposing variable constants, such as "Variable Speed of Light" (VSL) or "Variable Cosmological Constant" provide an intriguing parallel. For instance, within VSL theories, the speed of light c is postulated to vary over cosmological time scales. Certain hypothetical dynamical equations could dictate the dynamical evolution of these constants. Although these theories are quite speculative and do not form a part of mainstream physics, they offer an interesting perspective on the concept of metaplasticity and its potential implications for the dynamical evolution of neural mass models and the structure of their landscapes.

Psychedelics as wormholes in the neural landscape

In the parallel of general relativity and neural geometrodynamics, we see the effects of psychedelics as a deformation of the neural landscape (spacetime) that allows the brain state (of a particle or set of particles) to escape from a local minimum and transition to another location in the landscape (spacetime). Although transitions may be smooth and respect the topology of the landscape, deformations of the landscape may also be more extreme — sharp transitions through a *topological tipping point* of the dynamical landscape. This may be due to external inputs ($\eta(t)$), when our system is non-autonomous [66], e.g., from sensory or brain stimulation effect. And as we have discussed, it may be due to connectivity dynamics.

The creation of a wormhole in general relativity[4] can be viewed as a profound deformation of spacetime, bending and connecting distant parts of the universe in such a way that matter/energy, like an astronaut, can travel through vast distances in an instant. This change in the geometry and topology of spacetime can be likened to the effect of psychedelics on the human mind. Just as the wormhole alters the structure of spacetime, psychedelics may radically deform the dynamical landscape of neural dynamics, creating connections across distant landscape locations. In the same way that the astronaut uses the wormhole to bypass vast stretches of space, the deformation caused by psychedelics may allow the state of the brain to tunnel out and escape from a local minimum or stuck pattern of thought, providing access to new areas of the landscape — new perspectives and potentially unexplored territories of consciousness. Both phenomena are characterized by a fundamental transformation that enables traversal into otherwise inaccessible regions, whether in physical space or the brain's dynamical landscape (see Figure 5 for a sketch of this concept).

Characterizing the landscape

An important challenge in the program of neural geometrodynamics is to explore practical methods to characterize the landscape. Here again, we can draw inspiration from physics and mathematics.

The roots of this approach can be traced back to the 19th century when Carl Friedrich Gauss pioneered the field of differential geometry. Gauss's Theorema Egregium demon-

---

[4]Wormholes, a term due to John A. Wheeler [67], also known as Einstein-Rosen bridges, are solutions to the Einstein field equations of general relativity which some models suggest could exist under certain conditions. However, creating or stabilizing a traversable wormhole would likely require forms of exotic matter with properties not yet observed in the known universe.

Figure 5: Psychedelic wormhole. On the left, the landscape is characterized by a deep pathological attractor, where the neural state is trapped. After ingestion of psychedelics (middle), a radical transformation of the neural landscape takes place, with the formation of a wormhole connecting the pathological attractor to another, healthier attractor location and allowing the neural state to tunnel out. After the acute effects wear off (right panel), the landscape returns near its original topology and geometry, but activity-dependent plasticity reshapes it into a less pathological geometry.

strated that the curvature of a surface could be determined entirely by measurements within the surface, without any reference to the surrounding space [68]. This seminal insight has laid the groundwork for understanding manifolds in various contexts, including the theory of relativity. In the era of general relativity, the interplay between geometry and physics was further enriched. Differential geometry and algebraic topology — which comes into play when one is interested in the global properties of the manifold, such as its shape, connectedness, and the presence of holes [69, 70] — became essential in describing the fabric of spacetime itself. It enabled physicists to conceptualize how mass and energy warp the geometry of spacetime, thus influencing the motion of objects.

In our current endeavor, these ideas find application in characterizing the complex dynamical landscapes of neural data. Modern tools from deep learning, such as variational autoencoders, can be used to unravel the reduced spaces underlying neuroimaging or neurophysiological data [35, 36], while dynamical systems theory in concert with differential geometry, group theory, and algebraic topology data analysis [71] offer robust frameworks to understand and characterize them [66, 72–76]. The convergence of these mathematical techniques extends to neuroscience the tradition of linking geometry and topology to physical phenomena.

5 Conclusions

In this paper, we have used the umbrella of neural geometrodynamics to frame the coupling of state dynamics and their complexity, and the geometry and structure of reduced space, with plastic phenomena, and discussed it in the context of the acute and longer-lasting effects of psychedelics.

As a source of inspiration, we have established a parallel with other mathematical theories of nature, namely in general relativity, where dynamics and the "kinematic theatre" are intertwined (see the Appendix for a similar parallel with electrodynamics).

Although we can think of "geometry" in neural geometrodynamics as referring to the structure imposed by connectivity on state dynamics (paralleling the role of the metric in general relativity), it is more appropriate to think of it as the geometry of the reduced phase space (or invariant manifold) where state trajectories ultimately will lie: this is where the term reaches its fuller meaning. Since the fluid geometry and topology of the invariant manifolds underlying apparently complex neural dynamics may be strongly related to brain function and first-person (structured) experience [17], further research should focus on creating and characterizing these fascinating mathematical structures.

Acknowledgments

GR, ELS, and RST have received funding from the European Research Council (ERC Synergy Galvani) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 855109). GR, ELS, JV, and RST have received funding from European Union's Horizon 2020 research and innovation programme under grant agreement No 101017716 (Neurotwin).

References

[1] Gustavo Deco et al. "The dynamic brain: from spiking neurons to neural masses and cortical fields". In: *PLoS computational biology* 4.8 (2008), e1000092.

[2] Michael Breakspear. "Dynamic models of large-scale brain activity". In: *Nature neuroscience* 20.3 (2017), pp. 340–352.

[3] Joana Cabral, Morten L Kringelbach, and Gustavo Deco. "Functional connectivity dynamically evolves on multiple time-scales over a static structural connectome: Models and mechanisms". In: *NeuroImage* 160 (2017), pp. 84–96.

[4]   Gustavo Deco et al. "Whole-brain multimodal neuroimaging model using serotonin receptor maps explains non-linear functional effects of LSD". In: *Current biology* 28.19 (2018), pp. 3065–3074.

[5]   Morten L Kringelbach et al. "Dynamic coupling of whole-brain neuronal and neurotransmitter systems". In: *Proceedings of the National Academy of Sciences* 117.17 (2020), pp. 9566–9576.

[6]   Jacob S Aday et al. "Long-term effects of psychedelic drugs: A systematic review". In: *Neuroscience & Biobehavioral Reviews* 113 (2020), pp. 179–189.

[7]   David E Nichols. "Psychedelics". In: *Pharmacological reviews* 68.2 (2016), pp. 264–355.

[8]   Kimberley Van Elst et al. "Psychedelics and Neuroplasticity: A Systematic Review Unraveling the Biological Underpinnings of Psychedelics". In: *Frontiers in Psychiatry* 12 (2021), p. 725. DOI: 10.3389/fpsyt.2021.725061. URL: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8461007/.

[9]   Rafael Moliner et al. "Psychedelics promote plasticity by directly binding to BDNF receptor TrkB". en. In: *Nat. Neurosci.* 26.6 (June 2023), pp. 1032–1041.

[10]   Romain Nardou et al. "Psychedelics reopen the social reward learning critical period". en. In: *Nature* 618.7966 (June 2023), pp. 790–798.

[11]   R L Carhart-Harris et al. "Canalization and plasticity in psychopathology". en. In: *Neuropharmacology* 226.109398 (Mar. 2023), p. 109398.

[12]   R. L. Carhart-Harris and K. J. Friston. "REBUS and the Anarchic Brain: Toward a Unified Model of the Brain Action of Psychedelics". In: *Pharmacological Reviews* 71 (3 2019), pp. 316–344. DOI: 10.1124/pr.118.017160.

[13]   S Kirkpatrick, C D Gelatt Jr, and M P Vecchi. "Optimization by simulated annealing". en. In: *Science* 220.4598 (May 1983), pp. 671–680.

[14]   Giulio Ruffini et al. "LSD-induced increase of Ising temperature and algorithmic complexity of brain dynamics". en. In: *PLoS Comput. Biol.* 19.2 (Feb. 2023), e1010811.

[15]   G Ruffini and G Deco. "The 2D Ising model, criticality and AIT". Oct. 2021.

[16]   Robin L. Carhart-Harris. "The entropic brain - Revisited". In: *Neuropharmacology* (2018). DOI: 10.1016/j.neuropharm.2018.03.010..

[17]   Giulio Ruffini and Edmundo Lopez-Sola. "AIT foundations of structured experience". en. In: *J. AI. Consci.* 09.02 (Sept. 2022), pp. 153–191.

[18]   Mark A Geyer and Franz X Vollenweider. "Serotonin research: contributions to understanding psychoses". en. In: *Trends Pharmacol. Sci.* 29.9 (Sept. 2008), pp. 445–453.

[19] Robin L Carhart-Harris et al. "Neural correlates of the psychedelic state as determined by fMRI studies with psilocybin". en. In: *Proc. Natl. Acad. Sci. U. S. A.* 109.6 (Feb. 2012), pp. 2138–2143.

[20] Drummond E-Wen McCulloch et al. "Psychedelic resting-state neuroimaging: A review and perspective on balancing replication and novel analyses". en. In: *Neurosci. Biobehav. Rev.* 104689 (May 2022), p. 104689.

[21] Robin L Carhart-Harris et al. "The entropic brain: a theory of conscious states informed by neuroimaging research with psychedelic drugs". en. In: *Front. Hum. Neurosci.* 8 (Feb. 2014), p. 20.

[22] Robin L. Carhart-Harris et al. "Neural correlates of the LSD experience revealed by multimodal neuroimaging". In: *Proceedings of the National Academy of Sciences* 113.17 (2016), pp. 4853–4858. ISSN: 0027-8424. DOI: 10.1073/pnas.1518377113. eprint: https://www.pnas.org/content/113/17/4853.full.pdf. URL: https://www.pnas.org/content/113/17/4853.

[23] Selen Atasoy et al. "Common neural signatures of psychedelics: frequency-specific energy changes and repertoire expansion revealed using connectome-harmonic decomposition". In: *Progress in Brain Research* 242 (2018), pp. 97–120.

[24] Andrea I Luppi et al. "Connectome harmonic decomposition of human brain dynamics reveals a landscape of consciousness". In: *BioRxiv* 10.2020.08 (2020), pp. 10–244459.

[25] S Parker Singleton et al. "LSD and psilocybin flatten the brain's energy landscape: insights from receptor-informed network control theory". May 2021.

[26] Katrin H Preller et al. "Changes in global and thalamic brain connectivity in LSD-induced altered states of consciousness are attributable to the 5-HT2A receptor". en. In: *Elife* 7 (Oct. 2018).

[27] Enzo Tagliazucchi et al. "Increased global functional connectivity correlates with LSD-induced ego dissolution". en. In: *Curr. Biol.* 26.8 (Apr. 2016), pp. 1043–1050.

[28] Giulio Ruffini. "An algorithmic information theory of consciousness". In: *Neuroscience of Consciousness* 3.1 (2017).

[29] Jakub Vohryzek et al. "Understanding brain states across spacetime informed by whole-brain modelling". In: *Philosophical Transactions of the Royal Society A* 380.2227 (2022), p. 20210247.

[30] Arthur Juliani, Adam Safron, and Ryota Kanai. "Deep CANALs: A Deep Learning Approach to Refining the Canalization Theory of Psychopathology". In: (2023).

[31] David E Nichols, Matthew W Johnson, and Charles D Nichols. "Psychedelics as medicines: an emerging new paradigm". In: *Clinical Pharmacology & Therapeutics* 101.2 (2017), pp. 209–219.

[32] Min-Fang Kuo and Michael A Nitsche. "Effects of transcranial electrical stimulation on cognition". In: *Clinical EEG and neuroscience* 43.3 (2012), pp. 192–199.

[33] Lawrence Cayton. *Algorithms for manifold learning*. Tech. rep. UC San Diego: Department of Computer Science & Engineering, 2008. URL: https://escholarship.org/uc/item/8969r8tc.

[34] Nicholas Konz et al. "Correction to: The intrinsic manifolds of radiological images and their role in deep learning". In: *Lecture Notes in Computer Science*. Lecture notes in computer science. Cham: Springer Nature Switzerland, 2022, pp. C1–C1.

[35] Yonatan Sanz Perl et al. "Generative Embeddings of Brain Collective Dynamics Using Variational Autoencoders". In: *Physical Review Letters* 125.23 (Dec. 2020). DOI: 10.1103/physrevlett.125.238101. URL: https://doi.org/10.1103/physrevlett.125.238101.

[36] Yonatan Sanz Perl et al. "Model-based whole-brain perturbational landscape of neurodegenerative diseases". In: *eLife* 12 (Mar. 2023). Ed. by Muireann Irish, Timothy E Behrens, and Jordi A Matias-Guiu, e83970. ISSN: 2050-084X. DOI: 10.7554/eLife.83970. URL: https://doi.org/10.7554/eLife.83970.

[37] Steven H. Strogatz. *Nonlinear Dynamics and Chaos: With Applications to Physics, Biology, Chemistry, and Engineering*. 1st. Westview Press, 2001. ISBN: 978-0738204536.

[38] R L Carhart-Harris and K J Friston. "REBUS and the anarchic brain: Toward a unified model of the brain action of psychedelics". en. In: *Pharmacol. Rev.* 71.3 (July 2019), pp. 316–344.

[39] J G Jefferys. "Nonsynaptic modulation of neuronal activity in the brain: electric currents and extracellular ions". en. In: *Physiol. Rev.* 75.4 (Oct. 1995), pp. 689–723.

[40] Costas A Anastassiou et al. "Ephaptic coupling of cortical neurons". en. In: *Nat. Neurosci.* 14.2 (Feb. 2011), pp. 217–223.

[41] Giulio Ruffini et al. "Realistic modeling of mesoscopic ephaptic coupling in the human brain". en. In: *PLoS Comput. Biol.* 16.6 (June 2020), e1007923.

[42] Dimitris A Pinotsis and Earl K Miller. "In vivo ephaptic coupling allows memory network formation". en. In: *Cereb. Cortex* (July 2023).

[43] Donald Olding Hebb. *The organization of behavior: A neuropsychological theory*. New York, NY: John Wiley and Sons, 1949.

[44] Gina G. Turrigiano et al. "Activity-Dependent Scaling of Quantal Amplitude in Neocortical Neurons". In: *Nature* 391.6670 (Feb. 1998), pp. 892–896. ISSN: 1476-4687. DOI: 10.1038/36103. (Visited on 12/01/2023).

[45] Gina G Turrigiano and Sacha B Nelson. "Homeostatic plasticity in the developing nervous system". In: *Nature reviews neuroscience* 5.2 (2004), pp. 97–107.

[46] Arianna Maffei, Sacha B. Nelson, and Gina G. Turrigiano. "Selective Reconfiguration of Layer 4 Visual Cortical Circuitry by Visual Deprivation". In: *Nature Neuroscience* 7.12 (Dec. 2004), pp. 1353–1359. ISSN: 1546-1726. DOI: 10.1038/nn1351. (Visited on 12/01/2023).

[47] Keith B. Hengen et al. "Firing Rate Homeostasis in Visual Cortex of Freely Behaving Rodents". In: *Neuron* 80.2 (Oct. 2013), pp. 335–342. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2013.08.038. (Visited on 12/01/2023).

[48] Peter J. Hellyer et al. "Local Inhibitory Plasticity Tunes Macroscopic Brain Dynamics and Allows the Emergence of Functional Brain Networks". In: *NeuroImage* 124.Pt A (Jan. 2016), pp. 85–95. ISSN: 1053-8119. DOI: 10.1016/j.neuroimage.2015.08.069. (Visited on 12/01/2023).

[49] Zhengyu Ma et al. "Cortical Circuit Dynamics Are Homeostatically Tuned to Criticality In Vivo". In: *Neuron* 104.4 (Nov. 2019), 655–664.e4. ISSN: 0896-6273. DOI: 10.1016/j.neuron.2019.08.031. (Visited on 12/01/2023).

[50] Marina Chistiakova et al. "Heterosynaptic plasticity: multiple mechanisms and multiple roles". In: *The Neuroscientist* 20.5 (2014), pp. 483–498.

[51] Lauren Lepow, Hirofumi Morishita, and Rachel Yehuda. "Critical period plasticity as a framework for psychedelic-assisted psychotherapy". In: *Frontiers in neuroscience* 15 (2021), p. 710004.

[52] W C Abraham and M F Bear. "Metaplasticity: the plasticity of synaptic plasticity". en. In: *Trends Neurosci.* 19.4 (Apr. 1996), pp. 126–130.

[53] Linnea R Vose and Patric K Stanton. "Synaptic plasticity, metaplasticity and depression". en. In: *Curr. Neuropharmacol.* 15.1 (2017), pp. 71–86.

[54] Charles W Misner, Kip S Thorne, and John Archibald Wheeler. *Gravitation*. en. Princeton, NJ: Princeton University Press, Oct. 2017.

[55] Robin L Carhart-Harris and DJ Nutt. "Serotonin and brain function: a tale of two receptors". In: *Journal of psychopharmacology* 31.9 (2017), pp. 1091–1120.

[56] Robin Carhart-Harris et al. "Trial of psilocybin versus escitalopram for depression". en. In: *N. Engl. J. Med.* 384.15 (Apr. 2021), pp. 1402–1411.

[57] Rodrigo Andrade. "Serotonergic regulation of neuronal excitability in the prefrontal cortex". en. In: *Neuropharmacology* 61.3 (Sept. 2011), pp. 382–386.

[58] Michael M. Schartner et al. "Increased Spontaneous MEG Signal Diversity for Psychoactive Doses of Ketamine, LSD and Psilocybin". In: *Scientific Reports* 7.1 (Apr. 2017), p. 46421. ISSN: 2045-2322. DOI: 10.1038/srep46421. (Visited on 12/01/2023).

[59] Thomas F. Varley et al. "Serotonergic Psychedelics LSD & Psilocybin Increase the Fractal Dimension of Cortical Brain Activity in Spatial and Temporal Domains". In: *NeuroImage* 220 (Oct. 2020), p. 117049. ISSN: 1053-8119. DOI: 10.1016/j.neuroimage.2020.117049. (Visited on 12/01/2023).

[60] Zen J. Lau et al. "Brain Entropy, Fractal Dimensions and Predictability: A Review of Complexity Measures for EEG in Healthy and Neuropsychiatric Populations". In: *The European Journal of Neuroscience* 56.7 (Oct. 2022), pp. 5047–5069. ISSN: 0953-816X. DOI: 10.1111/ejn.15800. (Visited on 12/01/2023).

[61] D. E. Nichols. "Psychedelics". In: *Pharmacological Reviews* 68 (2 2016), pp. 264–355. DOI: 10.1124/pr.115.011478.

[62] Alex C. Kwan et al. "The neural basis of psychedelic action". In: *Nature Neuroscience* 25.11 (2022), pp. 1407–1419. ISSN: 1546-1726. DOI: 10.1038/s41593-022-01177-4. URL: https://doi.org/10.1038/s41593-022-01177-4.

[63] Abigail E Calder and Gregor Hasler. "Towards an understanding of psychedelic-induced neuroplasticity". en. In: *Neuropsychopharmacology* 48.1 (Jan. 2023), pp. 104–112.

[64] James JH Rucker, Jonathan Iliff, and David J Nutt. "Psychiatry & the psychedelic drugs. Past, present & future". In: *Neuropharmacology* 142 (2018), pp. 200–218.

[65] Tanya Calvey and Fleur M Howells. "An introduction to psychedelic neuroscience". In: *Progress in brain research* 242 (2018), pp. 1–23.

[66] Gisela D Charó et al. "Noise-driven topological changes in chaotic dynamics". en. In: *Chaos* 31.10 (Oct. 2021), p. 103115.

[67] Charles W Misner, Kip S Thorne, and Wojciech H Zurek. "John Wheeler, relativity, and quantum information". en. In: *Phys. Today* 62.4 (Apr. 2009), pp. 40–46.

[68] Manfredo P. do Carmo. *Differential Geometry of Curves and Surfaces*. Upper Saddle River, NJ, USA: Prentice-Hall, 1976. ISBN: 978-0-13-212589-5.

[69] Stephen W. Hawking and George F. R. Ellis. *The Large Scale Structure of Space-Time*. Cambridge: Cambridge University Press, 1973. ISBN: 978-0521099066.

[70] Jean-Pierre Luminet. "The Status of Cosmic Topology after Planck Data". In: *Universe* 2.1 (2016). ISSN: 2218-1997. DOI: 10.3390/universe2010001. URL: https://www.mdpi.com/2218-1997/2/1/1.

[71] Robert Gilmore. "Topological analysis of chaotic dynamical systems". In: *Rev. Mod. Phys.* 70 (4 Oct. 1998), pp. 1455–1529. DOI: 10.1103/RevModPhys.70.1455. URL: https://link.aps.org/doi/10.1103/RevModPhys.70.1455.

[72] Gunnar Carlsson. "Topological pattern recognition for point cloud data". en. In: *Acta Numer.* 23 (May 2014), pp. 289–368.

[73] Chad Giusti, Robert Ghrist, and Danielle S Bassett. "Two's company, three (or more) is a simplex : Algebraic-topological tools for understanding higher-order structure in neural data". en. In: *J. Comput. Neurosci.* 41.1 (Aug. 2016), pp. 1–14.

[74] Michael W Reimann et al. "Cliques of neurons bound into cavities provide a missing link between structure and function". en. In: *Front. Comput. Neurosci.* 11 (June 2017), p. 48.

[75] Jose A Perea. "Topological time series analysis". en. In: *Not. Am. Math. Soc.* 66.05 (May 2019), p. 1.

[76] Denisse Sciamarella and G B Mindlin. "Topological structure of chaotic flows from human speech data". In: *Phys. Rev. Lett.* 82.7 (Feb. 1999), pp. 1450–1453.

[77] Samuel E Gralla, Abraham I Harte, and Robert M Wald. "Rigorous derivation of electromagnetic self-force". en. In: *Phys. Rev.* 80.2 (July 2009).

[78] John David Jackson. *Classical Electrodynamics.* en. 3rd ed. Nashville, TN: John Wiley & Sons, July 1998.

[79] Gustavo Deco et al. "Awakening: Predicting external stimulation to force transitions between different brain states". In: *Proceedings of the National Academy of Sciences* 116.36 (2019), pp. 18088–18097.

[80] Morten L Kringelbach et al. "Toward naturalistic neuroscience: Mechanisms underlying the flattening of brain hierarchy in movie-watching compared to rest and task". In: *Science Advances* 9.2 (2023), eade6049.

A   Appendix

A.1   Classical dynamics of particles and fields

Here we provide the equations for other systems where one can think of part of the equation describing the geometry of a space-providing subsystem ("kinematic theatre") and another the subsystem moving in this space, influenced by the structure and then affecting its geometry in return. Several such examples can be found in physics.

Non-relativistic electrodynamics

The non-relativistic dynamics of N-charged particles and the associated electromagnetic field are described by (coupled) Newton's second law and Maxwell's equations. The charged particles are influenced by the electromagnetic field and at the same time, generate it. This can lead to problematic scenarios, such as the self-interaction problem: charged particles generate an electromagnetic field, and if one considers a particle's interaction with its own field, paradoxical or unphysical results can arise. This self-interaction leads to divergences in calculations and has been a longstanding challenge in classical electrodynamics until recently [77].

The motion of the ith electron is given by $m_i \frac{d^2 \mathbf{r}_i}{dt^2} = \mathbf{F}_i$, where $\mathbf{F}_i = q_i(\mathbf{E} + \mathbf{v}_i \times \mathbf{B})$ is the Lorentz force. The electromagnetic field obeys Maxwell's equations:

$$
\nabla \cdot \mathbf{E} = \frac{\rho}{\varepsilon_0},
$$
$$
\nabla \cdot \mathbf{B} = 0,
$$
$$
\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t},
$$
$$
\nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \varepsilon_0 \frac{\partial \mathbf{E}}{\partial t},
$$

where $\rho(\mathbf{r}, t) = \sum_i q_i \delta(\mathbf{r} - \mathbf{r}_i(t))$ and $\mathbf{J}(\mathbf{r}, t) = \sum_i q_i \mathbf{v}_i(t) \delta(\mathbf{r} - \mathbf{r}_i(t))$ are the charge and current densities.

Relativistic equations

Maxwell's equations are relativistic (they transform properly under Lorentz transformations), but Newton's is not. For the **relativistic version**, the field strength tensor $F^{\mu\nu}$ is defined in terms of the four-potential A^μ [78],

$$
F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu
$$

683 The dual tensor $\tilde{F}^{\mu\nu}$ is defined in terms of $F_{\alpha\beta}$ and the Levi-Civita symbol $\varepsilon^{\mu\nu\alpha\beta}$,

$$\tilde{F}^{\mu\nu} = \frac{1}{2}\varepsilon^{\mu\nu\alpha\beta}F_{\alpha\beta}$$

684 The homogeneous Maxwell's equations are given by:

$$\partial_\mu\tilde{F}^{\mu\nu} = 0$$

685 and the inhomogeneous Maxwell's equations are given by:

$$\partial_\mu F^{\mu\nu} = \mu_0 J^\nu$$

686 where J^ν is the four-current vector, which we now define for a particular case. Given a
687 distribution of N particles each with charge q_i and four-velocity u_i^μ, the four-current J^μ
688 at position $\mathbf{x}$ and time t is given by

$$J^\mu(\mathbf{x}, t) = \sum_{i=1}^{N} q_i u_i^\mu \delta^3(\mathbf{x} - \mathbf{x}_i(t))$$

689 Here, $\mathbf{x}_i(t)$ is the position of the i-th particle at time t, and δ^3 is the three-dimensional
690 Dirac delta function.

691 The equation of motion for a charged particle in an electromagnetic field, commonly
692 known as the Lorentz force equation, in its relativistic form is:

$$\frac{dp^\mu}{d\tau} = qF^{\mu\nu}u_\nu$$

693 Here, p^μ is the four-momentum of the particle, u^ν is the four-velocity of the particle,
694 $F^{\mu\nu}$ is the electromagnetic field tensor, and q is the charge of the particle. The equation
695 describes how the four-momentum of the particle changes with proper time τ under the
696 influence of the electromagnetic field.

A.2 Modeling plasticity in neural mass models

In this section, we provide a brief overview of plasticity mechanisms and how they relate to the terms in the formalism, namely the functions h and ψ. See Table 1 for a summary. Including plasticity in Neuronal Mass Models (NMMs) allows for the modeling of time-varying connectivity strengths that reflect the learning and adaptation processes observed in biological neuronal networks.

Functional Plasticity

The simplest and most common way to include synaptic plasticity is through Hebbian learning rules. Hebbian plasticity, a type of functional plasticity, follows the principle that "neurons that fire together wire together" [43]. It can be included in NMM using the equation

$$\dot{w}_{ij} = \eta x_i x_j - \delta w_{ij} \tag{7}$$

where w_{ij} is the synaptic strength from neuron j to neuron i, x_i and x_j are the neuronal activities, and η and δ are parameters controlling the learning and decay rates.

Homeostatic Plasticity

Homeostatic plasticity, a form of plasticity that adjusts synaptic strengths to keep the overall activity of a neuron or network within a certain range, can be included in an NMM using the equation [44–47],

$$\dot{w}_{ij} = \eta(x_i - x_{\text{target}})x_j - \delta w_{ij} \tag{8}$$

where x_{target} is the target activity level.

Structural Plasticity

Structural plasticity, where the actual number and dendrites and arrangement of synapses change over time, can be represented in NMMs by modifying (or even adding) rows and columns from the w adjacency matrix to represent the formation or elimination of synapses or even fibers.

Empirically-derived Structural Plasticity

NMMs can be used to infer the structural changes to plasticity without explicitly describing the plastic mechanism per se. For example, in the post-acute psychedelic state, NMMs can be used to infer the plastic changes to w_{ij} by optimizing the model functional

connectivity FS_{ij}^{model} to approximate the empirical functional connectivity FS_{ij}^{model} with a certain learning rate ϵ as in the following equation.

$$w_{ij} = w_{ij} + \epsilon(FS_{ij}^{empirical} - FS_{ij}^{model}) \tag{9}$$

Such optimization is, for example, computed through gradient descent methods with priors on the topology of structural connectivity between brain regions [79]. Recent methods have further extended this framework by adding time-shifted correlation [80] to the optimization as a better description of the overall brain state, as in this case, the post-acute psychedelic state.

Including these forms of plasticity in NMMs allows for more realistic modeling of neural systems in better capturing their adaptive nature and the impact of learning and experience on synaptic connections.

Type	Time Scale	Mechanism	Effects	State-dependence
Functional or dynamical Plasticity	Milliseconds to minutes	Changes in strength/efficiency of synapses (e.g., Hebbian plasticity, LTP, LTD)	Short and long-term memory, fine-tuning of connections	State-dependent (h)
Acute Psychedelic-induced Plasticity	Minutes to Hours	Targeting of serotonergic neuroreceptors especially the 5-HT$_{2A}$ with a result in overall excitability [11, 12, 61, 62]	Flattens or de-weights the dynamical landscape	State-independent (ψ)
Homeostatic Plasticity	Hours to days	Regulation of overall excitability to maintain stability (e.g., adjusting synapse strength for E/I balance)	Balances and stabilizes network	State-dependent (h)
Structural Plasticity (e.g., post-acute psychedelic-induced plasticity)	Hours to years	Larger physical changes in neurons (e.g., dendrite growth, synapse formation) through an increase of endogenous BDNF and via TrkB binding (the receptor of BDNF) [63]	Long-term memory, development	Depends on context
Metaplasticity	Various, often longer-term	Changes in mechanisms governing synaptic plasticity (e.g., modulation of thresholds/rules)	Regulates other forms of plasticity, "plasticity of plasticity" [52]	Depends on context

Table 1: Summary of Different Types of Neural Plasticity Phenomena. **State-dependent Plasticity** (h) refers to changes in neural connections that depend on the current state or activity of the neurons involved. For example, functional plasticity often relies on specific patterns of neural activity to induce changes in synaptic strength. **State-independent Plasticity** (ψ) refers to changes that are not directly dependent on the specific activity state of the neurons. For example, acute psychedelic-induced plasticity acts on the serotonergic neuroreceptors and thus acts on the brain networks regardless of specific activity patterns. Some forms of plasticity, like structural plasticity and metaplasticity, may exhibit characteristics of both state-dependent and state-independent plasticity, depending on the context and specific mechanisms involved. Finally, **metaplasticity** refers to the adaptability or dynamics of plasticity mechanisms.