Population interconnectivity shapes the distribution and complexity of chimpanzee cumulative culture

Authors: Cassandra Gunasekaram[1]*, Martin Kuhlwilm[2,3], Federico Battiston[4], Claudia Fontsere[5], Cecilia Padilla-Iglesias[1], Ammie K. Kalan[6], Maria A. van Noordwijk[1,7], Andrea Manica[8], Susana Carvalho[9,10], Jaume Betranpetit[11], Tomas Marques-Bonet[12], Andrew Whiten[13], Carel P. van Schaik[1,7], Lucio Vinicius[1], Andrea Bamberg Migliano[1]*

Affiliations:

[1]Department of Evolutionary Anthropology, University of Zurich, Zurich, Switzerland

[2]Department of Evolutionary Anthropology, University of Vienna, Vienna, Austria

[3]Human Evolution and Archaeological Sciences (HEAS), University of Vienna, Vienna, Austria

[4]Department of Network and Data Science, Central European University, Vienna, Austria

[5]Center for Evolutionary Hologenomics, The Globe Institute, University of Copenhagen, Copenhagen, Denmark

[6]Department of Anthropology, University of Victoria, Victoria BC, Canada

[7]Comparative Socioecology, Max Planck Institute of Animal Behavior, Konstanz, Germany

[8]Department of Zoology, University of Cambridge, Cambridge, UK

[9]Primate Models for Behavioural Evolution Lab, Institute of Human Sciences, University of Oxford, Oxfordshire, UK

[10]Interdisciplinary Center for Archaeology and Evolution of Human Behaviour ((ICArEHB), Universidade do Algarve, Faro, Portugal

[11]Institut de Biologia Evolutiva, CSIC - Universitat Pompeu Fabra, Barcelona, Spain

23   [12]Catalan Institution of Research and Advanced Studies (ICREA), Universitat Pompeu

24   Fabra, Barcelona, Catalunya, Spain

25   [13]Centre for Social Learning and Cognitive Evolution, School of Psychology and

26   Neuroscience, University of St Andrews, St Andrews, UK

27   *Corresponding author. Email: cassandra.gunasekaram@uzh.ch, andrea.migliano@uzh.ch

Abstract: While cumulative culture is a hallmark of hominin evolution, its origins can be traced back to our common ancestor with chimpanzees. Here we investigate the evolutionary origins of chimpanzee cumulative culture, and why it remained incipient. To trace cultural transmission among the four chimpanzee subspecies, we built between-populations networks based on genetic markers of recent migration and shared cultural traits. We show that limited levels of interconnectivity favoured the emergence of a few instances of cumulative culture in chimpanzees. As in humans, cultural complexification likely happened in steps, with between-community transmission promoting incremental changes and repurposing of technologies. We propose that divergence in social patterns led to increased between-group mobility in *Homo*, propelling our lineage towards a trajectory of irreversible dependence on cultural complexification.

One-Sentence Summary:

Population interconnectivity and migration explain the origins of chimpanzee cumulative culture and why it remained incipient

Culture, defined as arrays of behavioural traditions transmitted via social learning (Whiten and van Schaik, 2007), has been increasingly documented across animal species (Jesmer et al., 2018; Schuppli and van Schaik, 2019; Whiten, 2021). Human culture is nevertheless vastly more complex, diverse, and most crucially, cumulative (Henrich, 2017; Mesoudi and Thornton, 2018; Tomasello et al., 1993). Human cumulative culture provides numerous examples of cultural ratcheting (Tomasello et al., 1993; Tennie et al., 2009, 2020; Whiten 2022, 2023), an evolutionary process that builds cultural traits upon pre-existing ones. Eventually their production may thence become opaque and they can no longer be reinvented from scratch. Cultural ratcheting has only recently been investigated in other species (Rawlings et al., 2021). Chimpanzees, our closest living relatives, exhibit a vast repertoire of cultural behaviours that vary across sites and are only weakly tied to their habitat (Kalan et al., 2020; Kühl et al., 2019; Nishida et al., 2010; Whiten et al., 1999, 2001). Chimpanzee culture also includes possible instances of cultural accumulation, most evidently in the case of foraging toolsets or multi-step extractive processes (Boesch, 2012; McGrew, 1992, 2010; Nishida et al., 2010; Boesch et al., 2020). It remains to be explained what conditions favoured the emergence of cultural accumulation in the common ancestor of humans and chimpanzees, and why it remained incipient in chimpanzees in comparison to humans and other tool-making hominins.

Recent analyses have argued that a major role in the cultural differentiation between the two lineages emerged in the evolution of a hominin foraging niche characterised by higher levels of individual mobility and migration (Migliano and Vinicius, 2022). Group interconnectivity was shown to facilitate cultural accumulation by preventing erosion of cultural diversity (Henrich, 2004) and accelerate the recombination of innovations through between-group transfers (Derex and Boyd, 2016; Migliano et al., 2020). Demonstrating the long-term operation of partial connectivity in humans, genetic and archaeological studies have shown that Central African

hunter-gatherers have maintained large-scale networks through migration over the past 120,000 years (Padilla-Iglesias et al., 2022a), as evidenced by varied signs of cultural and transmission even between distinct linguistic groups across the whole Congo Basin (Padilla-Iglesias et al., 2022b).

Investigating levels of interconnectivity between communities may therefore provide relevant insights into cumulative cultural evolution in chimpanzees. Genetic analyses of evolutionary divergence and recent migration among the four chimpanzee subspecies (Fontsere et al., 2022) revealed a very different pattern of interconnectivity between populations relative to Central African hunter-gatherers. Since the early separation of the four subspecies between 139-633 kya (de Manuel et al., 2016), low levels of genetic connectivity between the Nigeria-Cameroon, Central, and Eastern subspecies were found during the late Pleistocene and Holocene (Fontsere et al., 2022). While shared nearly-private rare alleles (NePRAs) provide evidence for reduced connectivity between populations from 15,000 years ago, shared identical-by-descent (IBD) genetic segments suggest some level of migration between Central, Nigeria Cameron and Eastern regions in the past 5,000 years (Fontsere et al., 2022). Besides being relatively isolated on a regional level, chimpanzees live in large, polygynandrous communities with steep social hierarchies and strict male philopatry, which virtually eliminates male migration (Pusey & Packer, 1986). This contributes to a much lower frequency of between-group exchanges in chimpanzees compared to humans, further limiting cultural diffusion to migrating females (Luncz & Boesch, 2014). As a result, while an adult hunter-gatherer male may interact with around 300 other tool-making individuals throughout a lifespan, an adult male chimpanzee habitually observes only around 20 others (Hill et al., 2014), or around 40 in the case of females, who typically migrate once in a lifetime.

Given the different social structure of chimpanzees, here we investigate whether the

incipient cumulative culture observed in chimpanzees is associated with their relatively modest

levels of between-group connectivity and cultural transmission. We combined genetic

information on recent between-group migration with chimpanzee cultural data compiled by the

Pan African Program (Lester et al., 2021). The latter has built upon earlier surveys (Whiten et al.,

2001; Nishida et al., 2010) to catalogue cultural repertoires in 144 chimpanzee communities

across Africa (Kalan et al., 2020; Kühl et al., 2019), resulting in a growing dataset of

community-level cultural traits (Boesch 2012; Boesch et al., 2020; Carvalho et al., 2013; Sanz et

al., 2009). Within this dataset, 35 populations match the NePRA dataset (Fig. 1A and 1B) and 34

populations match the IBD dataset (Fig 1C), allowing us to trace group interconnectivity over the

past 15,000 years.

We used our data on recent genetic connectivity to predict the sharing of 15 foraging

behaviours between chimpanzee populations. We classified foraging behaviours into three

categories of increasing complexity (see SI for details): non-tool behaviours (Fig. 1D), 'simple',

unitary tool use (Fig. 1E), and 'complex' toolset behaviours (Fig 1F) (Kühl et al., 2019; Kalan et

al., 2020, Pradhan, et al., 2012). We tested the following predictions: (1) foraging without tools

and with simple tools should exhibit weak associations with genetic markers of group

interconnectivity: they should exhibit a broader distribution across the four subspecies and

regions due to a higher likelihood of independent invention at the evolutionary time scale; (2)

complex toolsets should be largely explained by genetic markers of recent migration: their

distribution should be more reliant on cultural transmission, and therefore more regionally

limited due to reduced group interconnectivity in comparison to humans; and (3) as in humans,

cumulative cultural evolution in chimpanzees is likely to be a stepwise process facilitated by

transmission across multiple individuals, generations and populations.

Figure 1. Cultural and genetic connections across chimpanzee populations. Links represent

genetic connections based on genetic markers of migration (A-C) or shared knowledge of

foraging behaviours (D-F). A) 35 chimpanzee populations included in this study, with habitat

and subspecies at each site. B) NePRA connections, with colours indicating proportion of shared

NePRAs. C) Mean of maximum IBD length (mega base pairs, Mbp) shared between two sites.

D) Shared non-tool behaviours. E) Shared simple tool behaviours. F) Shared complex toolset

behaviours.

Results

Complex toolsets rely on cultural transmission between populations

To analyse the effect of genetic connectivity on shared knowledge of behavioural traits, we applied Bayesian logistic regression models. We fitted models to predict the dyadic sharing of 15 behaviours (coded as a binary response variable) between populations. We fit separate models for the sharing of the three behavioural categories (non-tool, simple tool, complex toolset; see SI for details of regressions and full code). As the choice of predictors, all regressions included habitat (savannah or forest) and months of observation at each site as fixed effects; and both type of behaviour (e.g. ant eating, termite eating, and honey eating in regressions of non-tool sharing) and dyad ID as random intercept effects (see Fig. S3 for random intercept and slope models, which overall render similar results). Each regression included a single genetic or geographic predictor at a time, because genetic markers, region and geographical distances are intercorrelated; see SI for correlation tests).

Genetic markers of recent between-group migration revealed a clear pattern. Dyadic IBD sharing (binary variable; sharing any vs. no IBD segment) had a strong effect on complex toolset sharing (odds ratio (OR) = 3.19, 95% credible interval (CI): [1.39, 7.68]) but no effect on non-tool behaviour (OR = 0.77, CI: [0.29, 2.01]), and a weaker effect on simple tool sharing (OR = 1.41, CI: [0.98, 2.00]); similar results were obtained using shared IBD segment length (Fig. 2A). NePRA sharing also had the strongest effect on complex toolset sharing (binary: OR = 5.69, CI: [2.14, 17.15], proportions: OR = 3.15, CI: [1.34, 7.32]), a weak effect on simple tool sharing (binary: OR = 1.42, CI: [0.98, 2.04], proportions: OR = 1.45, CI: [1.31, 1.70]), and no effect on the sharing of non-tool behaviours (binary: OR = 1.00, CI: [0.46, 2.22], proportions: OR = 0.93, CI: [0.63, 1.41]) (Fig. 2B, see SI for details). Dyadic region (binary; same vs. different region) supported these results, with same-region populations sharing more complex toolsets (OR =

3.88, CI: [1.78, 8.51], Fig. S5). Geographical distance also confirmed the pattern with a strong

negative effect on complex toolset sharing (OR = 0.42, 95% CI: [0.28, 0.61], Fig. S5),

suggesting increased sharing of complex toolsets with shorter distances (Fig. 2C). The

associations between genetic markers and complex tool sharing were confirmed by Mantel tests

even after controlling for habitat or observation time (Fig. S7). Overall, the results demonstrate a

strong association between group interconnectivity and the sharing of complex toolsets.

Figure 2. Bayesian logistic regressions of shared cultural traits between chimpanzee

populations. Panels display log of the odds ratio estimates and the 65%, 85% and 95% credible

intervals. A) Effect of shared IBD segments (binary = sharing any vs. no IBD, IBD length =

population-average of longest IBD segment shared between individuals) on non-tool, simple

tool, and complex toolset behaviours. B) Effects of sharing any NePRA (binary; above or below

a noise threshold at a similarity of 0.0006, see SI) and proportion of shared NePRAs (see

Methods).

Migration is a key but not sufficient condition for complex toolset sharing

To further quantify the effects of recent migration on non-tool, simple tool and complex

toolset sharing, we applied reinforcement analysis (Battiston et al. 2014). The technique is

derived from social network analysis and compares two networks with the same nodes (in our

case, populations) but different structures to determine whether one is likely to be the condition

for the existence of the other. Specifically, if the links of network A strongly predict the presence

of their corresponding links in network B, then the links of network A are strongly conditioned

(or reinforced) by the presence of the links in network B (see SI for further details and

examples), providing a possible hint at the directionality of effects between two correlated

networks. We built two sets of networks, one with links representing the sharing of a genetic

marker (IBD segment or NePRA) between two populations (Fig. 1B, C), and the other with

shared cultural behaviours between two populations (non-tool, simple tool and toolset) as links

(Fig. 1D-F). We predicted that the sharing of a complex toolset (but not of non-tool or simple

tool behaviours) should strongly predict the presence of a genetic marker (as evidence that

complex tool sharing requires migration). Conversely, we did not expect a shared genetic marker

to strongly predict the sharing of any of the cultural behaviours (since the latter is not a condition

for migration, or alternatively, since cultural transmission depends not only on migration but on

other factors, as discussed below).

We first tested whether links representing shared behaviours of increasing cultural

complexity predict the presence of an underlying shared IBD segment or NePRA between the

same populations. We found that shared complex toolsets predict a shared IBD segment with

probabilities over 50%, and a strong NePRA similarity in over 80% of cases, pointing to the

dependence of complex toolsets on migration. Values for non-tool behaviours and simple tools

are much lower (Fig. 3A, B). Conversely, IBD segment sharing, or NePRA sharing (at three

increasing levels of similarity) predicted a very small fraction of cultural links at any complexity

level (Fig. 3C, D). Together, the results show that at least 80% of the instances of complex

toolset sharing are associated with a marker of migration in the past 15,000 years. The values are

much lower for simple tools and non-tool behaviours, suggesting that they are more likely to be

independently invented. On the other hand, only a small fraction of all genetic links reflecting

recent migration between populations results in cultural sharing of any behavioural type. This

confirms that the studied behaviours show no signs of genetic determination (Lycett et al., 2010),

and also indicates that successful cultural transmission through migration may be limited by

factors such as the skill levels of migrant females, conformity to community behaviours by low-

ranked migrants, or absence of ecological context for expression of the behaviour (Kendal et al.,

2015; Luncz & Boesch, 2014).

Figure 3. Reinforcement analyses of genetic and cultural sharing networks. A) Probability

of the presence of IBD segment links as a function of presence of the corresponding cultural

links for the three complexity levels; B) Probability of NePRA sharing at three increasing levels

of similarity as a function of cultural links; Panels C) and D) show the probability of presence of

links of each behavioural category as a function of IBD links and NePRA similarity links

respectively (D). Coloured symbols represent the distribution of values within each category.

Evolution of complexity and cumulative culture in chimpanzees

Interconnectivity plays a role in the evolution of cultural complexity in humans by

promoting cultural aggregation and recombination (Derex and Boyd, 2016; Migliano et al.,

2022). We therefore investigated possible signs of stepwise cultural evolution in networks of

chimpanzee populations. We assessed evidence of possible evolutionary links between the

simple and complex forms of the same underground foraging technology (indicating possible

stepwise aggregation), and of repurposing (or major shifts in the use of the technology).

Underground foraging tools exhibit distinctive traits of cumulative culture such as variability in

number and sequence of tools across populations (Boesch, 2012; Boesch et al., 2020; Sanz et al.,

2009), and opacity for naive individuals since target resources are hidden from direct observation

(Boesch, 2012). Interestingly, reinforcement analysis produced evidence of a role of migration

on complex toolset sharing when all underground toolsets were placed into a single category,

suggesting the possibility of an evolutionary link between them (Fig. 3). We compiled all

instances of underground foraging in the Central, Nigeria-Cameroon and Eastern regions,

including both simple tools (e.g. digging tool) and complex toolsets (e.g. digging stick followed

by a dipping tool); exploited resources (honey, termites and ants) from previously published

literature (Table S3); and combined this with information on recent genetic links between

populations where they are found (shared IBD segments and NePRAs). When we plotted all

populations where underground foraging is found and then added their genetic links, we

observed that no single population was disconnected from others in the resulting network (Fig. 4A & Fig. 4b). The IBD segment and NePRA networks extend across the three regions through several long links, resembling a pattern often found in hunter-gatherer and other human networks (Granovetter, 1973; Padilla-Iglesias et al., 2022b).

This observation further suggests that underground resource foraging may have evolved through incremental changes from use of a single honey digging tool, as found in the East (Bili-Uele and Nyungwe) and in the Central region (La Belgique), into complex toolsets found in the Central and Nigeria-Cameroon regions. For example, Bili-Uele (East), which exhibits only simple underground honey extraction, is directly linked in the shared IBD segment network to Gashaka (Nigeria-Cameroon), where an underground honey toolset is observed, possibly as an example of stepwise aggregation (or addition of a dipping tool to the existing digging tool).

Moreover, in the IBD segment network the two populations using underground honey toolsets (Gashaka in Nigeria-Cameroon, and Loango in the Central region) are connected to populations displaying complex underground termite foraging such and Korup (Nigeria-Cameroon), Goualougo and Mt. Cristal (Central). This observation and the lack of a simple underground termite foraging variant suggest a possible repurposing of a pre-existing honey toolset for underground termite foraging, implying a major innovation in resource exploitation. Since IBD segments are genetic markers of migration in the past 5,000 years, contact between the three chimpanzee subspecies and tool complexification may have happened recently. Alternatively, the NePRA network tracking a longer evolutionary period (the past 15,000 years) suggests an alternative route of complexification, this time from the simple underground honey tools in Nyungwe (East) to the toolset found in Loango (Central) through incremental addition, followed by repurposing of the technology (from honey to termite foraging) in Goualougo (Central). Although we cannot differentiate between the two possibilities, Mantel tests (Fig.

S7D) identified a correlation of underground toolset with NePRAs but not with the more recent

IBD segments, pointing to an earlier spread and complexification of the technology within the

three subspecies.

In summary, a comparison across sites suggests a variety of markers of cumulative

culture in chimpanzee populations: the association between sharing of cultural traits and

migration, increased group connectivity and cultural complexity in network positions

characterised by increased genetic and cultural exchanges, incremental addition of innovations

underlying the origin or complex toolkits, and possibly major innovations with the adaptation of

a toolset for new foraging purposes.

Figure 4. Distribution of underground foraging in chimpanzee populations, underlying genetic links, and putative instances of cultural accumulation. All displayed examples match the description of behaviours in Table S3, apart from the honey probing toolset at Bulindi, which includes a digging stick and an additional probing stick most likely to gather bee larvae rather than honey (McLennan, 2011). At Ngel Nyaki, underground dipping tools are reported, but it is unclear whether they are used as single tools or as components of a complex toolset (Dutton & Chapman, 2015). Lines connect populations sharing (A) IBD segments and (B) strong NePRAs links.

Discussion

     In this study we investigated the conditions favouring the emergence of cumulative culture in chimpanzees, and why it has remained incipient in comparison to humans and earlier tool-making hominins. By analysing cultural and genetic data from 35 populations representing the four chimpanzee subspecies, we showed that group interconnectivity can account to a large extent for differences in the distributions of cultural foraging behaviours. We showed that foraging without tools and with simple tools are weakly associated with genetic markers of recent migration between populations, and are broadly distributed across the four subspecies. We interpret this pattern as a sign that while simple cultural behaviours can be culturally transmitted between populations, they are also more likely to be repeatedly invented at the evolutionary time scale. We stress that our view is compatible with studies demonstrating the extensive role of social learning in culture (Lonsdorf, 2003; Schuppli and van Schaik, 2019) and the low probabilities of individual recreation of cultural traits by naive chimpanzees (van Schaik et al., 2019; Tennie et al., 2009, 2020; Whiten, 2023), since the evolutionary scale of generations and populations provides the opportunity for unlikely events at the individual level to emerge in the long run.

     Compared to simple tools, toolsets are less likely to be reinvented due to their complexity and opacity, and accordingly they were strongly associated with genetic markers of recent migration and between-group transmission in the past 15,000 years. The finding suggests that chimpanzees display enough within- and between-region migration to foster at least an incipient level of cultural accumulation. It is especially interesting that all nut-cracking populations are connected to each other through IBD segment links, pointing to a spread of the technology within the past 5,000 years, matching archaeological evidence confirming the oldest nut-cracking sites at least 4,300 years ago (Mercader et al., 2007).

We conclude that cumulative cultural evolution in chimpanzees, as in humans, is a stepwise process facilitated by transmission across multiple individuals, generations and populations. From this perspective, the diversity of complex toolsets in chimpanzees, with nut-cracking limited to Western populations, and honey and termite extraction found in the Central, Eastern and Nigeria-Cameroon regions, results from the gradual and stepwise accumulation of techniques shared across interconnected groups, rather than allopatric differentiation due to isolation. Remarkably, populations that genetically interconnect three chimpanzee subspecies (Mt. Cristal, Gashaka and Goualougo) exhibit some of the highest levels of cultural complexity, further indicating that demographic exchange across regions promotes cultural exchange and complexification. The case study of underground foraging provided possible instances of incremental change as well as major adaptive shifts through repurposing of existing toolsets. We expect that the rich cultural record of chimpanzees can provide many other examples of associations between cultural accumulation and cultural transmission through migration, but unfortunately our analyses are limited to the 35 populations where both cultural and genetic data are currently available. Our study expanded on concepts of between-community cultural transmission proposed in the first systematic analyses of cultural variation in nine chimpanzee communities (Whiten et al., 2001) to examine cultural transmission across sites and subspecies, and we hope that in the future a growing dataset of genetic and cultural data will allow for more extensive testing of our predictions.

Finally, the reliance of cumulative culture on between-group interconnectivity also offers a possible explanation for the divergence between chimpanzees and the hominin lineage. The human foraging niche hypothesis (Migliano and Vinicius, 2022) proposes that cumulative culture has evolved at a faster pace in the ancestral lineage of humans due to the emergence of more extensive social networks in earlier hunting and gathering hominins. The argument points to

differences in sociality (Migliano et al., 2017), dispersal systems (Dyble et al., 2015, 2016),

population sizes (Powell et al., 2009), network structures (Derex et al., 2018; Migliano et al.,

2020), social bonding patterns (Kappeler and van Schaik, 2001), and cooperative breeding

(Burkart et al., 2009; Page et al., 2017) which resulted in larger networks and increased

connectivity, mobility and cooperation between groups. Archaeological studies indicate that the

pattern of population substructuring and interconnectivity may have characterised earlier

hominins (Blegen, 2017; Brooks et al., 2018; Scerri et al., 2019) and facilitated cumulative

cultural evolution. Increased interconnectivity and dependency on cultural exchange may have

selected for increased teaching, cooperative learning and joint innovation, setting humans on a

unique gene-culture coevolutionary pathway (Laland, 2003).

Acknowledgements. We thank William McGrew for his insightful comments on the draft.

Funding: C.G., C.P.-I. and A.B.M. have been funded by the University of Zurich, Switzerland.

M.K. has been funded by the Vienna Science and Technology Fund (WWTF)

[10.47379/VRG20001]. F.B. acknowledges support from the Air Force Office of Scientific

Research under award number FA8655-22-1- 7025.

References and Notes

Battiston, F., Nicosia, V., & Latora, V. (2014). Structural measures for multiplex
networks. *Physical Review E*, *89*(3), 032804.

Blegen, N. (2017). The earliest long-distance obsidian transport: Evidence from the~ 200 ka
Middle Stone Age Sibilo School Road Site, Baringo, Kenya. *Journal of Human Evolution*, *103*,
1-19.

Boesch, C. (2012). *Wild cultures: a comparison between chimpanzee and human cultures*.
Cambridge University Press.

Boesch, C., Kalan, A. K., Mundry, R., Arandjelovic, M., Pika, S., Dieguez, P., Ayimisin, E. A.,
Barciela, A., Coupland, C., Egbe, V. E., Eno-Nku, M., Fay, M., Fine, D., Hernandez-Aguilar, A.,
Hermans, V., Kadam, P., Kambi, M., Llana, M., Maretti, G., Morgan, D., Murai, M., Neil, E.,
Nicholl, S., Ormsby, L. J., Orume, R., Pacheco, L., Piel, A., Sanz, C., Sciaky, L., Stewart, F. A.,

Tagg, N., Wessling, E. G., Willie, J., & Kühl, H. S. (2020). Chimpanzee ethnography reveals unexpected cultural diversity. *Nature Human Behaviour*, *4*(9), 910-916.

Brooks, A. S., Yellen, J. E., Potts, R., Behrensmeyer, A. K., Deino, A. L., Leslie, D. E., Ambrose, S. H., Ferguson, J. R., d'Errico, F., Zipkin, A. M., Whittaker, S., Post, J., Veatch, E. G., Foecke, K., & Clark, J. B. (2018). Long-distance stone transport and pigment use in the earliest Middle Stone Age. *Science (80-) 360*, 90–94. (doi:10.1126/science.aao2646)

Burkart, J. M., Hrdy, S. B., & Van Schaik, C. P. (2009). Cooperative breeding and human cognitive evolution. *Evolutionary Anthropology: Issues, News, and Reviews: Issues, News, and Reviews*, *18*(5), 175-186.

Carvalho, S., Matsuzawa, T., & McGrew, W. C. (2013). From pounding to knapping: how chimpanzees can help us to model hominin lithics. In *Tool Use in Animals: Cognition and Ecology*, 225-241.

de Manuel, M., Kuhlwilm, M., Frandsen, P., Sousa, V. C., Desai, T., Prado-Martinez, J., Hernandez-Rodriguez, J., Dupanloup, I., Lao, O., Hallast, P., Schmidt, J. M., Heredia-Genestar, J. M., Benazzo, A., Barbujani, G., Peter, B. M., Kuderna, L. F. K., Casals, F., Angedakin, S., Arandjelovic, M., Boesch, C., Kühl, H., Vigilant, L., Langergraber, K., Novembre, J., Gut, M., Gut, I., Navarro, A., Carlsen, F., Andrés, A. M., Siegismund, H. R., Scally, A., Excoffier, L., Tyler-Smith, C., Castellano, S., Xue, Y., Christina Hvilsom, C., & Marques-Bonet, T. (2016). Chimpanzee genomic diversity reveals ancient admixture with bonobos. *Science*, *354*(6311), 477-481.

Derex, M., & Boyd, R. (2016). Partial connectivity increases cultural accumulation within groups. *Proceedings of the National Academy of Sciences*, *113*(11), 2982-2987.

Derex, M., Perreault, C., & Boyd, R. (2018). Divide and conquer: intermediate levels of population fragmentation maximize cultural accumulation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *373*(1743), 20170062.

Dutton, P., & Chapman, H. (2015). New tools suggest local variation in tool use by a montane community of the rare Nigeria–Cameroon chimpanzee, Pan troglodytes ellioti, in Nigeria. *Primates*, *56*, 89-100.

Dyble, M., Salali, G. D., Chaudhary, N., Page, A., Smith, D., Thompson, J., Vinicius, L., Mace, R., & Migliano, A. B. (2015). Sex equality can explain the unique social structure of hunter-gatherer bands. *Science*, *348*(6236), 796-798.

Dyble, M., Thompson, J., Smith, D., Salali, G. D., Chaudhary, N., Page, A. E., Vinicius, L., Mace, R., & Migliano, A. B. (2016). Networks of food sharing reveal the functional significance of multilevel sociality in two hunter-gatherer groups. *Current Biology*, *26*(15), 2017-2021.

Fontsere, C., Kuhlwilm, M., Morcillo-Suarez, C., Alvarez-Estape, M., Lester, J. D., Gratton, P., Schmidt, J. M., Dieguez, P., Aebischer, T., Álvarez-Varona, P., Agbor, A., Angedakin, S., Assumang, A. K., Ayimisin, E. A., Bailey, E., Barubiyo, D., Bessone, M., Carretero-Alonso, A., Chancellor, R., Cohen, H., Danquah, E., Deschner, T., Dunn, A., Dupain, J., E. Egbe, V. E., Feliu, O., Goedmakers, A., Granjon, A.-C., Head, J., Hedwig, D., Hermans, V., Hernandez-Aguilar, R., A., Imong, I., Jones, S., 4 Junker, J., Kadam, P., Kaiser, M., Kambere, M., Vyalengerera, M. K., Kalan, A. K., Kienast, I., Kujirakwinja, D., Langergraber, K. Lapuente, J., Larson, B., Laudisoit, A., Lee, K., Llana, M., Llorente, M., Marrocoli, S., Morgan, D., Mulindahabi, F., Murai, M., Neil, E., Nicholl, S., Nixon, S., Normand, E., Orbell, C., Ormsby, L.

J., Pacheco, L., Piel, A., Riera, L., Robbins, M. M., Rundus, A., Sanz, C., Sciaky, L., Sommer, V., Stewart, F. A., Tagg, N., Tédonzong, L. R., Ton, E., van Schijndel, J., Vergnes, V., Wessling, E. G., Willie, J., Wittig, R. M., Yuh, Y. G., Yurkiw, K., Zuberbuehler, K., Hecht, J., Vigilant, L., Boesch, C., Andrés, A. M., Hughes, D. A., Kühl, H. S., Lizano, E., Arandjelovic, M., & & Marques-Bonet, T. (2022). Population dynamics and genetic connectivity in recent chimpanzee history. *Cell Genomics*, *2*(6).

Granovetter, M. S. (1973). The strength of weak ties. *American journal of sociology*, *78*(6), 1360-1380.

Henrich, J. (2004). Demography and cultural evolution: how adaptive cultural processes can produce maladaptive losses—the Tasmanian case. *American antiquity*, *69*(2), 197-214.

Henrich, J. (2017). *The secret of our success: How culture is driving human evolution, domesticating our species, and making us smarter.* Princeton University press.

Hill, K. R., Wood, B. M., Baggio, J., Hurtado, A. M., & Boyd, R. T. (2014). Hunter-gatherer inter-band interaction rates: Implications for cumulative culture. *PloS one*, *9*(7), e102806.

Jesmer, B. R., Merkle, J. A., Goheen, J. R., Aikens, E. O., Beck, J. L., Courtemanch, A. B., Hurley, M. A., McWhirter, D. E., Miyasaki, H. M., Monteith, K. L., & Kauffman, M. J. (2018). Is ungulate migration culturally transmitted? Evidence of social learning from translocated animals. *Science*, *361*(6406), 1023-1025.

Kalan, A. K., Kulik, L., Arandjelovic, M., Boesch, C., Haas, F., Dieguez, P., Barratt, C. D., Abwe, E. E., Agbor, A., Angedakin, S., Aubert, F., Ayimisin, E. A., Bailey, E., Bessone, M., Brazzola, G., Buh, V. E., Chancellor, R., Cohen, H., Coupland, C., Curran, B., Danquah, E., Deschner, T., Dowd, D., Eno-Nku, M., Fay, M., Goedmakers, A., Granjon, A.-C., Head, J., Hedwig, D., Hermans, V., Jeffery, K. J., Jones, S., Junker, J., Kadam, P., Kambi, M., Kienast, I., Kujirakwinja, D., Langergraber, K. E., Lapuente, J., Larson, B., Lee, K. C., Leinert, V., Llana, M., Marrocoli, S., Meier, A. C., Morgan, B., Morgan, D., Neil, E., Nicholl, S., Normand, E., Ormsby, L. J., Pacheco, L., Piel, A., Preece, J., Robbins, M. M., Rundus, A., Sanz, C., Sommer, V., Stewart, F., Tagg, N., Tennie, C., Vergnes, V., Welsh, A., Wessling, E. G., Willie, J., Wittig, R. M., Yuh, Y. G., Zuberbühler, K., & Kühl, H. S. (2020). Environmental variability supports chimpanzee behavioural diversity. *Nature Communications*, *11*(1), 4451.

Kappeler, P. M., & van Schaik, C. P. (2002). Evolution of primate social systems. *International journal of primatology*, *23*, 707-740.

Kendal, R., Hopper, L. M., Whiten, A., Brosnan, S. F., Lambeth, S. P., Schapiro, S. J., & Hoppitt, W. (2015). Chimpanzees copy dominant and knowledgeable individuals: implications for cultural diversity. *Evolution and human behavior*, *36*(1), 65-72.

Kühl, H. S., Boesch, C., Kulik, L., Haas, F., Arandjelovic, M., Dieguez, P., Bocksberger, G., McElreath, M. B., Agbor, A., Angedakin, S., Ayimisin, E. A., Bailey, E., Barubiyo, D., Bessone, M., Brazzola, G., Chancellor, R., Cohen, H., Coupland, C., Danquah, E., Deschner, T., Dowd, D., Dunn, A., Egbe, V. E., Eshuis, H., Goedmakers, A., Granjon, A.-C., Head, J., Hedwig, D., Hermans, V., Imong, I., Jeffery, K. J., Jones, S., Junker, J., Kadam, P., Kambere, M., Kambi, M., Kienast1, I., Kujirakwinja, D., Langergraber, K. E., Lapuente, J., Larson, B., Lee, K., Leinert, V., Llana19, M., Maretti, G., Marrocoli, S., Martin, R., Mbi, T. J., Meier, A. C., Morgan, B., Morgan, D., Mulindahabi, F., Murai, M., Neil, E., Niyigaba, P., Ormsby, L. J., Orume, R., Pacheco, L., Piel, A., Preece, J., Regnaut, S., Rundus, A., Sanz, C., van Schijndel, J., Sommer, V., Stewart, F., Tagg, N., Vendras, E., Vergnes, V., Welsh, A., Wessling, E. G., Willie, J.,

Wittig, R. M., Yuh, Y. G., Yurkiw, K., Zuberbühler, K., & Kalan, A. K. (2019). Human impact erodes chimpanzee behavioral diversity. *Science*, *363*(6434), 1453-1455.

Laland, K. N. (2003). Gene-culture coevolution. *Encyclopaedia of Cognitive Science*, *2*, 268-274.

Lester, J. D., Vigilant, L., Gratton, P., McCarthy, M. S., Barratt, C. D., Dieguez, P., Agbor, A., Álvarez-Varona, P., Angedakin, S., Ayimisin, E. A., Bailey, E., Bessone, M., Brazzola, G., Chancellor, R., Cohen, H., Danquah, E., Deschner, T., Egbe, V. E., Eno-Nku, M., Goedmakers, A., Granjon, A.-C., Head, J., Hedwig, D., Hernandez-Aguilar, R. A., Jeffery, K. J., Jones, S., Junker, J., Kadam, P., Kaiser, M., K. Kalan, A. K., Kehoe, L., Kienast, I., Langergraber, K. E., Lapuente, J., Laudisoit, A., Lee,K., Sergio Marrocoli, S., Mihindou, V., Morgan, D., Muhanguzi, G., Neil, E., Nicholl, S., Orbell, C., Ormsby, L. J., Pacheco, L., Piel, A., Robbins, M., M., Rundus, A., Sanz, C., Sciaky, L., Siaka, A. M., Städele, V., Stewart, F., Tagg, N., Ton, E., van Schijndel, J., Vyalengerera, M. K., M., Wessling, E. G., Willie, J., Wittig, R. M., Yuh, Y. G., Yurkiw, K., Zuberbuehler, K., Boesch, B., Kühl, H. S., & Arandjelovic, M. (2021). Recent genetic connectivity and clinal variation in chimpanzees. *Communications Biology*, *4*(1), 283.

Lonsdorf, E. V., Eberly, L. E., & Pusey, A. E. (2004). Sex differences in learning in chimpanzees. Nature, 428(6984), 715-716.

Luncz, L. V., & Boesch, C. (2014). Tradition over trend: Neighboring chimpanzee communities maintain differences in cultural behavior despite frequent immigration of adult females. *American Journal of Primatology*, *76*(7), 649-657.

Lycett, S. J., Collard, M., & McGrew, W. C. (2010). Are behavioral differences among wild chimpanzee communities genetic or cultural? An assessment using tool-use data and phylogenetic methods. *American Journal of Physical Anthropology*, *142*(3), 461-467.

Mercader, J., Barton, H., Gillespie, J., Harris, J., Kuhn, S., Tyler, R., & Boesch, C. (2007). 4,300-year-old chimpanzee sites and the origins of percussive stone technology. *Proceedings of the National Academy of Sciences*, *104*(9), 3043-3048.

McGrew, W. C. (1992). *Chimpanzee material culture: implications for human evolution*. Cambridge University Press.

McGrew, W. C. (2010). Chimpanzee technology. *Science*, *328*(5978), 579-580.

McLennan, M. R. (2011). Tool-use to obtain honey by chimpanzees at Bulindi: new record from Uganda. *Primates*, *52*, 315-322.Dutton & Chapman 2015

Mesoudi, A. & Thornton, A. (2018). What is cumulative cultural evolution?. *Proceedings of the Royal Society B*, *285*(1880), 20180712.

Migliano, A. B., Battiston, F., Viguier, S., Page, A. E., Dyble, M., Schlaepfer, R., Smith, D., Astete, L., NGales, M., Gómez-Gardeñes, J., Latora, V., & Vinicius, L. (2020). Hunter-gatherer multilevel sociality accelerates cumulative cultural evolution. *Science advances*, *6*(9), eaax5913.

Migliano, A. B., Page, A. E., Gómez-Gardeñes, J., Salali, G. D., Viguier, S., Dyble, M., Thompson, J., Chaudhary, N., Smith, D., Strods, J., Mace, R., Thomas, M. G., Latora, V., & Vinicius, L. (2017). Characterization of hunter-gatherer networks and implications for cumulative culture. *Nature Human Behaviour*, *1*(2), 0043.

Migliano, A. B. & Vinicius, L. (2022). The origins of human cumulative culture: from the foraging niche to collective intelligence. *Philosophical Transactions of the Royal Society B*, *377*(1843), 20200317.

Nishida, T., Zamma, K., Matsusaka, T., Inaba, A., & McGrew, W. C. (2010). *Chimpanzee behavior in the wild: an audio-visual encyclopedia*. Springer Science & Business Media.

Padilla-Iglesias, C., Atmore, L. M., Olivero, J., Lupo, K., Manica, A., Arango Isaza, E., Vinicius, L., & Migliano, A. B. (2022a). Population interconnectivity over the past 120,000 years explains distribution and diversity of Central African hunter-gatherers. *Proceedings of the National Academy of Sciences*, *119*(21), e2113936119.

Padilla-Iglesias, C., Blanco-Portillo, J., Ioannidis, A., Manica, A., Vinicius, L., & Migliano, A. (2022b). Cultural Evolution of Central African hunter-gatherers reflects a deep history of interconnectivity. *Preprint (Version 1)*, https://doi.org/10.21203/rs.3.rs-2205369/v1.

Page, A. E., Chaudhary, N., Viguier, S., Dyble, M., Thompson, J., Smith, D., Salali, G. D., Mace, R., & Migliano, A. B. (2017). Hunter-gatherer social networks and reproductive success. *Scientific Reports*, *7*(1), 1153.

Powell, A., Shennan, S., & Thomas, M. G. (2009). Late Pleistocene demography and the appearance of modern human behavior. *Science*, *324*(5932), 1298-1301.

Pradhan, G. R., Tennie, C., & van Schaik, C. P. (2012). Social organization and the evolution of cumulative technology in apes and hominins. *Journal of human evolution*, *63*(1), 180-190.

Pusey, A. E. & Packer, C. (1986). Dispersal and philopatry. In *Primate Societies, eds BB Smuts, DL Cheney, RM Seyfarth, RW Wrangham, TT Struhsaker*. Univ Chicago Press pp 250-266.

Rawlings, B. S., Legare, C. H., Brosnan, S. F., & Vale, G. L. (2021). Leveling the playing field in studying cumulative cultural evolution: Conceptual and methodological advances in nonhuman animal research. *Journal of Experimental Psychology: Animal Learning and Cognition*, *47*(3), 252.

Sanz, C., Call, J., & Morgan, D. (2009). Design complexity in termite-fishing tools of chimpanzees (Pan troglodytes). *Biology Letters*, *5*(3), 293-296.Boesch 2012

Scerri E. M. L., Chikhi L., & Thomas M. G. (2019). Beyond multiregional and simple out-of-Africa models of human evolution. Nat. Ecol. Evol.3, 1370–1372. (doi:10.1038/s41559-019-0992-1)

Schuppli, C. & Van Schaik, C. P. (2019). Animal cultures: how we've only seen the tip of the iceberg. *Evolutionary Human Sciences*, *1*, e2.

Tennie, C., Call, J., & Tomasello, M. (2009). Ratcheting up the ratchet: on the evolution of cumulative culture. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *364*(1528), 2405-2415.

Tennie, C., Hopper, L. M., & van Schaik, C. P. (2020). On the origin of cumulative culture: Consideration of the role of copying in culture-dependent traits and a reappraisal of the zone of latent solutions hypothesis. In L. M. Hopper & S. R. Ross (Eds.), *Chimpanzees in context: A comparative perspective on chimpanzee behavior, cognition, conservation, and welfare* (pp. 428–453). The University of Chicago Press.

Tomasello, M., Kruger, A. C., & Ratner, H. H. (1993). Cultural learning. *Behavioral and brain*
sciences, *16*(3), 495-511.

van Schaik, C. P., Pradhan, G. R., & Tennie, C. (2019). Teaching and curiosity: sequential
drivers of cumulative cultural evolution in the hominin lineage. *Behavioral ecology and*
sociobiology, *73*, 1-11.

Whiten, A. (2021). The burgeoning reach of animal culture. *Science*, *372*(6537), eabe6514.

Whiten, A. (2023). Social learning and culture in the great apes. In J. Tehrani, R. Kendal, J.
Kendal (Eds.) *Oxford Handbook of Cultural Evolution*. pp C23S1–C23S31. Oxford: Oxford
University Press.

Whiten, A., Goodall, J., McGrew, W. C., Nishida, T., Reynolds, V., Sugiyama, Y., Tutin, C. E.
G., Wrangham, R. W., Boesch, C. (1999). Cultures in chimpanzees. *Nature, 399*, 682–685.
doi:10.1038/21415

Whiten, A., Goodall, J., McGrew, W. C., Nishida, T., Reynolds, V., Sugiyama, Y., Tutin, C. E.
G., Wrangham, R. W., Boesch, C. (2001). Charting cultural variation in chimpanzees.
Behaviour, *138*(11-12), 1481-1516. https://doi.org/10.1163/156853901317367717

Whiten, A. & van Schaik, C. P. (2007). The evolution of animal 'cultures' and social
intelligence. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *362*(1480),
603-620.

Supplementary Information

Materials and Methods

Cultural data

To investigate cultural similarities between chimpanzee populations, we used a behavioural

database compiled from 144 chimpanzee communities in two previous studies (Kühl et al., 2019,

Kalan et al., 2020). The data for 106 communities was compiled from the published literature

and extended with data for 46 communities from the Pan African Program: the Cultured

Chimpanzee (for further details see Kühl et al., 2019, SI; and Kalan et al., 2020, Methods), and

eight of in the published literature. We extracted communities from the 35 sites that were also in

the genetic database and lumped communities at the same site together to match the genetic data.

To control for potential ecological factors, we further extracted data on mid-term environmental

variability from Kalan et al. (2020), who classified communities as predominantly savannah

woodland or forest (Kalan et al., 2020, Methods), as well as compiled information on

observation time in months at each site (population-average across multiple communities).

We selected foraging behaviours to have a comparative set of behaviours across different

complexity levels, and only included behaviours present in at least two populations to compare

shared knowledge (Table S1). We stratified behaviours into three groups according to the

number of material parts: non-tool, simple (single tool) and complex (multi-part tool). To

estimate cultural similarity, we further resolved tool use behaviours according to the type of

tools. This was specifically the case for honey foraging, where tools involved to extract above-

ground or underground substrates differed in the type of tools and in the process of extraction.

We divided both the simple honey tool as well as the complex honey toolset into above-ground

and underground by screening the published literature from the respective populations. We

repeated this procedure for the termite toolset; only one of the 35 populations (Goualougo,

Central) had evidence for an above-ground termite toolset (Sanz & Morgan 2007) and was

therefore excluded. Due to the similar process of underground extraction using a multipart

toolset for honey and termites (Boesch, 2012), we grouped these into one behavioural category to

increase statistical power.

Following Kühl et al. (2019) and Kalan et al., (2020) we coded presence of each behaviour

(1/0) in all 35 populations, so that direct or indirect observations of a behaviour at a site were

coded as present (1) and otherwise as not observed (0). We excluded cases that were coded as

'not applicable' by Kalan et al. (2020). We determined 15 foraging behaviours and classified

three as non-tool, nine as simple and three as complex. For statistical analysis of shared

knowledge of behaviours, we created a database comprising all possible pairs of the 35

populations. For each behaviour we coded shared occurrence (1/0) in all pairs of populations. We

coded a behaviour as shared (1) if it was present in both populations of a pair. If the behaviour

was only present in one or none of the populations, we coded not shared (0).

567
568   <u>Genetic connectivity data</u>
569
We applied two genetic markers to analyse the effect of recent migration on shared

knowledge. We used a genetic database from chromosome 21 data from Fontsere et al. (2022),

comprising of non-invasive faecal and shed hair samples from 828 individuals at 48 sampling

sites across all four chimpanzee subspecies as part of the PanAf Program. Coming from non-

invasive samples, only a subset of 240 individuals had high-coverage genotypes (at least 5-fold

coverage on average), while the data is patchy and of low coverage for most individuals, as

described by Fontsere et al., (2020).

To assess recent connectivity we used 'identical-by-descent'-like (IBDs) (Fontsere et al.,

2022; Browning & Browning, 2013; Thompson, 2013; Padilla-Iglesias et al., preprint) reflecting

a recent common ancestor between two individuals (see details of data processing and analysis in

Fontsere et al., 2022). For individuals from different sites this implies a recent migration event

within the past ~50 generations. The data only included samples with >5-fold coverage to

increase the power to detect IBD-like segments in the limited dataset, resulting in data from 34

sites that match the cultural data (no IBD data from Outamba-Kilimi). We performed all analyses

on IBD with these 34 populations. Due to exponential decay of IBD segments through

recombination, the length of shared IBD segments is correlated with the time of the migration

event, with longer IBDs representing more recent genetic exchange. Therefore, to assess the time

frame of the MRCA between two populations, we extracted the highest IBD length shared

between each pair of individuals of a population-pair and calculated the average length of this for

each pair of populations (henceforth IBD length). We refrained from calculating the number of

IBDs shared between a population-pair even though this could serve as a measure of connectivity

to avoid potential biases produced by filtering only samples with >5-fold coverage. The data

came from two IBD analysis events, with the second including only a subset of the Western

populations to control for differences in sampling density (for details, see Fontsere et al., 2022,

see SI). We calculated IBD length for only the first data set and both data sets combined (Fig.

S1). Since the IBD length from both data sets did not differ much and the combined dataset was

able to identify four additional IBD connections, we continued analysis using the combined

dataset.

To support the analysis of connectivity between populations and increase the time frame of

potential genetic and cultural exchange, we further used nearly private rare alleles (NePRAs)

(Fontsere et al., 2022; Schiffels et al., 2016) which represent potentially earlier migration from

one population to another (1.5 kya – 15 kya). We used data from Fontsere et al. based on

963,656 SNPs (on average 26,671 per site) obtained from 434 individuals (on average 11 per

site) with at least 1-fold coverage and 38 sampling sites, permissive for missing data. These

SNPs were classified as NePRAs if the allele frequency at a site was larger than zero and the

cumulative frequency of this allele was less than one at all other sites. We used a matrix from 35

sites that match the cultural data including binary data (1/0) for each SNP. If a NePRA was

detected in at least one sample at a site, the SNP was coded as present (1). If there were no cases

of this allele at the site, it was coded as not detected (0). In some cases, a SNP was not sequenced

in any of the samples at a given site which was coded as NA. To assess similarity between two

populations, we extracted the SNPs that were sequenced in both populations of each pair and

calculated the proportion of shared NePRAs. This gives a measure of genetic similarity between

populations and potential introgression over the past several thousand years. One limitation of

the definition of NePRAs is that populations that have only recently undergone separation, for

example in the case of Western chimpanzees that might have emerged from a recent population

bottleneck (de Manuel et al., 2016; Fontsere et al., 2022), have fewer NePRAs (rare allele

variants are at high frequencies in multiple populations) than populations that have diversified

earlier and where mutations had more time to occur and spread to other populations. Therefore,

genetically close populations might express lower levels of NePRA similarity. Due to

probabilistic mutation rate, NePRA similarity representing a very old common ancestor cannot

be excluded and is considered as noise. After visual inspection of the distribution of NePRA

similarity, we determined a shift in scale at a threshold of NePRA similarity at 0.0006 and

classified everything below this threshold as noise (Fig. S2).

Statistical analysis

To analyse the effect of genetic connectivity on shared knowledge of different behavioural

traits, we followed the methods of Kalan et al. (2020) by applying Bayesian Regression Models

(BRMs) with Bernoulli response distribution and logit link function. We conducted all analyses

in R (R Core Team (2022), version 4.2.1), and implemented the code in RStudio (RStudio Team

(2023), version 2023.3.1.446). We built separate models for non-tool, simple, and complex

behaviours, implementing shared occurrence (1/0) of each behaviour in the three complexity

categories as response variable.

We fitted an individual model for each of the genetic predictors since NePRA similarity and

IBD length were highly correlated (Pearson's r = 0.478, p-value < 0.001). To analyse whether

any exchange of NePRAs or IBD segments could predict shared culture, we fit separate models

for binary IBD-exchange (1 = sharing of at least one IBD segment, 0 = sharing no IBD

segments) between each population-pair as well as binary NePRA exchange (1/0); NePRA

similarity above 0.0006 was classified as potential exchange (1) and below was considered as

noise and classified as no exchange (0). To test whether the amount of contact could predict

sharing of behaviours, we included the continuous variable of NePRA similarity. Since detection

of IBD segments was highly sensitive to low-coverage data, we tested the population-average of

the maximum IBD length shared between each pair of individuals of a population-pair to analyse

the most recent contact. Due to a high correlation with genetic predictors, subspecies (Spearman

rho between NePRA similarity and subspecies = 0.660, p-value < 0.001, Spearman rho between

IBD length and subspecies = 0.0.80, p-value < 0.001) and geographic distance (Pearson r

between NePRA similarity and distance = -0.497, p-value < 0.001, Pearson r between IBD

length and distance = -0.476, p-value < 0.001), we performed two separate models with either

same subspecies (binary, 1 = same subspecies, 0 = different subspecies) or geographic distance

in km calculated from coordinates at the centre of the sites (Fontsere et al., 2022; Kalan et al.,

2020; Kühl et al., 2019)) between a pair of populations (Fig. S5). To control for variation in

observation time at different sites, we extracted the shorter observation time in months among

the population-pair and included it as a fixed effect (shared observation time). Additionally, we

included the binary variable of same habitat (1 if both populations in a pair have the same habitat

/ 0 if the habitat from the two populations are different). We added the population-pair and

behaviour as random effects (Schielzeth, 2010). Due to the small sample size, we did not include

random slope effects. However we did estimate random slopes of behaviour on the genetic or

geographic predictor, observation time and habitat (Fig. S3). After inspection of the distribution

of the continuous predictors, we log-transformed geographic distance and NePRA similarity and

square-root-transformed IBD length and z-transformed all three ($\beta \sim N(0,1)$) (Schielzeth 2010).

Following Kalan et al., we also z-transformed shared observation time.

We used the function *brm* from the R-package *brms* (version 2.18.0) (Bürkner 2017) to fit

all BRMs. By default, we run 2000 iterations including 1000 iterations for warm-up over four

MCMC chains, creating 8000 usable posterior samples. Model diagnosis of each model showed

that there were no divergent transitions after warm-up; all MCMC chains were stationary and

converged and all Rhat values were below 1.01 (Gelmann et al. 2014). We used weak priors with

a standard normal distribution with a mean of 0 and a standard deviation of 1. To assess to effect

of the priors, we also tested a wide prior ($\beta \sim N(0,10)$) (Fig. S4). We reported the odds ratios of

the mean estimate to compare the effect sizes of the predictors on each of the samples, and

assessed the significance of the results using the credible intervals at 65%, 85% and 95% (mean

estimate of the marginal posterior distribution, credible intervals around the mean estimate of the

posterior distribution) (Fig. 2, table S2). We tested the association between IBD and shared

culture using a combined dataset from two IBD sequencing events (the first including all

populations and the second using a subset of the first data); however, we checked how the first

dataset performed individually (Fig. S6).

We applied Mantel tests to assess correlations between cultural, geographical, ecological

and genetic distances. We calculated Jaccard distances between population-pairs for non-tool,

simple and complex using the *vegdist* function from the *vegan* R-package (version 2.6.2)

(Oksanen et al. 2022) and estimated behavioural similarity between population-pairs for each of

the three samples by subtracting the Jaccard distance from 1 for more intuitive analysis of the

results. We further applied the NePRA similarity and IBD length matrices as measures of genetic

similarity, and geographic proximity by dividing 1 by distance in km. Additionally, we used a

matrix of same habitat (1/0) as well as same subspecies (1/0) and shared observation time

(continuously increasing). We performed Mantel tests using the *ecodist* R-package (version

2.0.9) (Goslee et al. 2007) with the Spearman correlation coefficient between each pair of

similarity/proximity matrices being calculated from 10,000 resamples. We performed simple

Mantel tests in which each pair of matrices was tested (Fig. S7A), as well as partial Mantel tests

controlling for same habitat (Fig. S7B) as well as shared observation time (Fig. S7C). Further,

we performed a separate simple Mantel test correlating each complex behaviour with either IBD

length or NePRA similarity (Fig. S7D). When interpreting these result, the low sample size

within each complex behavioural category and the following weak statistical power of these

results must be considered and compared with other analyses.

Social network analysis

To investigate the relationship between the cultural and genetic transfers resulting from

between-group migration, we applied reinforcement analysis. The technique attempts to

determine the directionality of effects between two networks with similar nodes (in our case,

populations) but different links, by reciprocally predicting the presence of a link in a network

from the strength of the same link in the other. As an example, let us imagine a 'marriage'

network M where a link represents a married couple, and a 'lifespan' network L with the same nodes, but where a link is present if the two individuals were alive at the same time. In this case, links in the marriage network will always predict the presence of the same link in the lifespan network, since being married is conditioned upon the two individuals being contemporaneous. Reversely, links in the lifespan network will predict marriage links with much lower probability, since being contemporaneous is not conditioned upon being married, or alternatively, since many factors other than being alive at the same time are needed for marriage to happen. The rationale applies to networks characterised by links categorised at distinct levels. For example, assume a network C only including married couples with two link levels; no children, or with children. A second network T includes links representing time of marriage; less than six months, or over six months. We expect a dramatic increase in the proportion of 'over six months' links predicted by 'with children' links (since pregnancy length is nine months). Reversely, the increase in the proportion of predicted 'with children' links by 'over six months' links is expected to be less pronounced, since there are many childless couples married for longer than six months (due to factors other than duration of marriage behind the decision to have children).

We created an individual network of shared knowledge for each behavioural type, where a link represented sharing the respective behaviour. Next, we divided the IBD network into sharing-IBD and no-sharing, and the NePRA network into three levels of genetic similarity. The first bin (weak links, N = 322) contained all links that had a NePRA similarity < 0.0006 (below our defined threshold for noise). The second and third bin were divided at 0.002, creating two almost equally sized bins, with the second bin containing links of NePRA similarity between 0.0006 and 0.002 (medium links, N = 137), and the third bin comprising all links with a similarity larger than 0.002 (strong links, N = 136). The first bin in the analysis involving IBD length contained population-pairs that did not share any IBDs (no sharing, N = 428), while the

second bin contained links that shared at least one IBD (IBD sharing, N = 133). To analyse the

directionality of the association, we calculated the probability of links in each genetic

subnetwork that also displayed a link in each of the behavioural networks. To test for reverse

directionality, we calculated the proportion of links in each bin of the two genetic networks for

each of the behavioural network, and reported averaged results across behaviours for non-tool,

simple and complex (Fig. 3A-D). Long IBD segments (>0.5 mega base pairs (Mbp)) may provide

evidence for recent connectivity (less than 5 kya) whereas shorter IBD segments (<0.5 Mbp)

have been subjected to exponential decay of IBD length and therefore project older exchange

(more than 5 kya). Therefore, we separated IBDs into bins of no IBD sharing, short IBD (IBD

length <0.5 Mbp) and long IBD (>0.5 Mbp) and repeated reinforcement analysis for each

complex trait using these bins (Fig. S8).

Visual inspection of distribution of cultural traits

To examine how various forms of a technology may relate to each other and with patterns

of interconnectivity, we summarised all simple and complex instances of underground foraging

in the Central, Nigeria-Cameroon and Eastern regions using all 107 populations from the

compiled database from Kühl et al. and Kalan et al. (Fig. S9). We divided the behaviours

according to the resources extracted (honey, termites and ants) as well as the different steps by

extending the database by screening previously published data (Fig. 4, Table S3). We classified

the increase in complexity of a variant of the behaviour at a site into incremental change (adding

an additional tool) and repurposing (using the toolset for a new substrate) (Table S3). We

analysed the genetic links (IBD: Fig. 4A: NePRA, Fig. 4B) between the respective populations

for which we had genetic data and identified the changes in complexity between these

populations.

References

1. Kühl, H. S., Boesch, C., Kulik, L., et al. (2019). Human impact erodes chimpanzee behavioral diversity. Science, 363(6434), 1453-1455.

2. Kalan, A. K., Kulik, L., Arandjelovic, M., et al. (2020). Environmental variability supports chimpanzee behavioural diversity. Nature Communications, 11(1), 4451.

3. Sanz, C. M., & Morgan, D. B. (2007). Chimpanzee tool technology in the Goualougo Triangle, Republic of Congo. Journal of Human Evolution, 52(4), 420-433.

4. Boesch, C. (2012). Wild cultures: a comparison between chimpanzee and human cultures. Cambridge University Press.

5. Fontsere, C., Kuhlwilm, M., Morcillo-Suarez, C et al. (2022). Population dynamics and genetic connectivity in recent chimpanzee history. Cell Genomics, 2(6).

6. Thompson, E. A. (2013). Identity by descent: variation in meiosis, across genomes, and in populations. Genetics, 194(2), 301-326.

7. Browning, B. L., & Browning, S. R. (2013). Detecting identity by descent and estimating genotype error rates in sequence data. The American Journal of Human Genetics, 93(5), 840-851.

8. Schiffels, S., Haak, W., Paajanen, P., et al. (2016). Iron age and Anglo-Saxon genomes from East England reveal British migration history. Nature communications, 7(1), 10408.

9. R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

10. Posit team (2023). RStudio: Integrated Development Environment for R. Posit Software, PBC, Boston, MA. URL http://www.posit.co/.

11. Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients. Methods in Ecology and Evolution, 1(2), 103-113.

12. Bürkner, P.-C. (2017). Brms: an R package for bayesian multilevel models using stan. J. Stat. Softw. 80, 1–28.

13. Gelman, A., Carlin, J. B., Stern, H. S., et al. (2014). Bayesian data analysis 3rd edition. CRC Press.

14. Oksanen F. J., Simpson, G., Blanchet F., et al. (2022). Vegan: Community Ecology Package. R package Version 2.4-3. https://CRAN.R-project.org/package=vegan

15. Goslee, S. C. & Urban, D. L. (2007). The ecodist package for dissimilarity-based analysis of ecological data. J. Stat. Softw. 22, 1–19.

16. Csardi G, Nepusz T (2006). The igraph software package for complex network research. InterJournal, Complex Systems 1695. https://igraph.org

Fig. S1. Shared IBD length for 133 population-pairs that share at least one IBD. Blue shows IBD length from the main data set from Fontsere et al. 2022 including all 34 populations. Red shows the IBD length from two sequencing events combined. The stars show four population-pairs where IBD was only detected in the second analysis. This is the case for the following population-pairs (subspecies in brackets): 1. Goualougo (C) - La Belgique (C), 2. Comoe (W) - Mt. Sangbe (W), 3. Kabogo (E) - Ngogo (E), 4. Nyungwe (E) – Rubi-Télé (E). A Wilcoxon-Rank-Sum test showed no meaningful difference between the two data sets (P-value = 0.58).

Fig. S2. Distribution of NePRA similarity. The red dashed line indicates the noise-threshold at 0.0006. The maximum NePRA similarity is at 0.021 between Conkouati (Central) and Loango (Central).

Fig. S3. Analysis of random slope effects of behaviour. As expected, credible intervals are wider when including random slope effects but the mean estimate does not differ largely. The plot shows the mean of the marginal posterior distribution (dots) and the 67%, 87%, 97% credible intervals centered on the mean (line thickness corresponds decreasingly with size of credible intervals).

Fig. S4. Analysis of prior sensitivity. Interestingly, only the models testing complex tools seem to vary when implementing a wide prior (N(0,10)) compared to a weak prior (N(0,1)), specifically for the NePRA binary predictor where the distribution is wider. Still the mean estimates differ minimally. The plot shows the mean of the marginal posterior distribution (dots) and the 67%, 87%, 97% credible intervals centered on the mean (line thickness corresponds decreasingly with size of credible intervals).

Fig. S5. Bayesian logistic regression of effects of geographic distance and same region (1/0) on shared cultural traits between chimpanzee populations. The plot shows the mean of the marginal posterior distribution (dots) and the 67%, 87%, 97% credible intervals centered on the mean (line thickness corresponds decreasingly with size of credible intervals).

Fig. S6. Analysis of robustness of IBD data. As expected, there is no clear difference in results between the IBD data set 1 and the combined data set. The plot shows the mean of the marginal posterior distribution (dots) and the 67%, 87%, 97% credible intervals centered on the mean (line thickness corresponds decreasingly with size of credible intervals).

Fig. S7. Mantel correlation analysis with 10,000 resamples displaying Spearman correlation coefficients r as the size of the pies and significance showed through the stars (* = 0.05>*P*>0.01, ** 0.01>*P*>0.001, *** P<0.001) testing the null hypothesis that there is no correlation between the respective matrices. Panel A shows correlations from simple Mantel tests which show that while NePRA similarity, IBD length, (geographic) proximity and region (subspecies) are highly correlated, all are correlated only with complex tools, apart from NePRA similarity which also has a slight correlation with simple tools. These results correspond to the fits of the Bayesian regressions. Further, partial Mantel tests showed that controlling for habitat (B) or observation time (C) provide similar results. Panel D shows the results from the Mantel tests for each complex trait.

Fig. S8. Reinforcement analysis of short vs. long IBD segments shared in each of the complex
cultural networks. Results show that the proportion of links in the nut cracking and honey toolset
network that share IBD are all at least 0.5 Mbp long, whereas the underground toolset has a
higher proportion of short IBD links, suggesting that the underground toolset might have spread
earlier.

839
840
841

Fig. S9. Distribution of the 15 behaviours in all 107 populations from Kalan et al. 2020.
Communities (N = 144) at the same site were grouped together into one population.

Table S1. Definitions of the 15 foraging behaviours used in this study and described by Kühl et al. 2019. New definitions in this study (highlighted): Honey above-ground, Honey underground, Honey toolset and Underground toolset.

	Behaviour	Definition
Non-tool	Ant eating	eat ants with bare hands
	Honey eating	eat honey with bare hands
	Termite eating	use hands to break open termite mounds; no tooluse seen
Simple tool use	Algae fishing	fish algae with plant part from bottom of water source
	Ant dipping	collect army ants (Dorylus spp.) from trail or nest using plant part
	Ant fishing	use plant part to obtain non-Dorylus spp. ants
	Honey above-ground	extract honey from above-ground beehives using a plant part (e.g. dip or pound)
	Honey underground	extract honey from subterranean beehive using a digging tool
	Leaf sponge	use chewed wad of leaves as a sponge to soak up water and squeeze into mouth
	Marrow pick	pick brain matter or bone marrow using plant part
	Termite fishing	extract termites from above-ground mound using plant part
	Water dipping	chew on plant part to create a brush tip to soak-up water and ingest
Complex toolset	Honey toolset	extract honey from above-ground beehives using functionally different types of tools sequentially from the following tool types: pounding, lever, dipping, swabbing (REFs)
	Nut cracking	crack nuts using hammer and anvil from wood or stone
	Underground toolset	extract termites or honey from subterranean nest / beehive using a digging tool followed by a dipping tool

Table S2. Results of Bayesian Regression Models for non-tool, simple and complex behaviours. Shared knowledge included as response variable, using weak priors and testing genetic and geographic predictors in separate models. Estimate of the mean marginal posterior distribution, standard deviation of the estimate (sd), the 2.5% and 97.5% credible intervals centered on the mean (CI), proportion of the posterior distribution greater than zero and odds ratios of the mean estimate (OR) are reported. *Average across all six models per response category.

	Estimate	sd	CI 2.5%	CI 97.5%	Post. Dist.>0	OR
Non-tool behaviours (n = 35)						
Intercept*	-1.831	1.223	-4.438	0.441	0.06	0.16
NePRA binary	0.009	0.401	-0.767	0.800	0.51	1.009
NePRA similarity	-0.074	0.208	-0.469	0.345	0.357	0.929
IBD binary	-0.388	0.452	-1.309	0.491	0.197	0.678
IBD length	-0.309	0.266	-0.860	0.174	0.119	0.734
Distance	0.241	0.233	-0.202	0.727	0.856	1.273
Same subspecies	0.044	0.419	-0.801	0.833	0.547	1.045
Same habitat*	0.535	0.406	-0.260	1.341	0.907	1.708
Shared observation time (months)*	0.435	0.113	0.224	0.675	1.00	1.544
Simple behaviours (n = 35)						
Intercept*	-2.765	0.785	-4.08	-1.016	0.002	0.063
NePRA binary	0.529	0.163	0.211	0.848	1.00	1.697
NePRA similarity	0.368	0.083	0.206	0.533	1.00	1.445
IBD binary	0.321	0.18	-0.043	0.675	0.96	1.379
IBD length	0.142	0.076	-0.010	0.286	0.967	1.153
Distance	-0.103	0.085	-0.267	0.065	0.113	0.902
Same subspecies	0.022	0.178	-0.326	0.372	0.549	1.022
Same habitat	0.004	0.164	-0.318	0.325	0.509	1.005
Shared observation time (months)*	0.303	0.62	0.183	0.26	1.00	1.354
Complex behaviours (n = 35)						
Intercept*	-2.602	1.141	-4.91	-0.41	0.013	0.079
NePRA binary	1.739	0.537	0.762	2.842	1.00	5.691
NePRA similarity	1.148	0.304	0.567	1.744	1.00	3.152
IBD binary	0.943	0.401	0.137	1.735	0.989	2.568
IBD length	0.333	0.168	0.004	0.65	0.976	1.395
Distance	-0.91	0.209	-1.344	-0.521	0.00	0.403
Same subspecies	1.324	0.401	0.562	2.113	1.00	3.758
Same habitat	1.907	0.589	0.828	3.137	1.00	6.79
Shared observation time (months)*	0.352	0.113	0.124	0.567	0.997	1.424

Table S3. Description of type of underground foraging behaviours at 15 chimpanzee populations. Honey digging tool: Underground honey, beehive or bee larvae extraction using a digging tool to enlarge the subterranean beehive entrance; Honey toolset: Digging for a beehive followed by the use of a dipping tool to collect honey; Termite toolset: Use of a digging tool to perforate the ground and create a hole to the subterranean termite mound followed by a fishing tool to collect termites; Ant toolset: Digging tool and dipping tool found at the same subterranean ant nest. The changes related to cumulative culture from the simple variant are described (underground honey digging tool): Incremental change = increase in number of tools involved, Repurposing = change in resource type exploited. The reporting literature was cited.

Population	Subspecies	Tool types	Changes from simple variant	Reference
Bili-Uele	East	Honey digging tool	None	Hicks et al., 2019
Bulindi	East	Honey digging tool followed by probing tool	Incremental change	McLennan, 2011, 2015
Kahuzi-Biega	East	Honey digging tool	None	Yamagiwa et al., 1988
Nyungwe	East	Honey digging tool	None	Matthews et al., 2019
Gashaka	Nigeria-Cameroon	Honey toolset	Incremental change	Sommer et al., 2012
Ngel Nyaki	Nigeria-Caeroon	Honey digging tool Honey dipping tool Ant toolset	None Other tool Incremental change Repurposing	Dutton & Chapman, 2015
Korup	Nigeria-Cameroon	Termite toolset	Incremental change Repurposing	Boesch et al., 2020
Bai Hokou	Central	Honey digging tool	None	Fay & Caroll, 1994
Campo Ma'an	Central	Termite toolset	Incremental change Repurposing	Boesch et al., 2020
Goualougo	Central	Termite toolset	Incremental change Repurposing	Sanz & Morgan, 2007
La Belgique	Central	Honey digging tool Termite toolset	None Incremental change Repurposing	Deblauwe et al, 2006
Loango	Central	Honey toolset	Incremental change	Boesch et al., 2009
Mt. Cristal	Central	Termite toolset	Incremental change Repurposing	Boesch et al., 2020
Ngotto	Central	Honey digging tool	None	Hicks et al., 2005
Wonga-Wongue	Central	Termite toolset	Incremental change Repurposing	Boesch et al., 2020