1

The Pvc15 D2-Pnf SP Interaction Mediates AAA ATPase Activity,

Payload Stability, and Translocation into the PVC Tube Lumen.

Rhys Evans[1] and Nicholas R. Waterfield[2]*.

[1] School of Life Science, University of Warwick, Coventry, United Kingdom

[2] Warwick Medical School, University of Warwick, Coventry, United Kingdom

* Corresponding author.

10
Summary

This manuscript concerns elucidation of the mechanism by which the N-

terminal leader peptide sequence of the effector proteins of the *Photorhabdus*

PVC toxin delivery system, are targeted for loading into the lumen of the

nanosyringe structure, The Pvc15 AAA-ATPase is shown to be necessary for

this process and the interaction of leader sequences is investigated.

Abstract

The *Photorhabdus* virulence cassette (PVC) is an elegant, multi-protein,

contractile nanostructure which injects functionally bioactive polypeptides into

the eukaryotic cytosol. Signal peptides (SPs) are N-terminal amino acid motifs

from native 'payload' proteins which can also associate wide range of

heterologous proteins to the PVC tube lumen. In addition, Pvc15 is a classic

AAA ATPase encoded within the PVC operon which encodes an N-domain and

tandem AAA domains D1 and D2. This work finds that Pvc15's ATPase

activity is mediated by the E555 residue situated in the Walker B motif of the

sole functional ATPase domain, D2. Pvc15 multimerisation may be required to

translocate payload into the PVC tube lumen, which is facilitated by intact N

and D1 domains. In addition, ATPase activity requires the presence of other

PVC operon components which hints at a PVC-regulated mode of action for

loading. Furthermore, D2 is a chaperone for the native Pnf SP, though C-

terminal truncations of the Pnf SP confers Pvc15-independent stability to the

payload. These findings provide insight into the nuanced roles of the Pvc15-SP

interaction for payload-PVC association and the effects that SP length and type

has on payload stability and PVC-loading ability.

Introduction

The *Photorhabdus* virulence cassette (PVC) is one classic example of a class of bacterial bioactive protein delivery devices collectively termed extracellular contractile injection systems, or eCIS (Yang *et al.*, 2006; Hapeshi and Waterfield, 2017; Chen *et al.*, 2019). Devices of this nature are often large (>10MDa) multiprotein phage-like nanosyringes consisting of a rigid, hollow tube whose distal spike is used to puncture into the cell membrane of the target host cell, after contraction of the surrounding interlinked sheath proteins, to deliver a toxic effector 'cargo' or 'payload' (Kube and Wendler, 2015; Desfosses *et al.*, 2019; Jiang *et al.*, 2019). *Photorhabdus* species encode 5-6 PVC operons in their genomes, each differing only slightly in the conservation of the PVC core structural components (Yang *et al.*, 2006). Operons differ more greatly by the effector cargoes that are encoded up to ~6kb downstream of their respective PVC operons and which give namesake to the operon itself. Cargo loading is mediated, in part, by the presence of a 50-70 amino acid N-terminal "signal peptide" (SP) which is effective for loading even large heterologous proteins such as β-lactamase, mRFP, and *Renilla* luciferase into the 150-200nm-long tube lumen (Jiang *et al.*, 2022; Wang *et al.*, 2021).

The modular nature of PVC payloads to be associated with the PVC offers an exciting potential for these devices to be bioengineered for the delivery of therapeutic proteins into the cytosol (reviewed by Lee *et al.*, 2019). For example, the distal C-terminal domain of Pvc13 tail fibres can be replaced with epitope tags or adenovirus binding domains to retarget wild-type PVCs to various cell types (Kreitz *et al.*, 2023).

How the PVC-specific SPs target the cargo to the PVC for loading remains a mystery. Pvc15, an <u>A</u>TPase <u>a</u>ssociated with diverse cellular <u>a</u>ctivities (a AAA+ ATPase), is required for cargo loading and for the unfolding of SP-associated mRFP (Jiang *et al.*, 2022). In addition, the *Serratia entomophila* homologue,

anti-feeding prophage (Afp) are only produced when co-expressed with Afp15,

whereas Δ*pvc15* PVCs are intact but lack their associated cargo (Bhardwaj,

Mitra and Hurst, 2021).

To understand the relevance of SPs to eCIS loading, this work presents a

comprehensive analysis of eCIS effector diversity in lineage Ia and Ib operons,

and protein prediction is used to identify the ubiquity of SPs in these ORFs.

Protein structure prediction is also used to infer the function of Pvc15 domains

in its monomeric and potential hexameric assemblies (Snider, Thibault and

Houry, 2008). In addition, analysis of Western blot band intensities is used to

infer the effects of SP type (i.e., Pnf50 and Cif50) and length on cargo stability

and PVC loading when co-expressed with Pvc15 in *E. coli*. Finally, Pvc15

mutants are constructed and expressed *in trans* to the PVC*pnf*Δ*pvc15* operon

where the ATPase activity is then characterised to infer the importance of the

identified domains and residues to Pvc15's various roles.

Results

eCIS lineage analysis reveals diverse putative effector functionality.

To assess the diversity of functions of effectors at eCIS operons, the eCIS
database (dbeCIS; http://www.mgc.ac.cn/dbeCIS/) was used to identify lineage
Ia and lineage Ib eCIS (Sarris *et al.*, 2014; Chen *et al.*, 2019). It was
hypothesised that effectors associated with eCIS would be encoded up to 6kb
downstream of the final 'core' gene in the operon: usually the Afp16 'cap'
equivalent gene. By appending the sequence information to a custom database,
the function of each ORF was predicted by sequence and domain architecture
using BLASTp and HHPred, respectively. Occurrences were then counted using
regular expressions. The distributions of predicted functions in lineage Ia and Ib
are shown in FIG. 1a and b, respectively.
Photorhabdus eCIS-downstream ORFs are significantly more highly
represented in their use of rearrangement hotspot (Rhs) repeat-containing
proteins, particularly in *P. asymbiotica* and *P. luminescens*. eCIS-downstream
ORFs of the exclusively insect-pathogenic *P. temperata*, meanwhile, were more
commonly predicted to have deamidase and hydrolase activities. It should be
noted that the dbeCIS database does not include certain strains of
Photorhabdus: *P. asymbiotica* HIT and JUN European strains and the
Australian strain of *P. asymbiotica* Kingscliff (Wilkinson *et al.*, 2010).
The 6kb cut-off in this methodology reflects the approximate sizes of each
ORF within an eCIS operon as well as the distances between them. In total,
there are 88 and 240 ORFs encoded up to 6kb downstream of the final base of
the final core gene in the operon in lineages Ia and Ib, respectively. Of note, 24
ORFs across 13 species in lineage Ib (10% of all ORFs in lineage Ib) had either
sequence or domain architecture similarity to a highly conserved domain of
unknown function (DUF), DUF4157: a previously-termed PVC-
metallopeptidase, although its function has never been confirmed. It contains a

Zn-binding motif, HExxH, typical of metallopeptidases; authors have
speculated its function to be involved in toxin interactions such as for loading or
export (Geller *et al.*, 2021). This hypothesis may be reflected in the current
work since every instance of DUF4157 was encoded adjacent to another eCIS-
downstream ORF with the only exception being *Stigmatella aurantiaca* strain
DW4/3-1.
In addition, tertiary structures of ORFs of lineage Ia putative eCIS effectors
were predicted using AlphaFold2 (Jumper *et al.*, 2021) which was shown to be
an accurate predictor of effector structure (FIG. S2). Signal peptides (SPs) could
be identified by low-confidence N-terminal extensions (FIG. S2A and S2B).
ORFs encoding SPs were verified under a set of conditions: amino acids 1-50
possessing an average pLDDT < 40 and considerably lower than that of the rest
of the protein. Remarkably, SPs were found in 55.4% of ORFs up to 5kb
downstream of a lineage Ia eCIS operon (FIG. 1A). These SPs are localised
almost entirely to the PVC loci in *Photorhabdus* and the Afp loci in *Serratia*
and *Yersinia*; this implicates the disordered SP as an Afp-homologue-specific
feature of known eCIS cargoes. It also raises the question of whether other
mechanisms of eCIS cargo-loading exist outside of the use of N-terminal SPs in
the other genera.

Protein prediction enables analysis of potentially relevant residues and
supersecondary structures in the Pvc15 AAA domains.

AlphaFold was used to predict the structure of the Pvc15 encoded at the Pnf
operon: Pvc15$_{pnf}$ (Evans *et al.*, 2021; Jumper *et al.*, 2021; Mirdita *et al.*, 2021).
According to HHPred, Pvc15 is a putative AAA+ domain-containing protein
most homologous to the mycobacterial homologue of Cdc48/p97/VPC and
orthologue to *M. tuberculosis* Rv0435c: Msm0858 from *Mycolibacterium*
smegmatis (5E7P) (Unciuleac, Smith and Shuman, 2016). Like Msm0858,
Pvc15 has an N-domain, spanning residues 1-205, as well as two tandem AAA

domains, D1 (residues 206-434) and D2 (residues 435-698) (FIG. S2; S2A by

residue, S2B by secondary structure, S2C by inferred domains according to

alignment (bottom), and S2D by pLDDT).

A closer look at the structure indicates that the conserved Walker A and B

motifs are present in each of D1 and D2 (FIG. 2A). At the centre of the ATPase

domain is a classical β-α-β sandwich or Rossmannoid fold (Hanukoglu, 2015), a

hallmark of AAA+ proteins and ordered as β5-β1-β4-β3-β2 (Miller and

Enemark, 2016).

The ATPase domain supersecondary structure is typical of a clade 3 AAA+

ATPase (FIG. 2B; Miller and Enemark, 2016). In D2, the Walker A motif

consists of a glycine-rich loop situated between β1 and α1. The "GTGK" motif

forms a compound nest of the form "LRLR" which coordinates the β-phosphate

of ATP or ADP within the concave of the NH bonds, as in the case with Ras

(FIG. S4). Divalent cations such as Mg^{2+} may also coordinate the phosphate

groups (Bugreev and Mazin, 2004). Lys502 is likely responsible for nucleotide

binding; a role which can be abrogated by mutating to alanine in homologous

structures (Hanson and Whiteheart, 2005).

Walker B motifs tend to be less well conserved but often feature a set of up

to 4 hydrophobic amino acids followed by a 'DE'; in Pvc15 these residues are

part of a 'DEAD' motif, which is often found in proteins with helicase activity

(FIG. 2C) (Watson and Milner-White, 2002; Cordin *et al.*, 2004). The aspartate

residue (D554) may coordinate a divalent cation such as Mg^{2+} whilst the

glutamate residue (E555) is responsible for ATP hydrolysis. Mutation of the

latter to a glutamine is sufficient to impair this process in other ATPases such as

the Lon protease (Shin *et al.*, 2020) and SecA (Kim *et al.*, 2013). On the other

hand, Jiang and colleagues (2022) identified the Walker A and B motifs of

Pvc15*pnf* but apparently failed to abolish its unfolding activity upon SP-

associated fluorescent proteins by point mutations, indicating that Pvc15 may

have other roles outside of its ATPase activity.

Additionally, the second region of homology (SRH) is a defining

characteristic of AAA+ ATPases (FIG. 2B; right): the sensor 1 motif is located

in the loop between β4-α4; the sensor residue, Asn597 is likely responsible for

orientation of the nucleophilic water used in hydrolysis by the Walker B motif.

Finally, two arginine fingers, Arg608 and Arg609, are located in a position

which may enable them to stabilise the negative charge of the γ phosphate

during ATP hydrolysis within a neighbouring cleft (i.e., *in trans*) in the

hexameric structure.

Overall, protein structure prediction indicates that Pvc15 is a type II class 3

'classical' AAA+ ATPase. A search of the DALI server (Holm and Rosenström,

2010) using the N domain of Pvc15 indicates many hits for DNA-directed

transcription factor subunits, the best result being RNA polymerase II

transcription factor B from *S. cerevisiae* with a Z-score of 7.5 at 2.9Å r.m.s.d

(PDB: 5OQJ) (Schilbach *et al.*, 2017) and also includes the MepR

transcriptional regulator from *S. aureus* with a Z-score of 6.8 and 3.5Å r.m.s.d

over 90 aligned residues (PDB: 4LD5) (Birukou *et al.*, 2013).

In the close type II orthologue from *M. smegmatis*, Msm0858, both the D1

and D2 domains are capable of ATP hydrolysis despite having deviant residues

in the D1 domain Walker B box and sensor 1 residue (Unciuleac, Smith and

Shuman, 2016). Meanwhile, the Cdc48/p97 homologue's D1 domain is used for

stabilising the hexamer and binding ATP whilst the D2 domain is the primary

catalytic domain (Baek *et al.*, 2013). All Pvc15 homologues have even more

deviant residues than these homologues: Pvc15 lacks even an aspartate or

glutamate residue in the Walker B box, the Walker A nest lysine is replaced

with arginine, and it lacks any second region of homology (i.e., sensor 1 and R-

fingers). Altogether, these data indicate that D1 is likely deficient for ATPase

activity but is, nonetheless, required for oligomerisation of Pvc15 to form a

functional ATPase hexameric complex.

Pvc15 hexamer predictions suggest an asymmetrical arrangement with pore poops for substrate translocation.

MoLPC was used to predict the most likely complete hexameric assembly of Pvc15 (FIG. S5) (Bryant *et al.*, 2022). The resulting structure represents an asymmetrical structure with a pore diameter of 13.7Å at the D2 domains (FIG. S5A). N-terminal domains assume a more open conformation and each subunit neighbours the next at an angle of between 9° and 12°; this produces an open right-handed 'spiral staircase' arrangement. Two of the subunits at the 'top' and 'bottom' of the staircase are predicted to be adjacent at an angle of ~46°.

Pore loops 1 and 2 spiral in accordance with the larger domains and represent two 'faces' capable of contact with the translocating substrate (FIG. S5B). Lys528 and Tyr529 form a cation-π interaction, which suggests a strong binding to substrate, whilst Arg657 may play a role in contacting negatively charged amino acids in the substrate. This arrangement is similar to that of Vps4 (PDB: 6BMF) though the environment surrounding the aromatic residue can be specific for the ATPase's intended function; the grip would be looser when neighbouring glycine (e.g. ClpB D2 (PDB: 6OAY)) or made more hydrophobic when neighbouring aliphatic residues such as valine (e.g. YME1 (PDB: 6AZ0)) (Puchades, Sandate and Lander, 2020).

Moreover, Arg608 and Arg609 are predicted to be located in a locale where they may contact the ATP-binding site *in trans* to communicate macromolecular changes during hydrolysis to the adjacent subunit (FIG. S5C).

Different signal peptides have differential effects on cargo stability.

Jiang and colleagues (2022) characterised Pvc15 as being necessary for loading of cargo proteins to the PVC. To establish whether this effect was due to cargo stabilisation or loading efficiency, it was investigated what how the SP affects cell lysate abundance.

To first ascertain the optimal time to measure protein expression after

induction in NiCo21 cells, aliquots taken over time were analysed by Western

blot and normalised by sum of lanes in a Coomassie stain. 3-4 h post-induction

had the highest levels of protein expression (FIG. S6), so subsequent

experiments on cargo stabilisation were conducted at 3 h post-induction with

IPTG and arabinose to induce both the cargo and PVC*pnf* operon.

PVC*pnf* with either native Pnf or Pnf toxoid (C190A) were induced in the

presence or absence of Pvc15 and normalised using the FLAG-tagged Pvc16 in

Western Blot analysis (FIG. 3A). Normalised band intensities on these Western

blots showed a significant decrease in the stability of Pnf when Pvc15 was

removed ($t(4) = -5.55$, $p = 0.005$ **), indicating that Pvc15 does act to increase

protein abundance in the lysate. Surprisingly, however, truncation of the SP

from its C-terminus made the cargo more stable than when it was encoded with

the native SP, regardless of whether Pvc15 was absent ($t(4) = 3.30$, $p = 0.030$ *)

or present ($t(4) = 4.26$, p-value $= 0.013$ *).

Measuring protein content after the addition of kanamycin at 20 μg/mL to

block protein synthesis indicated that the effects on stability were due to

differences in protein degradation rather than translation (FIG. 3C). Altogether,

this indicates that Pnf alone is rapidly degraded and rescued only by the co-

expression of Pvc15; of note, strains encoding Pvc15 displayed this effect

regardless of whether Pvc15 was induced, indicating that Pvc15 may exert this

chaperoning effect at very low cytosolic levels. Given this interpretation, Pvc15

is predicted to rescue roughly 30% of cytosolic Pnf from degradation (95% CI

[15%, 43%]) when compared to the abundance of the Pvc15-independent SP,

Pnf10.

Conversely, Pvc15 had seemingly no effect on the stability of Pnf encoded

with either the native Cif50 SP or the C-terminal truncated Cif10, though the

latter displayed a decrease to the stability of the cargo compared to the longer

SP both with and without Pvc15 ($t(3) = -11.28$, $p = 0.0013$ **; $t(4) = -5.51$, $p =$

0.005 **) (FIG. 3D). This indicates that Pvc15 interacts differentially between

the different PVC cargo SPs. In addition, the difference in band intensity

observed between Pnf10 and Cif10 indicates that the 10 amino acid tip is,

indeed, an important factor affecting the cargo's stability. In contrast, the SP tip

appears to be dispensable for Pvc15-dependent stabilisation since only Pnf50

displayed a differential effect with or without the chaperone.

Signal peptides work with the Pvc15 ATPase for loading.

If native Pnf can be stabilised by encoding shorter SPs, presumably making it a

less desirable target for protein degradation machinery, then perhaps the longer

SPs which are encoded naturally are preferred in the context of loading.

Therefore, it was hypothesised that cargoes encoding C-terminal truncated SPs

would be far less capable of being loaded into the PVC*pnf* chassis; this, then,

would justify the bacterium having evolved longer, unstable signal peptides in

its native cargoes to optimise loading whilst also encoding a nearby chaperone

to mitigate the accompanying instability.

To compare loading efficiencies of the different SP-truncated cargoes,

PVC*pnf* syringes were purified using FLAG immunoprecipitation (FIG. 4).

There was a visual decrease in the relative level of Pnf in the final elution step

of the α-Myc Western Blot (FIG. 4A). Overall, either truncation of the signal

peptide or the deletion of Pvc15 resulted in a ~50-75% decrease in the loading

efficiency of Pnf (FIG. 4B). These results, therefore, confirm the hypothesis that

shorter SPs, whilst conferring stability to the Pnf cargo, fail to recruit cargo to

the PVC at comparable levels to that of the full-length SPs.

After further purification of the PVCs by anion exchange, the absence of

cargo was made even more apparent by Western Blot, though, as expected, the

PVCs purified from the PVC*pnf*Δ*pvc15* operon were intact when imaged in

transmission electron microscopy (TEM) (FIG. 4C).

As shown by recent data, PVCs can be artificially targeted to certain cell

receptors to ensure optimal payload injection and cell type specificity whereby

Pnf can be made to be cytotoxic to the target cells (Kreitz *et al.*, 2023).

However, some background binding to non-targeted cells is also possible. Pnf

acts similarly to *E. coli* cytotoxic necrotizing factor, CNF, by deamidating a

crucial residue in RhoA which subsequently causes the formation and

maintenance of stress fibres (Buetow *et al.*, 2001; Vlisidou *et al.*, 2019); the

cytoskeleton has been subsequently shown to interfere with mitochondrial

activity via voltage-dependent anion channels (VDAC) and mitochondrial

creatine kinase (MtCK) (Guzun *et al.*, 2009). Therefore, the effects of Pnf-

loaded FLAG-purified PVCs (with native Pvc13 tail fibres) on eukaryotic cell

respiration could be tested on Raji B cells by the addition of resazurin to

measure fluorescence from the mitochondria-reduced form: resorufin (FIG. 4D).

The optimal starting cell concentration for a 24 h time course was found to be 8

x 10^5 cells/mL (FIG. S7). Despite a decrease in the relative levels of resorufin,

this human cell line did not indicate a decrease of cell viability or number (data

not shown). Interestingly, PVCs purified more thoroughly by anion exchange

failed to replicate the effects of the FLAG-purified PVCs, perhaps indicating the

requirement for an ionic co-factor for functional contraction of the sheath.

Nonetheless, only the wild-type Pnf-loaded PVCs were capable of exerting a

decrease to the relative levels of cell respiration.

These results also find that the SP does not require cleavage to be loaded into

the PVC since purified samples resulted in Pnf molecular weights

corresponding to uncleaved SPs (~39kDa). Smaller bands observed in *E. coli*

lysates (~25kDa and ~17kDa) are also present when expressing Pnf on its own

without any other PVC components despite Pnf itself not being detectable,

indicating that these bands may be the result of Pnf degradation by the cell.

Pvc15 domain D2 is an effective chaperone for the Pnf SP and is the sole functional ATPase domain.

Guided by the bioinformatic analysis of Pvc15 structure, plasmids encoding either wild-type or mutant Pvc15 downstream of the same promoter as the Pnf payload, pVTRaDuet (FIG. S8), were made and transformed into *E. coli* BL21 (DE3) co-expressing PVC*pnf*Δ*pvc15*. Mutants include deletions of the N to D1 domains (Pvc15ΔN-D1), the D2 domain (Pvc15ΔD2), or abolishment of the catalytic E555 (Pvc15E555Q). Optimal Pnf abundance was measured by comparing Western Blot band intensities from lysates at 4 h post-induction which showed that deletion of domain D2 resulted in a depletion of Pnf payload, suggesting this domain is an effective chaperone (FIG. 5A). Unlike when Pvc15 is deleted, however, some Pnf can be observed which may indicate that domains N and/or D1 are also able to stabilise the payload somewhat.

To ascertain whether these mutants still had ATPase activity, a malachite green assay was used to measure a change in OD_{650} and normalise to a standard curve (FIG. S10) to calculate the amount of inorganic phosphate generated from lysates when ATPase substrate is introduced (FIG. 5B). Each of the mutants had activity comparable to the negative control (PVC*pnf*Δ*pvc15* + Pnf; 1.9 mU/mL) whilst wild-type Pvc15 co-expressed with the PVC*pnf*Δ*pvc15* operon and the Pnf payload had much larger ATPase activity (23.0 mU/mL). Notably, this ATPase activity was not replicated if lysates did not co-express the remaining PVC*pnf* operon proteins, suggesting that Pvc15's ATPase activity, conducted solely by the D2 domain but dependent on intact N-D1 domains, also requires interactions with other PVC*pnf* proteins or mature PVC to function. Altogether, this data suggests that the PVC*pnf*-dependent ATPase activity facilitates translocation of payload into the PVC tube lumen.

Indeed, FLAG-purified PVC*pnf* lacked co-purification with payload Pnf only if co-expressed with the ATPase-incompetent Pvc15 (FIG. 5C), confirming

that ATPase activity is required for translocating payload into the tube lumen of

the mature PVC.

The effect of SP mutations on cargo stability.

Jiang and collagues (2022) investigated the effects of point mutations within the

signal peptide to identify any single residues which may have altered the

association of Pnf with the PVC using alanine scanning and failed to identify

any key residues that alter loading of Pnf50-*Renilla* luciferase (Rluc). However,

this data does not reveal the nature of Pvc15-specific interactions nor how

stability of the native cargoes, such as Pnf, are affected by point mutations since

Rluc is stable regardless of the presence or absence of these N-terminal SPs.

It was investigated whether mutations to the SP would confer instability to

the cargo with or without Pvc15. To demonstrate each set of point mutations, a

pVTRA vector encoding both the native Pnf downstream of double

polyhistidine-tagged Pvc15 was constructed and termed pVTRaDuet (Figure

S8A). However, α-His Western Blot did not indicate successful expression of

Pvc15 at 4 h post-induction or overnight incubation (data not shown). Since the

previous pVTRA vector indicated that Pnf is highly unstable at 4 h post-

induction in the absence of Pvc15, however, the expression profile of the

pVTRaDuet vector (Figure S8B) shows that the stabilising effect of Pvc15 is

present despite not being expressed at detectable levels in Western Blot

analysis.

Interestingly, Pnf co-expressed with Pvc15 for 20 h post-induction results in

loss of detectable cargo; however, cargo levels can be reconstituted when also

co-expressed with the rest of the PVC*pnf* operon (lanes 4-6 compared to lanes

10-12); there was no difference in Pnf-Myc band intensity at 4 h post-induction

of the pVTRaDuet vector with or without the co-expression of the

PVC*pnf*Δ*pvc15* operon. This may suggest additional interactions mediate cargo

stability at late stages of PVC induction, or that translocation into the PVC tube

lumen is necessary for retaining payload abundance in the cytosol after long
incubations.

Conclusion

This work showed that each of Pvc15's domains are vital for associating SP-
associated cargo proteins with the PVC, and that SPs are ubiquitous in other
homologous eCIS putative cargoes such as the Afp in *Serratia* and *Yersinia*.
Despite their similar role in PVC maturation, these SPs act differentially with
the Pvc15 ATPase and chaperone; their groups of homology between similar
strains and operons in *Photorhabdus* may give clues as to whether certain
conserved motifs within SPs have similar roles or protein-protein interactions.
The N-domain was found to have homology to DNA-binding proteins, but
what role such activity has in Pvc15's role should be investigated further. The
D2 domain, in particular, plays significant roles in the ATPase activity, SP
stabilisation, and PVC loading, whilst the D1 domain may be involved in
stabilising the Pvc15 hexamer. Understanding the functional roles of the SP,
Pvc15, and interactions with other PVC components will enable researchers to
facilitate the potential future use of these devices for eukaryotic inter-membrane
trafficking of bioactive therapeutic proteins.

Materials and Methods

Plasmid constructs used in this work.

The pVTRA vector (Pérez-Martin and de Lorenzo, 1996) was used for cargo
expression by the addition of 0.1-1.0mM IPTG. The pBAD-PVC*pnf* vector was
constructed by synthesis of the PVC*pnf* operon encoding FLAG (DYKDDDK)
to the C-terminus of Pvc16 and can be induced with 0.2% arabinose (Guzman *et*
al., 1995). The pVTRaDuet vector, which encodes Pvc15 and Pnf under the
control of the same promoter, was generated by PCR of Pvc15 with overhangs

encoding hexahistidine tags and the relevant restriction enzyme sites into

pVTRA: KpnI and BamHI using the primers (Forward: 5'-

GGTACCTTGACCACAATGACTTAGTCTGAGTAAAAAATATGCACCAC

CACCACCACCACAATATATCGCCTGTTTTTTATGATTCATTG-3';

Reverse: 5'-

GGATCCTCCTCAATGATGATGATGATGATGAAATGTTAATCGTCCGA

CTTTAGC-3').

Analysing the dbeCIS effector dataset.

Using R within RStudio (R Core Team, 2020; RStudio Team, 2020), regular

expressions (via the *grepl()* function) were used to count the occurrences of

chosen keywords that described the most common predicted function entries in

the dataset. Of the entries that did not include the set of keywords, the phrases

't\\dss', 'toxin', or 'effector' were used to infer the ORF as a toxin, designated

as 'Other Toxin', or otherwise assigned as 'Other'. If different functions were

predicted for a single ORF, the one that appeared most commonly was

preferred.

Function counts were normalised by the number of strains within each

species as well as the number of operons in each genome. For lineage Ib, the

counts were also normalised by the number of species in the dataset. Thus, one

can infer the frequency of each function for the expected number of ORFs

encoded within 5kb downstream of any given eCIS operon from any given

strain in that genus.

Protein expression.

Transformed BL21(DE3) *E. coli* (New England Biolabs) (NiCo21) were

incubated overnight in a starter culture of 10mL at 180rpm and 37 °C with the

appropriate antibiotic(s); each biological replicate was grown into a separate

starter culture to minimise inter-sample interference. 5mL of this culture was

diluted 1:100 and grown again for 3-4 h to $OD_{600} = 1.0$ at which point 0.2%

arabinose was used to induce expression of the pBAD-PVC*pnf* vector and 1mM

IPTG was used to induce the pVTRA vector encoding the cargo. Cells were

then incubated at 200rpm and 25 °C.

For measuring cargo stability, aliquots were taken 3 h post-induction; for

measuring degradation of the cargo, kanamycin was added at a final

concentration of 20μg/mL and aliquots were taken after 4 h. For PVC

purification, cells were incubated overnight after induction then centrifuged at

3,000g for 20 minutes at 4 °C. Cells were then lysed using 10mL of lysis buffer

per gram of pellet, prepared as follows:

- 200μg/mL lysozyme
- 0.5% Triton X-100
- 1 tablet complete EDTA-free protease inhibitor per 50mL
- 70μL per 100mL DNase I

After resuspension, cells were homogenized using a cell homogenizer at

30kpsi (One Shot Model; Constant Systems LTD).

For making aliquots at each stage of the purification process, 25μL of

sample was taken and added to 25μL 2X LDS sample buffer with 50mM DTT

and boiled for 20 minutes at 95 °C. The homogenized sample was then spun at

12,000g for 35 minutes at 4 °C and the supernatant was filtered through a

syringe filter with 0.45μm pore size.

To purify the PVC syringes, the Pvc16 'cap' or tail tube terminator protein

was tagged with a FLAG peptide (DYKDDDDK) and a FLAG resin

(GenScript, anti-DYKDDDDK G1 Affinity resin, cat. no.: L00432) was used in

conjunction with a gravity flow column (Bio-Rad Econo-column®; cat. no.:

7371512) according to the manufacturer's instructions.

SDS-PAGE, Western Blot analysis, and quantification.

Sodium-Dodecyl-Sulphate (SDS) Polyacrylamide Gel Electrophoresis was

done in precast gels (Bio-Rad TGX Mini-Protean 4-15% gels with Dual Colour

Standard, cat. no.: 4561084DC) which were used as per the manufacturer's

instructions. Runs were conducted in a Mini-Protean gel tank using PAGE

buffer and run at 150V for 45 minutes.

After gel running, the extracted gel was loaded onto a PVDF membrane

using a Biorad Transblot Turbo electroblotter for 7 minutes. The Thermo Pierce

Fast Western Blot kit (cat. no.: 35055) was used according to the

manufacturer's instructions. Rabbit anti-FLAG (DYKDDDDK, Cell Signaling,

14793) or mouse anti-Myc (Cell Signaling, 2276) monoclonal antibodies were

used and visualised using the horseradish peroxidase (HRP)-bound secondary

antibody provided in the Pierce kit after adding luminol and peroxide buffer.

Bands were visualised using the chemiluminescence protocol in a G:BOX -F3

(Syngene) and using 2x2 binning for more than 300 seconds of exposure.

For quantification of band intensities, samples were normalised based on

their OD_{600} for cell lysates or protein concentration for purifications, as

determined using the BSA protocol on a spectronanophotometer. In addition,

for a given band, the mean background value of the blot was subtracted and a

corresponding loading control was used for that replicate either in the form of a

fixed point band which appeared in all samples, such as the FLAG-tagged

Pvc16 cap protein, or the mean of all points in the corresponding lane in the

Coomassie blue stain calculated using FIJI (ImageJ). After dividing the mean

grey values of each band by its corresponding loading control mean grey value,

test bands were relativised to their control samples; statistical analysis was done

using a Student's T-test.

Resazurin assay for cell respiration.

Raji cells incubated at 5% CO_2 and 37 °C for 2 days were diluted to 8 x 10^5

cells/mL (determined by first conducting a calibration curve experiment to

optimise the starting cell number) and seeded at 100μL in quadruplicate in a
black, clear-bottomed 96-well plate. Background fluorescence from the media
was subtracted from the mean fluorescence of the 9 media-only wells at each
time point. The plate was incubated for 1 h at 5% CO_2 at 37 °C and PBS or a
sample of 10μg of PVC was added (measured using a nanospectrophotometer)
and incubated for a further 4 h. Finally, resazurin (12.5mg/mL stock) was added
to each well in quick succession (~5μM final concentration). The plate was
incubated for a further 6 h and fluorescence was measured: excitation = 530-
570nm, emission = 580-620nm. Though these experiments used cells from the
same culture flask and passage, biological replicates of separate purifications of
payload or PVC protein were added such that samples were treated
independently.

ATPase activity

The colorimetric ATPase assay kit (Abcam, cat. no.: ab234055) was used with
cell lysates, as per the manufacturer's instructions. 15mL of *E. coli* NiCo21
cultures grown to OD600 = 1.0 were induced with 0.2% arabinose and 0.5mM
IPTG to co-express PVC*pnfΔpvc15* and the LS-associated payload with or
without Pvc15 for 16 h then processed according to the manufacturer's
instructions.

Acknowledgements

This work was supported by the Biotechnology and Biological Sciences
Research Council (BBSRC) as part of the Midlands Integrative Biosciences
Training Partnership (MIBTP); grant number: BB/M01116X/1.
Thanks go to Dr. John James for assisting with experimental design and
providing certain materials and cell lines and to the collaboration with
Nanosyrinx Ltd.

References

493 Baek, G.H. *et al.* (2013) 'Cdc48: A Swiss Army Knife of Cell Biology',

494 *Journal of Amino Acids*, 2013, p. 183421. Available at:

495 https://doi.org/10.1155/2013/183421.

496 Bhardwaj, P., Mitra, A.K. and Hurst, M.R.H. (2021) 'Investigating the

497 Process of Sheath Maturation in Antifeeding Prophage: a Phage Tail-Like

498 Protein Translocation Structure', *Journal of Bacteriology*, 203(20), pp. e00104-

499 21. Available at: https://doi.org/10.1128/JB.00104-21.

500 Birukou, I. *et al.* (2013) 'The Molecular Mechanisms of Allosteric Mutations

501 Impairing MepR Repressor Function in Multidrug-Resistant Strains of

502 Staphylococcus aureus', *mBio*, 4(5), pp. e00528-13. Available at:

503 https://doi.org/10.1128/mBio.00528-13.

504 Bryant, P. *et al.* (2022) 'Predicting the structure of large protein complexes

505 using AlphaFold and Monte Carlo tree search', *Nature Communications*, 13(1),

506 p. 6028. Available at: https://doi.org/10.1038/s41467-022-33729-4.

507 Buetow, L. *et al.* (2001) 'Structure of the Rho-activating domain of

508 Escherichia coli cytotoxic necrotizing factor 1', *Nature Structural Biology*, 8(7),

509 pp. 584–588. Available at: https://doi.org/10.1038/89610.

510 Bugreev, D.V. and Mazin, A.V. (2004) 'Ca2+ activates human homologous

511 recombination protein Rad51 by modulating its ATPase activity', *Proceedings*

512 *of the National Academy of Sciences*, 101(27), pp. 9988–9993. Available at:

513 https://doi.org/10.1073/pnas.0402105101.

514 Chen, L. *et al.* (2019) 'Genome-wide Identification and Characterization of a

515 Superfamily of Bacterial Extracellular Contractile Injection Systems', *Cell*

516 *Reports*, 29(2), pp. 511-521.e2. Available at:

517 https://doi.org/10.1016/j.celrep.2019.08.096.

518 Cordin, O. *et al.* (2004) 'The newly discovered Q motif of DEAD-box RNA

519 helicases regulates RNA-binding and helicase activity', *The EMBO Journal*,

520 23(13), pp. 2478–2487. Available at: https://doi.org/10.1038/sj.emboj.7600272.

Desfosses, A. *et al.* (2019) 'Atomic structures of an entire contractile

injection system in both the extended and contracted states', *Nature*

microbiology, 4(11), pp. 1885–1894. Available at:

https://doi.org/10.1038/s41564-019-0530-6.

Evans, R. *et al.* (2021) *Protein complex prediction with AlphaFold-*

Multimer. preprint. Bioinformatics. Available at:

https://doi.org/10.1101/2021.10.04.463034.

Geller, A.M. *et al.* (2021) 'The extracellular contractile injection system is

enriched in environmental microbes and associates with numerous toxins',

Nature Communications, 12(1), p. 3743. Available at:

https://doi.org/10.1038/s41467-021-23777-7.

Guzman, L.M. *et al.* (1995) 'Tight regulation, modulation, and high-level

expression by vectors containing the arabinose PBAD promoter.', *Journal of*

Bacteriology, 177(14), pp. 4121–4130.

Guzun, R. *et al.* (2009) 'Regulation of respiration controlled by

mitochondrial creatine kinase in permeabilized cardiac cells in situ: Importance

of system level properties', *Biochimica et Biophysica Acta (BBA) -*

Bioenergetics, 1787(9), pp. 1089–1105. Available at:

https://doi.org/10.1016/j.bbabio.2009.03.024.

Hanson, P.I. and Whiteheart, S.W. (2005) 'AAA+ proteins: have engine, will

work', *Nature Reviews Molecular Cell Biology*, 6(7), pp. 519–529. Available at:

https://doi.org/10.1038/nrm1684.

Hanukoglu, I. (2015) 'Proteopedia: Rossmann fold: A beta-alpha-beta fold at

dinucleotide binding sites', *Biochemistry and Molecular Biology Education*,

43(3), pp. 206–209. Available at: https://doi.org/10.1002/bmb.20849.

Hapeshi, A. and Waterfield, N.R. (2017) '*Photorhabdus asymbiotica* as an

Insect and Human Pathogen', in R.H. ffrench-Constant (ed.) *The Molecular*

Biology of Photorhabdus Bacteria. Cham: Springer International Publishing,

pp. 159–177. Available at: https://doi.org/10.1007/82_2016_29.

Holm, L. and Rosenström, P. (2010) 'Dali server: conservation mapping in 3D', *Nucleic Acids Research*, 38(Web Server issue), pp. W545–W549. Available at: https://doi.org/10.1093/nar/gkq366.

Jiang, F. *et al.* (2019) 'Cryo-EM Structure and Assembly of an Extracellular Contractile Injection System', *Cell*, 177(2), pp. 370-383.e15. Available at: https://doi.org/10.1016/j.cell.2019.02.020.

Jiang, F. *et al.* (2022) 'N-terminal signal peptides facilitate the engineering of PVC complex as a potent protein delivery system', *Science Advances*, 8(17), p. eabm2343. Available at: https://doi.org/10.1126/sciadv.abm2343.

Jumper, J. *et al.* (2021) 'Highly accurate protein structure prediction with AlphaFold', *Nature*, 596(7873), pp. 583–589. Available at: https://doi.org/10.1038/s41586-021-03819-2.

Kim, D.M. *et al.* (2013) 'ATPase active-site electrostatic interactions control the global conformation of the 100 kDa SecA translocase', *Journal of the American Chemical Society*, 135(8), pp. 2999–3010. Available at: https://doi.org/10.1021/ja306361q.

Kreitz, J. *et al.* (2023) 'Programmable protein delivery with a bacterial contractile injection system', *Nature*, pp. 1–8. Available at: https://doi.org/10.1038/s41586-023-05870-7.

Kube, S. and Wendler, P. (2015) 'Structural comparison of contractile nanomachines', *AIMS Biophysics*, 2(2), pp. 88–115. Available at: https://doi.org/10.3934/biophy.2015.2.88.

Lee, Y.-W. *et al.* (2019) 'Protein Delivery into the Cell Cytosol using Non-Viral Nanocarriers', *Theranostics*, 9(11), pp. 3280–3292. Available at: https://doi.org/10.7150/thno.34412.

Miller, J.M. and Enemark, E.J. (2016) 'Fundamental Characteristics of AAA+ Protein Family Structure and Function', *Archaea*, 2016, p. 9294307. Available at: https://doi.org/10.1155/2016/9294307.

578   Mirdita, M. *et al.* (2021) *ColabFold - Making protein folding accessible to*

579 *all*, p. 2021.08.15.456425. Available at:

580 https://doi.org/10.1101/2021.08.15.456425.

581   Pérez-Martin, J. and de Lorenzo, V. (1996) 'VTR expression cassettes for

582 engineering conditional phenotypes in Pseudomonas: activity of the Pu

583 promoter of the TOL plasmid under limiting concentrations of the XylR

584 activator protein', *Gene*, 172(1), pp. 81–86. Available at:

585 https://doi.org/10.1016/0378-1119(96)00193-X.

586   Puchades, C., Sandate, C.R. and Lander, G.C. (2020) 'The molecular

587 principles governing the activity and functional diversity of AAA+ proteins',

588 *Nature reviews. Molecular cell biology*, 21(1), pp. 43–58. Available at:

589 https://doi.org/10.1038/s41580-019-0183-6.

590   R Core Team (2020) 'R: A language and environment for statistical

591 computing.' Available at: http://www.r-project.org/index.html.

592   RStudio Team (2020) 'RStudio: Integrated Development for R'.

593   Sarris, P.F. *et al.* (2014) 'A Phage Tail-Derived Element with Wide

594 Distribution among Both Prokaryotic Domains: A Comparative Genomic and

595 Phylogenetic Study', *Genome Biology and Evolution*, 6(7), pp. 1739–1747.

596 Available at: https://doi.org/10.1093/gbe/evu136.

597   Schilbach, S. *et al.* (2017) 'Structures of transcription pre-initiation complex

598 with TFIIH and Mediator', *Nature*, 551(7679), pp. 204–209. Available at:

599 https://doi.org/10.1038/nature24282.

600   Shin, M. *et al.* (2020) 'Structural basis for distinct operational modes and

601 protease activation in AAA+ protease Lon', *Science Advances*, 6(21), p.

602 eaba8404. Available at: https://doi.org/10.1126/sciadv.aba8404.

603   Snider, J., Thibault, G. and Houry, W.A. (2008) 'The AAA+ superfamily of

604 functionally diverse proteins', *Genome Biology*, 9(4), p. 216. Available at:

605 https://doi.org/10.1186/gb-2008-9-4-216.

Unciuleac, M.-C., Smith, P.C. and Shuman, S. (2016) 'Crystal Structure and

Biochemical Characterization of a Mycobacterium smegmatis AAA-Type

Nucleoside Triphosphatase Phosphohydrolase (Msm0858)', *Journal of*

Bacteriology, 198(10), pp. 1521–1533. Available at:

https://doi.org/10.1128/JB.00905-15.

Vlisidou, I. *et al.* (2019) 'The *Photorhabdus asymbiotica* virulence cassettes

deliver protein effectors directly into target eukaryotic cells', *eLife*, 8, p.

e46259. Available at: https://doi.org/10.7554/eLife.46259.

Wang, X. *et al.* (2021) 'Characterization of *Photorhabdus* Virulence

Cassette as a causative agent in the emerging pathogen Photorhabdus

asymbiotica', *Science China Life Sciences* [Preprint]. Available at:

https://doi.org/10.1007/s11427-021-1955-4.

Watson, J.D. and Milner-White, E.J. (2002) 'A novel main-chain anion-

binding site in proteins: the nest. A particular combination of φ,ψ values in

successive residues gives rise to anion-binding sites that occur commonly and

are found often at functionally important regions11Edited by J. Thornton',

Journal of Molecular Biology, 315(2), pp. 171–182. Available at:

https://doi.org/10.1006/jmbi.2001.5227.

Wilkinson, P. *et al.* (2010) 'New plasmids and putative virulence factors

from the draft genome of an Australian clinical isolate of Photorhabdus

asymbiotica', *FEMS Microbiology Letters*, 309(2), pp. 136–143. Available at:

https://doi.org/10.1111/j.1574-6968.2010.02030.x.

Yang, G. *et al.* (2006) 'Photorhabdus virulence cassettes confer injectable

insecticidal activity against the wax moth', *Journal of Bacteriology*, 188(6), pp.

2254–2261. Available at: https://doi.org/10.1128/JB.188.6.2254-2261.2006.

Figure Legends

FIG. 1. Taxonomic distributions of putative functions of lineage Ia and Ib eCIS-downstream ORFs.

The dbeCIS database was used to identify structural operons of extracellular contractile injection system (eCIS) in lineage Ia **(A)** and Ib **(B)**, as defined by Chen *et al.*, 2019. Open reading frames (ORFs) up to 6kb downstream of the final structural gene in the operon were appended to a database where BLASTp and HHPred were used to identify sequence homology and most likely function. Regular expressions were used to count how many times keywords appeared using R scripts. ORFs which did not fit any of the assigned keywords were classed as either 'Other Toxin' or 'Other' depending on additional information found for that entry. Numbers presented at the top of each bar represent the total number of ORFs used to generate the stacked bar. Functions were chosen based on the most commonly predicted functions in the database. AlphaFold2 predictions of each amino acid sequence was done to observe whether low confidence (i.e., highly disordered) N-terminal signal peptides (SPs) were present. The abundance of these sequences is displayed on the right axis with bars shown as red borders. In **(B)**, 'Protease' and 'Nuclease' are treated as subsets of 'Hydrolase' and are preferred to the more ambiguous 'Hydrolase' term. Genera encoding only one eCIS operon with one or less downstream ORFs were excluded from analysis: lineage Ia: *Chania* and *Shewanella*; lineage Ib: *Gynuella*, *Minicystis*, *Sorangium*, *Moorea*, *Geobacter*, *Dyadobacter*, *Draconibacterium*, *Anabaena*, *Aulosira*, *Tolypothrix*, *Fremyella*, *Herbaspirillum*, *Tenderia*, and *Stigmatella*.

FIG. 2. C-terminal ATPase domain in Pvc15 contains conserved features.
(A) Walker A and B motifs are highly conserved ATP and GTP-binding sites and are found in the C-terminal region of Pvc15 homologues including Afp and ClpV/TssH. **(B)** Alignment of the α/β sandwich region of the C-terminal D2 domain of Pvc15. Notable motifs include the Walker A and B motifs which are

highly conserved sequences of amino acids in AAA+ ATPases; these include

the 'nest' residues Gly499-Lys502 in the Walker A motif, which coordinate

ATP, and Asp554-Glu555 in the Walker B motif which coordinate a divalent

cation to catalyse ATP hydrolysis. Other notable residues in the second region

of homology (SRH) include the sensor 1 (S1) residue Asn597, the sensor 2 (S2)

residue Arg659, and the more distant Arg-finger residues which coordinate

neighbouring protomers in the quaternary structure: Arg608 and Arg609. **(C)**

The Walker A motif "LRLR" nest with proposed coordination of phosphate as

documented for the P-loop of Ras by Watson and Milner-White (2002). Phi

bond angles are also shown; carbon atoms are shown in green, nitrogen in blue,

and oxygen in red; Amber-relaxed side chains are depicted as thin sticks.

FIG. 3. C-terminal truncations of signal peptides differentially affect the

stability of the cargo.

The abundance of Pnf-Myc was characterised using normalised band intensities

from α-Myc + α-β-RNA polymerase (β-RNAP) primary antibodies. **(A)** *E. coli*

BL21(DE3) cells were induced for 3 h at 25 °C with 0.2% arabinose and

0.5mM IPTG for PVC*pnf* and cargo induction, respectively. Error bars indicate

standard deviation of 3 biological replicates (n = 3); statistical inference was

done using a Student's T-test. Deletion of Pvc15 from the PVC*pnf* operon

results in a dramatic decrease in abundance of native Pnf50-Pnf-Myc from cell

lysates (t(4) = -5.55, p = 0.005 **). However, C-terminal truncation of the Pnf

SP to retain the 10 N-terminal amino acids (Pnf10) increases the abundance of

the cargo by over 100% when co-expressed with the full PVC*pnf* operon (t(4) =

3.30, p = 0.030 *) as well as with the Pvc15-deleted operon (t(4) = 4.26, p =

0.013 *). Pnf10-Pnf-Myc abundance was unaffected by the presence or absence

of the chaperone, implying a differential effect of the chaperone targeted

towards the native 50 amino acid SP. **(B)** Cells induced for 3 h were centrifuged

to remove IPTG and pelleted cells were resuspended in the same volume of LB

+ kanamycin at 20 μg/mL to halt both transcription and translation. Aliquots

were then taken every 40 minutes. Non-linear regressions were predicted using

the lm() function in R to compare coefficients and half-lives. C-terminal

truncated Pnf10-Pnf decays at roughly the same rate as that of native Pnf

regardless of its co-expression with the PVC ($t_{1/2}$ = 40-48 mins). **(C)** For co-

expression with the Cif SP (Cif50), truncation led to a decrease in abundance of

the Pnf cargo, regardless of the presence of Pvc15 (with: t(3) = -11.28, p =

0.0013 **; without: t(4) = -5.51, p = 0.005 **). **(D)** The Cif SP truncation

mutant (Cif10) displayed a significantly faster second-order degradation rate

($t_{1/2}$ = 43 mins), compared to the native Cif SP (Cif50) which displayed a first-

order degradation rate ($t_{1/2}$ = 157 mins).

FIG. 4. Full-length SPs and a Pvc15 chaperone are required for effective

PVC*pnf* cargo association.

pVTRa vectors expressing Myc-tagged Pnf with either the full-length (Pnf50)

or the C-terminal truncated SP (Pnf10) were co-expressed with pBADPVC*pnf*

or a mutant with deleted Pvc15 (PVC*pnf*Δ*pvc15*). PVC*pnf* nanosyringes were

immunoprecipitated using FLAG-resin. **(A)** For each of the 4 cell strains, band

intensities of the purification steps, supernatant (S), flow-through (FT), and

elution (E), were compared by Western blot using α-Myc or α-FLAG primary

antibodies. Band intensity for Pnf cargo in the elution step of the native system

(Pnf50+Pvc15) was visually more pronounced. **(B)** α-Myc band intensity in **(A)**

was normalised by the corresponding α-FLAG band. More than a 50% decrease

in loading efficiency was observed for all mutants with either a C-terminal

truncated SP (t(3) = -3.22, p-value = 0.049 *), or in the absence of Pvc15 (t(4) =

-4.23, p-value = 0.013 *). Error bars show standard deviation with n = 2-3

reiterations of purified nanosyringes; statistical analysis was done with a two-

sided Student's T-test. **(C)** Western blot and TEM were used to verify the

purification of intact PVC*pnf* after anion exchange. The PVC*pnf*Δ*pvc15* mutant

failed to co-purify with cargo. **(D)** The effect on respiration of Pnf-loaded

FLAG-purified PVC*pnf*, as well as Ni-NTA-purified Pnf cargo, was tested on

Raji B cells to measure resazurin reduction into the fluorescent resorufin. The

native Pnf-loaded PVC*pnf* impeded the reduction of resazurin by the cells (t(6)

= -9.95, p-value = 5.569e-05 ***) whilst other samples did not affect cell

viability, including Pnf with a C190A mutation of the active site. Pnf co-

expressed with the PVC*pnf*Δ*pvc15* operon similarly had no effect. Pnf-Myc

alone was insufficient to affect resorufin fluorescence, indicating the importance

of its PVC*pnf* vector for delivery.

FIG. 5. Pvc15 domain D2 stabilises the payload and has PVC*pnf*-dependent

ATPase activity for cargo loading.

pVTRaDuet, encoding Pvc15 and Pnf downstream of the same promoter, was

expressed in *E. coli* and used to measure whether Pvc15 mutations influenced

(A) Pnf cell lysate abundance, **(B)** ATPase activity, and **(C)** cargo loading.

(A) Pnf co-expressed with Pvc15 lacking domain D2 had depleted abundance

compared to the wild-type (t(4) = -8.66, p-value = 0.0010 ***). The Pvc15 D2

B-box mutant, E555Q, retained a high level of Pnf in lysates compared to when

the whole domain was deleted (t(4) = -4.68, p-value = 0.0094 **), implicating

D2 as an essential chaperoning domain whilst N-D1 are less efficient

chaperones.

(B) A colorimetric malachite green ATPase assay (Abcam; Cat. No.: ab234055)

was done in a 96-well plate to analyse *E. coli* lysates. Only the wild-type Pvc15

co-expressed with Pnf and the PVC*pnf*Δ*pvc15* operon was shown to have

significant ATPase activity. Comparing to the Pvc15+Pnf PVC*pnf*Δ*pvc15*[+]

strain: *E. coli*+Pnf: t(4) = -4.59, p-value = 0.010 *; Pvc15ΔD2: t(4) = -4.29, p-

value = 0.013 *; Pvc15ΔN-D1: t(4) = -4.08, p-value = 0.015 *; Pvc15E555Q:

t(4) = -4.37, p-value = 0.012 *. Mutants were not significantly different from

that of the *E. coli*+Pnf negative control, whilst the pVTRaDuet +

pBADPVC*pnf*Δ*pvc15* strain was not significantly different from the positive

control (t(3) = -1.05, p-value = 0.37). Furthermore, wild-type Pvc15 only

retained its level of ATPase activity when co-expressed with the

PVC*pnf*Δ*pvc15* operon (with vs without PVC*pnf*Δ*pvc15*: t(4) = -3.18, p-value =

0.033 *).

Error bars show **standard error of the mean**; statistical inference was done via

a two-sided Student's T-test; ns = not significant, * = p < 0.05. Data were

collected as 3 replicates of separately cultured *E. coli* lysates.

(C) Pvc16-FLAG-tagged PVC*pnf* was purified using the α-DYKDDDDK resin

whilst co-expressing Pvc15 mutants. PVC*pnf*Δ*pvc15* strains co-expressing

ATPase-incompetent Pvc15 mutants showed a lack of associated Pnf payload

compared to the wild-type, indicating that ATPase activity was required for

effective cargo loading. S = supernatant, FT = flow-through, E = elution

(purified PVC*pnf*). Data is representative of 3 reiterations of PVC*pnf*

purifications. **(D)** Quantification of **(C)** as well as 2 other reiterations of PVC*pnf*

purifications.

Supplementary Material

Figure S1.

PVC signal peptide homology is more aptly described in distinct groups than

all-vs-all alignments.

Figure S2.

AlphaFold2.0 predictions of native PVC cargo effectors display signal peptides

as disordered polypeptides.

Figure S3.

Tertiary overlay of Pvc15 homologues indicates high structural conservation,
particularly in the D1 and D2 domains.

Figure S4.

D2 Walker A P-loop likely coordinates phosphate in its 'LRLR' nest.

Figure S5.

A complete prediction of a Pvc15 hexamer reveals an open right-handed spiral
staircase.

Figure S6.

Pnf expression calibration curve indicates optimal payload expression at 3-4 h
post-induction.

Figure S7.

Raji B cell resazurin assay calibration curve of starting cell number.

Figure S8.

pVTRaDuet vector for simultaneous Pnf-Myc and Pvc15 expression may
indicate additional PVC core interactions.

Figure S9.

Induction of the PVC*pnf* operon does not affect the ability of wild-type Pvc15
to rescue Pnf abundance.

Figure S10.

Standard curve for conversion of OD_{650} to nmol phosphate at t=0.

FIG. 1.

A

B

FIG. 2.

A

B

C

N

PVCpnf-15_(PAU03339) 1 MNISPVFYDSLNQDNDRDLSFLFSELERIDLALQHHFYCVESQRSELLDEFLLTEAEVVTRLDKPLGKPHWINDDYLAISQKGNVS-----LMAASRLMDLIERFELTDFERDVLLLGLLPHFDSRYYRLF 126
Msm0858   1 --MSQGFDDEFG------------GADQYDTPLSHLRLTARLNTSAL-DSR-RGVVRLHPEVLAALGIREWDAVALTGTRTTAAVAGVAGPGVPAGTALLDDVTLSNAGVRENAAVLVSPVTVYGARS---- 112

PVCpnf-15_(PAU03339) 127 SLIQGGQQGRLP---SFALALELFCHSALEKQVQQASFLHRAPLMGCQ---LLSIDTSQKTLAWLQTPFIT----DSGV--------YHF---------LLGHHY----------------IMPALEHC 212
Msm0858   113 -VTVSGSRLATQSISPATLRMALLGKV--MTVGDTVSLLPRDLGPGTSTSAATSALASSVG ITWTSELLTVTAVDPPGTVS-VQPNSVVSWGTGTPEDPAPPPTGRHTVSPQRSEQPVSFDDVKVTHPQAVKL 241

D1

PVCpnf-15_(PAU03339) 213 AEWLTPTGIGCYPEGLKQVLGNVLLSDNDNIRPIVLL**GMAGSARAY**TITNMMASEGKQTLLVDISKLADSDEKNIILQIKHILRETRMHGACLL**RNFC**LLVEQNKQL-----------LDSLSELLNQPEL 334
Msm0858   242 DEWLRLSLD--EPELL-KTLGA-------TPHLGVLVS**GPAGVGKA**TMVRAVCASRRV--VELDGPEVGALQVDERLRSVTSAVAAVTESGGVLF**ADVD**ALLPAGNEMRPPEPVATLILAELRKAVATPGV 361

　　　　　　　　　　　　　　　　　　　　　　α0　　　　　β1　　　　　　α1　　　　β2　　*Clade 3 helix*　PL1_D1　　　α2　　　　β3　　　　　　　α3

PVCpnf-15_(PAU03339) 335 RIVCLIEPYSPLVLKKIPVLLIEMPLLTPAEKARLLIASLPDNC-------------SEDIDTITLSQ-RYTFNPETLPLILQEAQLYQQQ---RDPLDILQQCDIRQALNLRAQQNFGQLAQRIIPKR 447
Msm0858   362 AFIA---------TSAVPEN-VDARLRAPEVCDRELGLSLPDATARRSLLEMLLRGVPSEDLDLGDIADHTPGFVVADLAAVVREGALRAAARASSSDDDPVLRHADLEGALTVIRPLSRSASEEVSVGSV 482

　　　　　　β4　　　　　β5　　　　　α5　　　　　　　　　α6　　　　　　α7　　　　　　　　　α8　　　β0

D2

PVCpnf-15_(PAU03339) 448 SLKDLLVSDEIAQQLREILIAIKYREQVLAGGFKDKIAYGTGISALFY**GDSGTGKT**MAAEVIADHIGVDLIKVDLSTVVNKYIGETEKNLSRIFDLAEQDA-GVLFF**DEAD**ALFGKRSETKDSQDRHANIEV 578
Msm0858   483 DLDDV---GDMVETKRALTEAVLWPLQ--HPDTFSRLGIDPPGRVL**GPPGCGKT**FVVRALASSGRLSVHAVKGSELMDKWVGSSEKAVRELFARARDSAPSLVFLDE**DEID**ALAPRRGQNFDSGVTDK--VV 607

　　　　　　　　α0　　　　　β1　　　　α1　　　　β2　　*Clade 3 helix*　PL1_D2　　　α2　　　　β3　　　　　　　α3　　　　　　PL2

PVCpnf-15_(PAU03339) 579 SYLL---QRLENYPGLVILS**TNNRGHLD**SAFN**RR**FTFITRF---TYPDEKIRKKMWQEIWPRNIKISEDIDFNELAQRTSVTGANIRNIALLSS--FFASEQGNDEVSNENIEIALKREL------AKVGR 695
Msm0858   608 ASLLTELDGIEPLRDVVVLG**ATNRPDLI**DPALL**LR**PGRLERLVFVEPPDAAARRDIL-RTAGKSIPLADDVDLDSLADDLDGYSAA--DCVALLRESAMTAMRRSIDAADVTAADVAKARETVRPSLDPAQVES 737

　　　　　　β4　　　Sensor 1　　　α4　　β5　　　　α5　　　　　　α6　　　　　　α7　　　　　　α8

PVCpnf-15_(PAU03339) 696 L-TF---- 698
Msm0858   738 LREFAEKR 745

　　　　　　　　　Sensor 1 R Finger x2

D

Lid   β-α-β base

D1

P-loop
Walker A
N310
R257
R309
Walker B

α0
β5
β1
α6
α5
β4
α3
α1
β3
α8
β2
α7
β0
Clade 3 helix
α2
Pore loop 1

90°

D2

Sensor 1
P-loop
N597
Walker A
E555
K502
D554
R608
R609
Walker B

α0
β5
β1
R608
α6
α5
α4
R609
N597
β4
Pore loop 2
α1
β3
α3
α8
β2
α7
β0
α2
Clade 3 helix
Pore loop 1

90°

E

FIG. 3.

FIG. 4.

FIG. 5.