Individual genotype but not phenotype predicts river migration success in Atlantic salmon

Paolo Moccetti[1,2,3], Jonathan D. Bolland[2], Colin E. Adams[4], Jessica R. Rodger[5], Hannele M. Honkanen[4], Matthew Newton[4], Angus J. Lothian[5], Andy D. Nunn[2], Domino A. Joyce[1]

[1] Evolutionary and Ecological Genomics Group, School of Natural Sciences, University of Hull, UK

[2] Hull International Fisheries Institute, School of Natural Sciences, University of Hull, UK

[3] Energy and Environment Institute, University of Hull, UK

[4] Scottish Centre for Ecology and the Natural Environment, School of Biodiversity One Health & Veterinary Medicine, University of Glasgow, UK

[5] Atlantic Salmon Trust Fellow, Scottish Centre for Ecology and the Natural Environment, School of Biodiversity One Health & Veterinary Medicine, University of Glasgow, UK

Author for correspondence:

Paolo Moccetti[1,2,3]

School of Natural Sciences, University of Hull

Cottingham Road, Hull, HU6 7RX, UK

Email address: p.m.moccetti-2019@hull.ac.uk, moccetti.paolo@gmail.com

ORCID

PM, 0000-0002-3257-7993; JDB, 0000-0001-7326-5075; CEA, 0000-0003-2470-9754; JRR, 0000-0003-1913-6533; HMH, 0000-0003-0157-4359; AJL, 0000-0002-4121-3817; ADN, 0000-0001-8370-1221; DAJ, 0000-0002-4253-5623

Keywords: Animal migration; salmon; SNPs; genome scan; Geometric morphometrics; telemetry

Abstract

Migratory species typically undertake demanding long-distance journeys, across different habitat types during which they are exposed to multiple natural and anthropogenic stressors. Mortality during migration is typically high, and may be exacerbated by human-induced pressures. Understanding individual responses to these selection pressures is rarely attempted, because of the challenges of relating individual phenotypic and genetic data to migration success. Here we show distinct SNP sets unequivocally separated Atlantic salmon smolts making successful migrations to sea from those that failed to migrate, in two different rivers. In contrast morphological variation was not diagnostic of migration success. Populations from each river were genetically distinct, and while different genes were implicated in migration success in each river, they related to common biological processes (for example osmoregulation and immune response). Given that migration failure should quickly purge polymorphism at selected SNPs from a population, the question of how genetic diversity in these populations is maintained is an important one. Standing genetic variation could be maintained by different life history strategies and/or environmentally driven balancing selection. Our work highlights the importance of preserving genetic diversity to ensure evolutionary resilience at the population level, and has practical implications for management.

Introduction

Animal migration has evolved independently many times across the Animal Kingdom (Dingle & Drake, 2007; Bowlin et al., 2010; Shaw, 2016). Migration events can involve large numbers of individuals moving between different habitats and regions, and these events play a key ecological and socio-economic role in natural and human communities (Bauer & Hoye, 2014). Migratory species typically rely on multiple habitats to complete their life cycle and often undertake demanding long-distance journeys exposing themselves to numerous natural and anthropogenic stressors, such as predation, adverse weather conditions, pathogens, pollution, artificial constructions and harvesting (Alerstam, Hedenström & Åkesson, 2003). Mortality during migration is typically high and can be exacerbated by human-induced pressures, such that it impacts upon migratory populations and the ecosystems that depend on them (Wilcove & Wikelski, 2008; Harris et al., 2009; Middleton et al., 2013; Klaassen et al., 2014; Baker et al., 2020). With an ongoing global decline of migratory species (Wilcove & Wikelski, 2008), a better understanding of the factors causing mortality in migration is urgently required to predict responses of migratory populations to future environmental challenges and implement incisive conservation actions.

Recent advances in telemetry technology have made it possible to investigate migratory behaviours of species both temporally and spatially (Doherty et al., 2017; Thorup et al., 2023). This has enabled a better understanding of the exogenous factors directly influencing migration mortality (Thorstad et al., 2013; Hays et al., 2003; Palacín et al., 2017; Weinz et al., 2020). Organisms require a suite of specific morphological, physiological and behavioural adaptive features to successfully complete a migratory cycle (Justen & Delmore, 2022). Given the phenotypic and genetic variation found in most populations, it is reasonable to expect that some genetic or phenotypic traits are more likely to increase migration success than others. However which traits these might be remains poorly understood.

Genomic tools have recently been applied to identify factors regulating migratory behaviour at the population or species level. Several studies have discovered the genetic basis for migratory features such as migration timing and distance, orientation and propensity to migrate, with specific genomic regions linked to these traits (Zhu et al., 2009; Liedvogel, Åkesson & Bensch,

2011; Hecht et al., 2012; O'Malley et al., 2013; Hess et al., 2014; Hecht et al., 2015; Pritchard
et al., 2018; Waples, Naish & Primmer, 2020; Justen & Delmore, 2022). However,
understanding the genomic basis on which selection could act at an individual level to dictate
migration success has rarely been attempted (but see Bourret, Dionne & Bernatchez, 2014),
despite the fundamental insights it could provide into how populations might respond to
selection, and the implications for management of conservation genetics of migratory species.
An important phenotypic trait expected to influence migration success is body morphology
(Minias et al., 2013). Morphological variation (i.e. body shape and size) can affect behaviour,
resource use, survival and reproductive success of individuals (Wainwright 1994; Skulason &
Smith, 1995; Fruciano, Tigano & Ferrito, 2011). The effect of morphology on movement is
particularly evident in fish because of a direct link to swimming performance (Pakkasmaa &
Piironen, 2000; Fisher & Hogan, 2007; Drinan et al., 2012; Stelkens et al., 2012; Páez &
Dodson, 2017). Chapman et al. (2015) found a direct correlation between migration propensity
and body shape, while other studies have demonstrated an increased ability and 'motivation'
to pass river barriers in relation to size, fat content and morphology (Newton et al., 2018;
Lothian et al., 2020; Goerig et al., 2020). Nevertheless, research on migration survival and
mortality as a consequence of body shape variation (as opposed to size; Kennedy, Gale &
Ostrand, 2007; Hostetter et al., 2012; Romer et al., 2013; Furey et al., 2016; Lilly et al., 2022)
is still lacking.
Atlantic salmon (*Salmo salar*) is a migratory species of socio-economic importance that has
suffered substantial declines over the past 40 years (ICES, 2021) due to multiple abiotic and
biotic factors not yet fully understood (Forseth et al., 2017; Dadswell et al., 2022). The Atlantic
salmon has a complex life cycle, which includes two long distance migration stages; a long-
distance feeding migration from freshwater to sea as a juvenile (smolt) and an adult returning
spawning migration from sea to freshwater. In addition it is a philopatric species, accurately
homing to its natal spawning grounds (Thorstad et al., 2010). Fidelity to a specific river limits
gene flow among populations and has been shown to promote the evolution of local adaption
through natural selection, genetic drift, and bottlenecks (Garcia De Leaniz et al., 2007; Fraser
et al., 2011). The seaward migration of smolts constitutes a key life-stage for Atlantic salmon
and provides an ideal opportunity to study the genetic and phenotypic components that may
differentially affect the ability of individual animals to successfully complete their migration.
The identification of genetic and phenotypic traits could play a vital role in local management
of Atlantic salmon (Bernos, Jeffries & Mandrak, 2020)
114
115 Here, we analysed genomic and morphological data of migrating Atlantic salmon smolts in
116 two rivers. We wanted to test to what extent (I) Atlantic salmon populations in the two rivers
117 were genetically distinct, and (II) migration success by seaward migrating smolts could be
118 predicted by specific genomic regions and/or morphological traits.
119

Methods

121
Sampling, tagging and study design
123
The study reported here formed part of a wider acoustic telemetry study to examine migratory
behaviours and migration success in juvenile salmon (smolts) on their first migration from natal
rivers to sea (see Whelan, Roberts & Gray, 2019). Atlantic salmon were captured between 11
April and 3 May 2019 from the rivers Oykel (57°59.640' N, 4°48.282' W) and Spey (57°24.960'
N, 3°22.602' W), Scotland, using 1.5 m diameter rotary screw traps and a box trap (Fig. 1).
Fish were anaesthetised in MS222 and tagged with Vemco V7-2L acoustic transmitters (7mm

diameter, 19.5mm length, 1.5g in air, 137 dB re 1µPa @ 1m, acoustic transmission repeat cycle of 28 seconds ± 10 seconds, InnovaSea, Bedford, Nova Scotia, Canada). For more details on the tagging and release procedure see Lilly et al. (2022). Before being tagged, fish were measured (fork length, mm), weighed (g), and photographed. Photographs of the left side of each fish were taken from approximately 30 cm directly above the fish, with a Fujifilm FinePix XP130 Compact Digital Camera on a background reference scale. An adipose fin-clip was also taken from every fish and stored in 96% ethanol for later DNA extraction.

Two acoustic monitoring receivers (InnovaSea VR2Tx) were deployed in each river, one of which was immediately downstream of the tagging site (0.2 and 0.6 km in the rivers Oykel and Spey, respectively; Fig. 1). The second receiver was deployed at the river mouth (Fig. 1). Of all the salmon tagged and released in the two rivers (Oykel, n = 149, Spey n = 150), 91.9 and 96.7% respectively were detected by the first receiver after release. Of these, 78 (Oykel) and 82 (Spey) smolts were randomly selected for this study, distributed evenly between migratory outcomes ensuring a balanced design. Fish from both rivers were allocated into two groups based on their migratory outcome; 1) fish detected on the second and final river receiver were categorised as 'successful' river migrants, and 2) fish only detected on the first receiver were considered as 'unsuccessful' river migrants (Table 1, Fig. 1). To assess receiver detection efficiency, additional receivers deployed as a part of the broader telemetry study in the marine coastal waters of the Moray Firth were used. Since all smolts detected in marine waters were also detected by the two freshwater receivers, detection efficiency was determined to be 100%, meaning that no fish were wrongly miscategorised as unsuccessful migrants as a result of missed detections at the second river receiver.

To assess whether the study rivers harboured genetically differentiated Atlantic salmon populations, the genetic variation across rivers in the Moray Firth (Fig. 1) was investigated. In addition to fish from the study rivers Oykel and Spey, Atlantic salmon smolts from the rivers Findhorn (*n* = 3; 57°25.05′ N, 3°53.35′ W), Deveron (*n* = 4; 57°30.45′ N, 2°42.35′ W) and Ness (*n* = 4; 57°27.17′ N, 4°15.35′ W) were included in the analysis. To further contextualise the relative genetic diversity of these rivers, Atlantic salmon samples from the River Ure, England (*n* = 76; 54°16.19′, N 1°44.57′ W) were also included in the analysis (Fig. 1). Fish from the Findhorn, Deveron and Ness were sampled in Spring 2019 using rotary screw traps, while fish from the Ure were captured employing backpack electric fishing equipment (Electracatch 24 V DC input, 200-400 V, 100 W, 50 Hz Pulsed DC, variable pulse width output).

Table 1. Classification of smolts (*n*) in the rivers Oykel and Spey based on tracking results.

River	Unsuccessful	Successful	Total river migration distance
Oykel	36	42	30.5 km
Spey	35	47	50.1 km

166
Figure 1. The study area (*a*) and genetic structuring (*b, c*) of salmon populations from rivers
flowing into the Moray Firth (Scotland), with samples from the River Ure (England, *a*) top left
panel) for comparison. Tagging locations and acoustic receivers are shown in map (*a*). The
principal components analysis (PCA) and the ADMIXTURE analysis plots are based on 44,504
SNPs pruned for linkage disequilibrium. In the PCA scatterplot (*b*), dots represent individual
fish, and variance (%) explained by the first and second axes are shown. Colours correspond
to rivers. In the ADMIXTURE plot (K=3; *c*) each fish individual is represented by a vertical
bar. 'Dev' and 'Fin' are abbreviations for the rivers Deveron and Findhorn, respectively.

Genomic analyses

DNA extraction, genotyping and quality control
DNA was extracted from adipose fin samples employing a modified Mu-DNA: Tissue protocol
(Sellers et al., 2018) using a solid phase reversible immobilization (SPRI) magnetic bead
capture method (adapted from Rohland & Reich, 2012) to isolate high molecular weight DNA.
The DNA samples were sent to the Centre for Integrative Genetics (CIGENE, Ås, Norway) for
genotyping, including biological and technical replicates to ensure consistency across plates.
A custom 220,000 SNP (Single Nucleotide Polymorphism) Affymetrix Axiom array designed
for Atlantic salmon (see Barson et al., 2015 for details) was used for data generation. Following
the manufacturer's instructions, only SNPs categorised as PolyHighResolution and
NoMinorHom were used for analyses, while SNPs with unknown position were excluded from
the dataset, leaving 213,945 available loci for genomic investigation. We then performed
quality control (QC) and filtering of SNPs data in PLINK version 1.9 and 2.0 (www.cog-
genomics.org/plink/1.9/ and www.cog-genomics.org/plink/2.0/; Purcell et al., 2007; Chang et
al., 2015). SNPs were filtered for Hardy-Weinberg equilibrium (PLINK 1.9. command: --*hwe*
0.001), Minor allele frequency (--*maf 0.05*) and genotype missingness (--*geno 0.1*), and
individuals with a high rate of missing SNPs (--*mind 0.1*) were discarded from analyses. First-
degree relatives were also removed using PLINK 2.0 (--*king-cutoff 0.177*). In the migration
success analyses, these QC steps were performed separately for the rivers Oykel and Spey and
resulted in the retention of 198,336 SNPs and 82 individual fish from the River Oykel and
201,475 SNPs and 78 individuals from the River Spey. For consistency, only the 194,909 SNPs
overlapping between the Oykel and the Spey were retained for the analysis of river migration

success. The fish used in this analysis were the same employed for morphometric
investigations, but included two additional individuals which were not photographed. For the
regional population structure analyses (paragraph below), QC was performed on all rivers
together and SNPs in high linkage disequilibrium were pruned in PLINK 1.9 (*--indep 50 5 1.4*)
leaving 44,504 unlinked SNPs available for analysis.
204
Regional rivers genetic structuring
To investigate the genetic variation across rivers in the Moray Firth (Fig. 1), a principal
component analysis (PCA) was performed in PLINK 1.9 using individuals from the rivers
Oykel, Spey, Findhorn, Deveron, Ness and Ure. Using the same samples, ADMIXTURE v.1.3.
(Alexander, Novembre & Lange, 2009) was used to infer the most likely number of genetic
clusters (K, testing from K=1 to K=6), that was determined based on the lowest cross-validation
error.
212
Outlier analysis
To detect SNP markers with unusually high levels of allelic differentiation between successful
and unsuccessful migrants in each river, three different approaches were computed using the
unpruned 194,909 SNP dataset. In the first approach, the R (R Core Team, 2022) package
'OutFLANK' v. 0.2 (Whitlock & Lotterhos, 2015), which estimates among-groups Fst for each
locus was used. For the second, the software BayPass 2.31 (Gautier, 2015; Olazcuaga et al.,
2020), which utilises the *XtX* statistics which accounts for the demographic history of the
populations was implemented. Thirdly, the allele-based chi-squared association test in PLINK
1.9 (command: *--assoc*) was utilised. See code (https://tinyurl.com/salmon-migration-success)
for details about parameters used to run these analyses. Outlier loci in 'OutFLANK' were
identified by applying a q-value < 0.05 threshold, while outliers in BayPass and in the
association test in PLINK were determined as the top 0.1% SNPs ranked by *XtX* scores (in
BayPass) and *P*-values (in PLINK). Outliers were visualised using the 'qqman' v. 0.1.8 R
package (Turner, 2014). Outlier SNPs found by the three methods were then overlaid and only
the outliers detected by all three methods were retained for further analyses. The suite of outlier
SNPs that resulted from this analysis were then used to run a PCA in PLINK to compare
successful and unsuccessful migrating salmon smolts in the two rives.
230
Gene annotation and functional enrichment
For each river, the ten genes closest to each outlier SNP were extracted using the *closest*
function in the software bedtools 2.29.1 (Quinlan & Hall, 2010) and the genes within 10 kb
upstream and downstream of outlier SNPs were filtered in R (Wellband et al., 2019). To test
whether these candidate genes were significantly enriched for a specific gene ontology (GO)
biological process term, we used the R package 'topGO' v. 2.46.0 (Alexa & Rahnenfuhrer,
2016) employing genes from within 10 kb of all 194,909 SNPs of our array extracted with
bedtools as background reference. An alpha value of 0.01 was adopted as significance
threshold. For these analyses, the NCBI *Salmo salar* Annotation Release 100 (ICSASG_v2)
was used as a reference genome, while the GO terms for each gene were obtained through the
R package 'Ssa.RefSeq.db' v. 1.2 (Grammes, 2016).
242
243 **Morphometric analyses**
244
245 *Landmark digitisation*
246 Fish morphology was analysed using length (mm), weight (g), Fulton's condition factor (K;
247 Nash, Valencia & Geffen, 2006) and geometric morphometrics (GM). The GM analyses were

based on photographs of 158 salmon (Spey *n* = 77, Oykel *n* =81). The images of each fish were imported into tpsUtil v. 1.78 (Rohlf, 2019) and randomly shuffled using the *Randomly order specimens* function so that the operator was blind to the river-of-origin of the specimens. Nine fixed and 4 semi- landmarks (Fig. 2) were digitised on each image by a single operator using tpsDig v. 2.31 (Rohlf, 2017) using a subset of landmarks from the scheme proposed by Moccetti et al. (2023). Furthermore, five linear body measurements (Fig. 2) used as proxy of body slenderness which has been associated to swimming ability were also included (e.g., Pakkasmaa & Piironen, 2001; Drinan et al., 2012). Landmark coordinates were imported into R and analysed using the 'geomorph' and 'RRPP' v. 4.0.4 packages (Adams et al., 2021; Baken et al., 2021; Collyer & Adams, 2021). Preliminary analysis revealed body bending as a major source of shape variation in the dataset. This was corrected by employing landmarks 1, 14, 15 and 11 (see Moccetti et al., 2023 for details). All subsequent analyses were performed on landmarks 1-13 only. PCA plots were produced with the 'ggplot2' package (Wickham, 2016).

Figure 2. Fixed (blue circles) and semi (green diamonds) landmarks and linear measurements used for the geometric morphometrics analyses of Atlantic salmon smolts (image modified from Moccetti et al, 2023). Landmarks 14 and 15 (red stars) were used to correct for body arching. **(1)** Tip of snout; **(2)** Midpoint between 1 and 3; **(3)** Directly above middle of eye; **(4)** Perpendicular to 3; **(5)** Dorsal surface posterior of cranium; **(6)** Perpendicular to 11; **(7)** Anterior insertion point of dorsal fin; **(8)** Anterior insertion point of adipose fin; **(9)** Dorsal insertion point of caudal fin; **(10)** Versal insertion point of caudal fin; **(11)** Posterior midpoint of hypural plate; **(12)** Anterior insertion point of anal fin; **(13)** Anterior insertion point of ventral fin; **(14)** Lateral line - perpendicular to 7; **(15)** Lateral line - perpendicular to 12.

The following analyses were performed separately for each river. First, we tested whether successful and unsuccessful migrating fish in the two rivers were different in length (mm), weight (g) or Fulton's condition factor using *t*- and *Mann-Whitney U*- tests depending on the data distribution. Fulton's condition factor (K) was calculated as: $K = 100 \times W.L^{-3}$, where W = weight (g) and L = length (cm). We next tested for a difference in body shape between successful and unsuccessful fish, and whether these differences were consistent across rivers. A generalised Procrustes analysis (GPA) was performed to remove effects not related to body shape through translation, scaling and rotation of the landmark configurations (Rohlf & Slice, 1990). The residual effect of fish size on body shape was tested using Procrustes ANOVAs, with Procrustes coordinates used as a response variable and log centroid size used as an independent variable with a randomised residual permutation procedure (10,000 iterations). No significant effect of size on shape was found in either river ($p > 0.05$). To visualise body shape

of successful and unsuccessful fish, a PCA was performed on the Procrustes-aligned
coordinates of fish of each migration category from each river. Procrustes ANOVAs and *t*-tests
were subsequently used to test for differences in body shape and linear distances between fish
with different migratory outcomes. For linear distance and length, weight and condition factor
comparisons, significance values were Bonferroni corrected to limit the increased error rate
correlated with multiple testing (Rice, 1989).
291

Results

Genomic analyses

Regional rivers genetic structuring
The study rivers Spey and Oykel were genetically differentiated from one another (Fig. 1). The
second Principal Component (PC2) successfully separated the fish from the River Ness,
geographically located between the Oykel and the Spey, from all the other rivers (Fig. 1). Fish
from the rivers Deveron and Findhorn clustered with the spatially adjacent River Spey (Fig. 1).
PC1 separated the more geographically distant River Ure, situated in northern England, from
all Scottish rivers flowing into the Moray Firth (Fig. 1). Similarly, ADMIXTURE analysis
identified three different genetic clusters consisting of the rivers Oykel, Spey and Ure, with the
rivers Deveron, Findhorn clustering with the Spey, while the Ness was admixed with the rivers
Oykel and Spey.
306

Genomic regions linked to migratory outcome
One hundred and fifty-three outlier SNPs were detected across the genome by all three methods
('OutFLANK', BayPass and PLINK) for migrating fish in the River Oykel, and 159 for fish
from the River Spey (Supplementary files 1, 2). None of the outlier SNPs were found in fish
from both rivers. The PCA computed on this subset of outlier loci accordingly resulted in a
clear separation of successful and unsuccessful fish along PC1 axis in both rivers (Fig. 3).
313

Figure 3. The Manhattan plots (*a, b*) show all SNPs (dots) in each river and the outliers (in green) detected by 'OutFLANK'. *c* and *d*; Principal components analysis scatterplots based on 153 (Oykel) and 159 (Spey) outlier SNPs between successful (blue) and unsuccessful (red) migrant Atlantic salmon smolts. Each dot represents an individual fish. Variance (%) explained by the first and second axes is also shown.

Gene annotation and functional enrichment
There were 119 and 120 putative coding regions (hereafter genes) within 10 kb of the outlier SNPs' locations in the Oykel and Spey fish, respectively (Supplementary file 3). None of these genes were identified as outliers in fish from both rivers. In both Oykel and Spey samples, 25 genes contained more than one outlier SNP within the 10kb region (Supplementary file 3). The two genes enclosing the highest number of outlier SNPs were the *anion exchange protein 2-like* (encompassing 17 SNPs, River Oykel) and the *collagen alpha-1(I) chain-like* (encompassing 11 SNPs, River Spey). Gene set enrichment analysis identified 29 (Oykel) and 56 (Spey) significantly enriched GO terms. ($p < 0.001$; Supplementary file 4). None of these were over-represented in both river datasets.

Morphological differences between successful and unsuccessful salmon

There were no differences in length, weight or Fulton's condition factor between successful and unsuccessful migrating smolts ($p > 0.07$; Supplementary Figure 1, Supplementary Tables 1, 2). Procrustes ANOVAs based on the 13 landmark coordinates did not show significant differences in body shape between successful and unsuccessful smolts in either of the two rivers ($p > 0.31$; Fig. 4, Supplementary Table 3). Likewise, after Bonferroni correction (new *alpha* value = 0.005), comparison of body linear measurements did not show any significant difference between migrating groups ($p > 0.04$; Supplementary Tables 4, 5).

Figure 4. Mean body shape projections (top) and principal components analysis scatterplots (lower panel) show an absence of shape difference between the migratory groups. Procrustes-aligned coordinates of successful (blue) and unsuccessful (red) Atlantic salmon, where dots represent individual fish are shown below. Variance (%) explained by the first and second axes and 95% confidence ellipses are displayed. Projections show a complete overlap of the blue

348 (successful) and red (unsuccessful) lines in both rivers despite magnifying morphological
349 differences three times to aid visualisation.
350

Discussion

352
Our work shows that distinct SNP sets unequivocally separated Atlantic salmon smolts making
successful migrations to sea, from those that failed to migrate to sea in two different rivers. In
both rivers, the outlier SNPs predicting individual migration success were near several genes
that could be relevant for migration but we found no evidence of phenotypic differences in
body shape between successful and unsuccessful Atlantic salmon river migrants.
Categorising genes containing at least two outlier SNPs by biological function highlighted both
similar and different processes across the study rivers. In fish from both rivers, genes putatively
linked to osmoregulation, immunity, nervous, skeletal and cardiovascular system development
and activity, and morphogenesis of different organs were detected. The enrichment analysis
supported these findings, with GO terms linked to these functions overrepresented in the two
river datasets.
Candidate genes linked to general neuronal, cardiovascular and skeletal functions may play an
important role in migration, but a direct link to smolt migration success is hard to determine.
In contrast, osmoregulation and immune response are processes shown to play an important
part in salmonid migration. In the Oykel 38 (of 153) and in the Spey ten (of 159) outlier SNPs
were located near six and four candidate osmoregulatory genes, respectively. These genes were
associated with a range of processes including ion transmembrane and water transport,
response to salt stress and hyperosmotic response. Osmoregulation and individual ability to
undergo physiological changes required for seawater entry has been shown to be important to
increase chances of survival and predator avoidance in seaward migrating salmonid smolts
(Kennedy, Gale & Ostrand, 2007) and could play a role in migration success of Atlantic salmon
smolts in the last tidal kilometres of the Oykel and Spey where transitional zone between
freshwater and saltwater occurs.
Immunity related genes were also widely detected in association with the outlier SNPs
separating successful and unsuccessful river migrants in fish from both the Oykel (27 SNPs
near seven genes) and Spey (11 SNPs near 4 genes). Studies using proteomics in Pacific salmon
(*Oncorhynchus* spp.) have found significant correlation between migratory outcome,
expression of specific immune-related genes and viral and parasite-induced infection burden
(Miller et al., 2011; Jeffries et al., 2014; Furey et al., 2021; Mauduit et al., 2022). Our findings
now highlight the potentially important role of parasite driven selection in Atlantic salmon
migration success. An additional factor that requires further investigation, is the possibility that
there are individual differences in response to the tagging process, since there are immune
genes (Oykel=3, Spey=2) annotated with GO terms involved in blood coagulation, wound
healing and response to anaesthetic. Our study is only possible to carry out on tagged fish, so
we can't include untagged fish for comparison.
While particular SNP sets allowed us to clearly predict migratory outcome of Atlantic salmon
smolts in the Oykel and Spey, analyses of length, weight, body condition and body shape did
not find any significant difference between successful and unsuccessful migrants in either of
the rivers. This is somewhat surprising given the importance that morphology plays in
swimming performance in fish (Webb, 1978; Webb, 1984; Fisher & Hogan, 2007; Langerhans
& Reznick, 2010; Assumpção et al., 2012), the specific hydrodynamic characteristics required
to effectively migrate in running waters (Langerhans & Reznick, 2010; Brodersen et al., 2014),
and that size, body shape and condition may also be important in anti-predatory behaviour
(Domenici et al., 2007). However, we did find a difference in smolt morphology between rivers

(Moccetti et al., 2023) and there is genetic differentiation of the Oykel and Spey salmon
populations indicating reproductive isolation between these geographically close populations,
likely facilitated by fine-scale homing. Different evolutionary and demographic histories,
(evidenced by the geographic structure we find between river populations) combined with
different contemporary ecological selection pressures will therefore lead to different traits
being linked to migration success. For example, there were no genes that contained outlier
SNPs nor GO processes in common in Atlantic salmon from both the Oykel and Spey. Despite
candidate genes involved with similar processes being detected in both river datasets, clear
differences in gene types enclosing outlier SNPs between the Oykel and Spey were also found.
Genes potentially linked to sexual characters and maturation were more prevalent in the Oykel
dataset (42 outlier SNPs associated with seven genes *vs.* two SNPs in one gene in the Spey).
In contrast, several genes involved with vision, hearing and olfaction as well as genes involved
with muscular development, contraction and aerobic metabolism were detected exclusively in
the Spey. These processes might potentially indicate a sex-biased migration survival
(Thompson et al., 2016) in the Oykel, and a role in locomotion and response to external stimuli,
such as predators, in migration success in the Spey (Domenici et al., 2007; Popper & Hawkins,
2019).
Overall, we found that migratory outcome for individual salmon smolts in given rivers, in a
given year, can be predicted successfully from a subset of SNPs. We next need to understand
the ecological and environmental factors which determine those subsets. From an evolutionary
and conservation point of view, the mechanism through which the observed genetic diversity
is maintained needs to be identified, given that migration failure should theoretically quickly
purge polymorphism at selected SNPs from a population. We propose that variation in life
history could maintain standing genetic variation for environmentally driven balancing
selection (Mérot et al., 2020). 'Partial' migration (Shaw, 2016), where only a portion of the
population migrates, is common in several taxa and may be responsible for maintaining high
genetic diversity if migratory and resident individuals interbreed (Pulido, 2011). Within-
population differences in migratory strategies (e.g. timing, duration and routes) between age-
classes and sexes are also well known phenomena, particularly in birds (Cristol, Baker &
Carbone, 1999). These sub-groups are exposed to different biotic and abiotic conditions
potentially selecting for different genotypes that may maintain the gene pool diversity within
the population (Dingle & Drake, 2007; Wittmann et al., 2017; Briedis & Bauer, 2018).
Alternative migratory and reproductive tactics are well documented in Atlantic salmon
(Fleming, 1998; Thorstad et al., 2010; Birnie-Gauvin, Thorstad & Aarestrup, 2019), thus
individuals with different life histories experiencing temporally and spatially fluctuating
selection can interbreed and induce genetic mixing. Typically, the life history of salmon
involves seaward migration followed by a return to their natal river to spawn, but a number of
males become sexually reproductive in freshwater as morphological juveniles before migrating
to sea ('precocious male parr'; Lepais et al., 2017). Their contribution to paternity could be
substantial (ca. 60% in one study, Saura et al., 2008). The number of years spent in freshwater
before smolting, and at sea before upstream spawning migration can also vary considerably
(Thorstad et al., 2010). Finally, unlike Pacific salmon species, a non-negligible proportion of
Atlantic salmon survive reproduction (especially females), return to the ocean as 'kelt' and
spawn multiple times (Hedger et al., 2009). Weather and ecological conditions can change
dramatically among and within years inducing different selective pressures on migrating smolts
and other salmon life stages. For instance, variations in water discharge and temperature may
affect ecological factors such as migration timing (Thorstad et al., 2010), predation (Kennedy,
Gale & Ostrand, 2007; Hostetter et al., 2012) and pathogen infection (Wagner et al., 2005) as
well as passage of artificial migration barriers (Marschall et al., 2011). Clearly, all these
variables may differentially alter the allele frequencies under selection and help maintain

standing genetic variation. Straying between rivers could also be a source of genetic diversity
(Palstra, O'Connell & Ruzzante, 2007; Keefer & Caudill, 2014), although we found no
evidence of this in our study.
From a conservation point of view, understanding and predicting these selection pressures
could be invaluable in managing existing populations, and could inform stock selection where
hatchery-reared individuals are used to augment populations (Jepsen, Nielsen & Deacon, 2003;
Koed et al., 2020; Waples, Naish & Primmer, 2020).
Overall, our findings show that migration success could be linked to specific genotypic traits
and highlight the importance of preserving genetic diversity for conservation, to allow
populations to respond to heterogeneity between years, and the increased variability that long-
term climate change may produce. Our next challenge is to understand in detail the selection
pressures and associated genetic changes in populations.
459

Acknowledgments

461 We would like to thank the Spey Fisheries Board and the Kyle of Sutherland Fisheries Trust
462 for their help and support with this project. We would also like to thank Jonathan Archer,
463 Georgios Kyriakou, Fraser Brydon, Jessica Whitney, Mustafa Soganci and Rowan Smith for
464 their help collecting the data for this study.
465

Funding

467 This research was part of the Missing Salmon Project funded by the Atlantic Salmon Trust.
468 PM was supported by the Leeds-York-Hull Natural Environment Research Council (NERC)
469 Doctoral Training Partnership (DTP) Panorama under grant NE/S007458/1.
470

Ethics

For fish samples from the rivers Oykel, Spey, Deveron, Findhorn and Ness, the care and use
of experimental animals complied with the UK Home Office animal welfare laws, guidelines
and policies (UK Home Office Licence PPL 70/8794) and was approved by the University of
Glasgow Animal Welfare and Ethics Review Board (AWERB). Atlantic salmon from the River
Ure were treated in compliance with the UK ASPA (1986) Home Office project licence number
PD6C17B56.
478

Competing interests

480 The authors declare that they have no competing interests.

Data accessibility statement

482 Data and code used for analyses will be made public upon publication.

References
484

485 Adams DC, Collyer M, Kaliontzopoulou A, Baken E. 2021. "Geomorph: Software for
486 geometric morphometric analyses. R package version 4.0.2." https://cran.r-
487 project.org/package=geomorph.
488 Alerstam T, Hedenström A, Åkesson S. 2003. Long-distance migration: evolution and
489 determinants. *Oikos* 103:247–260. DOI: https://doi.org/10.1034/j.1600-

0706.2003.12559.x.

Alexa, A, Rahnenfuhrer, J. 2016. topGO: Enrichment Analysis for Gene Ontology. DOI: 10.18129/B9.bioc.topGO

Alexander DH, Novembre J, Lange K. 2009. Fast model-based estimation of ancestry in unrelated individuals. *Genome Research* 19:1655–1664. DOI: 10.1101/gr.094052.109.

Assumpção L de, Makrakis MC, Makrakis S, Wagner RL, Silva PS da, Lima AF de, Kashiwaqui EAL. 2012. The use of morphometric analysis to predict the swimming efficiency of two Neotropical long-distance migratory species in fish passage. *Neotropical Ichthyology* 10. DOI: https://doi.org/10.1590/S1679-62252012000400012

Baken EK, Collyer ML, Kaliontzopoulou A, Adams DC. 2021. geomorph v4.0 and gmShiny: Enhanced analytics and a new graphical interface for a comprehensive morphometric experience. *Methods in Ecology and Evolution* 12:2355–2363. DOI: https://doi.org/10.1111/2041-210X.13723.

Baker NJ, Boubée J, Lokman PM, Bolland JD. 2020. Evaluating the impact of hydropower on downstream migrating anguillid eels: Catchment-wide and fine-scale approaches to identify cost-effective solutions. *Science of The Total Environment* 748:141111. DOI: https://doi.org/10.1016/j.scitotenv.2020.141111.

Barson NJ, Aykanat T, Hindar K, Baranski M, Bolstad GH, Fiske P, Jacq C, Jensen AJ, Johnston SE, Karlsson S, Kent M, Moen T, Niemelä E, Nome T, Næsje TF, Orell P, Romakkaniemi A, Sægrov H, Urdal K, Erkinaro J, Lien S, Primmer CR. 2015. Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature* 528:405–408. DOI: 10.1038/nature16062.

Bauer S, Hoye BJ. 2014. Migratory animals couple biodiversity and ecosystem functioning worldwide. *Science* 344. DOI: 10.1126/science.1242552.

Bernos TA, Jeffries KM, Mandrak NE. 2020. Linking genomics and fish conservation decision making: a review. *Reviews in Fish Biology and Fisheries* 30:587–604. DOI: 10.1007/s11160-020-09618-8.

Birnie-Gauvin K, Thorstad EB, Aarestrup K. 2019. Overlooked aspects of the Salmo salar and Salmo trutta lifecycles. *Reviews in Fish Biology and Fisheries* 29:749–766. DOI: 10.1007/s11160-019-09575-x.

Bourret V, Dionne M, Bernatchez L. 2014. Detecting genotypic changes associated with selective mortality at sea in Atlantic salmon: polygenic multilocus analysis surpasses genome scan. *Molecular Ecology* 23:4444–4457. DOI: https://doi.org/10.1111/mec.12798.

Bowlin MS, Bisson IA, Shamoun-Baranes J, Reichard JD, Sapir N, Marra PP, Kunz TH, Wilcove DS, Hedenström A, Guglielmo CG, Åkesson S, Ramenofsky M, Wikelski M. 2010. Grand challenges in migration biology. *Integrative and Comparative Biology* 50:261–279. DOI: 10.1093/icb/icq013.

Briedis M, Bauer S. 2018. Migratory connectivity in the context of differential migration. *Biology Letters* 14:20180679. DOI: 10.1098/rsbl.2018.0679.

Brodersen J, Chapman BB, Nilsson PA, Skov C, Hansson LA, Brönmark C. 2014. Fixed and flexible: Coexistence of obligate and facultative migratory strategies in a freshwater fish. *PLoS ONE* 9 : e90294. DOI: 10.1371/journal.pone.0090294.

Chang CC, Chow CC, Tellier LCAM, Vattikuti S, Purcell SM, Lee JJ. 2015. Second-generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience* 4:s13742-015-0047–8. DOI: 10.1186/s13742-015-0047-8.

Chapman BB, Hulthén K, Brönmark C, Nilsson PA, Skov C, Hansson L-A, Brodersen J. 2015. Shape up or ship out: migratory behaviour predicts morphology across spatial scale in a freshwater fish. *Journal of Animal Ecology* 84:1187–1193. DOI:

https://doi.org/10.1111/1365-2656.12374.

Collyer ML, Adams DC. 2021. "RRPP: Linear Model Evaluation with Randomized Residuals in a Permutation Procedure, R package version 1.1.2." https://cran.r-project.org/package=RRPP.

Cristol DA, Baker MB, Carbone C. 1999. Differential Migration Revisited: latitudinal segregation by age and sex class. In: *Current Ornithology*. Nolan V, Ketterson ED, Thompson CF. (Eds). Springer US, Boston, MA. Pp. 33–88. DOI: 10.1007/978-1-4757-4901-4_2.

Dadswell M, Spares A, Reader J, McLean M, McDermott T, Samways K, Lilly J. 2022. The Decline and Impending Collapse of the Atlantic Salmon (Salmo salar) Population in the North Atlantic Ocean: A Review of Possible Causes. *Reviews in Fisheries Science & Aquaculture* 30:215–258. DOI: 10.1080/23308249.2021.1937044.

Dingle H, Drake VA. 2007. What Is Migration? *BioScience* 57:113–121. DOI: 10.1641/b570206.

Doherty PD, Baxter JM, Gell FR, Godley BJ, Graham RT, Hall G, Hall J, Hawkes LA, Henderson SM, Johnson L, Speedie C, Witt MJ. 2017. Long-term satellite tracking reveals variable seasonal migration strategies of basking sharks in the north-east Atlantic. *Scientific Reports* 7:42837. DOI: 10.1038/srep42837.

Domenici P, Turesson H, Brodersen J, Brönmark C. 2007. Predator-induced morphology enhances escape locomotion in crucian carp. *Proceedings of the Royal Society B: Biological Sciences* 275:195–201. DOI: 10.1098/rspb.2007.1088.

Drinan TJ, McGinnity P, Coughlan JP, Cross TF, Harrison SSC. 2012. Morphological variability of Atlantic salmon Salmo salar and brown trout Salmo trutta in different river environments. *Ecology of Freshwater Fish* 21:420–432. DOI: 10.1111/j.1600-0633.2012.00561.x.

Fisher R, Hogan JD. 2007. Morphological predictors of swimming speed: a case study of pre-settlement juvenile coral reef fishes. *Journal of Experimental Biology* 210:2436–2443. DOI: 10.1242/jeb.004275.

Fleming IA. 1998. Pattern and variability in the breeding system of Atlantic salmon (Salmo salar), with comparisons to other salmonids. *Canadian Journal of Fisheries and Aquatic Sciences* 55:59–76. DOI: 10.1139/d98-009.

Forseth T, Barlaup BT, Finstad B, Fiske P, Gjøsæter H, Falkegård M, Hindar A, Mo TA, Rikardsen AH, Thorstad EB, Vøllestad LA, Wennevik V. 2017. The major threats to Atlantic salmon in Norway. *ICES Journal of Marine Science* 74:1496–1513. DOI: 10.1093/icesjms/fsx020.

Fraser DJ, Weir LK, Bernatchez L, Hansen MM, Taylor EB. 2011. Extent and scale of local adaptation in salmonid fishes: review and meta-analysis. *Heredity* 106:404–420. DOI: 10.1038/hdy.2010.167.

Fruciano C, Tigano C, Ferrito V. 2011. Geographical and morphological variation within and between colour phases in Coris julis (L. 1758), a protogynous marine fish. *Biological Journal of the Linnean Society* 104:148–162. DOI: 10.1111/j.1095-8312.2011.01700.x.

Furey NB, Bass AL, Miller KM, Li S, Lotto AG, Healy SJ, Drenner SM, Hinch SG. 2021. Infected juvenile salmon can experience increased predation during freshwater migration. *Royal Society Open Science* 8:201522. DOI: 10.1098/rsos.201522.

Furey NB, Hinch SG, Bass AL, Middleton CT, Minke-Martin V, Lotto AG. 2016. Predator swamping reduces predation risk during nocturnal migration of juvenile salmon in a high-mortality landscape. *Journal of Animal Ecology* 85:948–959. DOI: https://doi.org/10.1111/1365-2656.12528

Garcia De Leaniz C, Fleming IA, Einum S, Verspoor E, Jordan WC, Consuegra S, Aubin-Horth N, Lajus D, Letcher BH, Youngson AF, Webb JH, Vøllestad LA, Villanueva B,

Ferguson A, Quinn TP. 2007. A critical review of adaptive genetic variation in Atlantic salmon: Implications for conservation. *Biological Reviews* 82:173–211. DOI: 10.1111/j.1469-185X.2006.00004.x.

Gautier M. 2015. Genome-Wide Scan for Adaptive Divergence and Association with Population-Specific Covariates. *Genetics* 201:1555–1579. DOI: 10.1534/genetics.115.181453.

Goerig E, Wasserman BA, Castro-Santos T, Palkovacs EP. 2020. Body shape is related to the attempt rate and passage success of brook trout at in-stream barriers. *Journal of Applied Ecology* 57:91–100. DOI: 10.1111/1365-2664.13497.

Grammes, F. 2016. Ssa.RefSeq.db: Annotation package for the Salmo salar RefSeq (ICSASG_v2). https://rdrr.io/github/FabianGrammes/Ssa.RefSeq.db/

Harris G, Thirgood S, Hopcraft JGC, Cromsigt JPGM, Berger J. 2009. Global decline in aggregated migrations of large terrestrial mammals . *Endangered Species Research* 7:55–76. DOI: https://doi.org/10.3354/esr00173.

Hays GC, Broderick AC, Godley BJ, Luschi P. 2003. Satellite telemetry suggests high levels of fishing-induced mortality in marine turtles . *Marine Ecology Progress Series* 262:305–309. DOI: 10.3354/meps262305.

Hecht BC, Hard JJ, Thrower FP, Nichols KM. 2015. Quantitative genetics of migration-related traits in rainbow and steelhead trout. *G3: Genes, Genomes, Genetics* 5:873–889. DOI: 10.1534/g3.114.016469.

Hecht BC, Thrower FP, Hale MC, Miller MR, Nichols KM. 2012. Genetic Architecture of Migration-Related Traits in Rainbow and Steelhead Trout, Oncorhynchus mykiss. *G3 Genes|Genomes|Genetics* 2:1113–1127. DOI: 10.1534/g3.112.003137.

Hess JE, Caudill CC, Keefer ML, McIlraith BJ, Moser ML, Narum SR. 2014. Genes predict long distance migration and large body size in a migratory fish, Pacific lamprey. *Evolutionary Applications* 7:1192–1208. DOI: https://doi.org/10.1111/eva.12203.

Hostetter NJ, Evans AF, Roby DD, Collis K. 2012. Susceptibility of Juvenile Steelhead to Avian Predation: the Influence of Individual Fish Characteristics and River Conditions. *Transactions of the American Fisheries Society* 141:1586–1599. DOI: 10.1080/00028487.2012.716011.

ICES. 2021. Working Group on North Atlantic Salmon (WGNAS). ICES Scientific Reports. 3:29. https://doi.org/10.17895/ices.pub.7923

Jeffries KM, Hinch SG, Gale MK, Clark TD, Lotto AG, Casselman MT, Li S, Rechisky EL, Porter AD, Welch DW, Miller KM. 2014. Immune response genes and pathogen presence predict migration survival in wild salmon smolts. *Molecular Ecology* 23:5803–5815. DOI: https://doi.org/10.1111/mec.12980.

Jepsen N, Nielsen EE, Deacon M. 2003. Linking individual migratory behaviour of Atlantic salmon to their genetic origin. In: *Aquatic telemetry: advances and applications. Proceedings of the Fifth Conference on Fish Telemetry held in Europe, Ustica, Italy* (pp. 9-13).

Justen H, Delmore KE. 2022. The genetics of bird migration. *Current Biology* 32:R1144–R1149. DOI: https://doi.org/10.1016/j.cub.2022.07.008.

Keefer ML, Caudill CC. 2014. Homing and straying by anadromous salmonids: a review of mechanisms and rates. *Reviews in Fish Biology and Fisheries* 24:333–368. DOI: 10.1007/s11160-013-9334-6.

Kennedy BM, Gale WL, Ostrand KG. 2007. Relationship between smolt gill Na+, K+ ATPase activity and migration timing to avian predation risk of steelhead trout (Oncorhynchus mykiss) in a large estuary. *Canadian Journal of Fisheries and Aquatic Sciences* 64:1506–1516. DOI: 10.1139/f07-117.

Klaassen RHG, Hake M, Strandberg R, Koks BJ, Trierweiler C, Exo K-M, Bairlein F,

Alerstam T. 2014. When and where does mortality occur in migratory birds? Direct evidence from long-term satellite tracking of raptors. *Journal of Animal Ecology* 83:176–184. DOI: https://doi.org/10.1111/1365-2656.12135.

Koed A, Birnie-Gauvin K, Sivebæk F, Aarestrup K. 2020. From endangered to sustainable: Multi-faceted management in rivers and coasts improves Atlantic salmon (Salmo salar) populations in Denmark. *Fisheries Management and Ecology* 27:64–76. DOI: https://doi.org/10.1111/fme.12385.

Langerhans RB, Reznick DN. 2010. Ecology and evolution of swimming performance in fishes: predicting evolution with biomechanics. In: *Fish locomotion: an eco- ethological perspective*. Domenici O, Kapoor BG. (Eds.). Science Publisher, Enfield, New Hampshire. Pp. 200-248.

Lepais O, Manicki A, Glise S, Buoro M, Bardonnet A. 2017. Genetic architecture of threshold reaction norms for male alternative reproductive tactics in Atlantic salmon (Salmo salar L.). *Scientific Reports* 7:1–13. DOI: 10.1038/srep43552.

Liedvogel M, Åkesson S, Bensch S. 2011. The genetics of migration on the move. *Trends in Ecology & Evolution* 26:561–569. DOI: https://doi.org/10.1016/j.tree.2011.07.009.

Lilly J, Honkanen HM, McCallum JM, Newton M, Bailey DM, Adams CE. 2022. Combining acoustic telemetry with a mechanistic model to investigate characteristics unique to successful Atlantic salmon smolt migrants through a standing body of water. *Environmental Biology of Fishes* 105:2045–2063. DOI: 10.1007/s10641-021-01172-x.

Lothian AJ, Schwinn M, Anton AH, Adams CE, Newton M, Koed A, Lucas MC. 2020. Are we designing fishways for diversity ? Potential selection on alternative phenotypes resulting from differential passage in brown trout. *Journal of Environmental Management* 262:110317. DOI: 10.1016/j.jenvman.2020.110317.

Marschall EA, Mather ME, Parrish DL, Allison GW, McMenemy JR. 2011. Migration delays caused by anthropogenic barriers: modeling dams, temperature, and success of migrating salmon smolts. *Ecological Applications* 21:3014–3031. DOI: https://doi.org/10.1890/10-0593.1.

Mauduit F, Segarra A, Mandic M, Todgham AE, Baerwald MR, Schreier AD, Fangue NA, Connon RE. 2022. Understanding risks and consequences of pathogen infections on the physiological performance of outmigrating Chinook salmon. *Conservation Physiology* 10:coab102. DOI: 10.1093/conphys/coab102.

Mérot C, Llaurens V, Normandeau E, Bernatchez L, Wellenreuther M. 2020. Balancing selection via life-history trade-offs maintains an inversion polymorphism in a seaweed fly. *Nature Communications* 11:670. DOI: 10.1038/s41467-020-14479-7.

Middleton AD, Kauffman MJ, McWhirter DE, Cook JG, Cook RC, Nelson AA, Jimenez MD, Klaver RW. 2013. Animal migration amid shifting patterns of phenology and predation: lessons from a Yellowstone elk herd. *Ecology* 94:1245–1256. DOI: https://doi.org/10.1890/11-2298.1.

Miller KM, Li S, Kaukinen KH, Ginther N, Hammil E, Curtis JMR, Patterson DA, Sierocinski T, Donnison L, Pavlidis P, Hinch SG, Hruska KA, Cooke SJ, English KK, Farrell AP. 2011. Genomic signatures predict migration and spawning failure in wild canadian salmon. *Science* 331:214–217. DOI: 10.1126/science.1196901.

Minias P, Kaczmarek K, Włodarczyk R, Janiszewski T. 2013. Wing Shape Influences Stopover Strategies in a Migratory Shorebird, The Common Snipe. *The Condor* 115:535–542. DOI: 10.1525/cond.2013.120137.

Moccetti P, Rodger JR, Bolland JD, Kaiser-Wilks P, Smith R, Nunn AD, Adams CE, Bright JA, Honkanen HM, Lothian AJ, Newton M, Joyce DA. 2023. Is shape in the eye of the beholder? Assessing landmarking error in geometric morphometric analyses on live fish. *bioRxiv*:2022.10.11.509915. DOI: 10.1101/2022.10.11.509915.

Nash RDM, Valencia AH, Geffen AJ. 2006. The origin of Fulton's condition factor - setting
the record straight. *Fisheries* 31:236–238.
Newton M, Dodd JA, Barry J, Boylan P, Adams CE. 2018. The impact of a small-scale
riverine obstacle on the upstream migration of Atlantic Salmon. *Hydrobiologia*
806:251–264. DOI: 10.1007/s10750-017-3364-3.
O'Malley KG, Jacobson DP, Kurth R, Dill AJ, Banks MA. 2013. Adaptive genetic markers
discriminate migratory runs of Chinook salmon (Oncorhynchus tshawytscha) amid
continued gene flow. *Evolutionary Applications* 6:1184–1194. DOI:
https://doi.org/10.1111/eva.12095.
Olazcuaga L, Loiseau A, Parrinello H, Paris M, Fraimout A, Guedot C, Diepenbrock LM,
Kenis M, Zhang J, Chen X, Borowiec N, Facon B, Vogt H, Price DK, Vogel H,
Prud'homme B, Estoup A, Gautier M. 2020. A Whole-Genome Scan for Association
with Invasion Success in the Fruit Fly Drosophila suzukii Using Contrasts of Allele
Frequencies Corrected for Population Structure. *Molecular Biology and Evolution*
37:2369–2385. DOI: 10.1093/molbev/msaa098.
Páez DJ, Dodson JJ. 2017. Environment-specific heritabilities and maternal effects for body
size, morphology and survival in juvenile Atlantic salmon (Salmo salar): evidence from
a field experiment. *Environmental Biology of Fishes* 100:209–221. DOI:
10.1007/s10641-016-0572-z.
Pakkasmaa S, Piironen J. 2001. Water velocity shapes juvenile salmonids. *Evolutionary*
Ecology 14:721–730. DOI: 10.1023/A:1011691810801.
Palacín C, Alonso JC, Martín CA, Alonso JA. 2017. Changes in bird-migration patterns
associated with human-induced mortality. *Conservation Biology* 31:106–115. DOI:
https://doi.org/10.1111/cobi.12758.
Palstra FP, O'Connell MF, Ruzzante DE. 2007. Population structure and gene flow reversals
in Atlantic salmon (Salmo salar) over contemporary and long-term temporal scales:
Effects of population size and life history. *Molecular Ecology* 16:4504–4522. DOI:
10.1111/j.1365-294X.2007.03541.x.
Popper AN, Hawkins AD. 2019. An overview of fish bioacoustics and the impacts of
anthropogenic sounds on fishes. *Journal of Fish Biology* 94:692–713. DOI:
https://doi.org/10.1111/jfb.13948.
Pritchard VL, Mäkinen H, Vähä JP, Erkinaro J, Orell P, Primmer CR. 2018. Genomic
signatures of fine-scale local selection in Atlantic salmon suggest involvement of sexual
maturation, energy homeostasis and immune defence-related genes. *Molecular Ecology*
27:2560–2575. DOI: 10.1111/mec.14705.
Pulido F. 2011. Evolutionary genetics of partial migration - the threshold model of migration
revis(it)ed. *Oikos* 120:1776–1783. DOI: 10.1111/j.1600-0706.2011.19844.x.
Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P,
de Bakker PIW, Daly MJ, Sham PC. 2007. PLINK: A Tool Set for Whole-Genome
Association and Population-Based Linkage Analyses. *The American Journal of Human*
Genetics 81:559–575. DOI: 10.1086/519795.
R Core Team. 2022. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org/.
Rohlf F. 2017. tpsUtil v. 1.78. Stony Brook, NY: Department of Ecology and Evolution,
State University of New York.
Rohlf F. 2019. tpsDig v. 2.31, Stony Brook, NY: Department of Ecology and Evolution, State
University of New York.
Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* 26:841–842. DOI: 10.1093/bioinformatics/btq033.
Hedger RD, Hatin D, Dodson JJ, Martin F, Fournier D, Caron F, Whoriskey FG. 2009.

Migration and swimming depth of Atlantic salmon kelts Salmo salar in coastal zone and marine habitats . *Marine Ecology Progress Series* 392:179–192.

Rice WR. 1989. Analyzing Tables of Statistical Tests. *Evolution* 43:223–225. DOI: 10.2307/2409177.

Rohland N, Reich D. 2012. Cost-effective, high-throughput DNA sequencing libraries for multiplexed target capture. *Genome research* 22:939–946. DOI: 10.1101/gr.128124.111.

Rohlf FJ, Slice D. 1990. Extensions of the Procrustes Method for the Optimal Superimposition of Landmarks. *Systematic Biology* 39:40–59. DOI: 10.2307/2992207.

Romer JD, Leblanc CA, Clements S, Ferguson JA, Kent ML, Noakes D, Schreck CB. 2013. Survival and behavior of juvenile steelhead trout (Oncorhynchus mykiss) in two estuaries in Oregon, USA. *Environmental Biology of Fishes* 96:849–863. DOI: 10.1007/s10641-012-0080-8.

Saura M, Caballero A, Caballero P, Morán P. 2008. Impact of precocious male parr on the effective size of a wild population of Atlantic salmon. *Freshwater Biology* 53:2375–2384. DOI: 10.1111/j.1365-2427.2008.02062.x.

Sellers GS, Di Muri C, Gómez A, Hänfling B. 2018. Mu-DNA: a modular universal DNA extraction method adaptable for a wide range of sample types. *Metabarcoding and Metagenomics* 2:e24556. DOI: https://doi.org/10.3897/mbmg.2.24556

Shaw AK. 2016. Drivers of animal migration and implications in changing environments. *Evolutionary Ecology* 30:991–1007. DOI: 10.1007/s10682-016-9860-5.

Skulason S, Smith TB. 1995. Resource polymorphisms in vertebrates. *Trends in Ecology & Evolution* 10:366–370. DOI: https://doi.org/10.1016/S0169-5347(00)89135-1.

Stelkens RB, Jaffuel G, Escher M, Wedekind C. 2012. Genetic and phenotypic population divergence on a microgeographic scale in brown trout. *Molecular Ecology* 21:2896–2915. DOI: https://doi.org/10.1111/j.1365-294X.2012.05581.x.

Thompson NF, Leblanc CA, Romer JD, Schreck CB, Blouin MS, Noakes DLG. 2016. Sex-biased survivorship and differences in migration of wild steelhead (Oncorhynchus mykiss) smolts from two coastal Oregon rivers. *Ecology of Freshwater Fish* 25:642–651. DOI: https://doi.org/10.1111/eff.12242.

Thorstad EB, Rikardsen AH, Alp A, Økland F. 2013. The use of electronic tags in fish research–an overview of fish telemetry methods. *Turkish Journal of Fisheries and Aquatic Sciences* 13:881–896. DOI: 10.4194/1303-2712-v13_5_13

Thorstad EB, Whoriskey F, Rikardsen AH, Aarestrup K. 2010. Aquatic Nomads: The Life and Migrations of the Atlantic Salmon. In: *Atlantic Salmon Ecology*. Aas O, Einum S, Klemetsen A, Skurdal J. (Eds.). Pp. 1–32. Wiley-Blackwell. DOI: https://doi.org/10.1002/9781444327755.ch1.

Thorup K, Tøttrup AP, Willemoes M, Klaassen RHG, Strandberg R, Vega ML, Dasari HP, Araújo MB, Wikelski M, Rahbek C. 2023. Resource tracking within and across continents in long-distance bird migrants. *Science Advances* 3:e1601360. DOI: 10.1126/sciadv.1601360.

Turner SD. 2014. qqman: an R package for visualizing GWAS results using Q-Q and manhattan plots. *bioRxiv*:5165. DOI: 10.1101/005165.

Wagner GN, Hinch SG, Kuchel LJ, Lotto A, Jones SRM, Patterson DA, Macdonald JS, Kraak G Van Der, Shrimpton M, English KK, Larsson S, Cooke SJ, Healey MC, Farrell AP. 2005. Metabolic rates and swimming performance of adult Fraser River sockeye salmon (Oncorhynchus nerka) after a controlled infection with Parvicapsula minibicornis. *Canadian Journal of Fisheries and Aquatic Sciences* 62:2124–2133. DOI: 10.1139/f05-126.

Wainwright, PC. 1994. Functional morphology as a tool in ecological research. In:

Ecological morphology: integrative organismal biology. Wainwright PC, Reilly SM.
(Eds). Pp. 42–59 University of Chicago Press, Chicago, IL.
Waples RS, Naish KA, Primmer CR. 2020. Conservation and Management of Salmon in the
Age of Genomics. *Annual Review of Animal Biosciences* 8:117–143. DOI:
10.1146/annurev-animal-021419-083617.
Webb PW. 1978. Fast-Start Performance and Body form In Seven Species of Teleost Fish.
Journal of Experimental Biology 74:211–226. DOI: 10.1242/jeb.74.1.211.
Webb PW. 1984. Body Form, Locomotion and Foraging in Aquatic Vertebrates. *American*
Zoologist 24:107–120. DOI: 10.1093/icb/24.1.107.
Weinz AA, Matley JK, Klinard N V, Fisk AT, Colborne SF. 2020. Identification of predation
events in wild fish using novel acoustic transmitters. *Animal Biotelemetry* 8:28. DOI:
10.1186/s40317-020-00215-x.
Wellband K, Mérot C, Linnansaari T, Elliott JAK, Curry RA, Bernatchez L. 2019.
Chromosomal fusion and life history-associated genomic variation contribute to within-
river local adaptation of Atlantic salmon. *Molecular Ecology* 28:1439–1459. DOI:
https://doi.org/10.1111/mec.14965.
Whelan K, Roberts D, Gray J. 2019. The SAMARCH Project International Salmonid Coastal
and Marine Telemetry Workshop. Available at https://www.samarch.org/wp-
content/uploads/2020/05/SAMARCH-Tracking-Conference-Nov-2019-
final_compressed.pdf (accessed 15 March 2023).
Whitlock MC, Lotterhos KE. 2015. Reliable Detection of Loci Responsible for Local
Adaptation: Inference of a Null Model through Trimming the Distribution of FST. *The*
American Naturalist 186:S24–S36. DOI: 10.1086/682949.
Wickham H. 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag, New
York. ISBN 978-3-319-24277-4.
Wilcove DS, Wikelski M. 2008. Going, going, gone: Is animal migration disappearing? *PLoS*
Biology 6:1361–1364. DOI: 10.1371/journal.pbio.0060188.
Wittmann MJ, Bergland AO, Feldman MW, Schmidt PS, Petrov DA. 2017. Seasonally
fluctuating selection can maintain polymorphism at many loci via segregation lift.
Proceedings of the National Academy of Sciences 114:E9932–E9941. DOI:
10.1073/pnas.1702994114.
Zhu H, Gegear RJ, Casselman A, Kanginakudru S, Reppert SM. 2009. Defining behavioral
and molecular differences between summer and migratory monarch butterflies. *BMC*
Biology 7:14. DOI: 10.1186/1741-7007-7-14.
824