Single-cell analysis reveals diversity of tumor-associated macrophages and their interactions with T lymphocytes in glioblastoma

Sai Batchu [a], Khalid Hanafy [b], and Ajith J Thomas [c]

[a] Cooper Medical School of Rowan University
[b] Neurological Institute, Cooper University Hospital
[c] Department of Neurosurgery, Cooper University Hospital

Abstract

Glioblastoma (GBM) is one of the most aggressive primary CNS malignancies. Despite advances in treatment protocols, clinical outcomes have not followed. Key obstacles remain - namely, the understanding of the tumor microenvironment (TME), in which tumor associated macrophages and their interactions with T cells have been of great interest. Although previous studies examining TAMs in GBM have shown that certain types are associated with certain clinical and/or pathologic features [2], these studies have used the outdated M1/M2 paradigm of macrophage polarization and failed to include the continuum of TAM states in GBM, and especially how they interact with T cells. Therefore, the present paper used single-cell RNA sequencing data from adult IDH-wildtype glioblastoma, with the primary aim of deciphering the cellular interactions of the 7 TAM subtypes with T cells inside the GBM TME. Furthermore, the interactions discovered herein are compared to IDH-mutated astrocytoma, allowing for focus on the cellular ecosystem unique to GBM. The resulting ligand-receptor interactions, signaling sources, and global communication patterns discovered provide a framework for future studies to explore methods in leveraging the immune system for treating GBM.

Introduction

Glioblastoma (GBM) is one of the most aggressive primary CNS malignancies. Despite advances in treatment protocols, clinical outcomes have not followed as evidenced by a 5-year survival rate suspended near 5% [1]. Key obstacles remain - namely, the understanding of the tumor microenvironment (TME). Composed of extracellular matrix proteins, various immune cells, soluble factors, and various cytokines that differ in combination compared to normal brain tissue, the TME is a highly heterogenous neighborhood with which GBM interacts. Recent focus has shifted towards the cellular relations in this system, mainly the surrounding stromal cells of which macrophages are the most prominent and significant in disease progression [2].

Macrophages and microglia (local CNS macrophage derived from yolk sac) are both important components of the immune response. However, while microglia have self-renewal capacity and maintain brain homeostasis with minimal immune stimulation[3, 4], bone-marrow-derived macrophages infiltrate the brain only in pathological conditions to facilitate phagocytosis of apoptotic cells or necrotic tissue [5]. Indeed, in GBM, the monocyte-derived macrophages may compose as much as 50% of the total mass [6]. However, unlike normal macrophages, these tumor-associated macrophages (TAMs) behave differently, altering the bridge between innate and adaptive immune responses and causing states of pathologic TAM-mediated immunosuppression.

A major component of the TAM mediated immunosuppression involved the lymphoid lineage, namely T cells. It has been observed TAMs may impede T cell functionality, promote T cell exhaustion, and impede T cell migration [7]. As macrophages exhibit high degrees of plasticity, they have historically been viewed in the M1 (pro-inflammatory) or M2 (anti-inflammatory) phenotypes, with the M1 macrophages associated with anti-tumorigenic effects. It is now accepted, however, that these dichotomous nomenclature oversimplifies TAM plasticity, especially in a dynamic setting such as the TME of GBM [8–10]. To truly appreciate the cellular interactions of TAMs and T cells in GBM, a framework that includes a continuum of TAM activation states must be used.

Recently, 7 TAM subtypes were found to be preserved in almost all cancer types and were characterized using single-cell multi-omics technologies. These 7 TAM subsets, based on signature genes and enriched pathways, have been categorized as follows: interferon-primed TAMs (IFN-TAMs), immune regulatory TAMs (reg-TAMs), inflammatory cytokine-enriched TAMs (inflam-TAMs), lipid-associated TAMs (LA-TAMs), pro-angiogenic TAMs (angio-TAMs), resident-tissue macrophage-like TAMs (RTM-TAMs) and proliferating TAMs (prolif-TAMs) [10].

Although previous studies examining TAMs in GBM have shown that certain types are associated with certain clinical and/or pathologic features [2], these studies have used the outdated M1/M2 paradigm of macrophage polarization and failed to include the continuum of TAM states in GBM, and especially how they interact with T cells. Therefore, the present paper used single-cell RNA sequencing data from adult IDH-wildtype glioblastoma, with the primary aim of deciphering the cellular interactions of the 7 TAM subtypes with T cells inside the GBM TME. Furthermore, the interactions discovered herein are compared to IDH-mutated astrocytoma, allowing for focus on the cellular ecosystem unique to GBM. The resulting ligand-receptor interactions, signaling sources, and global communication patterns discovered provide a framework for future studies to explore methods in leveraging the immune system for treating GBM.

Methods

Data acquisition
Single cell RNA sequencing data matrices and metadata comprising myeloid and T cells from 7 newly diagnosed IDH wildtype GBM samples and 1 IDH mutant astrocytoma sample derived from a previous study was used for the present analysis [11]. Another dataset including immune cells from GBM samples was examined separately for in silico validation [12]. Additionally, previously published GBM spatial transcriptomics data (sample 269_T) was computationally examined to explore the spatial distribution of gene expression of certain cell type patterns in the tissue extracted from these samples with respect to the barcode-spot's spatial dimensions [13]. 'AUCell' R package was used to annotate the TAM subpopulations from the original datasets. Gene signatures derived from previous literature [10] was used for TAM subpopulation classification. The R packages 'celldex' and 'SingleR' were used to classify the other immune cell populations, including T cells, into further subtypes using the 'MonacoImmuneData' reference index, which contains normalized expression values of bulk RNA-seq samples of 29 sorted immune cell populations, allowing for high resolution classification of immune cell transcriptomes [14].

Ligand-receptor analysis
Cell communication analysis was based on CellChat [15]. The CellChatDB was used as the reference database of literature-supported ligand-receptor interactions, containing more than 1,900 validated and manually curated interactions including paracrine/autocrine secreted signaling interactions. Using this database of known ligand-receptor interactions, CellChat initially identifies differentially over-expressed receptors and ligands in each provided cell type. Each interaction is subsequently associated with a probability value, modelled by the law of mass action based on the average expression values of a receptor of one receiver cell type and the average expression values of the ligand of the counterpart cell type.

CellChat was run using 'trimean', a robust statistical method for calculating the average gene expression per cell type. This method assembles fewer interactions but performs better for predicting stronger interactions which is advantageous when selecting those interactions for further experimental validations. Lastly, the significance of these interactions is identified by randomly permuting the cell type labels and subsequently recalculating the interaction probability [15]. Comparisons between IDH wildtype and mutant for identifying up-regulated and down-regulated signaling was performed by comparing the communication probability between two datasets for each ligand-receptor pair and each pair of cell groups. Interactions with p-value < 0.05 were considered significant.

Identifying major signaling sources and global communication patterns
To examine dominant senders and receivers within the interactome, the present study leveraged metrics derived from graph theory which were previously utilized for social network analysis [15, 16]. Specifically, weighted-directed networks and associated metrics, including out-degree and in-degree, were used to discover dominant intercellular interaction senders and receivers. In a weighted-directed network, the weights (i.e., strengths of interactions) are defined as the calculated communication probabilities based on the gene expression of ligands and receptors. Out-degree is calculated as the sum of these communication probabilities of outgoing signaling from an individual cell group. In-degree is computed as the sum of communication probabilities of incoming signaling to a cell group. These scores were used to identify the dominant cell senders and receivers, respectively, of intercellular signaling networks in the weighted-directed network [15]. Additionally, information flow was calculated for a

signaling pathway by summing the communication probabilities amongst all pairs of cell groups in the predicted network [15, 17].

To identify communication patterns among all signaling pathways, non-negative matrix factorization was carried out, with the number of patterns based on the Cophenetic and Silhouette metrics, which measure the stability for a particular number of patterns via hierarchical clustering of the derived consensus matrix [15, 18]. Plainly, outgoing communication patterns show how sender cell types coordinate with each other and how they work with certain signaling pathways to support communication. Correspondingly, incoming patterns reveal how target cell types (i.e., cells receiving signals) coordinate with each other to respond to incoming signaling.

Results

To predict significant ligand-receptor interactions between TAMs and T cells, a comprehensive dataset including single cell gene expression data from both IDH wildtype primary GBM and IDH mutant gliomas was used [11]. To understand which cell types are dominant senders (i.e., cell types sending signals) and which are dominant receivers (i.e., cell types receiving signals) with respect to each ligand-receptor interaction analyzed, incoming and outgoing interactions strengths (weights) were calculated for both GBM and glioma (Fig. 1A). In both IDH wildtype and mutant, Angio TAMs displayed robust incoming and outgoing signaling strength. In the IDH wildtype, all seven TAM types displayed strong outgoing interaction strength compared to IDH mutant, where only Inflam TAMs and LA TAMs showed strong outgoing interaction strength.

Next, we then sought to identify those signaling pathways that were conserved and those that were specific to GBM by calculating the information flow within signaling pathways, defined as the sum of communication probabilities amongst the total pairs of cell groups in the network (i.e., the amount of activity occurring in the pathway) (Fig. 1B). Compared to the IDH mutant astrocytoma, GBM exhibited increased information flow in multiple pathways including GALECTIN, COMPLEMENT, MIF, SPP, and PTN. When considering the absolute information flow in each pathway (Fig. 1C), greater information flow was observed in CCL, ANNEXIN, MIF, GALECTIN, COMPLEMENT, and SPP1 pathways. These flow states imply that although IDH wildtype GBM has increased activity in these pathways relative to IDH mutant astrocytoma, both types of gliomas have substantial activity occurring within these signaling pathways.

Since IDH-wildtype GBM displayed prominent activity of TAMs (Fig. 1A) and displayed specificity for certain pathways (Fig 1B, C), we then sought to identify the specific signaling changes of the different TAM populations between GBM and IDH-mutant (Fig. 1D). Angio TAMs and LA TAMs were observed to have shared signaling amongst both glioma types in SPP1 and MIF pathways. However, the other five TAM subtypes showed increased signaling in SPP1 specific to GBM (Fig. 1D). Overall, the seven TAM populations were observed to be contributing in an increased capacity to pathways such as SPP1, COMPLEMENT, GALECTIN and PTN compared to their counterparts in IDH mutant astrocytoma.

Further examination of each pathway yielded numerous significant ligand-receptor interactions (Fig. 2A, B). All types of TAMs exhibited MIF and SPP1 binding interactions with ligands on T cells such as CD44 and ITGA4+ITGB1 (Fig. 2A). Conversely, T cell ligands interacting with receptors on TAMs commonly included MIF-(CD74+CD44) and ANXA1-FPR1 (Fig 2A). To further verify these interactions, another dataset that included was analyzed independently in a similar fashion [12]. The ligand-receptor interactions were corroborated by the second dataset, as interactions including SPP1, MIF, LGALS9 between TAMs to T cells were found to be significant (Fig 3A, 3B).

Along with understanding ligand-receptor interactions in the tumor microenvironment, it is crucial to analyze how different cell types coordinate with each other in terms of signaling. To address this, a pattern recognition methodology was employed to identify global communication patterns of TAMs and T cells (see Methods). These communication patterns can be thought of as assigning cell types to signaling pathways for either outgoing signaling (considering cells as senders) or incoming signaling (considering cells as receivers) (Fig. 4A, B). The outgoing cell signaling patterns were broadly identified into 3 distinct categories. Most TAMs contribute to pattern 1, while pattern 2 was mainly contributed by Angio TAMs and T regulatory cells (Fig. 4A). Pattern 1 was mainly composed by pathways involving TNF, GALECTIN, SPP1, and COMPLEMENT while pattern 2 mainly composed of RESISTIN and ANGPTL signaling (Fig. 4A).

For the incoming signaling pathways in IDH wildtype GBM, 3 discernable patterns also emerged (Fig. 4B). Interestingly, Angio TAMs displayed pattern 3, as opposed to the other TAMs which displayed a pattern 1 incoming signaling. Pattern 3 is mostly contributed by VISFATIN, IL-1, ANGPTL and CD40.

These communication results imply that TAMs in GBM may depend on numerous signaling networks that may overlap in function while certain TAMs, such as Angio TAMs, rely on fewer and more heterogenous communication patterns. Additionally, leveraging these incoming and outgoing patterns may provide insight into autocrine vs. paracrine pathways for a given cell type in the GBM cellular landscape.

Given this interplay amongst TAMs and T cells, especially within the context of T cell exhaustion in GBM [19], a better understanding of the spatial interactions of these cell types should be explored. To this end, spatial transcriptomic data from a GBM dataset [13] was examined by transposing the gene signatures of T cell exhaustion and the TAMs analyzed in the present study (Fig. 5). The T cell signature used included persistent expression of multiple inhibitory receptors, such as *PD-1, LAG-3, TIM-3, and CTLA-4,* and TIGIT [19]. The spatial plots displayed a prominent overlap amongst TAM signatures and the T cell exhaustion gene signature, with a higher density and expression of T cell exhaustion related genes in spaces occupied by TAM signatures (Fig. 5)

Discussion

It is crucial to note, however, that the current paradigm of macrophage polarization into M1/ m2 activation states was largely defined in tissue culture without incorporating other polypeptide growth factors that could attenuate macrophage polarization. It is now known that the in-vivo profiles of macrophages are context-dependent, especially of TAMs in a dynamic TME. Therefore, unlike previous reports, this present study aimed to elucidate the interactions of T cells with TAMs while considering the continuum of TAM activation states.

IDH mutant gliomas have a distinct immune profile from IDH WT high grade gliomas and a diverse phenotype of TAMs.

All seven subsets of TAMs in both groups. Angio TAMs have an angiogenic signature associated with significant immune suppression and tumor cell proliferation. Though found in proportionally smaller numbers, they are the dominant receivers and senders in the population communicating with the lymphocyte population. Angio TAMs were rarely encountered in IDH mutant gliomas. Both IDH mutant and IDH WT gliomas have increased representation of myeloid cells with lymphocyte population increased in the IDH WT tumors [20]. RTM – TAMs formed the largest component of TAMs in the IDH mutant variety. RTM-TAMs are very similar to normal RTMs and a putative role for them in tumor invasiveness in lung cancer has been suggested [21]. LA TAMs and Proliferative TAMs were also seen in

larger numbers in the WT high grade gliomas. LA- TAMs express more lipid related and oxidative phosphorylation genes which include APOC1, APOE, ACP5 and FABP5. Lipid catabolism in macrophages is associated with IL-4 associated pathways and anti-inflammatory phenotype[22]. Inflammatory TAMs have an inflammatory signature and express inflammatory cytokines including IL1B, CXCL1/2/3/8, CCL3, and CCL3L1. Inflammatory TAMs play an active role in the recruitment of immune cells including granulocytes into the TME. Regulatory TAMs which are found more often in WT glioma seemed to have a profile consistent with microglia and seemed to play a more quiescent role in terms of activating lymphocytes. This is corroborated by Venteicher et al who demonstrated microglia in IDH mutant glioma TME associated with a less malignant phenotype [23]. Effector memory CD8 T cells were more often encountered in WT glioma while NK cells were more common in IDH mutant Glioma. Th1, Th2 and Th 17 cells were hardly seen in IDH Mutant in comparison to the WT tumors. Vd2 gd T cells were present abundantly in both groups.

TAMs in WT Glioblastomas secrete a common group of ligands which are immunomodulatory

The immune cells are responsible for secretion of cytokines in the GBM TME[6]. Galectin 1 is highly expressed in glioblastoma stem cells and are important for immunomodulation and angiogenesis[24, 25]. The signaling pathways associated with these CD 45 + cells GBM were Galectin, Complement, MIF, SPP and PTN. In addition, when considering the absolute information flow in each pathway, CCL and ANNEXIN also assumed importance. All these signaling pathways account for significant immune suppression and support tumor proliferation. Regardless of the subtype of TAM, SPP1(Osteopontin) seems to be very significant in outgoing and incoming signaling. SPP1 suppresses T cell function through an IRF8-OPN-CD 44 axis[26]. MIF (Macrophage Inhibitory Factor) also can act to reduce CD8+ Cytotoxic lymphocyte (CTL) activity [27, 28]. We found that there was significant upregulation of the MIF ligand - CD74/CD44 pair. Blocking this interaction in TAMs and Dendritic cells in melanoma decreases the immune suppression activity [29]. It can recruit macrophages and promote an immunosuppressive microenvironment. MIF also interacts with immune checkpoint molecules like PD-L1, contributing to T cell exhaustion and immune evasion. Specifically, in the intratumoral environment it suppresses cytotoxic T cells and NK cells. Complement secreted by TAMs is increasingly seen as immunosuppressive and promotes tumor growth. This immunosuppressive environment is characterized by high expression of immune checkpoints such as PD-1 and LAG-1.

Visfatin/NAMPT can act as a growth factor, enhancing tumor cell proliferation and survival. Additionally, lymphocyte proliferation, and promoting the differentiation of regulatory T cells (Tregs). IL-1 in glioblastoma is a pleiotropic cytokine that contributes to tumor growth, invasion, modulation of immune responses, and therapeutic resistance. Angptl2 is upregulated in glioblastoma and has been associated with tumor progression and poor prognosis. It is involved in promoting angiogenesis by inducing the proliferation, migration, and tube formation of endothelial cells. Angptl2 can also contribute to tumor cell invasion and metastasis by enhancing matrix metalloproteinase (MMP) expression and activity. Furthermore, it has been shown to promote the recruitment of TAMS which can foster an immunosuppressive microenvironment and support tumor growth.

Angptl4 has diverse functions in glioblastoma, including roles in angiogenesis, tumor cell survival, and metastasis. It can inhibit endothelial cell sprouting and angiogenesis, exerting anti-angiogenic effects. Angptl4 has also been reported to induce apoptosis (programmed cell death) in glioblastoma cells, suggesting a potential tumor-suppressive role. However, in certain contexts, Angptl4 expression has been associated with increased invasiveness and metastasis in glioblastoma. It can promote the epithelial-mesenchymal transition (EMT), a process associated with increased invasiveness and metastatic potential.

TAMs in WT modulate lymphocytes and NK cells through cell-to-cell contact.

T cells in glioblastoma which are cytotoxic express the NK Cell gene KLRB1. We found that most of the TAMs express the ligand CLEC2D on the surface which serves as an inhibitor for KLRB1 receptor[30].

TAMs are associated with T cell and NK cell exhaustion in the TME

 T lymphocytes exhaustion is severe in glioblastoma and the cells express several checkpoint receptors including PD-1, LAG-3 and CTLA -4[31]. The binding of TIM3 to its ligand, galectin-9 induces T cell apoptosis[32]. Indeed, we found complete absence of T lymphocytes in the neighborhood of Angio TAMs using spatial transcriptomics. The spatial distribution of TAMs within the glioblastoma tumor is not well-defined. Understanding their localization and interactions with tumor cells and immune cells in specific microenvironmental niches could provide valuable insights into their functional roles and immune-suppressive mechanisms.

TAMs interact with various immune cells within the tumor microenvironment, including CTLs. However, the precise mechanisms by which TAMs suppress the function of CTLs and evade their cytotoxic activity are not fully elucidated. TAMs in glioblastoma can suppress T lymphocytes, including cytotoxic T lymphocytes (CTLs), through various mechanisms. These include expression of Indoleamine 2,3-dioxygenase (IDO)which metabolizes tryptophan, an essential amino acid for T cell function. By depleting tryptophan, TAMs create an immunosuppressive environment that impairs CTL activity and proliferation. TAMs can produce Arg1, an enzyme that competes with CTLs for arginine, another essential amino acid. Reduced arginine availability hampers the function and proliferation of CTLs, weakening their cytotoxicity against tumor cells. TAMs can release nitric oxide, a molecule with immunosuppressive properties. Elevated levels of NO can inhibit T cell signaling and proliferation, impairing CTL function within the tumor microenvironment. TAMs can secrete various cytokines, such as interleukin-10 (IL-10) and transforming growth factor-beta (TGF-β), which promote an immunosuppressive milieu. These cytokines can inhibit T cell activation, dampen immune responses, and induce regulatory T cell (Treg) differentiation, further suppressing CTL activity. TAMs also express immune checkpoint molecules, such as programmed cell death-ligand 1 (PD-L1), which interacts with programmed cell death protein 1 (PD-1) on T cells. This interaction leads to T cell exhaustion and functional impairment, limiting the cytotoxicity of CTLs against tumor cells.

Future directions

The immune landscape in glioblastoma is complex and most importantly not static. It evolves along with the progression of tumor. Immunosuppressive macrophages dominate towards the later stages of glioma evolution. We have identified several subsets of macrophages among which angiogenic TAMs play a very significant role. The immunosuppressive molecules of galectin, SPP1, MIF and complement impede the T cell response.

Immunotherapeutic agents including checkpoint inhibitors and CAR T cell therapies have not shown encouraging utility in GBM, which is consistent with the immunologically 'cold' nature of GBM implying the microenvironment is pervaded by immunosuppressive TAMs [33, 34]. As these TAMs prevent T cell permeation, the possibility of re-educating these TAMs into a pro-inflammatory state is critical to explore if future immunotherapies should hope to have any substantial effect in outcomes.

Figure Legends

Figure 1
(A) Scatterplot displaying dominant senders and receiver cell type. The total outgoing and incoming communication probability associated with each cell group is displayed on x-axis or y-axis, respectively. Count is displayed by dot size which is proportional to the number of predicted signaling links associated with each cell group.
(B) Stacked bar plot and unstacked bar plot **(C)** showing conserved and context specific signaling pathways by comparing information flow for each signaling pathway.
(D) Scatterplot showing specific signaling changes of TAM populations. Positive values indicate increase in IDH wildtype GBM compared to IDH mutant astrocytoma.

Figure 2
(A) Chord diagram visualizing cell-cell communication for significant ligand-receptor interactions from TAM populations (senders, bottom arc) to T cell populations (receivers, top arc).
(B) Chord diagram visualizing significant ligand-receptor interactions from T cell populations (senders, bottom arc) to TAM populations (receivers, top arc).

Figure 3
Bubble plot showing all the significant interactions (L-R pairs) (y-axis) from TAM populations (senders) to T cell populations (receivers) (x-axis). The dot color and size represent the calculated communication probability (represents interaction strength) and p-values (represents statistical significance). Non-significant interactions represented by blank spaces.
(A) Abdelfattah et al., 2022 dataset.
(B) Additional independent analysis using Pombo Antunes et al., 2021 dataset.

Figure 4
Global communications analysis for **(A)** outgoing and **(B)** incoming communication patterns.
Cell patterns heatmap display which cell types contribute to which type of pattern. Communication patterns heatmap displays which signaling pathways contribute to the pattern. The dot plots show how each cell group contributed to the signaling pathways, where the dot size is proportional to the contribution score to show association between cell group and their enriched signaling pathways. A higher contribution score implies the signaling pathway is more enriched in the respective cell type.

Figure 5
Surface plot displaying the expression signatures of the T cell exhaustion cell state and seven TAM types derived from spatial transcriptomics. Brighter areas correspond to higher average expression of genes comprising the signature.

References

1. Alexander BM, Cloughesy TF (2017) Adult Glioblastoma. J Clin Oncol 35:2402–2409. https://doi.org/10.1200/JCO.2017.73.0119

2. Andersen JK, Miletic H, Hossain JA (2022) Tumor-Associated Macrophages in Gliomas—Basic Insights and Treatment Opportunities. Cancers 14:1319. https://doi.org/10.3390/cancers14051319

3. Wlodarczyk A, Holtman IR, Krueger M, Yogev N, Bruttger J, Khorooshi R, Benmamar□Badel A, de Boer□Bergsma JJ, Martin NA, Karram K, Kramer I, Boddeke EW, Waisman A, Eggen BJ, Owens T (2017) A novel microglial subset plays a key role in myelinogenesis in developing brain. EMBO J 36:3292–3308. https://doi.org/10.15252/embj.201696056

4. Hoeffel G, Ginhoux F (2015) Ontogeny of Tissue-Resident Macrophages. Front Immunol 6:486. https://doi.org/10.3389/fimmu.2015.00486

5. Murray PJ, Wynn TA (2011) Protective and pathogenic functions of macrophage subsets. Nat Rev Immunol 11:723–737. https://doi.org/10.1038/nri3073

6. Hambardzumyan D, Gutmann DH, Kettenmann H (2016) The role of microglia and macrophages in glioma maintenance and progression. Nat Neurosci 19:20–27. https://doi.org/10.1038/nn.4185

7. Woroniecka KI, Rhodin KE, Chongsathidkiet P, Keith KA, Fecci PE (2018) T-cell Dysfunction in Glioblastoma: Applying a New Framework. Clin Cancer Res Off J Am Assoc Cancer Res 24:3792–3802. https://doi.org/10.1158/1078-0432.CCR-18-0047

8. Xue J, Schmidt SV, Sander J, Draffehn A, Krebs W, Quester I, De Nardo D, Gohel TD, Emde M, Schmidleithner L, Ganesan H, Nino-Castro A, Mallmann MR, Labzin L, Theis H, Kraut M, Beyer M, Latz E, Freeman TC, Ulas T, Schultze JL (2014) Transcriptome-Based Network Analysis Reveals a Spectrum Model of Human Macrophage Activation. Immunity 40:274–288. https://doi.org/10.1016/j.immuni.2014.01.006

9. Szulzewsky F, Arora S, de Witte L, Ulas T, Markovic D, Schultze JL, Holland EC, Synowitz M, Wolf SA, Kettenmann H (2016) Human glioblastoma-associated microglia/monocytes express a distinct RNA profile compared to human control and murine samples. Glia 64:1416–1436. https://doi.org/10.1002/glia.23014

10. Ma R-Y, Black A, Qian B-Z (2022) Macrophage diversity in cancer revisited in the era of single-cell omics. Trends Immunol 43:546–563. https://doi.org/10.1016/j.it.2022.04.008

11. Abdelfattah N, Kumar P, Wang C, Leu J-S, Flynn WF, Gao R, Baskin DS, Pichumani K, Ijare OB, Wood SL, Powell SZ, Haviland DL, Parker Kerrigan BC, Lang FF, Prabhu SS, Huntoon KM, Jiang W, Kim BYS, George J, Yun K (2022) Single-cell analysis of human glioma and immune cells identifies S100A4 as an immunotherapy target. Nat Commun 13:767. https://doi.org/10.1038/s41467-022-28372-y

12. Pombo Antunes AR, Scheyltjens I, Lodi F, Messiaen J, Antoranz A, Duerinck J, Kancheva D, Martens L, De Vlaminck K, Van Hove H, Kjølner Hansen SS, Bosisio FM, Van der Borght K, De Vleeschouwer S, Sciot R, Bouwens L, Verfaillie M, Vandamme N, Vandenbroucke RE, De Wever O, Saeys Y, Guilliams M, Gysemans C, Neyns B, De Smet F, Lambrechts D, Van Ginderachter JA, Movahedi K (2021) Single-cell profiling of myeloid cells in glioblastoma across species and disease stage reveals macrophage competition and specialization. Nat Neurosci 24:595–610. https://doi.org/10.1038/s41593-020-00789-y

13. Ravi VM, Will P, Kueckelhaus J, Sun N, Joseph K, Salié H, Vollmer L, Kuliesiute U, von Ehr J, Benotmane JK, Neidert N, Follo M, Scherer F, Goeldner JM, Behringer SP, Franco P, Khiat M, Zhang J, Hofmann UG, Fung C, Ricklefs FL, Lamszus K, Boerries M, Ku M, Beck J, Sankowski R, Schwabenland M, Prinz M, Schüller U, Killmer S, Bengsch B, Walch AK, Delev D, Schnell O, Heiland DH (2022) Spatially resolved multi-omics deciphers bidirectional tumor-host interdependence in glioblastoma. Cancer Cell 40:639-655.e13. https://doi.org/10.1016/j.ccell.2022.05.009

14. Monaco G, Lee B, Xu W, Mustafah S, Hwang YY, Carré C, Burdin N, Visan L, Ceccarelli M, Poidinger M, Zippelius A, Pedro De Magalhães J, Larbi A (2019) RNA-Seq Signatures Normalized by mRNA Abundance Allow Absolute Deconvolution of Human Immune Cell Types. Cell Rep 26:1627-1640.e7. https://doi.org/10.1016/j.celrep.2019.01.041

15. Jin S, Guerrero-Juarez CF, Zhang L, Chang I, Ramos R, Kuan C-H, Myung P, Plikus MV, Nie Q (2021) Inference and analysis of cell-cell communication using CellChat. Nat Commun 12:1088. https://doi.org/10.1038/s41467-021-21246-9

16. Cordeiro M, Sarmento RP, Brazdil P, Gama J (2018) Evolving Networks and Social Network Analysis Methods and Techniques. In: Višňovský J, Radošinská J (eds) Social Media and Journalism - Trends, Connections, Implications. InTech

17. Landherr A, Friedl B, Heidemann J (2010) A Critical Review of Centrality Measures in Social Networks. Bus Inf Syst Eng 2:371–385. https://doi.org/10.1007/s12599-010-0127-3

18. Gaujoux R, Seoighe C (2010) A flexible R package for nonnegative matrix factorization. BMC Bioinformatics 11:367. https://doi.org/10.1186/1471-2105-11-367

19. Watowich MB, Gilbert MR, Larion M (2023) T cell exhaustion in malignant gliomas. Trends Cancer 9:270–292. https://doi.org/10.1016/j.trecan.2022.12.008

20. Klemm F, Maas RR, Bowman RL, Kornete M, Soukup K, Nassiri S, Brouland J-P, Iacobuzio-Donahue CA, Brennan C, Tabar V, Gutin PH, Daniel RT, Hegi ME, Joyce JA (2020) Interrogation of the Microenvironmental Landscape in Brain Tumors Reveals Disease-Specific Alterations of Immune Cells. Cell 181:1643-1660.e17. https://doi.org/10.1016/j.cell.2020.05.007

21. Casanova-Acebes M, Dalla E, Leader AM, LeBerichel J, Nikolic J, Morales BM, Brown M, Chang C, Troncoso L, Chen ST, Sastre-Perona A, Park MD, Tabachnikova A, Dhainaut M, Hamon P, Maier B, Sawai CM, Agulló-Pascual E, Schober M, Brown BD, Reizis B, Marron T, Kenigsberg E, Moussion C, Benaroch P, Aguirre-Ghiso JA, Merad M (2021) Tissue-resident macrophages provide a pro-tumorigenic niche to early NSCLC cells. Nature 595:578–584. https://doi.org/10.1038/s41586-021-03651-8

22. O'Neill LAJ, Kishton RJ, Rathmell J (2016) A guide to immunometabolism for immunologists. Nat Rev Immunol 16:553–565. https://doi.org/10.1038/nri.2016.70

23. Venteicher AS, Tirosh I, Hebert C, Yizhak K, Neftel C, Filbin MG, Hovestadt V, Escalante LE, Shaw ML, Rodman C, Gillespie SM, Dionne D, Luo CC, Ravichandran H, Mylvaganam R, Mount C, Onozato ML, Nahed BV, Wakimoto H, Curry WT, Iafrate AJ, Rivera MN, Frosch MP, Golub TR, Brastianos PK, Getz G, Patel AP, Monje M, Cahill DP, Rozenblatt-Rosen O, Louis DN, Bernstein BE, Regev A, Suvà ML (2017) Decoupling genetics, lineages, and microenvironment in IDH-mutant gliomas by single-cell RNA-seq. Science 355:eaai8478. https://doi.org/10.1126/science.aai8478

24. Yeo AT, Rawal S, Delcuze B, Christofides A, Atayde A, Strauss L, Balaj L, Rogers VA, Uhlmann EJ, Varma H, Carter BS, Boussiotis VA, Charest A (2022) Single-cell RNA sequencing reveals evolution of immune landscape during glioblastoma progression. Nat Immunol 23:971–984. https://doi.org/10.1038/s41590-022-01215-0

25. Ajarrag S, St-Pierre Y (2021) Galectins in Glioma: Current Roles in Cancer Progression and Future Directions for Improving Treatment. Cancers 13:5533. https://doi.org/10.3390/cancers13215533

26. Klement JD, Paschall AV, Redd PS, Ibrahim ML, Lu C, Yang D, Celis E, Abrams SI, Ozato K, Liu K An osteopontin/CD44 immune checkpoint controls CD8+ T cell activation and tumor immune evasion. J Clin Invest 128:5549–5560. https://doi.org/10.1172/JCI123360

27. Yan X, Orentas RJ, Johnson BD (2006) Tumor-derived macrophage migration inhibitory factor (MIF) inhibits T lymphocyte activation. Cytokine 33:188–198. https://doi.org/10.1016/j.cyto.2006.01.006

28. Regulation of the CTL Response by Macrophage Migration Inhibitory Factor | The Journal of Immunology | American Association of Immunologists. https://journals.aai.org/jimmunol/article/166/2/747/70562/Regulation-of-the-CTL-Response-by-Macrophage. Accessed 14 Aug 2023

29. Roumenina LT, Daugan MV, Noé R, Petitprez F, Vano YA, Sanchez-Salas R, Becht E, Meilleroux J, Clec'h BL, Giraldo NA, Merle NS, Sun C-M, Verkarre V, Validire P, Selves J, Lacroix L, Delfour O, Vandenberghe I, Thuilliez C, Keddani S, Sakhi IB, Barret E, Ferré P, Corvaïa N, Passioukov A, Chetaille E, Botto M, de Reynies A, Oudard SM, Mejean A, Cathelineau X, Sautès-Fridman C, Fridman WH (2019) Tumor Cells Hijack Macrophage-Produced Complement C1q to Promote Tumor Growth. Cancer Immunol Res 7:1091–1105. https://doi.org/10.1158/2326-6066.CIR-18-0891

30. Mathewson ND, Ashenberg O, Tirosh I, Gritsch S, Perez EM, Marx S, Jerby-Arnon L, Chanoch-Myers R, Hara T, Richman AR, Ito Y, Pyrdol J, Friedrich M, Schumann K, Poitras MJ, Gokhale PC, Castro LNG, Shore ME, Hebert CM, Shaw B, Cahill HL, Drummond M, Zhang W, Olawoyin O, Wakimoto H, Rozenblatt-Rosen O, Brastianos PK, Liu XS, Jones PS, Cahill DP, Frosch MP, Louis DN, Freeman GJ, Ligon KL, Marson A, Chiocca EA, Reardon DA, Regev A, Suvà ML, Wucherpfennig KW (2021) Inhibitory CD161 Receptor Identified in Glioma-infiltrating T cells by Single Cell Analysis. Cell 184:1281-1298.e26. https://doi.org/10.1016/j.cell.2021.01.022

31. Woroniecka K, Chongsathidkiet P, Rhodin K, Kemeny H, Dechant C, Farber SH, Elsamadicy AA, Cui X, Koyama S, Jackson C, Hansen LJ, Johanns TM, Sanchez-Perez L, Chandramohan V, Yu Y-RA, Bigner DD, Giles A, Healy P, Dranoff G, Weinhold KJ, Dunn GP, Fecci PE (2018) T Cell Exhaustion Signatures Vary with Tumor Type and are Severe in Glioblastoma. Clin Cancer Res Off J Am Assoc Cancer Res 24:4175–4186. https://doi.org/10.1158/1078-0432.CCR-17-1846

32. Sabatos CA, Chakravarti S, Cha E, Schubart A, Sánchez-Fueyo A, Zheng XX, Coyle AJ, Strom TB, Freeman GJ, Kuchroo VK (2003) Interaction of Tim-3 and Tim-3 ligand regulates T helper type 1 responses and induction of peripheral tolerance. Nat Immunol 4:1102–1110. https://doi.org/10.1038/ni988

33. Frederico SC, Hancock JC, Brettschneider EES, Ratnam NM, Gilbert MR, Terabe M (2021) Making a Cold Tumor Hot: The Role of Vaccines in the Treatment of Glioblastoma. Front Oncol 11:672508. https://doi.org/10.3389/fonc.2021.672508

34. Land CA, Musich PR, Haydar D, Krenciute G, Xie Q (2020) Chimeric antigen receptor T-cell therapy in glioblastoma: charging the T cells to fight. J Transl Med 18:428. https://doi.org/10.1186/s12967-020-02598-0