Intra-leaf modeling of *Cannabis* leaflet shape produces synthetic leaves that predict

genetic and developmental identities

Manica Balant [1,2,3], Teresa Garnatje [1], Daniel Vitales [1,2], Oriane Hidalgo [1,4], Daniel H.

Chitwood [3,5]

[1] Institut Botànic de Barcelona, IBB (CSIC-Ajuntament de Barcelona), Passeig del Migdia s.n.,

08038 Barcelona, Spain

[2] Laboratori de Botànica (UB), Unitat Associada al CSIC, Facultat de Farmàcia i Ciències de

l'Alimentació–Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona, Av.

Joan XXIII 27–31, 08028 Barcelona, Spain

[3] Department of Horticulture, Michigan State University, East Lansing, MI, USA

[4] Royal Botanic Gardens, Kew, Kew Green, Richmond TW9 3AE, UK

[5] Department of Computational Mathematics, Science & Engineering, Michigan State

University, East Lansing, MI, USA

Authors for correspondence:

Daniel H. Chitwood

Email: dhchitwood@gmail.com

Manica Balant

Email: manica.balant@ibb.csic.es

Summary

●   The iconic, palmately compound leaves of *Cannabis* have attracted significant

attention in the past. However, investigations into the genetic basis of leaf shape or

its connections to phytochemical composition have yielded inconclusive results. This

is partly due to prominent changes in leaflet number between leaves of a single

plant during development, which has until this point prevented the proper use of

common morphometric techniques.

●   Here we present a new method that overcomes the challenge of nonhomologous

landmarks in *Cannabis*. By modeling leaflet shape as a function of angle within a leaf,

we create synthetic leaves with comparable numbers of leaflets to which
morphometric analysis is applied.

● We analyze 339 leaves from 24 individuals from nine *Cannabis* accessions, including
both wild/feral accessions and cultivated varieties. Using 1,995 pseudo-landmarks in
synthetic leaves, we visualize a morphospace and accurately predict accession
identity, leaflet number, and relative node number.

● Intra-leaf modeling offers a rapid, cost-effective means of identifying *Cannabis*
accessions, making it a valuable tool for future taxonomic studies, cultivar
recognition, and possibly chemical content analysis and sex identification, in addition
to permitting the morphometric analysis of leaves in any species with variable
numbers of leaflets or lobes.

Key words: *Cannabis*, development, geometric morphometrics, intra-leaf modeling, leaf
morphology, leaf shape

INTRODUCTION

Cannabis sativa L. (hereafter referred to as *Cannabis*) is a versatile crop plant used by
humans for a variety of purposes throughout history. Although today it is commonly
associated with its psychoactive properties, traditional medicine has relied heavily on
Cannabis, and it is also a valuable source of food and fibers (Clarke & Merlin, 2013). Genetic
and archaeological evidence suggests that *Cannabis* was domesticated around 12,000 years
ago in East Asia, initially serving as a multipurpose crop before separate selections for fiber
and drug production emerged around 4,000 years ago (Ren *et al.*, 2021). Since then,
widespread cultivation has facilitated its global distribution. Throughout the 20th century,
Cannabis use was largely abandoned due to its illegal status in many parts of the world.
However, recent legalization for recreational and/or medicinal purposes in many countries
worldwide has led to a surge in the cannabis industry (*The Global Cannabis Report, 3rd*
Edition, 2022).

Extensive *Cannabis* use has resulted in the development of numerous cultivars and strains
that are well-suited to diverse uses and climates (Small, 2015). This significant

morphological and phytochemical diversity within the *Cannabis* genus poses challenges for

taxonomic classification. Over the past two centuries, various taxonomic approaches based

on genetics, morphology, and phytochemistry have been proposed (McPartland & Small,

2020). Some scientists advocated for a polytypic classification, recognizing the presence of

two (Lamarck & Poiret, 1783; Zhukovskii, 1971; Hillig, 2005a) or three (Emboden, 1974;

Schultes *et al.*, 1974; Hillig, 2005b; Clarke & Merlin, 2013) species with multiple subspecies,

while others argued for a monotypic genus, considering only a single species, *Cannabis*

sativa (Small & Cronquist, 1976; Sawler *et al.*, 2015; Small, 2015; McPartland, 2018;

McPartland & Small, 2020; Ren *et al.*, 2021). Hilling introduced a classification system based

on biotypes, considering molecular, morphological, and phytochemical data. He proposed

dividing *Cannabis* into two species, *C. sativa* and *C. indica* Lam., and six biotypes: *C. indica* as

narrow-leaflet drug (NLD), wide-leaflet drug (WLD), hemp and feral biotype, and *C. sativa* as

hemp and feral biotype (Hillig, 2005a). Recently, Lapierre *et al.* (2023) conducted a

comprehensive taxonomic review of the *Cannabis* genus and based on available genetic

data, strongly supported the theory that *Cannabis* is a highly diverse monotypic species.

Apart from taxonomic classification, *Cannabis* is often categorized based on its cultivation

purpose, morphology, and chemical composition. Fiber-type plants, commonly known as

hemp, are primarily grown for fiber and seed production. These plants contain less than

0.3% of the psychoactive compound THC (Δ9-tetrahydrocannabinol), while drug-type plants,

often referred to as marijuana and medicinal cannabis, can contain higher levels of THC

(Hurgobin *et al.*, 2021). *Cannabis* plants can also be separated based on the ratio of two

major cannabinoids THC and CBD (cannabidiol) into Type I (THC dominant), Type II

(balanced CBD/THC ratio), and Type III plants (CBD dominant) (Small & Beckstead, 1973). In

the medicinal and recreational cannabis industries plants are normally categorized as

'*sativa*', '*indica*', or '*hybrid*'. Taller plants with narrow leaflets and high THC percentage are

called 's*ativa*', while shorter and bushier plants with wider leaflets and high percentages of

both CBD and THC are called '*indica*'. Plants with intermediate characters are called '*hybrids*'

(McPartland & Guy, 2017). While the classification of *Cannabis* into *'indica'* and *'sativa'* is

not supported by genetic data, the visible differences in leaflet width have long been a

significant characteristic used to visually discriminate different types of *Cannabis*.

Cannabis arguably possesses one of the most iconic leaves among all plants. Its palmately

compound leaves with a varying number of leaflets are a popular culture symbol. *Cannabis*

exhibits a remarkable degree of phenotypic plasticity, further accentuated by selection

pressure during the domestication process (Small, 2015). Extensive variability in leaf

morphology has already been described by Quimby *et al.* (1973) and later Anderson (1980),

who was the first to quantify the width, length, and ratio of the central leaflet. This or

similar methods were then commonly used in studies investigating the morphological

characteristics of *Cannabis* species, subspecies, cultivars, biotypes and chemotypes (Small *et*

al., 1976; de Meijer *et al.*, 1992; de Meijer & Keizer, 1996; Hillig, 2005a; Clarke & Merlin,

2013; Lynch *et al.*, 2016; Karlov *et al.*, 2017; Parsons *et al.*, 2019; McPartland & Small, 2020;

Carlson *et al.*, 2021; Islam *et al.*, 2021; Jin *et al.*, 2021a; Vergara *et al.*, 2021; Buzna & Sala,

2022; Chen *et al.*, 2022; Murovec *et al.*, 2022), often with contradictory results. Leaf shape

has therefore played an important and sometimes controversial role in *Cannabis* taxonomy.

While researchers in previous *Cannabis* studies were aware of enormous plasticity and the

effect the environment has on leaf shape (Vergara *et al.*, 2021; Murovec *et al.*, 2022), they

very rarely paid attention to the effects of developmental processes, even though

heteroblastic changes (differences in leaf shape arising from juvenile-to-adult phase

transitions in the meristem) profoundly affect the arrangement and shape of *Cannabis*

leaves along the shoot. While some studies briefly mention the developmental changes of

leaves (Hillig, 2005a; Carlson *et al.*, 2021; Jin *et al.*, 2021a; Spitzer-Rimon *et al.*, 2022), the

only study focused on heteroblastic phase changes in leaves along the plant axis was done

by Heslop-Harrison and Heslop-Harrison (1958). In the lower part of the shoot *Cannabis*

leaves exhibit opposite phyllotaxy and one to three leaflets, transitioning to alternate

phyllotaxy and leaves with up to 11 to 13 leaflets in the upper section (Hillig, 2005a; Clarke

& Merlin, 2013; Small, 2015). Additionally, the changes in leaflet number are not uniform

between different *Cannabis* accessions (Hillig, 2005a). These changes during development

not only complicate categorization of plant accessions based on leaf shape, but also prevent

the use of morphometric techniques.

Morphometrics is the quantitative analysis of shape. It includes a wide range of methods,

from measuring allometric differences in dimensions like lengths, widths, and angles in

relation to size (Niklas, 1994), to geometric techniques that measure shape

comprehensively, like elliptical Fourier (EFDs; Kuhl & Giardina, 1982) and landmark-based

analyses (Bookstein, 1997). It can be used to classify species and to separate effects on

shape arising from genetic, developmental, and environmental mechanisms (Chitwood &

Sinha, 2016). Historically the field of ampelography (ἄμπελος, 'vine' + γράφος, 'writing';

Ravaz, 1902; Galet, 1952; Galet & trans. Morton, 1979) relied heavily on leaf shape to

distinguish grapevine varieties, which like *Cannabis*, is a culturally popular plant used for

psychoactive, recreational purposes with a charismatic leaf shape. Unlike *Cannabis,*

grapevine leaves have a consistent number of lobes, sinuses, and other associated

homologous points that can be used for both landmark-based and EFD morphometric

analysis (Chitwood *et al.*, 2014; Chitwood, 2021) to disentangle genetic (Demmings *et al.*,

2019), developmental (Chitwood *et al.*, 2016a; Bryson *et al.*, 2020; Migicovsky *et al.*, 2022),

and environmental effects (Chitwood *et al.*, 2016b, 2021) embedded in leaf shapes. The

variable number of leaflets in *Cannabis* leaves precludes analysis methods that rely on

homologous, comparable points to measure shape comprehensively. Vergara *et al.* (2021)

used a landmark-based approach but were limited to analyzing the central and two most

distal leaflets on each side, features that all *Cannabis* leaves except single-leaflet leaves

possess, but which excludes most of the shape variation within a leaf.

Here, we model leaflet shape as a function of angle within *Cannabis* leaves. The resulting

continuous models can be used to create synthetic leaves with any desired number of

leaflets. Analyzing 339 leaves from across the nodes of 24 individuals representing nine

accessions, we create synthetic leaves with five leaflets and 1,995 comparable pseudo-

landmarks. Dimension reduction separates leaf types along orthogonal axes and visualizes

an underlying morphospace. Linear discriminant analysis (LDA) predicts accession, leaflet

number, and relative node number with high accuracy. Intra-leaf modeling allows the

application of morphometric techniques to comprehensively measure leaf shape in

Cannabis, enabling future taxonomic and developmental studies, cultivar recognition, and

possibly chemical content analysis and sex identification, in addition to permitting the

morphometric analysis of leaves in any species with variable numbers of leaflets or lobes.

MATERIAL AND METHODS

Plant material and growing conditions

This study includes 24 individuals from nine accessions of *Cannabis sativa* (Table **1**; Fig. **1**),

encompassing both wild/feral accessions and cultivated varieties with a wide distribution

area. The plants were grown from seed in a growth chamber (Fitoclima D1200PLL, Aralab,

Portugal) to minimize the influence of the environment. Before sowing, the seeds were

sterilized overnight in a 5% H_2O_2 solution with the addition of Inex-A solution (Cosmocel,

Spain) at room temperature. Sterilized seeds were then transferred to Petri dishes and

placed in the growth chamber for germination. Once the first leaves emerged, the seedlings

were transferred to small peat pots with a pre-fertilized soil substrate (Kilomix Atami,

Spain). During this phase, the environmental conditions were set to 25°C, with an 18-hour

day and 6-hour night photoperiod, and a light intensity of 50 µmol $m^{-2}s^{-1}$ (Philips Master PL-

L 55W, Spain). After two weeks the surviving plants were transplanted to 3.5 l pots with the

same soil substrate. The light intensity was gradually increased to 300 µmol $m^{-2}s^{-1}$ over the

following week, without changing the photoperiod and temperature. After four weeks, the

photoperiod was changed to 12 hours of daylight and 12 hours of darkness, and the light

intensity was gradually increased to 700 µmol $m^{-2}s^{-1}$ over the following week, while keeping

the temperature at 25°C. The plants remained in these environmental conditions until the

flowering stage. Plants received daily irrigation with tap water, without any application of

nutrient or phytosanitary control.

Table 1 Accession details and number of leaves collected and analyzed in the study.

Accession ID	Accession type	Location/Cultivar name	Number of individuals	Number of leaves collected	Number of leaves analyzed
AM15	Wild/feral	Armenia, Sjunik marz, Goris town	5	90	72
BNG	Wild/feral	Bangladesh, Rangpur, Carmichael College Campus	1	14	9
FUT75	Cultivar	Futura 75	2	44	30
HU1	Wild/feral	Hungary, Nyírvasvári	4	83	65
IK	Landrace	India, Kerala	4	99	53
IKL	Landrace	India, Kullu	4	69	59
MAR	Landrace	Morocco, North Morocco	1	18	11
MN9	Wild/feral	Mongolia, Selenge aimag, Baruunburen sum	1	14	12
RO1	Wild/feral	Romania, Mangalija	2	36	28

Leaf sampling and imaging

A total of 467 leaves were sampled during the flowering stage, with the exception of

individuals from the accession IK, which did not begin to flower during the two-month

cultivation period. Leaves along the main axis of the plants were collected and immediately

scanned using a flatbed photograph scanner (Epson Perfection V370, Tokyo, Japan) at 1200

dpi resolution. A piece of velvet fabric was placed between the leaf and the scanner cover to

avoid any shadows. No adjustments to the angle of individual leaflets were made before

scanning. Each leaf was scanned with a scale and a label indicating the node it originated

from, followed by a sequential lowercase letter, since typically two leaves are present per

node. Starting at the base of the plant, the first two leaves were labeled as leaves "a" and

"b" from node number 1, and so on, until the shoot apex.

Cannabis leaves display a marked heteroblastic, or juvenile-to-adult, leaf shape progression.

Mature, juvenile leaves located on the first node at the base of the plant usually have a

simple, serrated leaf. As node number increases so does the leaflet number, reaching a

maximum of 9 to 13 leaflets in young, adult leaves at the growing tip. Eventually leaves

transition into an inflorescence type. During this transition, the number of leaflets per leaf

starts to decrease again until the top of the inflorescence. Leaves at the shoot base have

opposite phyllotaxy and transition to alternate phyllotaxy in the upper section on the stem

and inflorescence (Heslop-Harrison & Heslop-Harrison, 1958; Hillig, 2004; Potter, 2009;

Spitzer-Rimon *et al.*, 2022). To ensure that only stem leaves were included in our analysis,

we separated the two types (i.e., stem and inflorescence leaves) based on the point where

the decrease in the number of leaflets appeared. This point determined the "total node

number", the number of nodes per plant used for further analysis. Total node number

varied among individuals. To compare node positions, a relative node number was

calculated, which was defined by the node position divided by the total node number for

the individual plant, where zero is at the plant base and one at the last node included in the

analysis (Fig. **1**). Because of the nature of plant growth, the leaves at the base of the plant

were frequently too senesced to be incorporated in the analysis or were entirely lost.

Nevertheless, the nodes could still be identified, which allowed them to be taken into

account in the calculation of relative node number.

Fig. 1 Changes in the leaf shape and leaflet number during the development in nine *Cannabis* accessions. (a) Median values for all available leaflet number for each relative node number for the nine *Cannabis* accessions. (b) Changes in leaf shape between different developmental stages in different *Cannabis* accessions.

Image analysis and landmarking

After eliminating deformed leaves, simple leaves, leaves with even leaflet numbers, leaves with relative node values above one, inaccurately modeled leaves and outliers detected during dimension reduction, a total of 339 *Cannabis* leaves were analyzed. Photoshop was

used to separate petioles and leaflets smaller than 1 cm from the rest of the leaf. The leaf

outlines were then extracted and saved using Python modules NumPy (Harris *et al.*, 2020),

Matplotlib (Hunter, 2007) and OpenCV (Bradski, 2000). The code for extracting and plotting

the   leaf   outlines   can   be   found   on   GitHub

(https://github.com/BalantM/Cannabis_leaf_morpho). The *x* and *y* coordinates of blade

outlines and landmarks were extracted using ImageJ (Abràmoff *et al.*, 2004). The outline

was extracted using the *wand* tool (setting tolerance to 20 and including "smooth if

thresholded" option) and the landmarks were placed using the *multi-point* tool. Initially,

landmarks were placed at the beginning and end of each leaflet, starting from the lower left

side, and continuing to the lower right side of the leaf outline. Subsequently, landmarks

were placed in the same order on the tips of the leaflets. The final landmark was positioned

at the center of the petiolar junction (Fig. **2**). These landmarks delimit the boundaries of the

leaflets so that equidistant pseudo-landmarks can later be placed along the contour. The

number of landmarks per leaf ranged from 10 to 28, depending on the leaflet number. The

raw data containing the coordinates for leaf outlines and landmarks can be accessed on

GitHub (https://github.com/BalantM/Cannabis_leaf_morpho).

Fig. 2 The process of modeling new synthetic leaves for a leaf with (a) three leaflets from

accession AM15, (b) five leaflets from accession IKL, (c) seven leaflets from accession FUT75,

and (d) nine leaflets from accession IK. The first column shows the scans of the leaves, which

we use to extract the outline and place the landmarks on the tip, start, and end of each

leaflet and on the petiolar junction (second column). These coordinates are used to

generate 200 pseudo-landmarks on each side of each leaflet, sharing the landmark on the

tip of the leaflet and are converted into polar coordinates. Second degree polynomials are

then fitted through these pseudo-landmarks (third column). A synthetic leaf with five

leaflets defined by 1,995 pseudo-landmarks can then be modeled using the collection of

polynomial models for each leaf (fourth column).

Data analysis

To analyze leaves with different numbers of leaflets, pseudo-landmarks of each leaflet were

modeled as polynomial functions of angles within a leaf, in order to use the models to

construct a synthetic leaf with a desired number of leaflets. The Python code, presented as a

Jupyter   notebook   with   detailed   description,   is   available   on   GitHub

(https://github.com/BalantM/Cannabis_leaf_morpho). In brief, *x* and *y* coordinates of the

leaf outline were first interpolated to create an arbitrarily high number of coordinates to

increase resolution. The coordinates of manually selected landmarks were then compared

against the high-resolution coordinates of the leaf outline and the nearest neighboring point

of the high-resolution coordinates to each original landmark was identified and specified as

the new landmark point. Next, the outline and new landmark coordinates were rotated,

translated, and scaled so that the central leaflet had a length of one and pointed in the

same direction. The transformed points were then interpolated to generate 200 pseudo-

landmarks on each side of each leaflet, sharing the landmark on the tip of the leaflet (i.e., a

total of 399 pseudo-landmarks per leaflet). These pseudo-landmarks were then converted

to polar coordinates, where each point was defined by a radius and angle relative to the

petiolar junction and tip of the central leaflet.

Using the polar coordinates of each leaflet, a second-degree polynomial function using the

NumPy *polyfit* function was fitted through each of the corresponding pseudo-landmarks for

each leaflet, modeling radius as a function of angle. A synthetic leaf with five leaflets was

then modeled using the collection of 399 polynomial models for each leaf. The coordinates

defining the 1,995 pseudo-landmarks of the synthetic five-leaflet leaves were plotted and

visually inspected. Any inaccurately modeled leaves or plots containing noticeable errors or

discrepancies were eliminated. A principal component analysis (PCA) was performed using

scikit-learn (Pedregosa *et al.*, 2011) module in Python and proportions of explained variance

for each principal component and the cumulative variance was calculated. The

morphospace was calculated using the inverse transform of PC1 and PC2 scores and

visualized using seaborn (Waskom, 2021) and Matplotlib modules in Python. Theoretical

eigenleaves were plotted across equal intervals of the PCA morphospace. Points

representing the leaves were then plotted over and colored by the accession identity, leaflet

number, or relative node number (Fig. **3, 4**). To see which of the first two PCs explains most

of the leaf shape variation for accessions, leaflet number and relative node number Kruskal-

Wallis test were calculated using *stats.kruskal* function from the scipy.stats module

(Virtanen *et al.*, 2020). To visualize an average leaf for each accession, leaflet number, and

relative node number, the average coordinate values of synthetic leaves were calculated for

each of the categories and plotted using the Matplotlib module in Python.

Linear discriminant analysis (LDA) was applied to model accession, leaflet number, and

relative node number as the function of synthetic leaf coordinate values (Fig. **5**) using the

LinearDiscriminantAnalysis function from the scikit-learn module in Python. To test the

performance of the LDA model, the dataset was divided into two parts. Since most of the

analyzed leaves exhibit opposite phyllotaxy, wherein the nodes were represented by two

leaves (a and b) in the same developmental phase with the same number of leaflets, the

dataset was split into a training dataset (leaf a) comprising 177 leaves and a test dataset

(leaf b) containing 162 leaves. The *predict* function from *LinearDiscriminantAnalysis* in the

scikit-learn module was used to predict the accession identity, leaflet number, and relative

node number. The accuracy of the LDA model was calculated and visualized using the

function *confusion_matrix* from scikit-learn. Spearman Rank Correlation was calculated for

true and predicted results for relative node number with *spearmanr* function from the

scipy.stats module.

RESULTS

Heteroblastic changes in leaflet number along the main axis

Over 460 *Cannabis sativa* leaves were collected, scanned, and their leaflet number

recorded. The leaves exhibited a profound heteroblastic juvenile-to-adult progression along

the axis, but the changes were not uniform between the accessions (Fig. **1**). In the few rare

cases where the leaves in the lower nodes were present, the first nodes always started with

a simple serrated leaf. The second leaf usually had three leaflets and the most frequent

leaflet number in the third node was five. However, the leaflet number in the nodes above

varied dramatically between accessions. The number of nodes before the transition into the

inflorescence in each of the plants also varied. We therefore calculated relative node

number, a fractional number between 0 at the shoot base to 1 at the inflorescence

transition, to compare the node leaves between plants.

Principal component analysis (PCA) and morphospace

Using the outline and landmark coordinates of 339 leaves, we modeled new synthetic

leaves, all with five leaflets (Fig. **2**). Each synthetic leaf is defined by 1,995 pseudo-

landmarks, which overcomes the problems associated with variable leaflet numbers and

permits dimension reduction using PCA and the visualization of a *Cannabis* leaf

morphospace. The first and second PCs account for 69.46% and 10.06% of the shape

variation, respectively (Fig. **3, 4**). The eigenleaves plotted in the morphospace show that the

first PC mostly accounts for the variability in the width of the leaflets; with increasing values

along PC1, the leaflets become wider. The increasing values along PC2 result in eigenleaves

with wider and less acuminate leaflets. Wider, less acuminate leaflets represent high PC2

value eigenleaves that become narrower and more acuminate as leaflet and relative node

numbers increase and PC2 values decrease. A large portion of variance associated with PC1

corresponds to narrow vs. wide leaflets that are stereotypical descriptions of *sativa* vs.

indica or *wide-* vs. *narrow- leaflet* drug varieties.

Fig. 3 Principal component analysis (PCA) of the accessions performed on the synthetic

leaves using the 1,995 pseudo-landmarks (left) and average synthetic leaf shapes for each of

the nine analyzed accessions (right). The leaf shapes in the background of the PCA plot

represent the eigenleaves of the morphospace.

Fig. 4 Principal component analysis (PCA) and average leaf shapes showing the developmental changes caused by (a) leaflet number and (b) relative node number. The first PC explains 69.46% and the second 10.06% of variation.

Examining the distribution of factors within the morphospace using a Kruskal-Wallis test reveals that accession (Table **2**, Fig. **3**) and relative node number (Table **2**, Fig. **4b**)

significantly vary by both PC1 and PC2, while leaflet number significantly varies only by PC2

(Table **2**, Fig. **4a**). This indicates that the changes in leaf shape between accessions are not

independent from developmental variation. That a facet of variation in accession leaf shape

covaries with developmental variation across the shoot in leaflet and relative node number

suggests a heterochronic mechanism by which accession differences in leaf shape arise from

changes in developmental timing.

Table 2 Kruskal-Wallis test was used to test the leaf shape variation along PC1 and PC2 for

accessions, leaflet number and relative node number.

	PC1		PC2	
	H	**p value**	**H**	**p value**
Accession	78.60	**p < 0.0001**	84.43	**p < 0.0001**
Leaflet number	1.27	p = 0.7360	156.16	**p < 0.0001**
Relative node number	91.21	**p < 0.0001**	101.35	**p < 0.0001**

Linear discriminant analysis (LDA) and prediction of genetic and developmental identities

We used LDA to model each factor as a function of 1,995 pseudo-landmark points (Fig. **5a-**

c). Model accuracy was calculated by splitting leaves arising from the same nodes into

training (177 leaves) and testing (162 leaves) datasets. To evaluate model accuracy,

accession was treated as a categorical variable, as was leaflet number, as it not only has a

small number of levels (3, 5, 7, and 9 leaflets), but each level is well separated from the

others. To evaluate the accuracy of relative node number, we treated it as a continuous

variable, due to a high number of levels (9) that continuously overlap with each other (Table

3). Models for both accession and leaflet number are highly accurate (74.07% and 92.59%,

respectively). The model for relative node number is highly accurate as well, as inferred by a

highly significant Spearman's rank correlation coefficient value between actual and

predicted values (rho = 0.729, p < 0.0001).

Table 3 Predictive power of genetic and developmental identities using the LDA model.

	Correct prediction [n]	False prediction [n]	Prediction accuracy [%]	Correlation coefficient [rho]	p value
Accession	120	42	74.07	NA	NA
Leaflet number	150	12	92.59	NA	NA
Relative node number	NA	NA	NA	0.729	< 0.0001

A confusion matrix reveals that the LDA model in most cases had a high accuracy for

predicting accession identity (Fig. **5d**). Accessions IK, RO1, and MN9 show practically no

overlap in LDA space, while AM15, IKL, FUT75, BNG, HU1, and MAR show more overlap (Fig.

5a). The model showed an almost 93% success in determining leaflet number. Leaves with

only 3 leaflets are markedly different from the rest, and the prediction model consistently

classified them correctly (Fig. **5b**). Leaves with 5 to 9 leaflets showed less pronounced

differences in shape, resulting in a slightly lower accuracy of the prediction model for these

cases. However, an examination of the confusion matrix revealed that misclassifications

mainly occurred between leaves with neighboring leaflet numbers. Specifically, the model

tended to confuse leaves with 5 leaflets for those with 7 leaflets, but not for leaves with 9

leaflets (Fig. **5e**). The exceptional ability to predict leaves with 3 leaflets from the rest may

suggest that this developmental mechanism is biased towards variation at the base of the

shoot. Similar to leaflet number, the confusion matrix for the relative node model reveals

high rates of misclassification between the neighboring relative node numbers, as is

expected, and leaves from lower nodes were very rarely classified as those from higher

nodes (Fig. **5f**). A pronounced change in leaf shape occurs between the relative nodes 0.3

and 0.4, while the shape changes in later relative nodes are more gradual (Fig. **5c**).

Fig. 5 Accession, leaflet number and relative node numbers of *Cannabis* leaves can be predicted independently of each other. Linear discriminant analysis (LDA) plots for (a) accession, (b) leaflet number and (c) relative node number. In the lower row, the confusion matrices show the true and predicted identities for (d) accessions, (e) leaflet number, and (f) relative node number using the LDA model on the split test and train dataset.

DISCUSSION

To understand how leaf shape can both scientifically and practically aid our classification and biological understanding of *Cannabis*, we first turn to grapevines. Like *Cannabis*, both grapevine rootstocks and wine and table grape scions hold tremendous cultural and economic importance (Ravaz, 1902; Galet, 1952; Galet & trans. Morton, 1979). From enforcing appellation to identifying *Phylloxera*-resistant rootstock varieties, the berries and their associated sensory properties held little taxonomic value compared to leaf shape. Like other features of the shoot, leaf shape can easily be discerned at a glance, and even compared to genotype, it can possess higher power to distinguish varieties due to complicated pedigrees, high heterozygosity, and subtle clonal differences (Myles *et al.*,

2011; Zhou *et al.*, 2017). To deal with variation, early ampelographers focused on geometric

features that could be quantified, like the degree of lobing, length-to-width ratio, and the

angle of the petiolar sinus (Ravaz, 1902; Galet, 1952; Galet & trans. Morton, 1979). That

nearly every grapevine leaf has five lobes allowed the application of landmark-based

approaches (Chitwood *et al.*, 2014; Chitwood, 2021). With landmarks, leaf shape could be

measured comprehensively and statistically. One of the most fundamental properties of

grapevine leaf shape is that genetic and node-to-node variation are orthogonal to each

other and vary independently (Chitwood *et al.*, 2016a,b). Not only is this convenient

because it allows genetic and developmental identities to be independently predicted from

each other, but insightful, because it indicates that latent effects of distinct biological

mechanisms are embedded within each leaf and can be separated from each other.

Like grapevines, striking variation in leaf shape (Fig. **1**) has historically played a significant

role in taxonomic classification of *Cannabis*. Leaf shape and differences in phyllotaxy were

among the characters Lamarck used to describe a new *Cannabis* species (Lamarck & Poiret,

1783). Anderson (1980) introduced a quantitative approach by quantifying the length-to-

width ratio of the central leaflet. Further studies using different characters—including plant

height, stem diameter, achene shape, and phytochemical profiles—to characterize

accessions have only confirmed the importance of leaf characteristics (Small *et al.*, 1976;

Hillig, 2005a). The method measuring central leaflet width-to-length ratio has been adopted

by researchers as one of the main characters for determining species, subspecies, biotypes

and chemotypes of *Cannabis* (Hillig, 2005a; Clarke & Merlin, 2013; McPartland & Small,

2020). However, this method is only able to capture a limited aspect of leaf shape variation,

neglecting other important characteristics that we measure in this study, such as leaflet

outlines, serrations, angles, and relative changes in leaflet shape across the leaf. By

modeling leaflet shape as a function of angle, we synthetically create leaves with the same

number of leaflets for which high densities of corresponding pseudo-landmarks capture high

resolution shape features (Fig. **2**).

The method presented in this study can accurately determine accession based on leaf

shape, regardless of its developmental stage (Fig. **5a, d**). The method not only works

effectively on stabilized or cloned cultivar accessions but also on wild or feral accessions

cultivated from seed that can exhibit distinct plant phenotypes (Table **1**), indicating its

robustness and potential value in future germplasm classification. The variation in leaf

shape associated with PC1 (representing 69.64% of variation), mostly accounts for the

variability in the width of the leaflets (Fig. **3**). The eigenleaves representing PC1 vary by wide

and narrow leaflets, similar to stereotypical classifications of *sativa* and *indica* plants or

wide- vs. *narrow- leaflet* drug varieties. The reliance on these non-quantitative leaf shape

descriptors has led to numerous cultivars with unreliable names, inconsistent genetic

origins, and phytochemical profiles (Sawler *et al.*, 2015; Schwabe & McGlaughlin, 2019;

Watts *et al.*, 2021; Jin *et al.*, 2021b). Using this new method, any leaf can be projected onto

the morphospace, and the resulting PC values used to provide a quantitative measure of the

leaf shape would be useful for studies investigating correlation between shape and chemical

profiles or sex of the plant. For example, Jin *et al.* (2021a) conducted a study on clones of 21

cultivars and found a strong negative correlation between the width and length ratios of

central leaflets and CBD, and a positive correlation with THC; however, Vergara *et al.* (2021)

and Murovec *et al.* (2022) were unable to confirm these findings. All three studies used low-

resolution morphometric approaches. Sex of the plants also plays a crucial role in the

cannabis industry, where the presence of male plants and inevitable pollination leads to

decreases in cannabinoid production as plants shift the use of energy into seed

development. Several methods have been employed to differentiate between male and

female plants at early stages, but only genetic methods were successful so far (Toth *et al.*,

2020; Prentout *et al.*, 2020; Campbell *et al.*, 2021; Balant *et al.*, 2022; Torres *et al.*, 2022).

Our results quantify the variation in leaf shape between accessions that can potentially be

used to classify accessions and predict chemical profiles and plant sex faster and more

accurately.

Unlike grapevine, where developmental variance is orthogonal and separate from genetic

variance, in *Cannabis* these two factors are correlated. That the developmental source of

variation is colinear with accession identity suggests that part of the differences between

accession leaf shape is explained by shifts in developmental timing, or heterochrony.

Differences, especially in flowering time and growth rates between cultivars have been

observed before (de Meijer & Keizer, 1996; Hillig, 2005a; Spitzer-Rimon *et al.*, 2019; Carlson

et al., 2021; Naim-Feil *et al.*, 2021; Stack *et al.*, 2021; Chen *et al.*, 2022) and differences in

cannabinoid profiles, leaflet index and phenological development were proposed as
characteristics to discriminate between them (de Meijer & Keizer, 1996). Heterochronic
shifts are apparent in the differential rates that accessions increase leaflet number across
nodes, as well as maximum and average leaflet counts across accessions (Fig. **1**).
Remarkably, stages in developmental timing are conserved despite being shifted. For
example, a significant shape change exhibited between the leaves with 3 and leaves with 5
leaflets, with leaflets becoming more acuminate and narrower. In contrast, changes in shape
between leaves with a higher number of leaflets were more gradual. Additionally, we
observed a similar shift in leaf shape between the nodes 0.3 and 0.4, potentially indicating a
transition between the juvenile and adult phases of leaf development. Similar results were
obtained in previous research. Spitzer-Rimon *et al.* (2022) demonstrated that flowering buds
were initiated at node 7, while Moliterni *et al.* (2004) analyzing a different cultivar, found
developing flower buds in the 4th node, suggesting that transitions in growth phases are
conserved but not synchronized across cultivars.

Conclusions

In grapevine, leaf shape has long been utilized for variety identification. However, in the
case of *Cannabis*, previous attempts were hindered by the variability in leaflet numbers. In
this study, we present a pioneering method that successfully addresses this issue. By
generating synthetic leaves with customizable leaflet counts, we can now employ high-
resolution morphometric techniques to accurately classify different wild/feral and cultivated
Cannabis accessions. Through the use of 1,995 densely placed pseudo-landmarks, we were
able to predict the accession identity with over 74% accuracy. The method works well not
only on stabilized cultivars, but also on phenotypically more variable wild/feral accessions
grown from seed. Unifying the number of leaflets allowed us, for the first time, to make
comparisons among several leaves along the main axis, enabling us to investigate
developmental changes in leaf shape and detect heterochronic mechanisms influencing the
leaf shape in *Cannabis*. The implications of this new high-resolution method in both the
cannabis industry and research extend beyond its role in determining *Cannabis* accessions.
It also offers a promising tool for developmental studies, and for studying the correlation
between leaf shape and phytochemical profiles and the sex of the plants, where lower-
resolution methods provided inconclusive results so far. The method presented here offers

a fast, effective, robust and low-cost tool that can aid the future classification of *Cannabis*

germplasm. Furthermore, the use of this methodology extends beyond *Cannabis*, and can

be applied to numerous other plant species with varying numbers of lobes and leaflets

where the use of geometric morphometrics methods was not previously possible.

ACKNOWLEDGEMENTS

This research was supported by projects WECANN (CGL2017-84297-R, Ministerio de Ciencia,

Innovación y Universidades), and M. Balant FPI predoctoral contract of the Ministerio de

Ciencia, Innovación y Universidades (PRE2018- 083226). This work is also supported by NSF

Plant Genome Research Program awards IOS-2310355, IOS-2310356, and IOS-2310357. This

project was supported by the USDA National Institute of Food and Agriculture, and by

Michigan State University AgBioResearch. We would like to thank Joan Uriach Marsal from

Uriach Laboratories for additional financial support. We would also like to acknowledge

Cannaflos—Gesellschaft für medizinisches Cannabis mbH for providing seed of accessions

IK, IKL and MAR analyzed in the study, Carlos Sáez, Carlos Ribelles, Airy Gras, Joan Vallès,

Magsar Urgamal, Shagdar Tsooj, Marine Oganesian and Nina Stepanyan-Gandilyan for help

with the sample collection and the cultivation process, and Paula Bruna for helpful advice on

improving the quality of scanned images.

COMPETING INTERESTS

None declared.

AUTHOR CONTRIBUTIONS

MB and DHC conceived the study, with OH, TG and DV inputs in a preliminary design phase

of the project. MB, TG and DV cultivated the plants. MB and OH selected the method for

imaging *Cannabis* leaves. MB and TG scanned the leaves used in the study. MB and DHC

developed the morphometric method and MB analyzed the data. MB and DHC wrote the

first draft of the paper, that all authors read, commented, and edited.

DATA AVAILABILITY

The datasets and code for the method developed here are freely available on GitHub

(https://github.com/BalantM/Cannabis_leaf_morpho).

ORCID

Manica Balant https://orcid.org/0000-0002-4316-8511

Teresa Garnatje https://orcid.org/0000-0001-6295-6217

Daniel Vitales https://orcid.org/0000-0002-3109-5028

Oriane Hidalgo https://orcid.org/0000-0002-1547-8627

Daniel H. Chitwood https://orcid.org/0000-0003-4875-1447

SUPPORTING INFORMATION

The datasets and code generated and analyzed in this study are available on GitHub

(https://github.com/BalantM/Cannabis_leaf_morpho).

REFERENCES

Abràmoff MD, Magalhães PJ, Ram SJ. **2004**. Image processing with ImageJ Part II.

Biophotonics International **11**: 36–43.

Anderson LC. **1980**. Leaf Variation among Cannabis Species from a Controlled Garden.

Botanical Museum leaflets, Harvard University **28**: 61–69.

Balant M, González Rodríguez R, Garcia S, Garnatje T, Pellicer J, Vallès J, Vitales D, Hidalgo

O. **2022**. Novel Insights into the Nature of Intraspecific Genome Size Diversity in *Cannabis*

sativa L . *Plants* **11**: 2736.

Bookstein FL. **1997**. *Morphometric Tools for Landmark Data: Geometry and Biology*.

ambrige: Cambrige University.

Bradski G. **2000**. The OpenCV Library. *Dr. Dobb's Journal: Software Tools for the Professional*

Programmer **25**: 120–123.

Bryson AE, Wilson Brown M, Mullins J, Dong W, Bahmani K, Bornowski N, Chiu C, Engelgau

P, Gettings B, Gomezcano F, *et al.* 2020. Composite modeling of leaf shape along shoots

discriminates *Vitis* species better than individual leaves. *Applications in Plant Sciences* **8**:

e11404.

Campbell LG, Peach K, Wizenberg SB. **2021**. Dioecious hemp (*Cannabis sativa* L.) plants do

not express significant sexually dimorphic morphology in the seedling stage. *Scientific*

Reports **11**: 1–8.

Carlson CH, Stack GM, Jiang Y, Taşklran B, Cala AR, Toth JA, Philippe G, Rose JKC, Smart

CD, Smart LB. **2021**. Morphometric relationships and their contribution to biomass and

cannabinoid yield in hybrids of hemp (*Cannabis sativa*). *Journal of Experimental Botany* **72**:

7694–7709.

Chen X, Guo H, Zhang Q, Wang L, Guo R, Zhan Y, Lv P, Xu Y, Guo M, Zhang Y, *et al.* 2022.

Whole-genome resequencing of wild and cultivated cannabis reveals the genetic structure

and adaptive selection of important traits. *BMC Plant Biology* **22**: 371.

Chitwood DH. **2021**. The shapes of wine and table grape leaves: An ampelometric study

inspired by the methods of Pierre Galet. *Plants People Planet* **3**: 155–170.

Chitwood DH, Klein LL, O'Hanlon R, Chacko S, Greg M, Kitchen C, Miller AJ, Londo JP.

2016a. Latent developmental and evolutionary shapes embedded within the grapevine leaf.

New Phytologist **210**: 343–355.

Chitwood DH, Mullins J, Migicovsky Z, Frank M, VanBuren R, Londo JP. **2021**. Vein-to-blade

ratio is an allometric indicator of leaf size and plasticity. *American Journal of Botany* **108**:

571–579.

Chitwood DH, Ranjan A, Martinez CC, Headland LR, Thiem T, Kumar R, Covington MF,

Hatcher T, Naylor DT, Zimmerman S, *et al.* 2014. A modern ampelography: A genetic basis

for leaf shape and venation patterning in grape. *Plant Physiology* **164**: 259–272.

Chitwood DH, Rundell SM, Li DY, Woodford QL, Yu TT, Lopez JR, Greenblatt D, Kang J,

Londo JP. **2016b**. Climate and developmental plasticity: Interannual plasticity in grapevine

leaf morphology. *Plant Physiology* **170**: 1480–1491.

Chitwood DH, Sinha NR. **2016**. Evolutionary and Environmental Forces Sculpting Leaf

Development. *Current Biology* **26**: R297–R306.

Clarke R, Merlin M. **2013**. Cannabis*: Evolution and ethnobotany*. Berkley, Los Angeles,

London: University of California Press.

Demmings EM, Williams BR, Lee C-R, Barba P, Yang S, Hwang C-F, Reisch BI, Chitwood DH,

Londo JP. **2019**. Quantitative Trait Locus Analysis of Leaf Morphology Indicates Conserved

Shape Loci in Grapevine. *Frontiers in Plant Science* **10**: 1373.

Emboden WA. **1974**. *Cannabis* - a polytypic genus. *Economic Botany* **28**: 304–310.

Galet P. **1952**. *Précis d'Ampélographie Pratique*. Digitized by Google Books from Cornell
University.

Galet P, trans. Morton L. **1979**. *A Practical Ampelography: Grapevine Identification*. Cornell
University Press (English translation).

Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, Wieser E,
Taylor J, Berg S, Smith NJ, *et al.* **2020**. Array programming with NumPy. *Nature* **585**: 357–
362.

Heslop-Harrison J, Heslop-Harrison Y. **1958**. Studies on flowering-plant growth and
organogenesis-III. Leaf shape changes associated with flowering and sex differentiation in
Cannabis sativa. *Proceedings of the Royal Irish Academy. Section B: Biological, Geological,*
and Chemical Science **59**: 257–283.

Hillig KW. **2004**. A multivariate analysis of allozyme variation in 93 *Cannabis* accessions from
the VIR germplasm collection. *Journal of Industrial Hemp* **9**: 5–22.

Hillig KW. **2005a**. *A Systematic Investigation of Cannabis [PhD Thesis]*. Indiana University.

Hillig KW. **2005b**. Genetic evidence for speciation in *Cannabis* (Cannabaceae). *Genetic*
Resources and Crop Evolution **52**: 161–180.

Hunter JD. **2007**. Matplotlib: a 2D graphics environment. *Computing in Science and*
Engineering **9**: 90–95.

Hurgobin B, Tamiru-Oli M, Welling MT, Doblin MS, Bacic A, Whelan J, Lewsey MG. **2021**.
Recent advances in *Cannabis sativa* genomics research. *New Phytologist* **230**: 73–89.

Jin D, Henry P, Shan J, Chen J. **2021a**. Identification of phenotypic characteristics in three
chemotype categories in the genus *Cannabis*. *HortScience* **56**: 481–490.

Jin D, Henry P, Shan J, Chen J. **2021b**. Classification of cannabis strains in the Canadian
market with discriminant analysis of principal components using genome-wide single
nucleotide polymorphisms (T-Y Chiang, Ed.). *PLOS ONE* **16**: e0253387.

Kuhl FP, Giardina CR. **1982**. Elliptic Fourier features of a closed contour. *Computer Graphics*
and Image Processing **18**: 236–258.

Lamarck J-BPA de M de, Poiret J-L-M. **1783**. *Encyclopédie méthodique. Botanique*.
Paris,Liège, Panckoucke;Plomteux.

Lapierre É, Monthony AS, Torkamaneh D. **2023**. Genomics-based taxonomy to clarify
cannabis classification. *Genome*.

McPartland JM. **2018**. *Cannabis* Systematics at the Levels of Family, Genus, and Species.

Cannabis and Cannabinoid Research **3**: 203–212.

McPartland JM, Guy GW. **2017**. Models of Cannabis Taxonomy, Cultural Bias, and Conflicts

between Scientific and Vernacular Names. *Botanical Review* **83**: 327–381.

McPartland JM, Small E. **2020**. A classification of endangered high-THC cannabis (Cannabis

sativa subsp. indica) domesticates and their wild relatives. *PhytoKeyes* **144**: 81–112.

de Meijer EPM, Keizer LCP. **1996**. Patterns of diversity in Cannabis. *Genetic Resources and*

Crop Evolution **43**: 41–52.

Migicovsky Z, Swift JF, Helget Z, Klein LL, Ly A, Maimaitiyiming M, Woodhouse K, Fennell

A, Kwasniewski M, Miller AJ, *et al.* 2022. Increases in vein length compensate for leaf area

lost to lobing in grapevine. *American Journal of Botany* **109**: 1063–1073.

Moliterni VMC, Cattivelli L, Ranalli P, Mandolino G. **2004**. The sexual differentiation of

Cannabis sativa L.: A morphological and molecular study. *Euphytica* **140**: 95–106.

Murovec J, Eržen JJ, Flajšman M, Vodnik D. **2022**. Analysis of Morphological Traits,

Cannabinoid Profiles, THCAS Gene Sequences, and Photosynthesis in Wide and Narrow

Leaflet High-Cannabidiol Breeding Populations of Medical Cannabis. *Frontiers in Plant*

Science **13**: 786161.

Myles S, Boyko AR, Owens CL, Brown PJ, Grassi F, Aradhya MK, Prins B, Reynolds A, Chia

JM, Ware D, *et al.* 2011. Genetic structure and domestication history of the grape.

Proceedings of the National Academy of Sciences of the United States of America **108**: 3530–

3535.

Naim-Feil E, Pembleton LW, Spooner LE, Malthouse AL, Miner A, Quinn M, Polotnianka

RM, Baillie RC, Spangenberg GC, Cogan NOI. **2021**. The characterization of key physiological

traits of medicinal cannabis (*Cannabis sativa* L.) as a tool for precision breeding. *BMC Plant*

Biology **21**: 1–15.

Niklas KJ. **1994**. *Plant allometry: the scaling of form and process*. University of Chicago

Press.

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M,

Prettenhofer P, Weiss R, Dubourg V, *et al.* 2011. Generating the blood exposome database

using a comprehensive text mining and database fusion approach. *Journal ofMachine*

Learning Research **12**: 2825–2830.

Potter DJ. **2009**. *Optimisation Of* Cannabis sativa *L. As A Phytopharmaceutical [PhD Thesis]*.

London: King's College London, Department of Pharmaceutical Science Research.

Prentout D, Razumova O, Rhoné B, Badouin H, Henri H, Feng C, Käfer J, Karlov G, Marais

GAB. **2020**. An efficient RNA-seq-based segregation analysis identifies the sex chromosomes

of *Cannabis sativa*. *Genome Research* **30**: 164–172.

Quimby MW, Doorenbos NJ, Turner CE, Masoud A. **1973**. Mississippi-Grown Marihuana -

Cannabis sativa Cultivation and Observed Morphological Variations. *Economic Botany* **27**:

117–127.

Ravaz L. **1902**. *Les vignes américaines: Porte-greffes et producteurs directs*. Goulet,

Montpellier and Paris: Digitized by Google Books from Cornell University.

Ren G, Zhang X, Li Y, Ridout K, Serrano-Serrano ML, Yang Y, Liu A, Ravikanth G, Nawaz MA,

Mumtaz AS, *et al.* 2021. Large-scale whole-genome sequencing unravels the domestication

history of *Cannabis sativa*. *Science Advances* **7**: eabg2286.

Sawler J, Stout JM, Gardner KM, Hudson D, Vidmar J, Butler L, Page JE, Myles S. **2015**. The

Genetic Structure of Marijuana and Hemp. *PLOS ONE* **10**: e0133292.

Schultes RE, Klein WM, Plowman T, Lockwood TE. **1974**. Cannabis: An example of

taxonomic neglect. *Botanical Museum Leaflets, Harvard University* **23**: 337–367.

Schwabe AL, McGlaughlin ME. **2019**. Genetic tools weed out misconceptions of strain

reliability in *Cannabis sativa*: implications for a budding industry. *Journal of Cannabis*

Research **1**: 3.

Small E. **2015**. Evolution and Classification of *Cannabis sativa* (Marijuana, Hemp) in Relation

to Human Utilization. *Botanical Review* **81**: 189–294.

Small E, Beckstead HD. **1973**. Cannabinoid Phenotypes in *Cannabis sativa*. *Nature* **245**: 147–

148.

Small E, Cronquist A. **1976**. A Practical and Natural Taxonomy for Cannabis. *Taxon* **25**: 405–

435.

Small E, Jui PY, Lefkovitch LP. **1976**. A Numerical Taxonomic Analysis of Cannabis with

Special Reference to Species Delimitation. *Systematic Botany* **1**: 67–84.

Spitzer-Rimon B, Duchin S, Bernstein N, Kamenetsky R. **2019**. Architecture and Florogenesis

in Female *Cannabis sativa* Plants. *Frontiers in Plant Science* **10**: 350.

Spitzer-Rimon B, Shafran-Tomer H, Gottlieb GH, Doron-Faigenboim A, Zemach H,

Kamenetsky-Goldstein R, Flaishman M. **2022**. Non-photoperiodic transition of female

cannabis seedlings from juvenile to adult reproductive stage. *Plant Reproduction* **35**: 265–

277.

Stack GM, Toth JA, Carlson CH, Cala AR, Marrero-González MI, Wilk RL, Gentner DR,

Crawford JL, Philippe G, Rose JKC, *et al.* 2021. Season-long characterization of high-

cannabinoid hemp (*Cannabis sativa* L.) reveals variation in cannabinoid accumulation,

flowering time, and disease resistance. *GCB Bioenergy* **13**: 546–561.

The Global Cannabis Report, 3rd Edition. **2022**.

(https://prohibitionpartners.com/reports/the-global-cannabis-report-edition-3/) [accessed

13 August 2023]

Torres A, Pauli C, Givens R, Argyris J, Allen K, Monfort A, Gaudino RJ. 2022.

High-throughput methods to identify male *Cannabis sativa* using various genotyping

methods. *Journal of Cannabis Research* **4**: 57.

Toth JA, Stack GM, Cala AR, Carlson CH, Wilk RL, Crawford JL, Viands DR, Philippe G, Smart

CD, Rose JKC, *et al.* 2020. Development and validation of genetic markers for sex and

cannabinoid chemotype in *Cannabis sativa* L. *GCB Bioenergy* **12**: 213–222.

Vergara D, Feathers C, Huscher EL, Holmes B, Haas JA, Kane NC. 2021. Widely assumed

phenotypic associations in *Cannabis sativa* lack a shared genetic basis. *PeerJ* **9**: e10672.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E,

Peterson P, Weckesser W, Bright J, *et al.* 2020. SciPy 1.0: fundamental algorithms for

scientific computing in Python. *Nature Methods* **17**: 261–272.

Waskom M. 2021. seaborn: statistical data visualization. *Journal of Open Source Software* **6**:

3021.

Watts S, McElroy M, Migicovsky Z, Maassen H, van Velzen R, Myles S. 2021. *Cannabis*

labelling is associated with genetic variation in terpene synthase genes. *Nature Plants* **7**:

1330–1334.

Zhou Y, Massonnet M, Sanjak JS, Cantu D, Gaut BS. 2017. Evolutionary genomics of grape

(*Vitis vinifera* ssp. *vinifera*) domestication. *Proceedings of the National Academy of Sciences*

of the United States of America **114**: 11715–11720.

Zhukovskii PM. 1971. *Cultivated plants and their wild relatives*. Leningrad, USSR, Kolos.

718