Intra-leaf modeling of *Cannabis* leaflet shape produces leaf models that predict genetic

and developmental identities

Manica Balant [1,2,3], Teresa Garnatje [1,4], Daniel Vitales [1], Oriane Hidalgo [1,5], Daniel H.

Chitwood [3,6]

[1] Institut Botànic de Barcelona, IBB (CSIC-CMCNB), Passeig del Migdia s.n., 08038 Barcelona,

Spain

[2] Laboratori de Botànica (UB), Unitat Associada al CSIC, Facultat de Farmàcia i Ciències de

l'Alimentació, Universitat de Barcelona, Av. Joan XXIII 27–31, 08028 Barcelona, Spain

[3] Department of Horticulture, Michigan State University, East Lansing, MI, USA

[4] Jardí Botànic Marimurtra - Fundació Carl Faust, pg. Carles Faust, 9, 17300 Blanes, Spain

[5] Royal Botanic Gardens, Kew, Kew Green, Richmond TW9 3AE, UK

[6] Department of Computational Mathematics, Science & Engineering, Michigan State

University, East Lansing, MI, USA

Authors for correspondence:

Daniel H. Chitwood

Email: dhchitwood@gmail.com

Manica Balant

Email: manica.balant@ibb.csic.es

Summary

● The iconic, palmately compound leaves of *Cannabis* have attracted significant

attention in the past. However, investigations into the genetic basis of leaf shape or

its connections to phytochemical composition have yielded inconclusive results. This

is partly due to prominent changes in leaflet number within a single plant during

development, which has so far prevented the proper use of common morphometric

techniques.

● Here we present a new method that overcomes the challenge of nonhomologous

landmarks in palmate, pinnate and lobed leaves, using *Cannabis* as an example. We

model corresponding pseudo-landmarks for each leaflet as angle-radius coordinates

and model them as a function of leaflet to create continuous polynomial models,

bypassing the problems associated with variable number of leaflets between leaves.

● We analyze 341 leaves from 24 individuals from nine *Cannabis* accessions. Using

3,591 pseudo-landmarks in modeled leaves, we accurately predict accession identity,

leaflet number, and relative node number.

● Intra-leaf modeling offers a rapid, cost-effective means of identifying *Cannabis*

accessions, making it a valuable tool for future taxonomic studies, cultivar

recognition, and possibly chemical content analysis and sex identification, in addition

to permitting the morphometric analysis of leaves in any species with variable

numbers of leaflets or lobes.

Key words: *Cannabis*, development, geometric morphometrics, intra-leaf modeling, leaf

morphology, leaf shape

INTRODUCTION

Cannabis sativa L. (hereafter referred to as *Cannabis*) is a versatile crop plant used by

humans for a variety of purposes throughout history. Although today it is commonly

associated with its psychoactive properties, traditional medicine has relied heavily on

Cannabis, and it is also a valuable source of food and fibers (Clarke & Merlin, 2013). Genetic

and archaeological evidence suggests that *Cannabis* was domesticated around 12,000 years

ago in East Asia, initially serving as a multipurpose crop before separate selections for fiber

and drug production emerged around 4,000 years ago (Ren *et al.*, 2021). Since then,

widespread cultivation has facilitated its global distribution. Throughout the 20th century,

Cannabis use was largely abandoned due to its illegal status in many parts of the world.

However, recent legalization for recreational and/or medicinal purposes in many countries

worldwide has led to a surge in the cannabis industry (*The Global Cannabis Report, 3rd*

Edition, 2022).

Extensive *Cannabis* use has resulted in the development of numerous cultivars and strains

that are well-suited to diverse uses and climates (Small, 2015). This significant

morphological and phytochemical diversity within the *Cannabis* genus poses challenges for

taxonomic classification. Over the past two centuries, various taxonomic approaches based

on genetics, morphology, and phytochemistry have been proposed (McPartland & Small,

2020). Some scientists advocated for a polytypic classification, recognizing the presence of

two (Lamarck & Poiret, 1783; Zhukovskii, 1971; Hillig, 2005a) or three (Emboden, 1974;

Schultes *et al.*, 1974; Hillig, 2005b; Clarke & Merlin, 2013) species with multiple subspecies,

while others argued for a monotypic genus, considering only a single species, *Cannabis*

sativa (Small & Cronquist, 1976; Sawler *et al.*, 2015; Small, 2015; McPartland, 2018;

McPartland & Small, 2020; Ren *et al.*, 2021). Hillig (2005a) introduced a classification system

based on biotypes, considering molecular, morphological, and phytochemical data. He

proposed dividing *Cannabis* into two species, *C. sativa* and *C. indica* Lam., and six biotypes:

C. indica as narrow-leaflet drug (NLD), wide-leaflet drug (WLD), hemp and feral biotype, and

C. sativa as hemp and feral biotype. Recently, Lapierre *et al.* (2023) conducted a

comprehensive taxonomic review of the *Cannabis* genus and based on available genetic

data, strongly supported the theory that *Cannabis* is a highly diverse monotypic species.

Apart from taxonomic classification, *Cannabis* is often categorized based on its cultivation

purpose, morphology, and chemical composition. Fiber-type plants, commonly known as

hemp, are primarily grown for fiber and seed production. These plants contain less than

0.3% of the psychoactive compound THC (Δ9-tetrahydrocannabinol), while drug-type plants,

often referred to as marijuana and medicinal cannabis, can contain higher levels of THC

(Hurgobin *et al.*, 2021). *Cannabis* plants can also be separated based on the ratio of two

major cannabinoids THC and CBD (cannabidiol) into Type I (THC dominant), Type II

(balanced CBD/THC ratio), and Type III plants (CBD dominant) (Small & Beckstead, 1973). In

the medicinal and recreational cannabis industries plants are normally categorized as

'sativa', 'indica', or 'hybrid'. Taller plants with narrow leaflets and high THC percentage are

called 'sativa', while shorter and bushier plants with wider leaflets and high percentages of

both CBD and THC are called 'indica'. Plants with intermediate characters are called

'hybrids' (McPartland & Guy, 2017). While the classification of *Cannabis* into 'indica' and

'sativa' is not supported by genetic data, the visible differences in leaflet width have long

been a significant characteristic used to visually discriminate different types of *Cannabis*.

Cannabis arguably possesses one of the most iconic leaves among all plants. Its palmately

compound leaves with a varying number of leaflets are a popular culture symbol. *Cannabis*

exhibits a remarkable degree of phenotypic plasticity, further accentuated by selection

pressure during the domestication process (Small, 2015). Extensive variability in leaf

morphology has already been described by Quimby *et al.* (1973) and later Anderson (1980),

who was the first to quantify the width, length, and ratio of the central leaflet. This or

similar methods were then commonly used in studies investigating the morphological

characteristics of *Cannabis* species, subspecies, cultivars, biotypes and chemotypes (Small *et*

al., 1976; de Meijer *et al.*, 1992; de Meijer & Keizer, 1996; Hillig, 2005a; Clarke & Merlin,

2013; Lynch *et al.*, 2016; Karlov *et al.*, 2017; Parsons *et al.*, 2019; McPartland & Small, 2020;

Carlson *et al.*, 2021; Islam *et al.*, 2021; Jin *et al.*, 2021a; Vergara *et al.*, 2021; Buzna & Sala,

2022; Chen *et al.*, 2022; Murovec *et al.*, 2022), often with contradictory results. Leaf shape

has therefore played an important and sometimes controversial role in *Cannabis* taxonomy.

While researchers in previous *Cannabis* studies were aware of enormous plasticity and the

effect the environment has on leaf shape (Vergara *et al.*, 2021; Murovec *et al.*, 2022), they

very rarely paid attention to the effects of developmental processes, even though

heteroblastic changes (differences in leaf shape arising from juvenile-to-adult phase

transitions in the meristem) profoundly affect the arrangement and shape of *Cannabis*

leaves along the shoot. While some studies briefly mention the developmental changes of

leaves (Hillig, 2005a; Carlson *et al.*, 2021; Jin *et al.*, 2021b; Spitzer-Rimon *et al.*, 2022), the

only two studies focusing on heteroblastic phase changes in leaves along the plant axis were

done by Heslop-Harrison and Heslop-Harrison (1958) and Hesami *et al*. (2023). In the lower

part of the shoot *Cannabis* leaves exhibit opposite phyllotaxy and one to three leaflets,

transitioning to alternate phyllotaxy and leaves with up to 11 or 13 leaflets in the upper

section (Hillig, 2005a; Clarke & Merlin, 2013; Small, 2015). Additionally, the changes in

leaflet number are not uniform between different *Cannabis* accessions (Hillig, 2005a). These

changes during development not only complicate categorization of plant accessions based

on leaf shape, but also prevent the use of morphometric techniques.

Morphometrics is the quantitative analysis of shape. It includes a wide range of methods,

from measuring allometric differences in dimensions like lengths, widths, and angles in

relation to size (Niklas, 1994), to geometric techniques that measure shape

comprehensively, like elliptical Fourier (EFDs; Kuhl & Giardina, 1982) and landmark-based

analyses (Bookstein, 1997). It can be used to classify species and to separate effects on

shape arising from genetic, developmental, and environmental mechanisms (Chitwood &

Sinha, 2016). Historically the field of ampelography (ἄμπελος, 'vine' + γράφος, 'writing';

Ravaz, 1902; Galet, 1952; Galet & trans. Morton, 1979) relied heavily on leaf shape to

distinguish grapevine varieties. Unlike *Cannabis,* grapevine leaves have a consistent number

of lobes, sinuses, and other associated homologous points that can be used for both

landmark-based and EFD morphometric analysis (Chitwood *et al.*, 2014; Chitwood, 2021) to

disentangle genetic (Demmings *et al.*, 2019), developmental (Chitwood *et al.*, 2016a; Bryson

et al., 2020; Migicovsky *et al.*, 2022), and environmental effects (Chitwood *et al.*, 2016b,

2021) embedded in leaf shapes.

The variable number of leaflets in *Cannabis* (and several other species with lobed, pinnate

and palmate compound leaves) precludes analysis methods that rely on homologous,

comparable points to measure shape comprehensively. Methods to automatically isolate

individual leaflets (Failmezger *et al.*, 2018) or to model developmental trajectories, such as

heteroblastic series (Biot *et al.*, 2016) were proposed previously for morphometrical analysis

in such cases. In *Cannabis,* Vergara *et al.* (2021) used a landmark-based approach but were

limited to analyzing the central and two most distal leaflets on each side, features that all

Cannabis leaves except single-leaflet leaves possess, but which excludes most of the shape

variation within a leaf.

Here, we seek to build on these works and conceptually extend our framework of

continuously modeling leaflets within a palmate leaf. We model corresponding pseudo-

landmarks for each leaflet as angle-radius coordinates relative to the petiolar junction and

model angle and radius as a function of leaflet number to create continuous polynomial

models that bypass the problems associated with variable numbers of leaflets between

leaves. This enabled us to compare leaves with different numbers of leaflets within a plant

and to discern differences between genotypes rather than the heteroblastic series.

Analyzing over 300 *Cannabis* leaves, we model theoretical leaves with nine leaflets and

3,591 comparable pseudo-landmarks. Linear discriminant analysis (LDA) predicts accession,

leaflet number, and relative node number with high accuracy. Intra-leaf modeling allows the

application of morphometric techniques to comprehensively measure leaf shape in

Cannabis, enabling future taxonomic and developmental studies, cultivar recognition, and
possibly chemical content analysis and sex identification, in addition to permitting the
morphometric analysis of leaves in any species with variable numbers of leaflets or lobes.

MATERIAL AND METHODS

Plant material and growing conditions

This study includes 24 individuals from nine accessions of *Cannabis sativa* L. (Table **1**; Fig. **1**),
encompassing both wild/feral accessions and cultivated varieties with a wide distribution
area. The plants were grown from seeds in a growth chamber (Fitoclima D1200PLL, Aralab,
Portugal) to minimize the influence of the environment. Before sowing, the seeds were
sterilized overnight in a 5% H_2O_2 solution with the addition of Inex-A solution (Cosmocel,
Spain) at room temperature. Sterilized seeds were then transferred to Petri dishes and
placed in the growth chamber for germination. Once the first leaves emerged, the seedlings
were transferred to small peat pots with a pre-fertilized soil substrate (Kilomix Atami,
Spain). During this phase, the environmental conditions were set to 25°C, with an 18-hour
day and 6-hour night photoperiod, and a light intensity of 50 µmol $m^{-2}s^{-1}$ (Philips Master PL-
L 55W, Spain). After two weeks the surviving plants were transplanted to 3.5 l pots with the
same soil substrate. The light intensity was gradually increased to 300 µmol $m^{-2}s^{-1}$ over the
following week, without changing the photoperiod and temperature. The onset of flowering
in some *Cannabis* accessions is photoperiod dependent, therefore after four weeks, the
photoperiod was changed to 12 hours of daylight and 12 hours of darkness, and the light
intensity was gradually increased to 700 µmol $m^{-2}s^{-1}$ over the following week, while keeping
the temperature at 25°C. The plants remained in these environmental conditions until the
flowering stage. Plants received daily irrigation with tap water, without any application of
nutrient or phytosanitary control.

Table 1 Accession details and number of leaves collected and analyzed in the study.

Accession ID	Accession type	Location/Cultivar name	Number of individuals	Number of leaves collected	Number of leaves analyzed
AM15	Wild/feral	Armenia, Sjunik marz, Goris town	5	90	74
BNG	Wild/feral	Bangladesh, Rangpur, Carmichael College Campus	1	14	10
FUT75	Cultivar	Futura 75	2	45	30
HU1	Wild/feral	Hungary, Nyírvasvári	4	83	68
IK	Landrace	India, Kerala	4	92	53
IKL	Landrace	India, Kullu	4	69	47
MAR	Landrace	Morocco, North Morocco	1	18	15
MN9	Wild/feral	Mongolia, Selenge aimag, Baruunburen sum	1	14	10
RO1	Wild/feral	Romania, Mangalija	2	36	34

Leaf sampling and imaging

A total of 461 leaves were sampled during the flowering stage, with the exception of

individuals from the accession IK, which did not begin to flower during the two-month

cultivation period. Leaves along the main axis of the plants were collected and immediately

scanned using a flatbed photograph scanner (Epson Perfection V370, Japan) at 1200 dpi

resolution. A piece of velvet fabric was placed between the leaf and the scanner cover to

avoid any shadows. No adjustments to the angle of individual leaflets were made before

scanning. Each leaf was scanned with a scale and a label indicating the node it originated

from, followed by a sequential lowercase letter, since typically two leaves are present per

node. Starting at the base of the plant, the first two leaves were labeled as leaves "a" and

"b" from node number 1, and so on, until the shoot apex.

Cannabis leaves display a marked heteroblastic, or juvenile-to-adult, leaf shape progression.

Mature, juvenile leaves located on the first node at the base of the plant usually have a

simple, serrated leaf. As node number increases so does the leaflet number, reaching a

maximum of 9 to 13 leaflets in young, adult leaves at the growing tip. Eventually leaves

transition into an inflorescence type. During this transition, the number of leaflets per leaf

starts to decrease again until the top of the inflorescence. Leaves at the shoot base have

opposite phyllotaxy and transition to alternate phyllotaxy in the upper section on the stem

and inflorescence (Heslop-Harrison & Heslop-Harrison, 1958; Hillig, 2004; Potter, 2009;

Spitzer-Rimon *et al.*, 2022). To ensure that only stem leaves were included in our analysis,

we separated the two types (i.e., stem and inflorescence leaves) based on the point where

the decrease in the number of leaflets appeared. This point determined the "total node

number", the number of nodes per plant used for further analysis. Total node number

varied among individuals. To compare node positions, a relative node number was

calculated, which was defined by the node position divided by the total node number for

the individual plant, where zero is at the plant base and one at the last node included in the

analysis (Fig. **1**). Because of the nature of plant growth, the leaves at the base of the plant

were frequently too senesced to be incorporated in the analysis or were entirely lost.

Nevertheless, the nodes could still be identified, which allowed them to be taken into

account in the calculation of relative node number.

Image analysis and landmarking

After eliminating damaged and deformed leaves (39), simple leaves (4), leaves with even

leaflet numbers (3) and leaves with relative node values above one (57), a total of 358

Cannabis leaves were used for image analysis and landmarking. Photoshop was used to

separate petioles and leaflets smaller than 1 cm from the rest of the leaf. The leaf outlines

were then extracted and saved using Python modules NumPy (Harris *et al.*, 2020),

Matplotlib (Hunter, 2007) and OpenCV (Bradski, 2000). The code for extracting and plotting

the   leaf   outlines   can   be   found   on   GitHub

(https://github.com/BalantM/Cannabis_leaf_morpho_updated). The *x* and *y* coordinates of

blade outlines and landmarks were extracted using ImageJ (Abràmoff *et al.*, 2004). The

outline was extracted using the *wand* tool (setting tolerance to 20 and including "smooth if

thresholded" option) and the landmarks were placed using the *multi-point* tool.

(a)

	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1
AM15	3	4	5	7	7	7	7	7	7	
BNG		3	3	5	5.5	7	8	7	8	
FUT75			6	7	9	9	9	10	9	
HU1	5	6.5	7	9	9	9	9	9	9	
IK			7	9	9	9	9	9	9	
IKL	1	5	5	7	8.5	9	10	10	11	
MAR		4	7	7.5	9	9	10	11	10	
MN9			5	7		7	7	7	7	
RO1	2.5	3	5	7	7	7	7	7	7	

shoot base
mature, juvenile leaves

Relative node number

shoot tip
young, adult leaves

(b)

Fig. 1 Changes in the leaf shape and leaflet number during the development in nine *Cannabis* accessions. (a) Median values for all available leaflet number for each relative node number for the nine *Cannabis* accessions. (b) Changes in leaf shape between different developmental stages in different *Cannabis* accessions.

Initially, landmarks were placed at the beginning and end of each leaflet, starting from the lower left side, and continuing to the lower right side of the leaf outline. Subsequently, landmarks were placed in the same order on the tips of the leaflets. The final landmark was positioned at the center of the petiolar junction (Fig. **2**, second column). These landmarks delimit the boundaries of the leaflets so that equidistant pseudo-landmarks can later be

placed along the contour. The number of landmarks per leaf ranged from 10 to 28,

depending on the leaflet number. The raw data containing the coordinates for leaf outlines

and   landmarks   can   be   accessed   on   GitHub

(https://github.com/BalantM/Cannabis_leaf_morpho_updated).

Fig. 2 The process of modeling theoretical leaves for a leaf with (a) three leaflets from

accession AM15, (b) five leaflets from accession IKL, (c) seven leaflets from accession FUT75,

and (d) nine leaflets from accession IK. The first column shows the scans of the leaves, which

we use to extract the outline and place the landmarks on the tip, start, and end of each

leaflet and on the petiolar junction (second column). These coordinates are used to

generate 200 equidistant pseudo-landmarks on each side of each leaflet, sharing the

landmark on the tip of the leaflet for a total of 399 pseudo-landmarks. These coordinates

are then converted into polar coordinates. Each transformed leaflet is defined with 399

equidistant pseudo-landmarks, with three landmarks, two at the base and one at the tip.

Large points are placed every 25 pseudo-landmarks to emphasize that leaflet outlines are

defined by points (third column). Second degree polynomials for angles and for radius from

petiolar junction are then fitted through these 399 pseudo-landmarks (fourth column). A

modeled theoretical leaf with nine leaflets defined by 3,591 pseudo-landmarks can then be

modeled using the collection of 798 polynomial models for each leaf (399 polynomial

models for angles and 399 for radius from petiolar junction) (fifth column) and visualized in

the cartesian coordinate system (sixth column).

Reconstruction of the new modeled leaves

To analyze leaves with different numbers of leaflets, pseudo-landmarks of each leaflet were

modeled as 2^{nd} degree polynomial models of angles and radius as functions of leaflet

number within a leaf, in order to use the models to construct a modeled theoretical leaf

with a desired number of leaflets. The Python code, presented as a Jupyter notebook with

detailed   description,   is   available   on   GitHub

(https://github.com/BalantM/Cannabis_leaf_morpho_updated). The *x* and *y* coordinates of

the leaf outline were first interpolated to create an arbitrarily high number of coordinates to

increase resolution of the leaf outline. The coordinates of manually selected landmarks

were then compared against the high-resolution coordinates of the leaf outline and the

nearest neighboring point of the high-resolution coordinates to each original landmark was

identified and specified as the new landmark point. Next, the outline and new landmark

coordinates were rotated, translated, and scaled so that the central leaflet had a length of

one and pointed in the same direction. The transformed points were then interpolated to

generate 200 pseudo-landmarks on each side of each leaflet (from the landmark at the

bottom until the tip of the leaflet), sharing the landmark on the tip of the leaflet (i.e., a total

of 399 pseudo-landmarks per leaflet). These pseudo-landmarks were then converted to

polar coordinates, where each point was defined by a radius and angle relative to the

landmark of the petiolar junction and tip of the central leaflet (Fig. **2**, third column).

Using the polar coordinates of each leaflet, 2^{nd} degree polynomial models for *x* (angle) and *y*

(radius from petiolar junction) values were fit through each of the 399 corresponding

pseudo-landmarks for each leaflet using the Python *scipy.optimize.curve_fit* function

(Virtanen *et al.*, 2020), modeling angle and radius as a function of leaflet number (Fig. **2**,

fourth column). Using the coefficients for 2^{nd} degree polynomial models, we then model

each pseudo-landmark as a function of leaflet number to reconstruct the new theoretical

leaf with an arbitrary number of leaflets. Meaning that for each leaflet, each of the 399 *x*

and *y* pseudo-landmarks (i.e., angle and radius from petiolar junction coordinates) was

calculated using the 2^{nd} degree polynomial function, with coefficients obtained from the

previous step, and the newly defined leaflet number (9 in this case). The optimal number of

reconstructed leaflets was tested for the best prediction accuracy in Linear discriminant

analysis modeling and the highest accuracy was achieved by reconstructing 9 leaflets (Table

S1). It is important to note that the reconstructions start with the first real leaflet and end

with the last real leaflet. These 9 reconstructed leaflets are then equally divided between

these two points.

Nine leaflets were reconstructed using the collection of coefficients of 789 2^{nd} degree

polynomial models for each leaf; the 399 models for angle were used to model theoretical x

(i.e., angle) and 399 models for radius were used to model theoretical y (i.e., radius from

petiolar junction) pseudo-landmarks as a function of nine leaflets.

The coordinates defining the 3,591 pseudo-landmarks for each of the modeled leaves (399

pseudo-landmarks for each of the 9 reconstructed leaflets) were then plotted and visually

inspected. We detected 17 inaccurately modeled leaves, most likely caused by the position

of the petiole landmark compared to the landmark marking the start and end landmarks of

the leaflet. A total of 341 *Cannabis* leaves were then used in the analysis.

Validation of the leaf modeling approach

To validate our modeling approach, we extracted the polar coordinates of the original

central leaflets (Fig. **3a**) and central leaflets of the modeled leaves (Fig. **3b**) and used them in

Procrustes analysis using *Procrustes* function from scipy.spatial module (Virtanen *et al.*,

2020). Procrustes analysis minimizes the distance between all points for a set of

landmarks/pseudo-landmarks between two samples through translation, rotation, and

scaling, and returns new points of the two sets, superimposed to each other (Fig. **3c**). We

then calculated the Procrustes distance between the original central leaflet (angle and

radius coordinates) to its corresponding modeled reconstruction, a measure of their

similarity. The mean distance was calculated and compared to that of simulated

bootstrapped mean values by resampling (10,000 resamples) through randomly sorting

original leaflet coordinates against coordinates of reconstructed leaflets.

Morphometric analysis of the central leaflet shape using previously established

methodologies

The width-to-length ratio (W/L ratio), first described by Anderson (1980), was frequently

used to describe the shape of *Cannabis* leaves or even differentiate between different

Cannabis taxa. With previously established morphometric methods, the shape analysis of

central leaflets (that all leaves share) would also be possible, using EFDs or pseudo-

landmarks approach. To evaluate the effectiveness of these two previous methods for the

shape analysis of *Cannabis* leaves, we first extracted the Cartesian coordinates of central

leaflets (Fig. **4a**), that were previously scaled, rotated and translated, so that they were all

pointing in the same direction and had the length of one. We then interpolated 200 pseudo-

landmarks on each side of each leaflet, sharing the landmark on the tip of the leaflet (i.e., a

total of 399 pseudo-landmarks per leaflet).

To measure the W/L ratio, we calculated width of the leaf (as the leaves were already

normalized to length of one), calculating the minimum bounding rectangle. The distribution

of widths was then plotted using Python package *seaborn.kdeplot*. To see if the analyzed

accessions differed significantly in their W/L ratios, Kruskal-Wallis test was calculated using

stats.kruskal function from the scipy.stats module. To see which of the accessions differ in

W/L ratio, we calculated Dunn's Multiple Comparison Test with *scikit_posthocs* package in

Python (Terpilowski, 2019), using the *posthoc_dunn* function.

Linear discriminant analysis (LDA) was applied to model accession, leaflet number, and

relative node number as the function of central leaflet coordinate values, using the

LinearDiscriminantAnalysis function from the scikit-learn module in Python (Pedregosa *et*

al., 2011). To test the performance of the LDA model, the dataset was divided into two

parts. Since most of the analyzed leaves exhibit opposite phyllotaxy, wherein the nodes

were represented by two leaves (a and b) in the same developmental phase with the same

number of leaflets, the dataset was split into a training dataset (leaf a) comprising 180

leaves and a test dataset (leaf b) containing 161 leaves. The *predict* function from

LinearDiscriminantAnalysis in the scikit-learn module was used to predict the accession

identity, leaflet number, and relative node number, based on the central leaflet coordinate

values. The accuracy of the LDA model was calculated and visualized using the function

confusion_matrix from scikit-learn. Spearman Rank Correlation was calculated for true and

predicted results for relative node number with *spearmanr* function from the scipy.stats

module.

Data analysis of modeled leaves

A principal component analysis (PCA) was performed on the coordinates of the modeled

leaves using scikit-learn module in Python and proportions of explained variance for each

principal component and the cumulative variance was calculated. Points representing the

leaves were colored by the accession identity, leaflet number, or relative node number (Fig.

5). To see which of the first two PCs explains most of the leaf shape variation for accessions,

leaflet number and relative node number, Kruskal-Wallis test was calculated using

stats.kruskal function from the scipy.stats module. To visualize an average leaf for each

accession, leaflet number, and relative node number, the average coordinate values of

modeled leaves were calculated for each of the categories and plotted using the Matplotlib

module in Python (Fig. **5**).

To see if the modeled leaves can be used to model accession, leaflet number, and relative

node number, we followed the same steps as before for shape analysis of central leaflet.

Linear discriminant analysis (LDA) was applied to model accession, leaflet number, and

relative node number. The dataset was again split into a training and test dataset to see if

we were able to predict accession, leaflet number, and relative node number identity, based

on the coordinates of modeled leaves. The same was done on a combined dataset with

3990 coordinates, created by concatenating coordinates of modeled leaves and the

coordinates of the original central leaflets.

RESULTS

Heteroblastic changes in leaflet number along the main axis

Over 460 *Cannabis sativa* leaves were collected, scanned, and their leaflet number

recorded. The leaves exhibited a profound heteroblastic juvenile-to-adult progression along

the axis, but the changes were not uniform between the accessions (Fig. **1**). In the few rare

cases where the leaves in the lower nodes were present, the first nodes always started with

a simple serrated leaf. The second leaf usually had three leaflets and the most frequent

leaflet number in the third node was five. However, the leaflet number in the nodes above

varied dramatically between accessions. The number of nodes before the transition into the

inflorescence in each of the plants also varied. We therefore calculated relative node

number, a fractional number between 0 at the shoot base to 1 at the inflorescence

transition, to compare the node leaves between plants.

Validation of the leaf modeling approach

The modeling approach was validated by calculating the mean Procrustes distance of

modeled central leaflet coordinates to original central leaflet coordinates using 10,000

bootstrap replicas, assessing resampled means against the actual Procrustes mean value.

None of the 10,000 resamples yielded a mean lower than the observed Procrustes value,

confirming the robustness of the novel modeling approach (Fig. **3d**).

Fig. 3 Modeling approach validation using Procrustes analysis and bootstrap resampling. The

(a) original and (b) modeled central leaflets in polar coordinate system were superimposed

(c) and Procrustes distances calculated. (d) The resampled mean was plotted as a

distribution (green histogram) against the actual Procrustes mean (grey vertical line).

Width-to-length (W/L) ratio and central leaflet shape analysis

Our results indicate that the width-to-length (W/L) ratio of central leaflets is not able to

differentiate well between different *Cannabis* leaf accessions based on this information

alone (Fig. **4**). While the Kruskal-Wallis test did show overall significance between accessions

(Table **S2**), Dunn's post hoc indicated significance in leaf morphology for just one accession

(Table **S3**). The W/L ratio significantly differs from the rest only for the IK accession,

characterized by particularly narrow leaves (Table **S3**). The Kruskal-Wallis test was also

significant for leaflet numbers and relative node numbers (Table **S2**). Dunn's post hoc

revealed that while we can differentiate between leaflet numbers based on the W/L ratio of

central leaflet, we can only separate the lower and higher relative nodes (Table **S3**).

Fig. 4 Analysis of leaf shape using the approach adapted from Anderson (1980). (a) Visualization of the 341 central leaflets used in the analysis. W/L ratios plotted by (b) accession, (c) leaflet number and (d) relative node number.

To test whether the outline of the central leaflet can better predict the genetic and

developmental identity of *Cannabis* leaves, we used Linear discriminant analysis (LDA) to

model each factor as a function of 399 pseudo-landmark points defining the shape of

central leaflet (Fig. **5a-c**). To evaluate model accuracy, accession was treated as a categorical

variable, as was leaflet number, as it not only has a small number of levels (3, 5, 7, and 9

leaflets), but each level is well separated from the others. To evaluate the accuracy of

relative node number, we treated it as a continuous variable, due to a high number of levels

(9) that continuously overlap with each other. Models revealed low accuracy, as the

accession was correctly determined only in 47.20% (Table **2**). The LDA model for the shape

of central leaflet showed no overlap for the accessions IK and MN9, but the remaining

accessions showed significant overlap (Fig. **5a**). The confusion matrix revealed that only two

accessions were correctly identified more than half the time (AM15 – 53.13% and IK –

71.43% prediction accuracy) (Fig. **5d**). The LDA model showed better success when

identifying the leaflet number (57.76% overall accuracy) and relative node number, where

the true and predicted values show significant, but moderate correlation (rho = 0.629, p <

0.0001) (Fig**. 5b, c, e, f**; Table **2**).

Table 2 Predictive power of genetic and developmental identities using the LDA model on

the central leaflet shape.

	Correct prediction [n]	False prediction [n]	Prediction accuracy [%]	Correlation coefficient [rho]	p value
Accession	76	85	47.20	NA	NA
Leaflet number	93	68	57.76	NA	NA
Relative node number	NA	NA	NA	0.629	< 0.0001

Fig. 5 Accession, leaflet number and relative node numbers prediction of *Cannabis* leaves

using the outline of central leaflets. Linear discriminant analysis (LDA) plots for (a) accession,

(b) leaflet number and (c) relative node number. In the lower row, the confusion matrices

show the true and predicted identities for (d) accessions, (e) leaflet number, and (f) relative

node number using the LDA model on the split test and train dataset.

Principal component analysis on modeled leaves (PCA)

Using the outline and landmark coordinates of 341 leaves, we modeled new theoretical

leaves, all with nine leaflets. Each leaf is defined by 3,591 pseudo-landmarks, which

overcomes the problems associated with variable leaflet numbers and permits dimension

reduction using PCA (Fig. **6a-c**) and the visualization of average *Cannabis* leaves (Fig. **6d-f**).

The first and second PCs account for 85.85% and 7.25% of the shape variation, respectively

(Fig. **6a-c**). Examining the PC1 and PC2 with Kruskal-Wallis test reveals that that accession,

leaflet number and relative node number all vary significantly along the first PC axis. The

variation along the PC2 for accession and leaflet number is less pronounced, however still

significant, while PC2 values for relative node numbers do not vary significantly (Fig. **6**; Table

3). This indicates that the changes in leaf shape between accessions are not independent

from developmental variation. That a facet of variation in accession leaf shape covaries with

developmental variation across the shoot in leaflet and relative node number suggests a

heterochronic mechanism by which accession differences in leaf shape arise from changes

in developmental timing, and contrasts with the historical focus on changes in timing arising

from plasticity (Goebel, 1908; Ashby, 1948).

Table 3 Kruskal-Wallis test was used to test the leaf shape variation along PC1 and PC2 for

accessions, leaflet number and relative node number.

	PC1		PC2	
	H	**p value**	**H**	**p value**
Accession	112.64	**p < 0.0001**	18.57	**p < 0.05**
Leaflet number	204.36	**p < 0.0001**	10.75	**p < 0.05**
Relative node number	49.73	**p < 0.0001**	2.98	p > 0.05

The average modeled leaf shapes show that the most pronounced change in leaf shape

between the accessions and during the development corresponds to narrow vs. wide

leaflets that are stereotypical descriptions of *sativa* vs. *indica* or *wide-* vs. *narrow- leaflet*

drug varieties. Furthermore, the leaves with the lower number of leaflets have more acute

leaflet tips, that slowly transition into acuminate. Additionally, the outer leaflets in the

leaves from lower nodes (and in certain accessions) are longer, compared to the central

leaflet, and become shorter higher up (Fig. **6d-e**).

Fig. 6 Principal component analysis (PCA) of the accessions performed on modeled leaves using the 3,591 pseudo-landmarks (a-c). The first PC explains 85.58% and the second 7.25% of variation. The images on the right show the average modeled leaf shapes for each of the (d) nine analyzed accessions, (e) leaflet number and (f) relative node number.

LDA and prediction of genetic and developmental identities on modeled leaves

As in the analysis of central leaflet shape before, we used LDA to model accession, leaflet number and relative node number as a function of all 3,591 pseudo-landmark points defining the complete modeled leaves (Fig. **7**). Accuracy of the model was calculated on the split dataset, treating accession and leaflet number as categorical and relative node number as continuous variable. LDA models for both accession and leaflet number were highly

accurate (73.29% and 99.38%, respectively) (Table **4**), significantly improving the results

obtained by analyzing solely the outline of the central leaflet (Table **2)**. The model for

relative node number is highly accurate as well, as inferred by a highly significant

Spearman's rank correlation coefficient value between actual and predicted values (rho =

0.747, p < 0.0001) (Table **4**).

Table 4 Predictive power of genetic and developmental identities using the LDA model on

the modeled leaves.

	Correct prediction [n]	False prediction [n]	Prediction accuracy [%]	Correlation coefficient [rho]	p value
Accession	118	43	73.29	NA	NA
Leaflet number	160	1	99.38	NA	NA
Relative node number	NA	NA	NA	0.747	< 0.0001

A confusion matrix reveals that the LDA model in most cases had a high accuracy for

predicting accession identity (Fig. **7d**; Table **4**), much higher, as compared to the accuracy

achieved by using only the outline of the central leaflet (Fig. **5d,** Table **2)**. Accessions IK, RO1,

and MN9 show practically no overlap in LDA space, while AM15, BNG, FUT75, HU1, IKL and

MAR show more overlap (Fig. **7a**). The model showed an almost 100% success rate in

determining leaflet number, again, much higher than before.

Results of both methods revealed that leaves with only 3 leaflets are markedly different

from the rest, and the prediction model on theoretical leaves consistently classified them

correctly (Fig. **7e**). Leaves with 5 to 9 leaflets showed less pronounced differences in shape,

resulting in a slightly lower accuracy of the prediction model for these cases. However, an

examination of the confusion matrix revealed that misclassifications only occurred once

between leaves with neighboring leaflet numbers (7 and 9 leaflets) (Fig. **7e**). The marked

difference in shape of leaves with 3 leaflets from the rest may suggest that this

developmental mechanism is biased towards variation at the base of the shoot. Similar to

leaflet number, the confusion matrix for the relative node model reveals high rates of

misclassification between the neighboring relative node numbers, as is expected, and leaves

from lower nodes were very rarely classified as those from higher nodes (Fig. **5f**). A

pronounced change in leaf shape occurs between the relative nodes 0.3 and 0.4, while the

shape changes in later relative nodes are more gradual (Fig. **7c**).

Fig. 7 Accession, leaflet number and relative node numbers of *Cannabis* leaves can be

predicted independently of each other using modeled leaves. Linear discriminant analysis

(LDA) plots for (a) accession, (b) leaflet number and (c) relative node number. In the lower

row, the confusion matrices show the true and predicted identities for (d) accessions, (e)

leaflet number, and (f) relative node number using the LDA model on the split test and train

dataset.

Compared to only using the modeled leaves, the accuracy of the LDA model did not improve

significantly when using a combined dataset. A confusion matrix revealed that the LDA

model (Fig. **S1**) was slightly less successful in accession identity classification (71.43%) but

was higher for leaflet number (100%). The Spearman's rank correlation coefficient was

slightly higher and highly significant (rho = 0.748, p < 0.0001) (Table **5**).

Table 5 Predictive power of genetic and developmental identities using the LDA model on a

combined dataset.

	Correct prediction [n]	False prediction [n]	Prediction accuracy [%]	Correlation coefficient [rho]	p value
Accession	115	46	71.43	NA	NA
Leaflet number	161	0	100	NA	NA
Relative node number	NA	NA	NA	0.787	< 0.0001

DISCUSSION

Like grapevines, striking variation in leaf shape (Fig. **1**) has historically played a significant

role in taxonomic classification of *Cannabis*. Leaf shape and differences in phyllotaxy were

among the characters Lamarck used to describe a new *Cannabis* species (Lamarck & Poiret,

1783). Anderson (1980) introduced a quantitative approach by quantifying the length-to-

width ratio of the central leaflet. Further studies using different characters—including plant

height, stem diameter, achene shape, and phytochemical profiles—to characterize

accessions have only confirmed the importance of leaf characteristics (Small *et al.*, 1976;

Hillig, 2005a). The central leaflet width-to-length ratio has been adopted by researchers as

one of the main characters for determining species, subspecies, biotypes and chemotypes of

Cannabis (Hillig, 2005a; Clarke & Merlin, 2013; McPartland & Small, 2020). However, this

method is only able to capture a limited aspect of leaf shape variation, neglecting other

important characteristics that we measure in this study, such as leaflet outlines, serrations,

angles, and relative changes in leaflet shape across the leaf. By modeling leaflet shape as a

function of leaflet number, we model theoretical leaves with the same number of leaflets

for which high densities of corresponding pseudo-landmarks capture high resolution shape

features (Fig. **2**). To validate the modeling approach, we have compared the outline of the

original central leaflet and the outline of the modeled theoretical central leaflet. The

Procrustes analysis showed that the two leaflets are very similar in shape, and that the

modeling is even able to preserve the serration pattern to some degree (Fig. **3c**). The

modeling approach validated using 10,000 bootstrap replicas confirmed the robustness of

the novel modeling approach (Fig. **3d**). This method can be applied not only on palmately

composed leaves as in *Cannabis* but is also possible to use on pinnate and lobed leaves. To

demonstrate the proof of concept, we applied the method to a pinnate leaf of *Cardamine*

flexuosa With. and lobate leaf of *Quercus macrocarpa* Michx. (Fig. **8**), showing the method

could be applied in other leaf types. However, the method needs to be improved before

being applied to other species but shows the possible utility of intra-leaf modeling.

Fig. 8 Intra-leaf modeling of leaflets and lobes extended to pinnate leaves: Leaves from (a)

Cardamine flexuosa and (b) *Quercus marcocarpa*. Leaflets and lobes are defined by 100

equidistant pseudo-landmarks on each side, each defined by three landmarks, two at the

base and one at the tip. Large points are placed every 20 pseudo-landmarks to emphasize

that leaflet outlines are defined by points. The landmarks defining the base of each leaflet or

lobe are aligned to the rachis or midvein and the transformed leaflets and lobes have been

oriented parallel to the rachis, as defined by the landmarks at their base. The modeled

leaflets and lobes are created from 2nd degree polynomial models for each *x* and *y*

coordinate value for each pseudo-landmark as a function of leaflet or lobe number. From

these models, an equivalent number of modeled leaflets or lobes can be reconstructed (in

this case, five), permitting morphometric analysis.

The method presented in this study can accurately determine accession based on leaf

shape, regardless of its developmental stage (Fig. **7a, d**). The method not only works

effectively on stabilized or cloned cultivar accessions but also on wild or feral accessions

cultivated from seed that can exhibit distinct plant phenotypes (Table **1**), indicating its

robustness and potential value in future germplasm classification. Compared to the low

accuracy and prediction ability of the previously known methods (W/L ratio and shape

analysis of central leaflets), the newly proposed method demonstrates significantly

improved results (Table **2, 4, S2, S3**). The combined dataset of both, data for modeled leaves

and outline of the central leaflet, did not return significantly better results, further

confirming the effectiveness of the new modelling approach.

When observing the shape changes between averaged leaves for accessions and between

developmental stages, the most obvious are changes in leaflet widths, similar to

stereotypical classifications of *sativa* and *indica* plants or *wide-* vs. *narrow- leaflet* drug

varieties. However, other important changes in shape occur, such as transition from acute

to acuminate leaflet tip and changes in the relative length of outer most leaflets compared

to the central leaflet, that previous methods could not successfully capture (Fig. **6d-f**). The

reliance on the non-quantitative leaf shape descriptors in previous methods has led to

numerous cultivars with unreliable names, inconsistent genetic origins, and phytochemical

profiles (Sawler *et al.*, 2015; Schwabe & McGlaughlin, 2019; Jin *et al.*, 2021a; Watts *et al.*,

2021). For example, Jin *et al.* (2021b) conducted a study on clones of 21 cultivars and found

a strong negative correlation between the width and length ratios of central leaflets and

CBD, and a positive correlation with THC; however, Vergara *et al.* (2021) and Murovec *et al.*

(2022) were unable to confirm these findings. All three studies used low-resolution

morphometric approaches. Sex of the plants also plays a crucial role in the cannabis

industry, where the presence of male plants and inevitable pollination leads to decreases in

cannabinoid production as plants shift the use of energy into seed development. Several

methods have been employed to differentiate between male and female plants at early

stages, but only genetic methods were successful so far (Toth *et al.*, 2020; Prentout *et al.*,

2020; Campbell *et al.*, 2021; Balant *et al.*, 2022; Torres *et al.*, 2022). Our results quantify the

variation in leaf shape between accessions that can potentially be used to classify accessions

and predict chemical profiles and plant sex faster and more accurately.

Unlike grapevine, where developmental variance is orthogonal and separate from genetic

variance, in *Cannabis* these two factors are correlated. That the developmental source of

variation is colinear with accession identity suggests that part of the differences between

accession leaf shape is explained by shifts in developmental timing, or heterochrony.

Cannabis plants demonstrate extreme phenotypic plasticity depending on the

environmental conditions in which they grows (Small, 2015). Some *Cannabis* accessions are

photoperiod dependent and can remain in vegetative phase for longer periods of time

under long-day conditions (typically 18h darkness and 6h light), until the transition to short-

day (12h of darkness and 12h of light) induces the formation of the apical inflorescence.

Previous investigations showed that other morphological changes, such as decrease in leaf

area, number of leaflets per leaf and serration number, occur after the change in the

environmental conditions one or two nodes after (Heslop-Harrison & Heslop-Harrison, 1958;

Hesami *et al.*, 2023). However, differences, especially in flowering time and growth rates

between cultivars have been observed before (de Meijer & Keizer, 1996; Hillig, 2005a;

Spitzer-Rimon *et al.*, 2019; Carlson *et al.*, 2021; Naim-Feil *et al.*, 2021; Stack *et al.*, 2021;

Chen *et al.*, 2022) and differences in cannabinoid profiles, leaflet index and phenological

development were proposed as characteristics to discriminate between them (de Meijer &

Keizer, 1996). Heterochronic shifts are apparent in the differential rates in which accessions

increase leaflet number across nodes, as well as maximum and average leaflet counts across

accessions (Fig. **1**). Remarkably, stages in developmental timing are conserved despite being

shifted. For example, a significant shape change exhibited between the leaves with 3 and

leaves with 5 leaflets, with leaflets becoming more acuminate and narrower. In contrast,

changes in shape between leaves with a higher number of leaflets were more gradual.

Additionally, we observed a similar shift in leaf shape between the nodes 0.3 and 0.4,

potentially indicating a transition between the juvenile and adult phases of leaf

development. Similar results were obtained in previous research. Spitzer-Rimon *et al.* (2022)

demonstrated that flowering buds were initiated at node 7, while Moliterni *et al.* (2004)

analyzing a different cultivar, found developing flower buds in the 4th node, suggesting that

transitions in growth phases are conserved but not synchronized across cultivars. Due to the

differences in developmental timing between accessions, the use of continuous models

along the shoot could further improve the success predicting accession identity, as was the

case in grapevine (Bryson *et al.*, 2020).

Conclusions

In grapevine, leaf shape has long been utilized for variety identification. However, in the

case of *Cannabis*, previous attempts were hindered by the variability in leaflet numbers. In

this study, we present a pioneering method that successfully addresses this issue. By

generating theoretical leaves with customizable leaflet counts, we can now employ high-

resolution morphometric techniques to accurately classify different wild/feral and cultivated

Cannabis accessions. Through the use of 3,591 densely placed pseudo-landmarks, we were

able to predict the accession identity with almost 74% accuracy. The method works well not

only on stabilized cultivars, but also on phenotypically more variable wild/feral accessions

grown from seed. Unifying the number of leaflets allowed us, for the first time, to make

comparisons among several leaves along the main axis, enabling us to investigate

developmental changes in leaf shape and detect heterochronic mechanisms influencing the

leaf shape in *Cannabis*. The implications of this new high-resolution method in both the

cannabis industry and research extend beyond its role in determining *Cannabis* accessions.

It also offers a promising tool for developmental studies, and for studying the correlation

between leaf shape and phytochemical profiles and the sex of the plants, where lower-

resolution methods provided inconclusive results so far. The method presented here offers

a fast, effective, robust, and low-cost tool that can aid the future classification of *Cannabis*

germplasm. Furthermore, the use of this methodology extends beyond *Cannabis*, and can

be applied to numerous other plant species with palmate, pinnate, and lobate leaves with

varying numbers of lobes and leaflets where the use of geometric morphometrics methods

was not previously possible to this extent.

ACKNOWLEDGEMENTS

This research was supported by projects WECANN (CGL2017-84297-R, Ministerio de Ciencia,

Innovación y Universidades), Generalitat de Catalunya (grant number 2021SGR00315) and

M. Balant FPI predoctoral contract of the Ministerio de Ciencia, Innovación y Universidades

(PRE2018-083226). This work is also supported by NSF Plant Genome Research Program

awards IOS-2310355, IOS-2310356, and IOS-2310357. This project was supported by the

USDA National Institute of Food and Agriculture, and by Michigan State University

AgBioResearch. We would like to thank Joan Uriach Marsal from Uriach Laboratories for
additional financial support. We would also like to acknowledge Cannaflos—Gesellschaft für
medizinisches Cannabis mbH for providing seed of accessions IK, IKL and MAR analyzed in
the study, Carlos Sáez, Carlos Ribelles, Airy Gras, Joan Vallès, Magsar Urgamal, Shagdar
Tsooj, Marine Oganesian and Nina Stepanyan-Gandilyan for help with the sample collection
and the cultivation process, and Paula Bruna for helpful advice on improving the quality of
scanned images.

COMPETING INTERESTS

None declared.

AUTHOR CONTRIBUTIONS

MB and DHC conceived the study, with OH, TG and DV inputs in a preliminary design phase
of the project. MB, TG and DV cultivated the plants. MB and OH selected the method for
imaging *Cannabis* leaves. MB and TG scanned the leaves used in the study. MB and DHC
developed the morphometric method and MB analyzed the data. MB and DHC wrote the
first draft of the paper, that all authors read, commented, and edited.

DATA AVAILABILITY

The datasets and code for the method developed here are freely available on GitHub
(https://github.com/BalantM/Cannabis_leaf_morpho_updated,
https://github.com/DanChitwood/pinnate_leaf_modeling).

ORCID

Manica Balant https://orcid.org/0000-0002-4316-8511

Teresa Garnatje https://orcid.org/0000-0001-6295-6217

Daniel Vitales https://orcid.org/0000-0002-3109-5028

Oriane Hidalgo https://orcid.org/0000-0002-1547-8627

Daniel H. Chitwood https://orcid.org/0000-0003-4875-1447

SUPPORTING INFORMATION

The datasets and code generated and analyzed in this study are available on GitHub

(https://github.com/BalantM/Cannabis_leaf_morpho_updated,

https://github.com/DanChitwood/pinnate_leaf_modeling).

REFERENCES

Abràmoff MD, Magalhães PJ, Ram SJ. **2004**. Image processing with ImageJ Part II.

Biophotonics International **11**: 36–43.

Anderson LC. **1980**. Leaf Variation among *Cannabis* Species from a Controlled Garden.

Botanical Museum leaflets, Harvard University **28**: 61–69.

Ashby E. **1948**. Studies in the morphogenesis of leaves. I. An essay on leaf shape. *New*

Phytologist **47**: 153–176.

Balant M, González Rodríguez R, Garcia S, Garnatje T, Pellicer J, Vallès J, Vitales D, Hidalgo

O. **2022**. Novel Insights into the Nature of Intraspecific Genome Size Diversity in *Cannabis*

sativa L . *Plants* **11**: 2736.

Biot E, Cortizo M, Burguet J, Kiss A, Oughou M, Maugarny-Calès A, Gonçalves B, Adroher B,

Andrey P, Boudaoud A, *et al.* 2016. Multiscale quantification of morphodynamics:

MorphoLeaf, software for 2-D shape analysis. *Development* **143**: 3417–3428.

Bookstein FL. **1997**. *Morphometric Tools for Landmark Data: Geometry and Biology*.

Cambrige: Cambridge University Press.

Bradski G. **2000**. The OpenCV Library. *Dr. Dobb's Journal: Software Tools for the Professional*

Programmer **25**: 120–123.

Bryson AE, Wilson Brown M, Mullins J, Dong W, Bahmani K, Bornowski N, Chiu C, Engelgau

P, Gettings B, Gomezcano F, *et al.* 2020. Composite modeling of leaf shape along shoots

discriminates *Vitis* species better than individual leaves. *Applications in Plant Sciences* **8**:

e11404.

Campbell LG, Peach K, Wizenberg SB. **2021**. Dioecious hemp (*Cannabis sativa* L.) plants do

not express significant sexually dimorphic morphology in the seedling stage. *Scientific*

Reports **11**: 1–8.

Carlson CH, Stack GM, Jiang Y, Taşklran B, Cala AR, Toth JA, Philippe G, Rose JKC, Smart
CD, Smart LB. **2021**. Morphometric relationships and their contribution to biomass and
cannabinoid yield in hybrids of hemp (*Cannabis sativa*). *Journal of Experimental Botany* **72**:
7694–7709.

Chen X, Guo H, Zhang Q, Wang L, Guo R, Zhan Y, Lv P, Xu Y, Guo M, Zhang Y, *et al.* 2022.
Whole-genome resequencing of wild and cultivated cannabis reveals the genetic structure
and adaptive selection of important traits. *BMC Plant Biology* **22**: 371.

Chitwood DH. **2021**. The shapes of wine and table grape leaves: An ampelometric study
inspired by the methods of Pierre Galet. *Plants People Planet* **3**: 155–170.

Chitwood DH, Klein LL, O'Hanlon R, Chacko S, Greg M, Kitchen C, Miller AJ, Londo JP.
2016a. Latent developmental and evolutionary shapes embedded within the grapevine leaf.
New Phytologist **210**: 343–355.

Chitwood DH, Mullins J, Migicovsky Z, Frank M, VanBuren R, Londo JP. **2021**. Vein-to-blade
ratio is an allometric indicator of leaf size and plasticity. *American Journal of Botany* **108**:
571–579.

Chitwood DH, Ranjan A, Martinez CC, Headland LR, Thiem T, Kumar R, Covington MF,
Hatcher T, Naylor DT, Zimmerman S, *et al.* 2014. A modern ampelography: A genetic basis
for leaf shape and venation patterning in grape. *Plant Physiology* **164**: 259–272.

Chitwood DH, Rundell SM, Li DY, Woodford QL, Yu TT, Lopez JR, Greenblatt D, Kang J,
Londo JP. **2016b**. Climate and developmental plasticity: Interannual plasticity in grapevine
leaf morphology. *Plant Physiology* **170**: 1480–1491.

Chitwood DH, Sinha NR. **2016**. Evolutionary and Environmental Forces Sculpting Leaf
Development. *Current Biology* **26**: R297–R306.

Clarke R, Merlin M. **2013**. Cannabis*: Evolution and ethnobotany*. Berkley, Los Angeles,
London: University of California Press.

Demmings EM, Williams BR, Lee C-R, Barba P, Yang S, Hwang C-F, Reisch BI, Chitwood DH,
Londo JP. **2019**. Quantitative Trait Locus Analysis of Leaf Morphology Indicates Conserved
Shape Loci in Grapevine. *Frontiers in Plant Science* **10**: 1373.

Emboden WA. 1974. *Cannabis* - a polytypic genus. *Economic Botany* **28**: 304–310.

Failmezger H, Lempe J, Khadem N, Cartolano M, Tsiantis M, Tresch A. **2018**. MowJoe: a
method for automated-high throughput dissected leaf phenotyping. *Plant Methods* **14**: 27.

Galet P. **1952**. *Précis d'Ampélographie Pratique*. Digitized by Google Books from Cornell

780 University.

Galet P, trans. Morton L. **1979**. *A Practical Ampelography: Grapevine Identification*. Cornell University Press (English translation).

Goebel K. **1908**. *Einleitung in die experimentelle Morphologie der Pflanzen*. Leipzig.

Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, Wieser E, Taylor J, Berg S, Smith NJ, *et al.* 2020. Array programming with NumPy. *Nature* **585**: 357–362.

Hesami M, Pepe M, Jones AMP. **2023**. Morphological Characterization of *Cannabis sativa* L. Throughout Its Complete Life Cycle. *Plants* **12**: 3646.

Heslop-Harrison J, Heslop-Harrison Y. **1958**. Studies on flowering-plant growth and organogenesis-III. Leaf shape changes associated with flowering and sex differentiation in *Cannabis sativa*. *Proceedings of the Royal Irish Academy. Section B: Biological, Geological, and Chemical Science* **59**: 257–283.

Hillig KW. **2004**. A multivariate analysis of allozyme variation in 93 *Cannabis* accessions from the VIR germplasm collection. *Journal of Industrial Hemp* **9**: 5–22.

Hillig KW. **2005a**. A Systematic Investigation of *Cannabis*.

Hillig KW. **2005b**. Genetic evidence for speciation in *Cannabis* (Cannabaceae). *Genetic Resources and Crop Evolution* **52**: 161–180.

Hunter JD. **2007**. Matplotlib: a 2D graphics environment. *Computing in Science and Engineering* **9**: 90–95.

Hurgobin B, Tamiru-Oli M, Welling MT, Doblin MS, Bacic A, Whelan J, Lewsey MG. **2021**. Recent advances in *Cannabis sativa* genomics research. *New Phytologist* **230**: 73–89.

Jin D, Henry P, Shan J, Chen J. **2021a**. Classification of cannabis strains in the Canadian market with discriminant analysis of principal components using genome-wide single nucleotide polymorphisms. *PLOS ONE* **16**: e0253387.

Jin D, Henry P, Shan J, Chen J. **2021b**. Identification of phenotypic characteristics in three chemotype categories in the genus *Cannabis*. *HortScience* **56**: 481–490.

Kuhl FP, Giardina CR. **1982**. Elliptic Fourier features of a closed contour. *Computer Graphics and Image Processing* **18**: 236–258.

Lamarck J-B, Poiret J-L-M. **1783**. *Encyclopédie méthodique. Botanique*. Paris, Panckoucke; Liège, Plomteux.

Lapierre É, Monthony AS, Torkamaneh D. **2023**. Genomics-based taxonomy to clarify

cannabis classification. *Genome* **66**: 202–211.

McPartland JM. **2018**. *Cannabis* Systematics at the Levels of Family, Genus, and Species.

Cannabis and Cannabinoid Research **3**: 203–212.

McPartland JM, Guy GW. **2017**. Models of *Cannabis* Taxonomy, Cultural Bias, and Conflicts

between Scientific and Vernacular Names. *Botanical Review* **83**: 327–381.

McPartland JM, Small E. **2020**. A classification of endangered high-THC cannabis (*Cannabis*

sativa subsp. *indica*) domesticates and their wild relatives. *PhytoKeys* **144**: 81–112.

de Meijer EPM, Keizer LCP. **1996**. Patterns of diversity in *Cannabis*. *Genetic Resources and*

Crop Evolution **43**: 41–52.

Migicovsky Z, Swift JF, Helget Z, Klein LL, Ly A, Maimaitiyiming M, Woodhouse K, Fennell

A, Kwasniewski M, Miller AJ, *et al.* 2022. Increases in vein length compensate for leaf area

lost to lobing in grapevine. *American Journal of Botany* **109**: 1063–1073.

Moliterni VMC, Cattivelli L, Ranalli P, Mandolino G. **2004**. The sexual differentiation of

Cannabis sativa L.: A morphological and molecular study. *Euphytica* **140**: 95–106.

Murovec J, Eržen JJ, Flajšman M, Vodnik D. **2022**. Analysis of Morphological Traits,

Cannabinoid Profiles, THCAS Gene Sequences, and Photosynthesis in Wide and Narrow

Leaflet High-Cannabidiol Breeding Populations of Medical Cannabis. *Frontiers in Plant*

Science **13**: 786161.

Naim-Feil E, Pembleton LW, Spooner LE, Malthouse AL, Miner A, Quinn M, Polotnianka

RM, Baillie RC, Spangenberg GC, Cogan NOI. **2021**. The characterization of key physiological

traits of medicinal cannabis (*Cannabis sativa* L.) as a tool for precision breeding. *BMC Plant*

Biology **21**: 1–15.

Niklas KJ. **1994**. *Plant allometry: the scaling of form and process*. Chicago: University of

Chicago Press.

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, Blondel M,

Prettenhofer P, Weiss R, Dubourg V, *et al.* 2011. Scikit-learn: Machine Learning in Python.

Journal ofMachine Learning Research **12**: 2825–2830.

Potter DJ. **2009**. *Optimisation Of* Cannabis sativa *L. As A Phytopharmaceutical [PhD Thesis]*.

London: King's College London, Department of Pharmaceutical Science Research.

Prentout D, Razumova O, Rhoné B, Badouin H, Henri H, Feng C, Käfer J, Karlov G, Marais

GAB. **2020**. An efficient RNA-seq-based segregation analysis identifies the sex chromosomes

of *Cannabis sativa*. *Genome Research* **30**: 164–172.

Quimby MW, Doorenbos NJ, Turner CE, Masoud A. **1973**. Mississippi-Grown Marihuana -

Cannabis sativa Cultivation and Observed Morphological Variations. *Economic Botany* **27**:

117–127.

Ravaz L. **1902**. *Les vignes américaines: Porte-greffes et producteurs directs*. Goulet,

Montpellier and Paris: Digitized by Google Books from Cornell University.

Ren G, Zhang X, Li Y, Ridout K, Serrano-Serrano ML, Yang Y, Liu A, Ravikanth G, Nawaz MA,

Mumtaz AS, *et al.* 2021. Large-scale whole-genome resequencing unravels the

domestication history of *Cannabis sativa*. *Science Advances* **7**: eabg2286.

Sawler J, Stout JM, Gardner KM, Hudson D, Vidmar J, Butler L, Page JE, Myles S. **2015**. The

Genetic Structure of Marijuana and Hemp. *PLOS ONE* **10**: e0133292.

Schultes RE, Klein WM, Plowman T, Lockwood TE. **1974**. *Cannabis*: An example of

taxonomic neglect. *Botanical Museum Leaflets, Harvard University* **23**: 337–367.

Schwabe AL, McGlaughlin ME. **2019**. Genetic tools weed out misconceptions of strain

reliability in *Cannabis sativa*: implications for a budding industry. *Journal of Cannabis*

Research **1**: 3.

Small E. **2015**. Evolution and Classification of *Cannabis sativa* (Marijuana, Hemp) in Relation

to Human Utilization. *Botanical Review* **81**: 189–294.

Small E, Beckstead HD. **1973**. Cannabinoid Phenotypes in *Cannabis sativa*. *Nature* **245**: 147–

148.

Small E, Cronquist A. **1976**. A Practical and Natural Taxonomy for *Cannabis*. *Taxon* **25**: 405–

435.

Small E, Jui PY, Lefkovitch LP. **1976**. A Numerical Taxonomic Analysis of *Cannabis* with

Special Reference to Species Delimitation. *Systematic Botany* **1**: 67–84.

Spitzer-Rimon B, Duchin S, Bernstein N, Kamenetsky R. **2019**. Architecture and Florogenesis

in Female *Cannabis sativa* Plants. *Frontiers in Plant Science* **10**: 350.

Spitzer-Rimon B, Shafran-Tomer H, Gottlieb GH, Doron-Faigenboim A, Zemach H,

Kamenetsky-Goldstein R, Flaishman M. **2022**. Non-photoperiodic transition of female

cannabis seedlings from juvenile to adult reproductive stage. *Plant Reproduction* **35**: 265–

277.

Stack GM, Toth JA, Carlson CH, Cala AR, Marrero-González MI, Wilk RL, Gentner DR,

Crawford JL, Philippe G, Rose JKC, *et al.* 2021. Season-long characterization of high-

cannabinoid hemp (*Cannabis sativa* L.) reveals variation in cannabinoid accumulation,

flowering time, and disease resistance. *GCB Bioenergy* **13**: 546–561.

Terpilowski M. **2019**. scikit-posthocs: Pairwise multiple comparison tests in Python. *Journal*

of Open Source Software **4**: 1169.

The Global Cannabis Report, 3rd Edition. **2022**. London, Berlin, Barcelona: Prohibition

Partners.

Torres A, Pauli C, Givens R, Argyris J, Allen K, Monfort A, Gaudino RJ. **2022**.

High-throughput methods to identify male *Cannabis sativa* using various genotyping

methods. *Journal of Cannabis Research* **4**: 57.

Toth JA, Stack GM, Cala AR, Carlson CH, Wilk RL, Crawford JL, Viands DR, Philippe G, Smart

CD, Rose JKC, *et al.* 2020. Development and validation of genetic markers for sex and

cannabinoid chemotype in *Cannabis sativa* L. *GCB Bioenergy* **12**: 213–222.

Vergara D, Feathers C, Huscher EL, Holmes B, Haas JA, Kane NC. **2021**. Widely assumed

phenotypic associations in *Cannabis sativa* lack a shared genetic basis. *PeerJ* **9**: e10672.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E,

Peterson P, Weckesser W, Bright J, *et al.* 2020. SciPy 1.0: fundamental algorithms for

scientific computing in Python. *Nature Methods* **17**: 261–272.

Watts S, McElroy M, Migicovsky Z, Maassen H, van Velzen R, Myles S. **2021**. *Cannabis*

labelling is associated with genetic variation in terpene synthase genes. *Nature Plants* **7**:

1330–1334.

Zhukovskii PM. **1971**. *Cultivated plants and their wild relatives*. Leningrad, USSR, Kolos.

896