Flexible gating between subspaces by a disinhibitory motif: a neural network model of internally guided task switching

Yue Liu*, Xiao-Jing Wang*

New York University

Abstract

Behavioral flexibility relies on the brain's ability to switch rapidly between multiple tasks, even when the task rule is not explicitly cued but must be inferred through trial and error. The underlying neural circuit mechanism remains poorly understood. We investigated recurrent neural networks (RNNs) trained to perform an analog of the classic Wisconsin Card Sorting Test. The networks consist of two modules responsible for rule representation and sensorimotor mapping, respectively, where each module is comprised of a circuit with excitatory neurons and three major types of inhibitory neurons. We found that rule representation by self-sustained persistent activity across trials, error monitoring and gated sensorimotor mapping emerged from training. Systematic dissection of trained RNNs revealed a detailed circuit mechanism that is consistent across networks trained with different hyperparameters. The networks' dynamical trajectories for different rules reside in separate subspaces of population activity; they become virtually identical and performance was reduced to chance level when dendrite-targeting somatostatin-expressing interneurons were silenced, demonstrating that rule-based gating critically depends on the disinhibitory motif.

Introduction

A signature of cognitive flexibility is the ability to adapt to a changing task demand. Oftentimes, the relevant task is not explicitly instructed, but needs to be inferred from previous experiences. In laboratory studies, this behavioral flexibility is termed un-cued task switching. A classic task to evaluate this ability is the Wisconsin Card Sorting Test

*Corresponding author. Email: yl4317@nyu.edu, xjwang@nyu.edu

(WCST) [1]. During this task, subjects are presented with an array of cards, each with multiple features, and should respond based on the relevant feature dimension (i.e. the task rule) that changes across trials. Crucially, subjects are not instructed on when the rule changes, but must infer the currently relevant rule based on the outcome of previous trials. Intact performance on un-cued task switching depends on higher-order cortical areas such as the prefrontal cortex (PFC) [2, 3, 4, 5, 6], which has been proposed to represent the task rule and modulate the activity of other cortical areas along the sensorimotor pathway [7].

Four essential neural computations must be implemented by the neural circuitry underlying un-cued task switching. First, it should maintain an internal representation of the task rule across multiple trials when the rule is unchanged. Second, soon after the rule switches, the animal will inevitably make errors and receive negative feedback, since the switches are un-cued. This negative feedback should induce an update to the internal representation of the task rule. Third, the neural signal about the task rule should be communicated to the brain regions responsible for sensory processing and action selection. Fourth, this rule signal should be integrated with the incoming sensory stimulus to produce the correct action.

Prior work has identified neural correlates of cognitive variables presumed to underlie these computations including rule [8], feedback [8, 9, 10] and conjunctive codes for sensory, rule, and motor information [11]. In addition, different types of inhibitory neurons are known to play different functional roles in neural computation: while parvalbumin (PV)-expressing interneurons are suggested to underlie feedforward inhibition [12], interneurons that express somatostatin (SST) and vasoactive intestinal peptide (VIP) have been proposed to mediate top-down control [13, 14, 15, 16]. In particular, SST and VIP neurons form a disinhibitory motif [17, 18, 19] that has been hypothesized to instantiate a gating mechanism for flexible routing of information in the brain [20]. However, there is currently a lack of mechanistic understanding of how these neural representations and cell-type-specific mechanisms work together to accomplish un-cued task switching.

To this end, we used computational modeling to formalize and discover mechanistic hypotheses. In particular, we used tools from machine learning to train a collection of biologically informed recurrent neural networks (RNNs) to perform an analog of the WCST used in monkeys [21, 8, 9]. Training RNN [22] does not presume a particular circuit solution, enabling us to explore potential mechanisms. For this purpose, it is crucial that the model is biologically plausible. To that end, each RNN was set up to have two modules: a "PFC" module for rule maintenance and switching and a "sensorimotor" module for executing the sensorimotor transformation conditioned on the rule. To explore the potential functions of different neuronal types in this task, each module of our network consists of excitatory neurons with somatic and dendritic compartments as well as PV, SST and VIP inhibitory

neurons, where the connectivity between cell types is constrained by experimental data (Methods).

After training, we performed extensive dissection of the trained models to reveal that close interplay between local and across-area processing was essential for solving the WCST. First, we found that abstract cognitive variables were distinctly represented in the PFC module. In particular, two subpopulations of excitatory neurons emerge in the PFC module - one encodes the task rule and the other shows nonlinear mixed-selectivity for rule and negative feedback. Notably, neurons with similar response profiles have been reported in neurophysiological recordings of monkeys performing the same task [8, 9]. Second, we identified interesting structures in the local connectivity between different neuronal assemblies within the PFC module, which enabled us to compress the high-dimensional PFC module down to a low-dimensional simplified network. Importantly, the neural mechanism for maintaining and switching rule representation is readily interpretable in the simplified network. Third, we found that the rule information in the PFC module is communicated to the sensorimotor module via structured long-range connectivity patterns along the monosynaptic excitatory pathway, the di-synaptic pathway that involves PV neurons, as well as the trisynaptic disinhibitory pathway that involves SST and VIP neurons. In addition, different dendritic branches of the same excitatory neuron in the sensorimotor module can be differentially modulated by the task rule depending on the sparsity of the local connections from the dendrite-targeting SST inhibitory neurons. Fourth, single neurons in the sensorimotor module show nonlinear mixed selectivity to stimulus, rule and response, which crucially depends on the activity of the SST neurons. On the population level, the neural trajectories for the sensorimotor neurons during different task rules occupy nearly orthogonal subspaces, which is disrupted by silencing the SST neurons. Lastly, we found structured patterns of input and output connections for the sensorimotor module, which enables appropriate rule-dependent action selection. These results are consistent across dozens of trained RNNs with different types of dendritic nonlinearities and initializations, therefore pointing to a common neural circuit mechanism underlying the WCST.

Results

Training modular recurrent neural networks with different types of inhibitory neurons

We trained a collection of modular RNNs to perform the WCST. Each RNN consists of two modules: the "PFC" module receives an input about the outcome of the previous trial, and was trained to output the current rule; the "sensorimotor" module receives the sensory input and was trained to generate the correct choice (Figure 1b). The inputs and outputs were represented by binary vectors (Figure 1b, Methods) Each module was

endowed with excitatory neurons with somatic and two dendritic compartments, as well as three major types of genetically-defined inhibitory neurons (PV, SST and VIP). Different types of neurons have different inward and outward connectivity patterns constrained by experimental data in a binary fashion (Methods, Figure 1b). The somata of all neurons were modeled as standard leaky units with a rectified linear activation function. The activation of the dendritic compartments, which can be viewed as a proxy for the dendritic voltage, is a nonlinear sigmoidal function of the excitatory and inhibitory inputs they receive (Methods). The specific form of the nonlinearity is inspired by experiments showing that inhibition acts subtractively or divisively on the dendritic nonlinearity function depending on its relative location to the excitation along the dendritic branch [23]. Therefore, we trained a collection of RNNs, each with either subtractive or divisive dendritic nonlinearity, to explore the effect of dendritic nonlinearity on the network function.

The task we trained the network on is a WCST variant used in monkey experiments [21, 8, 9, 6] (Figure 1a). During each trial, a reference card with a particular color and shape is presented on the screen for 500 ms, after which three test cards appear around the reference card for another 500 ms. Each card can have one of the two colors (red or blue) and one of the two shapes (square or triangle). A choice should be made for the location that contains the test card that has the same relevant feature (color or shape) as the reference card, after which the outcome of the trial is given, followed by an inter-trial interval. The relevant feature to focus on, or the task rule, changes randomly every few trials. Critically, the rule changes were not cued, requiring the network to memorize the rule of the last trial using its own dynamics. Therefore, the network dynamics should be carried over between consecutive trials, rather than reset at the end of each trial as has been done traditionally [24, 25]. To this end, the network operated continuously across multiple trials during training, and the loss function was aggregated across multiple trials (Methods). We use supervised learning to adjust the strength of all the connections (input, recurrent and output) by minimizing the mean squared error between the output of both modules and the desired output (rule for the PFC module and response for the sensorimotor module). Notably, only the connections between certain cell types are non-zero and can be modified. This is achieved using a mask matrix, similar to [26] (Methods).

After training converged, we tested the models by running them continuously across 100 trials of WCST with 10 rule switches at randomly chosen trials. The networks made a single error after each rule switch, and were able quickly switch to the new rule and maintain good performance (Figure 1c, d). Correspondingly, single neurons from both modules exhibited rule-modulated persistent activity that lasted several trials (Supplementary Figure 1). Networks trained with different hyperparameters all had performance well above chance (Figure 1e).

In previous studies, the performance of monkeys during this task showed substantial
variability across different studies as well as different sessions with a study. The number
of error trials that monkeys take to switch to the new rule ranges from one trial to tens of
trials [21, 8, 9, 6]. Therefore, the model behavior does not match perfectly the wide range
of behavior exhibited by monkeys. This is due to the fact that we used supervise learning
to optimize the performance of the model, whereas monkeys may use a combination of
strategies during WCST. This point will be addressed further in the Discussion section.
Nevertheless, the mechanisms that the models used to accomplish the task should still shed
some light on the neural circuit mechanism underlying the task. In the following sections,
we will "open the black box" to understand the mechanism the networks used to perform
the WCST.

Two rule attractor states in the PFC module maintained by interactions between modules

We first dissected the PFC module, which was trained to represent the rule. Since
there are two rules in the WCST task we used, we expected that the PFC module might have
two attractor states corresponding to the two rules. Therefore, we examined the attractor
structure in the dynamical landscape of the PFC module by initializing the network at
states chosen randomly from the trial, and evolving the network autonomously (without any
input) for 500 time steps (which equals 5 seconds in real time). Then, the dynamics of the
PFC module during this evolution was visualized by applying principal component analysis
to the population activity. The PFC population activity settled into one of two different
attractor states depending on the rule that the initial state belongs to (Supplementary
Figure 2a). Therefore, there are two attractors in the dynamical landscape of the PFC
module, corresponding to the two rules.

Historically, persistent neural activity corresponding to attractor states were first
discovered in the PFC [27, 28, 29, 30]. However, more recent experiments found persistent
neural activity in multiple brain regions, suggesting that long-range connections between
brain regions may be essential for generating persistent activity [31, 32, 33, 34]. Inspired
by these findings, we wondered if the PFC module in our network could support the two
rule attractor states by itself, or that the long-range connections between the PFC and the
sensorimotor module are necessary to support them. To this end, we lesioned the inter-
modular connections in the trained networks and repeated the simulation. Interestingly, we
found that for the majority of the trained networks (42 out of 52), their PFC activity settled
into a trivial fixed point corresponding to an inactive state (Supplementary Figure 2b, c).
This result shows that the two rule attractor states in these networks are dependent on the
interactions between the PFC and the sensorimotor modules.

Figure 1: (Caption next page.)

Figure 1: (Previous page.) **Model setup and performance.**
a. The schematic of the WCST task. Subjects are required to choose the card that matches the reference card at the center in either shape or color, depending on a hidden rule that switches after a number of trials.
b. The RNN contains a "PFC" module and a "sensorimotor" module. The PFC module receives an input about the feedback of the previous trial, and was trained to produce the current rule. The sensorimotor module receives the sensory input and was trained to produce the correct choice. The input to the PFC module about the feedback is represented by a two-dimensional binary vector. The input to the sensorimotor module represents the features of the cards on the screen. Each card is represented by a four-dimensional binary vector, where the two non-zero entries represent the color and shape of the card. The target output of the PFC module about the correct rule is represented by a two-dimensional binary vector. The target output of the sensorimotor module about the correct choice is represented by a three-dimensional binary vector. Each module is endowed with excitatory neurons and three types of inhibitory neurons: PV, SST and VIP. The cell-type-specific connectivity is constrained by experimental data (Methods). Bottom panel shows the decomposition of the model architecture into the input and output connectivity (left, magenta. The dashed line from PFC to rule represents the fact that the PFC module was trained to represent the rule but there are no explicit rule output neurons in the model), the local recurrent connectivity (middle, black) and inter-modular connectivity (green, right). All connections were trained. Each excitatory neuron is modeled with a somatic and two dendritic compartments. Inset shows for the two types of dendritic nonlinearities used the relationship between the excitatory input onto the dendrite and the dendritic activity for different levels of inhibitory inputs.
c. The performance of the model during testing, for an example network. The network made one error after each rule switch (black vertical lines) and quickly recovered its performance.
d. Performance as a function of trial number after a rule switch, for the same example network as in **c**.
e. Testing performance for all trained networks.

Two emergent subpopulations of excitatory neurons in the PFC module

For the PFC module to keep track of the current rule in effect, the module should stay in the same rule attractor state after receiving positive feedback, but transition to the other rule attractor state after receiving negative feedback. We reasoned that this network function might be mediated by single neurons that are modulated by the task rule and negative feedback, respectively. Therefore, we set out to look for these single neurons.

In the PFC module of the trained networks, there are indeed neurons whose activity is modulated by the task rule in a sustained fashion (example neurons in Supplementary Figure 1 and Figure 2a, top). In contrast, there are also neurons that show transient activity only after negative feedback. Furthermore, this activity is also rule-dependent. In other words, their activity is conjunctively modulated by negative feedback and the task rule (example neurons in Supplementary Figure 1, red traces and Figure 2a, bottom). We

termed these two classes of neurons "rule neurons" and "conjunctive error x rule neurons" respectively.

We identified all the rule neurons and conjunctive error x rule neurons in the PFC module using a single neuron selectivity measure (see Methods for details). The two classes of neurons are clearly separable on the two-dimensional plane in Figure 2c, where the x axis is the input weight for negative feedback, and the y axis is the rule modulation, which is the difference in the mean activity between the two rules (for trials following a correct trial). As shown in Figure 2c, rule neurons receive negligible input about negative feedback, and many of them have activity modulated by rule. On the other hand, conjunctive error x rule neurons receive a substantial amount of input about negative feedback, yet their activity is minimally modulated by rule on trials following a correct trial (Figure 2b). This pattern was preserved when aggregating across trained networks (Figure 2c and Supplementary Figure 3). Interestingly, neurons with similar tuning profiles have been reported in the PFC and posterior parietal cortex of macaque monkeys performing the same WCST analog [8, 9].

Compared to excitatory neurons, a much smaller fraction of inhibitory neurons in the PFC were classified as conjunctive error x rule neurons. On average, 22.9% excitatory neurons were conjunctive error x rule neurons in each model, compared with 11.5% PV neurons and 5.2% SST neurons. Therefore, we focus only on the excitatory conjunctive error x rule neurons in the analysis below.

Maintaining and switching rule states via structured connectivity patterns between subpopulations of neurons within the PFC module

Given the existence of rule neurons and conjunctive error x rule neurons, what is the connectivity between them that enables the PFC module to stay in the same rule attractor state when receiving correct feedback, and switch to the other rule attractor state when receiving negative feedback?

To this end, we examined the connectivity between different subpopulations of neurons in the PFC module explicitly, by computing the mean connection strength between each pair of subpopulations. This analysis reveals that the excitatory rule neurons and PV rule neurons form a classic winner-take-all network architecture [35] with selective inhibitory populations [36, 37], where excitatory neurons preferring the same rule are more strongly connected, and they also more strongly project to PV neurons preferring the same rule (Figure 3a). On the other hand, PV neurons project more strongly to both excitatory neurons and other PV neurons with the opposite rule preference (Figure 3a). This winner-take-all network motif together with the excitatory drive from the sensorimotor module (Supplementary Figure 2) is able to sustain one of the two attractor states.

Figure 2: **Emergence of two subpopulations of excitatory neurons in the PFC module after training.**

a. Two example rule neurons (top) and conjunctive error x rule neurons (bottom). The solid traces represent the mean activity across trials that follows a correct trial, when those trials belong to color rule (blue) or shape rule (green) blocks. The dashed traces represent the mean activity after error trials, when those trials belong to color rule (blue) or shape rule (green) blocks. We use rule 1 and color rule, as well as rule 2 and shape rule interchangeably hereafter.

b. Rule neurons and conjunctive neurons are separable. The x axis represents the input weight for negative feedback, and the y axis is the difference between the mean activity over color rule trials and shape rule trials (for trials following a correct trial). As shown, the rule neurons (blue points) receive little input about negative feedback, but their activity is modulated by rule; The conjunctive error x rule neurons (red points) receive substantial input about negative feedback, but their activity is not modulated by rule (during trials following a correct trial).

c. The trend in **b** is preserved across a collection of trained networks. Here the results are shown for networks with subtractive dendritic nonlinearity. Networks with divisive dendritic nonlinearity show a similar pattern (Supplementary Figure 3).

Next, how are the rule neurons connected with the conjunctive error x rule neurons such that the sub-network formed by rule neurons can switch from one attractor to the other in the presence of the negative feedback input? Using the same method, we found that the connectivity between the rule neurons and the conjunctive error x rule neurons exhibited an interesting structure: the excitatory rule neurons more strongly target the conjunctive error x rule neurons that prefer the opposite rule; the PV rule neurons more strongly target conjunctive error x rule neurons that prefer the same rule (Figure 3b, top two panels). On the other hand, the conjunctive error x rule neurons more strongly target the excitatory and PV rule neurons that prefer the same rule (Figure 3b, bottom two panels).

This connectivity structure gives rise to a simple circuit diagram of the PFC module (Figure 3c), which leads to an intuitive explanation of the circuit mechanism underlying the switching of rule attractor state. For example, suppose the network is in the attractor state corresponding to color rule, and has just received a negative feedback and is about to switch to the attractor corresponding to the shape rule (Figure 3e, left). As shown in Figure 2b-c, the input current that represents the negative feedback mainly targets the conjunctive error x rule neurons. In addition, since the network is in the color rule state, the excitatory and PV neurons that prefer the color rule are more active than those that prefer the shape rule. According to Figure 3b (top two panels), the excitatory neurons that prefer the color rule strongly excite the error x shape rule neurons, and the PV neurons that prefer the color rule strongly inhibit the error x color rule neurons. Therefore, the error x shape rule neurons receive stronger total input than the error x color rule neurons, and will be more active (Figure 3e, middle). Their activation will in turn excite the excitatory neurons and PV neurons that prefer the shape rule (Figure 3b, bottom two panels). Finally, due to the winner-take-all connectivity between the rule populations (Figure 3a), the excitatory and PV neurons that prefer the color rule will be suppressed, and the network will transition to the attractor state for the shape rule (Figure 3e, right).

It is worth noting that the same mechanism can also trigger a transition in the opposite direction (from shape rule to color rule) in the presence of the same negative feedback signal. This is enabled by the biased connections between the rule and conjunctive error x rule populations.

Is the simplified circuit diagram (Figure 3c) consistent across trained networks, or different trained networks use different solutions? To examine this question, we computed a "connectivity bias" measure between each pair of populations for each trained network. This measure is greater than zero if the connectivity structure between a pair of populations is closer to the one in the simplified circuit diagram in Figure 3c than to the opposite (see Methods for details). Across trained networks, we found that the connectivity biases were mostly greater than zero (Figure 3d), indicating that the same circuit motif for rule

Figure 3: (Caption next page.)

maintenance and switching underlies the PFC module across different trained networks.

A similar circuit architecture exists between the excitatory neurons and the SST neurons in the PFC module (Supplementary Figure 5), where SST neurons receive stronger excitatory input from the conjunctive error x rule neurons that prefer the same rule, and also more strongly inhibit the error x rule neurons that prefer the same rule. In addition, they form a winner-take-all connectivity with the rule excitatory neurons by receiving stronger projections from the rule neurons that prefer the same rule and projecting back more strongly to the rule neurons that prefer the opposite rule. Therefore, they contribute to rule maintenance and switching in a similar way as the PV neurons.

Figure 3: (Previous page.) **An emergent circuit wiring diagram in the PFC module enables un-cued switching between rule attractor states.**
a. The connectivity matrix between different populations of rule neurons, for an example model. Text indicates the mean connection strength between two populations. The excitatory rule neurons project more strongly to, and receive more input from, neurons with the same preferred rule. The PV rule neurons project more strongly to and receive more input from neurons with the opposite rule preference. As a result, rule neurons form a classic winner-take-all connectivity with selective inhibitory populations that maintain the two rule attractor state.
b. The connectivity between rule neurons and conjunctive error x rule neurons, for an example model. Top left: excitatory rule neurons project more strongly to the conjunctive error x rule neurons that prefer the opposite rule; Top right: PV rule neurons project more strongly to conjunctive error x rule neurons that prefer the same rule; Bottom left: conjunctive error x rule neurons project more strongly to the excitatory rule neurons that prefer the same rule; Bottom right: conjunctive error x rule neurons project more strongly to the PV rule neurons that prefer the same rule.
c. The simplified circuit diagram between rule neurons and conjunctive neurons based on the result of **b**. The weaker connections are ignored. Rule 1 represents the color rule and rule 2 represents the shape rule
d. A connectivity bias was computed to describe the extent to which the connectivity pattern between each pair of subpopulations conform to the simplified diagram in **c**. A value greater than 0 indicates the connectivity structure is more similar to that in **c** than to the opposite. The connectivity biases across all trained models are mostly above 0, both for the connection among rule neurons (top) and the connection between rule neurons and conjunctive error x rule neurons (bottom). Here the results are shown for networks with subtractive dendritic nonlinearity. Networks with divisive dendritic nonlinearity show similar result (Supplementary Figure 4).
e. A schematic showing how the simplified circuit can switch from the rule 1 attractor state to the rule 2 attractor state after receiving the input about negative feedback. The conjunctive error x rule 2 neurons receive excitation from the currently-active rule 1 excitatory neurons (red arrow, left panel), and the conjunctive error x rule 1 neurons receive inhibition from the currently-active rule 1 PV neurons (blue arrow, left panel). This makes conjunctive error x rule 2 neurons more active than the conjunctive error x rule 1 neurons, even though the negative feedback input targets both error x rule 1 and error x rule 2 populations (left panel). The conjunctive error x rule 2 neurons then excite the rule 2 excitatory and PV neurons (red arrows, middle), which suppress the rule 1 excitatory and PV neurons due to the winner-take-all connectivity (blue arrows, middle) and eventually become more active (right).

Top-down propagation of the rule information through structured long-range connections

Given that the PFC module can successfully maintain and update the rule representation, how does it use the rule representation to reconfigure the sensorimotor mapping?

Figure 4: (Caption next page.)

First, we found that neurons in the sensorimotor module were tuned to rule (Supplementary Figure 6a), since they receive top-down input from the rule neurons in the PFC module. The PFC module exerts top-down control through three pathways: the monosynaptic pathway from the excitatory neurons in the PFC module to the excitatory neurons in the sensori-motor module, the tri-synaptic pathway that goes through the VIP and SST neurons in the sensorimotor module, and the di-synaptic pathway mediated by the PV neurons in the sensorimotor module (Figure 1b). We found that there are structured connectivity patterns along all three pathways. Along the monosynaptic pathway, excitatory neurons in the PFC module preferentially send long-range projections to the excitatory neurons in the sensori-motor module that prefer the same rule (Figure 4a). Along the tri-synaptic pathway, PFC excitatory neurons also send long-range projections to the SST and VIP interneurons in the sensorimotor module that prefer the same rule (Figure 4b-c). The SST neurons in turn send stronger inhibitory connections to the dendrite of the local excitatory neurons that prefer the opposite rule (Figure 4d). Along the di-synaptic pathway, the PV neurons are also more strongly targeted by PFC excitatory neurons that prefer the same rule (Figure 4e), and they inhibit local excitatory neurons that prefer the opposite rule (Figure 4f). These trends are preserved across trained networks (Supplementary Figure 7). Therefore, rule information is communicated to the sensorimotor module synergistically via the mono-synaptic excitatory pathway, the tri-synaptic pathway that involves the SST and VIP neurons, as well as the di-synaptic pathway that involves the PV neurons.

Figure 4: (Previous page.) **Structured top-down connections enable the propagation of the rule information.**

a. Each line represents the mean connection strength onto one excitatory neuron in the sensorimotor module, from the PFC excitatory neurons that prefer the same rule and the different rule. Bars represent mean across neurons. PFC excitatory neurons project more strongly to excitatory neurons in the sensorimotor module that prefer the same rule (Student's t test, $p < .001$).

b. Each line represents the mean connection strength onto one VIP neuron in the sensorimotor module, from the PFC excitatory neurons that prefer the same rule and the different rule. Bars represent mean across neurons. PFC excitatory neurons project more strongly to VIP neurons in the sensorimotor module that prefer the same rule (Student's t test, $p = .002$).

c. Each line represents the mean connection strength onto one SST neuron in the sensorimotor module, from the PFC excitatory neurons that prefer the same rule and the different rule. Bars represent mean across neurons. PFC excitatory neurons project more strongly to SST neurons in the sensorimotor module that prefer the same rule (Student's t test, $p < .001$).

d. Each line represents the mean connection strength onto one excitatory neuron of the sensorimotor module, from the local SST neurons that prefer the same rule and the different rule. Bars represent mean across neurons. Local SST neurons project more strongly to excitatory neurons in the sensorimotor module that prefer the opposite rule (Student's t test, $p < .001$).

e. Each line represents the mean connection strength onto one PV neuron in the sensorimotor module, from the PFC excitatory neurons that prefer the same rule and the different rule. Bars represent mean across neurons. PFC excitatory neurons project more strongly to PV neurons in the sensorimotor module that prefer the same rule (Student's t test, $p = 0.004$).

f. Each line represents the mean connection strength onto one excitatory neuron of the sensorimotor module, from the local PV neurons that prefer the same rule and the different rule. Bars represent mean across neurons. PV neurons in the sensorimotor module project more strongly to local excitatory neurons that prefer the opposite rule (Student's t test, $p < .001$).

g. The structure of the top-down connections as indicated by the results in **a-f**. The weaker connections are not shown.

Results in **a-f** are shown for an example network with subtractive dendritic nonlinearity. Networks with divisive and subtractive dendritic nonlinearity show similar patterns (Supplementary Figure 7).

Structured input and output connections of the sensorimotor module enable rule-dependent action selection

How does the sensorimotor module implement the sensorimotor transformation (from the cards to the response to one of the three spatial locations) given the top-down rule information from the PFC module? We sought to identify the structures in the input,

recurrent and output connections of the sensorimotor module that give rise to this function.

We start by observing that excitatory neurons in the sensorimotor module show a continuum of encoding strengths for task rule, response location and card features, and many neurons show conjunctive selectivity for these variables (Figure 5b, Supplementary Figure 6b). Therefore, we assigned each excitatory neuron in the sensorimotor module a preferred rule R, a preferred response location L and a preferred shared feature between the reference card and the test card at L, which we call F. For example, neurons with $R = color\ rule$, $L = 1$ and $F = blue$ would have the highest activity during color rule trials when the correct response is to choose the test card at location 1, and when that test card shares the blue color with the reference card (it belongs to the population in the sensorimotor module with the filled green color in Figure 5a). Intuitively, for this group of neurons to show such selectivity, they should receive strong input from the input neurons that encode the $F = blue$ feature of the test card at location $L = 1$ and the same feature for the reference card. This would enable them to detect when the test card at $L = 1$ and the reference card both have the feature $F = blue$.

In general, for neurons that prefer rule R, response location L and shared feature F, we can define their "preferred features" as the feature F of the reference card and the same feature F for the test card at location L. Across all neurons, we found that the weights from the input neurons encoding these preferred features were significantly stronger than those encoding other features (Figure 5c). In addition, there is also an intuitive structure in the output connections, where excitatory neurons in the sensorimotor module that prefer a given response location L send stronger output connections to the output neuron that prefers the same response location (Figure 5d). These structures were found to be consistent across trained networks (Supplementary Figure 8).

These structures in the input and output connections give rise to an intuitive explanation of how the sensorimotor module can perform rule-dependent action selection required for the WCST. Here we illustrate this mechanism with an example trial (Figure 5a), where the current rule is color, the reference card is a blue circle, and the test cards at locations 1 2 and 3 are blue triangle, red circle and red triangle, respectively. According to the rule of WCST, the correct response should be location 1, since the test card at that location matches the reference card in color. This choice can be generated as follows: the excitatory population in the sensorimotor module that prefers $R = color\ rule$, $L = 1$ and $F = blue$ will be most strongly activated since they not only receive strong top-down input from the PFC module, but also the strongest feedforward input. Therefore, they are the most strongly activated population (Figure 5a). And since they prefer response location 1, they will excite the output neuron that prefers response location $L = 1$, which is the correct choice.

Figure 5: (Caption next page.)

Figure 5: (Previous page.)   **Structures in the input and output weights of the sensorimotor module enable rule-dependent action selection.**
a. Excitatory neurons in the sensorimotor module were classified according to their preferred rule R, response location L and shared feature F. For example, neurons with $R = color\ rule$, $L = 1$ and $F = blue$ have the highest activity during color rule trials when the network chooses the test card at $L = 1$, and when that card shares the $F = blue$ feature with the reference card. For a neuron with a given R, L and F, its "preferred features" are defined as the feature F of the reference card and same feature of the test card at location L. For example, the preferred features for neurons with $R = color\ rule$, $L = 1$ and $F = blue$ are the blue feature of the reference card and the test card at $L = 1$.
b. The joint distribution of the selectivity for rule (R), response location (L) and shared feature (F) across all neurons in the sensorimotor module. Result is aggregated across all trained networks.
c. Excitatory neurons in the sensorimotor module receive stronger connections from the input neurons that encode their preferred features (as defined in **a**). Student's t-test, $p < .001$
d. Excitatory neurons in the sensorimotor module send stronger connections to the output neuron that represents their preferred response location. Student's t-test, $p < .001$.
Panel **a** shows an example trial illustrating how the sensorimotor module can generate the correct response. During this trial, the reference card is a blue circle, and the test cards at location 1, 2, 3 are blue triangle, red circle and red triangle, respectively. The current rule is color. Therefore the correct response location should be $L = 1$. The network can generate that response because (a) the PFC population that encode the color rule are active, which send strong top-down excitation to the $R = color\ rule$ population in the sensorimotor module; (b) the input neurons that encode the $F = blue$ feature of the reference card and the test card at $L = 1$ are both active, which provide strong feedforward input to the excitatory population in the sensorimotor module with $R = color\ rule$, $L = 1$ and $F = blue$. Therefore, this population will be most strongly activated. Since they send strong long-range excitations to the output neuron that represents $L = 1$, the correct output neuron will be activated.

Recurrent connectivity and dynamics within the sensorimotor module

Given that different populations of neurons in the sensorimotor module receive differential inputs about the external sensory stimuli and rule via the structured input and top-down connections, how are they recurrently connected to produce dynamics that lead to a categorical choice? To answer this, we first visualized the population neural dynamics in the sensorimotor module by using principal component analysis (Figure 6a-b). As shown in Figure 6a, neural trajectories during the inter-trial interval are clustered according to the task rule. During the response period, the neural trajectories are separable according to the response locations, albeit only in higher-order principal components (Figure 6b). In addition, the subspaces spanned by neural trajectories of different rules and response locations are more orthogonal to each other compared to randomly shuffled data (Figure 6c-d, Methods).

What connectivity structure gives rise to this signature in the population dynamics? To answer this, we examined the pattern of connection weights between excitatory and PV neurons that prefer different rules (R), response locations (L), and shared features (F) by computing the connectivity biases between populations of neurons that are selective to different rules (Figure 6e), response locations (Figure 6f) and shared features (Figure 6g). A greater than zero connectivity bias means populations that prefer different rules (or response locations or shared features) form a winner-take-all circuit structure analogous to the one observed between rule-selective populations in the PFC module (c.f. top panel of Figure 3d, details about how the connectivity biases were computed is described in Methods). We observed that many of the connectivity biases were significantly above zero (Figure 6e-g), especially for the ones that correspond to the inhibitory connections originating from the PV neurons. This indicates that populations of neurons in the sensorimotor module that are selective to different rules, response locations and shared features overall inhibit each other. This mutual inhibition circuit structure magnifies the difference in the amount of long-range inputs that different populations receive (Figure 5) and lead to a categorical choice.

SST neurons are essential to dendritic top-down gating

The previous sections elucidate the key connectivity structures that enable the network to perform the WCST. In this final section we are going to take advantage of the biological realism of the trained RNN and examine the function of SST neurons in this task.

It has been observed that different dendritic branches of the same neuron can be tuned to different task variables [38, 39, 40, 41]. This property may enable individual dendritic branches to control the flow of information into the local network [17, 20]. Given these

Figure 6: **Recurrent dynamics and connectivity within the sensorimotor module.**
a. Neural trajectories during the intertrial interval (ITI) for different task rules, visualized in the space spanned by the first three principal components. Black circles represent the start of the ITI.

b. Neural trajectories during the response period for different choices, visualized in the space spanned by higher order principal components. Black circles represent the start of the response period.

c. The principal angle between the subspaces spanned by neural trajectories during different task rules (gray distribution represents the principal angle obtained through shuffled data, see Methods). Each data point represents one trained network.

d. The principal angle between the subspaces spanned by neural trajectories during different responses (gray distribution represents the principal angle obtained through shuffled data, see Methods). Each data point represents one trained network.

e. The connectivity biases between different rule-selective populations across models.

f. The same as **e** but for different response location-selective populations.

g. The same as **e** but for different shared feature-selective populations.

h. The results in **e** - **g** show that neural populations selective for different rules, response locations and shared features mutually inhibit each other.

Data in **c-g** are shown for networks with subtractive dendritic nonlinearity. Networks with divisive dendritic nonlinearity show similar result (Supplementary Figure 9).

previous findings, we examined the coding of the top-down rule information at the level of individual dendritic branches. Since each excitatory neuron in our networks is modeled with two dendritic compartments, we examined the encoding of rule information by different dendritic branches of the same excitatory neuron in the sensorimotor module.

One strategy of gating is for different dendritic branches of the same neuron to prefer the same rule, in which case these neurons form distinct populations that are preferentially recruited under different task rules (population-level gating, Figure 7a, right). An alternative strategy is for different dendritic branches of the same neuron to prefer different rules, which would enable these neurons to be involved in both task rules (dendritic branch-specific gating, Figure 7a, left).

In light of this, we examined for our trained networks to what extent they adopt these strategies. We found that the rule selectivity between different dendritic branches of the same neuron were highly correlated (Figure 7b). This indicates that the trained networks are mostly using the population-level gating strategy, where different dendritic branches of the same neuron encode the same rule.

What factors might determine the extent to which the trained networks adopt these two strategies? Previous modeling work suggests that sparse connectivity from SST neurons to the dendrites of the excitatory neurons increases the degree of dendritic branch-specific gating, in the case where the connectivity is random [20]. To see if the same effect is present in our task-optimized network with structured connectivity, we re-trained networks with different levels of sparsity from 0 to 0.8 and studied its effect on the dendritic branch specificity of rule coding (Methods). We found that the degree of dendritic branch-specific encoding of the task rule increased with sparsity (see Figure 7c, d for subtractive dendritic nonlinearity; Supplementary Figure 10a for divisive dendritic nonlinearity). Intuitively, when the connection is sparse, a smaller number of SST neurons target each dendritic branch, making it more likely that the branch receives an uneven number of inputs from SST neurons selective for different rules. Taken together, we observed that the trained networks adopted a mixture of population-level and dendritic-level gating strategies for top-down control, and the balance between the two strategies depends on the sparsity of the connections from the SST neurons to the dendrites of excitatory neurons.

Indeed, SST neurons play a causal role in relaying the top-down rule information into the sensorimotor network and reconfiguring its dynamics according to the task rule. We simulated optogenetic inhibition by silencing the SST neurons in the sensorimotor module, which significantly impaired task performance (Figure 7e, see Methods section for details of the protocol). In addition, the principal angle between the subspaces for different rules (Figure 6c) significantly decreased after SST neurons in the sensorimotor module were silenced (Figure 7f). Silencing of the SST neurons in the sensorimotor module also significantly

diminished nonlinear mixed-selective coding of rule and stimulus among the excitatory neurons in the sensorimotor module (Figure 7g, Supplementary Figure 11, Methods), which has been proposed to be important for rule-based sensorimotor associations [42, 43, 44]. Taken together, these results highlight the role that SST neurons in the sensorimotor module play during top-down control. This analysis also shows that by combining artificial neural network with knowledge from neurobiology, it is possible to probe the functions of fine-scale biological components in cognitive behaviors.

Discussion

In this paper, we analyzed recurrent neural networks trained to perform a classic task involving un-cued task switching - the Wisconsin Card Sorting Test. The networks consist of a "PFC" module trained to represent the rule and interacts with a "sensorimotor" module that instantiates different sensorimotor mappings depending on the rule. In order to study the functions of dendritic computation and different neuronal types, each module is endowed with excitatory neurons with two dendritic branches as well as three major types of inhibitory neurons - PV, SST and VIP. After training, we dissected the trained networks to elucidate a number of intra-areal and inter-areal neural circuit mechanisms underlying WCST, as summarized in Figure 8.

Mapping between model components and brain regions

Different components of the trained network can be mapped to different brain regions (Figure 8). While single neurons in the dorsal-lateral PFC (DLPFC) are shown to encode the task rule [45], neurons in the anterior cingulate cortex (ACC) are thought to be important for performance monitoring [46], and have been shown to receive more input about the feedback [47, 48, 49, 50]. Therefore, the rule neurons and conjunctive error x rule neurons in the model correspond to the putative functions of the neurons in DLPFC and ACC. The input to the PFC module about negative feedback may come from subcortical areas such as the amygdala [51] or from the dopamine neurons in the substantia nigra pars compacta (SNc) and ventral tegmental area (VTA) [52, 53]. The sensorimotor module may correspond to parietal cortex or basal ganglia which have been shown to be involved in sensorimotor transformations [54, 55]. The neurons in the input layer that encode the color and shape of the card stimuli exist in higher visual areas such as the inferotemporal cortex [56, 57, 58]. The neurons in the output layer that encode different response locations could correspond to movement location-specific neurons in the motor cortex [59].

Figure 7: **Examining the role of SST neurons in the sensorimotor module in top-down gating.**

a. Two scenarios for top-down gating. Blue and green color represent dendritic branches that prefer one of the two rules. Different dendritic branches of the same neuron could have similar (right) or different (left) rule selectivity.

b. The rule selectivity of one dendritic branch against the other, aggregated across all models where the connections from the SST neurons to the excitatory neurons are all-to-all. The rule selectivity for different dendritic branches of the same neuron are highly correlated.

c. The rule selectivity of one dendritic branch against the other, aggregated across all models where the 80% of the connections from the SST neurons to the excitatory neurons are frozen at 0 throughout training. Note the the rule selectivity for different dendritic branches of the same neuron are less correlated than in **b**.

d. The degree of dendritic branch-specific encoding of the task rule is quantified as the difference in the rule selectivity between the two dendritic branches of the same excitatory neuron in the sensorimotor module. Across all dendritic branches, this quantity increases with the sparsity of the SST → dendrite connectivity.

e. Task performance drops significantly after silencing SST neurons in the sensorimotor module. Each line represents a trained network.

f. The principal angle between rule subspaces (c.f. Figure 6c) drops significantly after silencing SST neurons in the sensorimotor module. Each line represents a trained network.

g. The strength of conjunctive coding of rule and stimulus (as measured by the R^2 value in a linear model with conjunctive terms, see Methods) decreases after silencing SST neurons in the sensorimotor module (Student's t-test, $p < .001$). Each line represents one neuron. Results are aggregated across networks.

Results in **b-g** are for networks with subtractive dendritic nonlinearity. Networks with divisive dendritic nonlinearity show similar result (Supplementary Figure 10).

Figure 8: **A summary of the main results.**
Different components of the model can be mapped to different brain regions; The conjunctive error x rule neurons may reside in the anterior cingulate cortex; The rule neurons may be found in the dorsal-lateral PFC; The input to the PFC module about negative feedback may come from subcortical areas such as the amygdala or the midbrain dopamine neurons; The sensorimotor module may correspond to parietal cortex or basal ganglia which have been shown to be involved in sensorimotor transformations; Neurons in the input layer that encode the color and shape of the card stimuli exist in higher visual areas such as the inferotemporal cortex; Neurons in the output layer that encode different response locations could correspond to neurons in the motor cortex.

Attractor states supported by inter-areal connections.

We observed two rule attractor states in the dynamical landscape of the networks supported by the interaction between the two modules (Supplementary Figure 2). This is contrary to the traditional notion that local interactions within the frontal regions are sufficient for the maintenance of the persistent activity [27, 28, 29, 30]. Instead it suggests the possibility that the interactions between distributed brain regions underlie temporally-extended cognitive functions [31, 33, 60].

Circuit mechanism in the frontal-parietal network for rule maintenance and update

Two distinct types of responses among the excitatory neurons emerge in the PFC module as a result of training: neurons that only encode the rule, and neurons that conjunctively encode negative feedback and rule. Neurons that show conjunctive selectivity for rule and negative feedback have been reported in monkey prefrontal and parietal cortices while they perform the same WCST task [8, 9]. Theoretical work suggests that these mixed-selective neurons are essential if the network needs to switch between *different* rule attractor states after receiving the *same* input that signals negative feedback [61].

We further revealed the connectivity pattern between different populations of excitatory and PV neurons in the PFC module in order for the network to switch between rule attractor states (Figure 3c). In addition, this connectivity pattern is consistent across dozens of trained networks with different initializations and dendritic nonlinearities (Figure 3d and Supplementary Figure 4). This circuit mechanism bears resemblance to a previous circuit model of WCST [62]. In that model, the switching between different rule states is achieved by synaptic desensitization caused by the convergence of two signals - one that signals the recent activation of the synapse, and another that signals the negative feedback. However, that model does not predict the existence of neurons with conjunctive coding of negative feedback and rule, which has been observed experimentally [8, 9].

The simplified circuit for the PFC module in Figure 3c can be applied not only to rule switching, but to the switching between other behavioral states as well. For example, it resembles the head-direction circuit in fruit fly [63], where the offset in the connections between the neurons coding for head direction and those coding for the conjunction of angular velocity and head direction enables the update of the head-direction attractor state by the angular velocity input. In addition, this circuit structure may underlie the transition from staying to switching during patch foraging behavior. Indeed, in a laboratory task mimicking natural foraging for monkeys, it was found that neurons in the anterior cingulate cortex increase their firing rates to a threshold before animals switch to another food resource [64].

Connecting subspace to circuits

Methods that describe the representation and dynamics on the neuronal population level have gained increasing popularity and generated novel insights that cannot be discovered using single neuron analysis (e.g. [59, 65]). In the meantime, it would be valuable to connect population-level phenomena to their underlying circuit basis [66]. In our model, we found that silencing of the SST neurons has a specific effect on the population-level representation, namely, it decreases the angle between rule subspaces (Figure 7f). We also found that silencing the other types of inhibitory neurons has different effects (data not shown).

Silencing the PV neurons to an instability of the network dynamics, whereas silencing the VIP neurons caused an insignificant decrease of the network performance. The lack of effect after silencing the VIP neurons is due to the fact that the VIP neurons were largely inhibited by the SST neurons in the trained model. Future work could study the function of VIP neurons under different connectivity constraints between SST and VIP neurons.

Evidence accumulation across trials during rule switching

In this study, network models were built using supervised learning where the supervision signal was the correct response. This supervision signal optimized the network for task performance, which explains why they only made a single error after each rule switch. In the laboratory, the performance of monkeys can reach this optimal level (e.g. Figure 1D in Ref.[9], monkey W. See Figure 2 of Ref.[67] for a similar task), but most of the time the performance varies greatly and is not optimal [21, 8, 9, 6]. This can be due to a number of reasons, including random exploration [68], poor sensitivity to negative feedback [68], integration of reward history across multiple trials [69, 70, 10, 71], the gradual update of the value of the counterfactual rule [72] or the cost of cognitive control [73]. Future work can use the subjects' behavior (rather than the correct response) as the supervision signal, in order to study how negative feedback is combined with other internal signals across multiple trials to infer the relevant task rule.

WCST with more than two rules

The WCST task that our networks were trained on involves only two rules, whereas the WCST used clinically usually has three rules. In this case, the correct rule cannot be determined after a single error. In order to perform rationally in this task, neuronal processes with time constants on the order of several trials are essential. Future work could incorporate biological processes with longer time constants into the network model in order to investigate their functions in WCST.

In conclusion, our approach of incorporating neurobiological knowledge into training RNNs can provide a fruitful way to build circuit models that are functional, high-dimensional, and reflect the heterogeneity of biological neural networks. In addition, dissecting these networks can make useful cross-level predictions that connect biological ingredients with circuit mechanisms and cognitive functions.

Acknowledgements: This work was supported by James Simons Foundation Grant 543057SPI, the National Institutes of Health grant R01MH062349, and the ONR grant N00014-23-1-2040. YL thanks (alphabetically) Aldo Battista, Mark Buckley, Sage Chen,

Data availability statement: All computer code used to generate the model and results will be uploaded to Github upon acceptance of the manuscript.

Methods

Model setup

The RNN consists of two bidirectionally-connected modules, the PFC module and the sensorimotor module. Each module consists of 70 excitatory neurons and 30 inhibitory neurons. Each excitatory neuron has 2 dendritic compartments. The inhibitory neurons are evenly divided into three types: PV, SST and VIP. Different types of neurons have different connectivity, inspired by experimental findings [74]: PV neurons target the somatic compartment of excitatory neurons and other PV neurons, SST neurons target the dendritic compartment of excitatory neurons as well as PV and VIP neurons, and VIP neurons target SST neurons. Excitatory neurons target other excitatory neurons, PV and SST neurons. The connection strength between all other types of neurons were fixed at zero throughout training.

Only excitatory neurons send long-range projections to other modules. The long-range projections from the sensorimotor module to the PFC module target the dendritic compartment of the excitatory neurons and the PV neurons. This is inspired by the experimental evidence that PV neurons mediate feedforward inhibition [12]. The long-range top-down projections from the PFC to the sensorimotor module target the dendritic compartments of the excitatory neurons and all three types of inhibitory neurons. Finally, external inputs to both modules target the dendritic compartment of excitatory neurons and PV neurons.

The dynamics of the somata of the excitatory neurons in the RNN are described by

$$\tau \frac{dh_{\text{soma, exc}}}{dt} = -h_{\text{soma, exc}} + f_{\text{soma, exc}}(W_{\text{rec,eff}}h_{\text{soma, exc}} + \sum_{\text{dendrites}} h_{\text{dendrite}}), \qquad (1)$$

where $\tau = 100$ ms, $dt = 10$ ms. h_{dendrite} is the activity of the dendritic compartment. $f_{\text{soma, exc}}$ is the rectified linear activation function:

$$f_{\text{soma, exc}} = \begin{cases} x & x > 0 \\ 0 & \text{otherwise} \end{cases} \qquad (2)$$

528   The effective recurrent connectivity matrix $W_{\text{rec, eff}}$ is given by

$$529 \qquad W_{\text{rec, eff}} = W_{\text{rec}} * |M|, \tag{3}$$

530 where M is the mask matrix consisting of 1, 0 and -1 according to the cell-type-specific
531 connectivity described above. $*$ denotes element-wise product and $|.|$ denotes the absolute
532 value.

533   The dendritic activity is a nonlinear function of the excitatory and inhibitory inputs.

$$534 \qquad h_{\text{dendrite}} = f_{\text{dendrite}}(I_{\text{exc}}, I_{\text{inh}}). \tag{4}$$

I_{exc} is the total excitatory input to the dendrite. It consists of long-range inputs from the
input neurons (neurons that encode the feedback for the PFC module and neurons that
encode the stimulus for the sensorimotor module) as well as the long-range input from the
excitatory neurons in the other module. $I_{\text{exc}} = I_{\text{in}} + I_{\text{cross-module}}$. I_{inh} is the inhibitory
input to the dendrite from the local SST neurons. $I_{\text{inh}} = I_{\text{SST}\rightarrow\text{exc}}$. The functional form of
f_{dendrite} is described in the next section.

The inhibitory neurons are modeled as standard point neurons

$$542 \qquad \tau \frac{dh_{\text{soma, inh}}}{dt} = -h_{\text{soma, inh}} + f_{\text{soma, inh}}(W_{\text{rec,eff}}h_{\text{soma, inh}} + W_{\text{in}\rightarrow\text{PV}}u_{\text{in}}), \tag{5}$$

where $W_{\text{in}\rightarrow\text{PV}}$ is the weight matrix from the input neurons to the PV neurons and u_{in} is
the activity of the input neurons. The activation function $f_{\text{soma, inh}}$ is the same as the one
used for excitatory neurons $f_{\text{soma, exc}}$ (Equation 2).

Only the somata of excitatory neurons send output connections. The output y (rule
for the PFC module and choice for the sensorimotor module) is related to their activity
$h_{\text{exc, soma}}$ via simple linear transformation:

$$549 \qquad y = W_{\text{out}}h_{\text{exc, soma}}. \tag{6}$$

Variations in the model hyperparameters

Dendritic nonlinearities. We trained models with two types of dendritic nonlin-
earities f_{dendrite} - subtractive and divisive. They are inspired by in-vitro and computational
studies showing different types of inhibitory modulation on the dendritic activity depending
on the location of inhibition relative to excitation [23]. Both types of dendritic nonlineari-
ties are sigmoidal functions of the excitatory input. Under subtractive nonlinearity, as the
inhibitory input increases, the turning point of the sigmoid function moves to larger values,
consistent with the experimental observation when the inhibitory current is injected at the

same location or more distal than the excitation [23]. For the divisive nonlinearity, the turning point of the sigmoid is not affected by the level of inhibition, but the saturating level of the sigmoid function decreases with the level of inhibition, consistent with the experimental observation when the inhibitory current is injected close to the soma [23].

The equations of the different dendritic nonlinearities are given by:

$$f_{\text{subtractive}}(I_e, I_i) = \tanh\left(I_e - I_i\right)$$

$$f_{\text{divisive}}(I_e, I_i) = k_1(1 + \tanh(I_e - 1)) + k_2,$$

where $k_1 = \frac{1}{e^{I_i}}$ and $k_2 = -1 - \tanh(-1)$. The constant k_2 is introduced such that the value of the function is 0 when both excitatory and inhibitory inputs are 0.

Initializations. The connectivity matrices were initialized either using a normal distribution with mean 0 and standard deviation $\sqrt{\frac{2}{N}}$ (where N is the total number of recurrent units) or a uniform distribution between $-\sqrt{\frac{6}{N}}$ and $\sqrt{\frac{6}{N}}$.

Sparsity of the SST→dendrite connectivity in the sensorimotor module. To study how the degree of dendritic branch-specific rule encoding in the sensorimotor module is affected by the sparsity of the connections from SST neurons to the dendrite of excitatory neurons, we varied this sparsity by fixing a fraction of randomly chosen connections to be 0 throughout training. The sparsity levels used were 0, 0.2, 0.4, 0.6 and 0.8.

Random seeds. For each combination of the hyperparameter configuration introduced above (except the sparsity), we trained models using 50 random seeds for Pytorch (other random seeds were fixed). For each sparsity level other than 0, we trained models using 10 random seeds for Pytorch.

Task

The networks were trained on an analog of the Wisconsin Card Sorting Test (WCST) used for monkeys [21, 8, 9]. Each trial starts with the presentation of a "reference card" for 500 ms, after which three "test cards" appear around the reference card for 500 ms. Each card contains an object with a specific color (blue or red) and shape (circle or triangle). Among the three test cards, one of them matches the color of the reference card, another one matches the shape of the reference card, and the third card matches neither feature of the reference card. Depending on the rule (color or shape), the location where the test card has the same color or shape feature as the reference card should be chosen. The choice should be made during the 500 ms when both the reference card and the test cards are

presented. At the end of this period, a feedback signal is presented for 100 ms, indicating
whether the choice is correct or incorrect. This is followed by a 1 second inter-trial interval.
The task rule switches after a random number of trials, without informing the network.
Therefore, the network inevitably makes an error for the first trial after the rule switch since
it has not yet received the information that the rule has switched. The network should then
adjust its behavior to the new rule by utilizing the feedback signal.

Representation of inputs and outputs

Each card is represented as a four-dimensional binary vector, where different entries
represent the presence of the two colors and shapes. The feedback input is a two-dimensional
one-hot vector, where the two entries represent positive and negative feedback. The target
output for the sensorimotor module is a three-dimensional one-hot vector, where each entry
represents one response location on the screen. This target is non-zero only during the 500
ms response period when both the reference card and the test cards are presented. The
target output for the PFC module is a two-dimensional one-hot vector, where each entry
represents one rule. This target is non-zero during the entire trial.

Training method

606   During training, the networks ran continuously across 20 consecutive trials with 3
607 random rule switches. Importantly, the network dynamics were not reset during the inter-
608 trial interval. The loss function was aggregated across the 20 trials.

$$L = \sum_{\text{trial}=1}^{20} \sum_{\text{t}} \left(||y_{\text{pfc, trial}}(t) - \hat{y}_{\text{rule, trial}}(t)||^2 + ||y_{\text{sensorimotor, trial}}(t) - \hat{y}_{\text{choice, trial}}(t)||^2 \right),$$

(7)

where $y_{\text{pfc, trial}}(t)$ and $y_{\text{sensorimotor, trial}}(t)$ are the activity of the readout neurons for the PFC
and sensorimotor module at time t in a given trial, respectively. $\hat{y}_{\text{rule, trial}}(t)$ is the target
output for the PFC module which represents the rule of the current trial. It is a binary
vector of dimension 2 where each entry represents one rule. The activation of the entry
that represents the correct rule is 1 throughout the entire trial. $\hat{y}_{\text{choice, trial}}(t)$ is the target
output for the sensorimotor module which represents the correct choice for the current trial.
It is a binary vector of dimension 3 where each entry represents one of the three response
locations. The activation of the entry that represents the correct choice is 1 during the
response period (500 ms when both the reference card and the tests card are shown).
The standard backpropagation through time algorithm [75] with the Adam optimizer
[76] was used to update all connection weights.
We also used curriculum learning to speed up training. Initially, the stimulus and

choice of the previous trial together with the feedback were provided to the PFC module. This way all the information needed to perform the current trial is contained within the input, and the networks do not need to memorize past trials. After the networks reached 85% performance, the input about the previous stimulus was removed. When the networks reached 85% performance again, the information about the previous choice information was removed. The networks were then trained until they reached 95% performance.

Single neuron selectivity metric

The selectivity index (SI) for rule is defined as

$$SI_{\text{rule}} = \frac{h(\text{color}) - h(\text{shape})}{|h(\text{color})| + |h(\text{shape})|}, \tag{8}$$

where $h(\text{color})$ and $h(\text{shape})$ represent the trial-averaged single neuron activity during color rule and shape rule, respectively. Neural activity was first averaged over the inter-trial interval before further being averaged across trials.

The error selectivity is defined similarly

$$SI_{\text{error}} = \frac{h(\text{after error}) - h(\text{after correct})}{|h(\text{after error})| + |h(\text{after correct})|}, \tag{9}$$

where $h(\text{after error})$ and $h(\text{after correct})$ are the mean single neuron activity after error and correct trials, respectively. Neural activity was first averaged across the feedback presentation and inter-trial interval periods before being averaged across trials.

The selectivity for response location is defined as

$$SI_{\text{response}} = \frac{h(L^*) - h(\bar{L})}{|h(L^*)| + |h(\bar{L})|}, \tag{10}$$

where L^* the preferred response location of the neuron, and $h(L^*)$ represents the mean activity across trials when the network chooses location L^*. $h(\bar{L})$ represents the mean activity across trials when the choice of the network is not location L^*. Therefore, this selectivity index ranges from 0 to 1. We included neural activity during the response period when computing this selectivity index.

Neurons that prefer color/shape rule were further divided according to their preferred shared feature. The selectivity for the shared feature is defined as

$$SI_{\text{shared feature}} = \frac{h(\text{blue}) - h(\text{red})}{|h(\text{blue})| + |h(\text{red})|}, \tag{11}$$

for neurons that prefer the color rule, and

$$SI_{\text{shared feature}} = \frac{h(\text{circle}) - h(\text{triangle})}{|h(\text{circle})| + |h(\text{triangle})|}, \tag{12}$$

for neurons that prefer the shape rule. Here $h(\text{blue})$, $h(\text{red})$, $h(\text{circle})$, $h(\text{triangle})$ represent the mean activity of a neuron across trials when the reference card is blue, red (when the current rule is color), circle or triangle (when the current rule is shape). We included neural activity during the response period when computing this selectivity index.

Classification criteria for different neuronal populations

Each neuron in the PFC module was classified as a "rule neuron" if the absolute value of its rule selectivity was greater than 0.5 and the absolute value of its error selectivity was smaller than 0.5. It was classified as an "error neuron" if the absolute value of its rule selectivity was smaller than 0.5 and its error selectivity was greater than 0.5. Error neurons with greater mean activity during the color rule trials that follow an error trial were defined as error x color rule neurons, and the other error neurons were defined as error x shape rule neurons.

Each neuron in the sensorimotor module was assigned with a preferred rule, response location and shared feature according to the condition during which it has the highest activity. There was no threshold for this classification.

Connectivity bias

The connectivity bias (CB) was defined as the difference in the average connection weight between different sub-population of neurons. A positive value indicates an agreement with the simplified circuit diagram (Figures 3c, Figure 6h). For example, the connectivity bias from the PFC PV neurons to the PFC excitatory neurons is given by

$$\begin{aligned}
CB(\text{PFC PV} \longrightarrow \text{PFC E}) = \ &\bar{W}(\text{PFC PV rule1} \longrightarrow \text{PFC E rule2}) \\
&+ \bar{W}(\text{PFC PV rule2} \longrightarrow \text{PFC E rule1}) \\
&- \bar{W}(\text{PFC PV rule1} \longrightarrow \text{PFC E rule1}) \\
&- \bar{W}(\text{PFC PV rule2} \longrightarrow \text{PFC E rule2}),
\end{aligned} \tag{13}$$

where for example $\bar{W}(\text{PFC PV rule1} \longrightarrow \text{PFC E rule2})$ represents the average (unsigned) connection strength from the PFC PV neurons that prefer rule 1 to PFC excitatory neurons that prefer rule 2. Here rule 1 refers to color rule and rule 2 refers to shape rule.

675   The other connectivity biases were defined analogously.

$$
\begin{aligned}
CB(\text{PFC E} \longrightarrow \text{PFC E}) = {} & \bar{W}(\text{PFC E rule1} \longrightarrow \text{PFC E rule1}) \\
& + \bar{W}(\text{PFC E rule2} \longrightarrow \text{PFC E rule2}) \\
& - \bar{W}(\text{PFC E rule1} \longrightarrow \text{PFC E rule2}) \\
& - \bar{W}(\text{PFC E rule2} \longrightarrow \text{PFC E rule1})
\end{aligned}
\tag{14}
$$

$$
\begin{aligned}
CB(\text{PFC E} \longrightarrow \text{PFC PV}) = {} & \bar{W}(\text{PFC E rule1} \longrightarrow \text{PFC PV rule1}) \\
& + \bar{W}(\text{PFC E rule2} \longrightarrow \text{PFC PV rule2}) \\
& - \bar{W}(\text{PFC E rule1} \longrightarrow \text{PFC PV rule2}) \\
& - \bar{W}(\text{PFC E rule2} \longrightarrow \text{PFC PV rule1})
\end{aligned}
\tag{15}
$$

$$
\begin{aligned}
CB(\text{PFC PV} \longrightarrow \text{PFC PV}) = {} & \bar{W}(\text{PFC PV rule1} \longrightarrow \text{PFC PV rule2}) \\
& + \bar{W}(\text{PFC PV rule2} \longrightarrow \text{PFC PV rule1}) \\
& - \bar{W}(\text{PFC PV rule1} \longrightarrow \text{PFC PV rule1}) \\
& - \bar{W}(\text{PFC PV rule2} \longrightarrow \text{PFC PV rule2})
\end{aligned}
\tag{16}
$$

$$
\begin{aligned}
CB(\text{PFC E} \longrightarrow \text{PFC error x rule}) = {} & \bar{W}(\text{PFC E rule1} \longrightarrow \text{PFC error x rule2}) \\
& + \bar{W}(\text{PFC E rule2} \longrightarrow \text{PFC error x rule1}) \\
& - \bar{W}(\text{PFC E rule1} \longrightarrow \text{PFC error x rule1}) \\
& - \bar{W}(\text{PFC E rule2} \longrightarrow \text{PFC error x rule2})
\end{aligned}
\tag{17}
$$

$$
\begin{aligned}
CB(\text{PFC PV} \longrightarrow \text{PFC error x rule}) = {} & \bar{W}(\text{PFC PV rule1} \longrightarrow \text{PFC error x rule1}) \\
& + \bar{W}(\text{PFC PV rule2} \longrightarrow \text{PFC error x rule2}) \\
& - \bar{W}(\text{PFC PV rule1} \longrightarrow \text{PFC error x rule2}) \\
& - \bar{W}(\text{PFC PV rule2} \longrightarrow \text{PFC error x rule1})
\end{aligned}
\tag{18}
$$

$$
\begin{aligned}
CB(\text{PFC error x rule} \longrightarrow \text{PFC E}) = {} & \bar{W}(\text{PFC error x rule1} \longrightarrow \text{PFC E rule1}) \\
& + \bar{W}(\text{PFC error x rule2} \longrightarrow \text{PFC E rule2}) \\
& - \bar{W}(\text{PFC error x rule1} \longrightarrow \text{PFC E rule2}) \\
& - \bar{W}(\text{PFC error x rule2} \longrightarrow \text{PFC E rule1})
\end{aligned}
\tag{19}
$$

$$
\begin{aligned}
CB(\text{PFC error x rule} \longrightarrow \text{PFC PV}) = &\ \bar{W}(\text{PFC error x rule1} \longrightarrow \text{PFC PV rule1}) \\
&+ \bar{W}(\text{PFC error x rule2} \longrightarrow \text{PFC PV rule2}) \\
&- \bar{W}(\text{PFC error x rule1} \longrightarrow \text{PFC PV rule2}) \\
&- \bar{W}(\text{PFC error x rule2} \longrightarrow \text{PFC PV rule1})
\end{aligned}
\tag{20}
$$

The connectivity biases between the different response location-selective populations in the sensorimotor module (SM) are defined as

$$
\begin{aligned}
CB(\text{SM response E} \longrightarrow \text{SM response E}) = &\ \bar{W}(\text{SM E response 1} \longrightarrow \text{SM E response 1}) \\
&+ \bar{W}(\text{SM E response 2} \longrightarrow \text{SM E response 2}) \\
&+ \bar{W}(\text{SM E response 3} \longrightarrow \text{SM E response 3}) \\
&- \bar{W}(\text{SM E response 1} \longrightarrow \text{SM E response 2 and 3}) \\
&- \bar{W}(\text{SM E response 2} \longrightarrow \text{SM E response 1 and 3}) \\
&- \bar{W}(\text{SM E response 3} \longrightarrow \text{SM E response 1 and 2}).
\end{aligned}
\tag{21}
$$

In the last equation, for example, $\bar{W}(\text{SM response 1} \longrightarrow \text{SM response 2 and 3})$ represents the mean connection strength from excitatory neurons in the sensorimotor module that prefer response location 1 to those that prefer response locations 2 and 3.

The other connectivity biases were defined similarly

$$
\begin{aligned}
CB(\text{SM response E} \longrightarrow \text{SM response PV}) = &\ \bar{W}(\text{SM E response 1} \longrightarrow \text{SM PV response 1}) \\
&+ \bar{W}(\text{SM E response 2} \longrightarrow \text{SM PV response 2}) \\
&+ \bar{W}(\text{SM E response 3} \longrightarrow \text{SM PV response 3}) \\
&- \bar{W}(\text{SM E response 1} \longrightarrow \text{SM PV response 2 and 3}) \\
&- \bar{W}(\text{SM E response 2} \longrightarrow \text{SM PV response 1 and 3}) \\
&- \bar{W}(\text{SM E response 3} \longrightarrow \text{SM PV response 1 and 2}).
\end{aligned}
\tag{22}
$$

$$
\begin{aligned}
CB(\text{SM response PV} \longrightarrow \text{SM response E}) = {} & \bar{W}(\text{SM PV response 1} \longrightarrow \text{SM E response 2 and 3}) \\
& + \bar{W}(\text{SM PV response 2} \longrightarrow \text{SM E response 1 and 3}) \\
& + \bar{W}(\text{SM PV response 3} \longrightarrow \text{SM E response 1 and 2}) \\
& - \bar{W}(\text{SM PV response 1} \longrightarrow \text{SM E response 1}) \\
& - \bar{W}(\text{SM E response 2} \longrightarrow \text{SM PV response 2}) \\
& - \bar{W}(\text{SM E response 3} \longrightarrow \text{SM PV response 3}).
\end{aligned}
$$

691
(23)

$$
\begin{aligned}
CB(\text{SM response PV} \longrightarrow \text{SM response PV}) = {} & \bar{W}(\text{SM PV response 1} \longrightarrow \text{SM PV response 2 and 3}) \\
& + \bar{W}(\text{SM PV response 2} \longrightarrow \text{SM PV response 1 and 3}) \\
& + \bar{W}(\text{SM PV response 3} \longrightarrow \text{SM PV response 1 and 2}) \\
& - \bar{W}(\text{SM PV response 1} \longrightarrow \text{SM PV response 1}) \\
& - \bar{W}(\text{SM PV response 2} \longrightarrow \text{SM PV response 2}) \\
& - \bar{W}(\text{SM PV response 3} \longrightarrow \text{SM PV response 3}).
\end{aligned}
$$

692
(24)

693 The connectivity biases between the different rule-selective populations in the senso-
694 rimotor module are defined as

$$
\begin{aligned}
CB(\text{SM rule E} \longrightarrow \text{SM rule E}) = {} & \bar{W}(\text{SM E rule 1} \longrightarrow \text{SM E rule 1}) \\
& + \bar{W}(\text{SM E rule 2} \longrightarrow \text{SM E rule 2}) \\
& - \bar{W}(\text{SM E rule 1} \longrightarrow \text{SM E rule 2}) \\
& - \bar{W}(\text{SM E rule 2} \longrightarrow \text{SM E rule 1}).
\end{aligned}
$$

695
(25)

696 The other connectivity biases were defined similarly

$$
\begin{aligned}
CB(\text{SM rule E} \longrightarrow \text{SM rule PV}) = {} & \bar{W}(\text{SM E rule 1} \longrightarrow \text{SM PV rule 1}) \\
& + \bar{W}(\text{SM E rule 2} \longrightarrow \text{SM PV rule 2}) \\
& - \bar{W}(\text{SM E rule 1} \longrightarrow \text{SM PV rule 2}) \\
& - \bar{W}(\text{SM E rule 2} \longrightarrow \text{SM PV rule 1}).
\end{aligned}
$$

697
(26)

$$CB(\text{SM rule PV} \longrightarrow \text{SM rule E}) = \bar{W}(\text{SM PV rule 1} \longrightarrow \text{SM E rule 2})$$
$$+ \bar{W}(\text{SM PV rule 2} \longrightarrow \text{SM E rule 1})$$
$$- \bar{W}(\text{SM PV rule 1} \longrightarrow \text{SM E rule 1})$$
$$- \bar{W}(\text{SM PV rule 2} \longrightarrow \text{SM E rule 2}). \tag{27}$$

$$CB(\text{SM rule PV} \longrightarrow \text{SM rule PV}) = \bar{W}(\text{SM PV rule 1} \longrightarrow \text{SM PV rule 2})$$
$$+ \bar{W}(\text{SM PV rule 2} \longrightarrow \text{SM PV rule 1})$$
$$- \bar{W}(\text{SM PV rule 1} \longrightarrow \text{SM PV rule 1})$$
$$- \bar{W}(\text{SM PV rule 2} \longrightarrow \text{SM PV rule 2}). \tag{28}$$

The connectivity biases between the different shared feature-selective populations in the sensorimotor module are defined similarly. For the populations selective for the two colors

$$CB(\text{SM share feature (color) E} \longrightarrow \text{SM shared feature (color) E}) = \bar{W}(\text{SM E blue} \longrightarrow \text{SM E blue})$$
$$+ \bar{W}(\text{SM E red} \longrightarrow \text{SM E red})$$
$$- \bar{W}(\text{SM E blue} \longrightarrow \text{SM E red})$$
$$- \bar{W}(\text{SM E red} \longrightarrow \text{SM E blue}), \tag{29}$$

where for example $\bar{W}(\text{SM E blue} \longrightarrow \text{SM E blue})$ is the average connection strength within the neural population selective for the shared feature blue.

The other connectivity biases were defined similarly

$$CB(\text{SM shared feature (color) E} \longrightarrow \text{SM shared feature (color) PV}) = \bar{W}(\text{SM E blue} \longrightarrow \text{SM PV blue})$$
$$+ \bar{W}(\text{SM E red} \longrightarrow \text{SM PV red})$$
$$- \bar{W}(\text{SM E blue} \longrightarrow \text{SM PV red})$$
$$- \bar{W}(\text{SM E red} \longrightarrow \text{SM PV blue}). \tag{30}$$

$$CB(\text{SM shared feature (color) PV} \longrightarrow \text{SM shared feature (color) E}) = \bar{W}(\text{SM PV blue} \longrightarrow \text{SM E red})$$
$$+ \bar{W}(\text{SM PV red} \longrightarrow \text{SM E blue})$$
$$- \bar{W}(\text{SM PV blue} \longrightarrow \text{SM E blue})$$
$$- \bar{W}(\text{SM PV red} \longrightarrow \text{SM E red}).$$
$$(31)$$

$$CB(\text{SM shared feature (color) PV} \longrightarrow \text{SM shared feature (color) PV})$$
$$= \bar{W}(\text{SM PV blue} \longrightarrow \text{SM PV red})$$
$$+ \bar{W}(\text{SM PV red} \longrightarrow \text{SM PV blue})$$
$$- \bar{W}(\text{SM PV blue} \longrightarrow \text{SM PV blue})$$
$$- \bar{W}(\text{SM PV red} \longrightarrow \text{SM PV red}).$$
$$(32)$$

The connectivity biases between populations selective for different shared shapes were defined analogously by substituting blue and red with circle and triangle.

Simulation of the optogenetic inhibition

Optogenetic inhibition was simulated by clamping the activity of neurons at 0 throughout the entire trial and the inter-trial interval.

Principal angle between subspaces

The principal angle between two subspaces is a generalization of angle between lines and planes in Euclidean space to arbitrary dimensions [77]. It can be computed by iteratively finding pairs of unit length "principal vectors", one from each subspace, that have the greatest inner product, subject to the condition that the principal vectors are orthogonal to all previous principal vectors [78].

In computing the principal angles between different rule-selective and response-selective subspaces, we first determined the dimensionality of the subspaces using the participation ratio [79]. Then the principal angles were computed using the "subspace_angles" function from the Python package Scipy. The largest principal angle was used.

To obtain a shuffled distribution, we first evenly split all trials belonging to a particular rule or response into two halves. Then, we generated two subspaces from neural trajectories during the two group of trials. A principal angle between these two subspaces was then computed for each rule/response. The angles were then averaged across all rules/responses

729 to obtain a principal angle from shuffled data. This process was repeated 100 times to
730 generate a distribution of principal angles from shuffled data.

731 **Assessing the strength of non-linear mixed selectivity**

The extent to which neurons in the sensorimotor module encode the conjunction of
stimulus and rule in a non-linear fashion was evaluated using the coefficient of determination
of a linear regression model. To tease apart non-linear and linear mixed selectivity, we first
fitted the mean activity of each neuron during response period using a set of regressors that
represent either the rule or the stimulus alone:

$$FR(n, tr) = \sum_s \beta_{n,s} \mathbb{1}(\text{stim}(tr) = s) + \sum_r \beta_{n,r} \mathbb{1}(\text{rule}(tr) = r), \tag{33}$$

732 where $FR(n, tr)$ is the firing rate of neuron n during trial tr. $\mathbb{1}$ is the indicator function.
733 For example, $\mathbb{1}(\text{stim}(tr) = s) = 1$ if the stimulus during trial tr is s, and it is 0 otherwise.

Then, another linear regression model was fitted on the residual activity unexplained
by the linear regression model above, using the conjunction of rule and stimulus as regres-
sors:

$$\tilde{FR}(n, tr) = \sum_{s,r} \beta_{n,s,r} \mathbb{1}(\text{stim}(tr) = s, \text{rule}(tr) = r), \tag{34}$$

734 where $\tilde{FR}(n, tr)$ is the firing rate of neuron n during trial tr subtracted by the pre-
735 dicted firing rate from the model defined by Equation 33. The R^2 value of this regression
736 model was used to represent the strength of non-linear mixed selectivity.

References

[1] David A Grant and Esta Berg. A behavioral analysis of degree of reinforcement and ease of shifting to new responses in a weigl-type card-sorting problem. *Journal of experimental psychology*, 38(4):404, 1948.

[2] Brenda Milner. Effects of different brain lesions on card sorting: The role of the frontal lobes. *Archives of neurology*, 9(1):90–100, 1963.

[3] R Dias, TW Robbins, and AC Roberts. Primate analogue of the wisconsin card sorting test: effects of excitotoxic lesions of the prefrontal cortex in the marmoset. *Behavioral neuroscience*, 110:872, 1996.

[4] RE Passingham. Non-reversal shifts after selective prefrontal ablations in monkeys (macaca mulatta). *Neuropsychologia*, 10(1):41–46, 1972.

[5] Katsuyuki Sakai. Task set and prefrontal cortex. *Annu. Rev. Neurosci.*, 31:219–245, 2008.

[6] M. J. Buckley, F. A. Mansouri, H. Hoda, M. Mahboubi, P. G. Browning, S. C. Kwok, A. Phillips, and K. Tanaka. Dissociable components of rule-guided behavior depend on distinct medial and prefrontal regions. *Science*, 325:52–58, 2009.

[7] E. K. Miller and J. D. Cohen. An integrative theory of prefrontal cortex function. *Annual Review of Neuroscience*, 24:167–202, 2001.

[8] Farshad A Mansouri, Kenji Matsumoto, and Keiji Tanaka. Prefrontal cell activities related to monkeys' success and failure in adapting to rule changes in a wisconsin card sorting test analog. *Journal of Neuroscience*, 26(10):2745–2756, 2006.

[9] Tsukasa Kamigaki, Tetsuya Fukushima, and Yasushi Miyashita. Cognitive set reconfiguration signaled by macaque posterior parietal neurons. *Neuron*, 61(6):941–951, 2009.

[10] Morteza Sarafyazd and Mehrdad Jazayeri. Hierarchical reasoning by neural circuits in the frontal cortex. *Science*, 364(6441):eaav8911, 2019.

[11] Takuya Ito, Guangyu Robert Yang, Patryk Laurent, Douglas H Schultz, and Michael W Cole. Constructing neural network models from brain data reveals representational transformations linked to adaptive behavior. *Nature communications*, 13(1):673, 2022.

[12] Kristen Delevich, Jason Tucciarone, Z Josh Huang, and Bo Li. The mediodorsal thalamus drives feedforward inhibition in the anterior cingulate cortex via parvalbumin interneurons. *Journal of Neuroscience*, 35(14):5743–5753, 2015.

[13] Hyun-Jae Pi, Balázs Hangya, Duda Kvitsiani, Joshua I Sanders, Z Josh Huang, and Adam Kepecs. Cortical interneurons that specialize in disinhibitory control. *Nature*, 503(7477):521–524, 2013.

[14] Siyu Zhang, Min Xu, Tsukasa Kamigaki, Johnny Phong Hoang Do, Wei-Cheng Chang, Sean Jenvay, Kazunari Miyamichi, Liqun Luo, and Yang Dan. Long-range and local circuits for top-down modulation of visual cortex processing. *Science*, 345(6197):660–665, 2014.

[15] William Muñoz, Robin Tremblay, Daniel Levenstein, and Bernardo Rudy. Layer-specific modulation of neocortical dendritic inhibition during active wakefulness. *Science*, 355(6328):954–959, 2017.

[16] Andreas J Keller, Mario Dipoppa, Morgane M Roth, Matthew S Caudill, Alessandro Ingrosso, Kenneth D Miller, and Massimo Scanziani. A disinhibitory circuit for contextual modulation in primary visual cortex. *Neuron*, 108(6):1181–1193, 2020.

[17] X. J. Wang, J. Tegnér, C. Constantinidis, and P. S. Goldman-Rakic. Division of labor among distinct subtypes of inhibitory neurons in a cortical microcircuit of working memory. *Proceedings of the National Academy of Science, USA*, 101(5):1368–73, 2004.

[18] A. Kepecs and G. Fishell. Interneuron cell types are fit to function. *Nature*, 505:318–326, 2014.

[19] R. Tremblay, S. Lee, and B. Rudy. GABAergic interneurons in the neocortex: From cellular properties to circuits. *Neuron*, 91:260–292, 2016.

[20] Guangyu Robert Yang, John D Murray, and Xiao-Jing Wang. A dendritic disinhibitory circuit mechanism for pathway-specific gating. *Nature communications*, 7(1):1–14, 2016.

[21] Kiyoshi Nakahara, Toshihiro Hayashi, Seiki Konishi, and Yasushi Miyashita. Functional mri of macaque monkeys performing a cognitive set-shifting task. *Science*, 295(5559):1532–1536, 2002.

[22] G. R. Yang and X.-J. Wang. Artificial neural networks for neuroscientists: a primer. *Neuron*, 107:1048–1070, 2020.

[23] Monika Jadi, Alon Polsky, Jackie Schiller, and Bartlett W Mel. Location-dependent effects of inhibition on local spiking in pyramidal neuron dendrites. *PLoS computational biology*, 8(6):e1002550, 2012.

[24] Valerio Mante, David Sussillo, Krishna V Shenoy, and William T Newsome. Context-dependent computation by recurrent dynamics in prefrontal cortex. *nature*, 503(7474):78–84, 2013.

[25] Guangyu Robert Yang, Madhura R Joglekar, H Francis Song, William T Newsome, and Xiao-Jing Wang. Task representations in neural networks trained to perform many cognitive tasks. *Nature neuroscience*, 22(2):297–306, 2019.

[26] H Francis Song, Guangyu R Yang, and Xiao-Jing Wang. Training excitatory-inhibitory recurrent neural networks for cognitive tasks: a simple and flexible framework. *PLoS computational biology*, 12(2):e1004792, 2016.

[27] Joaquin M Fuster. Unit activity in prefrontal cortex during delayed-response performance: neuronal correlates of transient memory. *Journal of neurophysiology*, 36(1):61–78, 1973.

[28] Shintaro Funahashi, Charles J Bruce, and Patricia S Goldman-Rakic. Mnemonic coding of visual space in the monkey's dorsolateral prefrontal cortex. *Journal of neurophysiology*, 61(2):331–349, 1989.

[29] P. Goldman-Rakic. Cellular basis of working memory. *Neuron*, 14:477–85, 1995.

[30] Ranulfo Romo, Carlos D Brody, Adrián Hernández, and Luis Lemus. Neuronal correlates of parametric working memory in the prefrontal cortex. *Nature*, 399(6735):470–473, 1999.

[31] Thomas B Christophel, P Christiaan Klink, Bernhard Spitzer, Pieter R Roelfsema, and John-Dylan Haynes. The distributed nature of working memory. *Trends in cognitive sciences*, 21(2):111–124, 2017.

[32] Matthew L Leavitt, Diego Mendoza-Halliday, and Julio C Martinez-Trujillo. Sustained activity encoding working memories: not fully distributed. *Trends in Neurosciences*, 40(6):328–346, 2017.

[33] Zengcai V Guo, Hidehiko K Inagaki, Kayvon Daie, Shaul Druckmann, Charles R Gerfen, and Karel Svoboda. Maintenance of persistent activity in a frontal thalamocortical loop. *Nature*, 545(7653):181–186, 2017.

[34] Kartik K Sreenivasan and Mark D'Esposito. The what, where and how of delay activity. *Nature reviews neuroscience*, 20(8):466–481, 2019.

[35] Kong-Fatt Wong and Xiao-Jing Wang. A recurrent network mechanism of time integration in perceptual decisions. *Journal of Neuroscience*, 26(4):1314–1328, 2006.

[36] Farzaneh Najafi, Gamaleldin F Elsayed, Robin Cao, Eftychios Pnevmatikakis, Peter E Latham, John P Cunningham, and Anne K Churchland. Excitatory and inhibitory subnetworks are equally selective during decision-making and emerge simultaneously during learning. *Neuron*, 105(1):165–179, 2020.

[37] James P Roach, Anne K Churchland, and Tatiana A Engel. Choice selective inhibition drives stability and competition in decision circuits. *Nature Communications*, 14(1):147, 2023.

[38] Hongbo Jia, Nathalie L Rochefort, Xiaowei Chen, and Arthur Konnerth. Dendritic organization of sensory input to cortical neurons in vivo. *Nature*, 464(7293):1307–1312, 2010.

[39] Joseph Cichon and Wen-Biao Gan. Branch-specific dendritic ca2+ spikes cause persistent synaptic plasticity. *Nature*, 520(7546):180–185, 2015.

[40] Shannon K Rashid, Victor Pedrosa, Martial A Dufour, Jason J Moore, Spyridon Chavlis, Rodrigo G Delatorre, Panayiota Poirazi, Claudia Clopath, and Jayeeta Basu. The dendritic spatial code: branch-specific place tuning and its experience-dependent decoupling. *BioRxiv*, pages 2020–01, 2020.

[41] Jakob Voigts and Mark T Harnett. Somatic and dendritic encoding of spatial variables in retrosplenial cortex differs during 2d navigation. *Neuron*, 105(2):237–245, 2020.

[42] Mattia Rigotti, Omri Barak, Melissa R Warden, Xiao-Jing Wang, Nathaniel D Daw, Earl K Miller, and Stefano Fusi. The importance of mixed selectivity in complex cognitive tasks. *Nature*, 497(7451):585–90, 2013.

[43] Atsushi Kikumoto and Ulrich Mayr. Conjunctive representations that integrate stimuli, responses, and rules are critical for action selection. *Proceedings of the National Academy of Sciences*, 117(19):10603–10608, 2020.

[44] Atsushi Kikumoto, Ulrich Mayr, and David Badre. The role of conjunctive representations in prioritizing and selecting planned actions. *Elife*, 11:e80153, 2022.

[45] Jonathan D Wallis, Kathleen C Anderson, and Earl K Miller. Single neurons in prefrontal cortex encode abstract rules. *Nature*, 411(6840):953–956, 2001.

[46] Matthew M Botvinick, Jonathan D Cohen, and Cameron S Carter. Conflict monitoring and anterior cingulate cortex: an update. *Trends in cognitive sciences*, 8(12):539–546, 2004.

[47] René Quilodran, Marie Rothe, and Emmanuel Procyk. Behavioral shifts and action valuation in the anterior cingulate cortex. *Neuron*, 57(2):314–325, 2008.

[48] Nils Kolling, Marco K Wittmann, Tim EJ Behrens, Erie D Boorman, Rogier B Mars, and Matthew FS Rushworth. Value, search, persistence and model updating in anterior cingulate cortex. *Nature neuroscience*, 19(10):1280–1285, 2016.

[49] Farshad Alizadeh Mansouri, David J Freedman, and Mark J Buckley. Emergence of abstract rules in the primate brain. *Nature Reviews Neuroscience*, 21(11):595–610, 2020.

[50] Timothy Spellman, Malka Svei, Jesse Kaminsky, Gabriela Manzano-Nieves, and Conor Liston. Prefrontal deep projection neurons enable cognitive flexibility via persistent feedback monitoring. *Cell*, 184(10):2750–2766, 2021.

[51] C Daniel Salzman and Stefano Fusi. Emotion, cognition, and mental state representation in amygdala and prefrontal cortex. *Annual review of neuroscience*, 33:173–202, 2010.

[52] Clay B Holroyd and Michael GH Coles. The neural basis of human error processing: reinforcement learning, dopamine, and the error-related negativity. *Psychological review*, 109(4):679, 2002.

[53] Masayuki Matsumoto and Okihide Hikosaka. Two types of dopamine neuron distinctly convey positive and negative motivational signals. *Nature*, 459(7248):837–841, 2009.

[54] Richard A Andersen and He Cui. Intention, action planning, and decision making in parietal-frontal circuits. *Neuron*, 63(5):568–583, 2009.

[55] Bernard W Balleine and John P O'doherty. Human and rodent homologies in action control: corticostriatal determinants of goal-directed and habitual action. *Neuropsychopharmacology*, 35(1):48–69, 2010.

[56] Bevil R Conway. Color vision, cones, and color-coding in the cortex. *The neuroscientist*, 15(3):274–290, 2009.

[57] Rosa Lafer-Sousa and Bevil R Conway. Parallel, multi-stage processing of colors, faces and shapes in macaque inferior temporal cortex. *Nature neuroscience*, 16(12):1870–1878, 2013.

[58] Le Chang, Pinglei Bao, and Doris Y Tsao. The representation of colored objects in macaque color patches. *Nature communications*, 8(1):2064, 2017.

[59] Mark M Churchland, John P Cunningham, Matthew T Kaufman, Justin D Foster, Paul Nuyujukian, Stephen I Ryu, and Krishna V Shenoy. Neural population dynamics during reaching. *Nature*, 487(7405):51–56, 2012.

[60] Nicholas A Steinmetz, Peter Zatka-Haas, Matteo Carandini, and Kenneth D Harris. Distributed coding of choice, action and engagement across the mouse brain. *Nature*, 576(7786):266–273, 2019.

[61] Mattia Rigotti, Daniel Ben Dayan Rubin, Xiao-Jing Wang, and Stefano Fusi. Internal representation of task rules by recurrent dynamics: the importance of the diversity of neural responses. *Frontiers in computational neuroscience*, 4:24, 2010.

[62] Stanislas Dehaene and Jean-Pierre Changeux. The wisconsin card sorting test: Theoretical analysis and modeling in a neuronal network. *Cerebral cortex*, 1(1):62–79, 1991.

[63] Daniel Turner-Evans, Stephanie Wegener, Herve Rouault, Romain Franconville, Tanya Wolff, Johannes D Seelig, Shaul Druckmann, and Vivek Jayaraman. Angular velocity integration in a fly heading circuit. *Elife*, 6:e23496, 2017.

[64] Benjamin Y Hayden, John M Pearson, and Michael L Platt. Neuronal basis of sequential foraging decisions in a patchy environment. *Nature neuroscience*, 14(7):933–939, 2011.

[65] João D Semedo, Amin Zandvakili, Christian K Machens, M Yu Byron, and Adam Kohn. Cortical areas interact through a communication subspace. *Neuron*, 102(1):249–259, 2019.

[66] Christopher Langdon, Mikhail Genkin, and Tatiana A Engel. A unifying perspective on neural manifolds and circuits for cognition. *Nature Reviews Neuroscience*, pages 1–15, 2023.

[67] Ken-Ichiro Tsutsui, Takayuki Hosokawa, Munekazu Yamada, and Toshio Iijima. Representation of functional category in the monkey prefrontal cortex and its rule-dependent use for behavioral selection. *Journal of Neuroscience*, 36(10):3038–3048, 2016.

[68] Vishwa Goudar, Jeong-Woo Kim, Yue Liu, Adam JO Dede, Michael J Jutras, Ivan Skelin, Michael Ruvalcaba, William Chang, Adrienne L Fairhall, Jack J Lin, et al. Comparing rapid rule-learning strategies in humans and monkeys. *bioRxiv*, pages 2023–01, 2023.

[69] Steven W Kennerley, Mark E Walton, Timothy EJ Behrens, Mark J Buckley, and Matthew FS Rushworth. Optimal decision making and the anterior cingulate cortex. *Nature neuroscience*, 9(7):940–947, 2006.

[70] Braden A Purcell and Roozbeh Kiani. Hierarchical decision processes that operate over distinct timescales underlie choice and changes in strategy. *Proceedings of the national academy of sciences*, 113(31):E4531–E4540, 2016.

[71] Cheng Xue, Lily E Kramer, and Marlene R Cohen. Dynamic task-belief is an integral part of decision-making. *Neuron*, 110(15):2503–2511, 2022.

[72] Ido Ben-Artzi, Yoav Kessler, Bruno Nicenboim, and Nitzan Shahar. Computational mechanisms underlying latent value updating of unchosen actions. *Science Advances*, 9(42):eadi2704, 2023.

[73] Amitai Shenhav, Matthew M Botvinick, and Jonathan D Cohen. The expected value of control: an integrative theory of anterior cingulate cortex function. *Neuron*, 79(2):217–240, 2013.

[74] Xiaolong Jiang, Shan Shen, Cathryn R Cadwell, Philipp Berens, Fabian Sinz, Alexander S Ecker, Saumil Patel, and Andreas S Tolias. Principles of connectivity among morphologically defined cell types in adult neocortex. *Science*, 350(6264):aac9462, 2015.

[75] Paul J Werbos. Backpropagation through time: what it does and how to do it. *Proceedings of the IEEE*, 78(10):1550–1560, 1990.

[76] Diederik P Kingma and Jimmy Ba. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.

[77] Camille Jordan. Essai sur la géométrie à n dimensions. *Bulletin de la Société mathématique de France*, 3:103–174, 1875.

[78] ke Björck and Gene H Golub. Numerical methods for computing angles between linear subspaces. *Mathematics of computation*, 27(123):579–594, 1973.

[79] Peiran Gao, Eric Trautmann, Byron Yu, Gopal Santhanam, Stephen Ryu, Krishna Shenoy, and Surya Ganguli. A theory of multineuronal dimensionality, dynamics and measurement. *BioRxiv*, page 214262, 2017.

Supplementary Figure 1: **Activity of single neurons**.
Left: activity from example neurons across three consecutive trials with a rule switch. Blue and green traces represent neurons with higher activity during the color rule and shape rule respectively. Red traces for the PFC excitatory neurons represent neurons that are preferentially activated by negative feedback.
Right: Rule selectivity across neurons and task epochs.
VIP neurons in the PFC module do not receive any excitatory inputs, therefore are not shown.

Supplementary Figure 2: **Two attractor states in the PFC module supported by inter-modular connections.**

The neural trajectories that represent the autonomous dynamics of the trained RNN with intact inter-modular connections (**a**), when the connections from the PFC module to the sensorimotor module were lesioned (**b**), and when the connections from the sensorimotor module to the PFC module were lesioned (**c**). The networks started from random time point during color rule (blue) and shape rule (green) trials. Black points represent the final states of the networks after 500 timesteps (5 seconds).

Supplementary Figure 3: **The emergence of two populations of excitatory neurons in the PFC module of networks with divisive dendritic nonlinearity.** The rule modulation against input weight for negative feedback for all the rule neurons and conjunctive error x rule neurons in the PFC module of networks with divisive dendritic nonlinearity (c.f. Figure 2c).

Supplementary Figure 4: The connectivity biases between different subpopulations of excitatory and PV neurons in the PFC module of networks with divisive dendritic nonlinearity (c.f. Figure 3d).

Supplementary Figure 5: **The connectivity structure between the excitatory neurons and SST neurons in the PFC module. a**. The connectivity bias between the SST neurons and the conjunctive error x rule neurons in the PFC module, for networks with subtractive dendritic nonlinearity. **b**. The connectivity bias between the SST neurons and the rule neurons in the PFC module, for networks with subtractive dendritic nonlinearity. **c** - **d**. The same connectivity biases but for networks with divisive dendritic nonlinearity. **e** The connectivity structure between the SST and excitatory neurons in the PFC module as revealed by **a**- **d**. Only the stronger connections are plotted. A similar connectivity pattern exists between SST and excitatory neurons as compared to the connectivity pattern between PV and excitatory neurons (c.f. Figure 3c).

Supplementary Figure 6: **Single neurons in the sensorimotor module are conjunctively selective for rule, response and shared feature.**
a. The distribution of rule selectivity across different cell types in the sensorimotor module. Result is aggregated across all trained networks.
b. The trial-averaged activity of an example excitatory neuron in the sensorimotor module with preferred rule $R = color\ rule$, response location $L = 3$ and shared feature $F = circle$. Trials were sorted according to rule (left), response (middle) and shared feature (right).

.

Supplementary Figure 7: **Structure in the top-down projections, across all networks.**

a. Each line represents the mean connection strength onto one excitatory neuron in the sensorimotor module, from the PFC excitatory neurons that prefer the same rule and the different rule. Bars represent the mean across neurons. PFC excitatory neurons project more strongly to the excitatory neurons in the sensorimotor module that prefer the same rule (c.f. Figure 4a).

b. Each line represents the mean connection strength onto one VIP neuron in the sensorimotor module, from the PFC excitatory neurons that prefer the same rule and the different rule. Bars represent the mean across neurons. PFC excitatory neurons project more strongly to the VIP neurons in the sensorimotor module that prefer the same rule (c.f. Figure 4b).

c. Each line represents the mean connection strength onto one PV neuron in the sensorimotor module, from the PFC excitatory neurons that prefer the same rule and the different rule. Bars represent the mean across neurons. PFC excitatory neurons project more strongly to the PV neurons in the sensorimotor module that prefer the same rule (c.f. Figure 4c).

d. Each line represents the mean connection strength onto one SST neuron in the sensorimotor module, from the PFC excitatory neurons that prefer the same rule and the different rule. Bars represent the mean across neurons. PFC excitatory neurons project more strongly to local SST neurons that prefer the same rule (c.f. Figure 4d).

e. Each line represents the mean connection strength onto one excitatory neuron of the sensorimotor module, from the SST neurons in the sensorimotor module that prefer the same rule and the different rule. Bars represent mean across neurons. SST neurons in the sensorimotor module project more strongly to local excitatory neurons that prefer the opposite rule (c.f. Figure 4e).

f. Each line represents the mean connection strength onto one excitatory neuron of the sensorimotor module, from the PV neurons in the sensorimotor module that prefer the same rule and the different rule. Bars represent mean across neurons. PV neurons in the sensorimotor module project more strongly to local excitatory neurons that prefer the opposite rule (c.f. Figure 4f).

g - **l**. Same as **a** - **f** for networks with the divisive dendritic nonlinearity.

$p < 0.001$ for all panels, Student's t test.

Supplementary Figure 8: The structure in the input and output weights of the sensorimotor module. Data is aggregated across all trained networks with subtractive (**a**, **c**) and divisive (**b**, **d**) dendritic nonlinearity (c.f. Figure 5b, c). Student's t-test, $p < .001$ for all panels.

Supplementary Figure 9: **a-b**. The principal angle between different rule and response subspaces for networks trained with divisive dendritic nonlinearity (c.f. Figure 6c-d). **c-e**. The connectivity bias between different subpopulations of excitatory and PV neurons in the sensorimotor module of networks with divisive dendritic nonlinearity (c.f. Figure 6e-g).

Supplementary Figure 10: **a**. The relationship between the sparsity of the connection from SST to the dendrites of excitatory neurons in the sensorimotor module, for networks with divisive dendritic nonlinearity (c.f. Figure 7d). **b-d**. The performance (**b**), principal angle between rule subspaces (**c**) and the strength of conjunctive coding (**d**) decreased after silencing SST neurons in the sensorimotor module, for networks trained with divisive dendritic nonlinearity (for **d**, Student's t-test, $p < .001$. Each line represents one neuron. Results are aggregated across networks). c.f. Figure 7e-g.

Supplementary Figure 11: Example neurons in the sensorimotor module showing decreased conjunctive coding of rule and stimulus when SST neurons in the sensorimotor module are silenced. The strength of conjunctive coding is assessed by the R^2 value of a linear regression model where the independent variables are conjunctions of rule and stimulus and the dependent variable is the trial-to-trial residual neural activity unexplained by a model with only rule and stimulus as independent variables (see Methods for details).