Geometry of anisotropic contextual interactions in the visual cortex places fundamental limits on spatial vision.

Mitchell P. Morton[1,2], Sachira Denagamage[1,2], Nyomi V. Hudson[1], and Anirvan S. Nandy[1-4]

[1]Department of Neuroscience, Yale University, New Haven, CT 06510
[2]Interdepartmental Neuroscience Program, Yale University, New Haven, CT 06510
[3]Wu Tsai Institute, Yale University, New Haven, CT 06511
[4]Kavli Institute for Neuroscience, Yale University, New Haven, CT 06511

Correspondence: mitchell.morton@yale.edu, anirvan.nandy@yale.edu

ABSTRACT

Crowding, the impaired ability to accurately recognize a target stimulus among distractors, is a major bottleneck in visual perception. The spatial configuration of distractors in relation to the target profoundly influences perceptual fidelity. Notably, when a distractor is placed at a more eccentric point on the radial axis (termed 'radial-out crowding'), it exerts the strongest impairment. Despite the pronounced perceptual anisotropy, the prevalent assumption underlying our understanding of contextual interactions in the visual cortex assumes isotropy. We investigated how distractor stimuli in different spatial configurations impacted the representation of a target stimulus in laminar microcircuits in the primary visual cortex (V1). Our study reveals that radial-out crowding more strongly impacts the ability to decode the target orientation from V1 population activity compared to other spatial configurations. This effect was strongest among putative excitatory neurons in the superficial and input layers, which are the primary neural populations involved in feed-forward information propagation. Remarkably, the feedback pathway involving the deep cortical layers does not exhibit anisotropy. Mechanistically, the anisotropy is explained by a tuned suppression and

untuned facilitation of orientation responses, leading to an anisotropic broadening of

tuning curves in the feedforward pathway, but not in the feedback pathway. These results

underscore the non-uniform spatial integration of information by neurons in the visual

cortex, establishing the presence of anisotropic contextual interactions in the earliest

stages of cortical processing. By elucidating the distinct roles of feedforward and

feedback pathways in the context of crowding, this study advances our understanding of

the intricate interplay between spatial arrangement, neural circuitry, and the constraints

on perceptual fidelity during early visual processing.

INTRODUCTION

Our ability to perceive objects in the visual world is profoundly limited by the phenomenon of visual crowding, which decreases our ability to recognize target objects among visual clutter (*1*). Crowding not only limits both perception and visually guided behavior in normally sighted individuals but also impacts patients with conditions ranging from macular degeneration to dyslexia (*2*). The area in visual space over which distractor stimuli impact the perception of a central target is referred to as the crowding zone. A fundamental property of crowding zones is their anisotropy in visual space (for review, see *3*). Psychophysical studies of crowding have identified two major anisotropic characteristics of crowding in perception. First, distractor stimuli positioned on the radial axis exert a larger influence on perception than distractors on the tangential axis (*4*). Second, crowding is asymmetrical along the radial axis, such that distractors that are more eccentric than the target more strongly impair perception than distractors that are less eccentric than the target (*5*). Together, these results indicate that distractor stimuli positioned on the radial axis and more eccentric than the target (radial-out crowding) are more effective at inducing crowding effects than stimuli either on the radial axis but less eccentric than the target (radial-in crowding) or on the tangential axis (tangential crowding).

Psychophysical studies have suggested that crowding emerges in the cortex (*6, 7*), and various models (*8-12*) have been proposed to explain crowding effects based on the anatomy and connectivity within the primary visual cortex (V1). Despite these efforts, very little experimental evidence exists to validate or refute any proposed neural mechanisms. While the perceptual effects of crowding have been studied for many decades (*1, 3, 9, 13-15*), it was only recently that there has emerged direct evidence of

neural correlates underlying the phenomenon (*16, 17*). The question of exactly when and

how crowding effects emerge in the visual hierarchy remains unresolved, but recent work

has shown that crowding impairs feature representations as early as V1 (*16, 17*).

However, the neural mechanisms responsible for how the spatial geometry of stimuli

affects the strength of crowding remain poorly understood at the level of cortical

microcircuits composed of multiple laminae and cell classes (*18-21*).

We sought to understand how stimuli presented outside the receptive field center

could influence stimulus representations in the V1 laminar microcircuit in a spatially

anisotropic manner. We hypothesized that radial-out crowding would most strongly

impact the representation of target stimuli among excitatory neurons in the output layers

of V1. We further hypothesized that these anisotropic contextual interactions might be

most prevalent in the subpopulations involved in feed-forward information propagation,

particularly excitatory neurons in the input and superficial layers.

RESULTS

To study how the spatial configuration of stimuli influences neural responses in area V1,

we trained monkeys to perform a passive fixation task while we presented stimuli in

different geometric arrangements. Monkeys fixated on the center of the screen and

passively viewed stimuli presented either alone (target-only condition) or along with a

distractor in one of four locations (crowding). We defined three crowding conditions based

on the spatial position of the distractor relative to the target (Figure 1A, see Methods).

The radial-in and radial-out crowding conditions positioned the distractor on the radial

axis connecting the target to the fixation point. In the radial-in condition, the distractor was

placed between the target and the fixation point, while in the radial-out condition, the

distractor was past the target on the radial axis. For the tangential crowding condition, the

distractor was positioned on the tangential axis (the line orthogonal to the radial axis and

passing through the target) on either side of the target.

During the task, we performed laminar extracellular recordings (*22, 23*) of single

unit activity in V1 (Figure 1B, S1A-E). We positioned the stimuli such that the target

stimulus was presented at the center of the aggregate receptive field of the recording

sites. During our recordings, we collected data from 905 well-isolated single units (Figure

1B, 511 from monkey M, 394 from monkey D). We used established procedures (*22, 24-*

28) to classify these neurons based on their waveform shape into broad-spiking (putative

excitatory) and narrow-spiking (putative inhibitory) neurons (Figure 1B-C). Many

observed differences between broad- and narrow-spiking neurons suggest they

correspond well with excitatory and inhibitory neurons respectively (*22, 25, 26, 28-32*).

We used current source density (CSD) analysis to identify the superficial (II/III), input (IV),

and deep (V & VI) cortical layers (*22, 33*) (Figure S1D). This allowed us to analyze the

different impacts of distractor stimuli based on their spatial location in six subpopulations

of the laminar cortical circuit: broad- and narrow-spiking neurons in each of the three

layers. Each of these subpopulations was visually responsive and well-tuned to the

orientation of the target stimulus (Figure S2A-B).

To examine how contextual interactions among stimuli might impact information

coding in V1 in an anisotropic manner, we first determined how well we could decode the

identity (orientation and phase) of the target, alone or in each of the crowding conditions,

from simultaneously recorded V1 population activity using linear discriminant analysis

(Figure 2). All distractor stimuli impaired decoding performance, with radial-out crowding

showing the largest performance decrease (Figure 2).

We next sought to understand how distractor location differentially influences

information processing among laminar cortical circuit components. For each

subpopulation, we calculated the decoding performance for the target-only and crowding

conditions (Figure 3). Most crowding stimuli impair decoding performance compared to

the target-only in most subpopulations (Figure 3). In particular, the impairment of

decoding performance is worse in the radial-out condition compared to the tangential and

radial-in conditions among broad-spiking neurons in the input and superficial layers, the

two subpopulations that are key nodes in the feed-forward cortical information flow

pathway (Figure 3, middle right and top right). Radial-out crowding also impairs decoding

performance more than radial-in crowding in narrow-spiking neurons across all layers,

although radial-out crowding has a similar impact compared to tangential crowding in

these subpopulations (Figure 3, left column). Consistent with previous reports (*17*), we

also found that more effective crowding stimuli (radial-out crowding) most strongly

reduced spike-count correlations, and this result was strongest in the feed-forward

pathway (Figure S3).

Because distractor location has an anisotropic impact on the ability to decode

target orientation from population activity, we investigated whether differential modulation

in the tuning properties of V1 neurons could underlie these changes. We normalized the

tuning curve for each single unit based on its preferred target orientation (see Methods)

and plotted the average tuning curve of each subpopulation in response to the target only

and in the crowding conditions (Figure 4). Distractors broaden the tuning curves of V1

neurons in each of the subpopulations (Figure 4). All subpopulations showed substantially

reduced responses when distractors were presented alone (Figure S4).

To quantify how distractor location modulates tuning, we first calculated the firing

rates in response to preferred and orthogonal-to-preferred (orthogonal) target

orientations. Crowding stimuli reduced firing rates in response to the preferred target

orientation and increased the firing rates in response to the orthogonal target orientation

(Figure 5). This correspondingly resulted in a similar decrease in orientation selectivity

index (OSI) under different crowding conditions in each subpopulation (Figure S2C).

Although OSI was similar across conditions and subpopulations, there were distinct

differences in how flanker location modulated preferred and orthogonal firing rates. In

both narrow and broad-spiking neurons in the input layer, radial-out flankers resulted in

less suppression of firing to the preferred orientation, but more effectively facilitated firing

in response to the orthogonal orientation (Figure 5, middle row). This pattern was also

consistent among broad-spiking neurons in the superficial layers (Figure 5, top right).

To further characterize these results and obtain mechanistic insights into these

changes in tuning properties, we quantified the extent to which distractor location induced

untuned additive facilitation or tuned divisive suppression (*16, 17*) of V1 tuning curves.

We modeled each neuron's tuning curve in the target-only condition and then calculated

the extent to which the responses to stimuli under each of the crowding conditions were

best explained by additive and multiplicative shifts to the target-only tuning curve (Figure

6 & S5). Input layer neurons under radial-out crowding showed greater additive facilitation

compared to tangential or radial-in crowding. (Figure 6A, middle row). The radial-out

condition also showed greater divisive suppression in broad-spiking neurons in both the

input and superficial layers compared to the tangential and radial-in conditions (Figure

6B, top right and middle right). This effect was similar in narrow-spiking neurons in the

input and superficial layers, with radial-out crowding showing more divisive suppression

in both subpopulations compared to radial-in crowding and trending towards more

suppression compared to tangential crowding (Figure 6B, top left and middle left). Taken

together, these results show that radial-out crowding most effectively broadens tuning in

the feed-forward pathway through untuned facilitation and tuned suppression.

DISCUSSION

Natural vision activates neurons not in isolation, but in the context of other stimuli. The

extensive literature on contextual interactions in the visual cortex, particularly surround

modulation (*34-42*), has significantly enriched our understanding of visual processing.

However, a prevailing assumption in this literature has been the isotropic nature of the

surrounding context. This is evident in the experimental design of contextual stimuli

typically as annuli or as a circular constellation of flankers (*34, 35, 38, 43*). While several

studies (*44-46*) did explore the spatial distribution of contextual interactions and their

impact on neural responses, systematic geometrical arrangements of target and flanker

objects with respect to the center of gaze were not investigated. Furthermore, it has been

shown that intrinsic horizontal connections in V1 are anisotropic in a manner that could

facilitate contour processing, enriching connections among neurons with orientation

preferences and receptive fields that facilitate the detection of smooth contours (*47, 48*).

But here too the spatial geometries alluded to above have not been investigated.

In contrast, perceptual asymmetries in vision have garnered considerable interest

in the past several decades(*49-52*). In particular, the polar angle with respect to the center

of gaze has been shown to be especially relevant to perceptual performance (see review

by (*53*). Perceptual asymmetries are especially striking in the phenomenology of visual

crowding wherein distractors impair target identification over a distinctive elongated zone

in visual space. Cortical magnification results in radial-out distractors being represented

closer to the target in cortical space compared to radial-in distractors, potentially

explaining some of the anisotropy (*9, 10, 54*). However, it has been shown that

considerations of cortical magnification alone are not sufficient to explain the marked

elongation of the crowding zones along the radial axis defined by the center of gaze and

the target stimulus (*8*).

Macaques experience crowding similar to humans, impairing their ability to

discriminate the orientation of target stimuli (*15, 16*), and are thus a good model for

studying the neural mechanisms of crowding. Experiments in both macaques and

humans suggest that crowding can influence neural activity as early as V1 (*16, 55*),

including a radial-tangential anisotropy in BOLD responses in V1 (*56*). The present study

extends these findings by revealing the impact of distractor location on stimulus

representation in the laminar microcircuit of the primary visual cortex and the neural

mechanisms underlying these impacted representations.

Notably, the effects of radial-out crowding were more pronounced compared to

tangential and radial-in crowding across various measures. These effects included the

ability to decode target identity from simultaneously recorded populations, as well as from

putative excitatory neurons in the input and superficial layers. Anisotropies were also

evident in the modulation of tuning curves in these layers, with radial-out crowding

producing stronger broadening of tuning curves through untuned additive facilitation and

tuned divisive suppression. Additionally, radial-out distractors exhibited the most

substantial decrease in spike-count correlations in the input and superficial layers. These

anisotropies likely contribute to the impaired perception of target stimuli induced by radial-

out distractors. Importantly, the decreased ability to decode stimulus orientation from

neural activity at the subpopulation level was not solely attributable to changes in spike-

count correlations, as correlations were removed across all conditions for those analyses.

Surprisingly, we discovered that contextual interactions exhibit anisotropy even at

the earliest stages of processing in V1, specifically within the input layer. This finding

suggests a potential sub-cortical origin of the observed anisotropies (*57-61*). Additionally,

we observed the presence of anisotropy in the superficial layer of V1, implying that these

computations are partially inherited from the input layer. Broad-spiking neurons share

electrophysiological characteristics with excitatory pyramidal neurons (*26, 28, 32*). Given

that broad-spiking putative excitatory neurons in the superficial layer are likely projection

neurons to downstream ventral visual areas (*18, 62-64*), our findings suggest that the

anisotropic signaling is propagated along the feedforward pathway and could form the

basis of the perceptual anisotropies underlying visual crowding.

In contrast, we did not find the same anisotropy in the deep layers of V1. This

intriguing result suggests the emergence of novel contextual computations in the deep

layers, indicating that tuning and response properties are not merely inherited from input

and superficial layer projections to the deep layers. This is in agreement with previous

findings that neurons in the deep layers can have differences in orientation tuning

compared to the superficial layers (*65, 66*) and that intrinsic (horizontal) connections in

the deep layers are less extensive compared to the superficial layers (*67*). Moreover, as

deep-layer broad-spiking putative excitatory neurons are the likely sources of feedback

projections to subcortical areas, these feedback signals are likely not anisotropic. Despite

the likely feedforward propagation of anisotropic signals, the absence of anisotropy in the

feedback pathway suggests a mechanism of active cancellation in this important

modulatory pathway. Speculatively, this might play a role in maintaining a veridical

representation of the visual scene in the earliest stages of extra-retinal visual processing.

The implications of our findings extend to the integration of information from

outside the classical receptive field in the visual cortex. Orientation tuning properties and

their modulation are determined by the balance of excitation and inhibition to cortical

neurons (*68, 69*). Recurrent excitation (*70*) and lateral inhibition (*71*) are two key intra-

cortical motifs that determine this balance. Cross-orientation suppression (*72*), a form of

lateral inhibition likely plays an important role in shaping the cortical activity due to flanker-

target interactions examined in this study. Such suppressive activity may operate in

several ways according to orientation dependence and the strength of inhibitory

connections and have been described in various models of cortical activity including those

with attractor dynamics (*73, 74*), recurrence (*70*), balance (*75-77*), push-pull (*78, 79*) and

normalization (*80-82*). Our results imply that distractor location differentially impacts the

balance of excitation and inhibition in the laminar cortical circuit. Future empirical and

computational modeling studies of surround modulation should take this anisotropic and

layer-specific integration into account.

In conclusion, this study provides compelling evidence that contextual interactions

in the visual cortex exhibit anisotropy at the earliest stages of cortical processing of visual

signals. Distractors broaden V1 orientations tuning curves in a spatially anisotropic

manner. Radial-out distractors most effectively broaden tuning in the feedforward

pathway through untuned facilitation and tuned suppression. These findings contribute to

our understanding of the neural bases of visual crowding and offer mechanistic insights

into the processing of contextual signals in the visual cortex.

AUTHOR CONTRIBUTIONS

251   MPM and ASN conceptualized the project. MPM collected the data, with assistance from

252   SD and NVH. MPM analyzed the data. MPM and ASN wrote the paper.

ACKNOWLEDGEMENTS

254   This research was supported by NIH/NEI R01 EY032555, NARSAD Young Investigator

255   Grant, Ziegler Foundation Grant, and Yale Orthwein Scholar Funds to ASN, NIH/NINDS

256   training grants T32-NS007224 and T32-NS041228 to MPM, and by an NIH/NEI core grant

257   for vision research P30 EY026878 to Yale University. We would like to thank the

258   veterinary and husbandry staff at Yale for excellent animal care.

FIGURE LEGENDS

Figure 1. Anisotropic contextual interaction task and single-unit recordings.
(**A**) Anisotropic contextual interaction task schematic. Stimuli were presented alone at the
receptive field center (target only) or with a distractor at one of the four crowding locations.
The radial axis connects the target to the center of gaze, and the tangential axis is
orthogonal to it, also passing through the target. The top schematic shows stimuli at all
possible locations, while the bottom section shows the possible spatial configurations for
the presentation of an individual stimulus array. (**B**) Mean waveforms for all recorded
single units (blue=narrow-spiking, orange=broad-spiking). (**C**) Distribution of waveform
durations (trough-to-peak time) for all recorded single units across the two monkeys.
Neurons with waveform durations less than 350μs were classified as narrow-spiking
(blue), and neurons with waveform durations greater than 350μs were classified as broad-
spiking (orange).

273
274

Figure 2. Distractor stimuli impair decoding performance in simultaneously recorded populations in an anisotropic manner.
Example decoding performance from 4 simultaneously recorded populations (two in each monkey) in the target-only condition and each of the three crowding conditions. Points in the upper section of each plot show the decoding performance for each of the 20 different cross-validations. The lower section for each session shows the bootstrapped estimation of the difference between decoding performance in each crowding condition compared to the target-only condition. Half-violin plots show the bootstrapped distribution of the difference, and black dots and bars represent the mean and 95% confidence intervals of the difference in decoding performance.

Figure 3. Radial-out crowding most strongly impairs decoding in the feed-forward pathway.
Decoding performance from each subpopulation in the target-only condition and each of the three crowding conditions. Points in the upper section of each plot show the decoding performance for each of the 20 different cross-validations. The lower section for each session shows the bootstrapped estimation of the difference between decoding performance in each crowding condition compared to the target-only condition. Half-violin plots show the bootstrapped distribution of the difference, and black dots and bars represent the mean and 95% confidence intervals of the difference in decoding performance.

Figure 4. Crowding stimuli modulate the tuning curves of V1 neurons.
Normalized tuning curves for each subpopulation in the target-only and crowding conditions. Tuning curves are aligned with the preferred orientation at 0 degrees and duplicated to fill the circle. Tuning curves were normalized by the maximum firing rate in the target-only condition. Lines and shading represent mean +/- s.e.m. (black=target only, pink=tangential crowding, green=radial-in crowding, purple=radial-out crowding).

Figure 5. Crowding broadens tuning curves
Difference in firing rate in response to preferred (dark brown) and orthogonal (light tan) stimuli under each crowding condition in the six subpopulations. Colored half-violin plots show the bootstrapped estimation of the paired mean difference in firing rate for a given subpopulation, crowding condition, and stimulus orientation. Black dots and lines represent the mean and 95% confidence interval of the estimated difference.

Figure 6. Radial-out crowding causes stronger additive facilitation and divisive suppression of tuning curves
(**A**) Difference in the modeled (see Methods) untuned additive facilitation caused by each of the crowding conditions. Gray half-violin plots show the bootstrapped estimation of the paired mean difference in the modeled additive component for a given subpopulation and pair of crowding conditions. Black dots and lines represent the mean and 95% confidence interval of the estimated difference. (**B**) Same as (**A**) but for the multiplicative component, reflecting tuned divisive suppression.

Figure S1. Laminar recordings in V1.
(**A**) Section perspective schematic of the artificial dura recording chamber and laminar probe. (**B**) photo of the artificial dura chamber in one monkey. (**C**) Example receptive field contours vertically stacked for each recording site along the depth of each of the two shanks on the laminar probe. (**D**) Example current source density (CSD) map showing the early current sink (red) indicative of the input layer. (**E**) Data recorded from a single trial, including spikes (black vertical ticks) and local field potentials (black traces) collected from each channel on the two shanks. Data is arranged vertically based on the spatial location of the recording channel. Cortical layers are indicated on the left (green=superficial, gray=input, pink=deep). Vertical bars indicate the time of stimulus onsets.

Figure S2. Peri-stimulus time histograms, tuning curves of single units, and orientation selectivity index under crowding.
(**A**) Peri-stimulus time histogram (PSTH) of subpopulations in each condition following the onset of the preferred stimulus. The preferred orientation was calculated separately for each single unit. Spikes were counted in 30ms bins shifted by 5ms. Error bars show mean+/- s.e.m. (**B**) Example single unit tuning curves in the target-only condition. (**C**) Difference in orientation selectivity index (OSI) in each of the crowding conditions compared to the target-only condition. Colored half-violin plots show the bootstrapped estimation of the mean difference in OSI for a given subpopulation and crowding condition. Black dots and lines represent the mean and 95% confidence interval of the estimated difference.

Figure S3. Radial-out crowding decreases spike-count correlations in the feed-forward pathway
(**A**) Spike-count correlations from pairs of simultaneously recorded neurons in each subpopulation in the target-only condition and each of the three crowding conditions. In the top panels, error bars represent mean +/- 95% confidence intervals of the spike-count correlation in each condition. In the bottom panels, black dots and lines represent the mean and 95% confidence interval of the estimated difference from the target-only condition, and colored half-violin plots show the bootstrapped estimation of the mean difference in spike-count correlation for a given subpopulation and crowding condition. (**B**) Spike-count correlations for each pair of single units in each subpopulation. Each point represents the spike-count correlations of one pair of simultaneously recorded neurons in the same layer and cell class and each condition.

Figure S4. V1 neurons are not responsive to the distractor alone.
Normalized tuning curves for each subpopulation in the target-only and distractor-only conditions. Tuning curves are aligned with the preferred orientation at 0 degrees and duplicated to fill the circle. Tuning curves were normalized by the maximum firing rate in the target-only condition. The distractor-only condition is plotted relative to the orientation

of the distractor and includes data from all four distractor positions. Lines and shading represent mean +/- s.e.m. (black=target only, gold=distractor only).

Figure S5. Single-unit tuning model parameters in different crowding conditions
(**A**) Related to Figure 6A. Value of the fitted additive parameter for each neuron in each crowding condition. Each line represents the additive parameter for a single unit in across the two crowding conditions indicated at the bottom. (**B**) Related to Figure 6B. Same as (**A**) but for the multiplicative component.

MATERIALS & METHODS

Data Collection

Surgical procedures

Surgical procedures were similar to those described previously (*22, 23, 83*). We placed

low-profile titanium recording chambers in two rhesus macaques so that the chambers

allowed access to area V1 (right hemisphere in monkey D, left hemisphere in monkey M).

Chambers were targeted based on sulcus reconstructions created using preoperative

structural MRI. After chamber implantation, we removed the native dura mater and

replaced it with a transparent silicone artificial dura (AD). The AD allowed for the

visualization of cortical sites in V1 for probe targeting. All procedures were approved by

the Yale University Institutional Animal Care and Use Committee and conformed to NIH

guidelines.

Electrophysiology

Prior to a series of recordings, we electroplated (nanoZ, White Matter LLC) 64-channel

electrode arrays ("laminar probes," NeuroNexus Technologies, Inc., 2 shanks, 32

channels/shank, 70μm spacing between sites, 200μm between shanks) with PEDOT

(poly(3,4-ethylene dioxythiophene)). At the beginning of each recording session, we

inserted a laminar probe in V1. Laminar probes were attached to a titanium mounting

stage that was screwed into the chamber. We positioned the laminar probes using an

electronic micromanipulator (Narishige Inc.) and ensured that the probes were orthogonal

to the surface of the cortex by visual inspection through a surgical microscope (Leica

Microsystems). To position the probe within the brain, we first penetrated the AD,

arachnoid, and pia by moving the probe downward at a high speed (>100µm/s). After the

tip of the probe entered the cortex, we inserted the remainder of the probe at a slow speed

(2µm/s). Once the entire probe was in the brain, we slowly (2µm/s) relieved the pressure

on the brain by retracting the probe upward, relieving pressure on the brain but not moving

the probe relative to the cortex.

Electrical signals from the laminar probe were collected at 30kHz and digitized on

a 64-channel digital headstage and sent to the recording system (RHD Recording

System, Intan Technologies). Action potential waveforms were extracted offline using

Kilosort2 (*84, 85*) and manually sorted into single and multi-unit clusters (Phy; *84, 85*).

Multi-unit clusters were excluded from further analysis. Single-unit clusters were classified

based on their trough-to-peak waveform duration into broad- and narrow-spiking units

(*22, 24*). Neurons with waveform durations less than 350µs were classified as narrow-

spiking while those with waveform durations longer than 350µs were classified as broad-

spiking. Clusters with peaks preceding the trough were identified as axonal spikes and

excluded. Recordings were collected over the course of 22 sessions (9 in monkey D, 13

in monkey M) with hundreds of single units recorded in each subject (384 in monkey D,

521 in monkey M).

Behavioral Control and Eye Tracking

We controlled behavioral experiments using NIMH MonkeyLogic (*86*). Eye position and

pupil diameter were sampled at 120Hz (ETL-200, ISCAN Inc.) and sent to the behavioral

control system. Stimuli were presented on a monitor positioned 57cm from the monkey

with a 60Hz refresh rate. Trials were aborted if the eye position deviated more than 1.2-

1.5 degrees of visual angle (dva; 1.2 for monkey D, 1.5 for monkey M) from the central

fixation point.

Receptive Field Mapping

We mapped RFs of the column by presenting Gabor patch stimuli (2-4 cycles/degree,

0.25-1 degree Gaussian half-width, 100% luminance contrast) on a square grid spanning

the lower visual quadrant of interest (left in monkey D, right in monkey M) while the

monkey maintained fixation on the center of the screen. Grid spacing parameters were

optimized each session and ranged from 0.25-1 dva. A stimulus was presented at a

random location and orientation on the grid during each frame. We calculated the LFP

power for each recording channel 40-200ms after stimulus presentation in each location.

The LFP power at each location was smoothed using a Gaussian kernel (sigma = 0.75

dva), and the peak location averaged across all recording sites was defined as the RF

center. Spatial RF maps for each channel were plotted as stacked contours for each

shank for visualization.

Current Source Density Mapping

We used CSD mapping (*33*) to identify laminar boundaries in our recordings. While

monkeys maintained fixation on the screen, 100% luminance contrast white annular

stimuli were flashed for 32ms, positioned at the center of the RF. The CSD was calculated

as the second spatial derivative of the LFP following stimulus onset. CSD traces were

spatially smoothed using a Gaussian kernel (sigma = 140µm). The input layer was

identified by the boundaries of the early current sink characterizing feed-forward input into

layer IV. Channels above and below this sink were classified as superficial and deep

respectively.

Anisotropic Crowding Task

While monkeys maintained fixation, stimulus arrays were flashed on the screen for 100ms

and off the screen for 200ms. The arrays consisted of a target stimulus in the RF center

either in isolation (target only) or together with a distractor stimulus (crowding). There

were four different conditions (target only, tangential crowding, radial-in crowding, and

radial-out crowding) of stimulus arrays presented during a trial. This was repeated 4-6

times during each trial depending on the monkey's ability to hold fixation. The stimulus

conditions were randomly interleaved from flash to flash. The center of the target stimulus

was aligned with the center of the average of the RFs of each recording site. The four

distractor locations were positioned on the radial or tangential axes. The radial axis was

defined as the axis connecting the target (and RF) center and the fixation point at the

center of the monitor. The tangential axis was defined as the line orthogonal to the radial

axis and passing through the target center. The tangential crowding condition occurred

when the target was presented along with a distractor stimulus on the tangential axis,

either in the clockwise or counterclockwise direction. The radial-in crowding condition

occurred when the target was presented along with a distractor stimulus on the radial axis

and between the target center and the fixation point. The radial-out condition was similar

to the radial-in condition, except the distractor stimulus was placed on the radial axis

further from the fixation point than the target. Target-distractor spacing was identical

across stimulus conditions.

The target-only condition was presented on 10% of flashes. On 80% of flashes,

crowding stimuli were presented. In the crowding condition, the distractor stimulus was

positioned at a tangential, radial-in, or radial-out location with equal probability. If the

distractor was presented tangential to the target, it was positioned clockwise or

counterclockwise to the target with equal probability. In the remaining 10% of trials, a

stimulus was presented exclusively at one of the four distractor locations. Target stimuli

were sine Gabor patches (25% luminance contrast, 3.5 cycles/degree, 0.5-1.0 degree

Gaussian half-width) presented at 6 evenly spaced orientations and 2 opposing phases.

Flanking stimuli were presented at 100% luminance contrast but were otherwise identical

to the target stimuli. There was a 0.1-0.2 degree gap between the edges of the target and

flanking stimuli (edge defined as 2 standard deviations from the center of the Gaussian).

In the crowding condition, distractors were presented at the same orientation as the target

or orthogonal to the target with equal probability. In this study, we focused exclusively on

the condition in which the distractor was orthogonal to the target. When distractors were

presented alone, they were shown with the same orientation distribution as the targets.

Data Analysis

Wherever possible, we analyzed our data using the estimation statistics framework (*87,*

88) and show bootstrapped estimations of differences between conditions rather than

relying on null-hypothesis testing. The estimation statistics framework provides a

principled way to measure effect sizes coupled with estimates of uncertainty, yielding

interval estimates of uncertainty.

Tuning Curve Visualization

To visualize the modulation of tuning curves of each subpopulation by distractor stimuli,

we calculated the number of spikes for each neuron 0-300ms following stimulus onset in

each orientation and condition, collapsing across phases, and subtracted the baseline

firing rate (0-100ms before stimulus onset). For each neuron, we identified the orientation

with the highest firing rate in the target-only condition as the preferred orientation and

shifted the tuning curves to set this orientation to 0 degrees in each condition. We then

normalized the turning curves by dividing each neuron's response to each orientation and

condition by the firing rate for the preferred target-only orientation. For visualization, the

tuning curve was duplicated and rotated 180 degrees to completely fit around a circle.

We excluded neurons that did not respond in the target-only condition. To visualize the

tuning curves of individual single units, we followed the same procedure but did not

average across phases, instead plotting the response to each phase as the opposite

orientation. Additionally, single-unit tuning curves were not rotated based on the preferred

orientation.

Preferred and Orthogonal Firing Rate and OSI

OSI was calculated using the following equation:

500
$$OSI = \frac{FR_{pref} - FR_{orth}}{FR_{pref} + FR_{orth}}$$

Where FR_{pref} is the firing rate in response to the preferred orientation and FR_{orth} is the

firing rate in response to the stimulus orthogonal to the preferred orientation. Firing rates

were calculated 0-300ms following stimulus onset. The preferred orientation was

identified as the orientation with the highest firing rate. FR_{pref}, FR_{orth} , and OSI were

compared between the target-only condition and the crowding conditions by calculating

the bootstrapped estimation of the mean difference.

Modeling Tuning Curve Modulation

We investigated the components of single-unit tuning curve modulation using a previously

published approach to identify additive and multiplicative changes to tuning curves (*17*).

Briefly, we fit a Poisson model based on spike counts from 0-300ms after stimulus onset

in the target-only condition for each orientation. We then calculated the untuned additive

and tuned multiplicative modulation of the fitted target-only tuning curve that would best

account for the observed spike counts independently for each crowding condition (*17*).

We then compared the additive and multiplicative parameters for each neuron across

conditions by calculating a bootstrapped estimation of the paired mean difference of the

components.

Decoding Analysis

We trained a decoder using linear discriminant analysis (LDA) to determine the identity

(orientation and phase) of a target stimulus from simultaneously recorded single-unit

activity. Spike counts were calculated from 0-300ms following stimulus onset. The

decoder performance was determined using 20-fold cross-validation. The decoder was

trained on stimuli presented in the target-only condition and tested on neural responses

in each of the crowding conditions. This procedure was repeated for each recording

session. Performance in the tangential crowding condition was calculated separately for

the two tangential locations and averaged.

To calculate decoding performance at the subpopulation level, we first generated

a database of pseudo-population activity for each neuron in every combination of

conditions and target orientations. Spike counts were calculated from 0-300ms following

stimulus onset. To train the decoder for each subpopulation, we generated pseudo-

population activity by randomly drawing samples from this database, independently for

each neuron. We trained separate decoders for each subpopulation, training and testing

on responses from every neuron in each cell class and layer across all recording

sessions. Decoders were trained by sampling from data in the target-only condition. 20%

of each neuron's responses to each orientation were held out (the subset was

independent for each neuron) for cross-validation. This procedure was repeated 20 times

with a different random subset of responses used for cross-validation in the target-only

condition. The decoding performance on this 20% of left-out data was considered the

decoding performance in the target-only condition. We then tested each decoder on

pseudo-population activity generated from responses in the crowding conditions.

Peri-stimulus time histograms (PSTH)

PSTHs were generated based on spike counts in 30ms bins shifted by 5ms. PSTHs are

shown only for the preferred orientation as defined previously. The PSTH for each neuron

was calculated separately for each condition, and the average PSTH for each

subpopulation was plotted in Figure S2A. All times are relative to stimulus onset.

Spike-count correlations

Spike-count correlations were calculated for each pair of simultaneously recorded

neurons in the same layer and broad- or narrow-spiking cell class. Spike counts were

extracted from 100-300ms after stimulus onset and normalized by Z-score for each

stimulus orientation and condition. For each condition, the spike-count correlation was

calculated separately for each orientation from the normalized spike counts and was then

averaged across orientations. Thus, each pair of neurons had a single spike-count

correlation number for each stimulus condition.

REFERENCES

1. H. Bouma, Interaction Effects in Parafoveal Letter Recognition. *Nature* **226**, 177-178 (1970).
2. D. V. Meegan, S. P. Tipper, Reaching into cluttered visual environments: spatial and temporal influences of distracting objects. *Q J Exp Psychol A* **51**, 225-249 (1998).
3. D. Whitney, D. M. Levi, Visual crowding: a fundamental limit on conscious perception and object recognition. *Trends in Cognitive Sciences* **15**, 160-168 (2011).
4. A. Toet, D. M. Levi, The two-dimensional shape of spatial interaction zones in the parafovea. *Vision Res* **32**, 1349-1357 (1992).
5. Y. Petrov, A. V. Popple, S. P. McKee, Crowding and surround suppression: Not to be confused. *Journal of Vision* **7**, 12-12 (2007).
6. M. C. Flom, G. G. Heath, E. Takahashi, CONTOUR INTERACTION AND VISUAL RESOLUTION: CONTRALATERAL EFFECTS. *Science* **142**, 979-980 (1963).
7. S. P. Tripathy, D. M. Levi, Long-range dichoptic interactions in the human visual cortex in the region corresponding to the blind spot. *Vision Res* **34**, 1127-1138 (1994).
8. A. S. Nandy, B. S. Tjan, Saccade-confounded image statistics explain visual crowding. *Nature Neuroscience* **15**, 463-469 (2012).
9. D. G. Pelli, Crowding: a cortical constraint on object recognition. *Curr Opin Neurobiol* **18**, 445-451 (2008).
10. B. C. Motter, D. A. Simoni, The roles of cortical image separation and size in active visual search performance. *Journal of Vision* **7**, 6-6 (2007).
11. Y. Petrov, A. V. Popple, Crowding is directed to the fovea and preserves only feature contrast. *Journal of Vision* **7**, 8-8 (2007).
12. R. van den Berg, J. B. T. M. Roerdink, F. W. Cornelissen, A Neurophysiologically Plausible Population Code Model for Feature Integration Explains Visual Crowding. *PLOS Computational Biology* **6**, e1000646 (2010).
13. M. Manassi, D. Whitney, Multi-level Crowding and the Paradox of Object Recognition in Clutter. *Curr Biol* **28**, R127-r133 (2018).
14. D. G. Pelli, K. A. Tillman, The uncrowded window of object recognition. *Nat Neurosci* **11**, 1129-1135 (2008).
15. E. A. Crowder, C. R. Olson, Macaque monkeys experience visual crowding. *J Vis* **15**, 14 (2015).
16. C. A. Henry, A. Kohn, Spatial contextual effects in primary visual cortex limit feature representation under crowding. *Nature Communications* **11**, 1687 (2020).
17. C. A. Henry, A. Kohn, Feature representation under crowding in macaque V1 and V4 neuronal populations. *Current Biology* **32**, 5126-5137.e5123 (2022).
18. R. J. Douglas, K. A. C. Martin, NEURONAL CIRCUITS OF THE NEOCORTEX. *Annual Review of Neuroscience* **27**, 419-451 (2004).

596   19.   B. Tasic *et al.*, Shared and distinct transcriptomic cell types across neocortical
597   areas. *Nature* **563**, 72-78 (2018).
598   20.   H. Zeng, J. R. Sanes, Neuronal cell-type classification: challenges, opportunities
599   and the path forward. *Nature Reviews Neuroscience* **18**, 530-546 (2017).
600   21.   M. Migliore, G. M. Shepherd, An integrated approach to classifying neuronal
601   phenotypes. *Nature Reviews Neuroscience* **6**, 810-818 (2005).
602   22.   A. S. Nandy, J. J. Nassi, J. H. Reynolds, Laminar Organization of Attentional
603   Modulation in Macaque Visual Area V4. *Neuron* **93**, 235-246 (2017).
604   23.   J. J. Nassi, M. C. Avery, A. H. Cetin, A. W. Roe, J. H. Reynolds, Optogenetic
605   Activation of Normalization in Alert Macaque Visual Cortex. *Neuron* **86**, 1504-1517
606   (2015).
607   24.   J. F. Mitchell, K. A. Sundberg, J. H. Reynolds, Differential attention-dependent
608   response modulation across cell classes in macaque visual area V4. *Neuron* **55**,
609   131-141 (2007).
610   25.   B. W. Connors, M. J. Gutnick, Intrinsic firing patterns of diverse neocortical
611   neurons. *Trends Neurosci* **13**, 99-104 (1990).
612   26.   D. A. McCormick, B. W. Connors, J. W. Lighthall, D. A. Prince, Comparative
613   electrophysiology of pyramidal and sparsely spiny stellate neurons of the
614   neocortex. *J Neurophysiol* **54**, 782-806 (1985).
615   27.   Y. Kawaguchi, Groupings of nonpyramidal and pyramidal cells with specific
616   physiological and morphological characteristics in rat frontal cortex. *Journal of*
617   *Neurophysiology* **69**, 416-431 (1993).
618   28.   L. G. Nowak, R. Azouz, M. V. Sanchez-Vives, C. M. Gray, D. A. McCormick,
619   Electrophysiological classes of cat primary visual cortical neurons in vivo as
620   revealed by quantitative analyses. *J Neurophysiol* **89**, 1541-1566 (2003).
621   29.   D. Contreras, L. Palmer, Response to contrast of electrophysiologically defined
622   cell classes in primary visual cortex. *The Journal of neuroscience : the official*
623   *journal of the Society for Neuroscience* **23**, 6936-6945 (2003).
624   30.   R. C. Foehring, N. M. Lorenzon, P. Herron, C. J. Wilson, Correlation of
625   physiologically and morphologically identified neuronal types in human association
626   cortex in vitro. *J Neurophysiol* **66**, 1825-1837 (1991).
627   31.   N. V. Povysheva *et al.*, Properties of excitatory synaptic responses in fast-spiking
628   interneurons and pyramidal cells from monkey and rat prefrontal cortex. *Cerebral*
629   *cortex (New York, N.Y. : 1991)* **16**, 541-552 (2006).
630   32.   A. Hasenstaub *et al.*, Inhibitory postsynaptic potentials carry synchronized
631   frequency information in active cortical networks. *Neuron* **47**, 423-435 (2005).
632   33.   U. Mitzdorf, Current source-density method and application in cat cerebral cortex:
633   investigation of evoked potentials and EEG phenomena. *Physiol Rev* **65**, 37-100
634   (1985).
635   34.   A. Angelucci *et al.*, Circuits and Mechanisms for Surround Modulation in Visual
636   Cortex. *Annu Rev Neurosci* **40**, 425-451 (2017).
637   35.   J. R. Cavanaugh, W. Bair, J. A. Movshon, Nature and interaction of signals from
638   the receptive field center and surround in macaque V1 neurons. *J Neurophysiol*
639   **88**, 2530-2546 (2002).

640   36.   D. H. Hubel, T. N. Wiesel, RECEPTIVE FIELDS AND FUNCTIONAL
641   ARCHITECTURE IN TWO NONSTRIATE VISUAL AREAS (18 AND 19) OF THE
642   CAT. *J Neurophysiol* **28**, 229-289 (1965).
643   37.   C. Blakemore, E. A. Tobin, Lateral inhibition between orientation detectors in the
644   cat's visual cortex. *Experimental brain research* **15**, 439-440 (1972).
645   38.   L. Maffei, A. Fiorentini, The unresponsive regions of visual cortical receptive fields.
646   *Vision Research* **16**, 1131-IN1135 (1976).
647   39.   C. D. Gilbert, Laminar differences in receptive field properties of cells in cat primary
648   visual cortex. *J Physiol* **268**, 391-421 (1977).
649   40.   J. I. Nelson, B. J. Frost, Orientation-selective inhibition from beyond the classic
650   visual receptive field. *Brain Res* **139**, 359-365 (1978).
651   41.   M. P. Sceniak, M. J. Hawken, R. Shapley, Visual spatial characterization of
652   macaque V1 neurons. *J Neurophysiol* **85**, 1873-1887 (2001).
653   42.   G. Van den Bergh, B. Zhang, L. Arckens, Y. M. Chino, Receptive-field properties
654   of V1 and V2 neurons in mice and macaque monkeys. *The Journal of comparative
655   neurology* **518**, 2051-2070 (2010).
656   43.   S. Shushruth, J. M. Ichida, J. B. Levitt, A. Angelucci, Comparison of Spatial
657   Summation Properties of Neurons in Macaque V1 and V2. *Journal of
658   Neurophysiology* **102**, 2069-2083 (2009).
659   44.   M. K. Kapadia, M. Ito, C. D. Gilbert, G. Westheimer, Improvement in visual
660   sensitivity by changes in local context: parallel studies in human observers and in
661   V1 of alert monkeys. *Neuron* **15**, 843-856 (1995).
662   45.   M. K. Kapadia, G. Westheimer, C. D. Gilbert, Dynamics of spatial summation in
663   primary visual cortex of alert monkeys. *Proc Natl Acad Sci U S A* **96**, 12073-12078
664   (1999).
665   46.   M. K. Kapadia, G. Westheimer, C. D. Gilbert, Spatial distribution of contextual
666   interactions in primary visual cortex and in visual perception. *J Neurophysiol* **84**,
667   2048-2062 (2000).
668   47.   W. H. Bosking, Y. Zhang, B. Schofield, D. Fitzpatrick, Orientation selectivity and
669   the arrangement of horizontal connections in tree shrew striate cortex. *The Journal
670   of neuroscience : the official journal of the Society for Neuroscience* **17**, 2112-2127
671   (1997).
672   48.   K. E. Schmidt, R. Goebel, S. Löwel, W. Singer, The Perceptual Grouping Criterion
673   of Colinearity is Reflected by Anisotropies of Connections in the Primary Visual
674   Cortex. *European Journal of Neuroscience* **9**, 1083-1089 (1997).
675   49.   F. W. Campbell, J. J. Kulikowski, J. Levinson, The effect of orientation on the visual
676   resolution of gratings. *J Physiol* **187**, 427-436 (1966).
677   50.   J. A. Bauer, Jr., D. A. Owens, J. Thomas, R. Held, Monkeys show an oblique effect.
678   *Perception* **8**, 247-253 (1979).
679   51.   G. Westheimer, The distribution of preferred orientations in the peripheral visual
680   field. *Vision Res* **43**, 53-57 (2003).
681   52.   L. Liang, Y. Zhou, M. Zhang, Y. Pan, Revealing the Radial Effect on Orientation
682   Discrimination by Manual Reaction Time. *Front Neurosci* **11**, 638 (2017).
683   53.   M. M. Himmelberg, J. Winawer, M. Carrasco, Polar angle asymmetries in visual
684   perception and neural architecture. *Trends Neurosci* **46**, 445-458 (2023).

54. J. Rovamo, V. Virsu, Isotropy of cortical magnification and topography of striate
cortex. *Vision Res* **24**, 283-286 (1984).
55. R. Millin, A. C. Arman, S. T. L. Chung, B. S. Tjan, Visual Crowding in V1. *Cerebral*
Cortex **24**, 3107-3115 (2013).
56. M. Kwon, P. Bao, R. Millin, B. S. Tjan, Radial-tangential anisotropy of crowding in
the early visual areas. *J Neurophysiol* **112**, 2413-2422 (2014).
57. W. R. Levick, L. N. Thibos, Analysis of orientation bias in cat retina. *J Physiol* **329**,
243-261 (1982).
58. A. G. Leventhal, J. D. Schall, Structural basis of orientation sensitivity of cat retinal
ganglion cells. *The Journal of comparative neurology* **220**, 465-475 (1983).
59. J. D. Schall, V. H. Perry, A. G. Leventhal, Retinal ganglion cell dendritic fields in
old-world monkeys are oriented radially. *Brain Res* **368**, 18-23 (1986).
60. R. W. Rodieck, K. F. Binmoeller, J. Dineen, Parasol and midget ganglion cells of
the human retina. *The Journal of comparative neurology* **233**, 115-132 (1985).
61. E. L. Smith, 3rd, Y. M. Chino, W. H. Ridder, 3rd, K. Kitagawa, A. Langston,
Orientation bias of neurons in the lateral geniculate nucleus of macaque monkeys.
Vis Neurosci **5**, 525-545 (1990).
62. V. B. Mountcastle, The columnar organization of the neocortex. *Brain* **120 (Pt 4)**,
701-722 (1997).
63. R. Gattass, A. Sousa, C. Gross, Visuotopic organization and extent of V3 and V4
of the macaque. *Journal of neuroscience* **8**, 1831-1845 (1988).
64. K. D. Harris, T. D. Mrsic-Flogel, Cortical connectivity and sensory coding. *Nature*
503, 51-58 (2013).
65. R. Bauer, B. M. Dow, R. G. Vautin, Laminar distribution of preferred orientations in
foveal striate cortex of the monkey. *Experimental brain research* **41**, 54-60 (1980).
66. R. Bauer, B. M. Dow, A. Z. Snyder, R. Vautin, Orientation shift between upper and
lower layers in monkey visual cortex. *Experimental brain research* **50**, 133-145
(1983).
67. K. S. Rockland, J. S. Lund, Intrinsic laminar lattice connections in primate visual
cortex. *The Journal of comparative neurology* **216**, 303-318 (1983).
68. N. J. Priebe, Mechanisms of Orientation Selectivity in the Primary Visual Cortex.
Annu Rev Vis Sci **2**, 85-107 (2016).
69. K. A. Ferguson, J. A. Cardin, Mechanisms underlying gain modulation in the
cortex. *Nature reviews. Neuroscience* **21**, 80-92 (2020).
70. R. J. Douglas, C. Koch, M. Mahowald, K. A. Martin, H. H. Suarez, Recurrent
excitation in neocortical circuits. *Science* **269**, 981-985 (1995).
71. H. K. Hartline. (1949).
72. T. C. Freeman, S. Durand, D. C. Kiper, M. Carandini, Suppression without
inhibition in visual cortex. *Neuron* **35**, 759-771 (2002).
73. R. Ben-Yishai, R. L. Bar-Or, H. Sompolinsky, Theory of orientation tuning in visual
cortex. *Proc Natl Acad Sci U S A* **92**, 3844-3848 (1995).
74. D. C. Somers, S. B. Nelson, M. Sur, An emergent model of orientation selectivity
in cat visual cortical simple cells. *The Journal of neuroscience : the official journal*
of the Society for Neuroscience **15**, 5448-5465 (1995).
75. C. van Vreeswijk, H. Sompolinsky, Chaos in neuronal networks with balanced
excitatory and inhibitory activity. *Science* **274**, 1724-1726 (1996).

76. C. van Vreeswijk, H. Sompolinsky, Chaotic balanced state in a model of cortical
circuits. *Neural Comput* **10**, 1321-1371 (1998).
77. D. Hansel, C. van Vreeswijk, The mechanism of orientation selectivity in primary
visual cortex without a functional map. *The Journal of neuroscience : the official
journal of the Society for Neuroscience* **32**, 4049-4064 (2012).
78. T. W. Troyer, A. E. Krukowski, N. J. Priebe, K. D. Miller, Contrast-invariant
orientation tuning in cat visual cortex: thalamocortical input tuning and correlation-
based intracortical connectivity. *The Journal of neuroscience : the official journal
of the Society for Neuroscience* **18**, 5908-5927 (1998).
79. T. W. Troyer, A. E. Krukowski, K. D. Miller, LGN input to simple cells and contrast-
invariant orientation tuning: an analysis. *J Neurophysiol* **87**, 2741-2752 (2002).
80. G. C. DeAngelis, J. G. Robson, I. Ohzawa, R. D. Freeman, Organization of
suppression in receptive fields of neurons in cat visual cortex. *J Neurophysiol* **68**,
144-163 (1992).
81. D. J. Heeger, Normalization of cell responses in cat striate cortex. *Vis Neurosci* **9**,
181-197 (1992).
82. M. Carandini, D. L. Ringach, Predictions of a recurrent model of orientation
selectivity. *Vision Res* **37**, 3061-3071 (1997).
83. O. Ruiz *et al.*, Optogenetics through windows on the brain in the nonhuman
primate. *J Neurophysiol* **110**, 1455-1467 (2013).
84. C. Stringer *et al.*, Spontaneous behaviors drive multidimensional, brainwide
activity. *Science* **364**, eaav7893 (2019).
85. N. A. Steinmetz *et al.*, Neuropixels 2.0: A miniaturized high-density probe for
stable, long-term brain recordings. *Science* **372**, eabf4588 (2021).
86. J. Hwang, A. R. Mitz, E. A. Murray, NIMH MonkeyLogic: Behavioral control and
data acquisition in MATLAB. *Journal of Neuroscience Methods* **323**, 13-21 (2019).
87. R. J. Calin-Jageman, G. Cumming, Estimation for Better Inference in
Neuroscience. *eneuro* **6**, ENEURO.0205-0219.2019 (2019).
88. J. Ho, T. Tumkaya, S. Aryal, H. Choi, A. Claridge-Chang, Moving beyond P values:
data analysis with estimation graphics. *Nature Methods* **16**, 565-566 (2019).
761

Figure 1

A

Figure 1. Anisotropic contextual interaction task and single-unit recordings.
(**A**) Anisotropic contextual interaction task schematic. Stimuli were presented alone at the receptive field center (target only) or with a distractor at one of the four crowding locations. The radial axis connects the target to the center of gaze, and the tangential axis is orthogonal to it, also passing through the target. The top schematic shows stimuli at all possible locations, while the bottom section shows the possible spatial configurations for the presentation of an individual stimulus array. (**B**) Mean waveforms for all recorded single-units (blue=narrow-spiking, orange=broad-spiking). (**C**) Distribution of waveform durations (trough-to-peak time) for all recorded single-units across the two monkeys. Neurons with waveform durations less than 350μs were classified as narrow-spiking (blue), and neurons with waveform durations greater than 350μs were classified as broad-spiking (orange).

Figure 2

Figure 2. Distractor stimuli impair decoding performance in simultaneously recorded populations in an anisotropic manner.

Example decoding performance from 4 simultaneously recorded populations (two in each monkey) in the target-only condition and each of the three crowding conditions. Points in the upper section of each plot show the decoding performance for each of the 20 different cross-validations. The lower section for each session shows the bootstrapped estimation of the difference between decoding performance in each crowding condition compared to the target only condition. Half-violin plots show bootstrapped distribution of the difference, and black dots and bars represent the mean and 95% confidence intervals of the difference in decoding performance.

Figure 3

Figure 3. Radial-out crowding most strongly impairs decoding in the feed-forward pathway.
Decoding performance from each subpopulation in the target-only condition and each of the three crowding conditions. Points in the upper section of each plot show the decoding performance for each of the 20 different cross-validations. The lower section for each session shows the bootstrapped estimation of the difference between decoding performance in each crowding condition compared to the target only condition. Half-violin plots show bootstrapped distribution of the difference, and black dots and bars represent the mean and 95% confidence intervals of the difference in decoding performance.

Figure 4

Figure 4. Crowding stimuli modulate the tuning curves of V1 neurons.
Normalized tuning curves for each subpopulation in the target-only and crowding conditions. Tuning curves are aligned with the preferred orientation at 0 degrees and duplicated to fill the circle. Tuning curves were normalized by the maximum firing rate in the target-only condition. Lines and shading represent mean +/- s.e.m. (black=target only, pink=tangential crowding, green=radial-in crowding, purple=radial-out crowding).

Figure 5

Figure 5. Crowding broadens tuning curves

Difference in firing rate in response to preferred (dark brown) and orthogonal (light tan) stimuli under each crowding condition in the six subpopulations. Colored half-violin plots show the bootstrapped estimation of the paired mean difference in firing rate for a given subpopulation, crowding condition, and stimulus orientation. Black dots and lines represent the mean and 95% confidence interval of the estimated difference.

Figure 6

A
Additive component

B
Multiplicative component

Figure 6. Radial-out crowding causes stronger additive facilitation and divisive suppression of tuning curves
(**A**) Difference in the modeled (see Methods) untuned additive facilitation caused by each of the crowding conditions. Gray half-violin plots show the bootstrapped estimation of the paired mean difference in the modeled additive component for a given subpopulation and pair of crowding conditions. Black dots and lines represent the mean and 95% confidence interval of the estimated difference. (**B**) Same as (**A**) but for the multiplicative component, reflecting tuned divisive suppression.

Figure S1

Figure S1. Laminar recordings in V1.
(**A**) Section perspective schematic of the artificial dura recording chamber and laminar probe. (**B**) photo of the artificial dura chamber in one monkey. (**C**) Example receptive field contours vertically stacked for each recording site along the depth of each of the two shanks on the laminar probe. (**D**) Example current source density (CSD) map showing the early current sink (red) indicative of the input layer. (**E**) Data recorded from a single trial, including spikes (black vertical ticks) and local field potentials (black traces) collected from each channel on the two shanks. Data is arranged vertically based on the spatial location of the recording channel. Cortical layers are indicated on the left (green=superficial, gray=input, pink=deep). Vertical bars indicate the time of stimulus onsets.

Figure S2

Figure S2. Peri-stimulus time histograms, tuning curves of single units, and orientation selectivity index under crowding.

(**A**) Peri-stimulus time histogram (PSTH) of subpopulations in each condition following the onset of the preferred stimulus. The preferred orientation was calculated separately for each single unit. Spikes were counted in 30ms bins shifted by 5ms. Error bars show mean+/- s.e.m. (**B**) Example single unit tuning curves in the target-only condition. (**C**) Difference in orientation selectivity index (OSI) in each of the crowding conditions compared to the target-only condition. Colored half-violin plots show the bootstrapped estimation of the mean difference in OSI for a given subpopulation and crowding condition. Black dots and lines represent the mean and 95% confidence interval of the estimated difference.

Figure S3

Figure S3. Radial-out crowding decreases spike-count correlations in the feed-forward pathway
(**A**) Spike-count correlations from pairs of simultaneously recorded neurons in each subpopulation in the target-only condition and each of the three crowding conditions. In the top panels, error bars represent mean +/- 95% confidence intervals of the spike-count correlation in each condition. In the bottom panels, black dots and lines represent the mean and 95% confidence interval of the estimated difference from the target only condition, and colored half-violin plots show the bootstrapped estimation of the mean difference in spike-count correlation for a given subpopulation and crowding condition. (**B**) Spike-count correlations for each pair of single-units in each subpopulation. Each point represents the spike-count correlations of one pair of simultaneously recorded neurons in the same layer and cell class and in each condition.

Figure S4. V1 neurons are not responsive to the distractor alone.
Normalized tuning curves for each subpopulation in the target-only and distractor-only conditions. Tuning curves are aligned with the preferred orientation at 0 degrees and duplicated to fill the circle. Tuning curves were normalized by the maximum firing rate in the target-only condition. The distractor-only condition is plotted relative to the orientation of the distractor, and includes data from all four distractor positions. Lines and shading represent mean +/- s.e.m. (black=target only, gold=distractor only).

Figure S5

Figure S5. Single-unit tuning model parameters in different crowding conditions
(**A**) Related to Figure 6A. Value of the model fit additive parameter for each neuron in each crowding condition. Each line represents the additive parameter for a single unit in across the two crowding conditions indicated at the bottom. (**B**) Related to Figure 6B. Same as (**A**) but for the multiplicative component.