An anciently diverged family of RNA binding proteins maintain correct splicing of ultra-long exons through cryptic splice site repression.

Chileleko Siachisumo[1]*, Sara Luzzi[1]*, Saad Aldalaqan[1]*, Gerald Hysenaj[1], Caroline Dalgliesh[1], Kathleen Cheung[2], Matthew R Gazzara[3], Ivaylo D Yonchev[4], Katherine James[5], Mahsa Kheirollahi Chadegani[1], Ingrid Ehrmann[1], Graham R Smith[2], Simon J Cockell[2], Jennifer Munkley[1], Stuart A Wilson[4], Yoseph Barash[3] and David J Elliott[1]†

* These authors contributed equally to the paper as first authors
[1] Biosciences Institute, Faculty of Medical Sciences, Newcastle University, Newcastle, United Kingdom
[2] Bioinformatics Support Unit, Faculty of Medical Sciences, Newcastle University, Newcastle, United Kingdom.
[3] Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, United States.
[4] School of Biosciences, University of Sheffield, Sheffield, United Kingdom
[5] School of Computing, Newcastle University, Newcastle, United Kingdom
† To whom correspondence should be addressed. Email: david.elliott@newcastle.ac.uk

Abstract

We previously showed that the germ cell specific nuclear protein RBMXL2 represses cryptic splicing patterns during meiosis and is required for male fertility. It has remained unknown whether RBMXL2 evolved its role in splicing repression to deal with the transcriptionally permissive environment of meiosis or might fulfil a function required in all cells. RBMXL2 evolved from the X-linked RBMX gene, which is silenced during meiosis due to sex chromosome inactivation. Here we find that like RBMXL2, RBMX primarily operates as a splicing repressor in somatic cells, and specifically regulates a distinct class of exons that exceed the median human exon size. RBMX protein-RNA interactions are enriched within ultra-long exons, particularly within genes involved in genome stability, and RBMX represses the selection of cryptic splice sites that would compromise gene function. These similarities in overall function suggested that RBMXL2 during meiosis might replace the otherwise ubiquitous RBMX protein. To test this prediction we carried out inducible expression of RBMXL2 and the more distantly related RBMY protein in somatic cells, finding each could rescue aberrant patterns of RNA processing in response to RBMX depletion. The C-terminal disordered domain of RBMXL2 is sufficient to rescue proper splicing control after RBMX depletion. Our data indicate that RBMXL2 replaces RBMX during meiosis, and these proteins have maintained parallel roles that must have been conserved over at least 200 million years of mammalian evolution. We propose RBMX family proteins are important for the splicing inclusion of ultra-long exons because these are particularly susceptible to disruption by cryptic splice site selection.

Introduction.

Efficient gene expression in eukaryotes requires introns and exons to be correctly recognised by the spliceosome, the macromolecular machine that joins exons together. The spliceosome recognises short sequences that are present at exon-intron junctions within precursor mRNAs called splice sites. In higher organisms there is some flexibility in splice site recognition, as most genes produce multiple mRNAs by alternative splicing. However, aberrant "cryptic" splice sites that are weakly selected or totally ignored by the spliceosome occur frequently in the human genome and could function as decoys to interfere with gene expression (Aldalaqan et al., 2022; Sibley et al., 2016). Many cryptic splice sites are located amongst repetitive sequences within introns, where they are repressed by RNA binding proteins belonging to the hnRNP family (Attig et al., 2018). Cryptic splice sites can also be present within exons, particularly long exons where they can cause formation of exitrons (exon sequences that are removed as if they were introns) (Marquez et al., 2015).

The testis-specific nuclear RNA binding protein RBMXL2 was recently shown to repress cryptic splice site selection during meiosis, including within some ultra-long exons of genes involved in genome stability (I. Ehrmann et al., 2019). RBMXL2 is only expressed within the testis (Aldalaqan et al., 2022; I. Ehrmann et al., 2019), raising the question of how the cryptic splice sites controlled by RBMXL2 are repressed in other parts of the body. RBMXL2 is part of an anciently diverged family of RNA binding proteins. The *RBMXL2* gene evolved 65 million years ago following retrotransposition of the *RBMX* gene from the X chromosome to an autosome (I. Ehrmann et al., 2019). RBMX and RBMXL2 proteins (also known as hnRNP-G and hnRNP-GT) share 73% identity at the protein level and have the same modular structure comprising an N-terminal RNA Recognition Motif (RRM) and a C-terminal disordered region containing RGG repeats (Figure 1A). RBMX and RBMXL2 are also more distantly related to a gene on the long arm of the Y chromosome called *RBMY* that is deleted in some infertile men (with only ~37% identity between human RBMXL2 and RBMY) (Elliott et al., 1997; Ma et al., 1993). The role of RBMY in the germline is almost totally unknown, but RBMY protein has been implicated in splicing regulation (Elliott et al., 2000; Venables et al., 2000).

The X and Y chromosomes are inactivated during meiosis within a heterochromatic structure called the XY body (Turner, 2015; Wang, 2004). Meiosis is quite a long process, and to maintain cell viability during this extended period when the X chromosome is inactive, a number of autosomal retrogenes have evolved from essential X chromosome genes and are actively expressed during meiosis. However, it is unknown whether RBMXL2 and RBMX are functionally similar enough for RBMXL2 to provide a replacement function for RBMX during meiosis, or whether RBMXL2 evolved differently to control meiosis-specific patterns of expression. Suggesting somewhat different activities RBMX was recently shown to activate exon splicing inclusion, via a mechanism involving binding to RNA through

its C-terminal disordered domain facilitated by recognition of m6A residues and RNA polymerase II pausing (Liu et al., 2017; Zhou et al., 2019).

Here, we have used iCLIP and RNA-seq to analyse the binding characteristics and RNA processing targets of human RBMX. We identify a novel class of RBMX-dependent ultra-long exons connected to genome stability and transcriptional control, and find that RBMX, RBMXL2 and RBMY paralogs have closely related functional activity in repressing cryptic splice site selection. Our data reveal an ancient mechanism of gene expression control by RBMX family proteins that predates the radiation of mammals and provides a new understanding of how ultra-long exons are properly incorporated into mRNAs.

Results

RBMX primarily operates as a splicing repressor in somatic cells

We first set out to identify the spectrum of splicing events that are strongly controlled by RBMX across different human cell lines. We used RNA-seq from biological triplicate MDA-MB-231 cells treated with siRNA against RBMX (achieving >90% depletion, Figure 1B), followed by bioinformatics analysis using the SUPPA2 (Trincado et al., 2018) and MAJIQ (Vaquero-Garcia et al., 2016) splicing prediction tools. We identified 315 changes in RNA processing patterns in response to RBMX-depletion that were high enough amplitude to be visually confirmed on the IGV genome browser (Robinson et al., 2011) (Figure 1 – Figure supplement 1) (Figure 1 – Source Data 1). Analysis of these splicing events within existing RNA-seq data from HEK293 cells depleted for RBMX (GSE74085) (Liu et al., 2017) revealed 166 high amplitude events that are controlled by RBMX in both HEK293 and MDA-MB-231 cells (Figure 1C). We concentrated our downstream analysis on these splicing events (Figure 1 – Source Data 1). 92% of the splicing events regulated by RBMX in human somatic cells were already annotated on Ensembl, Gencode or Refseq (Figure 1D). Strikingly two thirds of these events are repressed by RBMX, meaning they were increasingly used in RBMX depleted cells compared to control, and include exon inclusion, alternative 5' and 3' splice sites, exitrons, and intron retention (Figure 1E). This means that RBMX primarily operates as a splicing repressor in human somatic cells.

Figure 1

Figure 1. RBMX primarily operates as a splicing repressor in human somatic cells. (A)
Schematic structure of RBMX family proteins (left side, cladogram) and amino acid similarity of
each domain between RBMX protein and two other members of this family, RBMXL2 and RBMY.
RRM, RNA recognition motif. CD, central domain important for recognition of nascent transcripts
and nuclear localisation. CTD, C-terminal domain, involved in RNA binding (Elliott et al., 2019). **(B)**
Western blot analysis shows efficient siRNA-mediated depletion of RBMX from MDA-MB-231 cells.
(C) Pie chart showing the percentages of events controlled by RBMX in both MDA-MB-231 (this
study) and HEK293 (Liu et al., 2017) cells. **(D)** Pie chart showing the percentages of events
controlled by RBMX in both MDA-MB-231 and HEK293 cells that have been previously annotated
(Refseq, Ensembl, Gencode), and those that are novel to this study. **(E)** Bar chart showing the
different types of alternative splicing events controlled by RBMX protein in both HEK293 and MDA-
MB-231 cells, summarising the proportion of splicing events that are activated by RBMX versus
those that are repressed.

Figure 1 - Figure Supplement 1

A

Figure 1 – Figure supplement 1 (A) Bar chart showing the different types of alternative splicing

events controlled by RBMX protein in MDA-MB-231 cells, summarising the proportion of splicing

events that are activated by RBMX versus those that are repressed.

Splicing control and sites of RBMX protein-RNA interaction are enriched within long internal exons

The above data indicated that RBMX has a major role in repressing cryptic splicing patterns in

human somatic cells. To further correlate splicing regulation to patterns of RBMX protein-RNA

interactions, we next mapped the distribution of RBMX-RNA binding sites in human somatic cells.

We engineered a stable human HEK293 cell line to express RBMX-FLAG fusion protein in

response to tetracycline addition. Western blotting showed that expression of RBMX-FLAG was

efficiently induced after tetracycline treatment. Importantly, levels of the induced RBMX-FLAG

protein were similar to those of endogenous RBMX (Figure 2A). We next used this inducible cell

line to carry out individual nucleotide resolution crosslinking and immunoprecipitation (iCLIP) – a

technique that produces a global picture of protein-RNA binding sites (Konig et al., 2011). After

crosslinking, RBMX-FLAG protein was immunoprecipitated, then infra-red labelled RNA-protein

adducts were isolated (Figure 2B) and subjected to library preparation. Following deep sequencing

of biological triplicate experiments, 5 to 10 million unique reads (referred to here as iCLIP tags,

representing sites of RBMX protein-RNA cross-linking) were aligned to the human genome. Each

individual iCLIP replicate showed at least 70% correlation with each of the others (Figure 2 –

Figure Supplement 1A). K-mer motif analysis revealed RBMX preferentially binds to AG-rich

sequences (Figure 2C and Figure 2 – Figure Supplement 1B).

In line with previous work on other RNA binding proteins (Van Nostrand et al., 2020), only 31% of

the RNA splicing events that are controlled by RBMX in both HEK293 cells and MDA-MB-231 cells

were identified by iCLIP as direct targets for RBMX binding (Figure 2 – Figure supplement 1C).

However, RBMX-responsive internal exons that did contain RBMX iCLIP tags were significantly

longer than the ones that are not bound by RBMX (Figure 2D). We therefore compared the length

of the exons regulated (identified by RNA-seq) and bound by RBMX (identified by iCLIP) within

protein-coding genes to all mRNA exons expressed in HEK293 (Liu et al., 2017). We reasoned that

larger exons might have a higher chance to be bound by RBMX merely because of their large size.

Because of this, we did not take into account the density of RBMX binding and considered all

exons that contained at least one iCLIP tag. Strikingly, we found that exons regulated and bound

by RBMX were significantly longer than the median size of HEK293 mRNA exons which is ~130 bp

(Figure 2E). This led us to test whether RBMX protein is preferentially associated with long exons.

For this we plotted the distribution of exons bound and regulated by RBMX together with all exons

expressed from HEK293 mRNA genes (Liu et al., 2017) (Figure 2 – Source Data 1). We found that

RBMX controls and binds two different classes of exons: the first have comparable length to the

average HEK293 exon, while the second were extremely long, exceeding 1000 bp in length (Figure

2F). We defined this second class as 'ultra-long exons', which represented the 20.8% of exons

regulated by RBMX and 15% of the ones that contained RBMX iCLIP tags. These proportions were

significantly enriched compared to the general abundance of ultra-long exons expressed from

HEK293 cells, which was only 3.9% (Figure 2G). Overall, this data revealed a function for RBMX in

the regulation of splicing of ultra-long exons.

Figure 2

Figure 2. Splicing control and sites of RBMX protein-RNA interaction are enriched within long internal exons. (A) Western blot showing levels of RBMX-FLAG protein, expressed after 24h treatment with tetracycline, compared to endogenous RBMX within HEK293 cells, both detected using α-RBMX antibody. α-GAPDH antibody was used as loading control. **(B)** RNAs cross-linked to RBMX-FLAG during iCLIP detected through the infrared adaptor (RBMX-RNA complexes). Lane 1, anti-FLAG pull-down from crosslinked HEK293 control cells not expressing RBMX-FLAG proteins, treated with 0.8 U/ml RNaseI. Lane 3, RBMX-FLAG pull-down crosslinked to RNA, treated with 2.5 U/ml RNaseI. Lanes 5-7, RBMX-FLAG pull-down crosslinked to RNA, treated with 0.8 U/ml RNaseI. Samples in lanes 5-7 were used for iCLIP library preparation. Lanes 2 and 4 are empty. **(C)** K-mer analysis shows the top 10 enriched motifs within sequences surrounding RBMX iCLIP tags. **(D)** Boxplot analysis shows sizes of exons containing splicing events regulated by RBMX, grouped by whether they contain CLIP tags or not. ****, p-value<0.0001 (Mann-Whitney test). **(E)** Boxplot analysis shows distribution of exon sizes relative to: exons contained in mRNA genes expressed in HEK293 cells (Liu et al., 2017); exons regulated by RBMX as identified by RNA-seq;

exons containing RBMX binding sites as identified by iCLIP, listed independently of iCLIP tag

density. Median sizes for each group are shown. ****, p-value<0.0001 (Wilcoxon rank test and

Kruskal-Wallis test). **(F)** Distribution plot of exon sizes for the groups shown in (E). Note the

increased accumulation of exons larger than 1000 bp (ultra-long exons) in RBMX bound and

regulated exons compared to all exons expressed in HEK293 (Liu et al., 2017). **(G)** Bar plot

indicating the proportion of ultra-long exons in the groups shown in (E, F). ****, p-value<0.0001

(Chi-squared test).

Figure 2 – Figure supplement 1

Figure 2 Figure Supplement 1. (A) Correlation analysis between three replicates for RBMX-FLAG

iCLIP. **(B)** K-mer analysis shows the top 50 enriched motifs within sequences surrounding RBMX

iCLIP binding. **(C)** Barplot showing percentage of exons regulated by RBMX that contain iCLIP

tags.

RBMX is important for proper splicing inclusion of full-length ultra-long exons within genes

involved in DNA repair and RNA polymerase II transcription

We next performed Gene Ontology analysis in order to determine whether RBMX regulates

splicing patterns for particular classes of genes. Both the genes bound by RBMX (detected using

iCLIP, Figure 3 – Figure supplement 1A and Figure 3 – Source Data 1) and regulated by RBMX in

both MDA-MB-231 and HEK293 cell lines (detected using RNA-seq, Figure 3 – Figure supplement 1B and Figure 3 – Source Data 1) each showed individual global enrichment in functions connected to genome stability and gene expression. Similarly, Gene Ontology analyses showed that genes that were both bound by and dependent on RBMX for correct splicing were enriched in pathways including transcription and gene expression, DNA replication and repair, and meiosis (Figure 3A and Figure 3 – Source Data 1). In comparison, Gene Ontology analyses of all ultra-long exons in the human genome revealed enrichment of gene expression but not genome stability pathways (Figure 3 – Figure supplement 1C and Figure 3 – Source Data 1), showing that regulation of ultra-long exons within genes involved in DNA metabolism and cell cycle is characteristic of RBMX. These data are consistent with published observations (Adamson et al., 2012; Munschauer et al., 2018; Zheng et al., 2020) that depletion of RBMX reduces genome stability. In addition, comet assays also detected increased levels of genome instability after RBMX depletion (Figure 3 - Figure Supplement 1D, E).

The above data indicated that RBMX-RNA binding interactions and splicing control by RBMX are particularly associated with long internal exons and enriched within classes of genes involved in genome stability. These exons included the 2.1 Kb exon 5 of the *ETAA1* (*Ewings Tumour Associated Antigen 1*) gene, where RBMX potently represses a cryptic 3' splice site that reduces the size of this exon from 2.1 Kb to 100 bp (Figure 3B and Figure 3 – Figure supplement 2A). RT-PCR analysis confirmed that RBMX depletion causes a much shorter version of *ETAA1* exon 5 to prevail, particularly in MDA-MB-231 and NCI-H520 cells, but less in MCF7 cells (Figure 3C). *ETAA1* encodes a replication stress protein that accumulates at sites of DNA damage and is a component of the ATR signalling response (Bass et al., 2016). Selection of RBMX-repressed cryptic 3' splice sites within *ETAA1* exon 5 removes a long portion of the open reading frame (Figure 3 – Figure supplement 2B). Consistent with the penetrance of this high amplitude *ETAA1* splicing defect being sufficiently high to affect protein production, no ETAA1 protein was detectable 72 hours after RBMX depletion from MDA-MB-231 cells (Figure 3D).

Another ultra-long exon is found within the *REV3L* gene that encodes the catalytic subunit of DNA polymerase ζ that functions in translesion DNA synthesis (Martin and Wood, 2019). RBMX similarly represses a cryptic 3' splice site within the ultra-long exon 13 of the *REV3L* gene (~4.2 Kb), that has an extremely high density of RBMX binding (Figure 3E). RT-PCR analysis confirmed a strong splicing switch to a cryptic splice site within *REV3L* exon 13 after RBMX was depleted from MDA-MB-231, MCF7 and NCI-H520 cells (Figure 3F).

We also detected extremely high density RBMX protein binding within exon 9 of the *ATRX* gene (3 Kb) that encodes a chromatin remodelling protein involved in mitosis. Depletion of RBMX results in expression of a shortened version of ATRX exon 9, caused by formation of an exitron through selection of cryptic 5' and 3' splice sites within exon 9 (Figure 3 – Figure supplement 2C).

RBMX protein-RNA interactions may insulate important splicing signals from the spliceosome.

The iCLIP data suggested a model where RBMX protein binding may insulate ultra-long exons so that cryptic splice sites cannot be accessed by the spliceosome. This model predicted that RBMX binding sites would be close to important sequences used for selection of cryptic splice sites. RBMX iCLIP tags mapped just upstream of the cryptic 3' splice sites within *ETAA1* exon 5 in HEK293 cells and MDA-MB-231 cells after RBMX depletion (Figure 3B), suggesting that RBMX may bind close to the branchpoints used for these cryptic splicing patterns. However, although usually located close to their associated 3' splice sites, in some cases branchpoints can be located far upstream (Gooding et al., 2006). We tested the prediction that RBMX may sterically interfere with components of the spliceosome by directly mapping the branchpoints associated with use of these cryptic *ETAA1* splice sites. To facilitate mapping of the branchpoint sequences used by the cryptic 3' splice site within *ETAA1* exon 5, we made a minigene by cloning the ultra-long *ETAA1* exon 5 and flanking intron sequences between constitutively spliced β-globin exons (Figure 3 – Figure supplement 2D). Confirming that this minigene recapitulated cryptic splicing patterns, after transfection into HEK293 cells we could detect splicing inclusion of both the full-length and shorter (cryptic) versions of *ETAA1* exon 5 *ETAA1* mRNA isoforms using multiplex RT-PCR (Figure 3 – Figure supplement 2E). We then used an RT-PCR assay (Figure 3 – Figure Supplement 2F) to monitor the position of branchpoints just upstream of the cryptic 3' splice sites of *ETAA1* exon 5 (Královičová et al., 2021). Sanger sequencing of the amplification product made in this assay confirmed that the branchpoint sequences used by these cryptic 3' splice sites are adjacent to RBMX binding sites (Figure 3G and Figure 3 – Figure supplement 2G).

Figure 3

Figure 3. RBMX protein is important for full-length splicing inclusion of ultra-long exons

involved in DNA repair and RNA polymerase II transcription. (A) Gene Ontology analysis of

genes regulated and bound by RBMX displaying significant Gene Ontology Biological Process

(GOBP) terms containing at least 5% of the total gene list. FDR, False Discovery Rate. Count,

number of genes in the GOBP group. GeneRatio, proportion of genes in the GOBP group relative

to the full list of RBMX-regulated genes. (B) Snapshot of RNA-seq merged tracks from MDA-MB-

231 cells and RBMX iCLIP tags from HEK293 cells from the IGV genome browser shows cryptic 3'

splice sites repressed by RBMX in *ETAA1* exon 5. At the bottom, schematic of PCR products

identified by RT-PCR in (C). **(C)** RT-PCR analysis shows splicing inclusion of *ETAA1* exon 5 upon

siRNA-mediated depletion of RBMX in the indicated cell lines. **(D)** Western blot analysis shows

ETAA1 protein expression is dependent on RBMX. Anti-Tubulin detection was used as loading

control. **(E)** Snapshot of RNA-seq merged tracks from MDA-MB-231 cells and RBMX iCLIP tags

from HEK293 cells from the IGV genome browser shows RBMX represses a cryptic 3' splice site

within the ultra-long exon 13 of *REV3L*. At the bottom, schematic of PCR products identified by RT-

PCR in (F). **(F)** RT-PCR analysis shows splicing inclusion of *REV3L* exon 13 upon siRNA-

mediated depletion of RBMX in the indicated cell lines. **(G)** Snapshot of RNA-seq merged tracks

from MDA-MB-231 cells and RBMX iCLIP tags from HEK293 cells from IGV genome browser. The

location of experimentally mapped branchpoints relative to RBMX binding is indicated.

Figure 3 – Figure supplement 1

A

B

C

D

E

Figure 3 – Figure supplement 1 (A) Gene Ontology analysis of genes bound by RBMX as

identified by iCLIP, displaying top 20 significant Gene Ontology Biological Process (GOBP).

Adjusted p-value were produced using the Benjamini-Hochberg method. Count, number of genes

in the GOBP group. GeneRatio, proportion of genes in the GOBP group relative to the full list of

RBMX-regulated genes. **(B)** Analysis as in (A) but relative to genes regulated by RBMX in both

MDA-MB-231 (this study) and HEK293 (Liu et al., 2017) as identified by RNA-seq. GOBP terms

containing at least 5% of the total gene list are shown. FDR, False Discovery Rate. **(C)** Analysis as

in (B) but relative to human genes containing ultra-long exons (equal or longer than 1 Kb, Ensembl

v.104). **(D)** Comet assay shows increased formation of DNA breaks in U2OS cells treated with

RBMX siRNA. Direction of comets is shown. Scale bars 200µm. **(E)** Quantification of percentage of

DNA in the tail of comets. n=58 cells in both conditions. ****, p<0.0001 (Mann-Whitney test).

Figure 3 – Figure supplement 2

Figure 3 – Figure supplement 2 (A) Snapshot of RNA-seq merged tracks from HEK293 cells (Liu et al., 2017) from the IGV genome browser shows cryptic 3' splice sites repressed by RBMX in *ETAA1* exon 5. **(B)** Schematic of ETAA1 protein in normal conditions and expected ETAA1 protein in RBMX-depleted cells. **(C)** Snapshot of RNA-seq merged tracks from MDA-MB-231 cells and RBMX iCLIP tags from HEK293 cells from the IGV genome browser shows an exitron repressed by RBMX in *ATRX* exon 9. **(D)** Schematic representation of *ETAA1* minigene cloned within pXJ41. **(E)** Example RT-PCR from *ETAA1* minigene shows detection of both the normal and cryptic version of exon 5. **(F)** Schematic of RT-PCR assay to detect branchpoints used during cryptic splicing of *ETAA1* exon 5. **(G)** RT-PCR analysis for mapping branchpoints used during *ETAA1* cryptic splicing (see Figure 3G). The distance from the relative cryptic splice site and from RBMX binding site as defined by iCLIP is indicated.

RBMXL2 and RBMY can replace the activity of RBMX in somatic cells

The above data showed that although RBMX can activate splicing of some exons, it predominantly operates as a splicing repressor in human somatic cells, and moreover has a key role in repressing cryptic splicing within ultra-long exons. This global pattern of RBMX activity is thus very similar to that previously reported for RBMXL2 in the germline, where RBMXL2 represses cryptic splice sites during meiosis. RBMXL2 is expressed during male meiosis when the X chromosome is silenced. To directly mimic this switch in protein expression we constructed a HEK293 RBMXL2-FLAG tetracycline-inducible cell line, from which we depleted RBMX using siRNA (Figure 4A). Western blots showed that RBMX was successfully depleted after siRNA treatment, and the RBMXL2-FLAG protein was strongly expressed after tetracycline induction, thus simulating their relative expression patterns in meiotic cells (Figure 4B). We then analysed cryptic splicing patterns. Confirming our previous finding, in the absence of tetracycline treatment depletion of RBMX led to increased selection of cryptic splice sites within *ETAA1* exon 5 and *REV3L* exon 13, and to formation of an exitron within *ATRX* exon 9 (Figure 4C-E, compare lanes 7-9 with lanes 10-12). Strikingly, tetracycline-induction of RBMXL2 was sufficient to repress production of each of these aberrant splice isoforms (Figure 4C-E, compare lanes 1-3 with lanes 4-6). We also globally investigated patterns of splicing in these rescue experiments by performing RNA-seq analysis of each of the experimental groups. For each target gene we analysed we found that RBMXL2 was able to replace the function of RBMX in splicing control (Figure 4 – Figure supplement 1B-D).

The above experiments showed that RBMXL2 is able to globally replace RBMX activity within somatic cells. RBMX and RBMXL2 are both more distantly related to the Y chromosome-encoded RBMY protein, with RBMX and RBMY diverging when the mammalian Y chromosome evolved (Figure 1A). RBMY has also been implicated in splicing control (Nasim et al., 2003; Venables et al., 2000), but its global functions are very poorly understood. We thus tested whether RBMY might also be performing a similar function to RBMX. Employing a HEK293 cell line containing

tetracycline-inducible, FLAG-tagged RBMY protein, we detected successful recovery of normal

splicing patterns of the ultra-long exons within the *ETAA1*, *REV3L* and *ATRX* genes within RBMX-

depleted cells 24 hours after tetracycline induction of RBMY (Figure 4 – Figure supplement 2).

These results indicate that even despite its more extensive divergence, RBMY can also

functionally replace RBMX in cryptic splice site control within long exons. Thus, splicing control

mechanisms by RBMX family proteins pre-date the evolution of the mammalian X and Y

chromosomes

Figure 4

9 **Figure 4. RBMXL2 can replace the activity of RBMX in ensuring proper splicing inclusion of**

10 **ultra-long exons. (A)** Schematic of the time-course experiment used to analyse RBMXL2 function

11 in RBMX-depleted HEK293 cells. All conditions were repeated in biological triplicates. **(B)** Western

12 blot analysis shows that RBMXL2-FLAG protein is stably expressed in HEK293 cells after 24 hours

13 of tetracycline induction, and RBMX protein is successfully depleted after 72 hours siRNA

treatment. **(C, E, G)** Capillary gel electrophoretograms show RNA processing patterns of

endogenous ultra-long exons within *ETAA1*, *REV3L* and *ATRX* controlled by RBMX and RBMXL2

analysed using isoform-specific RT-PCR. **(D, F, H)** Bar charts showing percentage splicing

inclusion (PSI) of cryptic isoforms from the endogenous *ETAA1*, *REV3L* and *ATRX* genes under

the different experimental conditions, relative to experiments in (C), (E) and (G) respectively. P-

values were calculated using unpaired t-test. *, p-value<0.05; ***, p-value<0.001. ****, p-

value<0.0001.

Figure 4 - Figure supplement 1

9 **Figure 4 – Figure supplement 1. RBMXL2 can replace the activity of RBMX in ensuring**

10 **proper splicing inclusion of ultra-long exons.** (A-C) Snapshot of RNA-seq merged tracks from

11 HEK293 cells from the IGV genome browser shows tetracycline-induced expression of RBMXL2

1 restores correct splicing patterns within *ETAA1* exon 5 (A), *REV3L* exon 13 (B) and *ATRX* exon 9

2 (C). Location of the splicing defects in RBMX-depleted cells is shown with red dotted lines.

Figure 4 - Figure supplement 2

Figure 4 – Figure supplement 2. RBMY can replace the activity of RBMX in ensuring proper

splicing inclusion of ultra-long exons. (A) Western blot analysis shows that RBMY-FLAG

protein is stably expressed in HEK293 cells after 24 hours of tetracycline induction, and RBMX

protein is successfully depleted after 72 hours siRNA treatment. All conditions were repeated in

biological triplicates. **(B, D, F)** Capillary gel electrophoretograms show RNA processing patterns of

endogenous ultra-long exons within *ETAA1*, *REV3L* and *ATRX* controlled by RBMX and RBMY

analysed using isoform-specific RT-PCR. **(C, E, G)** Bar charts showing percentage splicing

inclusion (PSI) of cryptic isoforms from the endogenous *ETAA1*, *REV3L* and *ATRX* genes under

the different experimental conditions, relative to experiments in (B), (D) and (F) respectively. P-

values were calculated using unpaired t-test. ***, p-value<0.001. ****, p-value<0.0001.

The disordered domain of RBMXL2 is required for efficient splicing control of ultra-long

exons

The above data showed that RBMX predominantly operates as a splicing repressor in somatic

cells, thus performing a functionally parallel role to RBMXL2 in the germline. Although RBMX

contains an RRM domain that is the most highly conserved region compared with RBMXL2 and

RBMY, splicing activation by RBMX depends on its C-terminal disordered domain that also binds

to RNA (Liu et al., 2017; Moursy et al., 2014). We thus reasoned that if RBMX and RBMXL2 were

performing equivalent molecular functions, rescue of splicing by RBMXL2 should be mediated by

the disordered region of RBMXL2 alone, independent of the RRM (Liu et al., 2017; Moursy et al.,

2014). To test this prediction, we created a new tetracycline-inducible HEK293 cell line expressing

the disordered region of RBMXL2 protein and not the RRM domain (RBMXL2ΔRRM, Figure 5B).

Tetracycline induction of this RBMXL2ΔRRM protein was able to rescue siRNA mediated depletion

of RBMX (Figures 5C-E), directly confirming that the C-terminal disordered domain of RBMXL2

protein is responsible for mediating cryptic splicing repression.

Figure 5

Figure 5. **The disordered domain of RBMXL2 is required to mediate splicing control of ultra-long exons in HEK293 cells.** (A) Western blot analysis shows that RBMXL2ΔRRM-FLAG protein is stably expressed in HEK293 cells after 24 hours of tetracycline induction, and RBMX protein is successfully depleted after 72 hours siRNA treatment. **(B, D, F)** Capillary gel electrophoretograms show RNA processing patterns of endogenous ultra-long exons within *ETAA1*, *REV3L* and *ATRX* analysed using isoform-specific RT-PCR. **(C, E, G)** Bar charts showing percentage splicing inclusion (PSI) of cryptic isoforms from the endogenous *ETAA1*, *REV3L* and *ATRX* genes under the different experimental conditions, relative to experiments in (B), (D) and (F) respectively. P-values were calculated using unpaired t-test. ****, p-value<0.0001.

Discussion.

We previously showed that the germ cell-specific RBMXL2 protein represses cryptic splice site selection during meiotic prophase. Here we find that this is part of a bigger picture, where the closely related but more ubiquitously expressed RBMX protein also provides a similar activity within somatic cells. Supporting this conclusion, both RBMX and RBMXL2 proteins most frequently operate as splicing repressors in their respective cell types. Furthermore, RBMXL2 and even the more diverged RBMY protein are able to provide a direct replacement for RBMX function within human somatic cells.

We further find that RBMX binds and is very important for proper splicing inclusion of a group of ultra-long exons, defined as exceeding 1 Kb in length. RBMXL2 similarly represses cryptic splice sites within ultra-long exons of genes involved in genome stability including *Brca2* and *Meioc* (I. Ehrmann et al., 2019). Although many of the splice sites we find to be repressed by RBMX within ultra-long exons are already annotated, they are not usually selected in the human cell lines we investigated and thus represent potential decoy splice sites that would interfere with full-length gene expression. Long human exons provide an enigma in understanding gene expression. Most human exons have evolved to be quite short (~130 bp) to facilitate a process called exon definition, in which protein-protein interactions between early spliceosome components bound to closely juxtaposed splice sites promote full spliceosome assembly (Black, 1995; Robberson et al., 1990). Exon definition also requires additional RNA binding proteins to recognise exons and flanking intron sequences. These include members of the SR protein family that bind to exonic splicing enhancers (ESEs) and activate exon inclusion, with exons typically having higher ESE content relative to introns. While the mechanisms that ensure proper splicing inclusion of long exons are not well understood, cryptic splice sites would be statistically more likely to occur within long exons compared to short exons, where they could prevent full-length exon inclusion. Cryptic splice sites within long exons could be particularly problematic (compared to an intronic location) since they would be embedded within a high ESE sequence environment. For example some long exons require interactions with the SR protein SRSF3 and hnRNP K and phase separation of transcription factors to be spliced (Kawachi et al., 2021).

Meiotic sex chromosome inactivation represses expression of a panel of essential genes, opening the need for alternative routes to fulfil their function (Turner, 2015). A number of essential X-linked genes have generated autosomal retrogenes that are expressed during meiosis however genetic inactivation of many of these retrogenes causes a phenotype that manifests outside of meiosis. (Wang, 2004). An exception is exemplified by the RPL10 and RPL10L proteins that are 95% identical: *RPL10* mutation causes meiotic arrest, and RPL10L has been shown to directly replace its X-linked ortholog RPL10 during meiosis (Jiang et al., 2017; Wang, 2004). RBMXL2 is the only other X-linked retrogene that has been shown to be essential for meiotic prophase (I. Ehrmann et

al., 2019). Here we show that ectopic expression of RBMXL2 can compensate for lack of RBMX in somatic cells. This is consistent with a recent model suggesting that RBMXL2 replaces RBMX function during meiosis because of transcriptional inactivation of the X chromosome (Aldalaqan et al., 2022). This general requirement for functionally similar RBMX family proteins across somatic and germ cells further suggest that RBMX-family functions in splicing control have been required for ~200 million years, since before the divergence of separate *RBMX* and *RBMY* genes early in mammalian evolution.

The iCLIP data reported here show a high density of RBMX binding within ultra-long exons, consistent with a model in which RBMX protein binding to RNA masks sequences required for cryptic splice sites selection. Such RBMX binding would block access to spliceosome components or splicing activator proteins (Figure 6). Our data show that the C-terminal disordered domain of RBMXL2 protein is sufficient to control splicing inclusion of ultra-long exons. This is exactly analogous to the mechanism of control of splicing activation by RBMX, which occurs via recognition of m6A modified RNA targets via the C-terminal disordered domain (Liu et al., 2017). Intriguingly, global studies have shown that m6A residues are enriched within some long internal exons (Dominissini et al., 2012), where they might help facilitate RBMX protein-RNA interactions. The C-terminal disordered region of RBMX is also reported to mediate protein-protein interactions therefore shorter exons that show defective splicing in RBMX-depleted cells but are not directly bound by RBMX could rely on different regulatory mechanisms. RBMY, RBMX and RBMXL2 directly interact with the SR protein Tra2β (Elliott et al., 2000; Venables et al., 2000) and have opposing functions during RNA binding and splicing regulation (Nasim et al., 2003; Venables et al., 2000). Hence it is still possible that RBMX family proteins counteract recognition by SR proteins of ESEs near cryptic splice sites via a protein-protein interaction mechanism.

Extensive literature shows that RBMX is important for genome stability, including being involved in replication fork activity (Munschauer et al., 2018; Zheng et al., 2020), sensitivity to genotoxic drugs (Adamson et al., 2012) and cell proliferation (https://orcs.thebiogrid.org/Gene/27316). Interestingly, many of the ultra-long exons controlled by RBMX are within genes important for genome stability, including *REV3L*, *ATRX* and *ETAA1*. This makes it likely that RBMX contributes to maintaining genome stability through ensuring full-length protein expression of genes important in this process. As an example, we show here that depletion of RBMX protein causes aberrant selection of a high amplitude cryptic splice site within *ETAA1* exon 5 which prevents detectable expression of ETAA1 protein, and contributes to genome instability (Bass et al., 2016). Cancer and neurological disorders are amongst the most common human diseases associated with defective DNA damage response (Jackson and Bartek, 2009). The double role of RBMX in genome maintenance via both direct participation in the DNA damage response and splicing regulation of genome stability genes could explain why mutations of *RBMX* are associated with an intellectual disability syndrome (Cai

1 et al., 2021; Shashi et al., 2015), and why RBMX has been identified as a potential tumour

2 suppressor (Adamson et al., 2012; Elliott et al., 2019).

Figure 6

Figure 6. Model of cryptic splice site repression within ultra-long exons by RBMX family proteins.

Ultra-long exons are intrinsically fragile as they contain cryptic splice sites within an environment

rich in Exonic Splicing Enhancers (ESEs). RBMX protein binding within ultra-long exons may

directly block access of spliceosome components to cryptic splice sites, and depletion of RBMX

from somatic cells activates selection of cryptic splice sites. This means a shorter version of the

originally ultra-long exon is included, that fits more easily with exon definition rules normally

followed for median size exons. During meiosis, lack of RBMX caused by X chromosome

inactivation is compensated by expression of RBMXL2 protein.

Materials and methods.

Cell culture and cell lines

MDA-MB-231 (ATCC® HTB-26™), MCF7 (ATCC® HTB-22™), U2OS (ATCC® HTB-96™) and NCI-H520 (ATCC® HTB-182™) cells were maintained in Dulbecco's Modified Eagle's Medium (DMEM) high glucose pyruvate medium (Gibco, #10569010), supplemented with 10% fetal bovine serum (FBS, Gibco, #21875034) and 1% Penicillin-Streptomycin (Gibco, #15140130). HEK293 (ATCC® CRL-1573) were maintained in DMEM plus 10% fetal bovine serum. Cell line validation was carried out using STR profiling was according to the ATCC® guidelines. All cell lines underwent regular mycoplasma testing.

Generation of tetracycline-inducible cell lines

RMBX, *RBMXL2*, *RBMY* and *RBMXL2ΔRRM* genes were cloned onto a FLAG-pcDNA5 vector and co-transfected with pOG44 plasmid into Flp-In HEK293 cells like previously described (Ehrmann et al., 2016). RBMX-FLAG, RBMXL2-FLAG, RBMY-FLAG and RBMXL2ΔRRM-FLAG expression was induced by the addition of 1μg/ml tetracycline (Sigma-Aldrich) to promote expression via a tetracycline-inducible promoter. The Flp-In HEK293 cells were cultured in high glucose pyruvate medium (Gibco, #10569010), supplemented with 10% FBS (Gibco, #21875034) and 1% Penicillin-Streptomycin (Gibco, #15140130).

siRNA knockdown and tetracycline induction

RBMX transient knockdown was established using two different pre-designed siRNAs targeting *RBMX* mRNA transcripts (hs.Ri.RBMX.13.1 and hs.Ri.RBMX.13.2, from Integrated DNA Technologies). Negative control cells were transfected with control siRNA (Integrated DNA Technologies, # 51-01-14-04). Cells were seeded onto 6-well plates forward transfected with Lipofectamine™ RNAiMAX transfection reagent (Invitrogen, # 13778150) according to manufacturer's instructions using 30 pmol of siRNA for 72h at 37°C before harvesting. For tetracycline-inducible cell lines, Flp-In HEK293 cells expressing either RBMXL2-FLAG, or RBMY-FLAG, or RBMXL2ΔRRM-FLAG genes were similarly seeded onto 6-well plates and treated with RBMX and control siRNAs for 72h at 37°C. 24h before harvesting 1μg/ml of tetracycline (Sigma-Aldrich) was added to half of the siRNA-treated samples to promote the expression of RBMXL2-FLAG and RBMY-FLAG.

RNA-seq

RNA was extracted from cells using RNeasy Plus Mini Kit (Qiagen #74134) following manufacturer's instructions and re-suspended in nuclease-free water. RNA samples were DNase treated (Invitrogen, AM1906). For siRNA treated MDA-MB-231 cells, paired-end sequencing was done initially for two samples, one of negative control and one of RBMX knock-down, using an

Illumina NextSeq 500 instrument. Adapters were trimmed using trimmomatic v0.32. Three

additional biological repeats of negative control and RBMX siRNA treated MDA-MB-231 cells were

then sequenced using an Illumina HiSeq 2000 instrument. The base quality of raw sequencing

reads was checked with FastQC (Andrews, 2010). RNA-seq reads were mapped to the human

genome assembly GRCh38/hg38 using STAR v.2.4.2 (Dobin et al., 2013) and subsequently

quantified with Salmon v. 0.9.1 (Patro et al., 2017) and DESeq2 v.1.16.1 (Love et al., 2014) on R

v.3.5.1. All snapshots indicate merged tracks produced using samtools (Li et al., 2009) and

visualised with IGV (Robinson et al., 2011). For HEK293 cells treated with either RBMX or control

siRNA, either in the absence or in the presence of tetracycline, RNAs were sequenced using an

Illumina NextSeq 500 instrument. Quality of the reads was checked with FastQC (Andrews, 2010).

Reads were then aligned to the human genome assembly GRCh38/hg38 to produce BAM files

using hisat2 v.2.2.1 (Kim et al., 2015) and samtools v.1.14 (Li et al., 2009) and visualised using

IGV (Robinson et al., 2011).

Identification of splicing changes

Initial comparison of single individual RNA-seq samples from RBMX-depleted and control cells was

carried out using MAJIQ (Vaquero-Garcia et al., 2016), which identified 596 unique local splicing

variations (LSV) at a 20% dPSI minimum cut off from 505 different genes potentially regulated by

RBMX. These LSVs were then manually inspected using the RNA-seq data from the second RNA

sequencing of biological replicates for both RBMX-depleted and control cells, by visual analysis on

the UCSC browser (Karolchik et al., 2014) to identify consistent splicing changes that depend on

RBMX expression. The triplicate RNA-seq samples were further analysed for splicing variations

using SUPPA2 (Trincado et al., 2018), which identified 6702 differential splicing isoforms with p-

value < 0.05. Predicted splicing changes were confirmed by visual inspection of RNA-seq reads

using the UCSC (Karolchik et al., 2014) and IGV (Robinson et al., 2011) genome browsers.

Identification of common splicing changes between RBMX-depleted MDA-MB-231 and HEK293

cells was done comparing data from this study with data from GSE74085 (Liu et al., 2017).

iCLIP

iCLIP experiments were performed on triplicate samples in RBMX-FLAG expressing Flp-In

HEK293 cells using the protocol described in (Huppertz et al., 2014). Briefly cells were grown in 10

cm tissue culture dishes and irradiated with 400 mJ cm−2 ultraviolet-C light on ice, lysed and

sonicated using Diagenode Bioruptor® Pico sonicator for 10 cycles with alternating 30 secs on/ off

at low intensity and 1 mg of protein was digested with 4 U of Turbo DNase (Ambion, AM2238) and

0.28 U/ml (low) or 2.5 U/ml (high) of RNAse I (Thermo Scientific, EN0602). The digested lysates

were immunoprecipitated with Protein G Dynabeads™ (Invitrogen, #10003D) and either 5 µg anti-

FLAG antibody (Sigma-Aldrich, F1804) or 5 µg IgG (Santa Cruz biotechnology, sc-2025).

Subsequently a pre-adenylated adaptor L3-IR-App (Zarnegar et al., 2016) was ligated to the 3' of

the RNA fragments. The captured Protein-RNA complexes were visualised using Odyssey LI-COR

CLx imager scanning in both the 700nm and 800nm channels. The RNA bound to the proteins was

purified, reverse transcribed with barcoded RT oligos complementary to the L3 adaptor. The

cDNAs were purified using Agencourt AMPure XP beads (Beckman Coulter™, A63880),

circularised and linearised by PCR amplification. The libraries were gel purified and sequenced on

Illumina NextSeq 500. All iCLIP sequencing read analysis was performed on the iMaps webserver

(imaps.goodwright.com) using standardised icount demultiplex and analyse work flow. Briefly,

reads were demultiplexed using the experimental barcodes, UMIs (unique molecular identifiers)

were used to remove PCR duplicates and reads were mapped to the human genome sequence

(version hg38/GRCh37) using STAR (Dobin et al., 2013). Crosslinked sites were identified on the

iMAPS platform and the iCount group analysis workflow was used to merge the replicate samples.

K-mer enrichment analysis

K-mer motif enrichment was performed with the z-score approach using the kmer_enrichment.py

script from the iCLIPlib suite of tools (https://github.com/sudlab/iCLIPlib). All transcripts for each

non-overlapping protein coding gene from the Ensembl v.105 annotation were merged into a single

transcript, used for this analysis, using cgat gtf2gtf --method=merge-transcripts (Sims et al., 2014).

Each crosslinked base from the merged replicate bam file was extended 15 nucleotides in each

direction. For every hexamer, the number of times a crosslink site overlaps a hexamer start

position was counted within the gene and then summed across all genes. This occurrence was

also calculated across 100 randomizations of the crosslink positions within genes. The z-score was

thus calculated for each hexamer as (occurrence – occurrence in randomized sequences) /

standard deviation of occurrence in randomized sequences.

Exon size analysis

Annotations of all human exons related to position and size were downloaded from Ensembl

Genes v.105 (http://www.ensembl.org/biomart/). Selection of exons expressed in HEK293 was

performed using data from control RNA-seq samples of the dataset GSE74085 (Liu et al., 2017),

subsequently filtered to focus on mRNA exons using biomaRt v.2.52.0 (Durinck et al., 2005). Size

of the mRNA exons containing RBMX-regulated splicing patterns was annotated using IGV

(Robinson et al., 2011). iCLIP tags were extended to 80 nt sequences centered at the crosslinked

site, and annotated within human exons using ChIPseeker v.1.32.0 (Yu et al., 2015) and Esembl

Genes v.105. iCLIP tags present in mRNA exons were filtered using biomaRt v.2.52.0 (Durinck et

al., 2005). iCLIP-containing exons were listed once, independently of the number of tags or tag

score. Plots were created using ggplot2 v.3.3.6. (Wickham, 2016) on R v.4.2.1. Statistical analyses

to compare exon length distributions between samples were performed by Wilcoxon Rank Sum

1. and Kruskal-Wallis tests using base R stats package v.4.2.1 and pseudorank v.1.0.1 (Happ et al.,
2. 2020). Significant enrichment of ultra-long exons in RBMX regulated and bound exons was
3. performed using the "N-1" Chi-squared test (Campbell, 2007) on the MedCalc Software version
4. 20.218.
5.

6. **Gene Ontology analyses**
7. Gene Ontology Analyses were performed in R v.4.2.1 using GOstats v.2.62.0 (Falcon and
8. Gentleman, 2007) except for [Figure 3 – Figure supplement 1A] for which clusterProfiler::enrichGO
9. v.4.4.4 (Yu et al., 2012) was used. Entrez annotations were obtained with biomaRt v.2.52.0
10. (Durinck et al., 2005). Read counts from control treated HEK293 cells (Liu et al., 2017) and the
11. Bioconductor annotation data package org.Hs.eg.db v.3.15.0 were used as background. P-values
12. were adjusted by false discovery rate using the base R stats package v.4.2.1, except for [Figure 3
13. – Figure supplement 1A] for which the default Benjamini-Hochberg method was used while running
14. enrichGO. Significantly enriched GOBP pathways were filtered with a p-value cut-off of 0.05.
15. Redundant terms identified with GOstats were removed using Revigo (Supek et al., 2011) with
16. SimRel similarity measure against human genes eliminating terms with dispensability score above
17. 0.5. The dot-plots were produced using ggplot2 v.3.3.6 (Wickham, 2016) focussing on
18. representative terms associated to at least 5% of the initial gene list. Full GOBP lists can be found
19. in Figure 3 – Source Data 1.
20.

21. **Comet assay**
22. The comet assay was performed using the Abcam Comet Assay kit (ab238544) according to
23. manufacturer's instructions. Briefly U2OS cells transfected with RBMX siRNA or control siRNA
24. were harvested after 72 hours, 1×10^5 cells were mixed with cold PBS. Cells in PBS were mixed
25. with low melting comet agarose (1/10) and layered on the glass slides pre-coated with low melting
26. comet agarose. The slides were lysed in 1x lysis buffer (pH10.0, Abcam Comet Assay kit) for 48
27. hours at 4°C, immersed in Alkaline solution (300 mM NaOH, pH>13, 1 mM EDTA) for 30 min at
28. 4°C in the dark and then electrophoresed in Alkaline Electrophoresis Solution (300 mM NaOH,
29. pH>13, 1 mM EDTA) at 300mA, 1volt/cm for 20 min. The slide was then washed in pre-chilled DI
30. H2O for 2 min, fixed in 70% ethanol for 5 min and stained with 1x Vista Green DNA Dye (1/10000
31. in TE Buffer (10 mM Tris, pH 7.5, 1 mM EDTA), Abcam Comet Assay kit) for 15 min and visualized
32. under fluorescence microscopy Zeiss AxioImager (System 3). Comet quantification was performed
33. using OpenComet (Gyori et al., 2014).
34.

35. **RNA extraction and cDNA synthesis for transcript isoform analysis.**
36. RNA was extracted using standard TRIzol™ RNA extraction (Invitrogen, #15596026) following
37. manufacturer's instructions. cDNA was synthesized from 500 ng total RNA using SuperScript™

VILO™ cDNA synthesis kit (Invitrogen #11754050) following manufacturer's instructions. To

analyse the splicing profiles of the alternative events primers were designed using Primer 3 Plus

(Untergasser et al., 2012) and the predicted PCR products were confirmed using the UCSC *In-*

Silico PCR tool. *ETAA1* transcript isoform containing the long exon 5 was amplified by RT-PCR

using primers 5'-GCTGGACATGTGGATTGGTG-3' and 5'-GTGCTCCAAAAAGCCTCTGG-3',

while *ETAA1* transcript isoform containing the short exon 5 was amplified using primers 5'-

GCTGGACATGTGGATTGGTG-3' and 5'-GTGGGAGCTGCATTTACAGATG-3'. RT-PCR with this

second primer pair could in principle amplify also a 2313 bp product from the *ETAA1* transcript

isoform containing the long exon 5, however PCR conditions were chosen to selectively analyse

shorter fragments. *REV3L* 5'-*TCACTGTGCAGAAATACCCAC*-3', 5'-

AGGCCACGTCTACAAGTTCA-3', 5'-*ACATGGGAAGAAAGGGCACT*-3'. *ATRX 5'-*

TGAAACTTCATTTTCAACCAAATGCTC-3' and *5'-ATCAAGGGGATGGCAGCAG-3'* All PCR

reactions were performed using GoTaq® G2 DNA polymerase kit from Promega following the

manufacturer's instructions. All PCR products were examined using the QIAxcel® capillary

electrophoresis system 100 (Qiagen). Statistical analyses were performed using GraphPad Prism

9.5.0.

Western blot analyses

Harvested cells treated with either control siRNA or siRNA against RBMX were resuspended in

100mM Tris-HCL, 200mM DTT, 4% SDS, 20% Glycerol, 0.2% Bromophenol blue, then sonicated

(Sanyo Soniprep 150) and heated to 95°C for 5 minutes. Protein separation was performed by

SDS-PAGE. Proteins were then transferred to a nitrocellulose membrane, incubated in blocking

buffer (5% Milk in 2.5% TBS-T) and stained with primary antibodies diluted in blocking buffer to the

concentrations indicated below, at 4°C over-night. After incubation the membranes were washed

three times with TBS-T and incubated with the secondary antibodies for 1 hour at room

temperature. Detection was carried out using the Clarity™ Western ECL Substrate (Cytiva,

RPN2232) and developed using medical X-ray film blue film in an X-ray film processor developer.

The following primary antibodies were used at the concentrations indicated: anti-RBMX (Cell

Signalling, D7C2V) diluted 1:1000, anti-ETAA1 (Sigma, HPA035048) diluted 1:1000, anti-Tubulin

(Abcam, ab18251) diluted 1:2000, anti-GAPDH (Abcepta, P04406) diluted 1:2000 and anti-FLAG

(Sigma, F1804) diluted 1:2000.

Minigene construction and validation

A genomic region containing *ETAA1* exon 5 and flanking intronic sequences were PCR amplified

from human genomic DNA using the primers

5'-AAAAAAAAACAATTGAGTTAAGACTTTTCAGCTTTTCTGA-3' and 5'-

AAAAAAAAACAATTGAGTGCTGGGAAAGAATTCAA

TGT-3'and cloned into pXJ41 (Bourgeois et al., 1999). Splicing patterns were monitored after

transfection into HEK293 cells. RNA was extracted with TRIzol™ (Invitrogen, #15596026) and

analysed using a One Step RT-PCR kit (Qiagen, #210210) following manufacturer's instructions.

RT–PCR experiments used 100 ng of RNA in a 5-µl reaction using a multiplex RT-PCR using

primers: 5'-GCTGGACATGTGGATTGGTG-3', 5'-GTGGGAGCTGCATTTACAGATG-3' and 5'-

GTGCTCCAAAAAGCCTCTGG-3 '. Reactions were analysed and quantified using the QIAxcel®

capillary electrophoresis system 100 (Qiagen).

Branchpoint analysis

Using the RNA from the long minigene transfections with RBMX and RBMXΔRRM, total RNA was

extraction using TRIZOL reagent (Life Technologies) following the standard manufacturer's

instructions, RNA concentration was quantified by NanoDrop UV-Vis spectrophotometer and

treated with DNase I (Invitrogen, Am1906). 1µg of purified RNA was reverse transcribed with a

SuperScript™ III Reverse Transcriptase (Invitrogen, #18080093) using ETAA1 DBR R1 RT-PCR

primer 5′-AAGTTCTTCTTCTTGACTTTGTGTT-3′ and treated with RNaseH (New England Biolabs,

M0297S). 1µl of the cDNA was used for PCR amplification reactions using using GoTaq® G2 DNA

Polymerase (Promega, #M7845) in the standard 25µl reaction following the manufacturer's

instructions. PCR amplification was carried out using 2 different primer sets ETAA1 DBR R2 5′-

GCTCTTGAATCACATCTAGCTCT-3′ and ETAA1 DBR F1 5′-AGCCAAACTAACTCAGCAACA-3′,

ETAA1 DBR R2 5′-GCTCTTGAATCACATCTAGCTCT-3′ and ETAA1 DBR F2 5′-

AGCATTTGAATCCAGGCAGC-3′ with 18 cycles of amplification at an annealing temperature of

56°C. PCR products were sub cloned into the pGEM-T-Easy vector (Promega, A1360) using

manufactures instructions. Plasmids were subjected to Sanger sequencing (Source BioScience)

and the sequences were checked with BioEdit (Hall, 1999) and aligned to the genome with UCSC

(Karolchik et al., 2014).

Accession numbers

Genomic data is deposited at the Gene Expression Omnibus accession GSE233498.

References.

Adamson B, Smogorzewska A, Sigoillot FD, King RW, Elledge SJ. 2012. A genome-wide homologous recombination screen identifies the RNA-binding protein RBMX as a component of the DNA-damage response. *Nat Cell Biol* **14**:318–328. doi:10.1038/ncb2426

Aldalaqan S, Dalgliesh C, Luzzi S, Siachisumo C, Reynard LN, Ehrmann I, Elliott DJ. 2022. Cryptic splicing: common pathological mechanisms involved in male infertility and neuronal diseases. *Cell Cycle* **21**:219–227. doi:10.1080/15384101.2021.2015672

Andrews S. 2010. FastQC - A quality control tool for high throughput sequence data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc/. *Babraham Bioinforma*. doi:citeulike-article-id:11583827

Attig J, Agostini F, Gooding C, Chakrabarti AM, Singh A, Haberman N, Zagalak JA, Emmett W, Smith CWJ, Luscombe NM, Ule J. 2018. Heteromeric RNP Assembly at LINEs Controls Lineage-Specific RNA Processing. *Cell* **174**:1067-1081.e17. doi:10.1016/j.cell.2018.07.001

Bass TE, Luzwick JW, Kavanaugh G, Carroll C, Dungrawala H, Glick GG, Feldkamp MD, Putney R, Chazin WJ, Cortez D. 2016. ETAA1 acts at stalled replication forks to maintain genome integrity. *Nat Cell Biol* **18**:1185–1195. doi:10.1038/ncb3415

Black DL. 1995. Finding splice sites within a wilderness of RNA. *RNA*.

Bourgeois CF, Popielarz M, Hildwein G, Stevenin J. 1999. Identification of a Bidirectional Splicing Enhancer: Differential Involvement of SR Proteins in 5′ or 3′ Splice Site Activation. *Mol Cell Biol* **19**:7347–7356. doi:10.1128/mcb.19.11.7347

Cai T, Cinkornpumin JK, Yu Z, Villarreal OD, Pastor WA, Richard S. 2021. Deletion of RBMX RGG/RG motif in Shashi-XLID syndrome leads to aberrant p53 activation and neuronal differentiation defects. *Cell Rep* **36**. doi:10.1016/j.celrep.2021.109337

Campbell I. 2007. Chi-squared and Fisher-Irwin tests of two-by-two tables with small sample recommendations. *Stat Med* **26**. doi:10.1002/sim.2832

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**:15–21. doi:10.1093/bioinformatics/bts635

Dominissini D, Moshitch-Moshkovitz S, Schwartz S, Salmon-Divon M, Ungar L, Osenberg S, Cesarkas K, Jacob-Hirsch J, Amariglio N, Kupiec M, Sorek R, Rechavi G. 2012. Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. *Nature* **485**:201–206. doi:10.1038/nature11112

Durinck S, Moreau Y, Kasprzyk A, Davis S, De Moor B, Brazma A, Huber W. 2005. BioMart and
Bioconductor: A powerful link between biological databases and microarray data analysis.
Bioinformatics **21**. doi:10.1093/bioinformatics/bti525

Ehrmann I., Crichton JH, Gazzara MR, James K, Liu Y, Grellscheid SN, Curk T, de Rooij D, Steyn
JS, Cockell S, Adams IR, Barash Y, Elliott DJ. 2019. An ancient germ cell-specific RNA-
binding protein protects the germline from cryptic splice site poisoning. *Elife* **8**.
doi:10.7554/eLife.39304

Ehrmann Ingrid, Crichton JH, Gazzara MR, James K, Liu Y, Grellscheid SN, Curk T, de Rooij D,
Steyn JS, Cockell S, Adams IR, Barash Y, Elliott DJ. 2019. An ancient germ cell-specific
RNA-binding protein protects the germline from cryptic splice site poisoning. *Elife* **8**.
doi:10.7554/eLife.39304

Ehrmann I, Gazzara MR, Pagliarini V, Dalgliesh C, Kheirollahi-Chadegani M, Xu Y, Cesari E,
Danilenko M, Maclennan M, Lowdon K, Vogel T, Keskivali-Bond P, Wells S, Cater H, Fort P,
Santibanez-Koref M, Middei S, Sette C, Clowry GJ, Barash Y, Cunningham MO, Elliott DJ.
2016. A SLM2 Feedback Pathway Controls Cortical Network Activity and Mouse Behavior.
Cell Rep **17**:3269–3280. doi:10.1016/j.celrep.2016.12.002

Elliott DJ, Bourgeois CF, Klink A, Stévenin J, Cooke HJ. 2000. A mammalian germ cell-specific
RNA-binding protein interacts with ubiquitously expressed proteins involved in splice site
selection. *Proc Natl Acad Sci U S A* **97**:5717–5722. doi:10.1073/pnas.97.11.5717

Elliott DJ, Dalgliesh C, Hysenaj G, Ehrmann I. 2019. RBMX family proteins connect the fields of
nuclear RNA processing, disease and sex chromosome biology. *Int J Biochem Cell Biol* **108**.
doi:10.1016/j.biocel.2018.12.014

Elliott DJ, Millar MR, Oghene K, Ross A, Kiesewetter F, Pryor J, McIntyre M, Hargreave TB,
Saunders PTK, Vogt PH, Chandley AC, Cooke H. 1997. Expression of RBM in the nuclei of
human germ cells is dependent on a critical region of the Y chromosome long arm. *Proc Natl*
Acad Sci. doi:10.1073/pnas.94.8.3848

Falcon S, Gentleman R. 2007. Using GOstats to test gene lists for GO term association.
Bioinformatics **23**:257–258. doi:10.1093/bioinformatics/btl567

Gooding C, Clark F, Wollerton MC, Grellscheid S-N, Groom H, Smith CWJ. 2006. A class of
human exons with predicted distant branch points revealed by analysis of AG dinucleotide
exclusion zones. *Genome Biol* **7**:R1. doi:10.1186/gb-2006-7-1-r1

Gyori BM, Venkatachalam G, Thiagarajan PS, Hsu D, Clement MV. 2014. OpenComet: An
automated tool for comet assay image analysis. *Redox Biol* **2**.

doi:10.1016/j.redox.2013.12.020

Hall TA. 1999. BIOEDIT: a user-friendly biological sequence alignment editor and analysis program
for Windows 95/98/ NT. *Nucleic Acids Symp Ser* **41**.

Happ M, Zimmermann G, Brunner E, Bathke AC. 2020. Pseudo-ranks: How to calculate them
efficiently in r. *J Stat Softw* **95**. doi:10.18637/jss.v095.c01

Huppertz I, Attig J, D'Ambrogio A, Easton LE, Sibley CR, Sugimoto Y, Tajnik M, König J, Ule J.
2014. iCLIP: Protein-RNA interactions at nucleotide resolution. *Methods* **65**.
doi:10.1016/j.ymeth.2013.10.011

Jackson SP, Bartek J. 2009. The DNA-damage response in human biology and disease. *Nature*.
doi:10.1038/nature08467

Jiang L, Li T, Zhang X, Zhang B, Yu C, Li Y, Fan S, Jiang X, Khan T, Hao Q, Xu P, Nadano D,
Huleihel M, Lunenfeld E, Wang PJ, Zhang Y, Shi Q. 2017. RPL10L Is Required for Male
Meiotic Division by Compensating for RPL10 during Meiotic Sex Chromosome Inactivation in
Mice. *Curr Biol* **27**:1498-1505.e6. doi:10.1016/j.cub.2017.04.017

Karolchik D, Barber GP, Casper J, Clawson H, Cline MS, Diekhans M, Dreszer TR, Fujita PA,
Guruvadoo L, Haeussler M, Harte RA, Heitner S, Hinrichs AS, Learned K, Lee BT, Li CH,
Raney BJ, Rhead B, Rosenbloom KR, Sloan CA, Speir ML, Zweig AS, Haussler D, Kuhn RM,
Kent WJ. 2014. The UCSC Genome Browser database: 2014 update. *Nucleic Acids Res* **42**.
doi:10.1093/nar/gkt1168

Kawachi T, Masuda A, Yamashita Y, Takeda J, Ohkawara B, Ito M, Ohno K. 2021. Regulated
splicing of large exons is linked to phase-separation of vertebrate transcription factors. *EMBO
J* **40**. doi:10.15252/embj.2020107485

Kim D, Langmead B, Salzberg SL. 2015. HISAT: A fast spliced aligner with low memory
requirements. *Nat Methods* **12**. doi:10.1038/nmeth.3317

Konig J, Zarnack K, Rot G, Curk T, Kayikci M, Zupan B, Turner DJ, Luscombe NM, Ule J. 2011.
ICLIP - transcriptome-wide mapping of protein-RNA interactions with individual nucleotide
resolution. *J Vis Exp*. doi:10.3791/2638

Královičová J, Borovská I, Pengelly R, Lee E, Abaffy P, Šindelka R, Grutzner F, Vořechovský I.
2021. Restriction of an intron size en route to endothermy. *Nucleic Acids Res* **49**:2460–2487.
doi:10.1093/nar/gkab046

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R.
2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*.

doi:10.1093/bioinformatics/btp352

Liu N, Zhou KI, Parisien M, Dai Q, Diatchenko L, Pan T. 2017. N6-methyladenosine alters RNA
structure to regulate binding of a low-complexity protein. *Nucleic Acids Res* **45**:6051–6063.
doi:10.1093/nar/gkx141

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* **15**. doi:10.1186/s13059-014-0550-8

Ma K, Inglis JD, Sharkey A, Bickmore WA, Hill RE, Prosser EJ, Speed RM, Thomson EJ, Jobling
M, Taylor K, Wolfe J, Cooke HJ, Hargreave TB, Chandley AC. 1993. A Y chromosome gene
family with RNA-binding protein homology: Candidates for the azoospermia factor AZF
controlling human spermatogenesis. *Cell* **75**:1287–1295. doi:10.1016/0092-8674(93)90616-X

Marquez Y, Höpfler M, Ayatollahi Z, Barta A, Kalyna M. 2015. Unmasking alternative splicing
inside protein-coding exons defines exitrons and their role in proteome plasticity. *Genome*
Res **25**. doi:10.1101/gr.186585.114

Martin SK, Wood RD. 2019. DNA polymerase ζ in DNA replication and repair. *Nucleic Acids Res.*
doi:10.1093/nar/gkz705

Moursy A, Allain FHT, Cléry A. 2014. Characterization of the RNA recognition mode of hnRNP G
extends its role in SMN2 splicing regulation. *Nucleic Acids Res* **42**:6659–6672.
doi:10.1093/nar/gku244

Munschauer M, Nguyen CT, Sirokman K, Hartigan CR, Hogstrom L, Engreitz JM, Ulirsch JC, Fulco
CP, Subramanian V, Chen J, Schenone M, Guttman M, Carr SA, Lander ES. 2018. The
NORAD lncRNA assembles a topoisomerase complex critical for genome stability. *Nature.*
doi:10.1038/s41586-018-0453-z

Nasim MT, Chernova TK, Chowdhury HM, Yue BG, Eperon IC. 2003. HnRNP G and Tra2β:
Opposite effects on splicing matched by antagonism in RNA binding. *Hum Mol Genet*
12:1337–1348. doi:10.1093/hmg/ddg136

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast and bias-aware
quantification of transcript expression. *Nat Methods* **14**:417–419. doi:10.1038/nmeth.4197

Robberson BL, Cote GJ, Berget SM. 1990. Exon definition may facilitate splice site selection in
RNAs with multiple exons. *Mol Cell Biol* **10**:84–94. doi:10.1128/mcb.10.1.84

Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP. 2011.
Integrative genomics viewer. *Nat Biotechnol.* doi:10.1038/nbt.1754

Shashi V, Xie P, Schoch K, Goldstein DB, Howard TD, Berry MN, Schwartz CE, Cronin K, Sliwa S,
Allen A, Need AC. 2015. The RBMX gene as a candidate for the Shashi X-linked intellectual
disability syndrome. *Clin Genet* **88**:386–390. doi:10.1111/cge.12511

Sibley CR, Blazquez L, Ule J. 2016. Lessons from non-canonical splicing. *Nat Rev Genet*.
doi:10.1038/nrg.2016.46

Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. Revigo summarizes and visualizes long lists of
gene ontology terms. *PLoS One* **6**. doi:10.1371/journal.pone.0021800

Trincado JL, Entizne JC, Hysenaj G, Singh B, Skalic M, Elliott DJ, Eyras E. 2018. SUPPA2: fast,
accurate, and uncertainty-aware differential splicing analysis across multiple conditions.
Genome Biol **19**:40. doi:10.1186/s13059-018-1417-1

Turner JMA. 2015. Meiotic Silencing in Mammals. *Annu Rev Genet* **49**:395–412.
doi:10.1146/annurev-genet-112414-055145

Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, Rozen SG. 2012.
Primer3-new capabilities and interfaces. *Nucleic Acids Res* **40**. doi:10.1093/nar/gks596

Van Nostrand EL, Freese P, Pratt GA, Wang X, Wei X, Xiao R, Blue SM, Chen JY, Cody NAL,
Dominguez D, Olson S, Sundararaman B, Zhan L, Bazile C, Bouvrette LPB, Bergalet J, Duff
MO, Garcia KE, Gelboin-Burkhart C, Hochman M, Lambert NJ, Li H, McGurk MP, Nguyen TB,
Palden T, Rabano I, Sathe S, Stanton R, Su A, Wang R, Yee BA, Zhou B, Louie AL, Aigner S,
Fu XD, Lécuyer E, Burge CB, Graveley BR, Yeo GW. 2020. A large-scale binding and
functional map of human RNA-binding proteins. *Nature* **583**. doi:10.1038/s41586-020-2077-3

Vaquero-Garcia J, Barrera A, Gazzara MR, Gonzalez-Vallinas J, Lahens NF, Hogenesch JB,
Lynch KW, Barash Y. 2016. A new view of transcriptome complexity and regulation through
the lens of local splicing variations. *Elife* **5**. doi:10.7554/eLife.11752

Venables JP, Elliott DJ, Makarova O V, Makarov EM, Cooke HJ, Eperon IC. 2000. RBMY, a
probable human spermatogenesis factor, and other hnRNP G proteins interact with Tra2beta
and affect splicing. *Hum Mol Genet* **9**:685–694. doi:10.1093/hmg/9.5.685

Wang PJ. 2004. X chromosomes, retrogenes and their role in male reproduction. *Trends*
Endocrinol Metab. doi:10.1016/j.tem.2004.01.007

Wickham H. 2016. ggplot2 Elegant Graphics for Data Analysis (Use R!), Springer.
doi:10.1007/978-0-387-98141-3

Yu G, Wang LG, Han Y, He QY. 2012. ClusterProfiler: An R package for comparing biological
themes among gene clusters. *Omi A J Integr Biol* **16**. doi:10.1089/omi.2011.0118

Yu G, Wang LG, He QY. 2015. ChIP seeker: An R/Bioconductor package for ChIP peak

annotation, comparison and visualization. *Bioinformatics* **31**.

doi:10.1093/bioinformatics/btv145

Zarnegar BJ, Flynn RA, Shen Y, Do BT, Chang HY, Khavari PA. 2016. IrCLIP platform for efficient

characterization of protein-RNA interactions. *Nat Methods* **13**. doi:10.1038/nmeth.3840

Zheng T, Zhou H, Li X, Peng D, Yang Y, Zeng Y, Liu H, Ren J, Zhao Y. 2020. RBMX is required for

activation of ATR on repetitive DNAs to maintain genome stability. *Cell Death Differ*.

doi:10.1038/s41418-020-0570-8

Zhou KI, Shi H, Lyu R, Wylder AC, Matuszek Ż, Pan JN, He C, Parisien M, Pan T. 2019.

Regulation of Co-transcriptional Pre-mRNA Splicing by m6A through the Low-Complexity

Protein hnRNPG. *Mol Cell* **76**:70-81.e9. doi:10.1016/j.molcel.2019.07.005

13
14
15