1

Auxin Exposure Disrupts Feeding Behavior and Fatty Acid

Metabolism in Adult *Drosophila*

Sophie A. Fleck[1], Puja Biswas[2], Emily D. DeWitt[1], Rebecca L. Knuteson[1], Robert

C. Eisman[1], Travis Nemkov[3], Angelo D'Alessandro[3], Jason M. Tennessen[1],

Elizabeth J. Rideout[2], and Lesley N. Weaver[1]*

[1]Department of Biology, Indiana University, Bloomington, IN 47405, USA.

[2]Department of Cellular and Physiological Sciences, Life Sciences Institute, The

University of British Columbia, Vancouver, BC, V6T 1Z3, Canada.

[3]Department of Biochemistry and Molecular Genetics, Anschutz Medical Campus,

University of Colorado School of Medicine, Aurora, CO, 80045, USA.

Running title: [Auxin disruption of triglycerides]

*Corresponding author. Email: lnweaver@iu.edu

Keywords: auxin, *Drosophila*, toxicology, metabolism, oogenesis, triglyceride

ABSTRACT

The ease of genetic manipulation in *Drosophila melanogaster* using the

Gal4/UAS system has been beneficial in addressing key biological questions. Current

modifications of this methodology to temporally induce transgene expression require

temperature changes or exposure to exogenous compounds, both of which have been

shown to have detrimental effects on physiological processes. The recently described

auxin-inducible gene expression system (AGES) utilizes the plant hormone auxin to

induce transgene expression and is proposed to be the least toxic compound for genetic

manipulation, with no obvious effects on *Drosophila* development and survival in one

wild-type strain. Here we show that auxin delays larval development in another widely-

used fly strain, and that short- and long-term auxin exposure in adult *Drosophila* induces

observable changes in physiology and feeding behavior. We further reveal a dosage

response to adult survival upon auxin exposure, and that the recommended auxin

concentration for AGES alters feeding activity. Furthermore, auxin fed male and female

flies exhibit a significant decrease in triglyceride levels and display altered transcription

of fatty acid metabolism genes. Although fatty acid metabolism is disrupted, auxin does

not significantly impact adult female fecundity or progeny survival, suggesting AGES

may be an ideal methodology for studying limited biological processes. These results

emphasize that experiments using temporal binary systems must be carefully designed

and controlled to avoid confounding effects and misinterpretation of results.

INTRODUCTION

The intricate dissection of cell-type specific processes in *Drosophila* is largely

dependent on the yeast-derived *Gal4/UAS* binary system, which allows manipulation of

biological pathways in a spatial and temporal manner (BRAND AND PERRIMON 1993). This

methodology utilizes the Gal4 transcription factor that is under control of a tissue-

specific promoter to induce transgene expression downstream of an U̲pstream

A̲ctivating S̲equence (UAS). Conditional control of gene expression using changes in

temperature (MCGUIRE *et al.* 2003) or feeding of small molecules (ROMAN *et al.* 2001;

POTTER *et al.* 2010; MCCLURE *et al.* 2022) restricts Gal4 activity to specific

developmental timepoints. For example, use of a temperature sensitive *Gal80* mutant

transgene [*Gal80^{ts}*, an inhibitor of Gal4 (DOUGLAS AND HAWTHORNE 1966)] allows for

temporal control of Gal4 activity with a simple shift to the *Gal80^{ts}* restrictive temperature

[29°C, (MCGUIRE *et al.* 2003)]. Conversely, drug-inducible systems such as *GeneSwitch*

and the *Q-system* control temporal and reversible transgene expression with

administration of RU486 or quinic acid, respectively, without the need to rear flies at the

Gal80^{ts} permissive temperature (18°C). These modifications to the *Gal4/UAS* system

have improved the capability to characterize the roles of essential biological pathways in

a tissue-specific manner while avoiding lethality at key developmental stages.

Despite these advancements, each methodology has caveats that must be

considered. For example, rearing flies containing *Gal80^{ts}* at 18°C nearly doubles the

developmental time from egg to adult (POWSNER 1935). In addition, the relatively high

restrictive temperature needed to inactivate Gal80^{ts} has adverse effects on physiological

processes including circadian rhythm (PARISKY *et al.* 2016), aging (MIQUEL *et al.* 1976),

and progeny survival (GANDARA AND DRUMMOND-BARBOSA 2022). Similarly, use of

RU486 has been demonstrated to repress muscle-specific mitochondrial genes

(ROBLES-MURGUIA *et al.* 2019) and lipogenesis (MA *et al.* 2021), among other defects

(LANDIS *et al.* 2015; YAMADA *et al.* 2017), making the *GeneSwitch* system non-ideal for

certain experiments. Although designed to provide flexibility in experimental design and

remove temperature-related defects, the alterations in physiology, behavior, and

lifespan [some of which are not shared between the sexes (LANDIS *et al.* 2015)] due to

RU486 feeding impose difficulties in data interpretation.

The <u>a</u>uxin-inducible <u>g</u>ene <u>e</u>xpression <u>s</u>ystem (AGES) was recently developed as

an alternative method to induce transgene expression and is compatible with the

breadth of *Gal4* transgenic lines available (MCCLURE *et al.* 2022). In this system, *Gal80*

is fused to auxin-inducible degron tags that target Gal80 for degradation, allowing for

Gal4-mediated transgene induction upon auxin consumption. This system poses

substantial advantages. For example, flies can be reared at the optimal temperature for

development (25^{o}C) and transgene expression is strictly induced with supplementation

of auxin to the media. In addition, both control and experimental animals are genetically

identical (similarly to the *GeneSwitch* and *Q-system*), thus minimizing differences that

may arise due to genetic variation.

However, there is emerging evidence that insects can synthesize auxin

(YOKOYAMA *et al.* 2017; TOKUDA *et al.* 2022), suggesting that there may be important

biological roles for this hormone in *Drosophila*. While exposure to 10 mM 1-

naphthaleneacetic acid (the widely employed synthetic hormone in the auxin family and

hereafter referred to as "auxin") has been reported to have no effect on insect

development, survival, or movement in a wild-type *Drosophila* strain (MCCLURE *et al.*

2022), it remains unclear whether auxin exposure affects development in other

commonly used *Drosophila* genetic background strains. It is also unknown whether

increased auxin exposure in adult *Drosophila* results in subtle defects in physiological

processes that may confound experimental interpretations. In this study we sought to

test whether auxin affects larval development in additional strains, and to determine

whether auxin exposure in adults leads to defects in metabolism, the transcriptome, and

oogenesis. We found that recommended concentrations of auxin for AGES disrupt

feeding behavior, whereas increasing levels of auxin results in physiological changes

and lethality. Additionally, we found that auxin exposure delays larval development,

decreases triglyceride levels and alters the transcriptomic profile of fatty acid

metabolism genes in both sexes. Finally, despite the decrease in circulating lipids, auxin

does not severely disrupt processes of oogenesis or progeny survival, suggesting

AGES may be an appropriate method to use for some studies. Our results highlight

changes in development and physiology that should be considered when using auxin to

manipulate gene and protein expression in larval and adult *Drosophila,* as well as other

factors researchers should account for in experimental design using temporal control of

the *Gal4/UAS* system.

MATERIALS AND METHODS

Drosophila strains and culture

Drosophila stocks were maintained on Bloomington *Drosophila* Stock Center (BDSC) Cornmeal Food that consists of 15.9 g/L inactive yeast, 9.2 g/L soy flour, 67 g/L yellow cornmeal, 5.3 g/L agar, 70.6 g/L light corn syrup, 0.059 M propionic acid at 22-25^{o}C. Medium was supplemented with inactive wet yeast paste for all experiments, unless otherwise noted. For a subset of experiments, where indicated, flies were reared on yeast-sugar-cornmeal food (LEWIS 1960) that consists of 20.5 g/L white sugar, 70.9 g/L D-glucose, 48.5 g/L cornmeal, 30.3 g/L yeast, 0.5 g/L $CaCl_2$-$2H_2O$, 0.5 g/L $MgSO_4$-$7H_2O$, 4.6 g/L agar, 4.9 ml/L propionic acid, 0.49 ml/L phosphoric acid. For experiments using this diet, the food was not supplemented with inactive wet yeast paste. The previously described *w^{1118}; tubP-TIR1-T2A-Gal80.AID* ["*AGES*", (MCCLURE *et al.* 2022)] line was used and obtained from the Bloomington *Drosophila* Stock Center (BDSC 92470; https://bdsc.indiana.edu). *Oregon-R* (BDSC 25211) and *y^1 w^{1118}; VK00040/TM6B* (BDSC 9755) lines were used as controls. Flies from the *w^{1118}* strain (BDSC 3605) were used to monitor larval development and adult triglyceride levels. Balancer chromosomes and other genetic elements are described in Flybase (www.flybase.org).

For all ovarian experiments, 0-to-2-day-old females were mated with *AGES* males and incubated at 25^{o}C for up to 15 days at ≥70% humidity on medium containing either 1 mM NaOH [0 mM 1-naphthaleneacetic acid (referred to as "auxin" throughout the manuscript)] or 10 mM auxin dissolved in 1 mM NaOH [the recommended

concentration of auxin (McClure *et al.* 2022)]. Medium was supplemented with inactive

wet yeast containing either 1 mM NaOH or 10 mM auxin daily, except where noted.

Larval development time

w^{1118} females laid eggs on grape plates supplemented with yeast paste during a

3-hr collection period. Newly-hatched larvae were transferred to the yeast-sugar-

cornmeal diet at a density of 50 larvae per 10 ml food. For the auxin-containing medium,

auxin was added to the recommended concentration of 5 mM to cooled fly food

immediately prior to filling vials (McClure *et al.* 2022). Percent pupation was calculated

by comparing the number of pupae at each 12 hr interval to the total pupae in the vial.

Auxin exposure dose response curves

Dose response curves were performed as previously described (Holsopple *et al.*

2023). Briefly, zero-to-two-day old flies were transferred to bottles containing fresh

BDSC food and aged for two days. Flies were sorted by sex in groups of 20 per vial and

aged for an additional 48 hours at 25°C to allow recovery from CO_2 anesthesia. Flies

were then transferred to starvation vials containing sterile milli-Q water for 16 hours at

25°C. Following starvation, flies were transferred to exposure vials containing liquid food

[4% sucrose (m/v), 1.5% yeast extract (m/v), 1 mM NaOH] and 0, 2, 4, 6, 10, 15, 20, 25,

90, 125, or 350 mM auxin. The number of dead female or male flies per vial were

counted in six replicates per concentration and the percentage of dead flies was

subjected to mathematical modeling through the Benchmark Dose Software (BMDS)

published by the U.S. Environmental Protection Agency (EPA;

https://www.epa.gov/bmds).

Fly Liquid-Food Interaction Counter (FLIC) assay

The Fly Liquid-Food Interaction Counter (FLIC) system was used to determine

differences in feeding behavior in male and female flies exposed to food with or without

auxin as previously described (Ro *et al.* 2014). FLIC *Drosophila* Feeding Monitors

(DFMs, Sable Systems International, models DFMV2 and DFMV3) were used in the

single choice configuration and each chamber was loaded with 0 mM auxin liquid food

solution [4% sucrose (m/v), 1.5% yeast extract (m/v), 1 mM NaOH] or the

recommended dose of 10 mM auxin liquid food [4% sucrose (m/v), 1.5% yeast extract

(m/v), 10 mM auxin]. Four-to-six-day old flies were briefly anesthetized and aspirated

into the DFM chambers. Feeding behavior was measured for 24 hours. Each FLIC

experiment contains pooled data from at least 30 flies for each genotype, sex, and auxin

concentration condition. FLIC data were analyzed using previously described custom R

code (Ro *et al.* 2014), which is available at

https://github.com/PletcherLab/FLIC_R_Code. Default thresholds were used for

analysis except for the following: minimum feeding threshold = 10, tasting threshold =

(0,10). Animals that did not participate (i.e., returned zero values), whose DFM returned

an unstable baseline signal, or who produced extreme outliers (i.e., exceeding twice the

mean of the population) were excluded from analysis. Data were subjected to a Mann-

Whitney *U*-test.

173 Ultra High-pressure Liquid Chromatography-Mass Spectrometry (UHPLC-MS)-

174 based metabolomics and analysis

Adult male and female flies were exposed to 0 mM or the recommended dose of

10 mM auxin as described above for the dose response curve assays and were flash

frozen in liquid nitrogen. Analyses were performed at the University of Colorado

Anschutz Medical Campus, as previously described (NEMKOV *et al.* 2019; NEMKOV *et al.*

2022) with minor modifications. Briefly, the analytical platform employs a Vanquish

UHPLC system (ThermoFisher Scientific) coupled online to a Q Exactive mass

spectrometer (ThermoFisher Scientific). The (semi)polar extracts were resolved over a

Kinetex C18 column, 2.1 x 30 mm, 1.7 µm particle size (Phenomenex) using a high-

throughput 1 minute gradient. Solvents were supplemented with 0.1% formic acid for

positive mode runs and 10 mM ammonium acetate +0.1% ammonium hydroxide for

negative mode runs. The Q Exactive mass spectrometer (ThermoFisher Scientific) was

operated independently in positive or negative ion mode, scanning in Full MS mode (2

µscans) from 60 to 900 m/z at 70,000 resolution, with 4 kV spray voltage, 45 sheath

gas, 15 auxiliary gas. Calibration was performed prior to analysis using the Pierce^{TM}

Positive and Negative Ion Calibration Solutions (ThermoFisher Scientific). Metabolomics

raw data were processed using El-Maven (AGRAWAL *et al.* 2019) and analyzed using

Metaboanalyst 5.0 (PANG *et al.* 2021), with the data first preprocessed using log

normalization and Pareto scaling.

Triglyceride assays

Zero-to-two-day old adult males and females were collected and maintained on

BDSC food for 2 days. Two-to-four-day old adults were starved for 16 hours and then

exposed to 0 mM or the recommended dose of 10 mM auxin liquid food for 48 hours as

described above for the dose response assays. Whole bodies from five animals of each

sex and genotype were washed in phosphate-buffered saline (PBS), pH 7.0 and flash

frozen in liquid nitrogen. Samples were homogenized in 100 µl cold PBS + 0.05%

Tween 20 (PBST) and heat-treated for 10 minutes at 90°C. The resulting homogenate

was assayed for triglyceride (TAG) and soluble protein levels as previously described

(TENNESSEN *et al.* 2014). TAG amounts were normalized to protein amounts and

expressed as µg/ml TAG per µg/ml protein. Data were subjected to a paired Student's *t*-

test.

For TAG assays on w^{1118} flies, newly hatched larvae were transferred to yeast-

sugar-cornmeal food and reared at a density of 50 larvae per 10 ml food at 25°C. Male

and female pupae were separated as late pupae according to sex combs. Two

experimental designs were used. For one design, virgin male and female flies were kept

at a density of 20 flies per 10 ml food (+/- recommended dose of 10 mM auxin) from

eclosion until 5 days of age. Five-day-old male and female flies were collected, snap

frozen at -80°C, weighed, and finally subjected to a TAG assay. In the second

experimental design, newly-eclosed flies were subjected to one of three protocols: (1)

maintained continuously on yeast-sugar-cornmeal food with no auxin, where flies were

collected, frozen, weighed, and subjected to a TAG assay at 5 and 10 days of age; (2)

maintained on yeast-sugar-cornmeal food with 10 mM auxin for 5 days and shifted to

food with no auxin for a further 5 days, where flies were collected, frozen, weighed, and

subjected to a TAG assay at 5 and 10 days of age; and (3) maintained continuously on

yeast-sugar-cornmeal food with the recommended 10 mM dose of auxin, with flies

collected, frozen, weighed, and subjected to a TAG assay at 5 and 10 days of age. Flies

were flipped every two days in both experimental designs. One biological replicate of

three or five flies was homogenized in 150 µl or 350 µl of 0.1% Tween in 1X phosphate-

buffered saline (PBS) using 50 µl of glass beads agitated at 8 m/s for 5 sec. TAG assay

was performed using the Stanbio Triglyceride Liquicolor kit (#SB2100430) according to

manufacturer's protocol. TAG is expressed as percent body fat as previously described

(WAT *et al.* 2020). Each experiment includes four biological replicates, and each

experiment was repeated twice for a total of eight biological replicates per sex; data

were analyzed using either a Student's *t*-test, one-way ANOVA, or two-way ANOVA, as

indicated.

RNA isolation, RNA sequencing, and data analysis

Twenty whole adult animals of each genotype and sex were exposed to 0 mM or

the recommended dose of 10 mM auxin, as described above for the dose response

curves, and flash frozen in liquid nitrogen. Tissue was lysed in 500 µl lysis buffer from

the RNAqueous-4PCR DNA-free RNA isolation for RT-PCR kit (Ambion). RNA was

extracted from all samples following manufacturer's instructions. Three independent

experiments were performed for RNA sequencing.

cDNA library construction, Illumina sequencing, and differential expression

analysis was performed by Novogene Bioinformatics Technology Co., Ltd (Beijing,

China). The cDNA libraries were prepared using the NEBNext Ultra RNA Library Prep

Kit for Illumina (New England Biolabs) according to the manufacturer's instructions. The

cDNA library for each sample was quality assessed using an Agilent Bioanalyzer 2100,

and library preparations were sequenced on a NovaSeq6000 platform with PE150 read

lengths.

Reads obtained from sequencing were aligned to the *D. melanogaster* reference

genome using the TopHat read alignment tool (TRAPNELL *et al.* 2009) for each of the

sequencing datasets. The reference sequences were downloaded from the Ensembl

project website (useast.ensembl.org). TopHat alignments were used to generate read

counts for each gene using HTSeq (ANDERS *et al.* 2015), which were subsequently used

to generate the differential expression results using the DESeq2 R package (ANDERS *et*

al. 2015). RNA sequencing produced an average of 40,448,108 reads across the 36

sequencing libraries, ranging from 37,632,234 to 58,099,256 reads per sample

(representing an average of 96.2% mapped to the *Drosophila* genome). Enriched genes

with a corrected *P* value less than 0.05 were considered significant.

cDNA synthesis and quantitative reverse-transcriptase polymerase chain reaction

(qRT-PCR)

cDNA was synthesized from 500 ng of total RNA described above for each

sample using Superscript II Reverse Transcriptase (Thermo Fisher Scientific) according

to the manufacturer's instructions. PowerUp SYBR Green Master Mix (Thermo Fisher

Scientific) was used for RT-qPCR. The reactions for three independent biological

replicates were performed in triplicate using LightCycler 96 (Roche). Amplification

fluorescence threshold was determined by LightCycler 96 software, and ddCT were

calculated using Microsoft Excel. Fold change of transcript levels was calculated in

Excel as described (TAYLOR *et al.* 2019). The primers used for all PCR reactions are

listed in Table S8. *Rp49* and *Act5C* transcript levels were used as references.

Egg laying and hatching assays

Egg production was measured as previously described (WEAVER AND DRUMMOND-

BARBOSA 2019) by maintaining five experimental females mated with five *AGES* males

in perforated plastic bottles capped with molasses/agar plates smeared with 0 mM auxin

or the recommended concentration of 10 mM auxin inactive yeast paste. Molasses/agar

plates were changed twice daily. The number of eggs laid per day was counted in five

replicates per genotype and results were subjected to a paired Student's *t*-test.

Egg hatching was measured by transferring up to 30 eggs from molasses/agar

plates to fresh molasses/agar plates containing 0 mM auxin inactive yeast paste in the

center every 2 days and the number of eggs that has hatched were counted 24 hours

after the transfer. The number of eggs hatched per day was counted in five replicates

per genotype and results were subjected to a paired Student's *t*-test.

Adult female ovary immunostaining and fluorescence microscopy

Ovaries and carcasses were dissected in Grace's Insect Medium (Gibco), fixed,

and washed as previously described (WEAVER AND DRUMMOND-BARBOSA 2019). Samples

were blocked for at least 3 hours in 5% normal goat serum (NGS; Jackson

ImmunoResearch) and 5% bovine serum albumin (BSA; Sigma) in phosphate-buffered

saline [PBS; 10 mM NaH_2PO_4/$NaHPO_4$, 175 mM NaCl (pH 7.4)] containing 0.1% Triton

X-100 (PBST). Samples were incubated overnight at 4°C in primary antibodies diluted in

blocking solution as follows: mouse monoclonal anti-alpha-spectrin (Developmental

Studies Hybridoma Bank; DSHB, 3 µg/ml), mouse monoclonal anti-Lamin C (DSHB, 0.8

µg/ml), and rat monoclonal anti-Vasa (DSHB, 2.15 µg/ml). Samples were washed in

PBST and incubated at room temperature for 2 hours with 1:200 Alexa Fluor 488- or

568-conjugated goat-species specific secondary antibodies (Molecular Probes) in

blocking solution. Samples were then washed three times for 15 minutes and mounted

in Vectashield containing 1.5 µg/ml 4',6-diamidino-2-phenylindole (DAPI; Vector

Laboratories) and imaged using a Leica SP8 confocal.

Cap cells and GSCs were identified as described (WEAVER AND DRUMMOND-

BARBOSA 2019), and two-way ANOVA with interaction (GraphPad Prism) was used to

calculate the statistical significance of any differences among genotypes in the rate of

cap cell or GSC loss from at least three independent experiments, as described

(ARMSTRONG *et al.* 2014). Progression through vitellogenesis was assessed using DAPI

staining, as described (WEAVER AND DRUMMOND-BARBOSA 2019). Three independent

experiments were performed and subjected to a Student's *t*-test for statistical analysis.

ApopTag assays

To detect dying germline cysts, the ApopTag Indirect *In Situ* Apoptosis Detection

Kit (Millipore Sigma) was used according to the manufacturer's instructions as

previously described (WEAVER AND DRUMMOND-BARBOSA 2019). Briefly, fixed and teased

ovaries were rinsed in equilibration buffer twice for 5 minutes each at room temperature.

Samples were incubated in 100 µl TdT solution at 37°C for 1 hour with mixing at 15-

minute intervals. Ovaries were washed three times in 1X PBS followed by incubation in

anti-digoxigenin conjugate for 30 min at room temperature protected from light. Samples

were washed four times in 1X PBS and processed for immunofluorescence as

described above.

RESULTS AND DISCUSSION

Adult *Drosophila* males have increased sensitivity to auxin compared to females.

We sought to utilize the AGES expression system in our laboratory for controlling

gene expression and began by determining whether we could recapitulate Gal4

expression at levels like that of *Gal80^{ts}*. We recombined the auxin-inducible degron line

("*AGES*") with *3.1Lsp2-Gal4* (*3.1Lsp2-Gal4^{AGES}*) to drive the expression of *UAS-*

nucGFP in adult female adipocytes compared to the previously described *3.1Lsp2-Gal4*

recombined with *Gal80^{ts}* (*3.1Lsp2Gal4^{ts}*; ARMSTRONG *et al.* 2014). Zero-to-two-day old

females of each genotype were fed inactive yeast paste for two days (pre-treatment)

prior to Gal4-induction. Gal4 expression was induced in females carrying the *AGES*

transgene by feeding solid food supplemented with inactive wet yeast paste containing

0 mM to the recommended dose of 10 mM auxin (inactive yeast was used to prevent

auxin metabolism by live yeast) for two days; whereas females carrying the *Gal80^{ts}*

transgene were shifted from 18°C to 29°C for two days and the relative expression of

GFP transcripts were measured (**Figure S1A**). There was minimal "leaky" Gal4

expression with *3.1Lsp2-Gal4^{AGES}* fed food without auxin (0 mM); whereas 5 mM and 10

mM auxin induced *nucGFP* expression at levels similar to that of *3.1Lsp2-Gal4^{ts}*.

Furthermore, we found that removal of auxin using *3.1Lsp2-Gal4^{AGES}* inactivated Gal4

activity at a faster rate than *3.1Lsp2-Gal4^{ts}* (shifted back to 18°C; **Figure S1B**).

Therefore, Gal4 expression using AGES can be recapitulated as previously described

(MCCLURE *et al.* 2022) and is faster at repressing transgene activation compared to

Gal80^{ts}.

Because the wild-type strain *Canton-S* was previously used to monitor

development and survival in flies reared on auxin-containing food (MCCLURE *et al.*

2022), we monitored larval development in flies from the *w^{1118}* genotype, a widely-used

genetic background strain in *Drosophila* biology. When we measured the time between

egg-laying and pupariation in a mixed-sex group of larvae reared on yeast-sugar-

cornmeal food supplemented with (5 mM) or without (0 mM) auxin, we found that the

time to pupariation was longer in auxin-fed larvae than in larvae raised without auxin

(**Figure S1C**). This suggests that the recommended dose of auxin delays larval

development in at least one widely-used *Drosophila* strain, indicating that genetic

background is an important consideration when using the AGES system for studies at

the larval stage of development.

Given these larval phenotypes, we next wanted to determine whether auxin

exposure induces physiological changes in adults. Because genetic variation can

influence physiology (SHORTER *et al.* 2015; EVANGELOU *et al.* 2019; DAMSCHRODER *et al.*

2020), we used multiple strains to test the effects of auxin on physiology. We first

compared the *AGES* line and the strain into which the *AGES* transgene was introduced

(*VK00040*) in our experiments. We also used *Oregon-R* [the wild-type strain used for

the ModENCODE project (CONSORTIUM *et al.* 2010)] as an additional control. We

incubated four-to-six-day old adult males and females of each genotype in vials

supplemented with chromatography paper soaked with liquid food containing 0 mM to

350 mM auxin and measured lethality after 48 hours of exposure (**Figure 1A-C; Figure**

S2A). Compared to the 0 mM control, both males and females showed increased

susceptibility to moderate concentrations of auxin (20-30 mM; **Figure 1B,C; Figure**

S2A). In addition, we observed sex differences in sensitivity to auxin exposure, with

females able to tolerate higher concentrations relative to males. For example, the

benchmark dose (BMD; the concentration that produces a change in the response) in

males was lower than that of females in each genotype (**Table 1**). Interestingly, both

AGES males and females had a higher BMD upon auxin exposure relative to the

VK00040 injection and *Oregon-R* lines, suggesting that the *AGES* transgene may

confer resistance to auxin toxicity. These results suggest that exposure to auxin in both

males and females results in lethality, where males have a higher susceptibility to auxin

exposure compared to females.

For the remainder of our analyses we used the recommended working

concentration of auxin for AGES (10 mM), which as previously described, did not cause

lethality in the *AGES* line for either sex [**Figure 1B,C**; **Table 1**; (MCCLURE *et al.* 2022)].

To determine whether feeding behavior could account for sex differences in the survival

response to auxin, we analyzed the food consumption of adult male and female flies

exposed to 0 mM or 10 mM auxin using the Fly Liquid-Food Interaction Counter [FLIC,

(RO *et al.* 2014)] over a 24-hour period (**Figure 1D-G, Figure S2B-E**). In both feeding

conditions, females exhibited more feeding events relative to males in all tested

genotypes (**Table 2**). For example, females fed 0 mM auxin in both the *VK00040* and

AGES lines had significantly more engagements with the interaction counter compared

to *VK00040* and *AGES* males (compare **Figure 1D** to **Figure 1F**). In addition, genetic

background confers differences in feeding behavior as seen by the difference between

VK00040 and *AGES*. Surprisingly, compared to the 0 mM food, females had fewer

interactions and slightly decreased event durations when fed the 10 mM auxin food;

whereas males only decreased the amount of time they interacted with the food when

exposed to 10 mM auxin (**Figure 1G**). These results are consistent with previous

reports that females eat more than males on normal food diets with yeast (WONG *et al.*

2009). However, our results also suggest that females exposed to auxin may have an

increased survival due to fewer interactions with the food relative to adult males,

resulting in differential auxin sensitivity. Therefore, future studies should investigate the

female-specific avoidance of auxin, since decreased feeding behavior could confound

interpretation of results.

Fatty acid metabolism is decreased in adults exposed to auxin.

To determine whether auxin exposure altered metabolism in adult *Drosophila*, we

performed metabolomics to compare adult males and females of each genotype

exposed to 10 mM auxin relative to the 0 mM control (**Table S1**). Partial Least Squares

Discriminant Analysis (PLS-DA) showed that each genotype clustered in distinct groups

(**Figure S3**). Notably, 0 mM controls clustered separately from 10 mM auxin samples in

all genotypes for both sexes along the Component 1 axis, which describes 33.4% or

34.4% of the variance for females and males, respectively. We further analyzed the

data using MetaboAnalyst (PANG *et al.* 2021), which revealed that the levels of

acylcarnitine and fatty acid metabolites were the most significantly decreased

metabolites in females exposed to 10 mM auxin regardless of genotype (**Figure 2A;**

Figures S4-6). Adult males exposed to 10 mM auxin also exhibited decreased fatty acid

and acylcarnitine metabolites (**Figure 2B**); however, amino acid levels were the most

significantly altered metabolites in males in response to auxin (**Figures S4-6**).

To confirm that fatty acid metabolism is altered in response to auxin, we exposed

adult males and females of each genotype to 0 mM or 10 mM auxin and measured the

levels of triacylglycerol (TAG) after 48 hours of exposure. Both males and females

exposed to 10 mM auxin of each genotype had significantly less TAG relative to the 0

mM control (**Figure 2C,D; Figure S4G**). We note that the lack of sex difference in lipid

levels is likely due to exposing adults to auxin prior to the onset of male-female

differences in TAG (WAT *et al.* 2020). To determine whether auxin influences TAG in

additional contexts, we transferred newly-eclosed virgin w^{1118} males and females to a

yeast-sugar-cornmeal diet supplemented with either 0 mM or the recommended dose of

10 mM auxin and measured TAG levels after 5 days. We found a significant decrease in

whole-body fat storage in both male and female flies exposed to 10 mM auxin

compared with flies transferred to food with 0 mM auxin (**Figure S4H**); however, the

magnitude of the auxin-induced decrease in body fat was greater in females than in

males (sex:diet interaction $p = 0.001$; two-way ANOVA). This suggests that auxin has a

stronger effect on female TAG levels, in line with our data showing a female-biased

reduction in food interactions on auxin-supplemented medium.

We next asked whether this effect on TAG levels was reversible. We exposed 0-

day-old male and female flies to diets with 0 mM or 10 mM auxin for five days, and then

measured TAG levels after shifting the flies to food supplemented with no auxin for five

additional days. We found that females but not males showed a strong trend toward

recovery of whole-body TAG levels after shifting the flies back to food with no auxin

(Figure S4I), suggesting the effect of auxin feeding on body fat were reversible only in

females. Furthermore, we found that flies exposed to food supplemented with 10 mM

auxin for 10 days show no additional reduction in whole-body TAG levels compared with

flies exposed to auxin for 5 days in either sex (Figure S4I). Taken together, these

results suggest that auxin exposure disrupts fatty acid metabolism, resulting in

decreased circulating lipids in adult males and females. While these changes to fat

metabolism are partially reversible after withdrawal of auxin supplementation in females,

changes to whole-body TAG levels persisted even after auxin withdrawal in males.

Auxin exposure induces global transcriptomic changes in adult *Drosophila*.

To determine whether auxin exposure induced changes in gene expression, we

exposed adult males and females of each genotype to 0 mM and 10 mM auxin for 48

hours and performed RNA sequencing analysis of whole animals. In both males and

females of each tested genotype, at least 150 transcripts were significantly altered in

response to auxin exposure (**Figures 3A-D and S7-8; Tables S2-7**). Genes involved in

drug metabolism [e.g., glutathione-S-transferases (GSTs) and uridine diphosphate-

glucuronosyltransferases (UGTs), phase II enzymes required to increase hydrophobicity

of compounds (YU 2008)] were significantly up-regulated in response to auxin exposure

in both males and females of all genotypes (**Figure 3E; Figures S7-8**), suggesting that

auxin induces a xenobiotic response at this concentration (YU 2008). Surprisingly,

genes involved in fatty acid metabolism (e.g., Fad2, which encodes a desaturase) were

also significantly up-regulated; whereas lipases (enzymes that break down fatty acids)

were significantly down-regulated in each genotype and sex (**Figure 3F; Figure S7D;**

Figure S8C,D). Notably, the transcription of enzymes associated with lipolysis such as

the lipase *brummer* (GRONKE *et al.* 2005) or the perilipins *Lsd-1* and *Lsd-2* (BELLER *et al.*

2010) were unaltered in response to auxin exposure (**Tables S2-7**). Therefore, it is

possible that in response to auxin, triglycerides are mobilized by an unknown

mechanism resulting in upregulation of fatty acid metabolism and attenuation of lipid

breakdown. It is also possible that decreased interactions or feeding time with auxin-

containing food promotes fasting, resulting in decreased TAG and lipid synthesis.

Collectively, these results suggest that auxin exposure in adults significantly alters the

transcriptome to adjust to decreased fatty acid metabolism and lipid stores.

Auxin is an essential hormone for plant growth and development (DU *et al.* 2020;

WOJCIK *et al.* 2020; GOMES AND SCORTECCI 2021), and has been widely used to

manipulate gene expression in multiple organisms (ZHANG *et al.* 2015; TROST *et al.*

2016; CHEN *et al.* 2018; LI *et al.* 2019; SHETTY *et al.* 2019; YESBOLATOVA *et al.* 2020;

MACDONALD *et al.* 2022). In *Arabidopsis*, auxin is a known regulator of fatty acid

synthesis in plants (HE *et al.* 2020) and has been shown to induce lipid synthesis to

control vacuole trafficking (LI *et al.* 2015). Our results suggest that auxin has the

opposite effect on lipid metabolism in adult *Drosophila* and decreases triglyceride levels.

Given its widespread use, the role of auxin in potentially regulating lipid metabolism

should be further characterized in additional model organisms using auxin to manipulate

gene expression (at their respective recommended dosages), such as *C. elegans*

(ZHANG *et al.* 2015) and mice (MACDONALD *et al.* 2022).

Auxin exposure in adult females does not influence oogenesis.

Although many tissues maintain and regulate lipid stores in *Drosophila* (including

muscle and gut [reviewed in (HEIER AND KUHNLEIN 2018)]), the adult adipose tissue is a

major lipid storage depot [reviewed in (CHATTERJEE AND PERRIMON 2021)] and impacts

peripheral tissue function such as adult *Drosophila* oogenesis (ARMSTRONG *et al.* 2014;

MATSUOKA *et al.* 2017; ARMSTRONG AND DRUMMOND-BARBOSA 2018; WEAVER AND

DRUMMOND-BARBOSA 2018; WEAVER AND DRUMMOND-BARBOSA 2019). Thus, we sought

to examine if the metabolic effects of auxin described above impact oogenesis.

Each ovary consists of 16-20 ovarioles composed of progressively older follicles

that ultimately give rise to a mature egg chamber [**Figure 4A,B**; (DRUMMOND-BARBOSA

2019)]. Oogenesis is maintained by 2-3 germline stem cells (GSCs) that reside in the

anterior germarium of each ovariole and can be identified based on their proximity to the

stem cell niche, which is primarily composed of somatic cap cells. GSCs self-renew and

give rise to early GSC progeny that differentiate to produce follicles that bud from the

germarium and complete oogenesis.

To determine if auxin exposure influences fecundity or progeny survival, we

performed egg laying and larval hatching analyses. Adult females (*Oregon-R*, *VK00040*,

or *AGES*) were maintained with *AGES* males (due to their increased resistance to

auxin) on molasses plates supplemented with wet inactive yeast containing either 0 mM

auxin or 10 mM auxin for 15 days. Relative to 0 mM auxin control, exposure to 10 mM

auxin did not significantly influence the number of eggs laid per female in any tested

genotype (**Figure S9A,C,E**). We note that the number of eggs laid under 0 mM auxin

conditions is relatively low; however, this is likely due to the decrease in egg production

in females fed inactive yeast compared to active yeast paste (**Figure S10**). In addition,

the relative hatching percentages of each genotype also were not affected by auxin

exposure (**Figure S9B,D,F**). These results suggest that auxin exposure in adult females

does not significantly influence the number of laid eggs or oocyte quality.

We next analyzed specific processes of oogenesis that are sensitive to metabolic

changes including GSC maintenance, early germline cyst survival, and survival of

vitellogenic follicles (DRUMMOND-BARBOSA 2019). We detected GSCs by the morphology

and location of the fusome relative to the GSC-niche (**Figure 4C**). GSC numbers in all

genotypes exposed to 10 mM auxin were comparable to those in 0 mM auxin controls at

each time point (**Figure 4D-E; Figure S11A,D**). Likewise, there was no change in the

number of cap cells over time (**Figure S11E**). Furthermore, there were no effects of

auxin exposure on the survival of early germline cysts [as determined by ApopTag

TUNEL labeling (Drummond-Barbosa and Spradling, 2001)] (**Figure 4F-H; Figure**

S11B) or survival of vitellogenic egg chambers (**Figure 4I-K; Figure S11C**). Based on

these data, we conclude that the recommended auxin concentration for AGES in adult

females does not adversely affect oogenesis, despite significant decreases in whole-

body fatty acid metabolism and lipid composition.

Consistent with our results, it was recently shown that obese adult *Drosophila*

females do not have reduced fecundity, but that fertility defects manifest only when

combined with a high sugar diet (NUNES AND DRUMMOND-BARBOSA 2023). Collectively,

these results suggest that lipid content alone (lean or obese) is not sufficient to regulate

distinct processes of oogenesis in adult *Drosophila* females. Therefore, AGES may be

suitable for the study of some processes in adult oogenesis.

Conclusions and suggestions for future studies.

The *Gal4/UAS* system has revolutionized the ability to perform tissue specific

manipulations in *Drosophila*. However, finding the ideal conditions to manipulate tissues

temporally without causing significant alterations in physiology or impacting organism

behavior presents a challenge. For example, using *Gal4/UAS* in conjunction with

Gal80^{ts} for adult specific manipulations has the adverse effect of decreasing adult

female fecundity, making this system unsuitable for aging studies on oogenesis

(GANDARA AND DRUMMOND-BARBOSA 2022). Similarly, controlling transgene expression

using RU486 inducible *GeneSwitch* drivers has not only been found to be less than

ideal for feeding and aging studies (LANDIS *et al.* 2015; YAMADA *et al.* 2017), but also

presents a workplace hazard to pregnant researchers (AVRECH *et al.* 1991). Likewise,

our findings herein indicate that methods requiring auxin for temporal transgene

induction or protein degradation may work well to study many biological processes, but

may not be the ideal system for the study of lipogenesis or other metabolic processes

dependent on lipid content. Indeed, our data suggests that auxin-induced changes to

metabolism persist even after its withdrawal from the diet, indicating that even short-

term auxin treatments may have unwanted physiological effects.

Each modification to the *Gal4/UAS* system designed to provide temporal

regulation of transgene expression has strengths and weaknesses and should be

carefully vetted to ensure the most ideal experimental design is balanced with the

caveats of the methodology. Regardless of the system used, researchers should

determine whether their experimental manipulation alters Gal4 expression patterns

compared to control conditions [i.e., thoroughly analyze Gal4 expression pattern across

developmental time and tissues (WEAVER *et al.* 2020)]. For example, the expression of

transgenes using numerous neuronal Gal4 drivers with AGES is significantly weaker

than Gal4-induced expression at 30ºC (HAWLEY *et al.* 2023). Finally, researchers should

ensure that non-specific transgene controls used in the same conditions as the

experimental (e.g., 0 mM versus 10 mM auxin) do not have phenotypes for

physiological outputs of interest and ensure that changes in physiology that could

confound result interpretations are accounted for.

DATA AVAILABILITY

Drosophila strains can be purchased from the Bloomington *Drosophila* Stock

Center. The data and analyses in this paper are described in the main figures. The raw

data and processed data files are available through the NCBI GEO accession number

GSE237283 and are also provided as supplemental figures and tables. Additional raw

data is available upon request.

ACKNOWLEDGEMENTS

We thank the Bloomington Stock Center (National Institutes of Health

P40OD018537) for *Drosophila* stocks. We are thankful to Flybase (www.flybase.org), an

essential *Drosophila* research resource (NIH 5U41HG000739). The authors would like

to thank the Indiana University Light Microscopy Imaging Center (LMIC) for access to

microscopy facilities. We are grateful for Kristina J. Weaver and Scott D. Pletcher for

assistance with the FLIC code. We are also grateful to Elizabeth T. Ables, Brian R.

Calvi, and Deepika Vasudevan for critical reading of the manuscript. This work was

supported by the Canadian Institutes for Health Research grant PJT-153072 (E.J.R.

and P.B.), the National Institutes of Health (NIH) grants R35 GM119557 (J.M.T), R00

GM127605 (L.N.W.), and R35 GM150517 (L.N.W).

FUNDING

This work was supported by the Canadian Institutes for Health Research grant

PJT-153072 (E.J.R and PB), the National Institutes of Health (NIH) grants R35

GM119557 (J.M.T), R00 GM127605 (L.N.W.), and R35 GM150517 (L.N.W).

CONFLICT OF INTEREST

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

S.A.F., P.B., E.D.D., R.L.K., R.C.E., T.N., A.D'A., and L.N.W. performed experiments,

analyzed, and interpreted the data. L.N.W., P.B., and E.J.R. wrote the manuscript and

J.M.T provided edits.

REFERENCES

Agrawal, S., S. Kumar, R. Sehgal, S. George, R. Gupta *et al.*, 2019 El-MAVEN: A Fast,

589   Robust, and User-Friendly Mass Spectrometry Data Processing Engine for

590   Metabolomics. Methods Mol Biol 1978**:** 301-321.

591 Anders, S., P. T. Pyl and W. Huber, 2015 HTSeq--a Python framework to work with

592   high-throughput sequencing data. Bioinformatics 31**:** 166-169.

593 Armstrong, A. R., and D. Drummond-Barbosa, 2018 Insulin signaling acts in adult

594   adipocytes via GSK-3beta and independently of FOXO to control Drosophila

595   female germline stem cell numbers. Dev Biol 440**:** 31-39.

596 Armstrong, A. R., K. M. Laws and D. Drummond-Barbosa, 2014 Adipocyte amino acid

597   sensing controls adult germline stem cell number via the amino acid response

598   pathway and independently of Target of Rapamycin signaling in Drosophila.

599   Development 141**:** 4479-4488.

600 Avrech, O. M., A. Golan, Z. Weinraub, I. Bukovsky and E. Caspi, 1991 Mifepristone

601   (RU486) alone or in combination with a prostaglandin analogue for termination of

602   early pregnancy: a review. Fertil Steril 56**:** 385-393.

Beller, M., A. V. Bulankina, H. H. Hsiao, H. Urlaub, H. Jackle *et al.*, 2010 PERILIPIN-

dependent control of lipid droplet structure and fat storage in Drosophila. Cell

Metab 12**:** 521-532.

Brand, A. H., and N. Perrimon, 1993 Targeted gene expression as a means of altering

cell fates and generating dominant phenotypes. Development 118**:** 401-415.

Chatterjee, N., and N. Perrimon, 2021 What fuels the fly: Energy metabolism in

Drosophila and its application to the study of obesity and diabetes. Sci Adv 7.

Chen, W., M. Werdann and Y. Zhang, 2018 The auxin-inducible degradation system

enables conditional PERIOD protein depletion in the nervous system of

Drosophila melanogaster. FEBS J 285**:** 4378-4393.

Consortium, m., S. Roy, J. Ernst, P. V. Kharchenko, P. Kheradpour *et al.*, 2010

Identification of functional elements and regulatory circuits by Drosophila

modENCODE. Science 330**:** 1787-1797.

Damschroder, D., K. Richardson, T. Cobb and R. Wessells, 2020 The effects of genetic

background on exercise performance in Drosophila. Fly (Austin) 14**:** 80-92.

Douglas, H. C., and D. C. Hawthorne, 1966 Regulation of genes controlling synthesis of

the galactose pathway enzymes in yeast. Genetics 54**:** 911-916.

Drummond-Barbosa, D., 2019 Local and Physiological Control of Germline Stem Cell

Lineages in Drosophila melanogaster. Genetics 213**:** 9-26.

Du, M., E. P. Spalding and W. M. Gray, 2020 Rapid Auxin-Mediated Cell Expansion.

Annu Rev Plant Biol 71**:** 379-402.

Evangelou, A., A. Ignatiou, C. Antoniou, S. Kalanidou, S. Chatzimatthaiou *et al.*, 2019

Unpredictable Effects of the Genetic Background of Transgenic Lines in

Physiological Quantitative Traits. G3 (Bethesda) 9**:** 3877-3890.

Gandara, A. C. P., and D. Drummond-Barbosa, 2022 Warm and cold temperatures

have distinct germline stem cell lineage effects during Drosophila oogenesis.

Development 149.

Gomes, G. L. B., and K. C. Scortecci, 2021 Auxin and its role in plant development:

structure, signalling, regulation and response mechanisms. Plant Biol (Stuttg) 23**:**

894-904.

Gronke, S., A. Mildner, S. Fellert, N. Tennagels, S. Petry *et al.*, 2005 Brummer lipase is

an evolutionary conserved fat storage regulator in Drosophila. Cell Metab 1**:** 323-

330.

Hawley, H. R., C. J. Roberts and H. L. Fitzsimons, 2023 Comparison of neuronal GAL4

drivers along with the AGES (auxin-inducible gene expression system) and

TARGET (temporal and regional gene expression targeting) systems for fine

tuning of neuronal gene expression in Drosophila. MicroPubl Biol 2023.

He, M., C. X. Qin, X. Wang and N. Z. Ding, 2020 Plant Unsaturated Fatty Acids:

Biosynthesis and Regulation. Front Plant Sci 11**:** 390.

Heier, C., and R. P. Kuhnlein, 2018 Triacylglycerol Metabolism in Drosophila

melanogaster. Genetics 210**:** 1163-1184.

Holsopple, J. M., S. R. Smoot, E. M. Popodi, J. K. Colbourne, J. R. Shaw *et al.*, 2023

Assessment of Chemical Toxicity in Adult Drosophila Melanogaster. J Vis Exp.

Landis, G. N., M. P. Salomon, D. Keroles, N. Brookes, T. Sekimura *et al.*, 2015 The

progesterone antagonist mifepristone/RU486 blocks the negative effect on life

span caused by mating in female Drosophila. Aging (Albany NY) 7**:** 53-69.

Lewis, E. B., 1960 A new standard food medium. Drosophila Information Service**:** 118-

119.

Li, R., R. Sun, G. R. Hicks and N. V. Raikhel, 2015 Arabidopsis ribosomal proteins

control vacuole trafficking and developmental programs through the regulation of

lipid metabolism. Proc Natl Acad Sci U S A 112**:** E89-98.

Li, S., X. Prasanna, V. T. Salo, I. Vattulainen and E. Ikonen, 2019 An efficient auxin-

inducible degron system with low basal degradation in human cells. Nat Methods

16**:** 866-869.

Ma, P., Y. Zhang, Q. Liang, Y. Yin, S. Wang *et al.*, 2021 Mifepristone (RU486) inhibits

dietary lipid digestion by antagonizing the role of glucocorticoid receptor on lipase

transcription. iScience 24**:** 102507.

Macdonald, L., G. C. Taylor, J. M. Brisbane, E. Christodoulou, L. Scott *et al.*, 2022

Rapid and specific degradation of endogenous proteins in mouse models using

auxin-inducible degrons. Elife 11.

Matsuoka, S., A. R. Armstrong, L. L. Sampson, K. M. Laws and D. Drummond-Barbosa,

2017 Adipocyte Metabolic Pathways Regulated by Diet Control the Female

Germline Stem Cell Lineage in Drosophila melanogaster. Genetics 206**:** 953-971.

McClure, C. D., A. Hassan, G. N. Aughey, K. Butt, A. Estacio-Gomez *et al.*, 2022 An

auxin-inducible, GAL4-compatible, gene expression system for Drosophila. Elife

11.

McGuire, S. E., P. T. Le, A. J. Osborn, K. Matsumoto and R. L. Davis, 2003

Spatiotemporal rescue of memory dysfunction in Drosophila. Science 302**:** 1765-

1768.

Miquel, J., P. R. Lundgren, K. G. Bensch and H. Atlan, 1976 Effects of temperature on

the life span, vitality and fine structure of Drosophila melanogaster. Mech Ageing

Dev 5**:** 347-370.

Nemkov, T., J. A. Reisz, S. Gehrke, K. C. Hansen and A. D'Alessandro, 2019 High-

Throughput Metabolomics: Isocratic and Gradient Mass Spectrometry-Based

Methods. Methods Mol Biol 1978**:** 13-26.

Nemkov, T., T. Yoshida, M. Nikulina and A. D'Alessandro, 2022 High-Throughput

Metabolomics Platform for the Rapid Data-Driven Development of Novel Additive

Solutions for Blood Storage. Front Physiol 13**:** 833242.

Nunes, R. D., and D. Drummond-Barbosa, 2023 A high-sugar diet, but not obesity,

reduces female fertility in Drosophila melanogaster. Development 150.

Pang, Z., J. Chong, G. Zhou, D. A. de Lima Morais, L. Chang *et al.*, 2021

MetaboAnalyst 5.0: narrowing the gap between raw spectra and functional

insights. Nucleic Acids Res 49**:** W388-W396.

Parisky, K. M., J. L. Agosto Rivera, N. C. Donelson, S. Kotecha and L. C. Griffith, 2016

Reorganization of Sleep by Temperature in Drosophila Requires Light, the

Homeostat, and the Circadian Clock. Curr Biol 26**:** 882-892.

Potter, C. J., B. Tasic, E. V. Russler, L. Liang and L. Luo, 2010 The Q system: a

repressible binary system for transgene expression, lineage tracing, and mosaic

analysis. Cell 141**:** 536-548.

Powsner, L., 1935 The Effects of Temperature on the Durations of the Developmental

Stages of Drosophila melanogaster. Physiological Zoology 8**:** 474-520.

Ro, J., Z. M. Harvanek and S. D. Pletcher, 2014 FLIC: high-throughput, continuous

analysis of feeding behaviors in Drosophila. PLoS One 9**:** e101107.

Robles-Murguia, M., L. C. Hunt, D. Finkelstein, Y. Fan and F. Demontis, 2019 Tissue-

specific alteration of gene expression and function by RU486 and the

GeneSwitch system. NPJ Aging Mech Dis 5**:** 6.

Roman, G., K. Endo, L. Zong and R. L. Davis, 2001 P[Switch], a system for spatial and

temporal control of gene expression in Drosophila melanogaster. Proc Natl Acad

Sci U S A 98**:** 12602-12607.

Shetty, A., N. I. Reim and F. Winston, 2019 Auxin-Inducible Degron System for

Depletion of Proteins in Saccharomyces cerevisiae. Curr Protoc Mol Biol 128**:**

e104.

Shorter, J., C. Couch, W. Huang, M. A. Carbone, J. Peiffer *et al.*, 2015 Genetic

architecture of natural variation in Drosophila melanogaster aggressive behavior.

Proc Natl Acad Sci U S A 112**:** E3555-3563.

Taylor, S. C., K. Nadeau, M. Abbasi, C. Lachance, M. Nguyen *et al.*, 2019 The Ultimate

qPCR Experiment: Producing Publication Quality, Reproducible Data the First

Time. Trends Biotechnol 37**:** 761-774.

Tennessen, J. M., W. E. Barry, J. Cox and C. S. Thummel, 2014 Methods for studying

metabolism in Drosophila. Methods 68**:** 105-115.

Tokuda, M., Y. Suzuki, S. Fujita, H. Matsuda, S. Adachi-Fukunaga *et al.*, 2022

Terrestrial arthropods broadly possess endogenous phytohormones auxin and

cytokinins. Sci Rep 12**:** 4750.

Trapnell, C., L. Pachter and S. L. Salzberg, 2009 TopHat: discovering splice junctions

with RNA-Seq. Bioinformatics 25**:** 1105-1111.

Trost, M., A. C. Blattner and C. F. Lehner, 2016 Regulated protein depletion by the

auxin-inducible degradation system in Drosophila melanogaster. Fly (Austin) 10**:**

35-46.

Wat, L. W., C. Chao, R. Bartlett, J. L. Buchanan, J. W. Millington *et al.*, 2020 A role for

triglyceride lipase brummer in the regulation of sex differences in Drosophila fat

storage and breakdown. PLoS Biol 18**:** e3000595.

Weaver, L. N., and D. Drummond-Barbosa, 2018 Maintenance of Proper Germline

Stem Cell Number Requires Adipocyte Collagen in Adult Drosophila Females.

Genetics 209**:** 1155-1166.

Weaver, L. N., and D. Drummond-Barbosa, 2019 The nuclear receptor seven up

functions in adipocytes and oenocytes to control distinct steps of Drosophila

oogenesis. Dev Biol 456**:** 179-189.

Weaver, L. N., T. Ma and D. Drummond-Barbosa, 2020 Analysis of Gal4 Expression

Patterns in Adult Drosophila Females. G3 (Bethesda) 10**:** 4147-4158.

Wojcik, A. M., B. Wojcikowska and M. D. Gaj, 2020 Current Perspectives on the Auxin-

Mediated Genetic Network that Controls the Induction of Somatic Embryogenesis

in Plants. Int J Mol Sci 21.

Wong, R., M. D. Piper, B. Wertheim and L. Partridge, 2009 Quantification of food intake

in Drosophila. PLoS One 4**:** e6063.

Yamada, R., S. A. Deshpande, E. S. Keebaugh, M. R. Ehrlich, A. Soto Obando *et al.*,

2017 Mifepristone Reduces Food Palatability and Affects Drosophila Feeding

and Lifespan. J Gerontol A Biol Sci Med Sci 72**:** 173-180.

Yesbolatova, A., Y. Saito, N. Kitamoto, H. Makino-Itou, R. Ajima *et al.*, 2020 The auxin-

inducible degron 2 technology provides sharp degradation control in yeast,

mammalian cells, and mice. Nat Commun 11**:** 5701.

Yokoyama, C., M. Takei, Y. Kouzuma, S. Nagata and Y. Suzuki, 2017 Novel tryptophan

metabolic pathways in auxin biosynthesis in silkworm. J Insect Physiol 101**:** 91-

96.

Yu, S. J., 2008 Detoxification Mechanisms in Insects, pp. 1187-1201 in *Encyclopedia of*

Entomology, edited by J. L. Capinera. Springer Netherlands, Dordrecht.

Zhang, L., J. D. Ward, Z. Cheng and A. F. Dernburg, 2015 The auxin-inducible

degradation (AID) system enables versatile conditional protein depletion in C. elegans.

Development 142**:** 4374-4384.

 FIGURE LEGENDS

 Figure 1. Increased auxin exposure in adult *Drosophila* decreases survival and

 alters feeding behavior.

(A) Cartoon schematic of auxin exposure feeding protocol. Cartoon was created using

BioRender. **(B,C)** Dose response curves for males and females exposed to increasing

concentrations of auxin in the *VK00040* transgene injection line (B) or the *AGES*

transgenic line (C). **(D)** The total number of feeding events for females in the *VK00040*

or *AGES* fly line exposed to 0 mM or 10 mM auxin. **(E)** The median time of feeding

activity of females in the *VK00040* or *AGES* fly lines exposed to 0 mM or 10 mM auxin.

(F) The total number of events for males in the *VK00040* or *AGES* fly lines exposed to 0

mM or 10 mM auxin. **(G)** The median time of feeding activity of males in the *VK00040* or

AGES fly lines exposed to 0 mM or 10 mM auxin. Data shown as mean ± SEM. *$P <$

0.05, Mann-Whitney *U*-test.

Figure 2. Auxin exposure decreases acylcarnitine, fatty acid, and triglyceride

levels in adult *Drosophila*.

(A,B) Box plots illustrating the relative abundance of acylcarnitine or fatty acid

metabolites in adult females (A) or males (B) exposed to 0 mM or 10 mM auxin. All box

plots were generated using MetaboAnalyst 5.0 as described in the methods. Black dots

represent individual samples, the horizontal bar in the middle represents the median,

and the purple diamond represents the mean concentration. For all box plots, the

metabolite fold-change was >2-fold and $P < 0.05$. **(C,D)** TAG contents (mg TAG/mg

protein) in whole adult females (C) or males (D) in the *VK00040* or *AGES* fly lines

exposed to 0 mM or 10 mM auxin. Data shown as mean ± SEM. *$P < 0.05$, **$P < 0.01$,

two-tailed Student's *t*-test.

Figure 3. Lipid metabolism and detoxification transcripts are significantly altered

in adult *Drosophila* in response to auxin exposure.

(A-D) Volcano plots of differentially expressed genes graphing the statistical

significance [-$\log_{10}$(padj)] against the magnitude of differential expression ($\log_2$ Fold

Change) in females (A,B) or males (C,D) of the *VK00040* or *AGES* line. **(E)** Significantly

up-regulated genes involved in "drug metabolism" in adult females or males of the

VK00040 or *AGES* lines exposed to 10 mM auxin and compared to the 0 mM control.

(F) Significantly up- (purple) and down-regulated (green) genes involved in fatty acid

metabolism in adult females or males of the *VK00040* or *AGES* lines exposed to 10 mM

auxin and compared to the 0 mM control.

Figure 4. Auxin exposure does not significantly influence processes of

oogenesis. (A) Cartoon schematic of the adult *Drosophila* ovariole showing the anterior

germarium followed by developing egg chambers, which consist of 16 germ cells (15

nurse cells and one oocyte; green) that are surrounded by follicle cells (purple). **(B)**

Schematic of the germarium, which contains 2-3 germline stem cells (GSCs, dark

green) and somatic cells (gray and purple). Each GSC divides asymmetrically to self-

renew and generate a cystoblast that divides to form a 16-cell cyst. Early germline cysts

are surrounded by follicle cells (purple) to bud a new egg chamber. GSCs and their

progeny are identified based on the position and morphology of the fusome (pink), a

germline-specific organelle. **(C)** Germaria from adult females exposed to 10 mM auxin

for 10 days. α-spectrin (magenta), fusome; LamC (red), cap cell nuclear lamina; Vasa

(green), germ cells; DAPI (blue), nuclei. GSCs are outlined. Scale bar, 10 µm. **(D,E)**

Average number of GSCs per germarium over time in females exposed to 0 mM or 10

mM auxin in the *VK00040* (D) *AGES* (E) lines. Data shown as mean ± SEM. No

statistically significant differences, two-way ANOVA with interaction. **(F)** Germaria from

adult females exposed to 10 mM auxin. ApopTag (green), dying cells; DAPI (blue),

nuclei). **(G,H)** Average percentage of germaria containing ApopTag-positive germline

cysts in adult females in the *VK00040* (G) or *AGES* (H) lines exposed to 0 mM or 10

mM auxin. Data shown as mean ± SEM; Student's *t*-test. 100 germaria were analyzed

for each genotype and condition. **(I)** Ovarioles exposed to 10 mM auxin for 10 days

showing a healthy ovariole (top) and an ovariole with a dying vitellogenic egg chamber

(bottom). Arrowheads point to healthy or dying vitellogenic egg chambers. DAPI (white),

nuclei. Scale bar, 100 µm. **(J,K)** Average percentages of ovarioles containing dying

vitellogenic egg chambers in females exposed to 0 mM or 10 mM auxin in the *VK00040*

(J) or *AGES* (K) lines. Data shown as mean ± SEM, Student's *t*-test. 100 ovarioles were

analyzed for each genotype and condition.

Table 1. Benchmark dose of male and female *Drosophila* exposed to auxin.

Genotype	Male	Female
Oregon-R	8.89 mM	19.97 mM
VK00040	17.446 mM	14.208 mM
AGES	32.449 mM	57.291 mM

819 **Table 2. Average feeding events and durations in male and female _Drosophila_.**

Genotype	Male		Female	
	0 mM	10 mM	0 mM	10 mM
Oregon-R	342.8 ± 39.85 events 3.21 ± 0.26 sec duration	278.72 ± 26.64 events 3.21 ± 0.3 sec duration	519.14 ± 81.52 events 3.22 ± 0.24 sec duration	268.82 ± 28.65 events 3.3 ± 0.29 sec duration
_VK00040_	266.42 ± 26.27 events 3.17 ± 0.29 sec duration	208.56 ± 21.15 events 2.22 ± 0.19 sec duration	432.23 ± 64.82 events 3.0 ± 0.22 sec duration	213.03 ± 22.54 events 2.66 ± 0.24 sec duration
AGES	217.15 ± 23.09 events 3.58 ± 0.25 sec duration	169.63 ± 15.25 events 2.71 ± 0.28 sec duration	277.44 ± 31.19 events 3.42 ± 0.25 sec duration	167.49 ± 15.9 events 2.71 ± 0.26 sec duration

Fleck et al.
Figure 1

Fleck et al.
Figure 3