Automating an insect biodiversity metric using distributed optical sensors: an evaluation across Kansas, USA cropping systems

Klas Rydhmer[1,2], James O. Eckberg[3], Jonathan G. Lundgren[4], Samuel Jansson[2], Laurence Still[2], John E. Quinn[5], Ralph Washington Jr.[2], Jesper Lemmich[2], Thomas Nikolajsen[2], Nikolaj Sheller[2], Alex M. Michels[4], Michael M. Bredeson[4], Steven T. Rosenzweig[3], and Emily N. Bick[2,6,7*]

[1] Department of Geosciences and Natural Resource Management, Copenhagen University, Rolighedsvej 23, Fredriksberg C, 1958, Denmark.
[2] FaunaPhotonics, Støberigade 14, Copenhagen, 2450, Denmark.
[3] Agriculture and Food Solutions, General Mills, Minneapolis, MN 55427, United States.
[4] Ecdysis Foundation, 46958 188th St, Estelline, SD 57234, United States.
[5] Department of Biology, Furman University, 3300 Poinsett Hwy, Greenville, SC 29613, United States
[6] Department of Entomology, University of Wisconsin-Madison, 1630 Linden Dr, Madison, WI 53706, United States.
[7] Department of Plant and Environmental Sciences, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark.

*Emily N. Bick

Email: ebick@wisc.edu

Author Contributions: Klas Rydhmer aggregated the data, performed the analysis, produced figures and tables, and co-wrote the first draft of the manuscript. James O. Eckberg planned and supervised the experiment and co-wrote the introduction, discussion, and conclusion. Emily N. Bick drafted the introduction, discussion, and conclusion, contributed to the methods, and coordinated edits to the manuscript. Samuel Jansson contributed to the data analysis, the results, and the methods sections and co-wrote the draft of the manuscript. Laurence Still assisted in planning, editing, and contributed to the data analysis. Ralph Washington Jr. and Nikolaj Sheller deployed optical sensors. Jesper Lemmich and Thomas Nikolajsen assisted with planning and coordination. Mike Bredeson, Alex M. Michels, and Jonathan Lundgren collected and supervised the analysis of the invertebrate inventory data. Jonathan G. Lundgren and Steven T. Rosenzweig contributed to the planning of the experiment. Steven Rosenzweig coordinated the recruitment of cooperating farmers and farm fields to host the experiment. John E. Quinn assisted with manuscript framing and editing.

Competing Interest Statement: Klas Rydhmer, Samuel Jansson, Laurence Still, Ralph Washington Jr., Nikolaj Sheller, Jesper Lemmich, and Thomas Nikolajsen are or were affiliated with FaunaPhotonics, who developed the sensor used in the study, as employees or stakeholders. Emily N. Bick was funded in part by FaunaPhotonics as part of a Postdoctoral Fellowship granted by the Danish Innovation Fund.

Classification: Biological Sciences, Biophysics, and Computational Biology

Keywords: Biodiversity, Bioinventory, Insect monitoring, Optical sensing, Biophotonics

Abstract

Global ecosystems and food supply depend on insect biodiversity for key functions such as pollination and decomposition. High-resolution, accurate data on invertebrate populations and communities across scales are critical for informing conservation efforts. However, conventional data collection methodologies for invertebrates are expensive, labor intensive, and require substantial taxonomic expertise, limiting researchers, practitioners, and policymakers. Novel optical techniques show promise for automating such data collection across scales as they

operate unsupervised in remote areas. In this work, optical insect sensors were deployed in 20
agricultural fields in Kansas, USA. Measurements were compared to conventional assessments
of insect diversity from sweep nets and Malaise traps. Species richness was estimated on optical
insect data by applying a clustering algorithm to the optical insect sensor's signal features of
wing-beat frequency and body-to-wing ratio. Species richness correlated more strongly between
the optical richness estimate and each of the conventional methods than between the two
conventional methods, suggesting sensors can be a reliable indicator of invertebrate richness.
Shannon- and Simpson indices were calculated for all three methods but were largely
uncorrelated including between conventional methods. Although the technology is relatively new,
optical sensors may provide next-generation insight into the spatiotemporal dynamics of
invertebrate biodiversity and their conservation.

Significance Statement
The implications of this research extend from the field level to the regional level. Much of what
scientists understand about the decline of invertebrates comes from a small number of long-term
studies that can be coarse and correlational in nature. High-resolution biodiversity data sets on
fields to landscapes may provide the insight needed for the successful management and
accounting of biodiversity by practitioners and policymakers. Such high-resolution data has the
potential to support global efforts and coordination of biodiversity conservation.
66

Main Text

Introduction

Invertebrate biodiversity is fundamental to ecosystem processes, functions, and services (Yang & Gratton, 2014). Monitoring invertebrate populations and communities can inform management and policy at multiple scales. Such data are critical to agricultural production and sustainability (Landis, 2017). However, invertebrate biodiversity is difficult to quantify (Geiger et al., 2016; Shortall et al., 2009) and monitor at broad spatial and temporal scales (Sánchez-Bayo & Wyckhuys, 2019; Tilman et al., 1994). The difficulty is largely due to the necessity of skilled labor required for taxa identification on which biodiversity quantification relies (Wägele et al., 2022), and is both limited and prohibitively costly (Gardner et al., 2008). Common approaches to collecting insect inventories include sweep netting as well as Malaise-, pan-, and light traps. Each method has its own bias toward certain insect groups (Bick et al., 2020) often resulting in the concurrent use of techniques in studies and practice (LaCanne & Lundgren, 2018).

New technology is needed to monitor invertebrate biodiversity in real time for agricultural systems. Such a tool would provide data to support biodiversity-focused management at field to landscape scales (LaCanne & Lundgren, 2018) and allow for tracking of the impact of conservation measures, or the lack thereof. Automation of systems has the potential to reduce labor, time, costs, and human error. While many automated insect monitoring tools are available for agricultural pest monitoring (Bick et al., 2023; Silva et al., 2013), overall, these approaches are not suitable for assessing biodiversity as they focus on the identification of indicator species, not communities (J. G. Lundgren & Fausti, 2015a). The automatic quantification of invertebrate biodiversity could improve the data available for monitoring and evaluation of conservation efforts but currently, no method exists at scale (Wägele et al., 2022), despite calls for such data and analytics to inform the assessment and management of ecosystems (García et al., 2023).

Real-time data on invertebrate biodiversity likely would improve our understanding of insect population changes at a regional or even global scale, filling a gap in the tracking of insect change. The incorporation of 'big data' has been shown to help mitigate some methodological biases (Geiger et al., 2016). One such effort is the global malaise project that is using automated taxonomic identification from traps using DNA, addressing the most labor-intensive part of this method (Krishna Krishnamurthy & Francis, 2012). It is a highly promising 'big data' approach; unfortunately, the method over-represents known species, has an inherent sampling bias towards flying insects and emphasizes species with large mitochondrial differences. Optical entomological methods such as lidar, where an optical signal is recorded from insects flying through a beam of emitted light, can record large numbers of insect flights without using a lure. However, it is unclear how optical sensors compare to conventional methods in measuring populations and communities (Garcia et al., 2023; Rydhmer et al., 2022).

The goal of this study is to determine if the measurement of an insect biodiversity metric can be automated with the use of optical near-infrared insect sensors. In this work, we deployed sensors (Rydhmer et al., 2021) in 20 agricultural fields across six crops in Kansas, USA. The sensors were deployed alongside Malaise traps and the sites were sampled with sweep nets. Each site was evaluated on two different occasions to capture seasonal changes. Specifically, we compared conventional methods to each other and with the automated biodiversity metric utilizing unsupervised clustering of data collected by a lidar-based sensing method.

Materials and Methods
Data collection

Insect populations were monitored at 20 sites (Figure 1) in June and July of 2020 using sensors alongside conventional methods (Malaise traps and sweep nets). Representative agricultural crops of central Kansas were sampled including three corn, three sorghum, six soybean, one alfalfa, two pasture, and five complex cover crops. The complex cover crops consisted of approximately eight species of annual grass and forb cover crops. An autonomous near-infrared sensor (described in (Rydhmer et al., 2021) and produced by FaunaPhotonics ApS., Copenhagen SV, Denmark) was placed ~50 m from the field margin and was monitored continuously for two periods of three days in June and in July. The sensor uses light-emitting diodes to transmit infrared light (810 nm & 970 nm), creating a measurement volume between 5 and 70 L, depending on insect size (Rydhmer et al., 2021). Insects flying in front of the sensor back-scatter light, which is recorded by a photodiode as a time signal.

Figure 1. Map of 20 field site locations distributed around central Kansas. Fields are color-coded by crop type. Field dots are enlarged and shifted to maintain the anonymity of the participating farms. Map data from www.openstreetmap.org.

Insect recordings are automatically separated from noise originating from other sources (e.g. rain drops or plant interference) using proprietary cloud-based neural network software, as used in Bick et al., 2023 and Rydhmer et al., 2021. Additionally, observations without clearly identified wingbeats or body-to-wing ratios were discarded. A total of 1,057,115 observations were recorded, of which 106,083 remained after filtering and were included in the study. A recorded observation consists of time series data from which information pertaining to the physical features of the individual insect can be obtained (Rydhmer et al., 2021).

Sensors were compared with conventional sampling of invertebrates (sweep nets and Malaise traps) in the same fields. Foliar and low flying insects were captured using a sweep net (38 cm diam., Bioquip™, Rancho Dominguez, CA, USA). Insects were collected at 50, 100, and 150 m from the field edge along a linear transect. Sweeps (n = 50 per location) were performed perpendicular to the transect, parallel to the field edge. Insects were transferred to a sealed plastic bag and were frozen until processed. In the laboratory, insects were thawed, sorted from the plant material, and identified.

Malaise traps were deployed at each site to capture the aerial insect community. A single bi-directional, Townes-style trap (dimensions 1.8 long; 1.8 m at its tallest height, and 1.2 m at its shortest height) was placed 100 m from the margin and adjacent to the ecosystem service

156   sampling areas. The wall of the trap was parallel with the field margin. The traps were allowed to
157   operate for 24 h, and the insects captured in the collection vials were preserved in ethanol.
158
All specimens collected by sweep net and malaise traps were identified to the lowest possible
taxonomic unit (i.e., species or morphospecies). Due to a lack of species identification knowledge
and time limitations, thrips (Insecta: Thysanoptera) were not identified beyond the family level and
were not included in community metrics analyses (abundance, species richness, and diversity).
All immature insects were identified to family and grouped together, except for lepidopteran
larvae, which were categorized as morphospecies independent of the adult stage due to their
functional differences. All other specimens were identified to species using written and online
taxonomic keys. Specimens that were not able to be positively identified to species were
separated into distinct morphospecies. Voucher specimens of all taxa are housed in the Mark F.
Longfellow Biological Collection at Blue Dasher Farm, Estelline, SD.
169
Ecosystem services were evaluated for insect and weed seed predation. First, invertebrate
predators were isolated from both the soil and foliar communities. Additionally, predation rates in
each field were assessed using sentinel wax moths (*Galleria mellonella* L. [Lepidoptera:
Pyralidae]) larvae following the Lundgren et al., 2006 protocol, using 15 sentinels per plot
arranged in three 5 × 3 7.5 m grid orientations (n = 45 per field). Weed seed predation was
assessed from isolating soil and foliar granivore communities and their services using seed cards
as described in Lundgren et al., 2006. Granivore services were measured on three abundant
weed species (Johnsongrass (*Sorghum halapense* (L.) Pers.; Poaceae), lambsquarters
(*Chenopodium album* L.; Amaranthaceae) and redroot pigweed (*Amaranthus retroflexus* L.;
Amaranthaceae), V & J Seed Farms, Woodstock, IL, USA). Seeds were attached to 10 × 8 cm
plastic cards (Avery™ insertable plastic dividers; #11200; Brea, CA, USA) using 6 cm strips of
double-sided tape (Scotch, 3M, St Paul, MN, USA). Each species (n = 20 seeds of each species;
60 seeds per card) were placed on a 2 × 10 pattern on each card. Fine quartz sand was spread
over exposed areas of the tape to exclude visiting invertebrates. To exclude granivorous
vertebrates, a wire cage (14 × 12 cm cage, 1.4 × 1.4 cm mesh opening) was placed over the card
and placed >3 cm above it. Control cards were used to account for seed loss from environmental
factors such as wind or rain and contained 1.5 × 1.5 mm black glass beads (Cousin™ DIY,
#AJM61215021, Largo, FL, USA) of comparable size as the weed seeds (Lundgren et al., 2006).
Each plot received three seed cards and one control card (n = 9 seed cards and three control
cards per field), placed on the soil surface in the four corners of each plot. Granivory was
measured as the number of seeds removed or damaged per card after a 3-dayosure.
191
Data analysis
The wing-beat frequency (WBF) and body-to-wing ratio (BWR) were calculated from all observed
insect flights similarly to previous groups (Gebru et al., 2018; A. Genoud et al., 2020; A. P.
Genoud et al., 2019). The signal from the insect body (σ_b) and the diffuse and specular signal
contributions from the insect wings (σ_{dw} and σ_{sw}) are estimated and separated using sliding
minimum, sliding median and sliding maximum filters with a filter width corresponding to the wing
beat period of the insect. The BWR is defined as the closed ratio between the body and wing
contributions according to equation (1). An example of an insect signal is shown in Figure 1a.
200
201    $BWR = \sigma_b / (\sigma_{dw} + \sigma_b)$ (1)
202

203
Figure 2. Example of an insect event's signal and clustering. a) An example of an insect event
recorded from the sensor. The wing beats are visible as modulations on top of the signal. The
dashed red, solid magenta and dash-dotted blue curves show the body, diffuse- and specular
wing signals respectively. The BWR is the ratio between the magnitude of the body- and diffuse
wing signal. b) Clustered insect events recorded by the sensor in a soybean field (Field R) in July.
The grey events are too sparse to form clusters and are therefore discarded.

210
Insects of the same species exhibit similar physical properties, and therefore also similar signal
features (Kirkeby et al., 2021). Normalization of the feature space is a standard procedure prior to
clustering. While BWR values are bound between 0 and 1 by definition (equation 1), WBF values
frequencies typically vary between 20 Hz and 1 kHz (Jansson et al., 2019). WBFs were therefore
divided by 1000 to produce values falling predominantly between 0 and 1. For clustering, we used
the DBSCAN (Density-based spatial clustering of applications with noise) algorithm (Ram et al.,
2010) due to its suitability in identifying clusters without a Gaussian distribution assumption (Ester
et al., 1996). DBSCAN uses two parameters, the minimum number of insects needed to form a
cluster (min_samples) and the merge distance ϵ, to determine which observations to merge into
clusters. Data points too far away from any cluster and too sparsely distributed to form a new
cluster are defined as outliers. This method was used to calculate the number of clusters or
distinct groups (i.e. richness) and a diversity index of cluster groups based on Shannon and
Simpson indices.

224
225 All insects collected with Malaise traps and sweep nets were classified by order, family, and
226 species when possible. Then species richness (defined as the number of distinct taxonomic
227 species present, independent of abundance), Shannon index, and Simpson index were
228 calculated on the insect samples from both conventional methods for each field in June and July.

229
The data from the capture methods were randomly divided into two data sets: one used to
optimize the DBSCAN clustering algorithm, and the other used for testing. To have a sufficiently
large test set, the optimization set was limited to 30% of the data collected. During the
optimization, ϵ and min samples were tuned to maximize the Spearman correlation between
biodiversity metrics from the sensors and conventional metrics using stochastic gradual descent.
This process was repeated for the richness and Shannon and Simpson indices for each of the
trapping methods, plus an additional model fitted to the combined species richness from both
conventional methods. Shannon and Simpson indices were not calculated on the combined

dataset since these indices rely on the relative abundance of species, which are not comparable
between the two methods.
240
241 Optimal parameters could be found that produced significant correlation ($p < 0.05$) for four of the
242 seven comparative measures; however, no parameters could be found that satisfactorily modeled
243 the Shannon index from the sweep netting nor the Simpson index for either trapping method.
244
245 Spearman-rank correlations between the clustering results calculated from the optical sensor
246 data and the biodiversity measures obtained with the two physical insect field-sampling methods
247 were calculated. Additionally, Analyses of Variance (ANOVA)and TukeyHSD post hoc analyses
248 were conducted to evaluate the impact of sampling month, crop type, and field on richness
249 estimates.
250
251 **Results**
252
253 In total, 106,083 insect flight events were recorded by the sensors. The Malaise traps collected
254 14,641 insects, whereas sweep nets collected 15,858 insects (Figure 3). The optical sensors
255 recorded approximately one order of magnitude more insect flights compared to the number of
256 insects collected with each of the conventional methods (Figure 3). Insect abundance was
257 uncorrelated between all methods, including both conventional methods and the automated

method (Figure 4; Malaise trap counts and sweep net counts r = 0.25, p =0.16, sensor events and sweep net counts r = 0.05, p = 0.78, sensor events and Malaise trap counts r = 0.05, = <0.88).

Figure 3. The number of insects collected using sweep nets (top panel) and Malaise traps (middle panel), and insect flight events recorded with the sensor (bottom panel) per field. Insect

numbers are separated by month with insects observed in June visualized with blue bars and in July with red bars.

Figure 4. Scatter plots of measured insect abundance comparing the monitoring methods on a logarithmic scale. a) Scatter plot of the number of insects captured with sweep nets and Malaise traps. b) Scatter plot of the number of insects captured with sweep nets and the number of insect flight events recorded by the sensor. c) Scatter plot of the number of insects captured with Malaise traps and the number of insect flight events recorded by the sensor. No correlations were found on insect abundance for any of these methods.

Comparing the relative insect abundance between orders collected with conventional methods (Figure 5) depicts differences in capturing biases due to methodology. Diptera were most frequently collected from the Malaise traps, whereas Hemiptera, then Diptera and Psocoptera were more frequently captured with sweep nets. In general, less flight-active insects were more prominent in the sweep net data.

 Figure 5. The number of insects collected with sweep net sampling and Malaise trap monitoring,
286 aggregated by order.
287
There were no discernible differences in variation between time points from sensors (F = 6.091,
Pr = 0.0191) and sweep nets (F = 1.326, Pr = 0.258). However, Malaise trap abundance showed
significantly greater insect densities in July (μ = 76.9, F = 9.71, Pr = 0.0037) than June (μ = 43.3).
Crop type was found to impact sweep net abundance (F = 3.369, Pr = 0.008) but not sensor (F =
1.644, Pr = 0.17) or Malaise trap (F = 1.692, Pr = 0.152) abundance estimates. A series of
TukeyHSD post hoc analyses found no differences in abundance estimates between sample time
points for each field.
295
296 Insect species richness was estimated from sensor-recorded insect flight events using a set of
297 seven DBSCAN parameters (models) to cluster the held-out test data, yielding the number of
298 clusters per field – the novel richness metric (Figure 2b). The correlation between the number of
299 clusters and each of the comparative biodiversity metrics are shown in Table 1.
300

Fitting data	Correlation coefficients (R, p-value)						
	Richness			Shannon		Simpson	
	Malaise trap	Sweep net	Combined	Malaise trap	Sweep net	Malaise trap	Sweep net
Richness Malaise	**0.42, 0.050**	0.26, 0.246	0.37, 0.112	**0.44, 0.044**	-0.22, 0.359	0.35, 0.114	0.01, 0.964
Richness sweep net	**0.45, 0.035**	**0.44, 0.046**	**0.47, 0.035**	0.11, 0.612	0.20, 0.378	0.09, 0.702	0.16, 0.475
Richness combined	**0.52, 0.014**	**0.48, 0.028**	**0.55, 0.012**	0.21, 0.344	0.20, 0.394	0.14, 0.534	0.14, 0.537
Shannon Malaise	**0.42, 0.05**0	0.26, 0.246	0.37, 0.112	**0.44, 0.044**	-0.22, 0.359	0.35, 0.114	0.01, 0.964
Shannon sweep net	**0.45, 0.037**	0.37, 0.097	**0.46, 0.040**	0.27, 0.221	0.22, 0.339	0.17, 0.459	0.08, 0.733
Simpson Malaise	**0.44, 0.041**	0.32, 0.159	0.43, 0.061	-0.16, 0.484	0.03, 0.893	0.02, 0.913	0.06, 0.797
Simpson sweep net	0.05, 0.837	-0.13, 0.569	-0.07, 0.774	**-0.73, 0.005**	0.07, 0.817	-0.04, 0.846	0.12, 0.598

301 **Table 1.** Correlations between the automated biodiversity metrics calculated from sensed insect
302 data, and those obtained from Malaise trap and sweep net collections. Rows in the table denote
303 which data was used to fit the clustering algorithm, whereas columns indicate which parameters
304 the obtained correlations refer to. Correlations with a p-value below 0.05 are significant and
305 marked in bold.
306
Per field, the mean number of clusters that approximate richness was 41.1 (N = 34, SE = 3.29).
The Malaise traps had a mean field richness of 60.5 species (N = 37, SE = 6.43) containing 10
orders, 146 families, and 709 species. The mean richness observed in the sweep nets was 47.4
species (N = 35, SE = 5.53), containing 11 orders, 149 families, and 664 species. Combined, the
collected samples with both field-sampling methods contained 941 species distributed over 183
insect families in 11 orders.
313
314 The three models fit on species richness are generally comparable (fit on sweep net, Malaise trap,
315 and their combined data). Identical DBSCAN parameters were calculated when the models were
316 fit on the combined sweep net and Malaise trap richness. We therefore used this model termed the

317 'automated biodiversity metric' to evaluate the relationships between sensors, and the
318 conventionally measured species richness and ecosystem services.
319
All species richness metrics were correlated (Figure 6a-d). The weakest correlation was between
Malaise trap and sweep net richness metrics (R = 0.36, p = 0.046). The correlation between the
number of clusters calculated in the sensor data and the conventional models was strongest for
the combined richness, which was what the model was fitted on (R= 0.55, p = 0.012; Figure 6d).
Significant yet weaker correlations were also found between the model and the Malaise trap and
sweep net richness respectively (R = 0.52, p = 0.014; R = 0.48, p = 0.028).
326

327
Figure 6. Scatter plots and Spearman correlations for the species richness estimations across all
models. The sensor results are from the model fitted to the total richness in both Malaise traps
and sweep nets. a) Species richness calculated from Malaise traps vs. sweep net samples, b)
species richness calculated from sensor clusters vs. sweep net samples, c) species richness
calculated from sensor clusters vs. Malaise trap samples, and d) species richness calculated from
sensor clusters vs. total richness across traps and sweeps.
334
335
336 No correlations were found when comparing sensor richness to any ecosystem services (Table
337 2). Conventional sampling methods were typically not correlated with ecosystem services with

338 one exception. Sweep net species richness was positively correlated with the percent of
339 waxworms predated (Table 2).
340

Richness metrics	% Waxworms Predated	Total # of predators	Johnsongrass predation	Pigweed predation	Lambsquarter predation	All seed predation
	Spearman's correlation coefficient (R, p-value)					
sweep net	**0.49, 0.04**	0.34, 0.17	-0.37, 0.13	0.05, 0.85	-0.04, 0.89	-0.07, 0.78
Malaise trap	-0.16, 0.52	-0.13, 0.61	-0.16, 0.52	-0.36, 0.15	-0.40, 0.10	-0.45, 0.06
total - richness	0.21, 0.43	0.16, 0.56	-0.39, 0.14	-0.26, 0.33	-0.32, 0.23	-0.41, 0.11
sensor clusters	-0.11, 0.66	-0.26, 0.29	0.14, 0.56	-0.11, 0.66	-0.15, 0.54	-0.11, 0.66

341
342 **Table 2.** Spearman's correlation table between richness metrics calculated from sweep nets,
343 Malaise traps, combined conventional methods, and sensor clusters (automated biodiversity
344 metric) compared to ecosystem services of percent waxworm predation, total number of
345 predators, Johnsongrass predation, Pigweed predation, Lambsquarter predation, and all seed
346 predation.
347
348
349 **Discussion**
350
This work serves as the first field-validated insect biodiversity metric using autonomous
distributed optical sensors (Kouakou et al., 2020). The automated biodiversity metric was
calculated and validated from flight events of sufficiently high quality to be able to extract a
wingbeat frequency and body-wing ratio. Our results suggest that the sensor-derived metric is
correlated with conventional estimates of biodiversity. Specifically, the sensor-derived biodiversity
metric optimized for insect species richness is more correlated with each conventional sampling
method than these methods are to one other (Figure a-d) (Malaise trap – sweep net R = 0.36;
Malaise trap – sensor R = 0.52; sweep net – sensor R = 0.48). This indicates that metrics derived
from optical sensors are likely generalizable and thus have the potential to provide accurate and
autonomous measurements of insect species richness. Ecosystem generalizability was
demonstrated by deriving and testing the biodiversity metric across six major crops in central
Kansas. Still, future work is needed to evaluate the extent to which the metric may be generalized
across agroecosystems outside our study area and to other terrestrial ecosystems. Current
results indicate that the insect diversity metric may be calculated for a variety of functionally
different crops without the need to classify insects into taxonomic groups. This approach is
beneficial as such classification at present requires skilled labor and significant time. This metric
may provide new insight into the management of ecosystems as significant and growing evidence
suggests that biodiversity is correlated with greater ecosystem functions such as pest control
(Lundgren & Fausti, 2015). Future work may focus on characterizing the composition of insect
communities and species to address specific needs of managing ecosystems beyond
biodiversity.
372
The automated biodiversity metric and Malaise trap species richness significantly correlated
across all method iterations, save one (Table 1). Sweep net richness only correlated with two of
the method iterations. The stronger correlation between the sensors and Malaise traps is
hypothesized to result from these methods monitoring flying insects continuously, compared to
sweep net sampling. Results were less clear for the correlations with Shannon and Simpson
species diversity indices. The models fitted on Malaise trap richness also was significantly
correlated with for Malaise trap Shannon index (Table 1). This is likely due to the co-correlation

between the richness and Shannon index in the malaise trap (R=0.6, p=0.01, Supplementary
Table 3). Other similar curiosities, such as the negative correlation with the Malaise trap Shannon
index achieved when fitting on the sweep net Simpson index are also assumed to be the results
of co-correlations between the conventional methods. A full table of all co-correlations is included
as supplementary material (Supplementary Table 3). However, when fitting on Shannon index
from the sweep-net data no correlation was found. This is likely due to overfitting, where the
model performed well on the fitting data but did not generalize to the rest of the dataset. A larger
fitting dataset is needed to resolve this issue. No model resulted in significant correlations in
Simpson indices between any of the sampling methods. The lack of consistent correlations
between biodiversity metrics may also reflect the nature of the biodiversity indexes which
considers species evenness, a characteristic not fully accounted for in the DBSCAN algorithm
based on minimum thresholds for classification of clusters.
392
The sensor observed the greatest number of insects, recording almost one order of magnitude
more than both the Malaise traps and sweep nets (7.25 and 6.69 times, respectively). This
difference is likely explained first by the observational period and then by methodology. Both
sensors and Malaise traps continuously monitored each field, unlike sweep nets which collect
insects at discrete time points. The sensors' monitoring period was three times longer than the
Malaise traps. However, even after accounting for the greater measurement period, the sensor
methodology was still 2.42 times more efficient than Malaise traps at observing all insects. A
previous study reported sensors observed 19 times the number of insects compared to those
collected in water traps, another continuous monitoring method (Rydhmer et al., 2021). Specific
insect species are also detected more efficiently. For example, the sensor was reported to be
18.6 and 6.7 times more efficient than plant counts (discrete sampling) and water traps
(continuous monitoring) at observing pollen beetles (*Brassicogethes aeneus*; Bick et al., 2023).
Greater sampling efficiency may be associated with measurement volume, a potential correlate of
insect counts. Optical sensors with greater measurement volumes report recording tens of
thousands of insect flights per day (Brydegaard et al., 2020). A potential confounding effect of
sensors is the possible 'double counting' bias, as an individual insect can be recorded repeatedly.
While 'double counting' possibly explains the greater number of observed insects, such biases
are a common limitation of count-based inferences on population dynamics (Elphick, 2008).
Despite its potential limitations, our understanding of complex species and community dynamics
can benefit greatly from automation that significantly increases sampling intensity across space
and time.
414
The lack of correlation of abundance across all three methods (Figure 4) is surprising as previous
work has shown correlations between sensor measurements and water traps for insect
abundance (Rydhmer et al., 2021). Disparities in sampling timing may be contributing to the lack
of correlation. While sweep netting occurred in conjunction with setting up or taking down the
Malaise traps, these efforts were substantially less correlated with the setup of the optical
sensors: to the nearest 3 days in June and the nearest 22 days in July. The lack of correlation
between the Malaise traps and the sensors may be due to the long period between the
monitoring sessions at each site. Insect flight activity is heavily influenced by the weather, or the
seasonal differences between the beginning and end of July – both of which may also explain the
significance of the month on Malaise trap data. An additional factor may be the high noise
composition of the recorded signals due to plant interference. During cleaning of this dataset, it is
possible that variations in the relative degree of noise signals between fields (e.g. as a result of
different crop heights and stiffness) resulted in more data loss from noisier fields, thus introducing
a systematic error in abundance measurements for the sensor data. However, it should be noted
that we observed no statistical correlation in abundance collected with Malaise traps versus
sweep nets.
431
One challenge with the sensor's dataset was the high proportion of noise signals, thought to
result from plant interference. Of the total 1,057,115 signals recorded by the sensor, only ~10%

were classified as insects and included in the analysis. While we believe the noise classification
filter is highly accurate, misclassified events may alter the total count. The signals generated by
insects and plants moving through the sensor's measurement volume are very different. Most
non-insect events show no high-frequency components and are therefore correctly removed by
the noise filter. However, plants modulating in front of the sensor may appear to have a wing beat
frequency and would be misclassified as insects. It is also hypothesized that strong signals
generated by vegetation interference may obscure weaker signals generated by small insects.
Regardless, misclassification is likely low enough to substantially alter the representation of the
insect population or the automated biodiversity metric.
443
There is a need for 'big data' in entomology, and more particularly for measuring biodiversity to
inform conservation. Autonomous optical sensors, such as the ones used in this study, provide
one such solution that offers continuous, potentially real-time monitoring to support next
generation, big data insight to the field of entomology. Moreover, the success in leveraging
sensor data to calculate a biodiversity species richness metric indicates great promise in the use
of autonomous sensors to monitor biodiversity. The use of sensors for insect and insect diversity
monitoring is faster, in this case potentially more representative of richness, and likely cost-
effective due to a decrease in labor compared to conventional methods. Sensors complement
conventional sampling methods by allowing for real-time estimation of biodiversity, reducing time
lags associated with traditional species inventory. Automated methods presented in this paper,
once a generalizable calibration has been determined, offer faster estimates of biodiversity which
will support time-critical decision making and conservation planning efforts. Methods such as the
one described in this paper do not rely on identifying taxonomic groups and remove human error
(a major concern for insect identification). Furthermore, standardized sensors are not prone to
local and regional variations in sampling methods and may therefore be able to facilitate
comparative biodiversity monitoring on a global scale.
460
There is a need to scale up and scale down sensor monitoring to understand species dynamics.
The current technology complements the entomological radar group 'BioDAR' which is aiming to
use libraries of insect radar signals for functional group classifications for high flying migratory
insects at a regional scale (Rhodes et al., 2022). Similar estimates of insect functional groups
might be similarly inferred from optical sensor recordings for all flying insects on a field scale.
Similarly, vertical looking radar is used to classify insects into higher level taxonomic groups such
as Order or even Genus (Chapman et al., 2002; Stefanescu et al., 2013; Wood et al., 2009). It
seems likely that similar or even higher precision can be achieved by including taxonomic
information in clustering algorithms, such as specific orders (e.g. Lepidoptera, Coleoptera,
Diptera). Future work could focus on identifying these groups, determining functional biodiversity,
and quantifying their contribution to ecosystem services.
472
The current study shows a single instance of correlation between richness and a measure of
ecosystem services. Greater species richness does not always translate into an increase in
functional biodiversity or ecosystem services, as there is often ecological redundancy (Greenop
et al., 2018). The lack of a relationship may also reflect different ecological interactions among
species in the upper canopy versus above canopy level. These questions can be further explored
in future work when the sensor's ability to estimate functional biodiversity has been developed.
479
Conclusion
Conservation of biodiversity is gaining recognition as a global challenge with similar significance
to climate change. However, unlike global climate, species populations and biodiversity function
across different local to regional spatiotemporal scales. Detailed data on insect diversity across
these scales is needed to assess the decline and inform conservation efforts. There is a call to
automate the collection of *in situ* data and integrate such data with remote sensing- based
models to accelerate conservation of global ecosystems (Garcia et al 2023). These integrated
technology and big data approaches are especially needed for the conservation of invertebrates

and could expand upon and accelerate the long-term and detailed monitoring efforts of
invertebrate biodiversity (Sánchez-Bayo & Wyckhuys, 2019) The current study demonstrates
successful development of *in situ* data collection able to be integrated with remote sensing
models as described by Garcia et al 2023. Such approaches are poised to support the
Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services with rich,
real-time data and help inform global biodiversity models that have had to rely on coarse, low-
resolution data sets in some cases (Schipper et al., 2020).
495
Real-time feedback of simple metrics of biodiversity would greatly benefit agriculture by
demonstrating the association between management and biodiversity in real-time, without time
lag to process samples. The combination of pest detection with optical sensors (Bick et al., 2023;
Kirkeby et al., 2021) and monitoring of biodiversity may inform integrated pest management and
reduction of pesticide applications. Farming informed by such data analytics may deliver
significant benefits to farmers including substantial reductions in pests (Lundren and Fausti 2015)
and significant economic benefits at both farm and regional levels (Landis et al 2008, LaCanne
and Lundgren 2018). Networks of local monitoring sensors may scale up to infer regional
biodiversity and inform its management. The current technology could complement the global
Malaise trap initiative by increasing the number of observations by an order of magnitude and
providing earlier warning signs of regional pest movement or species decline. Our results suggest
AI supported human expertise may provide the most efficient, robust inference on biodiversity.
Thus, we are not advocating such technology replaces conventional monitoring, but rather that
this automation enhances the state of the art. Autonomous monitoring has the potential to
revolutionize the field of entomology by forming the basis for a next generation of species to
community insect models predicting the dynamics of invertebrates.
512
513

Acknowledgments
We thank the USDA Cheney Lake Conservation District (Lisa French), Understanding Ag (Ray
Archuleta) for logistical support and field assistance. Tom Rabaey (General Mills) provided helpful
feedback and advice to guide this project. We thank Kevin James Knagg and Mads Fogtmann
from FaunaPhotonics A/S for facilitating this work. We thank Ecdysis Foundation field and lab
technicians for collecting, processing, and Dr. Kelton D. Welch for identifying invertebrate
specimens. We thank the farmers cooperating in the General Mills regenerative Agriculture
Programs who granted us access and permission to sample their farm fields.

522
523 **References**
524

Bick, E., Dryden, D. M., Nguyen, H. D., & Kim, H. (2020). A Novel CO2-Based Insect Sampling
Device and Associated Field Method Evaluated in a Strawberry Agroecosystem. *Journal of*
Economic Entomology. https://doi.org/10.1093/jee/toz359

Bick, E., Sigsgaard, L., Torrance, M. T., Helmreich, S., Still, L., Beck, B., Rashid, R. El, Lemmich, J.,
Nikolajsen, T., & Cook, S. M. (2023). Dynamics of pollen beetle (Brassicogethes aeneus)
immigration and colonisation of oilseed rape (Brassica napus) in Europe. *Pest Management*
Science. https://doi.org/10.1002/ps.7538

Brydegaard, M., Jansson, S., Malmqvist, E., Mlacha, Y. P., Gebru, A., Okumu, F., Killeen, G. F., &
Kirkeby, C. (2020). Lidar reveals activity anomaly of malaria vectors during pan-African
eclipse. *Science Advances*, *6*(20). https://doi.org/10.1126/sciadv.aay5487

Chapman, J. W., Reynolds, D. R., Smith, A. D., Riley, J. R., Pedgley, D. E., & Woiwod, I. P. (2002).
High-altitude migration of the diamondback moth Plutella xylostella to the U.K.: a study

using radar, aerial netting, and ground trapping. *Ecological Entomology*, *27*(6), 641–650.
https://doi.org/10.1046/j.1365-2311.2002.00472.x

Elphick, C. S. (2008). How you count counts: The importance of methods research in applied
ecology. *Journal of Applied Ecology*, *45*(5), 1313–1320. https://doi.org/10.1111/J.1365-
2664.2008.01545.X

Ester, M., Kriegel, H., Sander, J., & Xu, X. (1996). A Density-Based Algorithm for Discovering
Clusters in Large Spatial Databases with Noise. *KDD'96: Proceedings of the Second*
International Conference on Knowledge Discovery and Data Mining, *96*(34), 226–231.

García, G. C., Bagstad, K. J., Brun, J., Chaplin-Kramer, R., Dhu, T., Murray, N. J., Nolan, C. J.,
Ricketts, T. H., Sosik, H. M., Sousa, D., Willard, G., & Halpern, B. S. (2023). The future of
ecosystem assessments is automation, collaboration, and artificial intelligence. *Environ.*
Res. Lett, *18*, 11003. https://doi.org/10.1088/1748-9326/acab19

Garcia, K., Olimpi, E. M., M'Gonigle, L., Karp, D. S., Wilson-Rankin, E. E., Kremen, C., & Gonthier,
D. J. (2023). Semi-natural habitats on organic strawberry farms and in surrounding
landscapes promote bird biodiversity and pest control potential. *Agriculture, Ecosystems &*
Environment, *347*, 108353. https://doi.org/10.1016/J.AGEE.2023.108353

Gardner, T. A., Barlow, J., Araujo, I. S., Ávila-Pires, T. C., Bonaldo, A. B., Costa, J. E., Esposito, M.
C., Ferreira, L. V., Hawes, J., Hernandez, M. I. M., Hoogmoed, M. S., Leite, R. N., Lo-Man-
Hung, N. F., Malcolm, J. R., Martins, M. B., Mestre, L. A. M., Miranda-Santos, R., Overal, W.
L., Parry, L., … Peres, C. A. (2008). The cost-effectiveness of biodiversity surveys in tropical
forests. *Ecology Letters*, *11*(2), 139–150. https://doi.org/10.1111/J.1461-
0248.2007.01133.X

Gebru, A., Jansson, S., Ignell, R., Kirkeby, C., Prangsma, J. C., & Brydegaard, M. (2018). Multiband
modulation spectroscopy for the determination of sex and species of mosquitoes in flight.
Journal of Biophotonics, *11*(8). https://doi.org/10.1002/jbio.201800014

Geiger, M. F., Moriniere, J., Hausmann, A., Haszprunar, G., Wägele, W., Hebert, P. D. N., & Rulik,
B. (2016). Testing the Global Malaise Trap Program – How well does the current barcode
reference library identify flying insects in Germany? *Biodiversity Data Journal*, *4*(4).
https://doi.org/10.3897/BDJ.4.E10671

Genoud, A., Gao, Y., Williams, G., & Thomas, B. (2020). A comparison of supervised machine
learning algorithms for mosquito identification from backscattered optical signals.
Ecological Informatics, *58*.

Genoud, A. P., Gao, Y., Williams, G. M., & Thomas, B. P. (2019). Identification of gravid
mosquitoes from changes in spectral and polarimetric backscatter cross sections. *Journal*
of Biophotonics, *12*(10). https://doi.org/10.1002/JBIO.201900123

Greenop, A., Woodcock, B. A., Wilby, A., Cook, S. M., & Pywell, R. F. (2018). Functional diversity
positively affects prey suppression by invertebrate predators: a meta-analysis. *Wiley Online*
Library, *99*(8), 1771–1782. https://doi.org/10.1002/ecy.2378

Jansson, S., Gebru, A., Ignell, R., & Abbott, J. (2019). Correlation of mosquito wing-beat
576   harmonics to aid in species classification and flight heading assessment. *European*
577   *Conference on Biomedical Optic*.

578   Kirkeby, C., Rydhmer, K., Cook, S., & Strand, A. (2021). Advances in automatic identification of
579   flying insects using optical sensors and machine learning. *Scientific Reports*, *11*(1), 1555.

580   Kouakou, B. K., Jansson, S., Brydegaard, M., & Zoueu, J. T. (2020). Entomological Scheimpflug
581   lidar for estimating unique insect classes in-situ field test from Ivory Coast. *OSA Continuum*,
582   *3*(9), 2362–2371.

583   Krishna Krishnamurthy, P., & Francis, R. A. (2012). A critical review on the utility of DNA
584   barcoding in biodiversity conservation. *Biodiversity and Conservation 2012 21:8*, *21*(8),
585   1901–1919. https://doi.org/10.1007/S10531-012-0306-2

586   LaCanne, C. E., & Lundgren, J. G. (2018). Regenerative agriculture: Merging farming and natural
587   resource conservation profitably. *PeerJ*, *2018*(2), e4428.
588   https://doi.org/10.7717/PEERJ.4428/SUPP-1

589   Landis, D. A. (2017). Designing agricultural landscapes for biodiversity-based ecosystem services.
590   *Basic and Applied Ecology*, *18*, 1–12. https://doi.org/10.1016/J.BAAE.2016.07.005

591   Lundgren, J. G., & Fausti, S. W. (2015a). Trading biodiversity for pest problems. *Science*
592   *Advances*, *1*(6). https://doi.org/10.1126/SCIADV.1500558

593   Lundgren, J. G., & Fausti, S. W. (2015b). Trading biodiversity for pest problems. *Science*
594   *Advances*, *1*(6). https://doi.org/10.1126/SCIADV.1500558/SUPPL_FILE/1500558_SM.PDF

595   Lundgren, J., Shaw, J., Zaborski, E., & Eastman, C. (2006). The influence of organic transition
596   systems on beneficial ground-dwelling arthropods and predation of insects and weed
597   seeds. *Renewable Agriculture and Food Systems*, *21*(4), 227–237.
598   https://pubag.nal.usda.gov/catalog/49820

599   Ram, A., Jalal, S., & Kumar, M. (2010). A density based algorithm for discovering density varied
600   clusters in large spatial databases. *International Journal of Computer Applications*, *3*(6), 1–
601   4. https://doi.org/10.13140/RG.2.1.4420.1448

602   Rhodes, M. W., Bennie, J. J., Spalding, A., ffrench-Constant, R. H., & Maclean, I. M. D. (2022).
603   Recent advances in the remote sensing of insects. *Biological Reviews*, *97*(1), 343–360.
604   https://doi.org/10.1111/BRV.12802

605   Rydhmer, K., Bick, E., Still, L., Strand, A., Luciano, R., Helmreich, S., Beck, B., Grønne, C.,
606   Malmros, L., Poulsen, K., Elbæk, F., Brydegaard, M., Lemmich, J., & Nikolajsen, T. (2021).
607   Automating insect monitoring using unsupervised near-infrared sensors. *Scientific Reports*,
608   *12*(1), 2603. https://arxiv.org/abs/2108.05435v1

609   Rydhmer, K., Prangsma, J., Brydegaard, M., Smith, H. G., Kirkeby, C., Kappel Schmidt, I., & Boelt,
610   B. (2022). Scheimpflug lidar range profiling of bee activity patterns and spatial
611   distributions. *Animal Biotelemetry*, *10*(1). https://doi.org/10.1186/S40317-022-00285-Z

Sánchez-Bayo, F., & Wyckhuys, K. A. G. (2019). Worldwide decline of the entomofauna: A review
of its drivers. *Biological Conservation*, *232*, 8–27.
https://doi.org/10.1016/J.BIOCON.2019.01.020

Schipper, A. M., Hilbers, J. P., Meijer, J. R., Antão, L. H., Benítez-López, A., de Jonge, M. M. J.,
Leemans, L. H., Scheper, E., Alkemade, R., Doelman, J. C., Mylius, S., Stehfest, E., van
Vuuren, D. P., van Zeist, W. J., & Huijbregts, M. A. J. (2020). Projecting terrestrial
biodiversity intactness with GLOBIO 4. *Global Change Biology*, *26*(2), 760–771.
https://doi.org/10.1111/GCB.14848

Shortall, R. C., Moore, A., Smith, E., Hall, J. M., Woiwod, P. I., & Harrington, R. (2009). Long-term
changes in the abundance of flying insects. *Insect Conservation and Diversity*, *2*(4), 251–
260. https://doi.org/10.1111/J.1752-4598.2009.00062.X

Silva, D. F., De Souza, V. M. A., Batista GEAPA, K. E., & Ellis, D. P. W. (2013). *Applying machine*
learning and audio analysis techniques to insect recognition in intelligent traps.
Proceedings—2013 12th International Conference on Machine Learning and Applications,
ICMLA 2013. 2013.

Stefanescu, C., Páramo, F., Åkesson, S., Alarcón, M., Ávila, A., Brereton, T., Carnicer, J., Cassar, L.
F., Fox, R., Heliölä, J., Hill, J. K., Hirneisen, N., Kjellén, N., Kühn, E., Kuussaari, M., Leskinen,
M., Liechti, F., Musche, M., Regan, E. C., … Chapman, J. W. (2013). Multi-generational long-
distance migration of insects: studying the painted lady butterfly in the Western
Palaearctic. *Wiley Online Library*, *140*(4), 474–486. https://doi.org/10.1111/j.1600-
0587.2012.07738.x

Tilman, D., May, R. M., Lehman, C. L., & Nowak, M. A. (1994). Habitat destruction and the
extinction debt. *Nature 1994 371:6492*, *371*(6492), 65–66.
https://doi.org/10.1038/371065a0

Wägele, J. W., Bodesheim, P., Bourlat, S. J., Denzler, J., Diepenbroek, M., Fonseca, V., Frommolt,
K. H., Geiger, M. F., Gemeinholzer, B., Glöckner, F. O., Haucke, T., Kirse, A., Kölpin, A.,
Kostadinov, I., Kühl, H. S., Kurth, F., Lasseck, M., Liedke, S., Losch, F., … Wildermann, S.
(2022). Towards a multisensor station for automated biodiversity monitoring. *Basic and*
Applied Ecology, *59*, 105–138. https://doi.org/10.1016/J.BAAE.2022.01.003

Wood, C. R., Reynolds, D. R., Wells, P. M., Barlow, J. F., Woiwod, I. P., & Chapman, J. W. (2009).
Flight periodicity and the vertical distribution of high-altitude moth migration over
southern Britain. *Bulletin of Entomological Research*, *99*, 525–535.
https://doi.org/10.1017/S0007485308006548

Yang, L. H., & Gratton, C. (2014). Insects as drivers of ecosystem processes. *Current Opinion in*
Insect Science, *2*, 26–32. https://doi.org/10.1016/J.COIS.2014.06.004

647
648

Supplementary information

Field name	Crop type	Sensor		Malaise trap		Sweep net	
		June	*July*	*June*	*July*	*June*	*July*
A	ALFALFA	5529	3437	38	1530	254	6579
B	COMPLEX COVER CROP	5739	2244	108	-	79	62
C	COMPLEX COVER CROP	5454	4290	51	828	169	546
D	COMPLEX COVER CROP	3718	704	493	93	525	172
E	COMPLEX COVER CROP	-	240	258	87	13	219
F	COMPLEX COVER CROP	6008	2281	243	303	626	583
G	CORN	-	912	761	278	12	68
H	CORN	-	1158	-	398	303	81
I	CORN	-	687	561	636	427	740
J	PASTURE	-	1993	195	1437	8	24
K	PASTURE	1980	1055	-	64	45	375
L	SORGHUM	8492	5545	68	743	32	304
M	SORGHUM	4845	3080	239	948	254	80
N	SORGHUM	2349	-	155	1176	708	167
O	SOYBEAN	7524	2310	144	101	358	-
P	SOYBEAN	1657	4162	7	225	14	-
Q	SOYBEAN	865	1349	3	1090	77	354
R	SOYBEAN	1349	4679	32	160	713	-
S	SOYBEAN	2696	1867	184	771	307	580
T	SOYBEAN	2443	3442	41	192	-	-

Supplementary Table 1. A table describing the crop type and number of insects observed in each field in June and July across all three methods.

	Sweep net	Malaise traps	Sensors
ALFALFA	N=2, μ=3416.5±4472.5	N=2, μ=784.0±1055.0	N=2, μ=4483.0±1479.3
COMPLEX COVER CROP	N=8, μ=434.4±232.2	N=10, μ=351.0±242.2	N=9, μ=3408.7±2157.2
CORN	N=6, μ=178.3±185.8	N=4, μ=375.5±339.6	N=3, μ=919.0±235.6
PASTURE	N=3, μ=411.0±274.4	N=3, μ=490.7±595.4	N=3, μ=1676.0±537.8
SORGHUM	N=6, μ=131.3±163.2	N=6, μ=451.0±545.2	N=5, μ=4862.2±2404.9
SOYBEANS	N=10, μ=245.9±245.3	N=12, μ=323.6±385.6	N=12, μ=2861.9±1872.7

Supplementary Table 2. Measured insect abundance per crop and monitoring method. Mean and standard deviation.

	Richness combined	Richness sweep net	Shannon sweepnet	Simpson sweepnet	Richness Malaise	Shannon Malaise	Simpson Malaise	No. clusters sensor	Shannon Sensor	Simpson Sensor
Richness combined		**0.73, 0.0**	**0.35, 0.05**	-0.06, 0.74	**0.85, 0.0**	**0.4, 0.02**	-0.23, 0.22	**0.55, 0.01**	0.41, 0.07	0.42, 0.07
Richness sweep net	**0.73, 0.0**		**0.58, 0.0**	0.02, 0.89	**0.36, 0.05**	0.16, 0.39	-0.21, 0.24	**0.48, 0.03**	0.3, 0.19	0.34, 0.13
Shannon sweepnet	**0.35, 0.05**	**0.58, 0.0**		**0.72, 0.0**	0.08, 0.68	-0.14, 0.45	-0.31, 0.09	0.36, 0.11	0.2, 0.39	0.18, 0.44
Simpson sweepnet	-0.06, 0.74	0.02, 0.89	**0.72, 0.0**		-0.14, 0.45	-0.33, 0.07	-0.17, 0.34	0.2, 0.4	0.2, 0.39	0.14, 0.54
Richness Malaise	**0.85, 0.0**	**0.36, 0.05**	0.08, 0.68	-0.14, 0.45		**0.59, 0.0**	-0.03, 0.87	**0.52, 0.01**	0.41, 0.06	**0.42, 0.05**
Shannon Malaise	**0.4, 0.02**	0.16, 0.39	-0.14, 0.45	-0.33, 0.07	**0.59, 0.0**		**0.61, 0.0**	0.26, 0.25	0.21, 0.34	0.28, 0.2

Simpson Malaise	-0.23, 0.22	-0.21, 0.24	-0.31, 0.09	-0.17, 0.34	-0.03, 0.87	**0.61, 0.0**		-0.05, 0.82	0.1, 0.65	0.14, 0.53
No. clusters sensor	**0.55, 0.01**	**0.48, 0.03**	0.36, 0.11	0.2, 0.4	**0.52, 0.01**	0.26, 0.25	-0.05, 0.82		**0.67, 0.0**	**0.62, 0.0**
Shannon Sensor	0.41, 0.07	0.3, 0.19	0.2, 0.39	0.2, 0.39	0.41, 0.06	0.21, 0.34	0.1, 0.65	**0.67, 0.0**		**0.95, 0.0**
Simpson Sensor	0.42, 0.07	0.34, 0.13	0.18, 0.44	0.14, 0.54	**0.42, 0.05**	0.28, 0.2	0.14, 0.53	**0.62, 0.0**	**0.95, 0.0**	

662 **Supplementary Table 3.** Co-correlations of all biodiversity metrics.
663
664

Fitting data	ε	min_samples
Richness Malaise	0.007467	7
Richness sweep net	0.009467	5
Richness combined	0.0092	5
Shannon Malaise	0.007467	7
Shannon sweep net	0.0104	7
Simpson Malaise	0.0084	3
Simpson sweep net	5.33E-04	3

665 **Supplementary Table 4:** Model parameters for each fitted metric.
666
667

 Supplementary Figure 1. A box plot depicting richness metrics from the Malaise traps, sweep
 nets, and sensors by field type.

673
674 **Supplementary Figure 2.** A scatterplot depicting the correlation of the species richness metrics
675 at each field, separated by the June and July timepoints.
676