1 **Robust Ultrasensitive Transcriptional Switch in Noisy Cellular**

2 **Environments**

3
4 **Authors**
5

6 Eui Min Jeong[1,2], and Jae Kyoung Kim[1,2,*]

7 [1]Biomedical Mathematics Group, Institute for Basic Science, Daejeon 34126, Republic of Korea

8 [2]Department of Mathematical Sciences, Korea Advanced Institute of Science and Technology, Daejeon

9 34141, Republic of Korea

10 [*]Correspondence: jaekkim@kaist.ac.kr

12
Abstract

Ultrasensitive transcriptional switches enable cells to make sharp transitions between on and

off states of transcription, essential for cells to respond to environmental cues with high fidelity. However,

conventional switches that rely on direct repressor-DNA binding are extremely sensitive to noise,

leading to unintended changes in gene expression. This raises doubts about the effectiveness of these

switches in noisy cellular environments. Here, we find an alternative avenue, where the combination of

different indirect transcriptional regulations generates a noise-resilient ultrasensitive switch. Specifically,

when a transcriptional repressor inhibits a transcriptional activator through the combination of

sequestration, blocking, and displacement instead of directly binding to DNA, a robust ultrasensitive

switch is achieved. Sequestration alone can generate an ultrasensitive transcriptional response, but the

switch remains vulnerable to noise because the undesired on or off state induced by noise can persist

for a long period. However, by jointly utilizing blocking and displacement, noise-induced transitions can

be rapidly restored to the original desired state. This transcriptional switch is effective in noisy cellular

contexts, which makes it suitable for robust synthetic system design. Our findings also provide insights

into the evolution of ultrasensitive switches in real cells. The concurrent use of seemingly redundant

multiple indirect transcriptional repression mechanisms in diverse biological systems appears to be a

strategy to improve the noise-resilience of ultrasensitive switches.

1. Introduction

Ultrasensitive responses are input/output relationships within biological systems, exhibiting a highly sensitive response to changes in input or stimulus [1]. In particular, transcriptional ultrasensitive switches are turned on or off depending on whether the number of the transcriptional factor exceeds or falls below a threshold. This distinctive property renders these switches valuable for systems requiring precise decision-making contingent on the number of input transcriptional factors. Moreover, these switches serve as catalysts for diverse cellular functions, including signal amplification and generation of bistability and oscillations [2, 3], thereby bestowing them with versatile utility across various biological contexts. The ultrasensitive transcriptional switch can be generated by the cooperative binding of transcriptional factors to multiple DNA binding sites [2, 3] or their titration via decoy sites [4]. However, recent studies have raised doubts about the effectiveness of these conventional transcriptional switches in noisy cellular environments [5, 6]. Specifically, when noise induces transitions between the transcriptional 'on' and 'off' irrespective of the number of the transcriptional factor, this undesired transition persists for a long time in these switches.

Alternatively, transcriptional repressors can suppress transcription indirectly by targeting transcriptional activator rather than directly binding to DNA [7, 8]. Specifically, repressors can sequester transcriptional activators, preventing their binding to DNA. For example, sigma factors or basic leucine zippers can be sequestered through interactions with anti-sigma factors or inhibitors, respectively [9-11]. Also, repressors can bind with activators already bound to DNA, leading to transcriptional inhibition. This blocking is utilized when PHO80, a kinase component in a signaling pathway of yeast, inhibits PHO4 [12]. In addition to sequestration and blocking, repressors can displace DNA-bound activators. An illustrative case is the IκB protein, which reduces NF-κB activator's binding affinity with DNA, displacing it from the DNA [13]. We recently found that the combination of the sequestration, blocking and displacement can generate ultrasensitive transcriptional switch [14]. Interestingly, these three indirect repression mechanisms synergistically function in systems like the NF-κB oscillator and circadian clocks [13, 15-19], where precise ultrasensitivity against noise is crucial. This implies that the ultrasensitive switch based on the combination of multiple indirect repression mechanisms could be robust against noise unlike the conventional switches relying on direct repression.

We found that the ultrasensitive switch based on the combination of the sequestration,

blocking, and displacement can stably sustain the transcriptional 'on' and 'off' states even in the

presence of noise. Specifically, our research has unveiled that the combined integration of these three

repression mechanisms immediately restores the original 'off' and 'on' states even when noise triggers

transitions to the undesired transcriptional states. In this switch, the sequestration generates a sharp

transition from the transcriptional 'on' state to the 'off' state as the molar ratio between repressors and

activators shifts from below one to above one. Furthermore, even if the transcriptional 'off' state transits

to the 'on' state due to noise, the blocking rapidly restores the original 'off' state when the molar ratio

surpasses one. In addition, the displacement ensures that the noise-induced transcriptional 'on' state

swiftly returns to the original 'off' state when the molar ratio falls below one. Taken together, we propose

an ultrasensitive transcriptional switch that can operate effectively even in noisy cellular environments

and can offer promising strategy for designing robust synthetic switches.

72   2. Results

73   2.1. Noise triggers both activation during TRP and repression during TAP of the

74   cooperative binding-based switch.

Ultrasensitivity can be generated through the cooperative binding of transcriptional repressors

to multiple DNA sites [3, 20]. To investigate this mechanism, we used a previously developed model [5]

(Fig. 1A). In the model, the DNA has three sites that can bind with the repressor (R) with a dissociation

constant of K_r. When one site of the DNA is occupied, R can bind to the other two sites with a

dissociation constant of cK_r. Similarly, when two sites of the DNA are occupied, then R can bind to the

remaining site with a dissociation constant of $c^2 K_r$. Therefore, in the case where $c = 1$, R binds

independently to three sites, but in the case where $c < 1$, R more favorably binds to the DNA where

one site or two sites is occupied than the DNA where all sites are unoccupied (i.e., cooperatively binds).

Because when all three sites are occupied by R, the transcription is suppressed (Fig. 1A, gray box),

the transcriptional activity is the probability that at least one site is unoccupied by R at the steady state.

The probability can be calculated from the chemical master equation (CME) governing the system with

respect to the effective total number of the repressor $\widetilde{R_T} = R_T/K_r$, where R_T is the total number of

repressors (Table S1; see the Methods).

The transcription is turned on with a high probability when $\widetilde{R_T}$ is low, referred to as the

transcriptional activation phase (TAP). As $\widetilde{R_T}$ increases, the probability decreases to zero, reaching to

the transcriptional repression phase (TRP; Fig. 1B). The sharpness of this transition from transcriptional

activation during TAP to repression during TRP depends on the cooperativity c. Specifically, when there

is no cooperativity (i.e., $c = 1$), the transition shows a similar sensitivity with the Michaelis-Menten

equation (Fig. 1B, red line). In contrast, when repressors bind cooperatively (i.e., $c < 1$), the sensitivity

of the transition increases (Fig. 1B, blue line). This sensitivity can be quantified using the effective Hill

coefficient $\log 81 / \log(EC_{10}/EC_{90})$, where EC_{90} and EC_{10} are the values of $\widetilde{R_T}$ at which the

transcriptional activity becomes 0.9 and 0.1, respectively [1]. The measured effective Hill coefficients in

the presence and absence of cooperativity are about 3 and 1.3, respectively (Fig. 1B).

Next, we simulated the time series of mRNA during TAP and TRP using the Gillespie algorithm.

We chose the values of $\widetilde{R_T}$ at EC_{90} and EC_{10}, for TAP and TRP, respectively. In the absence of

cooperativity, the timeseries of mRNA slightly fluctuate around their high average during TAP (Fig. 1C)

and around their low average during TRP (Fig. 1D). However, in the presence of cooperativity, the

timeseries of mRNA often drop to zero even during TAP (Fig. 1E), and conversely often lift up to high

levels even during TRP (Fig. 1F). These undesired transcriptional states, arising due to noise, persists

because transition between the transcriptional activation and repression occurs rarely due to

cooperative binding [5]. As a result, cooperativity increases the variance in the number of mRNA,

yielding a much higher Fano factor compared to the absence of cooperativity (Fig. 1G and Table S1).

2.2. Noise turns on the sequestration-based switch during TRP.

The transcriptional repression can occur not only through direct binding to DNA, but also by

indirectly inhibiting the transcriptional activator that promotes transcription. Among indirect repression

mechanisms, the sequestration has been identified to generate ultrasensitivity [21-24]. To investigate

whether this switch is robust to noise, we constructed a model describing the sequestration (Fig. 2A).

In the model, the free activator (A) binds to the free DNA (E_F) with a dissociation constant of K_a to form

the activated DNA (E_A). The transcription is inhibited as the free repressor (R) binds to A to form an

inactive complex (R_A) with a dissociation constant of K_s (i.e., sequestration; Fig. 2A, gray box). Thus,

the transcriptional activity is defined as the probability that the DNA is activated by the binding of A, not

sequestered by R at the steady state. We derived the probability as a function of $\widetilde{R_T} = R_T/A_T$, the

molar ratio between total numbers of the activator ($A_T = A + R_A + E_A$) and the repressor ($R_T = R + R_A$)

with two scaled parameters $\widetilde{K_a} = K_a/A_T$ and $\widetilde{K_s} = K_s/A_T$ (Table S1; see Method for details).

The transcription turns on and off, respectively, when the molar ratio $\widetilde{R_T}$ is either below (TAP)

or above one (TRP). The transition between transcriptional activation during TAP and repression during

TRP becomes more sensitive as the binding between A and the DNA becomes stronger (i.e., $\widetilde{K_a}$

decreases; Fig. 2B). This is because strong activator (i.e., small $\widetilde{K_a}$) can promote the transcription even

with a small number of activators, sustaining transcriptional activation during TAP as long as there are

more activators than repressors (i.e., $\widetilde{R_T} < 1$; Fig. 2B, blue line). However, when the molar ratio

exceeds one (i.e., $\widetilde{R_T} > 1$; Fig. 2B), the majority of A is sequestered by R. As a result, a sensitive

transition from TAP to TRP occurs near $\widetilde{R_T} = 1$ with the high effective Hill coefficient of about 50. On

the other hand, the transition occurs gradually with the effective Hill coefficient of about 2 in the case of

weak activator (i.e., large $\widetilde{K_a}$; Fig. 2B, red line).

Next, we simulated the timeseries of activated DNA and mRNA at the value of $\widetilde{R_T} = 10^{0.5}$ in

TRP (Fig. 2B, orange star). For a weak activator, the activated DNA indued by noise is swiftly restored

to the repressed state, keeping the number of mRNA consistently low around zero (Fig. 2C). On the

other hand, for a strong activator, this undesired activation of DNA during TRP persists over a long

period, leading to a significant increase in the number of mRNA (Fig. 2D). This disparity arises when a

sequestered A is released from R and binds with the DNA despite of being TRP (unexpected activation;

Fig. 2E). Then, a week activator quickly dissociates from the DNA itself and thus the transcription is

turned off to its original state (Fig. 2E, red arrow). Meanwhile, a strong activator remains bound to the

DNA for a long time (Fig. 2E, blue arrow). As a result, the undesired transcriptional activation persists

even during TRP, resulting in a considerable increase in the number of mRNA. This leads to a bimodal

distribution of mRNA molecules deviating substantially from a Poisson distribution, with a strong

activator (Fig. 2F, blue bars) unlike a weak activator (Fig. 2F, red bars). Thus, a strong activator yields

much higher transcriptional noise compared to a weak activator during TRP (Fig. 2G). In conclusion,

when the binding between the activator and DNA is tight, a sensitive transcriptional response can be

generated with the sole sequestration. However, this tight binding leads to high transcriptional noise as

the undesired transcriptional activation during TRP persists for a long period.

2.3. Noise turns off the sequestration and blocking-based switch during TAP

When transcription is regulated by the sole sequestration, high transcriptional noise occurs

during TRP due to the absence of repression for the activator bound on DNA. This DNA-bound-activator

can be suppressed when the repressor blocks the transcription by forming a complex with the DNA-

bound-activator (i.e., blocking) or pulls off the activator from the DNA (i.e., displacement). Indeed, these

repression mechanisms are commonly employed in addition to the sole sequestration in many biological

systems [7, 14]. We first added the blocking to the sole sequestration model, where the repressor (R)

can directly bind to the DNA-bound-activator (E_A) with a dissociation constant of K_b to form the

repressed DNA (E_R; Fig. 3A, gray box). Based on this model, we derived the transcriptional activity as

function of the molar ratio $\widetilde{R_T}$ with three scaled parameters $\widetilde{K_a} = K_a/A_T$, $\widetilde{K_s} = K_s/A_T$ and $\widetilde{K_b} = K_b/A_T$

(Table S1; see Method for details). The derived transcriptional activity exhibits a high sensitivity of

response to changes in $\widetilde{R_T}$ (i.e., the effective Hill coefficient of about 50; Fig. 3B). Furthermore, the

addition of blocking rapidly restores the undesired activation of DNA during TRP to the repressed state

(Fig. 3C, top panel). This is evident in the simulated timeseries of mRNA, which remain close to zero,

without sharp increase of mRNA (Fig. 3C, bottom right panel) unlike the sole sequestration (Fig. 3C,

bottom left panel). That is, the transcription triggered by noise-induced activator binding to DNA during

TRP is immediately inhibited by the repressor via blocking (Fig. 3D, blue arrow) in contrast to the sole

sequestration (Fig. 3D, red arrow). As a result, in the presence of the blocking, the distribution of mRNA

molecules is concentrated near zero (Fig. 3E, blue bars) rather than bimodal (Fig. 3E, red bars),

reducing the transcriptional noise during TRP (Fig. 3F and Table S1).

However, the addition of blocking dramatically increases the transcriptional noise during TAP

(Fig. 3F). To investigate the source of such high transcriptional noise during TAP, we simulated the

timeseries of non-activated DNA and mRNA at the value of $\widetilde{R_T} = 10^{-0.1}$ in TAP (Fig. 3C, gray star). In

the case of the sole sequestration, non-activated DNA is restored to the activated state rapidly (Fig. 3G,

top left panel), and thus the timeseries of mRNA maintain their high average during TAP (Fig. 3G,

bottom left panel). In contrast, when the blocking is added, this undesired repression during TAP

persists for a long period (Fig. 3G, top right panel), causing a dramatic reduction in the number of mRNA

(Fig. 3G, bottom right panel). This significant reduction arises when the repressor, released from the

activator due to noise, binds with the activator bound to DNA during TAP (unexpected repression; Fig.

3H). Then, the repressor remains bound until it unbinds from DNA-bound-activator itself (Fig. 3H). This

causes a dramatic reduction in the number of mRNA even during TAP, leading to a small additional

peak around zero in the mRNA distribution (Fig. 3I). Due to the bimodality, the transcriptional noise

surges during TAP (Fig. 3F).

181   2.4. The sequestration, blocking, and displacement-based switch is robust to noise

In the presence of blocking, the undesired repression during TAP stems from the unexpected

binding of the repressor to the activator on DNA. This can be prevented by the addition of displacement,

which facilitates dissociation of the DNA-bound repressor-activator complex from the DNA. To

investigate this, we incorporated the displacement into the blocking and sequestration model. In this

model, the repressor-activator complex (R_A) can dissociate from the DNA with a dissociation constant

of K_d (Fig. 4A, gray box). We derived the transcriptional activity as the function of the molar ratio $\widetilde{R_T}$

with four scaled parameters $\widetilde{K_a} = K_a/A_T$, $\widetilde{K_s} = K_s/A_T$, $\widetilde{K_b} = K_b/A_T$ and $\widetilde{K_d} = K_d/A_T$ (Table S1; see

Methods for details). The transcriptional activity can exhibit an ultrasensitive response to changes in

$\widetilde{R_T}$ with the effective Hill coefficient of about 200 (Fig. 4B, blue and red lines).

Furthermore, the addition of displacement rapidly restores the undesired repression of DNA

during TAP to the activated state (Fig. 4C, top panel). This is evident in the simulated timeseries of

mRNA, which do not drop to zero (Fig. 4C, bottom right panel) unlike the combination of the

sequestration and blocking. That is, when a repressor binds to the DNA-bound-activator due to noise

during TAP (unexpected repression; Fig. 4), repressor-activator complex is immediately released from

DNA via displacement (Fig. 4D, blue arrow). This does not happen without displacement (Fig. 4D, red

arrow). Taken together, the inclusion of displacement eliminates the additional peak near zero seen in

the sequestration and blocking model (Fig. 4E), leading to a reduction in transcriptional noise during

TAP (Fig. 4F and Table S1).

201 3. Discussion

Traditionally, ultrasensitive transcriptional responses are generated by cooperative binding of

transcriptional repressors to multiple DNA sites. This mechanism can trigger a sharp transcriptional

transition from transcriptional repression during TRP to activation during TAP as the number of

repressors surpasses a threshold. However, the presence of noise can lead to prolonged periods of

undesired repression and activation during TAP and TRP, respectively (Fig. 1). Thus, we have shifted

our focus from direct repression—where repressors directly bind to DNA, as seen in cooperative binding

to multiple DNA sites—to the realm of indirect transcriptional repression: sequestration, blocking, and

displacement. Sequestration alone can generate an ultrasensitive response that is attuned to changes

in the molar ratio between repressors and activators. Specifically, when the number of repressors falls

short of that of activators (i.e., the molar ratio is less than one), an unsequestered activator binds to

DNA, thereby promoting transcription. As the molar ratio surpasses one, a majority of activators become

sequestered by repressors, promoting a sharp transition to transcriptional inhibition (Fig. 2).

Furthermore, the transcriptional repression (activation) during TRP (TAP) can be stably maintained by

blocking (displacement) even in the presence of noise. Specifically, when an activator, unexpectedly

released from repressors, binds to DNA and initiates transcription during TRP (Fig. 2), the activator is

immediately blocked by a repressor, resulting in the prompt restoration of transcriptional repression (Fig.

3). When a repressor unexpectedly blocks an activator that is already bound to DNA during TAP, the

repressor-activator complex is displaced, swiftly reinitiating transcriptional activation (Fig. 4). Taken

together, our exploration has unveiled an ultrasensitive transcriptional switch endowed with resilience

against noise during both TAP and TRP. This breakthrough holds the potential to enhance the stability

and reliability of transcriptional regulation within intricate biological systems.

223   Interestingly, although two models with or without a particular repression mechanism may

appear similar in deterministic simulations, they can behave quite differently in stochastic simulations.

Specifically, in the deterministic ODE, where the transcription of mRNA is proportional to the

transcriptional activity, two models exhibiting similar transcriptional activity yield indistinguishable

behaviors of mRNA [14]. On the other hand, in the stochastic models, the distributions of mRNA can

be completely different despite resembling transcriptional activities. For example, all transcriptional

activity of each model shows similar responses with respect to changes in the molar ratio (Fig. 3B and

4B), whereas their Fano factors are completely different (Fig. 3F and 4F). Therefore, a comprehensive

understanding of the precise repression mechanism is essential to ensure unbiased modeling results

when considering the presence of noise.

233   Biological oscillators and biological switches require ultrasenstivity to generate strong rhythms

[24, 25] and bistable responses [2], respectively. In addition to the ultrasenstivity, these systems need

to exhibit robustness against intrinsic noise to ensure accurate timing and precise decision-making.

Indeed, such biological systems that require both ultrasensitivity and precision employ the multiple

repression mechanisms. For instance, the MDM2-p53 oscillator is known to utilize the blocking and

displacement to inhibit p53 by MDM2 [26, 27]. Similarly, the NF-kB oscillator, and circadian clocks in

mammals and Drosophila employ a combination of the sequestration, blocking, and displacement [13,

15-19]. Moreover, the Rb-EF bistable switches in the cell cycle incorporates the blocking and chromatin

structure modification [28-30]. Our work highlights the significance of the multiple repression

mechanisms in achieving robust and precise regulatory outcomes in these systems.

244 **4. Methods**

245   We used a approach established by A. Sanchez et al [5] to calculate the transcriptional activity

and the Fano factor for the number of mRNA (see SI for details). The results are summarized in Table

S1. The transcriptional activity is the probability that the transcription of DNA is activated. The

transcriptional activity and Fano factors of all models depending on the scaled repressor level ($\widetilde{R_T}$).

Specifically, the transcriptional activity ($TA(\widetilde{R_T})$; Fig. 1B) and Fano factor ($FF(\widetilde{R_T})$; Fig. 1G) for the three

binding sites model with cooperativity (Fig. 1A) depend on the effective total repressor $\widetilde{R_T} = R_T/K_r$,

where R_T is the total number of repressors and K_r is the dissociation constant between repressor and

binding sites on DNA:

$$TA(\widetilde{R_T}) = 1 - \frac{\widetilde{R_T}^3}{c^3 + 3c^3\widetilde{R_T} + 3c^2\widetilde{R_T}^2 + \widetilde{R_T}^3},$$

254 and

$$FF(\widetilde{R_T}) = 1 + \frac{\tilde{\alpha}\widetilde{R_T}^4\left[3(\tilde{\beta}+2)(\tilde{\beta}+3)\widetilde{R_T}^4 + [3c\tilde{\beta}^2 + (24c+3)\tilde{\beta} + 45c]\widetilde{R_T}^3 + [c\tilde{\beta}^2 + 3c(2c+3)\tilde{\beta} + 2c(9c+10)]\widetilde{R_T}^2 + c[(1+2c)\tilde{\beta} + (1+12c)]\widetilde{R_T} + 2c^2\right]}{\begin{array}{l}3(\tilde{\beta}+1)(\tilde{\beta}+2)(\tilde{\beta}+3)\widetilde{R_T}^8 + 3[c(3c+1)\tilde{\beta}^3 + (21c^2+8c+1)\tilde{\beta}^2 + (48c^2+17c+1)\tilde{\beta} + 6c(1+6c)]\widetilde{R_T}^7 \\ + c[6(1+36c^2+27c^3) + (20+36c+315c^2+135c^3)\tilde{\beta} + 3(3+5c+45c^2+9c^3)\tilde{\beta}^2 + (1+18c^2)\tilde{\beta}^3]\widetilde{R_T}^6 \\ + c[3c^2(3c+2)\tilde{\beta}^3 + (54c^4+90c^3+57c^2+2c+1)\tilde{\beta}^2 + (324c^4+252c^3+126c^2+12c+1)\tilde{\beta} + 18c^2(27c^2+6c+4)]\widetilde{R_T}^5 \\ + c^2[6c^2\tilde{\beta}^3 + 3c(9c^3+12c^2+19c+2)\tilde{\beta}^2 + (243c^4+198c^3+153c^2+9c+2)\tilde{\beta} + 18c^2(27c^2+9c+4)]\widetilde{R_T}^4 \\ + c^4[\tilde{\beta}^3 + 6(3c^2+2c+2)\tilde{\beta}^2 + (54c^3+153c^2+72c+29)\tilde{\beta} + 6(27c^3+54c^2+2)]\widetilde{R_T}^3 \\ + c^4[(3c^2+2c+1)\tilde{\beta}^2 + (36c^3+33c^2+18c+1)\tilde{\beta} + 54c^2(1+3c)]\widetilde{R_T}^2 + c^5[(6c^2+3c+2)\tilde{\beta} + 54c^2]\widetilde{R_T} + 6c^7\end{array}},$$

256 where $\tilde{\alpha}$ is the scaled production rate of mRNA, $\tilde{\beta}$ is the scaled production rate of mRNA, and c is

257 cooperativity between binding sites (Fig. 1B).

258   The transcriptional activity (Fig. 2B) and Fano factor (Fig. 2G) for the model describing the

259 sequestration (Fig. 2) depends on the molar ratio $\widetilde{R_T} = R_T/A_T$ between the total numbers of repressors

260 (R_T) and activators (A_T):

$$TA(\widetilde{R_T}) = \frac{\tilde{A}(\widetilde{R_T}, \widetilde{K_s})/\widetilde{K_a}}{1 + \tilde{A}(\widetilde{R_T}, \widetilde{K_s})/\widetilde{K_a}},$$

262 and

$$FF(\widetilde{R_T}) = 1 + \frac{\tilde{\alpha}}{\left(1 + \tilde{A}(\widetilde{R_T}, \widetilde{K_s})/\widetilde{K_a}\right)\left(\tilde{A}(\widetilde{R_T}, \widetilde{K_s}) + \widetilde{K_a} + \tilde{\beta}\right)},$$

264 where $\tilde{A}(\widetilde{R_T}, \widetilde{K_s}) = \frac{1 - \widetilde{R_T} - \widetilde{K_s} + \sqrt{(1 - \widetilde{R_T} - \widetilde{K_s})^2 + 4\widetilde{K_s}}}{2}$ is the fraction of free activator among total activator, and $\widetilde{K_s}$

265 and $\widetilde{K_a}$ are the scaled dissociation constants between repressor and activator and between activator

266 and DNA, respectively.

267   The transcriptional activity (Fig. 3B) and Fano factor (Fig. 3F) for the model describing the

268   sequestration and blocking (Fig. 3) depends on the molar ratio $\widetilde{R_T}$:

269
$$TA(\widetilde{R_T}) = \frac{\frac{\tilde{A}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_a}}}{1+\frac{\tilde{A}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_a}}+\frac{\tilde{A}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_a}}\frac{\tilde{R}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_b}}},$$

270   and

271
$$FF(\widetilde{R_T}) = 1 + \frac{\tilde{\alpha}\left[\widetilde{K_b}+\tilde{\beta}+(\tilde{A}(\widetilde{R_T},\widetilde{K_s})+\tilde{\beta})\frac{\tilde{A}(\widetilde{R_T},\widetilde{K_s})\tilde{R}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_a}}\right]}{\left(1+\frac{\tilde{A}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_a}}+\frac{\tilde{A}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_a}}\frac{\tilde{R}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_b}}\right)[\widetilde{K_a}(\widetilde{K_b}+\tilde{\beta})+(\tilde{A}(\widetilde{R_T},\widetilde{K_s})+\tilde{\beta})(\tilde{R}(\widetilde{R_T},\widetilde{K_s})+\widetilde{K_b}+\tilde{\beta})]},$$

272   where $\tilde{R}(\widetilde{R_T},\widetilde{K_s}) = \frac{\widetilde{R_T}-1-\widetilde{K_s}+\sqrt{(1-\widetilde{R_T}-\widetilde{K_s})^2+4\widetilde{K_s}}}{2}$ is the fraction of free repressor to the total activator and $\widetilde{K_b}$

273   is the scaled dissociation constant between repressor and DNA-bound-activator.

274   The transcriptional activity (Fig. 4B) and Fano factor (Fig. 4F) for the model describing the

275   sequestration, blocking, and displacement (Fig. 4) depends on the molar ratio $\widetilde{R_T}$:

276
$$TA(\widetilde{R_T}) = \frac{\tilde{I}(\widetilde{R_T},\widetilde{K_a},\widetilde{K_s})\frac{\tilde{A}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_a}}}{1+\tilde{I}(\widetilde{R_T},\widetilde{K_a},\widetilde{K_s})\frac{\tilde{A}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_a}}+\tilde{J}(\widetilde{R_T},\widetilde{K_a},\widetilde{K_s})\frac{\tilde{A}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_a}}\frac{\tilde{R}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_b}}},$$

277   and

278 $FF(\widetilde{R_T}) = 1 + \dfrac{\tilde{\alpha}\left[(\tilde{R}_A(\widetilde{R_T},\widetilde{K_s})+\widetilde{K_b}+\widetilde{K_d}+\tilde{\beta})+(1+\widetilde{K_d}+\tilde{\beta})\tilde{J}(\widetilde{R_T},\widetilde{K_a},\widetilde{K_s})\frac{\tilde{A}(\widetilde{R_T},\widetilde{K_s})\tilde{R}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_a}}\right]}{\left(1+\tilde{I}(\widetilde{R_T},\widetilde{K_a},\widetilde{K_s})\frac{\tilde{A}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_a}}+\tilde{J}(\widetilde{R_T},\widetilde{K_a},\widetilde{K_s})\frac{\tilde{A}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_a}}\frac{\tilde{R}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_b}}\right)[(\widetilde{K_b}+\widetilde{K_d}+\tilde{\beta})(\tilde{A}(\widetilde{R_T},\widetilde{K_s})+\widetilde{K_a}+\tilde{\beta})+(\widetilde{K_b}+\widetilde{K_a}+\tilde{\beta})\tilde{R}_A(\widetilde{R_T},\widetilde{K_s})+(1+\widetilde{K_d}+\tilde{\beta})\tilde{R}(\widetilde{R_T},\widetilde{K_s})]}$,

279   where $\tilde{I}(\widetilde{R_T},\widetilde{K_a},\widetilde{K_s}) = \frac{\widetilde{K_s}+\sigma\widetilde{K_a}+\tilde{R}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_s}+\sigma\widetilde{K_a}+\sigma\tilde{R}(\widetilde{R_T},\widetilde{K_s})}$, $\tilde{J}(\widetilde{R_T},\widetilde{K_a},\widetilde{K_s}) = \frac{\widetilde{K_s}+\sigma\widetilde{K_a}+\sigma\tilde{R}(\widetilde{R_T},\widetilde{K_s})}{\widetilde{K_s}+\sigma\widetilde{K_a}+\sigma\tilde{R}(\widetilde{R_T},\widetilde{K_s})}$, $\sigma = \frac{\widetilde{K_s}\widetilde{K_d}}{\widetilde{K_a}\widetilde{K_b}}$, and $\widetilde{K_d}$ is the scaled

280   dissociation constant between repressor-activator complex and DNA.

282 Data availability

283   The datasets used and/or analyzed during the current study available from the corresponding author

284   on reasonable request.

286 Code availability

287   The underlying code for this study is not publicly available but may be made available to qualified

288   researchers on reasonable request from the corresponding author.

290   **References**

291   1.   Ferrell JE, Jr., Ha SH. Ultrasensitivity part I: Michaelian responses and zero-order
292   ultrasensitivity. Trends Biochem Sci. 2014;39(10):496-503. Epub 2014/09/23. doi:
293   10.1016/j.tibs.2014.08.003. PubMed PMID: 25240485; PubMed Central PMCID:
294   PMCPMC4214216.
295   2.   Ferrell JE, Jr., Ha SH. Ultrasensitivity part III: cascades, bistable switches, and oscillators.
296   Trends Biochem Sci. 2014;39(12):612-8. Epub 2014/12/03. doi: 10.1016/j.tibs.2014.10.002.
297   PubMed PMID: 25456048; PubMed Central PMCID: PMCPMC4254632.
298   3.   Zhang Q, Bhattacharya S, Andersen ME. Ultrasensitive response motifs: basic amplifiers in
299   molecular signalling networks. Open Biol. 2013;3(4):130031. Epub 20130424. doi:
300   10.1098/rsob.130031. PubMed PMID: 23615029; PubMed Central PMCID:
301   PMCPMC3718334.
302   4.   Lu MS, Mauser JF, Prehoda KE. Ultrasensitive synthetic protein regulatory networks using
303   mixed decoys. ACS Synth Biol. 2012;1(2):65-72. doi: 10.1021/sb200010w. PubMed PMID:
304   22639735; PubMed Central PMCID: PMCPMC3358930.
305   5.   Sanchez A, Garcia HG, Jones D, Phillips R, Kondev J. Effect of promoter architecture on the
306   cell-to-cell variability in gene expression. PLoS Comput Biol. 2011;7(3):e1001100. Epub
307   20110303. doi: 10.1371/journal.pcbi.1001100. PubMed PMID: 21390269; PubMed Central
308   PMCID: PMCPMC3048382.
309   6.   Dey S, Soltani M, Singh A. Enhancement of gene expression noise from transcription factor
310   binding to genomic decoy sites. Sci Rep. 2020;10(1):9126. Epub 20200604. doi:
311   10.1038/s41598-020-65750-2. PubMed PMID: 32499583; PubMed Central PMCID:
312   PMCPMC7272470.
313   7.   Gaston K, Jayaraman PS. Transcriptional repression in eukaryotes: repressors and
314   repression mechanisms. Cell Mol Life Sci. 2003;60(4):721-41. Epub 2003/06/06. doi:
315   10.1007/s00018-003-2260-3. PubMed PMID: 12785719.
316   8.   Renkawitz R. Transcriptional repression in eukaryotes. Trends Genet. 1990;6(6):192-7. Epub
317   1990/06/01. doi: 10.1016/0168-9525(90)90176-7. PubMed PMID: 2196723.
318   9.   Benezra R, Davis RL, Lockshon D, Turner DL, Weintraub H. The protein Id: a negative
319   regulator of helix-loop-helix DNA binding proteins. Cell. 1990;61(1):49-59. doi: 10.1016/0092-
320   8674(90)90214-y. PubMed PMID: 2156629.
321   10.   Van Doren M, Ellis HM, Posakony JW. The Drosophila extramacrochaetae protein
322   antagonizes sequence-specific DNA binding by daughterless/achaete-scute protein
323   complexes. Development. 1991;113(1):245-55. doi: 10.1242/dev.113.1.245. PubMed PMID:
324   1764999.
325   11.   Chadsey MS, Karlinsey JE, Hughes KT. The flagellar anti-sigma factor FlgM actively
326   dissociates Salmonella typhimurium sigma28 RNA polymerase holoenzyme. Genes Dev.
327   1998;12(19):3123-36. doi: 10.1101/gad.12.19.3123. PubMed PMID: 9765212; PubMed
328   Central PMCID: PMCPMC317189.
329   12.   Jayaraman PS, Hirst K, Goding CR. The activation domain of a basic helix-loop-helix protein
330   is masked by repressor interaction with domains distinct from that required for transcription
331   regulation. EMBO J. 1994;13(9):2192-9. Epub 1994/05/01. PubMed PMID: 8187772; PubMed
332   Central PMCID: PMCPMC395073.
333   13.   Potoyan DA, Zheng W, Komives EA, Wolynes PG. Molecular stripping in the NF-
334   kappaB/IkappaB/DNA genetic regulatory network. Proc Natl Acad Sci U S A.
335   2016;113(1):110-5. Epub 2015/12/25. doi: 10.1073/pnas.1520483112. PubMed PMID:
336   26699500; PubMed Central PMCID: PMCPMC4711861.
337   14.   Jeong EM, Song YM, Kim JK. Combined multiple transcriptional repression mechanisms
338   generate ultrasensitivity and oscillations. Interface Focus. 2022;12(3):20210084. Epub

20220415. doi: 10.1098/rsfs.2021.0084. PubMed PMID: 35450279; PubMed Central PMCID:
PMCPMC9010851.
15. Bergqvist S, Alverdi V, Mengel B, Hoffmann A, Ghosh G, Komives EA. Kinetic enhancement
of NF-kappaBxDNA dissociation by IkappaBalpha. Proc Natl Acad Sci U S A.
2009;106(46):19328-33. Epub 2009/11/06. doi: 10.1073/pnas.0908797106. PubMed PMID:
19887633; PubMed Central PMCID: PMCPMC2780789.
16. Menet JS, Abruzzi KC, Desrochers J, Rodriguez J, Rosbash M. Dynamic PER repression
mechanisms in the Drosophila circadian clock: from on-DNA to off-DNA. Genes Dev.
2010;24(4):358-67. Epub 2010/02/18. doi: 10.1101/gad.1883910. PubMed PMID: 20159956;
PubMed Central PMCID: PMCPMC2816735.
17. Cao X, Yang Y, Selby CP, Liu Z, Sancar A. Molecular mechanism of the repressive phase of
the mammalian circadian clock. Proc Natl Acad Sci U S A. 2021;118(2). Epub 2021/01/15.
doi: 10.1073/pnas.2021174118. PubMed PMID: 33443219; PubMed Central PMCID:
PMCPMC7812753.
18. Chiou YY, Yang Y, Rashid N, Ye R, Selby CP, Sancar A. Mammalian Period represses and
de-represses transcription by displacing CLOCK-BMAL1 from promoters in a Cryptochrome-
dependent manner. Proc Natl Acad Sci U S A. 2016;113(41):E6072-E9. Epub 2016/10/01.
doi: 10.1073/pnas.1612917113. PubMed PMID: 27688755; PubMed Central PMCID:
PMCPMC5068302.
19. Ye R, Selby CP, Chiou YY, Ozkan-Dagliyan I, Gaddameedhi S, Sancar A. Dual modes of
CLOCK:BMAL1 inhibition mediated by Cryptochrome and Period proteins in the mammalian
circadian clock. Genes Dev. 2014;28(18):1989-98. Epub 2014/09/18. doi:
10.1101/gad.249417.114. PubMed PMID: 25228643; PubMed Central PMCID:
PMCPMC4173159.
20. Ferrell JE, Jr., Ha SH. Ultrasensitivity part II: multisite phosphorylation, stoichiometric
inhibitors, and positive feedback. Trends Biochem Sci. 2014;39(11):556-69. Epub 20141023.
doi: 10.1016/j.tibs.2014.09.003. PubMed PMID: 25440716; PubMed Central PMCID:
PMCPMC4435807.
21. Buchler NE, Cross FR. Protein sequestration generates a flexible ultrasensitive response in a
genetic network. Mol Syst Biol. 2009;5:272. Epub 2009/05/21. doi: 10.1038/msb.2009.30.
PubMed PMID: 19455136; PubMed Central PMCID: PMCPMC2694680.
22. Buchler NE, Louis M. Molecular titration and ultrasensitivity in regulatory networks. J Mol Biol.
2008;384(5):1106-19. Epub 2008/10/22. doi: 10.1016/j.jmb.2008.09.079. PubMed PMID:
18938177.
23. Kim JK, Forger DB. A mechanism for robust circadian timekeeping via stoichiometric balance.
Mol Syst Biol. 2012;8:630. Epub 2012/12/06. doi: 10.1038/msb.2012.62. PubMed PMID:
23212247; PubMed Central PMCID: PMCPMC3542529.
24. Kim JK. Protein sequestration versus Hill-type repression in circadian clock models. IET Syst
Biol. 2016;10(4):125-35. Epub 2016/07/23. doi: 10.1049/iet-syb.2015.0090. PubMed PMID:
27444022.
25. Novak B, Tyson JJ. Design principles of biochemical oscillators. Nat Rev Mol Cell Biol.
2008;9(12):981-91. Epub 2008/10/31. doi: 10.1038/nrm2530. PubMed PMID: 18971947;
PubMed Central PMCID: PMCPMC2796343.
26. Cross B, Chen L, Cheng Q, Li B, Yuan ZM, Chen J. Inhibition of p53 DNA binding function by
the MDM2 protein acidic domain. J Biol Chem. 2011;286(18):16018-29. Epub 2011/04/02.
doi: 10.1074/jbc.M111.228981. PubMed PMID: 21454483; PubMed Central PMCID:
PMCPMC3091211.
27. Chen J. The Cell-Cycle Arrest and Apoptotic Functions of p53 in Tumor Initiation and
Progression. Cold Spring Harb Perspect Med. 2016;6(3):a026104. Epub 2016/03/05. doi:
10.1101/cshperspect.a026104. PubMed PMID: 26931810; PubMed Central PMCID:
PMCPMC4772082.
28. Nevins JR. E2F: a link between the Rb tumor suppressor protein and viral oncoproteins.
Science. 1992;258(5081):424-9. Epub 1992/10/16. doi: 10.1126/science.1411535. PubMed
PMID: 1411535.
29. Yao G, Lee TJ, Mori S, Nevins JR, You L. A bistable Rb-E2F switch underlies the restriction
point. Nat Cell Biol. 2008;10(4):476-82. Epub 20080323. doi: 10.1038/ncb1711. PubMed
PMID: 18364697.
30. Dimova DK, Dyson NJ. The E2F transcriptional network: old acquaintances with new faces.
Oncogene. 2005;24(17):2810-26. doi: 10.1038/sj.onc.1208612. PubMed PMID: 15838517.

398
399
Acknowledgement

This study was funded by Institute for Basic Science (IBS-R029-C3) (J.K.K.), and the Human

Frontiers Science Program Organization (RGY0063/2017) (J.K.K.). The funder played no role in study

design, data collection, analysis and interpretation of data, or writing of this manuscript.

Authors Contribution

All authors provided conceptualization of the manuscript, performed visualization and formal analysis,

participated in investigation, and wrote the paper. E.M.J. contributed to the methodology. J.K.K.

acquired funding and supervised the work conducted here.

Competing Interests

All authors declare no financial or non-financial competing interests.

Supplementary Information

Supplementary tables: Table S1.

415
416

418
419
Figure 1. The transcriptional ultrasensitive switch based on cooperative binding suffers from
undesired switch transition due to noise. (A) Schematic diagram of the model describing the binding
of the repressor (R) to three sites on DNA. R can bind to an unoccupied site (E_{000}) with a dissociation
constant of K_r. When one site is occupied (E_{001}, E_{010}, or E_{100}), R can bind to the other two sites with
a dissociation constant of cK_r. When two sites are occupied (E_{110}, E_{101}, or E_{011}), R can bind to the
remaining site with a dissociation constant of c^2K_r. Accordingly, when $c < 1$, cooperativity is present,
while absent when $c = 1$. When all sites are occupied (E_{111}), the transcription is repressed (gray box);
otherwise, mRNA is transcribed with the rate of α and decays with the rate of β. (B) Consequently, as
the total effective number of the repressor ($\widetilde{R_T}$) increases, the probability that the transcription is
activated (i.e., transcriptional activity) decreases, transitioning from transcriptional activation phase
(TAP) to transcriptional repression phase (TRP). Such transition in the presence of cooperativity (blue
line) is more sensitive to changes in $\widetilde{R_T}$ (i.e., a shorter distance between EC_{90} and EC_{10}) compared to
in the absence of the cooperativity (red line). (C − D) In the absence of cooperativity, the simulated
timeseries of mRNA fluctuate around their high and low averages during TAP (i.e., when $\widetilde{R_T}$ is at EC_{90})
(C) and during TRP (i.e., when $\widetilde{R_T}$ is at EC_{10}) (D), respectively. (E − F) On the other hand, in the
presence of cooperativity, the number of mRNA can decrease to 0 even during TAP and can
significantly increase even during TRP (F), respectively. (G) As a result, in the presence of cooperativity,
the number of mRNA exhibits greater variance, resulting in a higher Fano factor compared to the
absence of cooperativity.

439
440
Figure 2. Transcriptional ultrasensitive switch based on the sole sequestration is sensitive to
noise during TRP due to undesired transcriptional activation. (A) Schematic diagram of the model
describing the sequestration. The binding of the activator (A) to the DNA (E_F) with a dissociation
constant of K_a forms the activated DNA (E_A) to promote the transcription of mRNA with the rate of α,
which decays with the rate of β. On the other hand, the binding of the repressor (R) to A with a
dissociation constant of K_s leads to the transcriptional repression (i.e., sequestration; gray box). (B)
Consequently, when the molar ratio $\widetilde{R_T} = R_T/A_T$ between the total activator (A_T) and repressor (R_T) is
less than one (TAP), unsequestered activators can turn on transcription. In contrast, when the molar
ratio exceeds one (TRP), almost all activators are sequestered by repressors, and thus transcription is
turned off. As a result, around $\widetilde{R_T} = 1$, a sensitive transition from TAP to TRP occurs with respect to
changes in $\widetilde{R_T}$. Notably, strong activators, which bind tightly to DNA, exhibit a more sensitive transition
compared to weak activators that bind loosely. (C – D) The undesired activation is restored during TRP
to the repressed state for a weak activator (C), unlike a strong activator (D). Here, the timeseries of
activated DNA and mRNA are simulated at $\widetilde{R_T} = 10^{0.5}$ (orange star in (B)). (E) When noise-induced
unbinding of A from the R_A complex occurs, the transcription can be turned on by binding for the A to
DNA (i.e, unexpected activation occurs). Such undesired activation during TRP persists for a long time
due to the tight binding of a strong activator (blue arrow), but not a weak activator (red arrow). This
sharply increases the number of mRNA. (F) As a result, the stationary distribution of mRNA molecules
shows an additional peak far from zero, leading to bimodality with a strong activator (blue bars) unlike
a weak activator (red bars). (G) Such bimodality deviates the distribution from a Poisson distribution,
resulting in a higher Fano factor for a strong activator compared to a weak activator during TRP.
462

463
464
Figure 3. Transcriptional ultrasensitive switch based on the combination of sequestration and
blocking is sensitive to noise during TAP due to undesired transcriptional repression. (A) The

blocking is added to the sole sequestration model. The repressor (R) binds to the DNA-bound-
activator (E_A) with a dissociation constant of K_b to block the transcription (i.e., blocking; gray box). (B)
The combination of the sequestration and blocking (blue line) generates an ultrasensitive switch
similar to the sole sequestration (red line). (C) However, unlike the sole sequestration, the
combination of the sequestration and blocking rapidly restores the undesired activation of DNA during
TRP to the repressed state. (D) When unexpected binding of activator to DNA occurs during TRP,
repressor rapidly inhibits the activator in the presence of blocking (blue arrow), but not in its absence
(red arrow). (E) As a result, the stationary distribution of mRNA molecules is concentrated around
zero (blue bars) without bimodality observed in the sole sequestration (red bars). (F) Thus, the noise
level dramatically reduces during TRP in the presence of blocking. However, the addition of blocking
to the sole sequestration increases the noise level during the TAP. (G) Such high transcriptional noise
is due to undesired repression persists for a long-time even during TAP when blocking is added.
Here, the timeseries of non-activated DNA (i.e., free or repressed DNA) and mRNA are simulated at
$\widetilde{R_T} = 10^{-0.1}$ (gray star in (B)). (H) When noise-induced binding of R to DNA-bound-activator occurs
(i.e, blocking occurs), due to their tight binding, the undesired repression during TAP persists for a
long time. This leads to the sharp decrease mRNA near to zero. (I) As a result, the stationary
distribution of mRNA molecules shows an additional peak around zero, leading to bimodality with
blocking (blue bars) unlike sole sequestration (red bars).
485

488
489
Figure 4. Transcriptional ultrasensitive switch based on the combination of sequestration,
blocking, and displacement is robust to noise during both TAP and TRP. (A) The displacement is
added to the combination of the blocking and sequestration model in Fig. 3(A). The repressor-activator
complex (R_A) dissociates from DNA (E_R) with a dissociation constant of K_d to inhibit the transcription
(i.e., displacement; gray box). (B) The combination of the sequestration, blocking and displacement
(blue line) generates an ultrasensitive switch similar to the sequestration and blocking (red line). (C)
With the addition of displacement, the undesired repression of DNA during TAP is rapidly restored to
the activated state. (D) When unexpected repression occurs during TAP, repressor-activator complex
dissociates from DNA for free activator to promote transcription immediately in the presence of
displacement (blue arrow), but not in its absence (red arrow). (E) As a result, an additional peak around
zero, observed in the combination of the sequestration and blocking (red bars), is removed from the
stationary distribution of mRNA (blue bars). (F) Thus, the addition of displacement leads to the reduction
in the noise level during TAP.