Title: Identifying signals of memory from observations of animal movements in Plato's cave

Authors

Dongmin Kim
kimx3725@umn.edu
Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul MN

Peter R Thompson
pt1@ualberta.ca
Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada

David Wolfson
wolfs064@umn.edu
Department of Fisheries, Wildlife and Conservation Biology, University of Minnesota, St. Paul MN

Jerod Merkle
jmerkle@uwyo.edu
Department of Zoology and Physiology, University of Wyoming, Laramie, WY

L. G. R. Oliveira-Santos
gu_tapirus@hotmail.com
Departmento de Ecologia, Universidade Federal do Mato Grosso do Sul, Campo Grande, Brazil

James D. Forester
jdforest@umn.edu
Department of Fisheries, Wildlife and Conservation Biology, University of Minnesota, St. Paul MN

Tal Avgar
tavgar@ualberta.ca
Wildlife Science Center, Biodiversity Pathways Ltd., and Department of Biology, University of British
Columbia – Okanagan, Kelowna, BC, Canada.

Mark A. Lewis
marklewis@uvic.ca
Department of Mathematics and Statistics, and Department of Biology, University of Victoria, Victoria,
BC, Canada

John Fieberg
jfieberg@umn.edu
Department of Fisheries, Wildlife and Conservation Biology, University of Minnesota, St. Paul MN

Abstract

1. Incorporating memory (i.e., some notion of familiarity or experience with the landscape) into models of animal movement is a rising challenge in the field of movement ecology. The recent proliferation of new methods offers new opportunities to understand how memory influences movement. However, there are no clear guidelines for practitioners wishing to parameterize the effects of memory on moving animals.

2. We review approaches for incorporating memory into Step-Selection Analyses (SSAs), a frequently used movement modeling framework. Memory-informed SSAs can be constructed by including spatial-temporal covariates (or maps) that define some aspect of familiarity (e.g., whether, how often, or how long ago the animal visited different spatial locations) derived from long-term telemetry data.

3. We demonstrate how various familiarity covariates can be included in SSAs using a series of coded examples in which we fit models to wildlife tracking data from a wide range of taxa. We discuss how these different approaches can be used to address questions related to whether and how animals use information from past experiences to inform their future movements.

4. By reviewing different approaches and providing code templates for their implementation, we hope to inspire practitioners to further investigate the importance of memory in animal movements using wildlife tracking data.

Keywords: animal movement, cognitive map, familiarity, memory, occurrence distribution, space use, step-selection analyses, wildlife tracking data

1. Introduction

Animal movement impacts processes at all ecological levels, including individual foraging efficiencies (Bonte et al., 2012; Van Moorter et al., 2009), population persistence (del Mar Delgado et al., 2018; Miles et al., 2020), species distributions (Fortin et al., 2005; Matthysen, 2005), connectivity (Doherty & Driscoll, 2018) and ecosystem functioning (Bauer & Hoye, 2014; Nathan et al., 2008; Subalusky et al., 2017). While much research has explored how animal movements are influenced by environmental conditions (Fortin et al., 2005), intra- and inter-specific social interactions (Matthysen, 2005), and internal states such as hunger levels (Hooten et al., 2019), the importance of past experience and memory is also increasingly recognized as a key component of animal movement (Piper, 2011; Fagan et al., 2013; Lewis et al., 2021). For example, by remembering the location and outcomes of previously visited locations, many species can increase energy intake rates (Janson & Byrne, 2007; Trapanese et al., 2019) and avoid areas that might increase mortality risk (Bracis et al., 2018). Further, by remembering average environmental conditions, such as the average timing of resource waves, animals can better time migratory movements (Abrahms et al., 2019; Bracis & Mueller, 2017).

The information animals gather, and perhaps memorize, while moving can be divided into three categories: 1) spatial information (i.e., locations animals have visited), 2) site attributes, including resource quality or quantity, and 3) temporal information (i.e., about how long-ago animals visited a previous site or when a site peaks in forage quality) (Fagan et al., 2013; Kashetsky et al., 2021; Lewis et al., 2021; Thompson, Lewis, et al., 2022; Tulving & Craik, 2000). Food-caching blue jays (*Cyanocitta cristata*) use all three kinds of information: they remember the locations of many caches, the type of seed in each cache, and how long it has been since making the cache (Darley-Hill & Johnson, 1981). Owing to the temporal variability present in most environments, it can be advantageous to rely more heavily on recent experiences and to discount memories from long ago (Tello-Ramos et al., 2019). For example, roe deer (*Capreolus capreolus*) primarily rely on recent experiences when making foraging decisions due to rapid changes in resource availability within their home ranges (Ranc et al., 2021). By revisiting sites, animals can update their knowledge of site attributes; optimal return times may depend on how quickly the reliability of past information decays due to environmental change as well as resource renewal rates (Ranc et al., 2021; Spencer, 2012). For example, wolves (*Canis lupus*) and brown bears (*Ursus arctos*) delay returning to previously visited kill sites so that prey numbers may recover (Gurarie et al., 2022; Selva et al., 2017).

Ecologists have developed theoretical models to explore how past experiences and memory might influence animal movements (Fagan et al., 2013; Kashetsky et al., 2021; Lewis et al., 2021; Wang & Salmaniw, 2023). The simplest encodable memory attribute is familiarity with a given location, either whether an individual has ever visited the site (Northrup et al., 2016; Rheault et al., 2021) or how frequently it has visited the site in the past (Bracis et al., 2018). When coupled to spatial movement models, preference for familiar locations is sufficient for the formation of stable home ranges (Gautestad et al., 2013; Merkle et al., 2017; Van Moorter et al., 2009). More complex memory attributes include locations of resources or past conflicts, allowing animals to integrate spatial and attribute memory (i.e., memory of where positive and negative experiences occurred). Attraction to previously discovered resources can lead to resource-driven patterns of nonterritorial spatial segregation (Aarts et al., 2021; Riotte-Lambert et al., 2015). By way of contrast, memory of and avoidance of locations where past conflicts with conspecifics occurred can give rise to spontaneous territorial pattern formation (Potts & Lewis, 2016).

One straightforward approach to integrating such complex types of memories into a model is to assume that past experiences can be encoded into a spatially-referenced system in the animal's brain (sometimes referred to as a "cognitive map"), which is then accessed during the retrieval phase to inform movements. Although hidden from direct observation, much like the objects casting shadows in Plato's cave, a

spatially-referenced map can be mathematically modeled as a surface changing dynamically over time as
memories are lost, reinforced, or replaced. These constructs are central to key empirical models for
memory, including "time since last visit" to a location (Schlägel & Lewis, 2014) as a determinant of wolf
movement (Schlägel et al., 2017), and episodic returns of brown bears to ephemeral seasonal resources
(Thompson, Derocher, et al., 2022; Thompson, Lewis, et al., 2022). Here, the map is a latent, spatially-
referenced variable, whose dynamics are inferred indirectly from animal movement patterns.
Advances in animal tracking technology and statistical modeling approaches have motivated ecologists to
explore the potential for memory-informed movements in a wide range of animal taxa (although
terrestrial mammals, mainly ungulates, remain by far the most-studied group; see Supporting Information
1). By tracking individual animals over relatively long time frames, ecologists can identify whether
current movements can be explained from previously observed locations. Popular analytical frameworks,
such as integrated step-selection analyses (Avgar et al., 2016; Fieberg et al., 2021) can be used to identify
signals of memory from such observations (Merkle et al., 2019; Oliveira-Santos et al., 2016). However,
several challenges remain before these approaches can be widely adopted. These include technological
challenges associated with managing tracking data, creating models with different "familiarity" or
"memory" covariates, and drawing biologically accurate conclusions from these models. Here, we
provide an overview of current modeling approaches and their features to help guide practitioners wishing
to identify the effects of memory in animal movement. We provide several examples with annotated code
and then discuss the strengths and limitations of current approaches and future directions for memory-
informed movement research.

2. Exploring how memory influences animal movements using step-selection analyses.

Step-Selection Analyses (SSAs) are widely used to quantify how the environment influences animal
movement (Avgar et al., 2016; Fieberg et al., 2021; Fortin et al., 2005; Thurfjell et al., 2014). SSAs model
movement in discrete time using two model components: 1) a selection-free movement kernel describing
how animals move in the absence of habitat selection, defined using distributions of step-length and
turning angle, and 2) a selection function describing animal preferences with regards to the habitat
attributes at each step's endpoint (Box 1). Model parameters in step-selection analyses can be estimated
using commonly available statistical software that implements conditional logistic regression (Avgar et
al., 2016; Fieberg et al., 2021). Because they allow one to model and predict dynamic space-use patterns
using accessible and available software, step-selection analyses are attractive to movement ecologists and
are widely used to analyze animal tracking data (Signer et al., 2019).

Box 1. Step-Selection Analyses (SSAs)

Step-Selection Analyses (SSAs) model the conditional probability, $p(s_t \mid H_{t-1}; \beta_m, \beta_w)$, of finding an individual at a location s_t at the time t given a set of previously visited locations, H_{t-1}, using a selection-free movement kernel, $k(s_t \mid H_{t-1}; \beta_m)$, which describes how animals would move in the absence of habitat selection, and a movement-free habitat-selection function, $w(s_t; t, \beta_w)$, which describes the animals' preferences for certain environmental features (e.g., variables representing resources, risks, and or other conditions; Matthiopoulos et al. 2023):

$$p(s_t|H_{t-1}; \beta_m, \beta_w) = \frac{\overbrace{k(s_t|H_{t-1};\beta_m)}^{Selection-free\ Movement\ kernel} \cdot \overbrace{w(s_t;t,\beta_w)}^{Habitat\ selection\ function}}{\underbrace{\int_{s' \in U} k(s'|H_{t-1};\beta_m) \cdot w(s';\beta_w)ds'}_{Normalisation\ over\ the\ spatial\ domain\ U}} \qquad (1)$$

$$\underbrace{H_{t-1,}}_{History\ of\ the\ locations} = \{s_{t-1}, s_{t-2}, \quad \underbrace{\dots\ ,\ s_{t-i}, \dots}_{Further\ past\ locations} s_0\} \qquad (2)$$

β_m contains parameters in the step-length and turn angle distributions $(\beta_{m1}, \dots, \beta_{mq})$, and β_w contains resource-selection parameters that quantify the attractiveness of different locations using a vector of selection coefficients $(\beta_{w1}, \dots, \beta_{wp})$ for each environmental covariate $(r_1(s_t), \dots, r_p(s_t))$. $s' \in U$ describes all the locations within the spatial domain U. To calculate a step length, *sl*, two locations are required, (s_t, s_{t-1}). Similarly, a turning angle, *ta*, is calculated using the current and the past two locations (s_t, s_{t-1}, s_{t-2}).

141
A variety of spatiotemporal familiarity covariates can be included in an SSA to model the effects of
memory (Box 2; Box 3). For example, one may estimate an occurrence distribution (OD) describing the
relative use of the landscape over a specific period in the past (e.g., a continuous surface of either the
relative intensity of use or a binary presence/absence variable). By including the OD as a spatial predictor
in an SSA, one can evaluate whether an animal's current landscape usage is biased toward or away from
previously visited locations (Rheault et al., 2021). A notable limitation of this approach is the risk of
overestimating the importance of familiarity due to unaccounted-for habitat attributes (e.g. if the OD
reflects the distribution of an unobserved resource; see Picardi et al. 2023). The OD approach further
necessitates that users choose an appropriate time window in the past for calculating the OD (Figure 1).
Using a limited time window for calculating the familiarity covariate implies an abrupt memory decay
function where past locations are memorized for a fixed amount of time and then forgotten completely.
Alternatively, one could choose to continually update the OD from the first to the last location, which
would imply the animal never forgets its past experiences. With either approach, the OD effectively
weights all previously visited areas within the specified time window equally, regardless of how long ago
the animal visited the location. Another option is to allow more recent (or distant) memories to have more
influence on current movements by replacing the OD with a covariate representing the length of time
since the animal last visited a location (TSLV).

For migratory species that navigate relatively long distances, familiarity predictors could include: 1)
distances between the current and previous year's migratory trajectories, or 2) angular covariates
constructed by comparing the direction of an animal's movement in relation to the centroid of
observations from a previous year's seasonal range (Figure 2). These methods can describe an animal's
use of memory for navigation and capture its tendency to use familiar migration routes and consistent but
seasonally varying home ranges (Merkle et al., 2019).

Whatever familiarity predictor we choose, and unless we are comfortable assuming the animal was
completely naïve (lacked any familiarity) at the onset of the tracking, accounting for familiarity requires
'sacrificing' some early positional data that is used to estimate the familiarity predictor (rather than its
effect on animal movement). For many long-lived animals, and particularly in highly seasonal
environments, the 'memory build up' period may need to be long (a year or even more), unless there are
reasons to believe memory (or its effect) decays over much shorter time scales. For example, Avgar et al.
(2015) used a full year of 'memory build up' data to model woodland caribou (*Rangifer tarandus*
caribou) space-use patterns over a subsequent year but have found no indication of memory decay within
that time (indicating the need to use an even longer 'memory build up' period).

Box 2. Guide for the memory-informed movement modeling within SSF framework

Question	Biological example	Method
Do animals retain information from their past experience and for how long?	Roe deer primarily rely on recent experiences when making foraging decisions due to rapid changes in resource availability within their home ranges (Ranc et al., 2021).	Occurrence Distribution (OD)
What is the animal's optimal return time based on memorized information regarding past foraging or predation history or resource renewal rates?	Predators like wolves and brown bears delay returning to previously visited kill sites so that prey numbers may rebound to normal levels (Gurarie et al., 2022; Selva et al., 2017).	Time Since Last Visit (TSLV)
Do migratory animals use consistent migration routes and return to the same seasonal ranges each year?	Mule deer (*Odocoileus hemionus*) have been shown to use consistent migration routes and seasonal ranges, likely relying on their past experience to help them navigate (Merkle et al., 2019).	Distance and directional bias towards previously used areas

---

Box 3. Familiarity function in SSFs

Movement ecologists have described familiarity with different parts of the landscape that animals experienced using familiarity covariates, which we formalize via a familiarity function, *f*. Including an exponential familiarity function allows the attractiveness of different locations to be governed by both environmental and familiarity covariates within the traditional SSF framework ($e^{\beta_w r(s)} \cdot e^{\beta_f f(s)} = e^{\beta_w r(s) + \beta_f f(s)}$). Examples of familiarity covariates include occurrence distributions (ODs) reflecting the intensity of past space use in the study area, time since last visit (TSLV), migratory distances between current and previously used paths, and angular covariates used to capture bias toward previously used migratory ranges:

$$p(s_t|H_{t-1}; \beta_m, \beta_w, \beta_f) = \frac{\overbrace{k(s_t|H_{t-1}; \beta_m)}^{Selection-free\ Movement\ kernel} \cdot \overbrace{w(s_t; t, \beta_w)}^{Habitat\ selection\ function} \cdot \overbrace{f(s_t; t, \beta_f)}^{familiarity\ function}}{\underbrace{\sum_{s\prime \in U} k(s\prime|H_{t-1}, \beta_m) \cdot w(s\prime; t, \beta_w) \cdot f(s\prime; t, \beta_f)}_{Normalisation\ over\ the\ spatial\ domain\ U}} \quad (3)$$

$$f(s_t; t, \beta_f) = \{e^{\beta_f \cdot f(s_t)}\} \quad (4)$$

Figure 1. Spatial memory can be quantified using an animal's occurrence distribution (OD) measured over some prior period (colored areas: 5-day periods at time 6 and time 8 = orange vs green) which captures an animal's movement path and its uncertainty. A time-varying covariate can be constructed by updating the OD at regular time intervals. This updating step ensures that distant experiences are eventually forgotten and no longer play a role in driving animal movement. Users must choose an appropriate time window in the past for calculating the memory covariate and how often to update it.

Figure 2. Familiarity covariates used in applications involving migratory animals. The red bounding box in panel (a) displays the area zoomed into for boxes b and c. Memory can be quantified using a distance predictor calculated as the minimum distance between current and past migratory paths (panel b). Specifically, we can calculate the distances (dashed lines) between Year 2 (observed [navy], random [pink]) locations and Year 1 (green) migratory paths. Memory can also be included as a directional bias predictor by comparing whether the current year's steps are biased toward the previous season's range (panel c). This bias predictor can be calculated using the angles, θ_1, θ_2, between the step (observed [navy], random [pink]) and the centroid of the previous year's winter range.

192
3. Case studies

In this section, we review memory-informed movement models for animal tracking data using 4 case
studies. The first two examples, involving data from sandhill cranes (*Antigone canadensis*) and feral hogs
(*Sus scrofa*), demonstrate how one can use a spatial familiarity predictor calculated from a past OD to
explore whether animals retain information from their past experiences and for how long. The third
example considers the migratory movements of mule deer and illustrates the use of two familiarity
predictors formed using the minimum distance between the current and last year's migratory paths and
the cosine of the angle between the direction of a current movement step and the previous year's centroid
of locations. The last example, involving data from a brown bear, uses a spatiotemporal covariate to
quantify how long it has been since the individual last visited spatial locations, and thus, how memory
may influence revisitation rates. We provide a workflow to reproduce the main components of the habitat-
and memory-based SSF from all the case studies, replicating the importance of memory-based metrics
versus habitat metrics in an SSF of animal movement.

1. Sandhill Crane – 'fixed-time' OD

Sandhill cranes breed throughout North America during summer and migrate south for the winter. During
their first year, juveniles migrate to overwintering areas with their parents and then disperse from the
family group either during the spring migration or upon arrival to the natal territory the following spring
(Hayes, 2015; Tacha et al., 1989). During the first few years of independence, subadult cranes typically
make long-distance movements across the landscape during the summer; in contrast, the movements of
breeding adults are largely constrained to their breeding territories (Wolfson et al., 2020). Once cranes
become successful breeding adults, typically between 4-6 years old, they use their accumulated
knowledge of the landscape to return and nest in the same breeding area in subsequent years (Nesbitt,
1992; Tacha et al., 1989).

As an illustrative example of how a sandhill crane's space use constricts each year as the crane learns its
landscape and develops a breeding territory, we consider a 5-year dataset of global positioning system
(GPS) telemetry locations of a sandhill crane (with a 15-minute fix interval), starting from the time of
fledging (Wolfson, 2018; Wolfson et al., 2020). A visualization of summer locations shows that the
spatial coverage visited by the crane decreases each year as it ages, selecting locations it had previously
visited (Figure 3a). This pattern suggests that the crane may be using memory of its visited landscape to
decide where to establish a breeding territory.

To quantify the potential effects of familiarity on the crane's summer movements, we calculated areas
associated with the 95% contour of an OD estimated from the previous year's summer locations using the
probability density functions from a continuous-time movement model conditioned off the Brownian
bridge (Fleming et al., 2015). We then fit a separate SSF to each year of data, including the previous
year's OD as a predictor in the model. A positive coefficient associated with the previous year's OD
would suggest the crane prefers to revisit sites it visited in the previous year (compared to *equally*
accessible and otherwise identical sites that it did not visit during the previous year, where accessibility is
determined by the selection-free movement kernel; Fieberg et al. 2021). If the coefficient is negative, the
opposite interpretation holds.

Coefficients associated with the previous year's OD were positive in all years (Figure 3b), even though
the crane rarely revisited previously used locations in the first two years (Figure 3c). Interpretation of the
coefficients associated with categorical predictors can be difficult, as they reflect a ratio of ratios (use:
availability ratio for one class versus the use: availability ratio of a reference class; Fieberg et al. 2021).
The positive coefficients for the previous year's OD in 2016 and 2017 reflect the fact that the use-
availability ratio associated with grid cells visited in the previous year was higher than the use:
availability ratio for grid cells that were not visited in the previous year. Thus, we end up with a positive

coefficient for the OD predictor even though most of the areas encountered and used by the crane
represent areas that were not visited in the prior year. Recall our interpretation of a positive coefficient, *if*
presented with two equally accessible locations (same step length and turn angle required to reach both
locations) that only differ in whether they had been visited in the previous year (i.e., they had the same
landcover class), the crane would be more likely to select the location it was familiar with. A limitation of
this model and application is that the crane was rarely presented with this type of choice. In the first two
years, it was rarely found near sites it had previously visited whereas in years 3-5 it rarely visited areas it
was unfamiliar with. A second limitation is that the birds may be selecting for an unobserved resource,
one that we did not include in our analysis, which introduces a hidden correlation between their past and
current space use.

Figure 3. Visualization of Sandhill crane tracking data and coefficients from fitted step-selection functions with a memory covariate formed using an OD capturing previously visited locations **a)** Movement patterns of an individual sandhill crane during summer seasons (06/15-09/22) from 2015 to 2020. **b)** Parameter estimates including those for the memory covariate (Odprev: OD from the previous year). **c)** Distribution of each year's used and available locations for sites (i.e., grid cells) that were (OD = 1) and were not (OD = 0) visited in the previous year. Numbers at the top of the bars indicate the number of locations in each group.

2. Feral Hog – OD with temporal variation

Hogs were introduced to the Pantanal wetland about 300 years ago, and currently represent the highest wild mammal biomass in this region. They are crepuscular-nocturnal, social, long-lived, cooperative animals that forage at the edges of water bodies and in ephemeral pools that become increasingly rare during the winter dry season. Hogs lack sweat glands and behaviorally mitigate heat stress by spending the hot hours of the day resting in forest patches.

As animals move, they can access and update their reference memory (long-term acquisition) and working memory (short-term acquisition) to navigate through space. Still, the stored spatial information may or may not be used depending on the current animal needs and context, which generates temporal heterogeneity (e.g., within the day, and seasons) in the use of that information. Oliveira-Santos et al. (2016) modeled the hidden process underlying spatial memory acquisition using spatiotemporal covariates generated from Biased Random Bridge kernel density estimates based on residence time (Benhamou, 2011). Specifically, for each individual step, they used previous locations to build multiple spatial maps, formed using different time windows, that could affect the hogs' future movement decisions. These maps were then continuously updated as the individual moved through the landscape.

Oliveira-Santos et al. (2016) considered 4 conceptually different hypotheses regarding how hogs process and use spatial information, combining long- and short-term memory with differential use of stored memory within the daily cycle. To represent long-term memory, familiarity covariates were constructed and constantly updated with all previous locations as the individual moved, whereas short-term memory was represented using a spatiotemporal covariate that kept track of just the last 3 days. Additionally, long- and short-term familiarity covariates were also built considering only daytime or only nighttime locations, which they referred to as long-term temporal and recent-temporal memory, respectively. When fitting models for these last two cases, movement steps taken at night or during the day were paired with familiarity covariates generated from previous locations collected only at night or day, respectively.

All tracked hogs strongly selected for previously visited areas and mainly those associated with short-term memory. Most of these individuals (65% of the tracked hogs) appeared to use working memory as part of their movement process, (e.g., covariates generated from recent nocturnal locations were better at predicting future nocturnal use than covariates generated from all time periods). Importantly, the effect of familiarity also varied within the day, being more important during the daylight hours when individuals were sleeping in well-known places than at night when animals were foraging and were more willing to take risks by walking through less familiar areas. Although hogs are acknowledged to have high cognitive skills and memory retention, they relied mainly on recent spatial information because the distribution of prime food resources in the study area responds quickly to foraging pressure and changes in water levels.

3. Mule Deer – Migratory paths and angles

Many large ungulates are migratory, capitalizing on seasonal and spatial variation in food, predation, and inhospitable conditions (Avgar et al., 2013). Mule deer are a concentrate forager (i.e., prefer to consume high-quality food) that display some of the longest terrestrial migrations in North America (Joly et al., 2019), spending winters in arid, low-elevation sagebrush, grassland, and desert ecosystems, and then migrating (up to 400 km) into montane ecosystems at higher elevations for summer.

Unlike some other migratory species (e.g., Sierra Nevada Bighorn Sheep; Berger et al., 2022), Mule deer migrate well outside their perceptual range (e.g., what they see, hear, and smell at a given moment), yet they display relatively strong fidelity to seasonal ranges (Morrison et al., 2021) and migration routes (Sawyer et al., 2019). For instance, on average, mule deer migrate on the same path during spring year after year 81% of the time (Sawyer et al., 2019). In some cases, such migrations occur across relatively vast expanses of featureless deserts, rolling hills, and thick forests, where sensory abilities such as vision may not improve navigation.

Mule deer may navigate during migration by memorizing the path of their previous migration route and
the general location of the seasonal range they are attempting to navigate to. The relative influence of
these memorized spatial locations can be assessed in a movement model that first considers several other
habitat features that may influence ungulate movement and space use. Merkle et al. (2019) examined the
relative role of memory usage versus local variation in habitat on mule deer navigation during migration.
They found that variables indexing past experience (distance to the previous migration route and direction
to the previous seasonal range) were 2 to 28 times more predictive of migratory movements than local
variation in habitat.

4. Brown Bear – time since last visit (TSLV)
Brown bears are opportunistic omnivores found in North America, Europe, and Asia (Pasitschniak-Arts,
1993). Their life history strategies and dietary compositions vary greatly depending on the environment
they live in (Finnegan et al., 2021; García-Rodríguez et al., 2021; MacHutchon & Wellwood, 2003), but
their ability to navigate towards previously visited food patches is ubiquitous throughout their natural
range (Selva et al., 2017; Thompson, Lewis, et al., 2022; Wirsing et al., 2018). The "barren-ground
grizzly bears" found in the Canadian Arctic are unique for many reasons, including their foraging and
denning behavior (Edwards & Derocher, 2015; McLoughlin, 2002). These bears' food resources are only
available for a short, albeit predictable, portion of their active seasons (Edwards & Derocher, 2015), so in
addition to returning to the correct spatial location where food was present, bears must identify the
temporal pattern of this resource and revisit the patch at the correct time. Ecologists interested in
understanding how these bears incorporate memory into their movement patterns must implement models
that account for these nonlinear temporal dynamics.
327
We modified the model developed by Schlägel & Lewis (2014) so that it could be implemented in the
step-selection framework and fit using conditional logistic regression. This required using distributions
from the exponential family (gamma, von Mises) to model the distribution of step lengths and turn angles,
respectively (Avgar et al. 2016; Fieberg et al. 2021). In addition, we included both linear and quadratic
terms to model the influence of time since last visit (TSLV) on the movement patterns of an adult female
brown bear in the Mackenzie River Delta region of the Northwest Territories, Canada. This bear was
immobilized and fitted with a GPS collar that recorded its location every 4 hours during the active season
(the time in which the bear was out of its den).
336
We calculated TSLV as the difference between the current timestamp and the last time the bear visited a
series of 2 x 2 km grid cells, updating the map at each observation time. This familiarity covariate
allowed "revisitation" to occur when the animal was in the perceptual vicinity of an area it previously
visited (i.e., within the same grid cell), without necessitating that the animal returns exactly to its previous
coordinates (Schlägel & Lewis, 2014). We discarded the first year of location data as a "burn-in", and set
TSLV to 365 for any grid cells where TSLV was missing, effectively assuming that these cells had been
visited just prior to the first observed location. The need for a burn-in period is a limitation of this
approach but is necessary since we have no history of the bear's past visits prior to the start of data
collection. A sensitivity analysis can be performed to evaluate whether results change if the length of the
burn-in period is increased or decreased. To analyze how selection strength varied with TSLV, we
calculated the relative selection strength (RSS) at different TSLV values (Avgar, 2017). As with quadratic
terms in other habitat selection models, by including both linear and quadratic terms, we were able to
identify a non-linear response to TSLV with intermediate TSLV values of approximately 350 days
displaying the strongest selection (Figure 4). These results agree with those of Thompson et al. (2021),
who also found that bears tended to revisit sites seasonally.

Figure 4. Visualization of the relative attractiveness of previously visited locations as a function of time since last visit (TSLV) modeled using linear and quadratic terms in a step-selection analysis of brown bear location data with the distribution of used (green, bottom) and available (brown, top) locations shown along the x-axis.

4. Discussion

Modern GPS tracking systems generate massive telemetry datasets by following individual animals over a long time on a global scale. With this abundance of available data, it is now possible to develop models that evaluate how memory relates to the animal movement (Nathan et al., 2022), which has inspired the recent development and application of many such models (see Supporting Information 1). Our review focused on approaches that account for familiarity with different areas of the landscape by including spatial, spatiotemporal, or angular covariates as predictors in step-selection analyses. These frameworks can be implemented using available statistical software for fitting conditional logistic regression models, which we demonstrate using multiple tracking data sets and annotated code examples. Although some data development is necessary before fitting the models (e.g., to create the memory predictors and generate random steps), users can leverage R packages to make these steps easier. For example, the ctmm package (Calabrese et al., 2016) can be used to calculate ODs, and the amt package (Signer et al., 2019) can be used to generate random steps necessary for parameterizing the model.

Individual-based models of animal movement are increasingly used to inform conservation and management at the population or even species levels. Prime among these applications is the use of habitat-selection models to identify critical habitats, delineate range boundaries, and project spatial distribution across space or time. A strong selection for certain habitats or environmental features and conditions may occur with or without memory, and it is important to consider how models that incorporate familiarity covariates may alter inferences. On the one hand, we might expect mild to moderate collinearity between familiarity covariates and other important environmental drivers, which can make it challenging to quantify their unique contributions (Picardi et al., 2023). On the other hand, accounting for familiarity and memory effects should reduce bias associated with estimators of habitat-

selection strength. Consider an animal that is both attracted to certain habitats and to places it has used in
the past; if we ignore the latter, we will consequently overestimate the former, putting more weight on
habitat attraction than it truly has. Similarly, it is important to consider the effect of environmental drivers
when looking for influences of memory on animal movements (Van Moorter et al., 2013), and in fact,
memory-free movement models that allow for habitat selection have been suggested as null models for
evaluating evidence of memory (Picardi et al., 2023). An additional benefit to considering memory effects
in SSAs is the option to simulate these effects under various management scenarios. For example,
managers might be interested in assessing the likelihood of successfully relocating an animal and having
it establish a home range in a new site (Ranc et al., 2021; Berger et al., 2022). Explicitly modeling the
process of building and responding to increasing knowledge of the landscape may be critical for obtaining
reliable predictions.
391
Information gathering, an essential prerequisite for memory, is governed by sensory ecology (Dusenbery
1992, Dukas 1998). The information available to the animal about any given spatial location (the 'signal')
is a function of the animal's position in relation to that location (the source of the signal), the strength of
the signal (e.g., the intensity of odor, light, or sound), the overlap with similar signals coming from
elsewhere on the landscape, and the animal's sensory capacity to perceive and process the signal (which
in itself may be a complex function of the animal's morphology and physiology). Information can only be
committed into memory if it is being perceived. Moreover, it is very likely that the weight given to
memorized information (or its retention time) is modulated by the signal-to-noise ratio at which the
information was perceived. The models we described here assume, for the most part, that animals sense
and retain information from only 'visited localities', which are arbitrarily sized spatial units (typically
corresponding to a single pixel of the available environmental data). Furthermore, most SSFs only assess
what the animal can perceive within local ranges (i.e., what is nearby), though angular and distance-to
covariates allow modeling perception over larger spatial scales. Thus, SSFs are a useful, but extreme
simplification of the true underlying sensory ecology.
406
More mechanistically inclined frameworks for explicitly modeling the sensory processes involved in
memory buildup have been proposed (e.g., Avgar et al. 2013, 2015; Ranc et al. 2022; Thompson, Lewis,
et al. 2022); however, fitting these models to wildlife tracking data is still a challenge. These frameworks
typically include additional free parameters used to construct spatiotemporal covariates that determine
how perceived habitat quality varies over time (e.g., by modeling how perception decays with distance
from the individual, and memory decays temporally). As a result, it is not possible to fit these models
using standard statistical software developed for SSAs. Instead, the parameters governing the
spatiotemporal covariate must be estimated simultaneously with other habitat-selection and movement
parameters through a custom-written likelihood function that can be optimized using Markov chain
Monte Carlo or other numerical optimization methods. We include an example from Thompson et al.
(2021) in our supplementary material to demonstrate this approach. Methods that more realistically model
the process of memory formation should be pursued, but we suspect that most practitioners will continue
to explore the role of memory on animal movements using simpler models that can fit within the standard
SSA framework using conditional logistic regression.
421

Authors' Contributions

423 DK and JF conceived of the topic, and DK led the writing of the manuscript. Case studies were led by
424 DK, DW, and JF (sandhill crane), DK, PT, and JF (brown bear), JM (mule deer), and L O-S and JDF
425 (hog). All authors discussed, edited, read, and approved the final manuscript.
426

Funding Information

DK was supported by the National Science Foundation INTERN award 1654609, the Smithsonian Institution Fellowship Program (SIFP), and the University of Minnesota Doctoral Dissertation Fellowship (DDF). JF was supported by the National Aeronautics and Space Administration award 80NSSC21K1182. JF and JDF received partial salary support from the Minnesota Agricultural Experimental Station.

Conflict of Interest

The authors declare no conflict of interest.

Data Availability Statement

The datasets and code associated with the case studies would be archived in the Data Repository for the University of Minnesota (DRUM) if the manuscript is accepted.

ORCID

Dongmin Kim https://orcid.org/0000-0002-1508-1590
Peter R Thompson https://orcid.org/0000-0002-7590-8473
David Wolfson https://orcid.org/0000-0003-1098-9206
Jerod Merkle https://orcid.org/0000-0003-0100-1833
L. G. R. Oliveira-Santos https://orcid.org/0000-0001-9632-5173
James D. Forester https://orcid.org/0000-0002-5392-9556
Tal Avgar https://orcid.org/0000-0002-8764-6976
Mark A. Lewis https://orcid.org/0000-0002-7155-7426
John Fieberg https://orcid.org/0000-0002-3180-7021

Supporting Information

Additional supporting information can be found online in the Supporting Information section at the end of this article.

References

Aarts, G., Mul, E., Fieberg, J., Brasseur, S., van Gils, J. A., Matthiopoulos, J., & Riotte-Lambert, L. (2021). Individual-Level Memory Is Sufficient to Create Spatial Segregation among Neighboring Colonies of Central Place Foragers. *The American Naturalist*, *198*(2), E37–E52. https://doi.org/10.1086/715014

Abrahms, B., Hazen, E. L., Aikens, E. O., Savoca, M. S., Goldbogen, J. A., Bograd, S. J., Jacox, M. G., Irvine, L. M., Palacios, D. M., & Mate, B. R. (2019). Memory and resource tracking drive blue whale migrations. *Proceedings of the National Academy of Sciences*, *116*(12), 5582–5587. https://doi.org/10.1073/pnas.1819031116

Avgar, T. (2017). Relative selection strength: Quantifying effect size in habitat-and step-selection inference. *Ecology and Evolution*, *7*(14), 5322–5330.

Avgar, T., Baker, J. A., Brown, G. S., Hagens, J. S., Kittle, A. M., Mallon, E. E., McGreer, M. T., Mosser, A., Newmaster, S. G., Patterson, B. R., Reid, D. E. B., Rodgers, A. R., Shuter, J., Street, G. M., Thompson, I., Turetsky, M. J., Wiebe, P. A., & Fryxell, J. M. (2015). Space-use behavior of woodland caribou based on a cognitive movement model. *Journal of Animal Ecology*, *84*(4), 1059–1070. https://doi.org/10.1111/1365-2656.12357

Avgar, T., Deardon, R., & Fryxell, J. M. (2013). An empirically parameterized individual based model of animal movement, perception, and memory. *Ecological Modelling*, *251*, 158–172. https://doi.org/10.1016/j.ecolmodel.2012.12.002

Avgar, T., Potts, J. R., Lewis, M. A., & Boyce, M. S. (2016). Integrated step selection analysis: Bridging the gap between resource selection and animal movement. *Methods in Ecology and Evolution*, *7*(5), 619–630. https://doi.org/10.1111/2041-210X.12528

Bauer, S., & Hoye, B. J. (2014). Migratory Animals Couple Biodiversity and Ecosystem Functioning
Worldwide. *Science*, *344*(6179), 1242552. https://doi.org/10.1126/science.1242552
Benhamou, S. (2011). Dynamic Approach to Space and Habitat Use Based on Biased Random Bridges.
PLoS ONE, *6*(1), e14592. https://doi.org/10.1371/journal.pone.0014592
Berger, D. J., German, D. W., John, C., Hart, R., Stephenson, T. R., & Avgar, T. (2022). Seeing is be-
leaving: perception informs migratory decisions in Sierra Nevada Bighorn sheep (Ovis
canadensis sierrae). *Frontiers in Ecology and Evolution*, *10*.
Bonte, D., Van Dyck, H., Bullock, J. M., Coulon, A., Delgado, M., Gibbs, M., Lehouck, V., Matthysen,
E., Mustin, K., Saastamoinen, M., Schtickzelle, N., Stevens, V. M., Vandewoestijne, S., Baguette,
M., Barton, K., Benton, T. G., Chaput-Bardy, A., Clobert, J., Dytham, C., … Travis, J. M. J.
(2012). Costs of dispersal. *Biological Reviews*, *87*(2), 290–312. https://doi.org/10.1111/j.1469-
185X.2011.00201.x
Bracis, C., Bildstein, K. L., & Mueller, T. (2018). Revisitation analysis uncovers spatio-temporal patterns
in animal movement data. *Ecography*, *41*(11), 1801–1811. https://doi.org/10.1111/ecog.03618
Bracis, C., & Mueller, T. (2017). Memory, not just perception, plays an important role in terrestrial
mammalian migration. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1855),
20170449. https://doi.org/10.1098/rspb.2017.0449
Calabrese, J. M., Fleming, C. H., & Gurarie, E. (2016). ctmm: An R package for analyzing animal
relocation data as a continuous-time stochastic process. *Methods in Ecology and Evolution*, *7*(9),
1124–1132. https://doi.org/10.1111/2041-210X.12559
Darley-Hill, S., & Johnson, W. C. (1981). Acorn dispersal by the blue jay (Cyanocitta cristata).
Oecologia, *50*(2), 231–232. https://doi.org/10.1007/BF00348043
del Mar Delgado, M., Miranda, M., Alvarez, S. J., Gurarie, E., Fagan, W. F., Penteriani, V., di Virgilio,
A., & Morales, J. M. (2018). The importance of individual variation in the dynamics of animal
collective movements. *Philosophical Transactions of the Royal Society B: Biological Sciences*,
373(1746), 20170008. https://doi.org/10.1098/rstb.2017.0008
Doherty, T. S., & Driscoll, D. A. (2018). Coupling movement and landscape ecology for animal
conservation in production landscapes. *Proceedings of the Royal Society B: Biological Sciences*,
285(1870), 20172272. https://doi.org/10.1098/rspb.2017.2272
Dukas, R. (Ed.). (1998). *Cognitive ecology: the evolutionary ecology of information processing and
decision making*. University of Chicago Press.
Dusenbery, D. B. (1992). Sensory ecology: how organisms acquire and respond to information. *(No
Title)*.
Edwards, M. A., & Derocher, A. E. (2015). Mating-related behaviour of grizzly bears inhabiting marginal
habitat at the periphery of their North American range. *Behavioural Processes*, *111*, 75–83.
Fagan, W. F., Lewis, M. A., Auger-Méthé, M., Avgar, T., Benhamou, S., Breed, G., LaDage, L.,
Schlägel, U. E., Tang, W., Papastamatiou, Y. P., Forester, J., & Mueller, T. (2013). Spatial
memory and animal movement. *Ecology Letters*, *16*(10), 1316–1329.
https://doi.org/10.1111/ele.12165
Fieberg, J., Signer, J., Smith, B., & Avgar, T. (2021). A 'How to' guide for interpreting parameters in
habitat-selection analyses. *Journal of Animal Ecology*, *90*(5), 1027–1043.
https://doi.org/10.1111/1365-2656.13441
Finnegan, S. P., Svoboda, N. J., Fowler, N. L., Schooler, S. L., & Belant, J. L. (2021). Variable
intraspecific space use supports optimality in an apex predator. *Scientific Reports*, *11*(1), 21115.
https://doi.org/10.1038/s41598-021-00667-y
Fleming, C. H., Fagan, W. F., Mueller, T., Olson, K. A., Leimgruber, P., & Calabrese, J. M. (2015).
Rigorous home range estimation with movement data: A new autocorrelated kernel density
estimator. *Ecology*, *96*(5), 1182–1188. https://doi.org/10.1890/14-2010.1
Fortin, D., Beyer, H. L., Boyce, M. S., Smith, D. W., Duchesne, T., & Mao, J. S. (2005). WOLVES
INFLUENCE ELK MOVEMENTS: BEHAVIOR SHAPES A TROPHIC CASCADE IN

YELLOWSTONE NATIONAL PARK. *Ecology*, *86*(5), 1320–1330. https://doi.org/10.1890/04-
0953
García-Rodríguez, A., Selva, N., Zwijacz-Kozica, T., Albrecht, J., Lionnet, C., Rioux, D., Taberlet, P., &
De Barba, M. (2021). The bear-berry connection: Ecological and management implications of
brown bears' food habits in a highly touristic protected area. *Biological Conservation*, *264*,
109376. https://doi.org/10.1016/j.biocon.2021.109376
Gautestad, A. O., Loe, L. E., & Mysterud, A. (2013). Inferring spatial memory and spatiotemporal scaling
from GPS data: comparing red deer Cervus elaphus movements with simulation models. *Journal*
of animal Ecology, *82*(3), 572-586.
Gurarie, E., Bracis, C., Brilliantova, A., Kojola, I., Suutarinen, J., Ovaskainen, O., Potluri, S., & Fagan,
W. F. (2022). Spatial Memory Drives Foraging Strategies of Wolves, but in Highly Individual
Ways. *Frontiers in Ecology and Evolution*, *10*.
https://www.frontiersin.org/article/10.3389/fevo.2022.768478
Hayes, M. A. (2015). *Dispersal and population genetic structure in two flyways of sandhill cranes (Grus*
canadensis). The University of Wisconsin-Madison.
Hooten, M. B., Scharf, H. R., & Morales, J. M. (2019). Running on empty: Recharge dynamics from
animal movement data. *Ecology Letters*, *22*(2), 377–389. https://doi.org/10.1111/ele.13198
Janson, C. H., & Byrne, R. (2007). What wild primates know about resources: Opening up the black box.
Animal Cognition, *10*(3), 357–367. https://doi.org/10.1007/s10071-007-0080-9
Joly, K., Gurarie, E., Sorum, M. S., Kaczensky, P., Cameron, M. D., Jakes, A. F., Borg, B. L.,
Nandintsetseg, D., Hopcraft, J. G. C., Buuveibaatar, B., Jones, P. F., Mueller, T., Walzer, C.,
Olson, K. A., Payne, J. C., Yadamsuren, A., & Hebblewhite, M. (2019). Longest terrestrial
migrations and movements around the world. *Scientific Reports*, *9*(1), 15333.
https://doi.org/10.1038/s41598-019-51884-5
Kashetsky, T., Avgar, T., & Dukas, R. (2021). The Cognitive Ecology of Animal Movement: Evidence
From Birds and Mammals. *Frontiers in Ecology and Evolution*, *9*, 724887.
https://doi.org/10.3389/fevo.2021.724887
Lewis, M. A., Fagan, W. F., Auger-Méthé, M., Frair, J., Fryxell, J. M., Gros, C., Gurarie, E., Healy, S. D.,
& Merkle, J. A. (2021). Learning and Animal Movement. *Frontiers in Ecology and Evolution*, *9*,
681704. https://doi.org/10.3389/fevo.2021.681704
Lima, S. L., & Zollner, P. A. (1996). Towards a behavioral ecology of ecological landscapes. *Trends in*
Ecology & Evolution, *11*(3), 131-135.
MacHutchon, A. G., & Wellwood, D. W. (2003). *Grizzly Bear Food Habits in the Northern Yukon,*
Canada.
Matthiopoulos, J., J.Fieberg, and G. Aarts. (2023). Species-Habitat Associations: Spatial data, predictive
models, and ecological insights, 2nd Edition. University of Minnesota Libraries
Publishing. Retrieved from the University of Minnesota Digital
Conservancy, https://hdl.handle.net/11299/217469.
Matthysen, E. (2005). Density-dependent dispersal in birds and mammals. *Ecography*, *28*(3), 403–416.
https://doi.org/10.1111/j.0906-7590.2005.04073.x
McLoughlin, C. (2002). Learner Support in Distance and Networked Learning Environments: Ten
Dimensions for Successful Design. *Distance Education*, *23*(2), 149–162.
https://doi.org/10.1080/0158791022000009178
Merkle, J. A., Potts, J. R., & Fortin, D. (2017). Energy benefits and emergent space use patterns of an
empirically parameterized model of memory-based patch selection. *Oikos*, *126*(2).
https://doi.org/10.1111/oik.03356
Merkle, J. A., Sawyer, H., Monteith, K. L., Dwinnell, S. P. H., Fralick, G. L., & Kauffman, M. J. (2019).
Spatial memory shapes migration and its benefits: Evidence from a large herbivore. *Ecology*
Letters, *22*(11), 1797–1805. https://doi.org/10.1111/ele.13362

Miles, D., Stedman, M., & Heald, A. (2020). LIVING WITH COVID-19: BALANCING COSTS AGAINST BENEFITS IN THE FACE OF THE VIRUS. *National Institute Economic Review*, *253*, R60–R76. https://doi.org/10.1017/nie.2020.30

Morrison, T. A., Merkle, J. A., Hopcraft, J. G. C., Aikens, E. O., Beck, J. L., Boone, R. B., Courtemanch, A. B., Dwinnell, S. P., Fairbanks, W. S., Griffith, B., Middleton, A. D., Monteith, K. L., Oates, B., Riotte-Lambert, L., Sawyer, H., Smith, K. T., Stabach, J. A., Taylor, K. L., & Kauffman, M. J. (2021). Drivers of site fidelity in ungulates. *Journal of Animal Ecology*, *90*(4), 955–966. https://doi.org/10.1111/1365-2656.13425

Nathan, R., Getz, W. M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D., & Smouse, P. E. (2008). A movement ecology paradigm for unifying organismal movement research. *Proceedings of the National Academy of Sciences*, *105*(49), 19052–19059. https://doi.org/10.1073/pnas.0800375105

Nathan, R., Monk, C. T., Arlinghaus, R., Adam, T., Alós, J., Assaf, M., Baktoft, H., Beardsworth, C. E., Bertram, M. G., Bijleveld, A. I., Brodin, T., Brooks, J. L., Campos-Candela, A., Cooke, S. J., Gjelland, K. Ø., Gupte, P. R., Harel, R., Hellström, G., Jeltsch, F., … Jarić, I. (2022). Big-data approaches lead to an increased understanding of the ecology of animal movement. *Science*, *375*(6582), eabg1780. https://doi.org/10.1126/science.abg1780

Nesbitt, S. A. (1992). First Reproductive Success and Individual Productivity in Sandhill Cranes. *The Journal of Wildlife Management*, *56*(3), 573. https://doi.org/10.2307/3808874

Northrup, J. M., Anderson Jr, C. R., & Wittemyer, G. (2016). Environmental dynamics and anthropogenic development alter philopatry and space-use in a North American cervid. *Diversity and Distributions*, *22*(5), 547–557. https://doi.org/10.1111/ddi.12417

Oliveira-Santos, L. G. R., Forester, J. D., Piovezan, U., Tomas, W. M., & Fernandez, F. A. S. (2016). Incorporating animal spatial memory in step selection functions. *Journal of Animal Ecology*, *85*(2), 516–524. https://doi.org/10.1111/1365-2656.12485

Pasitschniak-Arts, M. (1993). Ursus arctos. *Mammalian Speices*, *439*, 1–10.

Picardi, S., Abrahms, B., Gelzer, E., Morrison, T. A., Verzuh, T., & Merkle, J. A. (2023). Defining null expectations for animal site fidelity. *Ecology Letters*, *26*(1), 157–169. https://doi.org/10.1111/ele.14148

Piper, W. H. (2011). Making habitat selection more "familiar": a review. *Behavioral Ecology and Sociobiology*, *65*, 1329-1351.

Potts, J. R., & Lewis, M. A. (2016). How memory of direct animal interactions can lead to territorial pattern formation. *Journal of The Royal Society Interface*, *13*(118), 20160059. https://doi.org/10.1098/rsif.2016.0059

Ranc, N., Cagnacci, F., & Moorcroft, P. R. (2022). Memory drives the formation of animal home ranges: Evidence from a reintroduction. *Ecology Letters*, *25*(4), 716–728. https://doi.org/10.1111/ele.13869

Ranc, N., Moorcroft, P. R., Ossi, F., & Cagnacci, F. (2021). Experimental evidence of memory-based foraging decisions in a large wild mammal. *Proceedings of the National Academy of Sciences*, *118*(15), e2014856118. https://doi.org/10.1073/pnas.2014856118

Rheault, H., Anderson, C. R., Bonar, M., Marrotte, R. R., Ross, T. R., Wittemyer, G., & Northrup, J. M. (2021). Some Memories Never Fade: Inferring Multi-Scale Memory Effects on Habitat Selection of a Migratory Ungulate Using Step-Selection Functions. *Frontiers in Ecology and Evolution*, *9*. https://www.frontiersin.org/article/10.3389/fevo.2021.702818

Riotte-Lambert, L., Benhamou, S., & Chamaillé-Jammes, S. (2015). How Memory-Based Movement Leads to Nonterritorial Spatial Segregation. *The American Naturalist*, *185*(4), E103–E116. https://doi.org/10.1086/680009

Sawyer, H., LeBeau, C. W., McDonald, T. L., Xu, W., & Middleton, A. D. (2019). All routes are not created equal: An ungulate's choice of migration route can influence its survival. *Journal of Applied Ecology*, 1365-2664.13445. https://doi.org/10.1111/1365-2664.13445

Schlägel, U. E., & Lewis, M. A. (2014). Detecting effects of spatial memory and dynamic information on animal movement decisions. *Methods in Ecology and Evolution*, *5*(11), 1236–1246. https://doi.org/10.1111/2041-210X.12284

Schlägel, U. E., Merrill, E. H., & Lewis, M. A. (2017). Territory surveillance and prey management: Wolves keep track of space and time. *Ecology and Evolution*, *7*(20), 8388–8405. https://doi.org/10.1002/ece3.3176

Selva, N., Teitelbaum, C. S., Sergiel, A., Zwijacz-Kozica, T., Zięba, F., Bojarska, K., & Mueller, T. (2017). Supplementary ungulate feeding affects movement behavior of brown bears. *Basic and Applied Ecology*, *24*, 68–76. https://doi.org/10.1016/j.baae.2017.09.007

Signer, J., Fieberg, J., & Avgar, T. (2019). Animal movement tools (amt): R package for managing tracking data and conducting habitat selection analyses. *Ecology and Evolution*, *9*(2), 880–890. https://doi.org/10.1002/ece3.4823

Spencer, W. D. (2012). Home ranges and the value of spatial information. *Journal of Mammalogy*, *93*(4), 929–947. https://doi.org/10.1644/12-MAMM-S-061.1

Subalusky, A. L., Dutton, C. L., Rosi, E. J., & Post, D. M. (2017). Annual mass drownings of the Serengeti wildebeest migration influence nutrient cycling and storage in the Mara River. *Proceedings of the National Academy of Sciences*, *114*(29), 7647–7652. https://doi.org/10.1073/pnas.1614778114

Tacha, T. C., Haley, D. E., & Vohs, P. A. (1989). Age of Sexual Maturity of Sandhill Cranes from Mid-Continental North America. *The Journal of Wildlife Management*, *53*(1), 43. https://doi.org/10.2307/3801303

Tello-Ramos, M. C., Branch, C. L., Kozlovsky, D. Y., Pitera, A. M., & Pravosudov, V. V. (2019). Spatial memory and cognitive flexibility trade-offs: To be or not to be flexible, that is the question. *Animal Behaviour*, *147*, 129–136. https://doi.org/10.1016/j.anbehav.2018.02.019

Thompson, P. R., Derocher, A. E., Edwards, M. A., & Lewis, M. A. (2022). Detecting seasonal episodic-like spatio-temporal memory patterns using animal movement modelling. *Methods in Ecology and Evolution*, *13*(1), 105–120. https://doi.org/10.1111/2041-210X.13743

Thompson, P. R., Lewis, M. A., Edwards, M. A., & Derocher, A. E. (2022). Time-dependent memory and individual variation in Arctic brown bears (Ursus arctos). *Movement Ecology*, *10*(1), 18. https://doi.org/10.1186/s40462-022-00319-4

Thurfjell, H., Ciuti, S., & Boyce, M. S. (2014). Applications of step-selection functions in ecology and conservation. *Movement Ecology*, *2*(1), 4. https://doi.org/10.1186/2051-3933-2-4

Trapanese, C., Meunier, H., & Masi, S. (2019). What, where and when: Spatial foraging decisions in primates: Spatial foraging decisions in primates. *Biological Reviews*, *94*(2), 483–502. https://doi.org/10.1111/brv.12462

Tulving, E., & Craik, F. I. M. (2000). *The Oxford Handbook of Memory*. Oxford University Press.

Van Moorter, B., Visscher, D., Benhamou, S., Börger, L., Boyce, M. S., & Gaillard, J.-M. (2009). Memory keeps you at home: A mechanistic model for home range emergence. *Oikos*, *118*(5), 641–652. https://doi.org/10.1111/j.1600-0706.2008.17003.x

Van Moorter, B., Visscher, D., Herfindal, I., Basille, M., & Mysterud, A. (2013). Inferring behavioural mechanisms in habitat selection studies getting the null-hypothesis right for functional and familiarity responses. *Ecography*, *36*(3), 323-330.

Wang, H., & Salmaniw, Y. (2023). Open problems in PDE models for knowledge-based animal movement via nonlocal perception and cognitive mapping. *Journal of Mathematical Biology*, *86*(5), 71. https://doi.org/10.1007/s00285-023-01905-9

Wirsing, A. J., Quinn, T. P., Cunningham, C. J., Adams, J. R., Craig, A. D., & Waits, L. P. (2018). Alaskan brown bears (*Ursus arctos*) aggregate and display fidelity to foraging neighborhoods while preying on Pacific salmon along small streams. *Ecology and Evolution*, *8*(17), 9048–9061. https://doi.org/10.1002/ece3.4431

Wolfson, D. W. (2018). *Migratory ecology and movement patterns of mid-continent and eastern sandhill cranes*. University of Minnesota.

676   Wolfson, D. W., Fieberg, J. R., & Andersen, D. E. (2020). Juvenile Sandhill Cranes exhibit wider ranging
677   and more exploratory movements than adults during the breeding season. *Ibis*, *162*(2), 556–562.
678   https://doi.org/10.1111/ibi.12786

Appendix: Supporting Information 1

Table 1. Empirical modeling for memory

Paper	Species	Memory representation	Types	Methods	Model fit	Habitat-selection predictors
Avgar et al. 2015	Woodland caribou	Dynamics in animals' habitat selection over landscape and time	Forage	Likelihood of animals moving with cognitive selection functions	Bayesian (MCMC)	1) Foraging qualities 2) Predator density (Wolf) 3) Moose habitat 4) Snow depth
Rheault et al. 2021	Mule deer	Differences between animals' occurrences throughout time	Forage	SSF framework with an OD as a response variable in the model	Step Selection Functions (SSFs) with conditional logistic regression	1) Terrain ruggedness index 2) elevation 3) distance to treed edges 4) snow depth 5) land covers 6) NDVI 7) distance to well pads 8) distance to roads 9) previous ODs (past yr) 10) current ODs (prev 7 days)
Falcon-Cortes et al. 2021	Elk	Number of visits to the foraging path	Forage	A Markovian model with the probability of moving to patches	Bayesian (MCMC)	memory decay & memory use probability based on... 1) distance between patches 2) time since last visit to the patches
Thompson et al. 2022	Brown Bear	Current resource quality + time since the last visit	Forage	Hidden Markov Model with SSF Framework Likelihood of animals moving in non-stationary states with a cognitive map	Maximum-likelihood estimates (MLE)	1) distance from turbid water to riparian areas where bears selected 2) vegetation classes 3) density of ground squirrels and alpine sweet vetch 4) distance from human settlements
Oliveira-Santos et al. 2016	Feral hog	Utilization distribution of an animal's trajectory	Home range	SSF framework with an OD as a response variable in the model	Step Selection Functions (SSFs) with conditional logistic regression	1) Land cover types 2) Time of day 3) BRB densities 4) Residence time
Northrup et al. 2016	Mule deer	Differences between animals' occurrences throughout time	Home range	SSF framework with an OD as a response variable in the model	Hierarchical Bayesian regression (MCMC)	1) Tree 2) density of drilling well pads 3) average NDVI difference 4) major road density

						5) terrain ruggedness 6) fat 7) age 8) natural gas facilities' density 9) total snow depth difference 10) pipeline density difference
Schlägel et al. 2017	Wolf	Time since the last visit	Territorial defense	Spatially explicit random walk model similar framework of... + selection-free movement kernel + selection functions	Maximum-likelihood estimates (MLE)	1) Prey density 2) Distance to territory boundary
Ranc et al. 2021	Roe deer	Food availability	Home range	Likelihood of animals moving with cognitive selection functions	Maximum-likelihood estimates (MLE)	1) Min. daily temperature 2) within state resource access 3) between-state resource access 4) illumination index (dawn & dusk) 5) changes in the illumination index
Ranc et al. 2022	Roe deer	Surrounding available habitats	Home range	Likelihood of animals moving with cognitive selection functions	Maximum-likelihood estimates (MLE)	1) Landcover - reforested 2) Landcover - agriculture 3) step lengths 4) Memory 5) Memory decay
Gurarie et al. 2022	Wolf	Predation success & Territorial marking	Territorial movement	A discrete choice modeling framework	Maximum-likelihood estimates (MLE)	1) selected zone per each trip 2) predation quality 3) boundary coverage 4) mass of prey item 5) time since predation events 6) number of kills 7) boundary sizes
Merkle et al. 2019	Mule deer	Differences between animals' occurrences throughout time	Migration	SSF framework with a distance and turning angle bias as parameters in the model	Step Selection Functions (SSFs) with conditional logistic regression	1) Elevation 2) % tree cover 3) Distance to roads 4) Terrain position index 5) Integrated NDVI 6) Rate of Green-up 7) Distance to the previous route 8) Direction to the previous range

682