A Computational Approach for the Identification of Novel L1 Transcriptional Regulators

Juan I. Bravo[1,2], Chanelle R. Mizrahi[1,3], Seungsoo Kim[4], Lucia Zhang[1,5], Yousin Suh[4,6], Bérénice A. Benayoun[1,7,8,9,10]

[1] Leonard Davis School of Gerontology, University of Southern California, Los Angeles, CA 90089, USA.

[2] Graduate program in the Biology of Aging, University of Southern California, Los Angeles, CA 90089, USA.

[3] USC Gerontology Enriching MSTEM to Enhance Diversity in Aging Program, University of Southern California, Los Angeles, CA 90089, USA.

[4] Department of Obstetrics and Gynecology, Columbia University Irving Medical Center, New York, NY 10032, USA.

[5] Quantitative and Computational Biology Department, USC Dornsife College of Letters, Arts and Sciences, Los Angeles, CA 90089, USA.

[6] Department of Genetics and Development, Columbia University Irving Medical Center, New York, NY 10032, USA.

[7] Molecular and Computational Biology Department, USC Dornsife College of Letters, Arts and Sciences, Los Angeles, CA 90089, USA.

[8] Biochemistry and Molecular Medicine Department, USC Keck School of Medicine, Los Angeles, CA 90089, USA.

[9] USC Norris Comprehensive Cancer Center, Epigenetics and Gene Regulation, Los Angeles, CA 90089, USA.

[10] USC Stem Cell Initiative, Los Angeles, CA 90089, USA.

KEYWORDS: LINE1, transposons, regulators, eQTLs, screen, aging, GWAS

ABSTRACT

Long interspersed element 1 (L1) are a family of autonomous, actively mobile transposons that occupy ~17% of the human genome. The pleiotropic effects L1 induces in host cells—promoting genome instability, inflammation, or cellular senescence—are established, and L1's associations with aging and aging diseases are widely recognized. However, because of the cell type-specific nature of transposon control, the catalogue of L1 regulators remains incomplete.

Here, we employ an eQTL approach leveraging transcriptomic and genomic data from the GEUVADIS and 1000Genomes projects to computationally identify new candidate regulators of L1 expression in lymphoblastoid cell lines. To cement the role of candidate genes in L1 regulation, we experimentally modulate the levels of top candidates *in vitro*, including *IL16, STARD5, HSDB17B12,* and *RNF5*, and assess changes in TE family expression by Gene Set Enrichment Analysis (GSEA). Remarkably, we observe subtle but widespread upregulation of TE family expression following *IL16* and *STARD5* overexpression. Moreover, a short-term 24 hour exposure to recombinant human IL16 was sufficient to transiently induce subtle but widespread upregulation of *L1* subfamilies. Finally, we find that many L1 expression-associated genetic variants are co-associated with aging traits across genome-wide association study databases.

Our results expand the catalogue of genes implicated in L1 transcriptional control and further suggest that L1 contributes to aging processes. Given the ever-increasing availability of paired genomic and transcriptomic data, we anticipate this new approach to be a starting point for more comprehensive computational scans for transposon transcriptional regulators.

BACKGROUND

Transposable elements (TEs) constitute ~45% of the human genome [1]. Among these, the long interspersed element-1 (LINE-1 or L1) family of transposons is the most abundant, accounting for ~16-17% [1, 2], and remains autonomously mobile, with humans harboring an estimated 80-100 retrotransposition-competent L1 copies [3]. These retrotransposition competent L1s belong to evolutionarily younger L1PA and L1Hs subfamilies, are ~6 kilobases long, carry an internal promoter in their 5'-untranslated region (UTR), and encode two proteins — L1ORF1p and L1ORF2p — that are necessary for transposition [4]. The remaining ~500,000 copies are non-autonomous or immobile because of the presence of inactivating mutations or truncations [1] and include L1 subfamilies of all evolutionary ages, including the evolutionarily older L1P and L1M subfamilies. Though not all copies are transposition competent, L1s can nevertheless contribute to aspects of aging [5, 6] and aging-associated diseases [7-10].

Though mechanistic studies characterizing the role of L1 in aging and aging-conditions are limited, it is clear that its effects are pleiotropic. L1 can contribute to genome instability via insertional mutagenesis and an expansion of copy number with age [11] and during senescence [12]. L1 can also play a contributing role in shaping inflammatory and cellular senescence phenotypes. The secretion of a panoply of pro-inflammatory factors is a marker of senescent cells, called the senescence associated secretory phenotype (SASP) [13]. Importantly, the SASP is believed to stimulate the innate immune system and contribute to chronic, low-grade, sterile inflammation with age, a phenomenon referred to as "inflamm-aging" [13, 14]. During deep senescence,

L1 are transcriptionally de-repressed and consequently generate cytosolic DNA that

initiates an immune response consisting of the production and secretion of pro-

inflammatory interferons [15]. Finally, L1 is causally implicated in aging-associated

diseases like cancer. L1 may contribute to cancer by (i) serving as a source for

chromosomal rearrangements that can delete tumor-suppressor genes [16] or (ii)

introducing its promoter into normally-silenced oncogenes [17]. Thus, because of the

pathological effects L1 can have on hosts, it is critical that hosts maintain precise control

over L1 activity.

Eukaryotic hosts have evolved several pre- and post-transcriptional mechanisms

for regulating TEs [18, 19]. Nevertheless, our knowledge of regulatory genes remains

incomplete because of cell type-specific regulation and the complexity of methods

required to identify regulators. Indeed, one clustered regularly interspaced short

palindromic repeats (CRISPR) screen in two cancer cell lines for regulators of L1

transposition identified >150 genes involved in diverse biological functions, such as

chromatin regulation, DNA replication, and DNA repair [20]. However, only about ~36%

of the genes identified in the primary screen exerted the same effects in both cell lines

[20], highlighting the cell type-specific nature of L1 control. Moreover, given the

complexities of *in vitro* screens, especially in non-standard cell lines or primary cells, *in*

silico screens for L1 regulators may facilitate the task of identifying and cataloguing

candidate regulators across cell and tissue types. One such attempt was made by

generating gene-TE co-expression networks from RNA sequencing (RNA-seq) data

generated from multiple tissue types of cancerous origin [21]. Although co-expression

modules with known TE regulatory functions, such as interferon signaling, were

correlated with TE modules, it is unclear whether other modules may harbor as of now

uncharacterized TE-regulating properties, since no validation experiments were carried

out. Additionally, this co-expression approach is limited, as no mechanistic directionality

can be assigned between associated gene and TE clusters, complicating the

prioritization of candidate regulatory genes for validation. Thus, there is a need for the

incorporation of novel "omic" approaches to tackle this problem. Deciphering the

machinery that controls TE activity in healthy somatic cells will be crucial, in order to

identify checkpoints lost in diseased cells.

The 1000Genomes Project and GEUVADIS Consortium provide a rich set of

genomic resources to explore the mechanisms of human TE regulation *in silico*. Indeed,

the 1000Genomes project generated a huge collection of genomic data from thousands

of human subjects across the world, including single nucleotide variant (SNV) and

structural variant (SV) data [22, 23]. To accomplish this, the project relied on

lymphoblastoid cell lines (LCLs), which are generated by infecting resting B-cells in

peripheral blood with Epstein-Barr virus (EBV). Several properties make them

advantageous for use in large-scale projects, e.g. they can be generated relatively

uninvasively, they provide a means of obtaining an unlimited amount of a subject's DNA

and other biomolecules, and they can serve as an *in vitro* model for studying the effects

of genetic variation with any phenotype of interest [24, 25]. Indeed, the GEUVADIS

Consortium generated transcriptomic data for a subset of subjects sampled by the

1000Genomes Project, and used their genomic data to define the effects genetic

variation on gene expression [26]. Together, these resources provide a useful toolkit for

investigating the genetic regulation of TEs, generally, and L1, specifically.

In this study, we (i) develop a pipeline to identify novel candidate regulators of L1

expression in lymphoblastoid cell lines, (ii) provide experimental evidence for the

involvement of top candidates in L1 expression control, and (iii) expand and reinforce

the catalogue of diseases linked to L1.

RESULTS

In silico scanning for L1 subfamily candidate regulators by eQTL analysis

To identify new candidate regulators of L1 transcription, we decided to leverage publicly available human "omic" datasets with both genetic and transcriptomic information. For this analysis, we focused on samples for which the following data was available: (i) mRNA-seq data from the GEUVADIS project, (ii) SNVs called from whole-genome sequencing data overlayed on the hg38 human reference genome made available by the 1000Genomes project, and (iii) repeat structural variation data made available by the 1000Genomes project. This yielded samples from 358 European and 86 Yoruban individuals, all of whom declared themselves to be healthy at the time of sample collection (**Figure 1A**). Using the GEUVADIS data, we obtained gene and TE subfamily expression counts using TEtranscripts [27]. As a quality control step, we checked whether mapping rates segregated with ancestry groups, which may bias results. However, the samples appeared to cluster by laboratory rather than by ancestry (**Figure S1A**). As additional quality control metrics, we also checked whether the SNV and SV data segregated by ancestry following principal component analysis (PCA). These analyses demonstrated that the top two and the top three principal components from the SNV and SV data, respectively, segregated ancestry groups (**Figure S1B, Figure S1C**).

We then chose to do a three-part integration of the available "omic" data (**Figure 1B**). Since TEtranscripts quantifies TE expression aggregated at the TE subfamily level

and discards TE position information, we chose to carry out a *trans*-eQTL analysis

against global expression of each L1 subfamily. We reasoned that there would have to

be factors (i.e., miRNAs, proteins, non-coding RNAs) mediating the effects of SNVs on

L1 subfamily expression. Thus, to identify candidate genic mediators, we searched for

genes with *cis*-eQTLs that overlapped with L1 *trans*-eQTLs. As a final filter, we

reasoned that for a subset of regulators, L1 subfamily expression would respond to

changes in the expression of those regulators. Consequently, we chose to quantify the

association between L1 subfamily expression and candidate gene expression by linear

regression. We hypothesized that this three-part integration would result in

combinations of significantly correlated SNVs, genes, and L1 subfamilies (**Figure 1B**).

The *trans*-eQTL analysis against every expressed L1 subfamily led to the

identification of 499 *trans*-eQTLs distributed across chromosomes 6, 11, 12, 14, and 15

that passed genome-wide significance (**Figure 1C, Supplementary Table S1A**). The

cis-eQTL analysis led to the identification of 845,260 *cis*-eQTLs that passed genome-

wide significance (**Supplementary Figure S2, Supplementary Table S1B**). After

integrating the identified *cis*- and *trans*-eQTLs and running linear regression, we

identified 1,272 SNV-Gene-L1 trios that fulfilled our three-part integration approach

(**Supplementary Table S1C**). Among this pool of trios, we identified 7 unique protein-

coding genes including *IL16, STARD5, HLA-DRB5, HLA-DQA2, HSD17B12, RNF5, and*

FKBPL (**Figure 1C**). We note that although *EHMT2* did not pass out screening

approach, it does overlap *EHMT2-AS1*, which did pass our screening thresholds

(**Figure 1C**). We also note that several other unique non-coding genes, often

overlapping the protein-coding genes listed, were also identified (**Figure 1C**). For

simplicity of interpretation, we focused on protein-coding genes during downstream

experimental validation.

Next, to define first and second tier candidate regulators, we clumped SNVs in

linkage disequilibrium (LD) by L1 *trans*-eQTL p-value to identify the most strongly

associated genetic variant in each genomic region (**Figure 2A, Supplementary Figure**

S3A). LD-clumping identified the following index SNVs (*i.e.* the most strongly associated

SNVs in a given region): rs11635336 on chromosome 15, rs9271894 on chromosome

6, rs1061810 on chromosome 11, rs112581165 on chromosome 12, and rs72691418

on chromosome 14 (**Supplementary Table S1D**). Genes linked to these SNVs were

considered first tier candidate regulators and included *IL16*, *STARD5*, *HLA-DRB5*, *HLA-*

DQA2, and *HSD17B12* (**Figure 2B, Supplementary Table S1E**). The remaining genes

were linked to clumped, non-index SNVs and were consequently considered second tier

candidates and included *RNF5*, *EHMT2-AS1*, and *FKBPL* (**Supplementary Figure**

S3B). Additionally, for simplicity of interpretation, we considered only non-*HLA* genes

during downstream experimental validation, since validation could be complicated by

the highly polymorphic nature of *HLA* loci [28] and their involvement in multi-protein

complexes.

Finally, to computationally determine whether candidate genes may causally

influence L1 subfamily expression, we carried out mediation analysis on all SNV-gene-

L1 trios (**Supplementary Figure S4A**). Interestingly, 868 out of the 1,272 (68.2%) trios

exhibited significant (FDR < 0.05) mediation effects (**Supplementary Table S1F**).

Among the 1st tier candidate regulators, significant, partial, and consistent mediation

effects could be attributed to *STARD5*, *IL16*, *HSD17B12*, and *HLA-DRB5*

(**Supplementary Figure S4B, Supplementary Table S1F**). To note, while significant

mediation could be attributed to the index SNV for *STARD5*, significant mediation could

only be attributed to clumped SNVs for *IL16* and *HSD17B12*. Given that *STARD5* and

IL16 share *cis*-eQTL SNVs, this suggests that *STARD5* may be the more potent

mediator. Among the 2nd tier candidate regulators, significant, partial, and consistent

mediation effects could be attributed to *RNF5*, *EHMT2-AS1*, and *FKBPL*

(**Supplementary Figure S4C, Supplementary Table S1F**). These results suggest that

candidate genes may mediate the effects between linked SNVs and L1 subfamilies.

In silico scanning for L1 subfamily candidate regulators in an African population

We next sought to assess the cross-ancestry regulatory properties of candidate

genes by repeating our scan using the Yoruban samples as a smaller but independent

replication cohort. Here, rather than conduct a genome-wide scan for *cis*- and *trans*-

associated factors, we opted for a targeted approach focusing only on gene *cis*-eQTLs

and L1 subfamily *trans*-eQTLs that were significant in the analysis with European

samples (**Supplementary Figure S5A**). The targeted *trans*-eQTL analysis led to the

identification of 227 significant (FDR < 0.05) *trans*-eQTLs distributed across

chromosomes 6 and 11 (**Supplementary Table S2A**). The targeted *cis*-eQTL analysis

led to the identification of 1,248 significant (FDR < 0.05) *cis*-eQTLs (**Supplementary**

Table S2B). After integrating the identified *cis*- and *trans*-eQTLs and running linear

regression, we identified 393 SNV-Gene-L1 trios that fulfilled our three-part integration

approach (**Supplementary Table S2C**). Among this pool of trios, we identified 2 unique

protein-coding genes—*HSD17B12 and HLA-DRB6*—as well as several unique non-

coding genes (**Supplementary Table S2C**). Again, we clumped SNVs in linkage

disequilibrium (LD) by L1 *trans*-eQTL p-value. LD-clumping identified the following index

SNVs: rs2176598 on chromosome 11 and rs9271379 on chromosome 6

(**Supplementary Table S2D**). Genes linked to these SNVs were considered first tier

candidate regulators and included both *HSD17B12* and *HLA-DRB6* (**Supplementary**

Figure S5B, Supplementary Table S2E). Finally, we carried out mediation analysis on

all SNV-gene-L1 trios; however, no significant (FDR < 0.05) mediation was observed

(**Supplementary Table S2F**). These results implicate *HSD17B12* and the *HLA* loci as

candidate, cross-ancestry L1 expression regulators.

To assess why some candidate genes did not replicate in the Yoruba cohort, we

manually inspected *cis*- and *trans*-eQTL results for trios with those genes

(**Supplementary Figure S6A**). Interestingly, we identified rs9270493 and rs9272222 as

significant (FDR < 0.05) *trans*-eQTLs for L1MEb expression. However, those SNVs

were not significant *cis*-eQTLs for *RNF5* and *FKBPL* expression, respectively. For trios

involving *STARD5, IL16,* and *EHMT2-AS1*, neither the *cis*-eQTL nor the *trans*-eQTL

were significant. We note that for most of these comparisons, although the two

genotypes with the largest sample sizes were sufficient to establish a trending change

in *cis* or *trans* expression, this trend was often broken by the third genotype with

spurious sample sizes. This suggests that replication in the Yoruba cohort may be

limited by the small cohort sample size in the GEUVADIS project.

TE families and known TE-associated pathways are differentially regulated across L1

trans-eQTL variants

Though our eQTL analysis identified genetic variants associated with the

expression of specific, evolutionarily older L1 subfamilies, we reasoned that there may

be more global but subtle differences in TE expression profiles among genotype groups,

given that TE expression is highly correlated [21]. Thus, for each gene-associated index

SNV identified in the European eQTL analysis, we carried out differential expression

analysis for all expressed genes and TEs (**Supplementary Table S3A-S3C**; **Figure**

3A). At the individual gene level, we detected few significant (FDR < 0.05) changes: 4

genes/TEs varied with rs11635336 genotype (*IL16*/*STARD5*), 4 genes/TEs varied with

rs9271894 genotype (*HLA*), and 5 gene/TEs varied with rs1061810 genotype

(*HSD17B12*) (**Supplementary Table S3A-S3C**). Importantly, however, these

genes/TEs overlapped the genes/TEs identified in the *cis*- and *trans*-eQTL analyses,

providing an algorithmically independent link among candidate SNV-gene-TE trios.

In contrast to gene-level analyses, Gene Set Enrichment Analysis (GSEA)

provides increased sensitivity to subtle but consistent/widespread transcriptomic

changes at the level of gene sets (*e.g.* TE families, biological pathways, etc.). Thus, we

leveraged our differential expression analysis in combination with GSEA to identify

repeat family and biological pathway gene sets impacted by SNV genotype in the

GEUVADIS dataset (**Supplementary Table S3D-S3O**; **Figure 3A**). Interestingly,

changes in the genotype of rs11635336 (*IL16*/*STARD5*), rs9271894 (*HLA*), and

rs1061810 (*HSD17B12*) were associated with an upregulation, upregulation, and

downregulation, respectively, of multiple TE family gene sets (**Figure 3B,**

Supplementary Table S3P). Differentially regulated TE family gene sets included DNA

transposons, such as the hAT-Charlie family, and long terminal repeat (LTR)

transposons, such as the endogenous retrovirus-1 (ERV1) family (**Figure 3B,**

Supplementary Table S3P). Noteworthy, the L1 family gene set was the only TE gene

set whose expression level was significantly altered across all three SNV analyses

(**Figure 3B, Supplementary Table S3P**). Consistent with their relative significance in

the L1 *trans*-eQTL analysis, the L1 family gene set was most strongly upregulated by

alternating the *IL16*/*STARD5* SNV (NES = 3.74, FDR = 6.43E-41), intermediately

upregulated by alternating the *HLA* SNV (NES = 1.90, FDR = 7.19E-5), and least

strongly changed by alternating the *HSD17B12* SNV (NES = -1.57, FDR = 2.11E-2)

(**Figure 3C**). We briefly note here that rs9270493, a clumped SNV linked to *RNF5*, was

also linked to upregulation of the L1 family gene set (**Supplementary Table S3Q-S3R**).

These results suggest that TE subfamily *trans*-eQTLs are associated with subtle but

global differences in TE expression beyond a lone TE subfamily.

Next, we asked if other biological pathways were regulated concomitantly with

TE gene sets in response to gene-linked index SNVs, reasoning that such pathways

would act either upstream (as regulatory pathways) or downstream (as response

pathways) of TE alterations. GSEA with the MSigDB Hallmark pathway gene sets [29,

30] identified 5 gene sets fitting this criterion, including "oxidative phosphorylation",

"mTORC1 signaling", "fatty acid metabolism", "adipogenesis", and "cholesterol

homeostasis" (**Figure 3D, Supplementary Table S3S**). Interestingly, several of these

pathways or genes in these pathways have been implicated in TE regulation before.

Rapamycin, which acts through mTORC1, has been shown to alter the expression of L1

and other repeats [31, 32]. Estrogens, which are involved in cholesterol and lipid

metabolism, have been found to drive changes in repeat expression, and the receptors

for both estrogens and androgens are believed to bind repeat DNA [32, 33].

Pharmacological inhibition of the mitochondrial respiratory chain and pharmacological

reduction of endogenous cholesterol synthesis have also been shown to induce

changes in L1 protein levels or repeat expression more broadly [34, 35]. GSEA with the

GO Biological Process gene sets (**Figure 3E, Supplementary Table S3T**) and the

Reactome gene sets (**Figure 3F, Supplementary Table S3U**) also identified several

metabolism-related pathways including "ATP metabolic process", "Generation of

precursor metabolites and energy", and "metabolism of amino acids and derivatives".

These results add to the catalogue of pathways associated with differences in L1

expression.

In our eQTL analysis, we also identified two orphan index SNVs, rs112581165

and rs72691418, to which we could not attribute a protein-coding gene mediator. To

determine whether these SNVs also regulate any transposon families or biological

pathways, we repeated the differential expression analysis (with all expressed genes

and TEs) (**Supplementary Table S4A-S4B**) and the GSEA (**Supplementary Table**

S4C-S4J) with these SNVs (**Supplementary Figure S7A**). At the individual gene level,

we detected 3193 genes/TEs that varied significantly (FDR < 0.05) with rs112581165

genotype and 1229 genes/TEs that varied significantly with rs72691418 genotype

(**Supplementary Table S4A-S4B**). Similar to above, we next carried out GSEA to

identify changes in functionally relevant gene sets. Like the gene-linked index SNVs,

changes in the genotype of rs112581165 and rs72691418 were both associated with a

downregulation and upregulation, respectively, of 10 TE families (**Supplementary**

Figure S7B, Supplementary Table S4K). Noteworthy, the L1 family gene set was

among the most strongly dysregulated TE family gene sets for both rs112581165 (NES

= -4.32, FDR = 5.18E-89) and rs72691418 (NES = 4.01, FDR = 5.38E-79)

(**Supplementary Figure S7C**). These results suggest that TE subfamily *trans*-eQTLs

are associated with subtle differences in TE expression beyond the lone TE subfamily,

even in the absence of a protein-coding gene *cis*-eQTL.

Like before, we asked if other biological pathways were regulated concomitantly

with TE gene sets in response to orphan index SNVs. The top 10 Hallmark pathway

gene sets identified by GSEA included gene sets that were previously identified

("oxidative phosphorylation", "fatty acid metabolism", and "mTORC1 signaling"), as well

as several new pathways (**Supplementary Figure S7D, Supplementary Table S4L**).

Among the new pathways, "DNA repair" [20] and the "P53 pathway" [36, 37] have also

been linked to L1 control, and proteins in the "Myc targets v1" gene set interact with L1

ORF1p [38]. GSEA with the GO Biological Process gene sets (**Supplementary Figure**

S7E, Supplementary Table S4M) and the Reactome gene sets (**Supplementary**

Figure S7F, Supplementary Table S4N) identified several metabolism-related pathways and several translation-related pathways, such as "cytoplasmic translation", "eukaryotic translation initiation", and "eukaryotic translation elongation". Importantly, proteins involved in various aspects of proteostasis have been shown to be enriched among L1 ORF1p-interacting proteins [38]. Again, these results add to the catalogue of pathways associated with differences in TE expression, even in the absence of a candidate *cis* mediator.

Modulation of top candidate gene activity in a lymphoblastoid cell line induces small but widespread TE expression changes

We decided to validate the L1 regulatory properties of top candidate genes associated with L1 *trans*-eQTLs. For experimental purposes, we selected the GM12878 lymphoblastoid cell line, because (i) it is of the same cell type as the transcriptomic data used here for our eQTL analysis, and (ii) its epigenomic landscape and culture conditions have been well well-characterized as part of the ENCODE project [39, 40]. For validation purposes, we selected *IL16*, *STARD5*, *HSD17B12*, and *RNF5* out of the 7 protein-coding gene candidates. We chose these genes for validation because the first 3 are associated with top *trans*-eQTL SNVs and the fourth one had very strong predicted mediation effects. To note, although GM12878 was part of the 1000Genomes Project, it was not included in the GEUVADIS dataset. However, based on its genotype, we can predict the relative expression of candidate regulators (**Supplementary Figure S8A**), which suggest that GM12878 may be most sensitive to modulations in *IL16* and

STARD5 expression, given their relatively low endogenous expression. Interestingly,

examination of the ENCODE epigenomic data in GM12878 cells [39] demonstrated that

the region near the *IL16*/*STARD5*-linked index SNV (rs11635336) was marked with

H3K4Me1 and H3K27Ac, regulatory signatures of enhancers (**Supplementary Figure**

S8C). Similarly, the region near the *HLA*-linked index SNV (rs9271894) was marked

with H3K4Me1, marked with H3K27Ac, and accessible by DNase, suggesting regulatory

properties of the region as an active enhancer (**Supplementary Figure S8C**). These

results further highlight the regulatory potential of the *IL16*-, *STARD5*-, and *HLA*-linked

SNVs.

First, we decided to test the transcriptomic impact of overexpressing our top

candidates in GM12878 LCLs. Cells were electroporated with overexpression plasmids

(or corresponding empty vector), and RNA was isolated after 48h (**Figure 4A,**

Supplementary Figure S9A). Differential expression analysis comparing control and

overexpression samples confirmed the overexpression of candidate genes

(**Supplementary Figure S9B, Supplementary Table S5A-S5D**). Intriguingly, we

observed that *IL16* was significantly upregulated following *STARD5* overexpression

(**Supplementary Figure S9C, Supplementary Table S5B**), although the inverse was

not observed (**Supplementary Figure S9C, Supplementary Table S5A**), suggesting

that *IL16* may act downstream of *STARD5*. We note here that, consistent with the use

of a high expression vector, the *IL16* upregulation elicited by *STARD5* overexpression

($\log_2$ fold change = 0.45) was weaker than the upregulation from the *IL16*

overexpression ($\log_2$ fold change = 1.89) (**Supplementary Table S5A-S5B**).

To further assess the biological relevance of each overexpression, we carried out GSEA using the GO Biological Process, Reactome pathway, and Hallmark pathway gene sets (**Supplementary Table S5E-S5P**). Importantly, GSEA using GO Biological Process and Reactome pathway gene sets highlighted differences that were consistent with the known biology of our candidate genes. Firstly, *IL16* is involved in regulating T-cell activation, B-cell differentiation, and functions as a chemoattractant [41-46]. Moreover, it modulates macrophage polarization by regulating *IL-10* expression [47]. *IL16* overexpressing cells showed upregulation for "phagocytosis recognition" and "positive chemotaxis", downregulation for "negative regulation of cell differentiation", and downregulation for "Interleukin 10 signaling" (**Figure 4B-4C**). Secondly, *STARD5* encodes a cholesterol transporter and is upregulated in response to endoplasmic reticulum (ER) stress [48-50]. *STARD5* overexpressing cells showed downregulation of various cholesterol-related gene sets such as "sterol biosynthetic process", "sterol metabolic process", and "regulation of cholesterol biosynthesis by SREBP (SREBF)" (**Figure 4D-4E**). Thirdly, *HSD17B12* encodes a steroid dehydrogenase involved in converting estrone into estradiol and is essential for proper lipid homeostasis [51-53]. *HSD17B12* overexpressing cells showed downregulation of cholesterol-related gene sets, including "sterol biosynthetic process" and "regulation of cholesterol biosynthesis by SREBF (SREBP)" (**Supplementary Figure S9D-S9E**). Finally, *RNF5* encodes an ER and mitochondrial-bound E3 ubiquitin-protein ligase that ubiquitin-tags proteins for degradation [54-57]. *RNF5* overexpressing cells demonstrated alterations in gene sets involved in proteostasis and ER biology, including upregulation of "ERAD pathway",

"response to endoplasmic reticulum stress", and "intra-Golgi and retrograde Golgi-to-ER

traffic" (**Supplementary Figure S9F-S9G**). These results suggest that our approach

leads to biological changes consistent with the known biological impact of the genes

being overexpressed.

Next, we sought to determine whether modulation of candidate genes had any

impact on TE expression in general, and L1 in particular. Although there were no

significant changes for individual TE subfamilies following *IL16* and *STARD5*

overexpression (**Supplementary Table S5A-S5B**), we identified subtle but widespread

upregulation of various TE families across both conditions by GSEA (**Figure 4F,**

Supplementary Table S5Q-S5R). Interestingly, 7 families, including L1, ERV1, ERVL-

MaLR, Alu, ERVL, TcMar-Tigger, and hAT-Charlie families, were commonly

upregulated under both conditions (**Figure 4F**). In contrast, cells overexpressing

HSD17B12 or *RNF5* did not drive widespread changes in L1 family expression, as

assessed by GSEA (**Supplementary Table S5S-S5T**). Noteworthy, the L1 family gene

set was more strongly upregulated following *STARD5* overexpression (NES = 2.25,

FDR = 6.14E-7) compared to *IL16* overexpression (NES = 2.24, FDR = 2.40E-5)

(**Figure 4G, Supplementary Table S5Q-S5R**). Since *IL16* is upregulated in response to

STARD5 overexpression, this suggests that *STARD5* may synergize with *IL16* for the

regulation of L1 transcription.

Then, we decided to further characterize the impact of IL16 activity on TEs, since

(i) its overexpression led to a global upregulation of TE transcription, and (ii) it was itself

upregulated in response to *STARD5* overexpression, which also led to increased TE

expression. Thus, since IL16 is a soluble cytokine, we independently assessed its

regulatory properties by exposing GM12878 cells to recombinant human IL16 peptide

[rhIL16] for 24 hours (**Figure 5A, Supplementary Figure S10A**). Differential gene

expression analysis (**Supplementary Table S6A**) and comparison with the *IL16*

overexpression results demonstrated that differentially expressed genes were weakly

but significantly correlated (**Supplementary Figure S10B**). Additionally, we carried out

GSEA using the GO Biological Process, Reactome pathway, and Hallmark pathway

gene sets (**Supplementary Table S6B-S6E**) and compared those results with the

GSEA from the *IL16* overexpression (**Supplementary Table S6F-S6H**). Consistent with

the known biology of *IL16*, GSEA highlighted a downregulation of many immune cell-

related gene sets, including "leukocyte differentiation", "mononuclear cell differentiation",

and "Interleukin-10 signaling" (**Figure 5B-5C, Supplementary Table S6F-S6H**). Like

the overexpression results, exposure of GM12878 to rhIL16 for 24 hours led to an

upregulation of an L1 family gene set by GSEA, although the effect was less

pronounced than with the overexpression (**Figure 5D**). Even though treatment of

GM12878 with rhIL16 for 48 hours exhibited known features of IL16 biology

(**Supplementary Figure S10B-S10D, Supplementary Table S6J-S6Q**), the L1

upregulation was no longer detectable, though other TEs remained upregulated

(**Supplementary Figure S10E, Supplementary Table S6Q**). These results further

support the notion that *IL16* acts as a modulator of L1 expression.

Finally, we sought to define the biological pathways regulated concomitantly with

the L1 family gene set under all experimental conditions where it was upregulated (i.e.,

IL16 overexpression, *STARD5* overexpression, and 24 hours of rhIL16 exposure)

(**Figure 6A, Figure 6B, Supplementary Table S7A**). Again, we reasoned that such

pathways would act either upstream (as regulatory pathways) or downstream (as

response pathways) of TE alterations. GSEA with the Hallmark pathway gene sets

identified 7 gene sets fitting this criterion, including "TNFα signaling via NF-KB", "IL2

STAT5 signaling", "inflammatory response", "mTORC1 signaling", "estrogen response

early", "apoptosis", and "UV response up" (**Figure 6C, Supplementary Table S7B**).

GSEA with the GO Biological Process gene sets (**Figure 6D, Supplementary Table**

S7C) and the Reactome pathway gene sets (**Figure 6E, Supplementary Table S7D**)

also identified MAPK signaling, virus-related pathways like "HCMV early events",

pathways involved in cell differentiation, and pathways involved in cholesterol and

steroid metabolism like "signaling by nuclear receptors". These results further cement

the catalogue of pathways associated with differences in TE expression.

L1 trans-eQTLs are co-associated with aging traits in GWAS databases.

Although TE de-repression has been observed broadly with aging and age-

related disease [5, 58], whether this de-repression acts as a causal driver, or a

downstream consequence, of aging phenotypes remains unknown. We reasoned that if

increased TE expression at least partially drives aging phenotypes, L1 *trans*-eQTLs

should be enriched for associations to aging traits in genome-wide association studies

[GWAS] or phenome-wide association studies [PheWAS].

To test our hypothesis, we queried the Open Targets Genetics platform with our

499 *trans*-eQTL SNVs, mapped traits to standardized MeSH IDs, and then manually

curated MeSH IDs related to aging-related traits (**Figure 7A**). Consistent with our

hypothesis, a large proportion of L1 *trans*-eQTL SNVs (222/499 or 44.5%) were either

(i) associated with an aging MeSH trait by PheWAS or (ii) LD-linked to a lead variant

associated with an aging MeSH trait (**Figure 7B**). Moreover, among the 222 SNVs with

significant aging-trait associations, we observed frequent mapping to more than a single

age-related trait by PheWAS, with many SNVs associated with 10-25 traits (**Figure 7C,**

Supplementary Table S8A). Additionally, many of the 222 SNVs mapped to 1-5 aging

traits through a proxy lead variant (**Figure 7D, Supplementary Table S8A**). Among the

most frequently associated or linked traits, we identified type 2 diabetes mellitus,

hyperparathyroidism, thyroid diseases, coronary artery disease, hypothyroidism, and

psoriasis, among many others (**Figure 7E, Supplementary Table S8B**).

As a parallel approach, we queried the Open Targets Genetics platform with our

L1 *trans*-eQTL SNVs, as well as 500 combinations of random SNVs sampled from all

SNVs used in the eQTL analyses. We then leveraged broader phenotype categories

annotated by the platform, including 14 disease categories that we considered aging-

related, to determine whether L1 eQTL associations were enriched for any disease

categories (**Supplementary Figure S11A**). L1 eQTL associations were significantly

enriched (FDR < 0.05 and ES > 1) for 13 out of 14 disease categories, including cell

proliferation disorders, immune system diseases, and musculoskeletal diseases

(**Supplementary Figure S11B-N**). The cardiovascular diseases category was the only

disease category for which we did not observe a significant enrichment

(**Supplementary Figure S11O**). The enrichment for cell proliferation disorders is

consistent with the associations of L1 activity with cellular senescence [12, 15] and

cancer [59, 60]. The enrichment for immune system diseases is consistent with the role

of L1 as a stimulator of the interferon pathway, inflammation, and senescence [15], as

well as the more general notion that transposons can mimic viruses and stimulate

immune responses from their hosts [61]. The enrichment for musculoskeletal diseases

is consistent with an increase in L1 expression and copy number with age in muscle

tissue from aging mice [11]. These results reinforce the notion that L1 activity is strongly

and non-randomly associated with an assortment of age-related diseases.

Intriguingly, a large fraction of co-associated SNVs were on chromosome 6 near

the HLA locus, which has previously been shown to be a hotspot of age-related disease

traits [62]. Despite its association to our strongest L1 trans-eQTL SNV, little is known

about the regulation and impact of IL16 during aging. One study, however, found that

IL16 expression increases with age in ovarian tissue, and the frequency of *IL16*

expressing cells is significantly higher in ovarian tissue from women at early and late

menopause, compared to premenopausal women [63]**.** Given these findings, and since

L1 expression levels and copy number have been found to increase with age [5], we

asked whether circulating IL16 levels may also change with age, using C57BL/6JNia

mice as a model (**Figure 7F, Supplementary Table S8C**). Consistent with the notion

that increased IL16 levels may, at least partially, drive age-related TE de-repression, we

observed a significant increase in circulating IL16 levels in female mice with age, and a

trending increase with age in male mice (although the levels showed more animal-to-

animal variability). By meta-analysis, circulating IL16 levels changed significantly with

age across sexes (**Figure 7F**). These results further support the hypothesis that *IL16* is

involved in L1 biology and may modulate L1 age-related changes. In sum, our results

provide one of the first pieces of evidence of a causal link between L1 expression levels

and age-related decline.

DISCUSSION

In this work, we developed a pipeline to computationally identify candidate L1 transcriptional regulators by eQTL analysis. We provide experimental evidence for the involvement of top candidates in regulating L1 expression, demonstrating as a proof-of-principle that this approach can be broadly used on other large "omic"-characterized cohorts with human (i.e. GTEx [64, 65] or HipSci [66]) or mouse (i.e. DO mice [67]) subjects to identify other regulators of L1 activity. These datasets, combined with our approach, could be utilized to rigorously characterize conserved or group-specific TE regulatory mechanisms on multiple layers, such as across TE families (like Alu or ERVs), across cell or tissue types, across ancestry groups, and across species. This approach, which leverages existing datasets to perform *in silico* screening, could be a powerful method to expand our knowledge of TE regulation in non-diseased cells and tissues.

While we believe this approach can readily be applied to other datasets, we would like to note potential limitations with the approach implemented here, some of which were simply beyond the scope of this paper. Firstly, though it is common to use probabilistic estimation of expression residuals (PEER) [68] to enhance detection of *cis*-eQTLs, PEER was not implemented in our analysis as a precautionary measure, in order to avoid potentially blurring global TE signals, which likely led to a more conservative list of candidate *cis* gene mediators. Second, given the technical complexity in generating the vast amount of mRNA-seq data used for the eQTL analysis, it is possible that technical covariates introduced non-linear effects that would

not be easily removed by approaches like PEER or SVA [69]. For that reason, we opted

to supplement our computational predictions with experimental data. Third, the L1 *trans*-

eQTLs identified were specific to older L1 subfamilies (L1P and L1M) and were not

shared across subfamilies. One factor that may partially explain this is the heightened

difficulty of quantifying the expression of evolutionarily younger L1 subfamilies using

short-read sequencing [70]. More generally, significant single gene differences are often

difficult to reproduce across studies, and it is for this reason that methods like GSEA

were developed, to robustly identify broader changes in sets of genes [29]. Consistently,

GSEA suggests that many TE families, beyond the single L1 subfamilies identified in

the eQTL analysis, are differentially regulated among samples with different genotypes

for *trans*-eQTL SNVs and among samples where *IL16*/IL16 and *STARD5* were

manipulated. We note that although *HLA* and *HSD17B12* loci were significant in both

the European and African cohorts, we were not able to independently identify all of the

same candidate regulators. This is likely due to a combination of small sample size for

the African cohort and the existence of population-specific L1 regulation. Future studies

with larger sample sizes may be useful for expanding the catalogue of loci that are

biologically meaningful for L1 expression across more than one population. Importantly,

our computational scan is limited to loci exhibiting genomic variation among tested

individuals. This will vary with factors like the ancestry groups of the populations being

studied. Moreover, variants that confer extreme fitness defects may not exist at a

sufficiently high level in a population so as to allow for an assessment of their

involvement as eQTLs. Also, a potential "blindspot" of our current approach is that it

does not distinguish between TE reads of intronic or intergenic origin. Intronic TE RNA

is often considered less interesting from a biological perspective since their presence

may be attributed to readthrough transcription [71]. However, all analyses carried out in

this study relied on polyA-selected libraries, which should be enriched for mature gene

and transposon transcripts, minimizing the presence of readthrough transcribed L1

RNA. Additionally, even if L1-containing transcripts were quantified, these have been

implicated in tightly controlled functions like T-cell quiescence maintenance [31],

suggesting that intronic L1 RNA possesses biologically meaningful properties. Thus,

this pipeline should help expand our understanding of L1 biology in either case. Finally,

although we focused on protein-coding candidate regulators, it is possible that the non-

coding genes identified in our scan may also causally drive differences in L1

expression. Though not explored here, other regulatory factors like small RNAs may

also act as partial mediators. Since the GEUVADIS Consortium generated small RNA

data in parallel to the mRNA data used in this study [26], in the future, our pipeline could

be adapted to scan for *cis* small RNA mediators relatively easily. These unexplored

factors may explain the associations between orphan SNV genotypes and TE family

gene set changes.

Despite potential limitations, our approach identified *IL16, STARD5, HLA-DRB5,*

HLA-DQA2, HSD17B12, RNF5, FKBPL, and *EHMT2-AS1* as candidate L1 regulators in

the European cohort. Moreover, the upregulation of several TE gene sets following

*IL16/*IL16 and *STARD5* manipulation highlights a causal role for these genes in TE

control. Between these two genes, multiple lines of evidence suggest that *STARD5* is

the more potent mediator. First, the three-part integration statistics are more significant

for *STARD5* than for *IL16*. Second, the index SNV on chromosome 15 exhibited

significant mediation effects through *STARD5* but not *IL16*; the most significant

mediation effect for *IL16* was linked to a clumped SNV. Third, the upregulation for the

L1 family gene set was stronger under *STARD5* overexpression than under *IL16*

overexpression. Importantly, we observed that *IL16* was upregulated following *STARD5*

overexpression, but this upregulation was less than the upregulation from the *IL16*

overexpression. This suggests that *IL16* can participate in the L1-regulating properties

exerted by *STARD5*. Moving forward, it will be informative to assess the effects of *IL16*

and *STARD5* on L1 expression in other cell types. Interestingly, other genes like

EHMT2 have previously been linked to retrotransposons [72, 73]. For the remaining

genes that we experimentally tested, we note that GM12878 is predicted to have

relatively high endogenous *HSD17B12* expression and intermediate *RNF5* expression,

based on the GM12878 genotypes at *cis*-eQTLs for these genes. Given these

expression patterns, GM12878 may not be sensitive to overexpression of *RNF5*, and

especially insensitive to *HSD17B12* overexpression. For these two candidates, cells

may be more sensitive to knockdown- or knockout-based approaches. Indeed,

HSD17B12 is essential for mouse development, *HSD17B12* knockout in adult mice

results in reduced body weight and liver toxicity, and knockdown of the *Caenorhabditis*

elegans ortholog for *HSD17B12* reduces lipid stores and blocks induction of the

unfolded protein response of the endoplasmic reticulum [52, 74, 75]. Given that TEs are

often derepressed when homeostasis is challenged [61], such as following *HSD17B12*

knockout/knockdown, it remains possible that *HSD17B12* possesses L1 regulatory

properties that were not detectable by our approach. Future work could avert such

technical limitations by testing the impact of candidate genes by both up- and down-

regulation, or by selecting LCL samples with endogenous target gene expression levels

that would be most sensitive to our overexpression approach.

As another, theoretical line of evidence for the potential involvement of candidate

genes in L1 regulation, we highlight known interactions between tested candidate genes

and viral infections, which may be relevant under conditions where transposons are

recognized as viral mimics [61]. Indeed, *IL16* has been extensively studied for its ability

to inhibit human immunodeficiency virus (HIV) replication, partly by suppressing mRNA

expression [76-78]. Additionally, but in contrast to its HIV-suppressive properties, *IL16*

can enhance the replication of influenza A virus (IAV) and facilitate its infection of hosts,

potentially through its repression of type I interferon beta and interferon-stimulated

genes [79]. *IL16* can also contribute to the establishment of lifelong gamma herpesvirus

infection [80]. *STARD5* is another candidate implicated in the influenza virus replication

cycle [81]. *HSD17B12* promotes the replication of hepatitis C virus via the very-long-

chain fatty acid (VLCFA) synthesis pathway and the production of lipid droplets

important for virus assembly [82, 83]. Additionally, HSD17B12 has been found

interacting with the coronavirus disease 2019 (COVID-19) protein nonstructural protein

13 (NSP13), which is thought to antagonize interferon signaling [84]. Finally, *RNF5* has

been implicated in both promoting and antagonizing severe acute respiratory syndrome

coronavirus 2 (SARS-CoV-2) by either stabilizing the interactions of membrane protein

(M) [85] or inducing degradation of structural protein envelope (E) [86], respectively.

Fundamentally, *RNF5* regulates virus-triggered interferon signaling by targeting the

stimulator of interferon genes (STING) or mitochondrial antiviral signaling protein

(MAVS) for ubiquitin-mediated protein degradation [56, 57]. These studies reinforce the

roles of tested candidate regulators in virus-associated processes, including interferon-

mediated signaling.

Consistent with the notion that L1 is associated with aging and aging phenotypes

[5, 58], we observed that L1 *trans*-eQTL SNVs were associated with aging phenotypes

in GWAS/PheWAS databases. This is very surprising, but interesting, given that all

1000Genomes Project participants declared themselves to be healthy at the time of

sample collection. Assuming this to be true, our results suggest that L1 expression

differences exist in natural, healthy human populations, and these expression

differences precede onset of aging diseases. Though it is often unclear whether L1 mis-

regulation is a consequence or driver of aging phenotypes, our results suggest that L1

levels may drive aging phenotypes. As we continue to expand the catalogue of L1

regulators, especially in healthy cells and tissues, the L1 regulatory processes that are

disrupted over the course of aging will become increasingly clear. To that end, this work

may serve as a guide for conducting more comprehensive scans for candidate TE

regulators.

CONCLUSIONS

We developed an eQTL-based pipeline that leverages genomic and transcriptomic data to scan the human genome for novel candidate regulators of L1 subfamily expression. Though the initial scan identified genetic variants associated with expression differences in specific L1 subfamilies, secondary analyses by GSEA suggest that genetic variants are associated with subtle but global differences in the expression of many TE families. Our pipeline identified candidate genes, including *HSD17B12* and *HLA* genes, that likely play a conserved role in L1 regulation across human populations of different ancestries. Though some top candidates from the European cohort scan, such as *IL16, STARD5*, and *RNF5*, were not significant in the African cohort analysis, it is likely that some of these genes would appear in cross-ancestry scans with larger samples sizes. We detected subtle but global differences in L1 family expression following *IL16* overexpression, *STARD5* overexpression, and rhIL16 treatment for 24 hours, further suggesting that some candidate genes have regulatory potential. We generate a list of pathways, such as mTORC1 signaling and cholesterol metabolism, that may act upstream of L1 expression. Finally, the co-association of some genetic variants with both L1 expression differences and various age-related diseases suggests that L1 differences may precede and contribute to the onset of disease. Our results expand the potential mechanisms by which L1 expression is regulated and by which L1 may influence aging-related phenotypes.

METHODS

Publicly available data acquisition

The eQTL analysis was carried out on 358 European (EUR) individuals and 86 Yoruban (YRI) individuals for which paired single nucleotide variant, structural variant, and transcriptomic data were available from Phase 3 of the 1000 Genomes Project [22, 23] and from the GEUVADIS consortium [26]. Specifically, Phase 3 autosomal SNVs called on the GRCh38 reference genome were obtained from The International Genome Sample Resource (IGSR) FTP site (http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/data_collections/1000_genomes_project/relea se/20190312_biallelic_SNV_and_INDEL/). Structural variants were also obtained from the IGSR FTP site (http://ftp.1000genomes.ebi.ac.uk/vol1/ftp/phase3/integrated_sv_map/). mRNA-sequencing fastq files generated by the GEUVADIS consortium were obtained from ArrayExpress under accession E-GEUV-1.

Aggregating and pre-processing genotype data for eQTL analyses

To prepare SNVs for association analyses, all SNVs were first annotated with rsIDs from dbSNP build 155 using BCFtools v1.10.2 [87]. VCFtools v0.1.17 [88] was then used to remove indels and keep variants with the following properties in each of the two populations: possessed a minimum and maximum of two alleles, possessed a minor allele frequency (MAF) of at least 1%, passed Hardy-Weinberg equilibrium thresholding at $p < 1e-6$, with no missing samples, and located on an autosome. We

note here that sex chromosomes were not included in the analysis since (i) Y

chromosome SNVs were not available and (ii) analyses with X chromosome SNVs

require unique algorithms and cannot simply be incorporated into traditional association

pipelines [89, 90]. VCF files containing these filtered SNVs were then converted to

PLINK BED format using PLINK v1.90b6.17 [91], keeping the allele order. PLINK BED

files were subsequently used to generate preliminary 0/1/2 genotype matrices using the

'--recodeA' flag in PLINK. These preliminary matrices were manipulated in terminal,

using the gcut v9.0 function to remove unnecessary columns and datamash v1.7 to

transpose the data, to generate the final 0/1/2 matrices used for the eQTL analyses.

Finally, PLINK was used to prune the list of filtered SNVs, using the "--indep-pairwise 50

10 0.1" flag, and to generate principal components (PCs) from the pruned genotypes.

To control for inter-individual differences in genomic transposon copy number

load, we applied 1 of 2 approaches, depending on the analysis. For approach 1, the net

number of L1 and Alu insertions was quantified across the 444 samples. We chose to

aggregate the L1 and Alu copy numbers, since Alu relies on L1 machinery for

mobilization [92], and so the aggregate number may provide a finer view of L1-

associated copy number load. Briefly, VCFTools was used to extract autosomal

structural variants from the 1000Genomes structural variant calls. L1 and Alu insertions

and deletions were then extracted with BCFtools by keeping entries with the following

expressions: 'SVTYPE="LINE1"', 'SVTYPE="ALU"', 'SVTYPE="DEL_LINE1"', and

'SVTYPE="DEL_ALU"'. The resulting VCF files were then transformed to 0/1/2 matrices

in the same manner as the SNVs. A net copy number score was obtained for each

sample by adding the values for the L1 and Alu insertions and subtracting the values for

the L1 and Alu deletions. For approach 2, the complete structural variant matrix was

filtered with VCFtools using the same parameters as with the SNV matrices. The filtered

structural variant matrix was then pruned with PLINK, and these pruned structural

variant genotypes were used to generate principal components, in the same fashion as

with the SNV matrix. The net copy number score or the structural variant principal

components, depending on the analysis, were included as covariates.

mRNA-seq read trimming, mapping, and quantification

Fastq files were first trimmed using fastp v0.20.1 [93] with the following

parameters: detect_adapter_for_pe, disable_quality_filtering, trim_front1 17, trim_front2

17, cut_front, cut_front_window_size 1, cut_front_mean_quality 20, cut_tail,

cut_tail_window_size 1, cut_tail_mean_quality 20, cut_right, cut_right_window_size 5,

cut_right_mean_quality 20, and length_required 36. Read quality was then inspected

using fastqc v0.11.9.

Next, the GRCh38 primary human genome assembly and comprehensive gene

annotation were obtained from GENCODE release 33 [94]. Since LCLs are generated

by infecting B-cells with Epstein-Barr virus, the EBV genome (GenBank ID V01555.2)

was included as an additional contig in the human reference genome. The trimmed

reads were aligned to this modified reference genome using STAR v2.7.3a [95] with the

following parameters: outFilterMultimapNmax 100, winAnchorMultimapNmax 100, and

outFilterMismatchNoverLmax 0.04. Finally, the TEcount function in the TEtranscripts

v2.1.4 [27] package was employed to obtain gene and TE counts, using the GENCODE

annotations to define gene boundaries and a repeat GTF file provided on the Hammell

lab website (downloaded on February 19 2020 from

https://labshare.cshl.edu/shares/mhammelllab/www-

data/TEtranscripts/TE_GTF/GRCh38_GENCODE_rmsk_TE.gtf.gz) to define repeat

boundaries.

Gene *cis*-eQTL and L1 t*rans*-eQTL analyses

Gene and TE count files were loaded into R v4.2.1. Lowly expressed genes were

first filtered out if 323/358 European samples and 78/86 Yoruban samples did not have

over 0.44 counts per million (cpm) or 0.43 cpm, respectively. These fractions were

selected because they corresponded to expression in ~90% of samples and thus helped

maintain maximal statistical power by focusing on genes ubiquitously expressed across

each entire population. The cpm thresholds were selected because they corresponded

to 10 reads in the median-length library within each set of samples.

Then, counts underwent a variance stabilizing transformation (vst) using DESeq2

v1.36.0 [96]. The following covariates were regressed out from vst normalized

expression data using the 'removeBatchEffect' function in Limma v3.52.2 [97]: lab,

population category, principal components 1-2 of the pruned SNVs, biological sex, net

L1/Alu copy number, and EBV expression levels. Since the Yoruban samples were all

from the same population, the population variable was omitted in their batch correction.

Here, we note several things. First, EBV expression was included as a covariate

because heightened TE expression is often a feature of viral infections [98]. Secondly,

although PEER [68] is often used to remove technical variation for *cis*-eQTL analysis,

this can come at the expense of correcting out genome-wide biological effects. This can

be problematic in some settings, such as *trans*-eQTL analysis. Thus, PEER factors

were not included. The batch-corrected data underwent a final inverse normal

transformation (INT), using the RankNorm function in the R package RNOmni v1.0.1, to

obtain normally distributed gene expression values.

The INT expression matrices were split into genes and L1 subfamilies, which

were used to identify gene *cis*-eQTLs and L1 subfamily *trans*-eQTLs in the European

superpopulation using MatrixEQTL v2.3 [99]. For gene *cis*-eQTLs, SNVs were tested for

association with expressed genes within 1 million base pairs. We opted to use a *trans*-

eQTL approach using aggregate subfamily-level TE expression since the *trans*

approach should allow us to identify regulators of many elements rather than one. The

Benjamini-Hochberg false discovery rate (FDR) was calculated in each analysis, and we

used the p-value corresponding to an FDR of < 5% as the threshold for eQTL

significance. In addition, the *cis*-eQTL and *trans*-eQTL analyses were also repeated

using 20 permuted expression datasets in which the sample names were scrambled,

and the p-value corresponding to an average empirical FDR of < 5% was used as a

secondary threshold. To note, we calculated the average empirical FDR at a given p-

value p_i by (i) counting the total number of null points with $p \leq p_i$, (ii) dividing by the

number of permutations, to obtain an average number of null points with $p \leq p_i$, and (iii)

dividing the average number of null points with $p \leq p_i$ by the number of real points with p

$\leq p_i$. eQTLs were called as significant if they passed the stricter of the two thresholds.

SNV-gene and SNV-L1 associations that were significant in the European

superpopulation were then targeted and tested in the Yoruban population using R's

built-in linear modelling functions. In this case, only the Benjamini-Hochberg FDR was

calculated, and significant eQTLs were called if they possessed an FDR < 5%.

Defining SNV-gene-L1 trios and mediation analysis

For each population, the *cis-* and *trans*-eQTL results were integrated to identify

SNVs associated with both gene and L1 subfamily expression. We reasoned that L1

expression would respond to differences in expression of *bona fide* regulators.

Consequently, gene expression and L1 subfamily expression associations were

assessed by linear regression, and the p-values from this analysis were Benjamini-

Hochberg FDR-corrected. Candidate SNV-gene-L1 trios were defined as those with *cis-*

eQTL, *trans*-eQTL, and expression regression FDRs < 5%. To identify top, index SNVs

in regions of linkage disequilibrium (LD), SNVs within 500 kilobases of each other with

an R^2 > 0.10 were clumped together by *trans*-eQTL p-value using PLINK v1.90b6.17.

Mediation analysis was carried out using the 'gmap.gpd' function in eQTLMAPT v0.1.0

[100] on all candidate SNV-gene-L1 trios. Empirical p-values were calculated using

30,000 permutations, and Benjamini-Hochberg FDR values were calculated from

empirical p-values. Mediation effects were considered significant for trios with FDR <

5%.

Differential expression analysis across *trans*-eQTL SNV genotypes

Transcriptomic changes associated with alternating the allele of each SNV of

interest were evaluated using DESeq2 v1.36.0. Using the same filtered counts prepared

for the eQTL analysis, a linear model was constructed with the following covariates for

each SNV: SNV genotype in 0/1/2 format, biological sex, lab, population category,

principal components 1-2 of the pruned SNVs, and principal components 1-3 of the

pruned SVs (to account for structural variant population structure). As before, the

population label was omitted from the Yoruban population analysis. Significant genes

and TEs were those with an FDR < 5%.

Functional enrichment analyses

We used the Gene Set Enrichment Analysis (GSEA) paradigm as implemented

in the R package clusterProfiler v4.4.4 [101]. Gene Ontology, Reactome, and Hallmark

pathway gene sets were obtained from the R package msigdbr v7.5.1, an Ensembl ID-

mapped collection of gene sets from the Molecular Signature Database [29, 30].

Additionally, TE subfamilies were aggregated into TE family gene sets using the TE

family designations specified in the TE GTF file (downloaded on February 19 2020 from

https://labshare.cshl.edu/shares/mhammelllab/www-

data/TEtranscripts/TE_GTF/GRCh38_GENCODE_rmsk_TE.gtf.gz) used during the

RNA-seq quantification step. The DESeq2 v1.36.0 Wald-statistic was used to generate

a combined ranked list of genes and TEs for functional enrichment analysis. All gene

sets with an FDR < 5% were considered significant. For plots with a single analysis, the

top 5 downregulated and top 5 upregulated gene sets were plotted, at most. For plots

with multiple analyses, shared gene sets with the desired expression patterns in each

individual analysis were first identified. Then, the p-values for shared gene sets were

combined using Fisher's method, and this meta-analysis p-value was used to rank

shared gene sets. Finally, the top 5 gene sets with one expression pattern and the top 5

gene sets with the opposite expression pattern were plotted. If there were less than 5

gene sets in either group, those were replaced with gene sets exhibiting the opposite

regulation, in order to plot 10 shared gene sets whenever possible.

Cell lines and cell culture conditions.

GM12878 (RRID: CVCL_7526) lymphoblastoid cells were purchased from the

Coriell Institute. We opted to use GM12878 as a well-characterized representative cell

line for candidate validation, given that (i) it is of the same cell type as the transcriptomic

data used here for our eQTL analysis, and (ii) its epigenomic landscape and culture

conditions are well-characterized as part of the ENCODE project [39, 40].

GM12878 cells were maintained in RPMI (Corning cat. 15-040-CV) containing

15% FBS and 1X Penicillin-Streptomycin-Glutamine (Corning cat. 30-009-CI). Cells

were cultured in a humidified incubator at 37°C and 5% CO_2, subculturing cells 1:5 once

cells reached a density of ~10^6 mL^{-1}. All cells used were maintained below passage 30

and routinely tested for mycoplasma contamination using the PlasmoTest Mycoplasma

Detection Kit (InvivoGen).

Plasmids

The empty pcDNA3.1(+) backbone (Invitrogen cat. V79020) was a kind gift from

the lab of Dr. Changhan David Lee at the University of Southern California Leonard

Davis School of Gerontology. Overexpression vectors for *IL16* (CloneID OHu48263C),

STARD5-FLAG (CloneID OHu07617D), *HSD17B12*-FLAG (CloneID OHu29918D), and

RNF5-FLAG (CloneID OHu14875D) on a pcDNA3.1 backbone were purchased from

GenScript. Plasmid sequences were verified for accuracy using Plasmidsaurus's whole

plasmid sequencing service.

Transfections

Escherichia coli were cultured in LB Broth (ThermoFischer Scientific)

supplemented with 50 μg/mL carbenicillin to an optical density 600 (OD_{600}) of 2 − 4.

Plasmid extractions were carried out using the Nucleobond Xtra Midi Plus EF kit

(Macherey-Nagel) following manufacturer recommendations. Plasmids were aliquoted

and stored at -20°C until the time of transfection. On the day of transfection, GM12878

cells were collected in conical tubes, spun down (100xG, 5 minutes, room temperature),

resuspended in fresh media, and counted by trypan blue staining using a Countess II FL

automated cell counter (Thermo Fisher). The number of cells necessary for the

experiment were then aliquoted, spun down, and washed with Dulbecco's phosphate-

buffered saline (DPBS)(Corning, cat. #21-031-CV).

GM12878 cells were transfected by electroporation using the Neon Transfection

System (Invitrogen) with the following parameters: 1200 V, 20 ms, and 3 pulses for

GM12878 cells in Buffer R. Per reaction, we maintained a plasmid mass:cell number

ratio of 10 µg : $2*10^6$ cells. For mRNA-sequencing, $8*10^6$ GM12878 cells were

independently transfected for each biological replicate, with 4 replicates per

overexpression condition, and cultured in a T25 flask. Immediately after transfection,

cells were cultured in Penicillin-Streptomycin-free media for ~24 hours.

Afterwards, to promote selection of viable and healthy transfected GM12878

cells, we enriched for viable cells using the EasySep Dead Cell Removal (Annexin V)

Kit (STEMCELL Technologies) before seeding $2*10^6$ live cells in the same media used

for cell maintenance. After another 24 hours, cell viability was measured by trypan blue

staining on a Countess automated cell counter and cells were spun down (100xG, 5

min, room temperature) and lysed in TRIzol Reagent (Invitrogen) for downstream total

RNA isolation (see below).

Recombinant human IL16 (rhIL16) peptide treatment

Human rIL16 was obtained from PeproTech (cat. #200-16) and resuspended in

0.1% bovine serum albumin (BSA) solution (Akron, cat. #AK8917-0100). GM12878 cells

were seeded at a concentration of 500,000 live cells per mL of media on 6-well

suspension plates with 3 independent replicates per condition. Cells were exposed to 0,

24, or 48 hours of 100 ng mL^{-1} of rhIL16. To replace or exchange media 24 hours after

seeding, cells were transferred to conical tubes, spun down (100xG, 5 min, room

temperature), resuspended in 5 mL of the appropriate media, and transferred back to 6-

well suspension plates. After 48 hours, cell viability was measured by trypan blue

staining and cells were spun down (100xG, 5 min, room temperature) and lysed in

TRIzol Reagent (Invitrogen).

RNA extractions and mRNA sequencing

RNA was extracted using the Direct-zol RNA Miniprep kit (Zymo Research)

following manufacturer recommendations. The integrity of RNA samples was evaluated

using an Agilent High Sensitivity RNA ScreenTape assay (Agilent Technologies),

ensuring that all samples had a minimum eRIN score of 8 before downstream

processing. We then submitted total RNA samples to Novogene (Sacramento,

California) for mRNA library preparation and sequencing on the NovaSeq 6000 platform

as paired-end 150 bp reads.

Analysis of overexpression and rhIL16 exposure mRNA-seq

mRNA-seq reads were trimmed, mapped, and quantified like for the eQTL

analysis, except for the overexpression sample data. For this data, one modification

was made: the EBV-inclusive reference genome was further modified to include the

pcDNA3.1 sequence as an additional contig. Lowly expressed genes were filtered using

a cpm threshold as in the eQTL processing, but that cpm threshold had to be satisfied

by as many samples as the size of the smallest biological group. For the overexpression

data, surrogate variables were estimated with the 'svaseq' function [69] in the R

package 'sva' v3.44.9, and they were regressed out from the raw read counts using the

'removeBatchEffect' function in the R package Limma v3.52.2. DESeq2 was used to

identify significantly (FDR < 5%) differentially expressed genes and TEs between

groups. Functional enrichment analysis was carried out as previously described.

PheWAS analysis

To gather the known associated traits for the 499 TE-related SNVs, we used

Open Targets Genetics (https://genetics.opentargets.org/), a database of GWAS

summary statistics [102]. First, we queried the database using the 499 TE-related SNVs

and collected traits that were directly associated (with $P < 5\times10^{-8}$) with the SNVs, as well

as traits associated with lead variants that were in linkage disequilibrium (LD) with the

queried SNPs (with $R^2 > 0.6$). For age-related traits (ARTs), we used the

comprehensive list of 365 Medical Subject Headings (MeSH) terms reported by [103]

(downloaded from https://github.com/kisudsoe/Age-related-traits). To identify known

age-related traits, the known associated traits were translated into the equivalent MeSH

terms using the method described by [103]. Then, the MeSH-translated known

associated traits for the 499 TE-related SNVs were filtered by the MeSH terms for age-

related traits.

As a parallel approach, we mapped the RsIDs for all SNVs used during the eQTL

analyses to their corresponding bi-allelic Open Targets variant IDs, when available. The

variant IDs corresponding to L1 *trans*-eQTL SNVs were extracted, and 500 different

equal-length combinations of random SNVs were generated. Next, we queried the Open

Targets database using the lists of L1-associated and random SNVs and collected the

associated traits (with $P < 5 \times 10^{-8}$). Importantly, the database assigns traits to broader

categories, including 14 disease categories that we considered age-related. We

counted the number of L1-associated or random SNVs mapping to each category, and

we used the random SNV counts to generate an empirical cumulative distribution

function (ecdf) for each category. We calculated enrichment p-values using the formula

p = 1- ecdf(mapped eQTLs) and then Benjamini-Hochberg FDR-corrected all p-values.

An enrichment score (ES) was also calculated for each category using the formula ES =

number of mapped L1 eQTLs / median number of randomly mapping SNVs. Categories

with an ES > 1 and FDR < 0.05 were considered significantly enriched.

Quantification of mouse serum IL16 by ELISA

Serum was collected from male and female C57BL/6JNia mice (4-6 and 20-24

months old) obtained from the National Institute on Aging (NIA) colony at Charles

Rivers. All animals were euthanized between 8-11 am in a "snaking order" across all

groups to minimize batch-processing confounds due to circadian processes. All animals

were euthanized by CO_2 asphyxiation followed by cervical dislocation. Circulating IL16

levels were quantitatively evaluated from mouse serum by enzyme-linked

immunosorbent assay (ELISA). Serum was diluted 1/10 before quantifying IL16

concentrations using Abcam's Mouse IL-16 ELISA Kit (ab201282) in accordance with

manufacturer instructions. Technical replicates from the same sample were averaged to

one value before statistical analysis and plotting. P-values across age within each sex

were calculated using a non-parametric 2-sided Wilcoxon test, and p-values from each

sex-specific analysis were combined using Fisher's method.

DECLARATIONS

Ethics approval and consent to participate

All animals were treated and housed in accordance with the Guide for Care and Use of Laboratory Animals. All experimental procedures were approved by the University of Southern California's Institutional Animal Care and Use Committee (IACUC) and are in accordance with institutional and national guidelines. Samples were derived from animals on approved IACUC protocol #20770.

Availability of data and materials

Sequencing data generated in this study is accessible through the Sequence Read Archive (SRA) under BioProject PRJNA937306. All code is available on the Benayoun lab GitHub (https://github.com/BenayounLaboratory/TE-eQTL_LCLs). Analyses were conducted using R version 4.2.1. Code was re-run independently on R version 4.3.0 to check for reproducibility.

Competing interests

The authors declare that they have no competing interests.

Funding

This material is based upon work supported by the National Science Foundation Graduate Research Fellowship Program (NSF GRFP) under Grant No. DGE-1842487 (to J.I.B.), the National Institute on Aging (NIA) under Grant No. T32 AG052374 (to

1015   J.I.B.), the University of Southern California with a Provost Fellowship (to J.I.B.), the

1016   USC Gerontology Enriching MSTEM (GEMSTEM) to Enhance Diversity in Aging

1017   Program under Grant No. R25 AG076400 (to C.R.M.), and the National Institute of

1018   General Medical Sciences (NIGMS) under Grant No. R35 GM142395 (to B.A.B).

1019

1020

Authors' contributions

1022   J.I.B. and B.A.B designed the study. J.I.B., L.Z., and S.K. performed data

1023   analyses, with guidance from Y.S. and B.A.B. J.I.B. and C.R.M. carried out

1024   experiments. J.I.B., B.A.B., S.K., and Y.S. wrote the manuscript. All authors contributed

1025   to the editing of the manuscript.

1026

1027

Acknowledgements

1029   We would like to thank Prof. Rachel Brem for her feedback and insights on the

1030   eQTL analyses. We would also like to thank Dr. Minhoo Kim for her feedback on the

1031   manuscript. Some panels were created with BioRender.com.

1032

1033

1034

REFERENCES

1035

1. Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, Dewar K,

Doyle M, FitzHugh W, et al: **Initial sequencing and analysis of the human genome.**

Nature 2001, **409:**860-921.

2. Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, Sutton GG, Smith HO, Yandell M,

Evans CA, Holt RA, et al: **The Sequence of the Human Genome.** *Science* 2001,

291:1304-1351.

3. Brouha B, Schustak J, Badge RM, Lutz-Prigge S, Farley AH, Moran JV, Kazazian HH:

Hot L1s account for the bulk of retrotransposition in the human population.

Proceedings of the National Academy of Sciences 2003, **100:**5280-5285.

4. Moran JV, Holmes SE, Naas TP, DeBerardinis RJ, Boeke JD, Kazazian HH, Jr.: **High**

Frequency Retrotransposition in Cultured Mammalian Cells. *Cell* 1996, **87:**917-927.

5. Bravo JI, Nozownik S, Danthi PS, Benayoun BA: **Transposable elements, circular**

RNAs and mitochondrial transcription in age-related genomic regulation.

Development 2020, **147**.

6. Della Valle F, Reddy P, Yamamoto M, Liu P, Saera-Vila A, Bensaddek D, Zhang H,

Prieto Martinez J, Abassi L, Celii M, et al: *LINE-1* **RNA causes heterochromatin**

erosion and is a target for amelioration of senescent phenotypes in progeroid

syndromes. *Science Translational Medicine* 2022, **14:**eabl6057.

7. Simon M, Van Meter M, Ablaeva J, Ke Z, Gonzalez RS, Taguchi T, De Cecco M,

Leonova KI, Kogan V, Helfand SL, et al: **LINE1 Derepression in Aged Wild-Type and**

SIRT6-Deficient Mice Drives Inflammation. *Cell Metabolism* 2019, **29:**871-885.e875.

8. Zhao Y, Oreskovic E, Zhang Q, Lu Q, Gilman A, Lin YS, He J, Zheng Z, Lu JY, Lee J, et

al: **Transposon-triggered innate immune response confers cancer resistance to**

the blind mole rat. *Nature Immunology* 2021, **22:**1219-1230.

9.   Zhao H, Ji Q, Wu Z, Wang S, Ren J, Yan K, Wang Z, Hu J, Chu Q, Hu H, et al:

Destabilizing heterochromatin by APOE mediates senescence. *Nature Aging* 2022,

2:303-316.

10.   Liu EY, Russ J, Cali CP, Phan JM, Amlie-Wolf A, Lee EB: **Loss of Nuclear TDP-43 Is**

Associated with Decondensation of LINE Retrotransposons. *Cell Reports* 2019,

27:1409-1421.e1406.

11.   De Cecco M, Criscione SW, Peterson AL, Neretti N, Sedivy JM, Kreiling JA:

Transposable elements become active and mobile in the genomes of aging

mammalian somatic tissues. *Aging (Albany NY)* 2013, **5:**867-883.

12.   De Cecco M, Criscione SW, Peckham EJ, Hillenmeyer S, Hamm EA, Manivannan J,

Peterson AL, Kreiling JA, Neretti N, Sedivy JM: **Genomes of replicatively senescent**

cells undergo global epigenetic changes leading to gene silencing and activation

of transposable elements. *Aging Cell* 2013, **12:**247-256.

13.   Campisi J: **Aging, Cellular Senescence, and Cancer.** *Annual Review of Physiology*

2013, **75:**685-705.

14.   Franceschi C, Garagnani P, Parini P, Giuliani C, Santoro A: **Inflammaging: a new**

immune–metabolic viewpoint for age-related diseases. *Nature Reviews*

Endocrinology 2018, **14:**576-590.

15.   De Cecco M, Ito T, Petrashen AP, Elias AE, Skvir NJ, Criscione SW, Caligiana A,

Brocculi G, Adney EM, Boeke JD, et al: **L1 drives IFN in senescent cells and**

promotes age-associated inflammation. *Nature* 2019, **566:**73-78.

16.   Rodriguez-Martin B, Alvarez EG, Baez-Ortega A, Zamora J, Supek F, Demeulemeester

J, Santamarina M, Ju YS, Temes J, Garcia-Souto D, et al: **Pan-cancer analysis of**

whole genomes identifies driver rearrangements promoted by LINE-1

retrotransposition. *Nature Genetics* 2020, **52:**306-319.

17.   Flasch DA, Chen X, Ju B, Li X, Dalton J, Mulder HL, Easton J, Wang L, Baker SJ,

Chiang J, Zhang J: **Somatic LINE-1 promoter acquisition drives oncogenic FOXR2**

activation in pediatric brain tumor. *Acta Neuropathologica* 2022, **143:**605-607.

18.   Levin HL, Moran JV: **Dynamic interactions between transposable elements and**

their hosts. *Nature Reviews Genetics* 2011, **12:**615-627.

19.   Rebollo R, Romanish MT, Mager DL: **Transposable Elements: An Abundant and**

Natural Source of Regulatory Sequences for Host Genes. *Annual Review of*

Genetics 2012, **46:**21-42.

20.   Liu N, Lee CH, Swigut T, Grow E, Gu B, Bassik MC, Wysocka J: **Selective silencing of**

euchromatic L1s revealed by genome-wide screens for L1 regulators. *Nature* 2018,

553:228-232.

21.   Chung N, Jonaid GM, Quinton S, Ross A, Sexton CE, Alberto A, Clymer C, Churchill D,

Navarro Leija O, Han MV: **Transcriptome analyses of tumor-adjacent somatic**

tissues reveal genes co-expressed with transposable elements. *Mobile DNA* 2019,

10:39.

22.   Auton A, Abecasis GR, Altshuler DM, Durbin RM, Abecasis GR, Bentley DR, Chakravarti

A, Clark AG, Donnelly P, Eichler EE, et al: **A global reference for human genetic**

variation. *Nature* 2015, **526:**68-74.

23.   Sudmant PH, Rausch T, Gardner EJ, Handsaker RE, Abyzov A, Huddleston J, Zhang Y,

Ye K, Jun G, Hsi-Yang Fritz M, et al: **An integrated map of structural variation in**

2,504 human genomes. *Nature* 2015, **526:**75-81.

24.   Sie L, Loong S, Tan EK: **Utility of lymphoblastoid cell lines.** *Journal of Neuroscience*

Research 2009, **87:**1953-1959.

25.   Hussain T, Mulherkar R: **Lymphoblastoid Cell lines: a Continuous in Vitro Source of**

Cells to Study Carcinogen Sensitivity and DNA Repair. *Int J Mol Cell Med* 2012,

1:75-87.

26.   Lappalainen T, Sammeth M, Friedländer MR, 't Hoen PAC, Monlong J, Rivas MA,

Gonzàlez-Porta M, Kurbatova N, Griebel T, Ferreira PG, et al: **Transcriptome and**

genome sequencing uncovers functional variation in humans. *Nature* 2013,

501:506-511.

27.   Jin Y, Tam OH, Paniagua E, Hammell M: **TEtranscripts: a package for including**

transposable elements in differential expression analysis of RNA-seq datasets.

Bioinformatics 2015, **31:**3593-3599.

28.   Williams TM: **Human Leukocyte Antigen Gene Polymorphism and the**

Histocompatibility Laboratory. *The Journal of Molecular Diagnostics* 2001, **3:**98-104.

29.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich

A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP: **Gene set enrichment analysis: A**

knowledge-based approach for interpreting genome-wide expression profiles.

Proceedings of the National Academy of Sciences 2005, **102:**15545-15550.

30.   Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P: **The**

Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst*

2015, **1:**417-425.

31.   Marasca F, Sinha S, Vadalà R, Polimeni B, Ranzani V, Paraboschi EM, Burattin FV,

Ghilotti M, Crosti M, Negri ML, et al: **LINE1 are spliced in non-canonical transcript**

variants to regulate T cell quiescence and exhaustion. *Nature Genetics* 2022,

54:180-193.

32.   Wahl D, Cavalier AN, Smith M, Seals DR, LaRocca TJ: **Healthy Aging Interventions**

Reduce Repetitive Element Transcripts. *The Journals of Gerontology: Series A* 2020,

76:805-810.

33.   Sampathkumar NK, Bravo JI, Chen Y, Danthi PS, Donahue EK, Lai RW, Lu R, Randall

LT, Vinson N, Benayoun BA: **Widespread sex dimorphism in aging and age-related**

diseases. *Hum Genet* 2020, **139:**333-356.

34.   Baeken MW, Moosmann B, Hajieva P: **Retrotransposon activation by distressed**
mitochondria in neurons. *Biochemical and Biophysical Research Communications*
2020, **525:**570-575.

35.   Valdebenito-Maturana B, Valdebenito-Maturana F, Carrasco M, Tapia JC, Maureira A:
Activation of Transposable Elements in Human Skeletal Muscle Fibers upon Statin
Treatment. *International Journal of Molecular Sciences* 2023, **24:**244.

36.   Tiwari B, Jones AE, Caillet CJ, Das S, Royer SK, Abrams JM: **p53 directly represses**
human LINE1 transposons. *Genes & Development* 2020, **34:**1439-1451.

37.   Ardeljan D, Steranka JP, Liu C, Li Z, Taylor MS, Taylor MS, Payer LM, Gorbounov M,
Sarnecki JS, Deshpande V, et al: **Cell fitness screens reveal a conflict between**
LINE-1 retrotransposition and DNA replication. *Nature structural & molecular*
biology 2020, **27:**168-178.

38.   Luqman-Fatah A, Watanabe Y, Uno K, Ishikawa F, Moran JV, Miyoshi T: **The interferon**
stimulated gene-encoded protein HELZ2 inhibits human LINE-1 retrotransposition
and LINE-1 RNA-mediated type I interferon induction. *Nature Communications* 2023,
14:203.

39.   The EPC: **A User's Guide to the Encyclopedia of DNA Elements (ENCODE).** *PLOS*
Biology 2011, **9:**e1001046.

40.   **An integrated encyclopedia of DNA elements in the human genome.** *Nature* 2012,
489:57-74.

41.   Center DM, Cruikshank W: **Modulation of lymphocyte migration by human**
lymphokines. I. Identification and characterization of chemoattractant activity for
lymphocytes from mitogen-stimulated mononuclear cells. *J Immunol* 1982,
128:2563-2568.

42.   Cruikshank WW, Center DM, Nisar N, Wu M, Natke B, Theodore AC, Kornfeld H:
Molecular and functional analysis of a lymphocyte chemoattractant factor:

association of biologic function with CD4 expression. *Proc Natl Acad Sci U S A* 1994, **91:**5109-5113.

43. Theodore AC, Center DM, Nicoll J, Fine G, Kornfeld H, Cruikshank WW: **CD4 ligand IL-16 inhibits the mixed lymphocyte reaction.** *J Immunol* 1996, **157:**1958-1964.

44. Cruikshank WW, Lim K, Theodore AC, Cook J, Fine G, Weller PF, Center DM: **IL-16 inhibition of CD3-dependent lymphocyte activation and proliferation.** *J Immunol* 1996, **157:**5240-5248.

45. Center DM, Kornfeld H, Cruikshank WW: **Interleukin 16 and its function as a CD4 ligand.** *Immunol Today* 1996, **17:**476-481.

46. Wilson KC, Center DM, Cruikshank WW: **Mini ReviewThe Effect of Interleukin-16 and its Precursor on T Lymphocyte Activation and Growth.** *Growth Factors* 2004, **22:**97-104.

47. Huang Y, Du KL, Guo PY, Zhao RM, Wang B, Zhao XL, Zhang CQ: **IL-16 regulates macrophage polarization as a target gene of mir-145-3p.** *Mol Immunol* 2019, **107:**1-9.

48. Rodriguez-Agudo D, Malacrida L, Kakiyama G, Sparrer T, Fortes C, Maceyka M, Subler MA, Windle JJ, Gratton E, Pandak WM, Gil G: **StarD5: an ER stress protein regulates plasma membrane and intracellular cholesterol homeostasis.** *Journal of Lipid Research* 2019, **60:**1087-1098.

49. Rodriguez-Agudo D, Calderon-Dominguez M, Medina MA, Ren S, Gil G, Pandak WM: **ER stress increases StarD5 expression by stabilizing its mRNA and leads to relocalization of its protein from the nucleus to the membranes.** *J Lipid Res* 2012, **53:**2708-2715.

50. Soccio RE, Adams RM, Maxwell KN, Breslow JL: **Differential Gene Regulation of StarD4 and StarD5 Cholesterol Transfer Proteins: ACTIVATION OF StarD4 BY STEROL REGULATORY ELEMENT-BINDING PROTEIN-2 AND StarD5 BY**

ENDOPLASMIC RETICULUM STRESS *. *Journal of Biological Chemistry* 2005,

280:19410-19418.

51.   Luu-The V, Tremblay P, Labrie F: **Characterization of type 12 17beta-hydroxysteroid**

dehydrogenase, an isoform of type 3 17beta-hydroxysteroid dehydrogenase

responsible for estradiol formation in women. *Mol Endocrinol* 2006, **20:**437-443.

52.   Heikelä H, Ruohonen ST, Adam M, Viitanen R, Liljenbäck H, Eskola O, Gabriel M,

Mairinoja L, Pessia A, Velagapudi V, et al: **Hydroxysteroid (17β) dehydrogenase 12 is**

essential for metabolic homeostasis in adult mice. *Am J Physiol Endocrinol Metab*

2020, **319:**E494-e508.

53.   Nagasaki S, Miki Y, Akahira J, Suzuki T, Sasano H: **Transcriptional regulation of**

17beta-hydroxysteroid dehydrogenase type 12 by SREBP-1. *Mol Cell Endocrinol*

2009, **307:**163-168.

54.   Didier C, Broday L, Bhoumik A, Israeli S, Takahashi S, Nakayama K, Thomas SM,

Turner CE, Henderson S, Sabe H, Ronai Ze: **RNF5, a RING Finger Protein That**

Regulates Cell Motility by Targeting Paxillin Ubiquitination and Altered

Localization. *Molecular and Cellular Biology* 2003, **23:**5331-5345.

55.   Tcherpakov M, Delaunay A, Toth J, Kadoya T, Petroski MD, Ronai ZeA: **Regulation of**

Endoplasmic Reticulum-associated Degradation by RNF5-dependent

Ubiquitination of JNK-associated Membrane Protein (JAMP) *. *Journal of Biological*

Chemistry 2009, **284:**12099-12109.

56.   Zhong B, Zhang Y, Tan B, Liu TT, Wang YY, Shu HB: **The E3 ubiquitin ligase RNF5**

targets virus-induced signaling adaptor for ubiquitination and degradation. *J*

Immunol 2010, **184:**6249-6255.

57.   Zhong B, Zhang L, Lei C, Li Y, Mao A-P, Yang Y, Wang Y-Y, Zhang X-L, Shu H-B: **The**

Ubiquitin Ligase RNF5 Regulates Antiviral Responses by Mediating Degradation

of the Adaptor Protein MITA. *Immunity* 2009, **30:**397-407.

58.   Lai RW, Lu R, Danthi PS, Bravo JI, Goumba A, Sampathkumar NK, Benayoun BA:

Multi-level remodeling of transcriptional landscapes in aging and longevity. *BMB*

Rep 2019, **52:**86-108.

59.   Sato S, Gillette M, de Santiago PR, Kuhn E, Burgess M, Doucette K, Feng Y, Mendez-

Dorantes C, Ippoliti PJ, Hobday S, et al: **LINE-1 ORF1p as a candidate biomarker in**

high grade serous ovarian carcinoma. *Scientific Reports* 2023, **13:**1537.

60.   Rodić N, Sharma R, Sharma R, Zampella J, Dai L, Taylor MS, Hruban RH, Iacobuzio-

Donahue CA, Maitra A, Torbenson MS, et al: **Long Interspersed Element-1 Protein**

Expression Is a Hallmark of Many Human Cancers. *The American Journal of*

Pathology 2014, **184:**1280-1286.

61.   Lindholm HT, Chen R, De Carvalho DD: **Endogenous retroelements as alarms for**

disruptions to cellular homeostasis. *Trends in Cancer* 2023, **9:**55-68.

62.   Jeck WR, Siebold AP, Sharpless NE: **Review: a meta-analysis of GWAS and age-**

associated diseases. *Aging Cell* 2012, **11:**727-731.

63.   Ramirez J, Bitterman P, Basu S, Barua A: **Changes in IL-16 Expression in the Ovary**

during Aging and Its Potential Consequences to Ovarian Pathology. *J Immunol Res*

2022, **2022:**2870389.

64.   Lonsdale J, Thomas J, Salvatore M, Phillips R, Lo E, Shad S, Hasz R, Walters G, Garcia

F, Young N, et al: **The Genotype-Tissue Expression (GTEx) project.** *Nature Genetics*

2013, **45:**580-585.

65.   Carithers LJ, Ardlie K, Barcus M, Branton PA, Britton A, Buia SA, Compton CC, DeLuca

DS, Peter-Demchok J, Gelfand ET, et al: **A Novel Approach to High-Quality**

Postmortem Tissue Procurement: The GTEx Project. *Biopreservation and*

Biobanking 2015, **13:**311-319.

66.   Streeter I, Harrison PW, Faulconbridge A, The HipSci Consortium, Flicek P, Parkinson

H, Clarke L: **The human-induced pluripotent stem cell initiative—data resources for**

cellular genetics. *Nucleic Acids Research* 2016, **45:**D691-D697.

67.   Chick JM, Munger SC, Simecek P, Huttlin EL, Choi K, Gatti DM, Raghupathy N,

Svenson KL, Churchill GA, Gygi SP: **Defining the consequences of genetic variation**

on a proteome-wide scale. *Nature* 2016, **534:**500-505.

68.   Stegle O, Parts L, Piipari M, Winn J, Durbin R: **Using probabilistic estimation of**

expression residuals (PEER) to obtain increased power and interpretability of

gene expression analyses. *Nat Protoc* 2012, **7:**500-507.

69.   Leek JT: **svaseq: removing batch effects and other unwanted noise from**

sequencing data. *Nucleic Acids Research* 2014, **42:**e161-e161.

70.   Savytska N, Heutink P, Bansal V: **Transcription start site signal profiling improves**

transposable element RNA expression analysis at locus-level. *Frontiers in Genetics*

2022, **13**.

71.   Lanciano S, Cristofari G: **Measuring and interpreting transposable element**

expression. *Nature Reviews Genetics* 2020, **21:**721-736.

72.   Zeng TB, Pierce N, Liao J, Szabó PE: **H3K9 methyltransferase EHMT2/G9a controls**

ERVK-driven noncanonical imprinted genes. *Epigenomics* 2021, **13:**1299-1314.

73.   Jiang Q, Ang JYJ, Lee AY, Cao Q, Li KY, Yip KY, Leung DCY: **G9a Plays Distinct**

Roles in Maintaining DNA Methylation, Retrotransposon Silencing, and Chromatin

Looping. *Cell Reports* 2020, **33:**108315.

74.   Rantakari P, Lagerbohm H, Kaimainen M, Suomela J-P, Strauss L, Sainio K, Pakarinen

P, Poutanen M: **Hydroxysteroid (17β) Dehydrogenase 12 Is Essential for Mouse**

Organogenesis and Embryonic Survival. *Endocrinology* 2010, **151:**1893-1901.

75.   Garcia G, Zhang H, Moreno S, Tsui CK, Webster BM, Higuchi-Sanabria R, Dillin A:
Lipid homeostasis is essential for a maximal ER stress response. *eLife* 2023,
12:e83884.

76.   Baier M, Werner A, Bannert N, Metzner K, Kurth R: **HIV suppression by interleukin-**
16. *Nature* 1995, **378:**563.

77.   Zhou P, Goldstein S, Devadas K, Tewari D, Notkins AL: **Human CD4+ cells**
transfected with IL-16 cDNA are resistant to HIV-1 infection: inhibition of mRNA
expression. *Nat Med* 1997, **3:**659-664.

78.   Idziorek T, Khalife J, Billaut-Mulot O, Hermann E, Aumercier M, Mouton Y, Capron A,
Bahr GM: **Recombinant human IL-16 inhibits HIV-1 replication and protects against**
activation-induced cell death (AICD). *Clin Exp Immunol* 1998, **112:**84-91.

79.   Jia R, Jiang C, Li L, Huang C, Lu L, Xu M, Xu J, Liang X: **Interleukin 16 Enhances the**
Host Susceptibility to Influenza A Virus Infection. *Frontiers in Microbiology* 2021, **12**.

80.   Liu S, Lei Z, Li J, Wang L, Jia R, Liu Z, Jiang C, Gao Y, Liu M, Kuang L, et al:
Interleukin 16 contributes to gammaherpesvirus pathogenesis by inhibiting viral
reactivation. *PLoS Pathog* 2020, **16:**e1008701.

81.   Watanabe T, Watanabe S, Kawaoka Y: **Cellular Networks Involved in the Influenza**
Virus Life Cycle. *Cell Host & Microbe* 2010, **7:**427-439.

82.   Germain MA, Chatel-Chaix L, Gagné B, Bonneil É, Thibault P, Pradezynski F, de
Chassey B, Meyniel-Schicklin L, Lotteau V, Baril M, Lamarre D: **Elucidating novel**
hepatitis C virus-host interactions using combined mass spectrometry and
functional genomics approaches. *Mol Cell Proteomics* 2014, **13:**184-203.

83.   Mohamed B, Mazeaud C, Baril M, Poirier D, Sow AA, Chatel-Chaix L, Titorenko V,
Lamarre D: **Very-long-chain fatty acid metabolic capacity of 17-beta-**
hydroxysteroid dehydrogenase type 12 (HSD17B12) promotes replication of
hepatitis C virus and related flaviviruses. *Scientific Reports* 2020, **10:**4040.

84.   Feng K, Min Y-Q, Sun X, Deng F, Li P, Wang H, Ning Y-J: **Interactome profiling**
1290   **reveals interaction of SARS-CoV-2 NSP13 with host factor STAT1 to suppress**
1291   **interferon signaling.** *Journal of Molecular Cell Biology* 2021, **13:**760-762.

1292   85.   Yuan Z, Hu B, Xiao H, Tan X, Li Y, Tang K, Zhang Y, Cai K, Ding B: **The E3 Ubiquitin**
1293   **Ligase RNF5 Facilitates SARS-CoV-2 Membrane Protein-Mediated Virion Release.**
1294   *mBio* 2022, **13:**e03168-03121.

1295   86.   Li Z, Hao P, Zhao Z, Gao W, Huan C, Li L, Chen X, Wang H, Jin N, Luo Z-Q, et al: **The**
1296   **E3 ligase RNF5 restricts SARS-CoV-2 replication by targeting its envelope protein**
1297   **for degradation.** *Signal Transduction and Targeted Therapy* 2023, **8:**53.

1298   87.   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane
1299   T, McCarthy SA, Davies RM, Li H: **Twelve years of SAMtools and BCFtools.**
1300   *GigaScience* 2021, **10**.

1301   88.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE,
1302   Lunter G, Marth GT, Sherry ST, et al: **The variant call format and VCFtools.**
1303   *Bioinformatics* 2011, **27:**2156-2158.

1304   89.   Gao F, Chang D, Biddanda A, Ma L, Guo Y, Zhou Z, Keinan A: **XWAS: A Software**
1305   **Toolset for Genetic Data Analysis and Association Studies of the X Chromosome.**
1306   *J Hered* 2015, **106:**666-671.

1307   90.   Keur N, Ricaño-Ponce I, Kumar V, Matzaraki V: **A systematic review of analytical**
1308   **methods used in genetic association analysis of the X-chromosome.** *Briefings in*
1309   *Bioinformatics* 2022, **23**.

1310   91.   Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar
1311   P, de Bakker PIW, Daly MJ, Sham PC: **PLINK: A Tool Set for Whole-Genome**
1312   **Association and Population-Based Linkage Analyses.** *The American Journal of*
1313   *Human Genetics* 2007, **81:**559-575.

92. Ahl V, Keller H, Schmidt S, Weichenrieder O: **Retrotransposition and Crystal Structure of an Alu RNP in the Ribosome-Stalling Conformation.** *Molecular Cell* 2015, **60:**715-727.

93. Chen S, Zhou Y, Chen Y, Gu J: **fastp: an ultra-fast all-in-one FASTQ preprocessor.** *Bioinformatics* 2018, **34:**i884-i890.

94. Frankish A, Diekhans M, Ferreira A-M, Johnson R, Jungreis I, Loveland J, Mudge JM, Sisu C, Wright J, Armstrong J, et al: **GENCODE reference annotation for the human and mouse genomes.** *Nucleic Acids Research* 2018, **47:**D766-D773.

95. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR: **STAR: ultrafast universal RNA-seq aligner.** *Bioinformatics* 2012, **29:**15-21.

96. Love MI, Huber W, Anders S: **Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.** *Genome Biology* 2014, **15:**550.

97. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK: **limma powers differential expression analyses for RNA-sequencing and microarray studies.** *Nucleic Acids Research* 2015, **43:**e47-e47.

98. Macchietto MG, Langlois RA, Shen SS: **Virus-induced transposable element expression up-regulation in human and mouse host cells.** *Life Sci Alliance* 2020, **3**.

99. Shabalin AA: **Matrix eQTL: ultra fast eQTL analysis via large matrix operations.** *Bioinformatics* 2012, **28:**1353-1358.

100. Wang T, Peng Q, Liu B, Liu X, Liu Y, Peng J, Wang Y: **eQTLMAPT: Fast and Accurate eQTL Mediation Analysis With Efficient Permutation Testing Approaches.** *Frontiers in Genetics* 2020, **10**.

101. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, Feng T, Zhou L, Tang W, Zhan L, et al: **clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.** *The Innovation* 2021, **2:**100141.

1340   102.   Ghoussaini M, Mountjoy E, Carmona M, Peat G, Schmidt EM, Hercules A, Fumis L,

1341   Miranda A, Carvalho-Silva D, Buniello A, et al: **Open Targets Genetics: systematic**

1342   **identification of trait-associated genes using large-scale genetics and functional**

1343   **genomics.** *Nucleic Acids Res* 2021, **49:**D1311-d1320.

1344   103.   Kim S-S, Hudgins AD, Gonzalez B, Milman S, Barzilai N, Vijg J, Tu Z, Suh Y: **A**

1345   **Compendium of Age-Related PheWAS and GWAS Traits for Human Genetic**

1346   **Association Studies, Their Networks and Genetic Correlations.** *Frontiers in*

1347   *Genetics* 2021, **12**.

1348

1349

1350

 FIGURE LEGENDS

 Figure 1. Overview of the pipeline developed to scan for L1 transcriptional

 regulators *in silico*.

 (A) An illustration of the samples and "omic" data used in this study. Of the 358

 European individuals, 187 were female and 171 were male. Of the 86 African

 individuals, 49 were female and 37 were male. (Note that Utah subjects are of Northern

 European ancestry, and thus part of the European cohort for analytical purposes). **(B)** A

 schematic illustrating how genetic variants, gene expression, and TE expression can be

 integrated to identify highly correlated SNV-Gene-TE trios. **(C)** A Manhattan plot for the

 L1 subfamily *trans*-eQTL analysis in the European cohort. The genes that passed our

 three-part integration approach are listed next to the most significant *trans*-eQTL SNV

 they were associated with in *cis*. The dashed line at $p = 3.44E\text{-}8$ corresponds to an

 average empirical FDR < 0.05, based on 20 random permutations. One such

 permutation is illustrated in the bottom panel. The solid line at $p = 2.31E\text{-}8$ corresponds

 to a Benjamini-Hochberg FDR < 0.05. The stricter of the two thresholds, $p = 2.31E\text{-}8$,

 was used to define significant *trans*-eQTLs. FDR: False Discovery Rate. Some panels

 were created with BioRender.com.

 Figure 2. Identification of 1[st] tier candidate L1 expression regulators in the

 European cohort.

 (A) A schematic for how 1[st] tier candidate genes were defined. In short, these were

 genes in trios with index SNVs that were at the top of their respective peak. **(B)** The

three-part integration results for three protein-coding genes—*STARD5*, *IL16*, *HSD17B12*—that we considered first tier candidates for functional, *in vitro* testing. In the left column are the *trans*-eQTLs, in the middle column are the *cis*-eQTLs, and in the right column are the linear regressions for gene expression against L1 subfamily expression. Expression values following an inverse normal transform (INT) are shown. The FDR for each analysis is listed at the top of each plot. FDR: False Discovery Rate.

Figure 3. L1 *trans*-eQTLs are associated with subtle, widespread differences in TE families and known TE-associated pathways.

(A) Scheme for functionally annotating gene-linked index SNVs by GSEA. **(B)** GSEA analysis for shared, significantly regulated TE family gene sets across genotypes for rs11635336 (*IL16/STARD5*), rs9271894 (*HLA*), and rs1061810 (*HSD17B12*). **(C)** GSEA plots for the L1 family gene set results summarized in **(B)**. For these plots, the FDR value is listed. GSEA analysis for top, shared, concomitantly regulated **(D)** MSigDB Hallmark pathway, **(E)** GO Biological Process, and **(F)** Reactome pathway gene sets across genotypes for rs11635336 (*IL16/STARD5*), rs9271894 (*HLA*), and rs1061810 (*HSD17B12*). Shared gene sets were ranked by combining p-values from each individual SNV analysis using Fisher's method. In each bubble plot, the size of the dot represents the $-\log_{10}$(FDR) and the color reflects the normalized enrichment score. FDR: False Discovery Rate.

Figure 4. Impact of *IL16* and *STARD5* overexpression on LCL gene and TE expression landscapes.

IL16 and *STARD5* overexpression induce changes consistent with their known biology,

as well as subtle but widespread upregulation of TE families. **(A)** Scheme for

experimentally validating the roles of *IL16* and *STARD5* in L1 regulation. GSEA analysis

for top, differentially regulated **(B)** GO Biological Process and **(C)** Reactome pathway

gene sets following *IL16* overexpression. GSEA analysis for top, differentially regulated

(D) GO Biological Process and **(E)** Reactome pathway gene sets following *STARD5*

overexpression. **(F)** GSEA analysis for shared, significantly regulated TE family gene

sets following *IL16* and *STARD5* overexpression. **(G)** GSEA plots for the L1 family gene

set results summarized in **(F)**. For these plots, the FDR value is listed. In each bubble

plot, the size of the dot represents the $-\log_{10}$(FDR) and the color reflects the normalized

enrichment score. FDR: False Discovery Rate. Some panels were created with

BioRender.com.

1409

Figure 5. rhIL16 treatment is sufficient to transiently upregulate an L1 family gene

set.

(A) Scheme for experimentally validating the role of rhIL16 in L1 regulation. GSEA

analysis for top, shared, concomitantly regulated **(B)** GO Biological Process and **(C)**

Reactome pathway gene sets following *IL16* overexpression and rhIL16 exposure for 24

hours. Shared gene sets were ranked by combining p-values from each individual

treatment analysis using Fisher's method. **(D)** GSEA analysis for top, differentially

regulated TE family gene sets following rhIL16 exposure for 24 hours. The GSEA plot

for the L1 family gene set result summarized in the bubble plot is also shown. For this

plot, the FDR value is listed. In each bubble plot, the size of the dot represents the -

$\log_{10}$(FDR) and the color reflects the normalized enrichment score. FDR: False

Discovery Rate. Some panels were created with BioRender.com.

1422

Figure 6. **Consistent cellular responses to *IL16* overexpression, *STARD5***

overexpression, and rhIL16 exposure for 24 hours.

IL16 overexpression, *STARD5* overexpression, and rhIL16 exposure for 24 hours are

associated with subtle but widespread differences in TE families and known TE-

associated pathways. **(A)** Scheme for assessing concordantly regulated TE family and

pathway gene sets across conditions where an L1 gene set is upregulated. GSEA

analysis for top, shared, concomitantly regulated **(B)** TE family, **(C)** MSigDB Hallmark

pathway, **(D)** GO Biological Process, and **(E)** Reactome pathway gene sets following

IL16 overexpression, *STARD5* overexpression, and rhIL16 exposure for 24 hours.

Shared gene sets were ranked by combining p-values from each individual treatment

analysis using Fisher's method. In each bubble plot, the size of the dot represents the -

$\log_{10}$(FDR) and the color reflects the normalized enrichment score. FDR: False

Discovery Rate.

1436

Figure 7. L1 *trans*-eQTLs are co-associated with aging traits in GWAS databases.

(A) Scheme for obtaining *trans*-eQTL SNV-associated aging phenotypes from the Open

Targets Genetics platform. **(B)** A pie chart representing the number of SNVs (222/499)

associated with an aging-related MeSH trait, either by PheWAS or indirectly linked to

the phenotype through a proxy lead SNP in LD with the SNV. **(C)** Histogram depicting

the distribution of number of aging MeSH traits associated with the 222/499 SNVs by

PheWAS. **(D)** Histogram depicting the distribution of number of aging MeSH traits linked

with the 222/499 SNVs through a proxy lead SNP in LD with the SNVs. **(E)** A diagram

highlighting the organ targets of the top 10 most frequently associated aging traits. **(F)**

The concentrations of circulating IL16 in aging mice of both sexes was assessed by

ELISA. Significance across age in each sex was assessed using a Wilcoxon test. The

p-values from each sex (females in pink and males in blue) were combined by meta-

analysis using Fisher's method. Any p-value < 0.05 was considered significant. Some

panels were created with BioRender.com.

1451

1452

Figure 1

A Samples and data used in this study

Lymphoblastoid Cell Lines [LCLs]

358 European [EUR]
- -86 Britain
- -92 Finland
- -91 Tuscan
- -89 Utah

86 African
- -86 Yoruba [YRI]

GEUVADIS Consortium
mRNA-sequencing

1000Genomes Project Phase 3
Structural Variants [SVs]
Single Nucleotide Variants [SNVs]

mRNA
373 EUR
89 YRI

→ structural variant calls available →

mRNA and SVs
358 EUR
87 YRI

→ single nucleotide variant calls available →

mRNA, SVs, and SNVs
358 EUR
86 YRI

B Scheme for *in silico* identification of candidate transposon-subfamily regulators

Gene expression

cis-eQTL   regression

Genome-wide
bi-allelic SNVs   *trans*-eQTL   L1 subfamily
expression

AA
AT
TT

AA   AT   TT

AA   AT   TT

C Manhattan plots for European L1 *trans*-eQTL analysis

trans-eQTLs
p = 2.31E-8

ENSG00000271725, STARD5, IL16 ←

HSD17B12, ENSG00000283375, ENSG00000283341, ENSG00000246250

HLA−DRB5, HLA−DQA2 ←

ENSG00000263756 ←   → RNF5
FKBPL ←   → EHMT2−AS1

$-\log_{10}(p)$

Chr 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22

Permutation (example)

$-\log_{10}(p)$

Chr 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22

Figure 2

A Scheme for defining 1st tier candidate genes

B *In silico* screen results for 1st tier candidate regulators

Figure 3

A Scheme for functionally annotating SNVs

Clump SNVs and identify index SNVs with *cis*-eQTLs → Differential gene expression analysis using DESeq2 → Rank genes/TEs by Wald statistic → Gene set enrichment analysis using clusterProfiler

B Index SNV associations with TE family gene sets

Differentially regulated TE families

C Index SNV associations with an L1 family gene set

Differentially regulated L1 gene set

D Index SNV associations with MSigDB Hallmark gene sets

Differentially regulated Hallmark pathways

E Index SNV associations with GO Biological Process gene sets

Differentially regulated GO BPs

F Index SNV associations with Reactome gene sets

Differentially regulated Reactome pathways

Figure 4

A Scheme for assessing the effects of candidate genes on L1 expression

IL16 or *STARD5*
CMV

Extract plasmid DNA → Electroporation with GM12878 LCLs → 1 Day → Remove dead cells → 1 Day → Pellet cells and extract RNA → Sequence mRNA (Genes AAA, TEs AAA) → Differential gene expression and GSEA

B Top enriched GO Biological Processes following *IL16* overexpression

Differentially regulated GO BPs

- Regulation of response to extracellular stimulus
- Phagocytosis recognition
- Positive chemotaxis
- Complement activation
- Interleukin-12 production
- MAPK cascade
- ERK1 and ERK2 cascade
- Negative regulation of cell differentiation
- Sterol biosynthetic process
- Gliogenesis

NES 5.0 / 2.5 / 0.0 / −2.5 / −5.0
−log$_{10}$(FDR): 1.5, 2.0, 2.5, 3.0

IL16 OE

C Top enriched Reactome gene sets following *IL16* overexpression

Differentially regulated Reactome pathways

- Creation of C4 and C2 activators
- SRP-dependent cotranslational protein targeting to membrane
- Initial triggering of complement
- Eukaryotic translation elongation
- Scavenging of heme from plasma
- Interleukin-10 signaling
- Class A/1 rhodopsin-like receptors
- HCMV early events
- Signaling by GPCR
- GPCR ligand binding

NES 5.0 / 2.5 / 0.0 / −2.5 / −5.0
−log$_{10}$(FDR): 2.4, 2.7, 3.0, 3.3

IL16 OE

D Top enriched GO Biological Processes following *STARD5* overexpression

Differentially regulated GO BPs

- Cytoplasmic translation
- Protein folding
- Positive regulation of mRNA splicing via spliceosome
- Post-translational protein targeting to ER membrane
- Oxidative phosphorylation
- Sterol biosynthetic process
- Steroid biosynthetic process
- Sterol metabolic process
- Negative regulation of cell differentiation
- Alcohol biosynthetic process

NES 5.0 / 2.5 / 0.0 / −2.5 / −5.0
−log$_{10}$(FDR): 2, 3, 4, 5

STARD5 OE

E Top enriched Reactome gene sets following *STARD5* overexpression

Differentially regulated Reactome pathways

- Translation
- SRP-dependent cotranslational protein targeting to membrane
- Regulation of expression of SLITs and ROBOs
- Eukaryotic translation elongation
- Eukaryotic translation initiation
- Signaling by GPCR
- Cholesterol biosynthesis
- Activation of gene expression by SREBF (SREBP)
- Regulation of cholesterol biosynthesis by SREBP (SREBF)
- GPCR ligand binding

NES 5.0 / 2.5 / 0.0 / −2.5 / −5.0
−log$_{10}$(FDR): 6, 8, 10, 12

STARD5 OE

F *IL16* and *STARD5* associations with TE family gene sets

Differentially regulated TE families

- *L1* subfamilies
- *Alu* subfamilies
- *ERV1* subfamilies
- *ERVL−MaLR* subfamilies
- *hAT−Charlie* subfamilies
- *ERVL* subfamilies
- *TcMar−Tigger* subfamilies
- *hAT−Tip100* subfamilies

NES 4 / 2 / 0 / −2 / −4
−log$_{10}$(FDR): 2, 3, 4, 5, 6

IL16 OE   *STARD5* OE

G *IL16* and *STARD5* associations with an L1 family gene set

L1 gene set enrichment

IL16 OE   ES / Wald stat   2.40E-5

STARD5 OE   ES / Wald stat   6.14E-7

Rank in Ordered Dataset

Figure 5

A Scheme for assessing the effects of IL16 protein on L1 expression

Seed GM12878 LCLs — Treat cells with recombinant human IL16 — Pellet cells and extract RNA — Sequence mRNA — Differential gene expression and GSEA

B Top shared GO Biological Processes

Differentially regulated GO BPs

MAPK cascade
Leukocyte differentiation
Mononuclear cell differentiation
Gliogenesis
Negative regulation of cell differentiation
Regulation of hemopoiesis
Positive regulation of cell death
Regulation of leukocyte differentiation
Positive regulation of MAPK cascade
Positive regulation of cell adhesion

IL16 OE   rhIL16 T24

NES
4
2
0
−2
−4

−log$_{10}$(FDR)
1.5
2.0
2.5
3.0
3.5

C Top shared Reactome gene sets

Differentially regulated Reactome pathways

Regulation of cholesterol biosynthesis by SREBP (SREBF)
Interleukin-10 signaling
HCMV infection
HCMV early events
Activation of gene expression by SREBF (SREBP)
Formation of tubulin folding intermediates by CCT/TriC
Metabolism of steroids
L1cam interactions
Interleukin-3, Interleukin-5, and GM-CSF signaling
Neuronal system

IL16 OE   rhIL16 T24

NES
4
2
0
−2
−4

−log$_{10}$(FDR)
1.5
2.0
2.5
3.0
3.5

D rhIL16 associations with TE family gene sets

Differentially regulated TE families

L1 subfamilies
Gypsy subfamilies
Alu subfamilies
ERVL subfamilies

rhIL16 T24

NES
5.0
2.5
0.0
−2.5
−5.0

−log$_{10}$(FDR)
1.4
1.5
1.6
1.7
1.8

L1 gene set enrichment
1.26E-2
ES
Wald stat
Rank in Ordered Dataset

Figure 6

A Scheme for assessing shared pathway changes in conditions with L1 regulation

GSEA for experimental conditions with L1 regulation

Identify common pathways

B Shared upregulated TE family gene sets

C Top shared Hallmark pathways

D Top shared GO Biological Processes

E Top shared Reactome pathways

Figure 7

A   Scheme for identifying TE eQTL co-associated aging traits

L1 *trans*-eQTLs

Generate a list of TE *trans*-eQTLs → Identify co-associated phenotypes catalogued on Open Targets Genetics → Group phenotypes by age-related MeSH trait IDs

B   Number of SNVs with an age-related MeSH trait

No 55.5%   Yes 44.5%

C   Number of associated age-related traits per SNV

# of SNVs — # of PheWAS MeSH IDs

D   Number of linked age-related traits per SNV

# of SNVs — # of Linked MeSH IDs

E   Top traits with the most number of associated SNVs

Hyperparathyroidism
Thyroid Diseases
Hypothyroidism
Hemoglobinopathies
Coronary Artery Disease
Diabetes Mellitus, Type 2
Diabetes Mellitus
Body mass index
Arthritis, Rheumatoid
Psoriasis

F   Mouse serum [IL16] with age

Young (4 - 6 months)   Old (20 - 24 months)

Serum

[IL16]

Meta−Analysis p = 0.0479

0.0402   0.205

IL16 pg/mL

Young   Old   Young   Old
Female   Male