1

Omissions of Threat Trigger Subjective Relief and Prediction Error-Like Signaling in the Human Reward and Salience Systems

Anne L. Willems*[a,b], Lukas Van Oudenhove[b,c], & Bram Vervliet[a,b]

[a] *Laboratory of Biological Psychology, Department of Brain & Cognition, KU Leuven, Belgium*

[b] *Leuven Brain Institute, KU Leuven, Belgium*

[c] *Laboratory for Brain-Gut Axis Studies (LaBGAS), Translational Research in GastroIntestinal Disorders (TARGID), Department of chronic diseases and metabolism, KU Leuven, Belgium*

*Corresponding author: Anne L. Willems

Tiensestraat 102 box 3714, 3000 Leuven, Belgium

anne.willems@kuleuven.be

Tel: +32 16 32 19 44 77

Declarations of interest: none

Abstract

The unexpected absence of danger constitutes a pleasurable event that is critical for the learning of

safety. Accumulating evidence points to similarities between the processing of absent threat and the

well-established reward prediction error (PE). However, clear-cut evidence for this analogy in humans

is scarce. In line with recent animal data, we showed that the unexpected omission of (painful) electrical

stimulation triggers activations within key regions of the reward and salience pathways and that these

activations correlate with the pleasantness of the reported relief. Furthermore, by parametrically

violating participants' probability and intensity related expectations of the upcoming stimulation, we

showed for the first time in humans that omission-related activations in the VTA/SN were stronger

following omissions of more probable and intense stimulations, like a positive reward PE signal.

Together, our findings provide additional support for an overlap in the neural processing of absent

danger and rewards in humans.

Key words: prediction error, fMRI, ventral tegmental area, substantia nigra, striatum, relief, reward

Introduction

We experience pleasurable relief when an expected threat stays away[1]. This relief indicates that the outcome we experienced ("nothing") was better than we expected it to be ("threat"). Such a mismatch between expectation and outcome is generally regarded as the trigger for new learning, and is typically formalized as the prediction error (PE) that determines how much there can be learned in any given situation[2]. Over the last two decades, the PE elicited by the absence of expected threat (threat omission PE) has received increasing scientific interest, because it is thought to play a central role in learning of safety. Impaired safety learning is one of the core features of clinical anxiety[3]. A better understanding of how the threat omission PE is processed in the brain may therefore be key to optimizing therapeutic efforts to boost safety learning. Yet, despite its theoretical and clinical importance, research on how the threat omission PE is computed in the brain is only emerging.

To date, the threat omission PE has mainly been studied using fear extinction paradigms that mimic safety learning by repeatedly confronting a human or animal with a threat predicting cue (conditional stimulus, CS; e.g. a tone) in the absence of a previously associated aversive event (unconditional stimulus, US; e.g., an electrical stimulation). These (primarily non-human) studies have revealed that there are striking similarities between the PE elicited by unexpected threat omission and the PE elicited by unexpected reward. In the context of reward, it is well-established that dopamine neurons in the ventral tegmental area (VTA) and substantia nigra (SN) of the midbrain increase their firing rate to unexpected rewards (positive PE), suppress their firing for unexpected reward omissions (negative PE) and show no change in firing in response to completely predicted rewards, in line with a formalized PE[4,5]. Likewise, in fear extinction, dopaminergic neurons in the VTA phasically increase their firing rates to early (unexpected), but not late (expected) US omissions [6–10], which consequently triggers downstream dopamine release in the nucleus accumbens (NAc) shell[6,9,10]. Furthermore, optogenetically blocking (or enhancing) the firing rate of these dopaminergic VTA neurons during US omissions impairs (or facilitates) subsequent fear extinction learning[6–8]. Notably, such dopaminergic VTA/NAc responses to threat omissions have also been observed in other experimental tasks, such as conditioned inhibition[11] and avoidance[12,13], confirming that these neural activations match a more general threat omission PE-signal.

In humans, reward-like PE responses to threat omissions during extinction and avoidance have

mainly been reported in projection regions of dopaminergic midbrain neurons, such as the ventral

striatum (more specifically, NAc and ventral putamen) and prefrontal areas (the ventromedial prefrontal

cortex, vmPFC)[14–19]. Activations in these regions correlate with computationally modeled PE

signals[14,15,17] and are modulated by pharmacological manipulation of dopamine receptors[17] and by

genetic mutations that are known to enhance striatal phasic dopamine release[14]. Furthermore,

connectivity analyses revealed that NAc activations during US omissions in fear extinction were

functionally coupled to VTA/SN activations, and that this connectivity was enhanced by the

administration of the dopamine precursor L-dopa prior to extinction[17]. The emerging picture is that

cortico-striatal activations to unexpected omissions of threat are triggered by dopaminergic inputs from

the VTA/SN, just like a positive reward PE, and that these activations play a central role in different

types of threat omission-induced learning such as fear extinction and avoidance learning[20,21]. Still, direct

observations of threat omission-related VTA/SN responses are currently lacking in humans.

As mentioned above, unexpected omissions of threat not only trigger neural activations that

resemble a reward PE, they are also accompanied by a pleasurable emotional experience: relief[1].

Because these feelings of relief coincide with the PE at threat omission, relief has been proposed to be

an emotional correlate of the threat omission PE[18,22]. Indeed, emerging evidence has shown that

subjective experiences of relief follow the same time-course as theoretical PE during fear extinction.

Participants in fear extinction experiments report high levels of relief pleasantness during early US

omissions (when the omission was unexpected and the theoretical PE was high) and decreasing relief

pleasantness over later omissions (when the omission was expected and the theoretical PE was low)[22,23].

Accordingly, preliminary fMRI evidence has shown that the pleasantness of this relief is correlated to

activations in the NAc at the time of threat omission[18]. In that sense, studying relief may offer important

insights in the mechanism driving safety learning.

However, is a correlation with the theoretical PE over time sufficient for neural activations/relief

to be classified as a PE-signal? In the context of reward, Caplin and colleagues proposed three necessary

and sufficient criteria all PE-signals should comply to, independent of the exact operationalizations of

expectancy and reward (the so-called axiomatic approach[24,25]; which has also been applied to aversive

PE[26–28]). Specifically, the magnitude of a PE signal should: (1) be positively related to the magnitude of

the reward (larger rewards trigger larger PEs); (2) be negatively related to likelihood of the reward (more

probable rewards trigger smaller PEs); and (3) not differentiate between fully predicted outcomes of

different magnitudes (if there is no error in prediction, there should be no difference in the PE signal).

The previously discussed fear conditioning and extinction studies have been invaluable for

clarifying the role of the threat omission PE within a learning context[14–17,22,23]. However, these studies

were not tailored to create the varying intensity and probability-related conditions that are required to

systematically evaluate the threat omission PE in the light of the PE axioms. First, these only included

one level of aversive outcome: the electrical stimulation was either delivered or omitted; but the intensity

of the stimulation was never experimentally manipulated within the same task. As a result, the

magnitude-related axiom could not be tested. Second, as safety learning progressively developed over

the course of extinction learning, the most informative trials to evaluate the probability axiom (i.e. the

trials with the largest PE) were restricted to the first few CS+ offsets of the extinction phase, and the

exact number of these informative trials likely differed across participants as a result of individually

varying learning rates. This limited the experimental control and necessary variability to systematically

evaluate the probability axiom. Third, because CS-US contingencies changed over the course of the task

(e.g. from acquisition to extinction), there was never complete certainty about whether the US would

(not) follow. This precluded a direct comparison of fully predicted outcomes. Finally, within a learning

context, it remains unclear whether brain responses to the threat omission are in fact responses to the

violation of expectancy itself, or whether they are the result of subsequent expectancy updating.

Based on these reasons, we recently developed the Expectancy Violation Assessment (EVA)

task[29] in order to study threat omission responses outside of a learning context. By providing verbal

instructions on the probability and intensity of an upcoming electrical stimulation, which are then

violated by not delivering the stimulation, we showed that the experienced pleasantness of the omission-

relief reflects the degree of fearful expectation violation, with omissions of more intense and more

probable stimulations eliciting more pleasurable feelings of relief, much like a PE-signal.

Here, we applied the EVA-task in the MRI scanner to investigate brain responses to unexpected

omissions of threat in greater detail, examine their similarity to reward PE-signals, and explore the link

with subjective relief. Specifically, participants received trial-by-trial instructions about the probability

(0%, 25%, 50%, 75% and 100%) and intensity (weak, moderate, strong) of a potentially painful

upcoming electrical stimulation, time-locked by a countdown clock (see Fig.1A). While stimulations

were always delivered on 100% trials and never on 0% trials, most of the other trials (25%-75%) did

not contain the expected stimulation and hence provoked an omission PE. We expected that (1)

expected-but-omitted stimulation would trigger increased activity within key areas of the reward circuit

(such as the VTA/SN, NAc, left ventral Putamen and vmPFC); that (2) this omission-related activity

would fit the three criteria of a positive reward PE, and that (3) this activity would be related to self-

reported relief. These hypotheses and the analysis approach were preregistered on Open Science

Framework (OSF, https://osf.io/ugkzf). Small deviations related to the approach are reported in the

supplementary material (Supplementary Note 5).

135   **Results**

Self-reported relief and omission SCR track omissions of threat in a PE-like manner.

The verbal instructions were effective at raising the expectation of receiving the electrical

stimulation in line with the provided probability and intensity levels. Anticipatory SCR, which we used

as a proxy of fearful expectation, increased as a function of the probability and intensity instructions

(see Supplemental Figure 3). Accordingly, post-experimental questions revealed that by the end of the

experiment participants recollected having received more stimulations after higher probability

instructions, and were willing to exert more effort to prevent stronger hypothetical stimulations (see

Supplemental Figure 2).

Replicating our previous findings[29], self-reported relief-pleasantness and omission SCR tracked

the PE signal during threat omission (see Fig.1B/C). Overall, unexpected (non-0%) omissions of threat

elicited higher levels of relief-pleasantness and omission SCR than fully expected omissions (0%),

evidenced by a main effect of Probability in the 4 (Probability: 0%, 25%, 50%, 75%) x 4 (Run: 1, 2, 3,

4) LMM (For relief pleasantness ($N = 31$): $F(3,1417) = 188.34$, $p < .001$, $\omega_p^2 = 0.28$; and for omission

SCR ($N = 26$): $F(3,1190) = 72.90$, $p < .001$, $\omega_p^2 = 0.15$, with responses to all non-0% probability levels

being significantly higher than responses to 0%, p's $< .001$, Bonferroni-Holm corrected). Furthermore,

relief-pleasantness and omission SCR to unexpected omissions (non-0% omissions) increased as a

function of instructed Probability and Intensity, in line with the first two PE axioms, evidenced by main

effects of Probability (for relief-pleasantness ($N = 31$): $F(2,1031) = 30.64$, $p < .001$, $\omega_p^2 = 0.05$, all

corrected pairwise comparisons, p's $< .005$; for omission SCR ($N = 26$): $F(2,862) = 5.15$, $p < .01$, ω_p^2

$= 0.01$, with corrected 75% to 25% comparison, $p < .01$), and Intensity (for relief-pleasantness ($N = 31$):

$F(2, 1031) = 623.79$, $p < .001$, $\omega_p^2 = 0.55$, all corrected pairwise comparisons, p's $< .001$; for omission

SCR ($N = 26$): $F(2,862.01) = 107.47$, $p < .001$, $\omega_p^2 = 0.20$, all corrected pairwise comparisons, p's $<$

.001) in a 3 (Probability: 25%, 50%, 75%) x 3 (Intensity: weak, moderate, strong) x 4 (Run: 1, 2, 3, 4)

LMM. Relief-pleasantness also showed a significant Probability x Intensity interaction ($F(4,1031) =$

3.76, $p < .005$, $\omega_p^2 = 0.01$), indicating that the effect of probability was most pronounced for omissions

of moderate stimulation (all p's $< .05$). Note that while there was a general drop in reported relief

pleasantness and omission SCR over time, the effects of Probability and Intensity remained present until

the last run (see Supplementary Figure 5). This further confirms that probability and intensity

instructions were effective until the end of the task.

Figure 1. **Experimental Design and Behavioral Results. A.** All trials started with instructions on the probability (ranging from 0% to 100%) and intensity (weak, moderate, strong) of a potentially painful electrical stimulation (1 second), followed by the addition of a countdown bar that indicated the exact moment of stimulation or omission. The duration of the countdown clock was jittered between 3-7 seconds. Then, the screen cleared and the electrical stimulation was either delivered or omitted. Most of the trials (48 trials) did not contain the anticipated electrical stimulation (omission trials). Following a delay of 4 to 8 seconds, a rating scale appeared, probing stimulation-unpleasantness on stimulation trials, and relief-pleasantness on omission trials. After 8 seconds, an ITI between 4 and 7 seconds started, during which a fixation cross was presented on the screen. The task consisted in total of 72 trials, divided equally over 4 runs (18 trials/run). Each run contained all probability (25, 50, 75) x intensity (weak, moderate, strong) combinations exactly once not followed by the stimulation (9 omission trials), three 0%-omission trials (without any intensity information), three 100%-stimulation trials (followed by the stimulation of the given intensity, one per intensity level per run), and three additional trials from the probability (25,50,75) x Intensity (weak, moderate, strong) matrix that were followed by the stimulation (per run each level of intensity and probability was once paired with the stimulation). For a detailed overview of the trials see Supplementary Figure 1. **B.** SCR were scored as the time integral of the deconvoluted phasic activity (using CDA in Ledalab) within a response window of 1-4s post omission. Responses were square root transformed. SCR were larger following omissions of more probable and more intense US omissions. **C.** The pleasantness of the relief elicited by US omissions was rated on a VAS-scale ranging from 0 (neutral) to 100 (very pleasant). Omission-relief was rated as being more pleasant following omissions of more intense and more probable US. In both graphs, individual data points are presented, with the group averages plotted on top. The error bars represent standard error of the mean.

Unexpected omissions of threat trigger activations in the VTA/SN and ventral Putamen, but deactivations in the vmPFC

In line with our hypothesis and similar to relief and omission SCR, unexpected (non-0%) omissions of threat elicited on average stronger fMRI activations than fully expected (0%) omissions in the VTA/SN ($t(30) = 4.48$, $p < .001$, $d = 0.81$) and left ventral putamen ($t(30) = 3.50$, $p < .005$, $d = 0.63$) ROIs (see Fig. 2A/B). Surprisingly, NAc showed no significant change in activation ($t(30) = -0.59$, $p = .56$) (Fig. 2D), and vmPFC showed a significant deactivation ($t(30) = -4.71$, $p < .001$, $d = -0.85$) (Fig. 2C). This apparent deactivation could indicate that omission-related responses were lower for unexpected omissions compared to expected omissions. However, it could also have resulted from lingering safety-related vmPFC activations to the 0%-instructions (corresponding to certainty that no stimulation will follow). Such safety-related vmPFC activations are indeed commonly observed during the presentation of CS- in Pavlovian fear conditioning[30]. To exclude this alternative hypothesis, we examined the non0% > 0% contrast during the instruction window. We found no significant difference in vmPFC activation between 0% and non-0% trials, either as ROI average ($t(30) = -1.69$, $p_{unc} = .1$) or voxel-wise, SVC within the vmPFC mask (see Supplementary Figures 7-9 and Supplementary Tables 3-5 for full description of the anticipatory fMRI activations). This follow-up analysis suggests that the deactivation to unexpected omissions only emerged after the instruction window, and could therefore not be explained by safety-related activation that were obtained during 0% trials.

Figure 2. **Omission-related activations in the a priori ROIs.** Unexpected omissions of stimulation (non0% > 0%) triggered significant fMRI responses in (**A**) the VTA/SN, and (**B**) left ventral putamen, but deactivations in (**C**) the vmPFC and no change in activation in (**D**) the NAc. Only for the VTA/SN did the activations increase with increasing Probability and Intensity of omitted stimulation. vmPFC responses decreased with increasing intensity of the omitted stimulation. Fully predicted stimulations (100%) elicited stronger activations than fully predicted omission (0%) in (**E**) the VTA/SN, no difference in activation for (**F**) the left Putamen and (**G**) the NAc, and stronger deactivations for fully predicted stimulation versus omission in (**H**) the vmPFC. In all figures, the unexpected omission maps were overlayed with the a priori ROI masks (in teal) and were displayed at threshold $p < .001$ (unc) for visualization purposes. The crosshairs represent the peak activation within each a priori ROI. The extracted beta-estimates in figures A-D represent the ROI averages from each non-0% > 0% contrast (i.e., 25%>0%; 50%>0%; and 75%>0% for the weak, moderate and strong intensity levels). Any positive beta therefore indicates a stronger activation in the given region compared to a fully predicted omission. Any negative beta indicates a weaker activation. The dots and error bars represent the mean and standard error of the mean.

Omission-related VTA/SN, but not striatal or vmPFC activations increased in a PE-like manner

We next assessed if the omission-related (de)activations could represent reward-like PE-signals

by testing the PE axioms for each ROI separately. For axiom 1 and 2, we contrasted omissions following

all intensity x probability combination with 0%-omissions and extracted ROI-specific beta averages.

These beta-estimates were then entered into linear mixed models that included instructed intensity and

probability as regressors of interest, and averaged US-unpleasantness as regressor of no-interest, in

addition to a subject-specific intercept. Axiom 3 was tested via a one-sample (two-sided) t-test over the

100%-stimulation versus 0%-omission contrast.

We found that only omission-related **VTA/SN** activations partially fit the profile of a positive

reward PE-signal (Fig. 2A). Activations were stronger following omissions of more intense (Axiom 1,

$F(2,240) = 6.14$, $p < .005$, $\omega_p^2 = .04$), and at trend level of more probable threat (Axiom 2, $F(2,240) =$

2.94, $p = .055$, $\omega_p^2 = .02$). However, fully predicted stimulations (100%-trials) elicited stronger

activations than fully predicted omissions (0%-trials) ($V_{wilcoxon} = 452$, $p < .001$, r = 0.72), contradicting

axiom 3 (Fig 2E).

Unlike previous findings[14,15,17,18], we found no evidence for striatal reward PE-like processing

of threat omissions. While **ventral putamen** responses were stronger following unexpected omissions

compared to expected omissions, these activations did not increase with increasing intensity (axiom 1,

$F(2,240) = 2.29$, $p = .10$) or probability (axiom 2, $F(2,240) = 0.57$, $p = .57$) (Fig 2B). Notably, we did

find anecdotal evidence that fully predicted stimulations and fully predicted omissions triggered similar

activations in the ventral putamen (axiom 3, $t(30) = 1.23$, $p = .46$, BF of 2.62 in favor of the null-

hypothesis, Fig. 2F). This indicates that ventral putamen activations were exclusively triggered by

unexpected threat omissions, and not by fully predicted outcomes, which is similar to a PE-signal.

In general, there was no evidence for omission-related **NAc** activations. Activations were not

affected by intensity (axiom 1, $F(2,240) = 1.88$, $p = . 16$) or probability instructions (axiom 2, $F(2,240)$

$= 0.75$, $p = .48$) (Fig 2D), and while there were no differences in activation between fully predicted

stimulation and fully predicted omission ($t(30) = 0.67$, $p = .51$, BF in favor of null hypothesis = 4.24),

this equivalence was most likely caused by an overall absence of activation (Fig 2G).

Finally, omission-related **vmPFC** deactivations were stronger for omissions of more intense

threat (axiom 1, $F(2,240) = 9.29$, $p < .001$, $\omega_p^2 = 0.06$), but were unaffected by probability instructions

(axiom 2, $F(2,240) = 1.78$, $p = .17$) (Fig 2C). Furthermore, responses were smaller for completely

predicted stimulation compared to completely predicted omission (axiom 3, $t(30) = -8.65$, $p < .001$, $d =$

-1.55, Fig 2H). Taken together, we found no evidence that vmPFC deactivations reflected a positive

reward PE-like signal.

A potential explanation for the absent probability effects in the putamen and vmPFC might be

that the effects were obscured by including participants who did not believe the probability instructions.

Indeed, the provided instructions did not map exactly onto the actually experienced probabilities, but

were all followed by stimulation in 25% on the trials (except for the 0% trials and the 100% trials). We

therefore reran our analyses on a subset of participants who showed probability-related increases in their

anticipatory SCR during the countdown clock (N = 21, larger SCR to 75% compared to 25%

instructions, see Supplementary Figure 4), which we used as a post-hoc index of actual probability-

related expectancy. This subgroup analysis revealed no additional effect of Probability for the ventral

putamen or the vmPFC, but it rendered the effect of Intensity for the ventral Putamen significant

($F(2,160) = 3.10$ $p < .05$, $\omega_p^2 = 0.03$). In addition, it increased the effect of probability for the VTA/SN

activation ($\omega_p^2 = .02$ to .05). Likewise, a post-hoc trial-by-trial analysis of the omission-related fMRI

activations confirmed that the Probability effect for the VTA/SN activations was stable over the course

of the experiment (no Probability x Run interaction) and remained present when accounting for the

Gambler's fallacy (i.e., the possibility that participants start to expect a stimulation more when more

time has passed since the last stimulation was experienced) (see supplemental note 6). Overall, these

post-hoc analyses further confirm the PE-profile of omission-related VTA/SN responses.

Anterior insula and dmPFC/aMCC clusters show increased activation for unexpected omissions

of threat in a PE-like fashion.

We then explored neural threat omission processing within a wider secondary mask that

combined our primary ROIs with additional regions that have previously been associated to reward, pain

and PE processing (such as the wider striatum, including the caudate nucleus, and putamen; midbrain

nuclei, including the periaqueductal gray (PAG), and red nucleus; medial temporal structures, including

the amygdala and hippocampus; midline thalamus, habenula and cortical regions, including the anterior

insula (aINS), orbitofrontal (OFC), dorsomedial prefrontal (dmPFC) and anterior cingulate (ACC)

cortices). Significant omission-processing clusters (contrast non0% > 0% omissions) were extracted

from the mask using a cluster-level threshold ($p < .05$, FWE-corrected), following a primary voxel-

threshold ($p < .001$) and included the bilateral anterior insula, bilateral putamen, and a medial cortical

cluster encompassing parts of the dorsomedial prefrontal cortex (dmPFC) and the anterior medical

cingulate cortex (aMCC) (Fig. 3A-D). The left putamen cluster bordered and minimally overlapped with

our predefined ventral putamen ROI (4 out of 82 voxels) which was based on the peak PE activity in

previous studies[14,15]. Exploratory analyses within a wider whole-brain grey-matter mask identified

several other omission-processing clusters (see Supplementary Figure 10, Supplementary Table 6).

Probability and Intensity related activity modulations of these clusters can be found in Supplementary

Figure 11.

Follow-up analysis of the bilateral **putamen** clusters confirmed that putamen activations did not fit

the profile of a positive reward PE. Activations in neither cluster increased with increasing intensity

(axiom 1, left: $F(2,240) = 0.18$, $p = .83$; right: $F(2,240) = 0.06$, $p = .94$) nor probability of threat (axiom

2, left: $F(2,240) = 1.41$, $p = .25$; right: $F(2,240) = 0.87$, $p = .42$), but like for the a priori ventral Putamen

ROI, activations were comparable for fully predicted outcomes, especially in the left cluster (axiom 3,

left: $V = 339$, $p = .46$, BF = 2.41, right: $t(30) = 1.83$, $p = .46$, BF = 1.20).

Positive reward PE-like responses were found in the bilateral **aINS** and **dmPFC/aMCC**, where

omission-related activations were stronger following omissions of more intense (left aINS: $F(2,240) =$

8.95, $p < .001$, $\omega_p^2 = 0.06$; right aINS: $F(2,240) = 13.49$, $p < .001$, $\omega_p^2 = 0.09$; dmPFC/aMCC: $F(2,240)$

$= 6.59$, $p < .005$, $\omega_p^2 = 0.04$), and at trend level of more probable threat (left aINS: $F(2,240) = 1.87$, p

$=.16$, right aINS: $F(2,240) = 2.78$, $p = 0.06$, $\omega_p^2 = 0.01$; dmPFC/aMCC: $F(2,240) = 2.48$, $p =.09$, $\omega_p^2 =$

0.01). Notably aINS and dmPFC/aMCC clusters extended beyond our predefined secondary mask, and

including all adjacent above-threshold voxels ($p < .001$) rendered the effects of probability significant

(see Supplementary Figure 11). However, fully predicted stimulations also elicited stronger activations

than fully predicted omissions (left aINS: $t(30) = 3.21$, $p < .05$, $d = 0.58$, right aINS: $t(30) = 5.26$, $p <$

$.001$, $d = 0.95$, mdPFC/aMCC: $t(30) = 5.57$, $p < .001$, $d = 1.00$).

Finally, in addition to the vmPFC deactivations (which fell entirely within our vmPFC mask), we

found trend-level deactivations for unexpected omission in the **right amygdala** ($p = .053$). These

deactivations were stronger for omissions of more intense ($F(2,240) = 3.26$, $p < .05$, $\omega_p^2 = 0.02$), but not

more probable threat ($F(2,240) = 0.43$, $p = .65$). Furthermore, fully predicted stimulations triggered

larger deactivation than fully predicted omissions ($t = -2.77$, $p = .07$, $BF = 0.21$).

Figure 3. **Omission-related activations in the secondary mask.** We extracted unexpected omission (non0% > 0%) processing clusters within our secondary mask, using a voxel threshold, $p < .001$, followed by a cluster-threshold (FWE-corrected) of $p < .05$. We found significant positive omission processing clusters in **(A)** the bilateral putamen, **(B)** bilateral aINS, **(C)** dmPFC/aMCC, and a trend-level negative omission processing cluster in **(D)** the right amygdala. Omission related activations in the bilateral aINS and dmPFC/aMCC increased with increasing probability and intensity of omitted threat, whereas amygdala activations decreased with omissions of increasingly intense threat. Nevertheless, fully predicted stimulations (100%) elicited stronger activations than fully predicted omission (0%) in **(F)** the bilateral aIns, and **(G)** the dmPFC/aMCC, and stronger deactivations in the **(H)** right amygdala, but no difference in activation in the **(E)** bilateral putamen. In all figures, the unexpected omission maps are displayed at threshold $p < .001$ (unc) for visualization purposes. The extracted beta-estimates in figures A-D represent the ROI averages from each non-0% > 0% contrast (i.e., 25%>0%; 50%>0%; and 75%>0% for the weak, moderate and strong intensity levels). Any positive beta therefore indicates a stronger activation in the given region compared to a fully predicted omission. Any negative beta indicates a weaker activation. The dots and error bars represent the mean and standard error of the mean.

Omission-related activations are related to self-reported relief-pleasantness

We then examined whether omission-related fMRI activations were related to self-reported relief-pleasantness on a trial-by-trial basis. In a pre-registered analysis, we entered z-scored relief-pleasantness ratings as a parametric modulator to the omission regressor in a separate GLM that did not distinguish between the different probability x intensity levels. We found that the VTA/SN ($t(30) = 3.26$, $p < .01$, $d = 0.59$) and ventral putamen ROI (at trend level, $t(30) = 2.22$, $p = .068$, $d = 0.40$) were positively modulated by relief-pleasantness ratings, whereas the vmPFC ROI was negatively modulated by relief-pleasantness ratings ($V = 46$, $p < .001$, $r = 0.71$) (Fig. 4A-D). The positive and negative modulations indicate that stronger omission-related activations in the VTA/SN and ventral putamen, and stronger deactivations in the vmPFC were associated with more pleasant relief-reports. Likewise, the bilateral aINS ($t > 6.70$, $p < .001$, $d > 1.20$), dmPFC/aMCC ($t(30) = 6.13$, $p < .001$, $d = 1.10$), and right putamen ($t(30) = 4.90$, $p < .001$, $d = 0.88$), and at trend level left putamen ($t(30) = 2.37$, $p = 0.097$,

Figure 4. **Relief modulation in the a priori and secondary ROIs.** Omission-related activations were modulated by trial-by-trial levels of relief-pleasantness in **(A)** the VTA/SN, **(B)** left ventral Putamen**,** and **(D)** vmPFC, but not in **(C)** the NAc. Omission-related activations in the secondary ROIs were modulated by trial-by-trial levels of relief-pleasantness in **(A)** the right putamen, **(B)** bilateral aINS**, (C)** dmPFC/aMCC, and the **(D)** right amygdala. In all figures, the relief modulation maps are displayed at threshold p < .001 (unc) for visualization purposes.

288   $d = 0.43$) clusters we identified from the secondary mask were positively modulated, and right amygdala

289   was negatively modulated by relief-pleasantness (V = 59, $p < .001$, r = 0.67) (Fig. 4E-H). Omission-

290   related NAc activation was unrelated to self-reported relief-pleasantness (V = 214, $p = 0.52$).

A neural signature for relief-pleasantness

The (mass univariate) parametric modulation analysis showed that omission-related fMRI

activity in our primary and secondary ROIs correlated with the pleasantness of the relief. However,

given that each voxel/ROI is treated independently in this analysis, it remains unclear how the

activations were embedded in a wider network of activation across the brain, and which regions

contributed most to the prediction of relief. To overcome these limitations, we trained a (multivariate)

LASSO-PCR model (Least Absolute Shrinkage and Selection Operator-Regularized Principle

Component Regression) in order to identify whether a spatially distributed pattern of brain responses

can predict the perceived pleasantness of the relief (or "neural signature" of relief)[31]. Because we used

the whole-brain pattern (and not only our a priori ROIs), this analysis is completely data driven and can

thus identify which clusters contribute most to the relief prediction. We trained the model using fivefold

cross-validation with trial-by-trial whole-brain omission-related activation-maps as predictors, and trial-

by-trial relief-pleasantness ratings as outcome.

Predicted and reported relief correlated significantly ($r = 0.28$, $p < .001$) (Fig. 5C), indicating

that part of the variance in reported relief-pleasantness could be explained by the neural relief signature

response (Fig. 5A). Follow-up bootstrap tests (5000 samples) identified a distributed pattern of positive

and negative predictive clusters across the brain (Fig. 5B, Table 1). Increased responses in these clusters

predicted increased/decreased relief-pleasantness, respectively. Notably, bootstrap tests indicated that

none of our a priori regions of interest significantly contributed to the signature. This was further

supported by a pre-registered virtual lesion analysis where we compared the predictive performance of

our LASSO-PCR model based on whole-brain data to separate models excluding voxels from our main

ROIs in each iteration (see Fig. 5D).

Figure 5. **LASSO-PCR based neural signature of relief-pleasantness. A.** Relief predictive signature map consisting of all
voxels within a grey matter mask. All weights were used for prediction. **B.** Thresholded signature map ($p < .001$), consisting
of clusters that contribute significantly to the relief prediction (all clusters < 65 voxels). **C.** Predicted and reported relief-
pleasantness correlated significantly, $r = 0.28$, $p < .001$. Each line/color represents data of a single participant. **D.** Correlations
between reported and predicted relief in all lesion models (in each model one of the ROIs was removed from the grey matter
mask). The stable correlation across models confirmed that none of our ROIs contributed significantly to the relief-predictive
signature model.

Table 1: Main clusters contributing to the relief signature identified via bootstrapping

Positive weight clusters

L/R	Region	K	MNI Coordinates (xyz)			Z (peak)
R	Cerebellum Crus 2	17	6	-85	-30	0.006
L	Cerebellum Crus 1	18	-41	-75	-28	0.005
L	Inferior orbitofrontal gyrus	10	-21	14	-23	0.005
R	Inferior temporal gyrus	10	56	-23	-15	0.006
L	Middle orbitofrontal gyrus	15	-25	54	-15	0.007
R	Middle orbitofrontal gyrus	13	38	58	-10	0.007
R	Caudate gyrus	12	4	14	-1	0.007
L	Superior temporal gyrus	11	-55	2	1	0.005
R	Rolandic operculum	19	58	10	3	0.006
R	Superior occipital gyrus	13	20	-103	7	0.006
L	Middle occipital gyrus	14	-23	-97	7	0.006
R	Calcarine gyrus	18	12	-79	10	0.007

		K	x	y	z	Z
L	Middle occipital gyrus	12	-31	-95	14	0.005
R	Middle temporal gyrus	19	54	-47	14	0.005
R	Cuneus	12	4	-79	25	0.006
R	Supramarginal gyrus	12	52	-37	29	0.005
L	Superior occipital gyrus	17	-25	-71	38	0.004
R	Precentral gyrus	15	50	6	43	0.005
L	Superior occipital gyrus	14	-13	-79	43	0.004
L	Mid cingulate gyrus	27	-9	-23	45	0.005
R	Mid cingulate gyrus	12	2	-13	45	0.005
Negative weight clusters						
R	Cerebellum Crus 1	12	30	-69	-28	-0.005
L	Cerebellum Crus 1	11	-17	-89	-19	-0.007
R	Cerebellum Crus 1	41	30	-85	-19	-0.014
R	Fusiform gyrus	10	28	-49	-8	-0.006
R	Superior temporal gyrus	10	52	-2	-6	-0.004
R	Inferior occipital gyrus	11	28	-89	-4	-0.006
R	Superior temporal gyrus	10	66	-17	-4	-0.006
L	Superior temporal gyrus	16	-63	-5	-1	-0.005
L	Caudate	10	-17	18	1	-0.006
L	Middle temporal gyrus	14	-57	-65	-1	-0.004
R	Middle occipital gyrus	12	34	-93	5	-0.005
L	Middle occipital gyrus	11	-39	-89	3	-0.004
R	Middle temporal gyrus	40	58	-57	10	-0.005
L	Calcarine gyrus	18	-11	-81	12	-0.007
L	Cuneus	20	-3	-89	21	-0.006
L	Angular gyrus	16	-45	-59	29	-0.005
L	Middle frontal gyrus	12	-33	46	34	-0.005
R	Postcentral gyrus	64	46	-31	49	-0.005
R	Middle frontal gyrus	19	26	32	45	-0.004
L	Postcentral gyrus	10	-49	-17	45	-0.005
R	Middle frontal gyrus	12	26	10	47	-0.004

323
324 *Note.* Contributing clusters are defined based on voxels-wise FDR-correction with $q < .05$, $k >10$; *L/R*
325 indicates if the cluster (or peak of the cluster) is part of the left or right hemisphere; *Region* name is
326 identified using the AAL atlas; *K* is the number of voxels in the cluster; *coordinates* are the MNI
327 coordinates of cluster peak; Z is the weight of the cluster peak.
328

Discussion

We examined whether brain reactions to unexpected omissions of threat qualify as positive

reward PE signals, and explored their link with subjective relief. We showed that, similar to an

unexpected reward, unexpected omissions of stimulation triggered fMRI activations within key regions

of the reward and salience pathways (such as the VTA/SN, putamen, dmPFC/aMCC and aINS), and that

the magnitude of these activations correlated with the pleasantness of the reported relief. Moreover,

omission-related activations in the VTA/SN, the primary reward PE-encoding region in animals[4,5] and

humans[32,33], also tracked the probability and intensity of omitted stimulation, in line with the first two

criteria of a positive reward-PE signal. In contrast, the NAc and the vmPFC, two other regions that have

previously been implicated in reward PE, threat omission processing, and the valuation of rewards and

relief[16–18,34–36], showed no (NAc) or even decreased activations (vmPFC) in response to omitted threat;

and no correlation (NAc) and a negative correlation (vmPFC) with subjective relief.

Overall, the observed activity pattern of the VTA/SN supports the hypothesis that unexpected

omissions of threat are processed as reward PE-like signals in the human brain. However, there are two

caveats to this interpretation. First, it remains unclear whether these activations reflect the activity of

dopamine cells in this region. The dopamine basis of the reward PE is well established[4,5], and similar

VTA dopaminergic responses to threat omissions have been found during fear extinction in rodents[6–

8,11]. Yet, the nature of fMRI measurements does not allow us to directly trace back the observed BOLD

responses to the phasic firing of dopamine cells at the time of threat omission. Still, the general location

of the omission-related VTA/SN activation is consistent with a dopaminergic basis[37,38], as the peak

activation falls within a more medial subregion of the SN, which is predominantly composed of

dopamine cells (>80% of all cells)[39]. In addition, a recent human fear extinction study found that

ingestion of the dopamine precursor L-Dopa increased functional coupling between NAc and VTA/SN

at the time of threat omission[17].

A second caveat to the PE-interpretation of VTA/SN activations in the light of the present results

is that fully predicted stimulations (100%) triggered stronger activations than fully predicted omissions

(0%), which violates the third PE axiom and therefore opposes a PE interpretation. Theoretically, the

third axiom states that a pure PE-signal would not differentiate between these fully predicted outcomes,

whatever the outcomes are. As such, the violation implies that the VTA/SN responses could not

represent PE-signals. Yet, we argue that this axiom should not be a decisive criterion when comparing

fully predicted threat to its fully predicted omission. Specifically, a wealth of studies has reported

VTA/SN responses to both salient events (even aversive) and PE[4,40], and showed that these responses

might be functionally and anatomically distinct[38,41,42]. Moreover, on a smaller scale, recent

electrophysiological studies suggest that dopaminergic PE-signals themselves consist of two

components: an initial unselective and short-lasting component that detects any event of sufficient

intensity, followed by a subsequent component that codes the prediction error[4]. Thus, given that we

could not control for the delivery of the stimulation in the 100% > 0% contrast (the delivery of the

stimulation completely overlapped with the contrast of interest), it is impossible to disentangle responses

to the salience of the stimulation from those to the predictability of the outcome. A fairer evaluation of

the third axiom would require outcomes that are roughly similar in terms of salience. When evaluating

threat omission PE, this implies comparing fully expected threat omissions following 0% instructions

to fully expected absence of stimulation at another point in the task (e.g. during a safe intertrial interval).

Beyond the midbrain, unexpected omissions of threat also elicited striatal activations that spread

across the bilateral putamen (as in[14,15]). Although these activations did not increase with increasing

probability and intensity of the omitted stimulation, they were correlated with the reported relief-

pleasantness and only occurred when the omission was unexpected (not when the outcome was fully

expected, 100%-stimulation or 0% omission). Yet, in contrast to our predictions, the activations did not

extend to the NAc. This was surprising, because the NAc is the main striatal projection area of the VTA,

and numerous rodent and human studies have attributed reward PE signaling to this region[25,43]. Likewise,

two human studies found that self-reported relief and modeled PE-estimates at the time of threat

omission covaried with NAc activations[17,18].

Nevertheless, a growing body of research now indicates that human PE encoding is not confined

to the NAc, but instead spreads across the striatum. Meta-analyses on reinforcement learning in humans

have identified the (left) putamen, and not the NAc as the most consistent reward PE-encoding region

across studies[44,45]. Arguably, this emerging functional divergence between rodent and human striatal

responses may be linked to neuroanatomical differences at the level of the midbrain. Specifically, where

the (medial) VTA and its NAc projections have revealed marked omission-processing in rodents, we

found the strongest omission-related activations in the (medial) SN and the putamen, which is a SN

projection target. In addition, task-related differences might have contributed to the absent NAc

activations. One of the meta-analyses found that the NAc was especially involved in PE-responses when

a response is required to obtain the reward/threat omission (e.g., in instrumental tasks)[44]. Arguably, as

the EVA task does not allow active control over the omission, it might not have engaged the NAc.

Together, our findings call for caution when directly extrapolating rodent findings to human fMRI

results.

We found omission-related *deactivations* in the vmPFC that were stronger following omissions

of more intense threat, and that correlated with the reported relief. This is again in contrast with our

predictions, and with previous studies that showed vmPFC activations during early US omissions in the

context of fear extinction[16] and positive associations between vmPFC activity and subjective

pleasure[35,36]. Instead, we found vmPFC deactivations for both omissions and stimulations. Interestingly,

these deactivations were not limited to the vmPFC, but spanned across key regions of the default mode

network[46], such as the PCC and precuneus (see Supplementary Figure 10 and Supplementary Table 6).

Furthermore, they were accompanied by widespread activations in key regions of the salience network[47]

(such as the VTA/SN, striatum, aINS, dmPFC/aMCC). One potential explanation is therefore that the

deactivation resulted from a switch from default mode to salience network, triggered by the salience of

the unexpected threat omission or by the salience of the experienced stimulation.

In addition to examining the PE-properties of neural omission responses in our a priori ROIs,

we trained a LASSO-PCR model to establish a signature pattern of relief. One interesting finding that

only became evident when we compared the univariate and multivariate approach was that none of our

a priori ROIs appeared to be an important contributor to the multivariate neural signature, even though

all of them (except NAc) were significantly modulated by relief in the univariate analysis. Instead, we

identified a spatially distributed pattern of brain responses that consisted of several small clusters (all <

65 voxels) across the brain. Some of these clusters fell within other important valuation and error-

processing regions in the brain (e.g., OFC, MCC, caudate nucleus). However, all were small (all < 28

voxels) and require further validation in out of sample participants. Still, these data-driven maps suggest

that other regions than our ROIs might have been especially important for the emotional experience of

relief and that examining these multivariate patterns can aid our understanding of emotional relief.

Finally, two limitations of the study need to be addressed. First, by aiming to provide a fine-

grained analysis of the reward PE-properties of human fMRI responses to threat omission, we focused

exclusively on the necessary and sufficient requirements of reward PE signaling, and thereby

disregarded another core aspect of PE: its teaching property. In the EVA task expectancies are instructed

and all learning is explicitly discouraged. As a result, this task assesses PE completely outside of a

learning context. It therefore remains unclear how the PE-signals we observed relate to actual learning.

It could for instance be that the observed responses mainly reflected the surprisingness of the outcome,

independent of subsequent learning. It therefore remains important to study how the activity patterns of

PE-encoding regions are related to expectancy updating in learning paradigms. Furthermore, it would

be interesting to see how the neural omission-PE signals are affected in clinical populations. Second,

although single unit recordings in rodents have revealed clear PE-like phasic increases in the firing of

dopamine cells at the time of threat omission[6–8], one could argue that fMRI measurements cannot capture

the same sub-second response. Indeed, it is generally difficult to disentangle prediction *error* responses

from the immediately preceding *prediction responses* in fMRI paradigms[48]. Nevertheless, there was no

multicollinearity between anticipation and omission regressors in the first-level GLMs (Variance

Inflation Factor, VIF < 4), making it unlikely that the omission responses purely represented

anticipation. Still, because of the slower timescale of fMRI measurements, we cannot conclusively

dismiss the alternative interpretation that we assessed (part of) expectancy instead.

In conclusion, by violating instructions about the probability and intensity of a potentially

painful stimulation, we found widespread activations in the reward and salience pathways that were

furthermore related to PE-like feelings of pleasurable relief. But, more importantly, we showed for the

first time in humans that unexpected threat omission triggered VTA/SN activations that partially met

the necessary and sufficient criteria of a positive reward PE signal. In doing so, we provided an important

missing link for the human translation of the reward-like threat omission processing in rodents.

440   **Methods**

Participants

Thirty-one healthy volunteers between the ages of 18 and 25 (mean = 20.65, 19 females) were recruited

to participate in our study. All were right-handed and non-smoking, and declared to be free of any

serious medical or psychiatric disorder and regular medication use (see Supplementary Notes 1 & 2 and

Supplementary Table 1 & 2 for an overview of the exclusion criteria we employed, the sample size

rationale and the resulting study sample). Upon their enrollment, participants were furthermore asked to

refrain from consuming any alcohol and/or caffeine and exerting any recreational physical exercise in

the 24 hours before the scan session. The study was approved by the Ethical committee UZ/KU Leuven

(S63852). All participants provided written informed consent and received either partial course credits

or a monetary compensation for their participation.

Stimuli

Expectancy Violation Assessment (EVA) Task. The task was an fMRI adaptation of the

previously validated EVA task[29] and was programmed in affect5 software[49]. In this task, probability

(0%, 25%, 50%, 75%, 100%) and intensity (weak, moderate, strong) information of an upcoming

electrical stimulation to the wrist was presented on each trial in the upper left and right corner of the

screen respectively. A countdown clock, visualized as a receding bar, indicated the exact moment of

stimulation or omission. Responses to the omissions/stimulations were measured.

Electrical Stimulation. The electrical stimulation consisted of a single 2 ms electro-cutaneous

pulse, generated by a Digitimer DS8R Bipolar Constant Current Stimulator (Digitimer Ltd, Welwyn

Garden City, UK), and delivered via two MR-compatible EL509 electrodes (Biopac Systems, Goleta,

CA, USA). Electrodes were filled with Isotonic recording gel (Gel 101; Biopac Systems, Goleta, CA,

USA), and were attached to the right wrist. To match the instructions, a total of three intensities were

individually selected at the start of the scanning session. The weak stimulus was calibrated to be "mildly

uncomfortable", the moderate stimulus to be "very uncomfortable, but not painful", and the strong

stimulus to be "significantly painful, but tolerable". Selected Weak ($M = 9.35$, $SD = 4.65$), moderate (M

$= 16.26$ mA, $SD = 8.13$), and strong ($M = 34.26$, $SD = 19.53$) differed significantly (Friedman test: $\chi2(2)$

$= 62$, $p < .001$, W = 1, all Bonferroni-Holm corrected pairwise comparisons, $p < .001$).

Experimental Procedure

Participants were invited to the lab for an intake session, during which exclusion criteria were extensively checked, experimental procedures were explained, and informed consent was obtained. All included participants then filled out the questionnaire battery (see Supplementary Table 2) and were familiarized with the task and rating scales.

At the start of the scanning session, participants were fitted with skin conductance and stimulation electrodes and stimulation intensities were calibrated using a standard workup procedure (for details, see Supplementary Note 3). Participants were then prepared for the scanner and task instructions were repeated. It was emphasized that all trials in the EVA task were independent of one another, meaning that the presence/absence of stimulations on previous trials could not predict the presence/absence of stimulation on future trials. Finally, the selected intensities were presented again and recalibrated if necessary.

In total, the EVA task comprised 72 trials (48 omission trials), divided equally over four runs of 18 trials/run (for a schematic overview of the trial numbers and types, see Supplementary Figure 1). Since we were mainly interested in how omissions of threat are processed, we wanted to maximize and balance the number of omission trials across the different probability and intensity levels, while also keeping the total number of presentations per probability and intensity instruction constant. Therefore, we crossed all non-0% probability levels (25, 50, 75, 100) with all intensity levels (weak, moderate, strong) (12 trials). The three 100% trials were always followed by the stimulation of the instructed intensity, while stimulations were omitted in the remaining nine trials. Six additional trials were intermixed in each run: Three 0% omission trials with the information that no electrical stimulation would follow (akin to 0% Probability information, but without any Intensity information as it does not apply); and three trials from the Probability x Intensity matrix that were followed by electrical stimulation (across the four runs, each Probability x Intensity combination was paired at least once, and at most twice with the electrical stimulation). Note that, based on previous research, we did not expect the inconsistency between the instructed and perceived reinforcement rate to have a negative effect on

496   the Probability manipulation (see Supplementary Note 4). Within each run trials were presented in a

497   pseudo-random order, with at most two trials with the same intensity or probability as the previous trial.

498   All trials started with the Probability and Intensity instructions for 1 second, followed by the

addition of the countdown bar to the middle of the screen, counting down for 3 to 7 seconds (see Fig.

1A). Then, the screen cleared and the electrical stimulation was either delivered or omitted. Following

a delay of 4 to 8 seconds (during which skin conductance and BOLD responses to the omission were

measured), the rating scale appeared (probing shock-unpleasantness on shock trials and relief-

pleasantness on omission trials). The scale remained on the screen for 8 seconds or until the participant

responded, followed by an intertrial interval between 4 and 7 seconds during which only a fixation cross

was shown. Note that all phases in the trial were jittered (duration countdown clock, duration outcome

window, duration intertrial interval). After the last run, participants were asked some control questions

regarding the intensity and probability instructions. Specifically, they were asked how much effort they

would exert to prevent future weak/moderate/strong stimulation (from 0 "no effort" to 100 "a lot of

effort"); and to estimate how many stimulations they thought they received following each probability

instruction.

Subjective ratings

513   Relief-pleasantness and shock-unpleasantness were probed on omission-and shock-trials using

514   Visual Analogue Scales (VAS) ranging from 0 (neutral) to 100 (very pleasant/unpleasant).

516   **Skin Conductance Responses (SCR)**

517   Fluctuations in skin conductance were measured between two disposable, EL509 electrodes

filled with Isotonic recording gel (Gel 101; Biopac Systems, Goleta, CA, USA) that were attached to

the hypothenar palm of the left hand. Data were recorded continuously at a 1000 Hz sampling rate via a

Biopac MP160 System (Biopac Systems, Goleta, CA, USA), and Acqknowledge software (version 5.0).

Raw data were low-passed filtered at 5 Hz (Butterworth, zerophased) and downsampled to 100 Hz in

Matlab (version R2020b), after which they were entered into a continuous decomposition analysis

(CDA) with two optimization runs (Ledalab, version 3.4.9[50]). Skin conductance responses (SCR) were

scored as the time integral of the deconvoluted phasic activity (integrated SCR) within response

windows of 1-4 sec after the onset (anticipatory SCR) and the offset (omission/stimulation SCR) of the

countdown clock. Above-threshold responses (>0.01 μS) were square root transformed to reduce the

skewness of the distribution. $N = 3$ participants had incomplete datasets because of a missing run ($N =$

2) or delayed recording ($N = 1$), and data of $N = 5$ were completely excluded for SCR analyses as a

result of data loss due to technical difficulties ($N = 1$) or because they were identified as anticipation

non-responder (i.e. participant with smaller average SCR to the clock on 100% than on 0% trials) (N =

4). This resulted in a total sample of $N = 26$ for all SCR analyses. For all other analyses, the full data set

($N = 31$) was used (see Supplementary Table 1 for detailed overview).

Statistical Analyses of Ratings and SCR

Rating and SCR data were analyzed in R 4.2.1 (R Core Team, 2022; https://www.R-project.org)

via linear mixed models that were fit using the lme4 package (v1.1.29[51]). All models included within-

subject factors of Intensity and/or Probability and their interaction as fixed effect and a subject-specific

intercept as random effect. The intensity factor always contained 3 levels (weak, moderate, strong). The

Probability factor depended on the outcome variable. For models of relief and omission SCR that looked

at Probability as well as Intensity, Probability had 3 levels (25%, 50%, 75%); but for models of relief

and omission SCR that only assessed Probability, Probability had 4 levels (0%, 25%, 50%, 75%). To

account for potential changes in the effects of Probability (and Intensity) over time, models included

Run (4 levels: 1, 2, 3, 4) and their interaction with Probability (and Intensity) as regressor of no-interest.

In addition, we controlled for individual differences in the perceived unpleasantness of the stimulation,

by calculating the average of the reported stimulation unpleasantness across the entire task, and entering

the resulting (mean-centered) scores as a between-subjects covariate. Inclusion of both regressors of no-

interest indeed increased model fit (lower AIC). Model assumptions were checked and influential

outliers were identified. Influential outliers were defined as data points above $q_{0.75} + 1.5 * IQR$ or below

$q_{0.25} -1.5*IQR$, with IQR the inter quartile range and $q_{0.25}$ and $q_{0.75}$ corresponding to the first and third

quartile, respectively; and with a cook's distance of greater than 4/ number of data points (calculated via

influence.ME package in R, v0.9-9[52]). To reduce the influence of these data points, they were rescored

to twice the standard deviation from the mean of all data points (corresponding approximately to either

.05 and .95 percentile). If results did not change, we report the model including the original data points.

If results changed, we report the model with adjusted data points.

Main and interaction effects were evaluated using F-tests and p-values that were computed via

type III analysis of Variance using Kenward-Rogers degrees of freedom method of the lmerTest package

(v3.1.3[53]), and omega squared were reported as an unbiased estimate of effect size (calculated via

effectsize package, v0.7.0[54]). All significant effects were followed up with Bonferroni-Holm corrected

pairwise comparisons of the estimated marginal means in order to assess the direction of the effect

(emmeans package, v1.7.5, Length, 2022). Results related to the regressors of no-interest

(Supplementary Figure 5), the anticipatory SCR (Supplementary Figure 3), the post-experimental

questions (Supplementary Figure 2), and the stimulation responses (Supplementary Figure 6) are

reported in the supplementary material for completeness.

fMRI Analyses

MRI Acquisition. MRI data were acquired on a 3 Tesla Philips Achieva scanner, using a 32-

channel head coil, at the Department of Radiology of the University Hospitals Leuven. The four

functional runs (226 volumes each) were recorded using an T2*-weighted echo planar imaging sequence

(60 axial slices; FOV = 224 x 224 mm; in-plane resolution = 2 x2 mm; interslice gap = 0.2 mm; TR =

2000 ms; TE = 30 ms; MB = 2; flip angle = 90°). In addition, a high resolution T1-weighted anatomical

image was acquired for each subject for co-registration and normalization of the EPI data using a MP-

RAGE sequence (182 axial slices; FOV = 256 x 240 mm; in-plane resolution = 1 x 1 mm; TR = shortest,

TE = 4.6 ms , flip angle = 8°); and a short reverse phase functional run (10 volumes) was acquired using

the exact same imaging parameters as the functional runs, but with opposite phase encoding direction. This reverse phase run was used to estimate the B0-nonuniformity map (or fieldmap) to correct for susceptibility distortion. Functional data of one run were missing for $N = 4$ participants as a result of technical difficulties during scanning. Whenever available, rating and SCR data was still included in the analyses.

fMRI Preprocessing. Prior to preprocessing, image quality was visually checked via quality reports of the anatomical and functional images generated through MRIQC[55]. MRI data were then preprocessed using a standard preprocessing pipeline in fMRIPrep 20.2.3[56]. A detailed overview of the preprocessing steps in fMRPrep can be found in Supplementary Methods 2. In line with our preregistration, spikes were identified and defined as volumes having a framewise displacement (FD) exceeding a threshold of 0.9 mm or DVARS exceeding a threshold of 2. No functional run had more than 15% spikes, and hence none of the runs had to be excluded from our analyses based on our preregistered criterium. Afterwards, the functional data were spatially smoothed with a 4 mm FWHM isotropic Gausian kernel within the "Statistical parametric Mapping" software (SPM12; https://www.fil.ion.ucl.ac.uk/spm/).

Subject Level Analysis. Three subject-level general linear models (GLM) were specified. In all models, we concatenated the functional runs and added run-specific intercepts to account for changes over time. The first model investigated the effects of instructed Probability and Intensity on neural omission processing and therefore included separate stick regressors (duration = 0) for omissions of each Probability x Intensity combination (10 regressors), and stimulations (2 regressors: one for non-100% stimulations, one for 100% stimulations), in addition to boxcar regressors (duration = total duration of event) for the instruction (1 regressor) and rating (1 regressor) windows. The second model assessed how omission fMRI data was related to trial-by-trial relief-pleasantness ratings, by only including a single stick regressor for all omissions (1 regressor) and shocks (1 regressor), and boxcar regressors for instructions (1 regressor) and ratings (1 regressor). Z-scored relief-pleasantness ratings were added as parametric modulator for the omission regressor. A final model estimated single-trial omission responses (48 stick regressors), in addition to a single stick regressor for stimulation and boxcar

regressors for instructions (1 regressor) and ratings (1 regressor) and was used for the LASSO-PCR

analyses. Regressors in all models were convolved with a canonical hemodynamic response function

and a high pass filter of 180 s was applied to remove low frequency drift. Additional non-task related

noise was modeled by including nuisance regressors of no-interest for global CSF signal, 24 head motion

regressors (consisting of 6 translation and rotation motion parameters and their derivatives, z-scored;

and their quadratic terms), and dummy spike regressors.

Group Level Univariate Analysis.

Omission processing. To test whether our regions of interest (ROI) were activated by the

unexpected omission of threat, we contrasted all non-0% omissions (unexpected omissions) with 0%

omissions (expected omissions) at subject-level. Mean activity of each ROI was extracted from the

resulting contrast map through marsbar (v0.45[57]), and was entered into group-level (two-sided) one-

sample *t*-tests (per ROI) that were Bonferroni-Holm corrected for the total number of ROIs (4 in the

main analyses, 10 in the secondary analyses) in R.

We then evaluated whether the observed activity qualified as a 'Prediction Error' by applying

an axiomatic testing approach. Specifically, we tested for each ROI if the omission-related activity

increased with increasing expected Intensity (axiom 1) and Probability (axiom 2) of threat, and whether

completely predicted outcomes (0% omission and 100% stimulations) elicited equivalent activation

(axiom 3). For axioms 1 and 2, we extracted mean activity of each ROI from separate omission contrasts

that contrasted each omission type (i.e., all possible Probability x Intensity combinations) separately

with 0% omissions. These were then entered into a LMM in R including Probability (3 levels: 25%,

50%, 75%) and Intensity (3 levels: weak, moderate, strong) as within-subject factors, and a between-

subject covariate of average stimulation unpleasantness (mean-centered) as fixed effects, in addition to

a subject-specific intercept. Note that we did not include Run as regressor of no-interest, as Run effects

were already accounted for by adding run-specific intercepts to the first level models. Main and

interaction effects within each model were followed up with Bonferroni-Holm corrected pairwise

comparisons of the estimated marginal means in order to assess the direction of the effect. Finally, to

fulfill all necessary and sufficient requirements of a prediction error signal, we contrasted completely

predicted omissions (0%) with completely predicted stimulations (100%), as these should trigger

equivalent activation in PE-encoding regions. Given that we would expect equivalent activation, Bayes

Factors were computed using the BayesFactor package in R (v0.9.12-4.4, Morey & Rouder, 2022) to

compare the evidence in favor of alternative and null hypotheses. Larger Bayes Factors indicated more

evidence in favor of the null hypotheses.

Parametric modulation of relief. We then tested if the omission-related activity was correlated to

self-reported relief-pleasantness on a trial-by-trial basis. To this end, we extracted the mean activity

from the modulation contrast for each ROI, and entered these averages in separate one-sample (two-

sided) t-test in R, again correcting for the number of ROIs (4 for main analysis, 10 for secondary

analyses). In a subsequent exploratory analysis, we entered z-scored omission SCR-responses as

parametric modulator to the omission regressor. Results related to this analysis are reported in

Supplementary Figure 13 and Supplementary Table 8.

Neural signature of relief. A LASSO-PCR model (Least Absolute Shrinkage and Selection

Operator-Regularized Principal Component Regression) as implemented in CANlab neuroimaging

analysis tools (see https://canlab.github.io/) was trained using trial-by-trial whole-brain omission-related

activation-maps as predictors, and trial-by-trial relief-pleasantness ratings as outcome (for other

applications of this approach, see e.g.[31,58–60]). The added value of this machine-learning technique is that

relief is predicted across a set of voxels instead of being predicted for each voxel separately (as in

standard univariate regression). Specifically, while each voxel in the activation maps is considered a

predictor of relief, the LASSO-PCR technique uses a combination of Principle Component Analyses

(PCA) and LASSO-regression to (1) group predictive information across individual voxels into larger

components (PCA), and (2) maximize the predictive weight of the most informative components by

shrinking the regression weight of the least informative components to zero (LASSO regression). Here,

we embedded the LASSO-PCR technique within a five-fold cross-validation loop that iteratively trained

a LASSO-PCR model in each loop on a different training and validation dataset, which we then averaged

across loops to obtain the final model. Model performance was then assessed by calculating the Pearson

correlation between reported and model predicted relief. Important features to the signature pattern were

identified using bootstrap tests (5000 samples). Furthermore, the contribution of our ROIs to the model's

performance was assessed using virtual lesion analysis that consisted of repeating the model training,

but excluding voxels within the ROIs and assessing model performance. Note that we also estimated a

neural signature of omission SCR, by applying a LASSO-PCR model to predict omission-related SCR

responses. Results related to these analyses are reported in Supplementary Figure 14 and Supplementary

Table 9.

Regions of Interest (ROI). Our main regions of interest consisted of key regions of the reward and

(relief) valuation pathways such as the Ventral Tegmental Area (VTA) /Substantia Nigra (SN), Nucleus

Accumbens (NAC), ventral Putamen, and ventromedial prefrontal cortex (vmPFC). The VTA/SN mask

was obtained from Esser and colleagues[17], but was originally defined by Bunzeck and Düzel (2006)[61].

The ventral Putamen ROI was defined as a 6 mm sphere centered around the peak voxel (MNI

coordinates: -32,8-6) in the left ventral putamen identified by Raczka et al. (2011)[14], as in Thiele et al.

(2021)[15]. However, we overlayed this sphere with a Putamen mask obtained from a high-resolution

anatomical atlas for subcortical nuclei defined by Pauli et al. (2018)[62] to assure the mask was restricted

to voxels of the putamen and did not extend to the adjacent anterior Insula. Likewise, a bilateral NAc

mask was obtained from the Pauli et al. (2018)[62] atlas. The vmPFC mask was obtained by selecting

specific parcels from the atlas vmPFC cortex of AFNI (area 14, 32, 24, bilateral). Since we consider the

pleasantness of relief from omission of a threat a type of reward, the selection of parcels was made on

the base of an activation likelihood estimation (ALE) meta-analysis of 87 studies (1452 subjects)

comparing the brain responses to monetary, erotic and food reward outcomes[63].

In addition to our main ROIs, we specified a wider secondary mask that extended our main ROIs

with additional regions that have previously been associated to reward, pain and PE processing (such as

the wider striatum, including the nucleus caudatus, and putamen; midbrain nuclei, including the

periaqueductal gray (PAG), habenula, and red nucleus; limbic structures, including the amygdala and

hippocampus; midline thalamus and cortical regions, including the anterior insula (aINS), medial

orbitofrontal (OFC), dorsomedial prefrontal (dmPFC) and anterior cingulate (ACC) cortices). Masks for

these regions were obtained from CANlab combined atlas (2018) (see https://canlab.github.io/), and

Pauli atlas[62]. Whole brain voxel-wise analyses were restricted to the grey matter mask sparse (from

CANlab tools), extended with midbrain voxels (including VTA/SN, RN, habenula). All masks were

registered to functional space before analyses and are freely available online at OSF

(https://osf.io/ywpks/).

Data availability

Raw data files are not freely accessible because one participant did not consent to publish their

anonymized data online. Data are available on request by contacting ALW or BV.

Code availability

Analyses code and anatomical masks are freely available at OSF (https://osf.io/ywpks/).

Acknowledgements

This research was supported by FWO project grant G078929N (National Research Fund Flanders,

Belgium) and C1 project grant C16/19/002 (KU Leuven, Belgium) awarded to BV. ALW was supported

by the Internal Funds KU Leuven. LVO is a research professor funded by the KU Leuven Special

Research Fund. We would like to thank Mathijs Franssen and Ronald Peeters for their technical support;

and Silvia Papalini, Anraoi Rooney, Lieselotte Claes and Anamarija Banjac for their assistance during

fMRI data collection.

Author Contribution

AW: Conceptualization, methodology, formal analysis, investigation, data curation, writing – original

draft, visualization

LVO: Conceptualization, methodology, formal analysis, writing – review & editing

BV: Conceptualization, methodology, data curation, supervision, writing – original draft, funding

acquisition

705   **References**

1.   Deutsch, R., Smith, K. J. M., Kordts-Freudinger, R. & Reichardt, R. How absent negativity
relates to affect and motivation: an integrative relief model. *Front. Psychol.* **6**, (2015).

2.   Rescorla, R. & Wagner, A. A theory of Pavlovian conditioning: The effectiveness of
reinforcement and non-reinforcement. *Class. Cond. Curr. Res. Theory* (1972).

3.   Beckers, T. *et al.* Understanding clinical fear and anxiety through the lens of human fear
conditioning. *Nat. Rev. Psychol.* **2**, 233–245 (2023).

4.   Schultz, W. Dopamine reward prediction-error signalling: A two-component response. *Nat.*
Rev. Neurosci. **17**, 183–195 (2016).

5.   Watabe-Uchida, M., Eshel, N. & Uchida, N. Neural circuitry of reward prediction error. *Annu.*
Rev. Neurosci. **40**, 373–394 (2017).

6.   Luo, R. *et al.* A dopaminergic switch for fear to safety transitions. *Nat. Commun.* **9**, 2483
(2018).

7.   Salinas-Hernández, X. I. *et al.* Dopamine neurons drive fear extinction learning by signaling
the omission of expected aversive outcomes. *Elife* **7**, e38818 (2018).

8.   Cai, L. X. *et al.* Distinct signals in medial and lateral VTA dopamine neurons modulate fear
extinction at different times. *Elife* **9**, e54936 (2020).

9.   de Jong, J. W. *et al.* A neural circuit mechanism for encoding aversive stimuli in the
mesolimbic dopamine system. *Neuron* **101**, 133–151 (2019).

10. Badrinarayan, A. *et al.* Aversive stimuli differentially modulate real-time dopamine
transmission dynamics within the nucleus accumbens core and shell. *J. Neurosci.* **32**, 15779–
15790 (2012).

11. Yau, J. O. Y. & McNally, G. P. Brain mechanisms controlling pavlovian fear conditioning. *J.*
Exp. Psychol. Anim. Learn. Cogn. **44**, 341–357 (2018).

12. Oleson, E. B., Gentry, R. N., Chioma, V. C. & Cheer, J. F. Subsecond dopamine release in the
nucleus accumbens predicts conditioned punishment and its successful avoidance. *J. Neurosci.*
32, 14804–14808 (2012).

13. Wenzel, J. M. *et al.* Phasic dopamine signals in the nucleus accumbens that cause active
avoidance require endocannabinoid mobilization in the midbrain. *Curr. Biol.* **28**, 1392–1404
(2018).

14. Raczka, K. A. *et al.* Empirical support for an involvement of the mesostriatal dopamine system
in human fear extinction. *Transl. Psychiatry* **1**, e12 (2011).

15. Thiele, M., Yuen, K. S. L., Gerlicher, A. V. M. & Kalisch, R. A ventral striatal prediction error
signal in human fear extinction learning. *Neuroimage* **229**, 117709 (2021).

16. Lange, I. *et al.* Neural responses during extinction learning predict exposure therapy outcome
in phobia: results from a randomized-controlled trial. *Neuropsychopharmacology* **45**, 534–541
(2020).

17. Esser, R., Korn, C. W., Ganzer, F. & Haaker, J. L-DOPA modulates activity in the vmPFC,
nucleus accumbens, and VTA during threat extinction learning in humans. *Elife* **10**, e65280
(2021).

18. Leknes, S., Lee, M., Berna, C., Andersson, J. & Tracey, I. Relief as a reward: Hedonic and
neural responses to safety from pain. *PLoS One* **6**, e17870 (2011).

19. Boeke, E. A., Moscarello, J. M., LeDoux, J. E., Phelps, E. A. & Hartley, C. A. Active
avoidance: Neural mechanisms and attenuation of pavlovian conditioned responding. *J.*
Neurosci. **37**, 4808–4818 (2017).

20. Papalini, S., Beckers, T. & Vervliet, B. Dopamine: from prediction error to psychotherapy.
Transl. Psychiatry **10**, 164 (2020).

21. Kalisch, R., Gerlicher, A. M. V. & Duvarci, S. A dopaminergic basis for fear extinction.
Trends Cogn. Sci. **23**, 274–277 (2019).

22. Vervliet, B., Lange, I. & Milad, M. R. Temporal dynamics of relief in avoidance conditioning
and fear extinction: Experimental validation and clinical relevance. *Behav. Res. Ther.* **96**, 66–
78 (2017).

23. Papalini, S., Ashoori, M., Zaman, J., Beckers, T. & Vervliet, B. The role of context in
persistent avoidance and the predictive value of relief. *Behav. Res. Ther.* **138**, 103816 (2021).

24. Caplin, A. & Dean, M. Axiomatic methods, dopamine and reward prediction error. *Curr. Opin.*
Neurobiol. **18**, 197–202 (2008).

25. Rutledge, R. B., Dean, M., Caplin, A. & Glimcher, P. W. Testing the reward prediction error
hypothesis with an axiomatic model. *J. Neurosci.* **30**, 13525–13536 (2010).

26. Jepma, M., Roy, M., Ramlakhan, K., van Velzen, M. & Dahan, A. Different brain systems
support learning from received and avoided pain during human pain-avoidance learning. *Elife*
11, e74149 (2022).

27. Roy, M. *et al.* Representation of aversive prediction errors in the human periaqueductal gray.
Nat. Neurosci. **17**, 1607–1612 (2014).

28.   Ojala, K. E., Tzovara, A., Poser, B. A., Lutti, A. & Bach, D. R. Asymmetric representation of
aversive prediction errors in Pavlovian threat conditioning. *Neuroimage* **263**, 119579 (2022).

29.   Willems, A. L. & Vervliet, B. When nothing matters : Assessing markers of expectancy
violation during omissions of threat. *Behav. Res. Ther.* **136**, 103764 (2021).

30.   Fullana, M. A. *et al.* Neural signatures of human fear conditioning: An updated and extended
meta-analysis of fMRI studies. *Mol. Psychiatry* **21**, 500–508 (2016).

31.   Wager, T. D. *et al.* An fMRI-based neurologic signature of physical pain. *N. Engl. J. Med.* **368**,
1388–1397 (2013).

32.   D'Ardenne, K., McClure, S. M., Nystrom, L. E. & Cohen, J. D. BOLD responses reflecting
dopaminergic signals in the human ventral tegmental area. *Science (80-.).* **319**, 1264–1267
(2008).

33.   Zaghloul, K. A. *et al.* Human substantia nigra neurons encode unexpected financial rewards.
Science (80-.). **323**, 1496–1499 (2009).

34.   Gerlicher, A. M. V, Tüscher, O. & Kalisch, R. Dopamine-dependent prefrontal reactivations
explain long-term benefit of fear extinction. *Nat. Commun.* **9**, 4294 (2018).

35.   Kim, H., Shimojo, S. & O'Doherty, J. P. Is avoiding an aversive outcome rewarding? Neural
substrates of avoidance learning in the human brain. *PLOS Biol.* **4**, e233 (2006).

36.   Berridge, K. C. & Kringelbach, M. L. Pleasure systems in the brain. *Neuron* **86**, 646–664
(2015).

37.   Düzel, E. *et al.* Functional imaging of the human dopaminergic midbrain. *Trends Neurosci.* **32**,
321–328 (2009).

38.   Zhang, Y., Larcher, K. M. H., Misic, B. & Dagher, A. Anatomical and functional organization
of the human substantia Nigra and its connections. *Elife* **6**, e26653 (2017).

39.   Root, D. H. *et al.* Glutamate neurons are intermixed with midbrain dopamine neurons in
nonhuman primates and humans. *Sci. Rep.* **6**, 30615 (2016).

40.   Diederen, K. M. J. & Fletcher, P. C. Dopamine, prediction error and beyond. *Neuroscientist* **27**,
30–46 (2021).

41.   Matsumoto, M. & Hikosaka, O. Two types of dopamine neuron distinctly convey positive and
negative motivational signals. *Nature* **459**, 837–841 (2009).

42.   Pauli, W. M. *et al.* Distinct contributions of ventromedial and dorsolateral subregions of the
human substantia nigra to appetitive and aversive learning. *J. Neurosci.* **35**, 14220–14233

(2015).

43.   Hart, A. S., Rutledge, R. B., Glimcher, P. W. & Phillips, P. E. M. Phasic dopamine release in
the rat nucleus accumbens symmetrically encodes a reward prediction error term. *J. Neurosci.*
34, 698–704 (2014).

44.   Garrison, J., Erdeniz, B. & Done, J. Prediction error in reinforcement learning: A meta-analysis
of neuroimaging studies. *Neurosci. Biobehav. Rev.* **37**, 1297–1310 (2013).

45.   Chase, H. W., Kumar, P., Eickhoff, S. B. & Dombrovski, A. Y. Reinforcement learning models
and their neural correlates: An activation likelihood estimation meta-analysis. *Cogn. Affect.*
Behav. Neurosci. **15**, 435–459 (2015).

46.   Raichle, M. E. The brain's default mode network. *Annu. Rev. Neurosci.* **38**, 433–447 (2015).

47.   Seeley, W. W. The salience network: A neural system for perceiving and responding to
homeostatic demands. *J. Neurosci.* **39**, 9878–9882 (2019).

48.   Linnman, C., Rougemont-Bücking, A., Beucke, J. C., Zeffiro, T. A. & Milad, M. R.
Unconditioned responses and functional fear networks in human classical conditioning. *Behav.*
Brain Res. **221**, 237–245 (2011).

49.   Spruyt, A., Clarysse, J., Vansteenwegen, D., Baeyens, F. & Hermans, D. Affect 4.0: A free
software package for implementing psychological and psychophysiological experiments. *Exp.*
Psychol. **57**, 36–45 (2009).

50.   Benedek, M. & Kaernbach, C. A continuous measure of phasic electrodermal activity. *J.*
Neurosci. Methods **190**, 80–91 (2010).

51.   Bates, D., Maechler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models using
lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

52.   Nieuwenhuis, R., te Grotenhuis, M. & Pelzer, B. influence.ME: Tools for detecting influential
data in mixed effects models. *R J.* **4**, 38–47 (2012).

53.   Kuznetsova, A., Brockhoff, P. B. & B, C. R. H. lmerTEst package: Tests in linear mixed
effects models. *J. Stat. Softw.* **82**, 1–26 (2017).

54.   Ben-Shachar, M., Lüdecke, D. & Makowski, D. effectsize: Estimation of effect size indices
and standardized parameters. *J. Open Source Softw.* **5**, 2815 (2020).

55.   Esteban, O. *et al.* MRIQC: Advancing the automatic prediction of image quality in MRI from
unseen sites. *PLoS One* **12**, e0184661 (2017).

56.   Esteban, O. *et al.* fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat. Methods*

16, 111–116 (2019).

57.   Brett, M., Anton, J.-L., Valabregue, R. & Poline, J.-B. Region of interest analysis using an
SPM toolbox. Presented at the 8th International Conference on Functional Mapping of the
Human Brain, June 2-6, 2002, Sendai, Japan at https://marsbar-toolbox.github.io/. in (2002).

58.   Wager, T. D., Atlas, L. Y., Leotti, L. A. & Rilling, J. K. Predicting individual differences in
placebo analgesia: Contributions of brain activity during anticipation and pain experience. *J.*
Neurosci. **31**, 439–452 (2011).

59.   Speer, S. P. H. *et al.* A multivariate brain signature for reward. *Neuroimage* **271**, 119990
(2023).

60.   Zhou, F. *et al.* A distributed fMRI-based signature for the subjective experience of fear. *Nat.*
Commun. **12**, 6643 (2021).

61.   Bunzeck, N. & Düzel, E. Absolute coding of stimulus novelty in the human substantia
nigra/VTA. *Neuron* **51**, 369–379 (2006).

62.   Pauli, W. M., Nili, A. N. & Tyszka, J. M. A high-resolution probabilistic in vivo atlas of
human subcortical brain nuclei. *Sci. Data* **5**, 180063 (2018).

63.   Sescousse, G., Caldú, X., Segura, B. & Dreher, J. C. Processing of primary and secondary
rewards: A quantitative meta-analysis and review of human functional neuroimaging studies.
Neurosci. Biobehav. Rev. **37**, 681–696 (2013).