Complete genome-centric analysis reveals functional divergence and adaptive

evolution of uncultured bacteria in anammox community

Yi-Cheng Wang,[1,†] Yanping Mao,[2,†] Hui-Min Fu,[1,3] Jin Wang,[1] Xun Weng,[1] Zi-Hao

Liu,[1] Xiao-Wei Xu,[1] Peng Yan,[1] Fang Fang,[1] Jin-Song Guo,[1] Yu Shen[3] and You-Peng

Chen[1,*]

[1] Key Laboratory of the Three Gorges Reservoir Region's Eco-Environments of MOE,

Chongqing University, Chongqing 400045, China

[2] College of Chemistry and Environmental Engineering, Shenzhen University,

Shenzhen 518071, Guangdong, China

[3] National Research Base of Intelligent Manufacturing Service, Chongqing

Technology and Business University, Chongqing 400067, China

[*] Corresponding author: You-Peng Chen; Key Laboratory of the Three Gorges

Reservoir Region's Eco-Environments of MOE, Chongqing University, Chongqing

400045, China; E-mail address: ypchen@cqu.edu.cn.

[†] These authors contributed equally to this work.

Running title: Complete genome-centric analysis of anammox community

Abstract

Anaerobic ammonium-oxidation (anammox) bacteria play a crucial role in global nitrogen cycling and wastewater nitrogen removal, but they depend on mutualistic symbiosis with various other microorganisms for their survival. Although shotgun metagenomics based on short reads has been widely used in anammox research, metagenome-assembled genomes (MAGs) are often discontinuous and highly contaminated, which limits in-depth analyses of anammox communities. Here, for the first time, we performed Pacific Biosciences high-fidelity (HiFi) sequencing on the anammox granule sludge sample, and obtained 30 accurate and complete metagenome-assembled genomes (cMAGs). These cMAGs were obtained by selecting high-quality circular contigs from initial assemblies of long reads generated by HiFi sequencing, eliminating the need for Illumina short reads, binning, and reassembly. cMAG-centric analysis revealed divergences in general and nitrogen metabolism among members of the anammox community. Furthermore, we identified mobile genetic elements (MGEs) and putative horizontal gene transfer (HGT) events within these cMAGs to explore the adaptive evolution of the community. The results suggest that MGEs and HGT events, particularly transposons containing *tnpA* in anammox bacteria, might play important roles in the adaptive evolution of this anammox community. The cMAGs generated in the present study could be used to establish of a comprehensive database for anammox bacteria and associated microorganisms. Our findings highlight the advantages of HiFi sequencing for the studies of complex cultures such as anammox communities and advance our understanding of anammox communities.

Keywords: anammox community; adaptive evolution; mobile genetic elements; horizontal gene transfer; HiFi metagenomic sequencing

Introduction

Anaerobic ammonium-oxidation (anammox) plays a key role in the global nitrogen cycle and is a cost-effective technique for treating ammonium-rich wastewaters due to no external carbon source requirement, lower oxygen consumption and sludge production (Peeters and van Niftrik, 2019; Adams et al., 2022). Approximately 100 anammox wastewater treatment plants are currently operating worldwide (Lackner et al., 2014). Anammox bacteria affiliated with the phylum Planctomycetota (also known as Planctomycetes) are responsible for catalyzing anammox process, in which anammox bacteria utilize nitrite as electron acceptor to oxidize ammonium to dinitrogen gas (N_2) under anoxic conditions (Kartal et al., 2013). Although anammox bacteria have been discovered for over 20 years, and substantial efforts have been made to isolate them, yet pure cultures of these bacteria are still unavailable to date (Peeters and van Niftrik, 2019; Kuenen, 2020). It is well known that anammox bacteria hardly survive alone and share symbiotic relationships with various other microorganisms affiliated with the phyla Acidobacteriota, Armatimonadota, Bacteroidota, Chloroflexota, Proteobacteria, Patescibacteria, etc. (Bhattacharjee et al., 2017; Lawson et al., 2017; Feng et al., 2018; Zhao et al., 2018a). The co-existing bacteria can provide anammox bacteria with growth factors or stable nitrite under specific C/N ratios, and prevent the inhibition of anammox bacteria by organic matter (Zhao et al., 2018b; Cao et al., 2021). Therefore, comprehensive analysis of the diversity, functional divergence, and adaptive evolution of anammox communities can contribute to advance our understanding of anammox systems and promote the widespread application of this energy-efficient treatment process.

Shotgun metagenomics employing next-generation sequencing (NGS) has been widely used in studies of anammox systems, and the genomes of different anammox bacteria and associated bacteria have been successfully retrieved from samples from

various habitats (Speth et al., 2016; Lawson et al., 2017; Suarez et al., 2022; Yang et

al., 2022). However, due to the short reads and PCR requirement of NGS represented

by the Illumina platform, leading to some shortcomings such as fragmented assembly

and high genome contamination (Meziti et al., 2021). The recovered

metagenome-assembled genomes (MAGs) generated by NGS are generally

discontinuous because of repetitive, mobile, and conserved sequences, which limits

their application in further analyses (Yuan et al., 2015; Sangwan et al., 2016; Meziti et

al., 2019; Maguire et al., 2020). Long-read third-generation sequencing (TGS)

developed by Pacific Biosciences (PacBio) and Oxford Nanopore Technologies (ONT)

is a promising approach for addressing the above problems. The complete circular

genomes of anammox bacteria from highly enriched cultures of *Candidatus* Kuenenia

stuttgartiensis (> 87% of the total biomass) were previous recovered using only long

reads generated by PacBio RS II sequencing (Frank et al., 2018; Ding and Adrian,

2020). Additionally, Liu et al. (2020) employed a combination of Illumina and ONT

sequencing to develop an iterative hybrid assembly (IHA) method, which enabled the

retrieval of eight circular genomes from one sample of a partial nitrification anammox

reactor. However, nucleotide accuracy and assembly bubbles were not reported for

these eight circular genomes. Consequently, a simpler and more efficient approach for

recovering numerous accurate circular genomes from anammox systems is currently

lacking. Furthermore, whether TGS can be employed alone to retrieve accurate and

complete prokaryotic genomes, beyond those of dominant species, from anammox

communities remains unclear.

Recently, PacBio high-fidelity (HiFi) sequencing, which can yield high-accuracy

long reads (accuracy > 99.5% and average read length of 10-25 kb) (Hon et al., 2020),

has become popular for the assembly of plant and animal genomes (Hon et al., 2020;

Nurk et al., 2022), as well as the analysis of complex microbiomes (from human gut,

sheep fecal, chicken cecum, and saline lake sediment samples) (Bickhart et al., 2022;

Feng et al., 2022; Kim et al., 2022; Tao et al., 2023). HiFi sequencing has been shown

to greatly improve the quality of metagenome assembly (Tao et al., 2023). Most

importantly, HiFi sequencing enables the acquisition of accurate and complete

prokaryotic genomes from complex microbiomes, without the need for Illumina short

reads, binning, and reassembly. Kim et al. (2022) assembled the HiFi reads derived

from five human gut samples, and obtained 102 complete metagenome-assembly

genomes (cMAGs). One of these cMAGs is almost identical to the genome of pure

culture, with an average nucleotide identity (ANI) value exceeding 99.99%. Taken

together, HiFi sequencing provides a highly suitable approach for the accurate and

comprehensive analysis of environmental microbiomes, such as anammox

communities. The utilization of HiFi sequencing enables the retrieval of more reliable

cMAGs compared with MAGs. These cMAGs can serve as a valuable resource for

subsequent downstream analyses, including the identification of novel species,

exploration of genes associated with specific pathways (Parks et al., 2017; Galambos

et al., 2019), investigation of functional divergence, and study of the adaptive

evolutionary mechanisms of prokaryotes (Douglas and Langille, 2019).

To address the aforementioned research gaps and facilitate in-depth analysis of

anammox communities, we performed HiFi sequencing on one granular sludge

sample obtained from a lab-scale anammox reactor for the first time. The long-read

metagenome obtained was *de novo* assembled, and non-redundant 30 cMAGs,

including a genome of a novel anammox species and three genomes from novel

families, were filtered out without any binning and reassembly processes. Using these

reliable cMAGs, functional divergence and adaptive evolution within the anammox

community were further explored. Overall, the results of this study provide a valuable

workflow for analyzing complex aquatic microbiomes as well as new insights into

bacterial interactions and evolution within the anammox community.

Materials and methods

Sample collection and DNA extraction

The anammox granular sludge sample was collected from a lab-scale expanded

granular sludge blanket (EGSB) reactor (**Fig. S1**), which has been operated for two

years. The composition of the culture medium is shown in **Table S1** and **Table S2**.

The EGSB reactor was initially inoculated with commercially purchased anammox

granular sludge, and it was used as the parent reactor to generate an ongoing supply of

anammox granular sludge in our laboratory. Total genomic DNA was extracted using

the E.Z.N.A.® Soil DNA Kit (Omega Bio-Tek, USA) according to the manufacturer's

protocol. The purity and quantity of genomic DNA were assessed using the NanoDrop

2000 (ThermoFisher Scientific, USA) and the Qubit 3.0 Fluorometer (Invitrogen,

USA) respectively, and its integrity was verified using agarose gel electrophoresis.

The DNA sample with a high molecular weight and sufficient quantity was used in

subsequent procedures.

Metagenomic sequencing

Metagenomic sequencing libraries were prepared and sequenced by Biozeron

Biotechnology Co. Ltd. (Shanghai, China). For PacBio HiFi sequencing, 5 μg of DNA

was used to construct a SMRTbell library with the SMRTbell Prep Kit (Pacific

Biosciences, USA) according to the manufacturer's protocol. Damaged

double-stranded DNA in the initial DNA sample was repaired using the PreCR®

Repair Mix Kit (New England Biolabs, USA) before library construction. The

repaired DNA was then size-selected to obtain molecules larger than 3 kb. The

SMRTbell library was sequenced on the PacBio Sequel IIe platform using a single

SMRT 8M cell. HiFi reads were generated with the CCS module (–min-passes = 3,

and –min-rq = 0.99) of SMRT Link v11.0 (Pacific Biosciences, USA). For Illumina

sequencing, the sequencing library was constructed with a fragment length of ~450 bp

and sequenced using the PE150 strategy on the Illumina NovaSeq 6000 platform.

Basic information of both the HiFi and Illumina reads is shown in **Table S3** and **Table**

S4, respectively.

De novo **assembly and quality assessment**

De novo assembly of HiFi reads was performed using three different assemblers: hifiasm-meta v0.3-r063.2 (Feng et al., 2022), Flye v2.9.1-b1780 (Kolmogorov et al., 2020), and Canu v2.2 (Nurk et al., 2020). Both the hifiasm-meta and Flye assembly were conducted using the recommended PacBio HiFi meta mode. The Canu assembly was conducted using the PacBio HiFi mode with recommended parameters. The assembly quality was assessed using QUAST v5.2.0 (Gurevich et al., 2013), and the results are shown in **Table S5**.

Filtering, taxonomic classification, and functional annotation of cMAGs

The HiFi cMAGs were filtered from initial assemblies using an adapted workflow published before (Kim et al., 2022). The workflow is briefly described as follows: (1) the circular contigs without assembly bubbles (assessed by BubbleGun v1.1.6 (Dabbaghie et al., 2022) as required) and repeats were filtered from the three assemblies; (2) contigs with sequence lengths shorter than 100 kb were removed; (3) the completeness and taxonomic classification of remaining contigs were assessed using GTDB-Tk v2.1.1 (Parks et al., 2022), and contigs with bacterial marker genes (bac120) less than 80 were removed (at this step, no archaea genome was found); (4) the SSU rRNA of the remaining contigs was predicted using barrnap v0.9 (https://github.com/tseemann/barrnap), and contigs missing 5S, 16S or 23S rRNA were removed; (5) the tRNA of remaining contigs was identified using tRNAscan-SE v2.0.11 (Chan et al., 2021), and contigs with tRNA types fewer than 18 were filtered out; and (6) ANI values of contigs were calculated using FastANI v1.33 (Jain et al., 2018). Redundant contigs were those with ANI values ≥ 99% and alignment percentages ≥ 95%, and the contigs with the maximal sum of ANI values were selected as the representative sequences. The contigs obtained through above six steps were considered non-redundant cMAGs. The cMAG abundances represented by transcripts per million (TPM) were calculated using salmon v1.10.1 (Patro et al., 2017), by mapping Illumina reads onto these genomes.

Gene prediction of cMAGs was performed using Prodigal v2.6.3 (Hyatt et al.,

2010), and the predicted genes were comprehensively annotated using MEBS v1.2

(De Anda et al., 2017), eggNOG database v5.0.2 (e-value < 1e-5) (Huerta-Cepas et al.,

2019), NCycDB (e-value < 1e-5, identity > 40%, and coverage > 40%) (Tu et al.,

2019), and KofamKOALA (Aramaki et al., 2020). The gene abundances represented

by TPM were also calculated using salmon v1.10.1 (Patro et al., 2017).

Identification of mobile genetic elements (MGEs) and horizontal gene transfer

(HGT)

MGEs of cMAGs were identified by alignment (--escore = 1e-6, --pidentvalue = 80,

and –queryscore = 75) against mobileOG-db beatrix-1.6 (Brown et al., 2022). The

HGT events among all cMAGs were identified using MetaCHIP v1.10.12 (Song et al.,

2019) with default parameters. All putative transferred genes were further annotated

by alignment (e-value < 1e-5) against both the eggNOG database v5.0.2

(Huerta-Cepas et al., 2019) and NCBI non-redundant (NR) database.

Comparative genomic analyses

Reference genomes (**Table S6**) of 26 anammox bacteria in eight genera (*Ca.*

Anammoxibacter, *Ca.* Anammoxoglobus, *Ca.* Anammoximicrobium, *Ca.* Brocadia,

Ca. Jettenia, *Ca.* Kuenenia, *Ca.* Loosdrechtia, and *Ca.* Scalindua) were downloaded

from both the NCBI and EMBL-EBI databases. JSpeciesWS (Richter et al., 2015) was

used to calculate the ANI values (based on blast) between cMAGs representing

anammox bacteria and other reference genomes of *Ca.* Jettenia. DNA-DNA

hybridization (DDH) *in silico* was performed using GGDC v3.0 (Meier-Kolthoff et al.,

2022) with recommended parameters. All reference anammox bacterial genomes were

also comprehensively annotated like cMAGs.

Results

Overview of cMAGs from an anammox community

Through HiFi sequencing and assembly, a total of 541 circular contigs were obtained

from one anammox granular sludge sample (**Fig. 1**). These contigs were filtered based

on biological features and ANI values, yielding 30 non-redundant contigs known as

cMAGs (**Fig. 1**). The number of cMAGs from the assembly of hifiasm-meta, Canu

and Flye was 19, 7, and 4, respectively. The size of these cMAGs ranged from 0.6 to

4.4 Mb, and the number of coding sequences (CDSs) ranged from 629 to 4,388 (**Table**

1). According to the taxonomic classification by GTDB-Tk, only 16 cMAGs were

assigned to known species (**Fig. 2**). Three cMAGs could not be assigned to known

families, and four cMAGs could not be assigned to known genera. These cMAGs

represented nine phyla, with Patescibacteria having the highest number of cMAGs.

Notably, more than 75% and 50% of Patescibacteria genomes were not assigned at the

species and genus level, respectively. The genomes of Patescibacteria bacteria are

very compact, with far fewer CDSs than the genomes of bacteria affiliated with other

phyla. The phyla Actinobacteriota, Armatimonadota, Bdellovibrionota, and

Proteobacteria each had only one member, but the abundances of cMAG.1

(Actinobacteriota) and cMAG.30 (Proteobacteria) were not low compared with other

cMAGs. cMAG.27, cMAG.1, and cMAG.25 were the most abundant in the

community of 30 members, and the abundances of nine species (cMAG.10, cMAG.17,

cMAG.5, cMAG.26, cMAG.18, cMAG.19, cMAG.2, cMAG.23, and cMAG.13) were

less than one-fiftieth of that of cMAG.27. Pure cultures are not available for none of

the 16 species assigned at the species level at present, indicating that all cMAGs

might represent uncultured species.

A total of three anammox bacteria, namely cMAG.27, cMAG.28, and cMAG.29,

were identified and they were all assigned to the genus *Ca.* Jettenia. According to the

classification by GTDB-Tk, cMAG.28 was unclassified at the species level, and cMAG.27 and cMAG.29 were assigned to the species *Ca*. Jettenia sp. AMX2 and *Ca*. Jettenia caeni, respectively. Therefore, we downloaded the 11 available *Ca*. Jettenia genomes from the NCBI database and calculated ANI values among the above three cMAGs and other *Ca*. Jettenia genomes. As shown in **Fig. S2**, ANI values between cMAG.29 and *Ca*. Jettenia caeni (GCA_000296795.1), and between cMAG.27 and *Ca*. Jettenia sp. AMX2 (GCA_900696655.1) were 99.70% and 99.45%, respectively. These results indicate that cMAG.29 and cMAG.27 are same species of *Ca*. Jettenia caeni and *Ca*. Jettenia sp. AMX2, respectively. The ANI value between cMAG.28 and *Ca*. Jettenia sp. AM49 (GCA_021650895.1) was 95.83%, which was just within the threshold of 95–96% species boundary (Richter and Rosselló-Móra, 2009; Palmer et al., 2020). Consequently, we calculated the DDH value between cMAG.28 and *Ca*. Jettenia sp. AM49 (GCA_021650895.1), and the resulting DDH value (66.60%) is less than the species boundary threshold of 70% (Meier-Kolthoff et al., 2013), indicating that cMAG.28 could represent a novel species of *Ca*. Jettenia different from *Ca*. Jettenia sp. AM49.

Functional divergence in general metabolism

Metabolic inference revealed that, aside from members of Patescibacteria, other bacteria in the anammox system might play a crucial role in nitrogen (N) and iron (Fe) cycles (**Fig. 3**). Additionally, members of phyla such as Planctomycetota and Chloroflexota were found to be important for the carbon (C) cycle in this anammox system. These inferences were further supported by the presence of important metabolic genes or pathways in their genome. For instance, members of the phyla Planctomycetota and Chloroflexota harbor a relatively high abundance of carbohydrate enzymes, including glycoside hydrolases and glycosyl transferases (**Fig. S3**).

 In this community, there might be 13 species capable of autotrophic growth that utilize three different carbon fixation pathways. Conversely, no known carbon fixation

pathways could be identified in the remaining 17 species, suggesting that they might

be heterotrophic. In detail, all the members of the phyla Chloroflexota and

Planctomycetota (cMAG.27, cMAG.28, and cMAG.29) harbor the key genes

encoding CO dehydrogenase and acetyl-CoA synthase, which are essential for carbon

fixation via the Wood-Ljungdahl (WL) pathway. As for the only Proteobacteria

bacterium (cMAG.30), the presence of a nearly complete Arnon-Buchanan cycle

indicates its potential for autotrophic growth. Moreover, the presence of the complete

Embden–Meyerhof pathway (EMP) and the non-oxidative pentose phosphate pathway

(PPP) pathways in cMAG.30 reflect its capability for heterotrophic growth. Besides,

the presence of genes encoding cytochrome c oxidase (cbb$_3$-type), cytochrome bc$_1$

ubiquinol oxidase, and succinate dehydrogenase in cMAG.30 suggests its capability

for aerobic respiration. Three members of Patescibacteria (cMAG.18, cMAG.19, and

cMAG.26) were found to carry the RubisCO-encoding gene (*rbcL*), suggesting that

they might perform carbon fixation through the Calvin-Benson-Bassham cycle (CBB).

The presence of complete EMP and the non-oxidative PPP pathways in three *Ca.*

Jettenia species (cMAG.27, cMAG.28, and cMAG.29), as well as a complete

tricarboxylic acid cycle (TCA cycle) and *pdhA* encoding pyruvate dehydrogenase,

suggest their capabilities for fermentation.

Functional divergence in nitrogen metabolism

To further explore functional divergence in nitrogen metabolism of these bacteria, 30

cMAGs were comprehensively annotated using a curated nitrogen cycle database

(NCycDB). A total of 24 species were identified as potentially participating in key

nitrogen conversion processes, such as anammox, denitrification, nitrification, and

dissimilatory nitrate reduction to ammonia (DNRA) (**Fig. 4A** and **B**). Three members

of *Ca.* Jettenia harbor the full set of anammox genes, including *nirK*/*nirS*, *hzs*, and

hdh, but they all lack *nxrA* encoding nitrite oxidoreductase (**Fig. 4A**). In addition, they

all harbor *napA, narG*, and *nrfA*, suggesting their capacities for DNRA. Similarly,

among the 11 *Ca.* Jettenia species from other habitats, none of them carry the *nxrA*

gene (**Fig. S4**). Only three members, i.e. two members of *Ca.* Jettenia caeni and one member of *Ca.* Jettenia sp. AMX2, harbor both *napA* and *narG*. Nearly all of these *Ca.* Jettenia members (10 out of 11) may be capable of DNRA. Members belonging to the genera *Ca.* Jettenia and *Ca.* Brocadia possess a richer repertoire of nitrogen metabolism marker genes compared with anammox bacteria in other genera (**Fig. S4**).

A total of 20 non-anammox bacteria community members were found to carry denitrification genes (**Fig. 4A**). However, none of them carry the full set of denitrification genes, indicating a potential for cooperative denitrification. In addition, seven members carry the key gene *nrfA* for DNRA, while only five of them harbor the complete DNRA pathway. The Armatimonadota bacterium (cMAG.2) exclusively harbors *nrfA*. According to gene abundance, the denitrification pathway might have not outcompeted both the anammox and DNRA pathways (**Fig. 4B**). Of note, the majority of Patescibacteria members were not been identified with important known nitrogen metabolism genes, suggesting that they might play other roles in the system. No complete nitrification pathway was identified in any of these cMAGs, and the gene encoding ammonia monooxygenase (*amo*) was not found.

Adaptive evolution

To gain insights into the adaptive evolution of the anammox community, an analysis was conducted to identify MGEs in all cMAGs using the latest MGE database (mobileOG-db) with stringent parameters. Among these cMAGs, eight members were found to carry MGEs, with a total of 63 MGEs identified (**Fig. 5A**). Notably, cMAG.4, cMAG.7, cMAG.28, and cMAG.30 were found to harbor sequences related to phages. The majority of the predicted MGEs were associated with integration or excision processes. Of note, one of the anammox bacteria (cMAG.29) was predicted to harbor a discrete distribution of 28 transposons, all of which contained the transposase-encoding gene *tnpA* (**Fig. S5**). Another anammox bacterium (cMAG.28) also harbors three transposons that containing *tnpA*, while the remaining members (including cMAG.27) did not carry *tnpA*. The same identification procedure was

performed on the other 26 reference genomes of anammox bacteria. Approximately

half of the anammox bacteria were predicted to carry *tnpA*, with gene counts no less

than 18 in *Ca.* Jettenia caeni (only one) and all members of *Ca.* Kuenenia

stuttgartiensis, and others carried no more than eight *tnpA* genes (**Fig. 5B**).

Furthermore, the upstream and downstream genes of *tnpA* in cMAG.29 were found to

be different (**Fig. 6**), indicating that they represent the simple transposon (insertion

sequence) without additional functional genes.

In addition to MGE analysis, HGT events were identified on all cMAGs. A total of

50 putative HGT events along with their transfer direction were identified, involving 6

phyla and 14 species (**Fig. 7A**). Most of the putative HGT events occurred within the

phylum Chloroflexota, accounting for 66% of the total. Putative HGT events within

the phylum Bacteroidetes accounted for 34% of the total, with 24% assumed to occur

with members of different phyla. At the species level, putative HGT events were the

most abundant in one member of the genus UBA12294 (cMAG.13), with 25 donors

and 5 recipients. Surprisingly, a putative HGT event was observed between an

anammox bacterium (cMAG.27) and a denitrifying bacterium affiliated with

Bacteroidota, suggesting that cMAG.27 might have received a gene from cMAG.8.

The donor and recipient genes for putative HGT events were identified. A total of

26 genes (> 50% of the total) in both donated and received genes could be annotated

using the Clusters of Orthologous Groups (COG) database (**Fig. 7B** and **C**). The

donated genes are primarily responsible for energy production and conversion (C, 4

genes), posttranslational modification, protein turnover, chaperones (O, 4 genes),

while four genes were annotated with unknown functions. For example, cMAG.4

donated four genes (*sucD*, *fumC*, *atpA*, and *atpD*) to cMAG.9. Interestingly, the

denitrifying bacterium cMAG.13 donated 10 genes to another denitrifying bacterium,

cMAG.16, and reciprocally, cMAG.16 donated 5 genes to cMAG.13. It should be

noted that cMAG.27 might receive a gene encoding IS200 family transposase. No

transfer genes directly related to nitrogen removal were identified. However, nearly

half of the genes in both donors and recipients were either unannotated or had

unknown functions; thus, additional investigation is required.

Discussion

In this study, a total of 30 complete circular prokaryotic genomes were successfully

recovered from an anammox community sample using HiFi sequencing. We made

slight adjustments to the filtering criteria compared with a previous study (Kim et al.,

2022) to include circular contigs that did not meet all the proposed criteria but

exhibited other desirable features. According to the predictions by PlasClass (Pellow

et al., 2020) (**Table S7**), predicted values of these substandard contigs were far lower

than the threshold (0.5) for plasmids. More importantly, most of these contigs were

assigned to Patescibacteria, which are also referred to as candidate phyla radiation

bacteria and are known to have much smaller genomes compared with other

prokaryotes (Luef et al., 2015; Castelle and Banfield, 2018; Tian et al., 2020). They

might have fewer marker genes than other conventional bacteria. These substandard

contigs were therefore retained for downstream analysis. In strict terms, some of the

recovered cMAGs from this study or previous studies might represent complete

chromosomes rather than complete genomes, as the existence of associated MGEs

(e.g., plasmids) remains uncertain. For complex cultures, such as anammox

community, accurately resolving the link of MGEs to their host complete

chromosomes poses a major challenge. To address this problem, we propose that a

combination of the recently developed high-throughput single-microbe sequencing

(Microbe-seq) (Zheng et al., 2022) with HiFi sequencing would offer an effective

strategy.

Although hifiasm-meta did not generate the highest number of circular contigs, it

yielded cMAGs that accounted for 63% of the total. This is consistent with previous

findings and highlights the advantages of hifiasm-meta for the assembly of HiFi reads

(Kim et al., 2022). These cMAGs represent anammox bacteria and associated bacteria,

including those with low abundance, and ~47% of them may represent novel species,

including a novel species of anammox bacteria. This suggests that a considerable

number of anammox-associated bacteria remain poorly characterized. The reliability

of the recovered cMAGs was supported by the high ANI values with conspecific

genomes from previous studies. Our approach outperformed the IHA method (Liu et

al., 2020) in terms of using fewer long reads (26.7 Gb vs. 69.4 Gb) and no need for

short reads while ensuring the accuracy of circular genome reconstruction,

demonstrating the potential for HiFi sequencing to accurately recover prokaryotic

genomes from complex environmental microbiomes.

cMAGs provide comprehensive and accurate information on the metabolic

potential and genetic interactions within the anammox community. Unlike previous

MAGs (Lawson et al., 2017; Liu et al., 2020; Kallistova et al., 2022) that were

affected by discontinuity and contamination, these cMAGs offer a more complete

picture of the community. The anammox system was thought to be autotrophic due to

the absence of additional organic carbon sources, but there are still many heterotrophs

in this system. Similar phenomena have been widely observed in previous studies

(Speth et al., 2016; Lawson et al., 2017; Zhao et al., 2018b). In light of our culture

conditions, these co-existing heterotrophic bacteria might be responsible for granular

aggregation, supplying growth factors to anammox bacteria, and consuming organic

matter.

Approximately 70% of the species in the community were potentially engaged in

nitrogen removal processes, either through anammox or denitrification pathways. The

remaining 30% might not be directly involved in such nitrogen conversion processes.

Nevertheless, some of them could perform carbon fixation, which provides organic

matter to other co-existing bacteria. These members of Patescibacteria might also play

roles in the degradation of poly-N-acetylglucosamine produced by anammox bacteria

(Hosokawa et al., 2021). Three anammox bacteria affiliated with *Ca.* Jettenia could

conduct DNRA and carbon fixation through the WL pathway, which is consistent with

the results of previous studies (Strous et al., 2006; Lawson et al., 2017; Mardanov et

al., 2019). Intriguingly, our results indicate that these three anammox bacteria were

capable of respiratory ammonification with oxidation acetate, and similar results were

observed for the two other anammox bacteria affiliated with *Ca.* Jettenia (Ali et al.,

2015; Mardanov et al., 2019). The DNRA capability of anammox bacteria could be a

valuable strategy for survival, as they would no longer dependent on other

microorganisms for the supply of ammonium and nitrite (Kartal et al., 2007).

Furthermore, not all anammox bacteria are capable of carrying out the DNRA

pathway, and it could be speculated that these anammox bacteria with the DNRA

pathway might be better adapted to environments with a low concentration of

ammonium and a high concentration of nitrite or nitrate.

Adaptive evolution plays a crucial role in the adaptation of bacteria to different

habitats, and both MGEs and HGT contribute to this process (Brito, 2021; Arnold et

al., 2022; Shi et al., 2022). We identified abundant MGEs and putative HGT events

within the anammox community. Certain members of the community showed

evidence of past phage infections, which led to mutations in their genomes. It is well

known that transposons can provide a pool of pre-evolved genes that help bacteria

adapt quickly to their environment (Dahlberg and Hermansson, 1995). The

transposons containing the *tnpA* gene were prevalent in some anammox bacteria,

indicating they might play important roles in their environmental adaptation. As

mentioned above, anammox bacteria are not available for pure culture to date; thus,

molecular biology methods based on monoclones cannot be used for the identification

of metabolic pathways or bioaugmentation. Many questions about the basic metabolic

pathways of anammox bacteria have not yet been addressed (Peeters and van Niftrik,

2019; Kuenen, 2020), challenges (such as the fast start-up and operation under

non-optimal conditions) during the full-scale operation of anammox also require

attention (Adams et al., 2020; Kouba et al., 2022), and this will require the

development of methods for the genome editing of anammox bacteria. We believe that

the transposon containing *tnpA* might be one of the breakthroughs for the genome

editing of anammox bacteria in mixed cultures.

A considerable number of putative HGT events were predicted, suggesting that

HGT might be another important strategy for promoting the adaptive evolution of

bacteria in this anammox community. The transferred genes are involved in multiple

metabolic pathways rather than antibiotic resistance, showing that these putative HGT

events might contribute to the metabolic specialization and competitive advantage of

these species (Seeleuthner et al., 2018; Wang et al., 2022). Particularly, the putative

HGT events mainly occurred in denitrifying bacteria in the phyla Chloroflexota and

Bacteroidota. Similar findings were observed in the previous study of a full-scale

anammox systems (Wang et al., 2022). It could be inferred that denitrifying bacteria

were probably very active in genetic interactions within anammox communities. The

putative HGT event between cMAG.27 and cMAG.8 shows that HGT events could

occur between anammox bacteria and co-existing bacteria affiliated with distantly

related lineages. Moreover, the transferred gene is a transposase-coding gene, again

suggesting that some specific transposons could be utilized for the genome editing of

anammox bacteria.

In summary, our study demonstrates that HiFi sequencing enables the

reconstruction of accurate cMAGs from the complex anammox community without

the need for Illumina short reads, binning, and reassembly. These cMAGs represent

not only high-abundance species, but also low-abundance species in this anammox

community. Additionally, these cMAGs provide valuable insights into microbial

diversity, metabolic divergence, and genetic interactions within the community, and

they may lay a core foundation for a comprehensive database for anammox bacteria

and associated microorganisms. The presence of MGEs and putative HGT events

suggests ongoing evolution within the community, and they may play an important

role in the domestication of the anammox culture. To our knowledge, this is the first

study to describe the functional divergence and adaptive evolution of uncultured

bacteria in an anammox community at the complete genome level. Further studies are

warranted to confirm the functional divergence and genetic interactions within

anammox communities, employing a combination of HiFi sequencing and other

techniques.

Conflict of interest

461　The authors declare no conflict of interest.

Data availability

463　The 30 cMAGs generated in this study have been deposited in the GenBank under the

464　accession number PRJNA983097. The raw sequencing data have been deposited in

465　the Sequence Read Archive under the accession number PRJNA981667 and

466　PRJNA981689.

Acknowledgments

468　This study was supported by the National Natural Science Foundation of China

469　(21876016, 52070031, and 42207022) and the Natural Science Foundation of

470　Guangdong Province (2023A1515012019).

References

Adams M., J. Xie, A. w. J. Kabore, Y. Chang, J. Xie, M. Guo, *et al.* (2022) Research
advances in anammox granular sludge: A review. *Crit. Rev. Environ. Sci.*
Technol. 52: 631-674.
Adams M., J. X. Xie, J. W. Xie, Y. F. Chang, M. L. Guo, C. J. Chen, *et al.* (2020) The

effect of carrier addition on Anammox start-up and microbial community: a
review. *Rev. Environ. Sci. Biotechnol.* 19: 355-368.

Ali M., M. Oshiki, T. Awata, K. Isobe, Z. Kimura, H. Yoshikawa*, et al.* (2015)
Physiological characterization of anaerobic ammonium oxidizing bacterium
"*Candidatus* Jettenia caeni". *Environ. Microbiol.* 17: 2172-2189.

Aramaki T., R. Blanc-Mathieu, H. Endo, K. Ohkubo, M. Kanehisa, S. Goto*, et al.*
(2020) KofamKOALA: KEGG Ortholog assignment based on profile HMM
and adaptive score threshold. *Bioinformatics* 36: 2251-2252.

Arnold B. J., I. T. Huang, W. P. Hanage (2022) Horizontal gene transfer and adaptive
evolution in bacteria. *Nat. Rev. Microbiol.* 20: 206-218.

Bhattacharjee A. S., S. Wu, C. E. Lawson, M. S. M. Jetten, V. Kapoor, J. W. S.
Domingo*, et al.* (2017) Whole-community metagenomics in two different
anammox configurations: Process performance and community structure.
Environ. Sci. Technol. 51: 4317-4327.

Bickhart D. M., M. Kolmogorov, E. Tseng, D. M. Portik, A. Korobeynikov, I.
Tolstoganov*, et al.* (2022) Generating lineage-resolved, complete
metagenome-assembled genomes from complex microbial communities. *Nat.*
Biotechnol. 40: 711-719.

Brito I. L. (2021) Examining horizontal gene transfer in microbial communities. *Nat.*
Rev. Microbiol. 19: 442-453.

Brown C. L., J. Mullet, F. Hindi, J. E. Stoll, S. Gupta, M. Choi*, et al.* (2022)
mobileOG-db: a manually curated database of protein families mediating the
life cycle of bacterial mobile genetic elements. *Appl. Environ. Microbiol.* 88:
e0099122.

Cao S. B., R. Du, Y. Zhou (2021) Coupling anammox with heterotrophic
denitrification for enhanced nitrogen removal: A review. *Crit. Rev. Environ.*
Sci. Technol. 51: 2260-2293.

Castelle C. J., J. F. Banfield (2018) Major new microbial groups expand diversity and
alter our understanding of the tree of life. *Cell* 172: 1181-1197.

Chan P. P., B. Y. Lin, A. J. Mak, T. M. Lowe (2021) tRNAscan-SE 2.0: improved
detection and functional classification of transfer RNA genes. *Nucleic Acids*
Res. 49: 9077-9096.

Dabbaghie F., J. Ebler, T. Marschall (2022) BubbleGun: enumerating bubbles and
superbubbles in genome graphs. *Bioinformatics* 38: 4217-4219.

Dahlberg C., M. Hermansson (1995) Abundance of Tn3, Tn21, and Tn501 transposase
(*tnpA*) sequences in bacterial community DNA from marine environments.
Appl. Environ. Microbiol. 61: 3051-3056.

De Anda V., I. Zapata-Peñasco, A. C. Poot-Hernandez, L. E. Eguiarte, B.
Contreras-Moreira, V. Souza (2017) MEBS, a software platform to evaluate
large (meta)genomic collections according to their metabolic machinery:
unraveling the sulfur cycle. *GigaScience* 6: 1-17.

Ding C., L. Adrian (2020) Comparative genomics in "*Candidatus* Kuenenia

stuttgartiensis" reveal high genomic plasticity in the overall genome structure, CRISPR loci and surface proteins. *BMC Genomics* 21: 851.

Douglas G. M., M. G. I. Langille (2019) Current and promising approaches to identify horizontal gene transfer events in metagenomes. *Genome Biol. Evol.* 11: 2750-2766.

Feng X., H. Cheng, D. Portik, H. Li (2022) Metagenome assembly of high-fidelity long reads with hifiasm-meta. *Nat. Methods* 19: 671-674.

Feng Y., Y. P. Zhao, Y. Z. Guo, S. T. Liu (2018) Microbial transcript and metabolome analysis uncover discrepant metabolic pathways in autotrophic and mixotrophic anammox consortia. *Water Res.* 128: 402-411.

Frank J., S. Lücker, R. H. A. M. Vossen, M. S. M. Jetten, R. J. Hall, H. J. M. Op den Camp, *et al.* (2018) Resolving the complete genome of *Kuenenia stuttgartiensis* from a membrane bioreactor enrichment using Single-Molecule Real-Time sequencing. *Sci. Rep.* 8: 4580.

Galambos D., R. E. Anderson, J. Reveillaud, J. A. Huber (2019) Genome-resolved metagenomics and metatranscriptomics reveal niche differentiation in functionally redundant microbial communities at deep-sea hydrothermal vents. *Environ. Microbiol.* 21: 4395-4410.

Gurevich A., V. Saveliev, N. Vyahhi, G. Tesler (2013) QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29: 1072-1075.

Hon T., K. Mars, G. Young, Y. C. Tsai, J. W. Karalius, J. M. Landolin, *et al.* (2020) Highly accurate long-read HiFi sequencing data for five complex genomes. *Sci. Data* 7: 399.

Hosokawa S., K. Kuroda, T. Narihiro, Y. Aoi, N. Ozaki, A. Ohashi, *et al.* (2021) Cometabolism of the superphylum Patescibacteria with anammox bacteria in a long-term freshwater anammox column reactor. *Water* 13: 208.

Huerta-Cepas J., D. Szklarczyk, D. Heller, A. Hernández-Plaza, S. K. Forslund, H. Cook, *et al.* (2019) eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res.* 47: D309-D314.

Hyatt D., G. L. Chen, P. F. LoCascio, M. L. Land, F. W. Larimer, L. J. Hauser (2010) Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11: 119.

Jain C., L. M. Rodriguez-R, A. M. Phillippy, K. T. Konstantinidis, S. Aluru (2018) High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat. Commun.* 9: 5114.

Kallistova A., Y. Nikolaev, V. Grachev, A. Beletsky, E. Gruzdev, V. Kadnikov, *et al.* (2022) New insight into the interspecies shift of anammox bacteria *Ca.* "Brocadia" and *Ca.* "Jettenia" in reactors fed with formate and folate. *Front. Microbiol.* 12: 802201.

Kartal B., N. M. de Almeida, W. J. Maalcke, H. J. M. Op den Camp, M. S. M. Jetten, J. T. Keltjens (2013) How to make a living from anaerobic ammonium

oxidation. *FEMS Microbiol. Rev.* 37: 428-461.

Kartal B., M. M. M. Kuypers, G. Lavik, J. Schalk, H. den Camp, M. S. M. Jetten*, et al.* (2007) Anammox bacteria disguised as denitrifiers: nitrate reduction to dinitrogen gas via nitrite and ammonium. *Environ. Microbiol.* 9: 635-642.

Kim C. Y., J. Ma, I. Lee (2022) HiFi metagenomic sequencing enables assembly of accurate and complete genomes from human gut microbiota. *Nat. Commun.* 13: 6367.

Kolmogorov M., D. M. Bickhart, B. Behsaz, A. Gurevich, M. Rayko, S. B. Shin*, et al.* (2020) metaFlye: scalable long-read metagenome assembly using repeat graphs. *Nat. Methods* 17: 1103-1110.

Kouba V., C. Bachmannova, T. Podzimek, P. Lipovova, M. C. M. van Loosdrecht (2022) Physiology of anammox adaptation to low temperatures and promising biomarkers: A review. *Bioresour. Technol.* 349: 126847.

Kuenen J. G. (2020) Anammox and beyond. *Environ. Microbiol.* 22: 525-536.

Lackner S., E. M. Gilbert, S. E. Vlaeminck, A. Joss, H. Horn, M. C. M. van Loosdrecht (2014) Full-scale partial nitritation/anammox experiences—An application survey. *Water Res.* 55: 292-303.

Lawson C. E., S. Wu, A. S. Bhattacharjee, J. J. Hamilton, K. D. McMahon, R. Goel*, et al.* (2017) Metabolic network analysis reveals microbial community interactions in anammox granules. *Nat. Commun.* 8: 15416.

Liu L., Y. Wang, Y. Che, Y. Chen, Y. Xia, R. Luo*, et al.* (2020) High-quality bacterial genomes of a partial-nitritation/anammox system by an iterative hybrid assembly method. *Microbiome* 8: 155.

Luef B., K. R. Frischkorn, K. C. Wrighton, H.-Y. N. Holman, G. Birarda, B. C. Thomas*, et al.* (2015) Diverse uncultivated ultra-small bacterial cells in groundwater. *Nat. Commun.* 6: 6372.

Maguire F., B. F. Jia, K. L. Gray, W. Y. V. Lau, R. G. Beiko, F. S. L. Brinkman (2020) Metagenome-assembled genome binning methods with short reads disproportionately fail for plasmids and genomic Islands. *Microb. Genomics* 6: 000436.

Mardanov A. V., A. V. Beletsky, N. V. Ravin, E. A. Botchkova, Y. V. Litti, A. N. Nozhevnikova (2019) Genome of a novel bacterium "*Candidatus* Jettenia ecosi" reconstructed from the metagenome of an anammox bioreactor. *Front. Microbiol.* 10: 2442.

Meier-Kolthoff J. P., A. F. Auch, H. P. Klenk, M. Goker (2013) Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14: 60.

Meier-Kolthoff J. P., J. S. Carbasse, R. L. Peinado-Olarte, M. Göker (2022) TYGS and LPSN: a database tandem for fast and reliable genome-based classification and nomenclature of prokaryotes. *Nucleic Acids Res.* 50: D801-D807.

Meziti A., L. M. Rodriguez-R, J. K. Hatt, A. Pena-Gonzalez, K. Levy, K. T. Konstantinidis (2021) The reliability of metagenome-assembled genomes

(MAGs) in representing natural populations: Insights from comparing MAGs
against isolate genomes derived from the same fecal sample. *Appl. Environ.*
Microbiol. 87: e02593-02520.

Meziti A., D. Tsementzi, L. M. Rodriguez-R, J. K. Hatt, H. Karayanni, K. A. Kormas,
et al. (2019) Quantifying the changes in genetic diversity within
sequence-discrete bacterial populations across a spatial and temporal riverine
gradient. *ISME J.* 13: 767-779.

Nurk S., S. Koren, A. Rhie, M. Rautiainen, A. V. Bzikadze, A. Mikheenko, *et al.*
(2022) The complete sequence of a human genome. *Science* 376: 44-53.

Nurk S., B. P. Walenz, A. Rhie, M. R. Vollger, G. A. Logsdon, R. Grothe, *et al.* (2020)
HiCanu: accurate assembly of segmental duplications, satellites, and allelic
variants from high-fidelity long reads. *Genome Res.* 30: 1291-1305.

Palmer M., E. T. Steenkamp, J. Blom, B. P. Hedlund, S. N. Venter (2020) All ANIs are
not created equal: implications for prokaryotic species boundaries and
integration of ANIs into polyphasic taxonomy. *Int. J. Syst. Evol. Microbiol.* 70:
2937-2948.

Parks D. H., M. Chuvochina, C. Rinke, A. J. Mussig, P. A. Chaumeil, P. Hugenholtz
(2022) GTDB: an ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-based
taxonomy. *Nucleic Acids Res.* 50: D785-D794.

Parks D. H., C. Rinke, M. Chuvochina, P. A. Chaumeil, B. J. Woodcroft, P. N. Evans,
et al. (2017) Recovery of nearly 8,000 metagenome-assembled genomes
substantially expands the tree of life. *Nat. Microbiol.* 2: 1533-1542.

Patro R., G. Duggal, M. I. Love, R. A. Irizarry, C. Kingsford (2017) Salmon provides
fast and bias-aware quantification of transcript expression. *Nat. Methods* 14:
417-419.

Peeters S. H., L. van Niftrik (2019) Trending topics and open questions in anaerobic
ammonium oxidation. *Curr. Opin. Chem. Biol.* 49: 45-52.

Pellow D., I. Mizrahi, R. Shamir (2020) PlasClass improves plasmid sequence
classification. *PLoS Comp. Biol.* 16: e1007781.

Richter M., R. Rosselló-Móra (2009) Shifting the genomic gold standard for the
prokaryotic species definition. *PNAS* 106: 19126-19131.

Richter M., R. Rosselló-Móra, F. Oliver Glöckner, J. Peplies (2015) JSpeciesWS: a
web server for prokaryotic species circumscription based on pairwise genome
comparison. *Bioinformatics* 32: 929-931.

Sangwan N., F. F. Xia, J. A. Gilbert (2016) Recovering complete and draft population
genomes from metagenome datasets. *Microbiome* 4: 8.

Seeleuthner Y., S. Mondy, V. Lombard, Q. Carradec, E. Pelletier, M. Wessner, *et al.*
(2018) Single-cell genomics of multiple uncultured stramenopiles reveals
underestimated functional diversity across oceans. *Nat. Commun.* 9: 310.

Shi A., F. Fan, J. R. Broach (2022) Microbial adaptive evolution. *J. Ind. Microbiol.*
Biotechnol. 49: kuab076.

Song W., B. Wemheuer, S. Zhang, K. Steensen, T. Thomas (2019) MetaCHIP:
community-level horizontal gene transfer identification through the
combination of best-match and phylogenetic approaches. *Microbiome* 7: 36.
Speth D. R., M. H. in 't Zandt, S. Guerrero-Cruz, B. E. Dutilh, M. S. M. Jetten (2016)
Genome-based microbial ecology of anammox granules in a full-scale
wastewater treatment system. *Nat. Commun.* 7: 11172.
Strous M., E. Pelletier, S. Mangenot, T. Rattei, A. Lehner, M. W. Taylor, *et al.* (2006)
Deciphering the evolution and metabolism of an anammox bacterium from a
community genome. *Nature* 440: 790-794.
Suarez C., P. Dalcin Martins, M. S. M. Jetten, S. Karačić, B. M. Wilén, O. Modin, *et
al.* (2022) Metagenomic evidence of a novel family of anammox bacteria in a
subsea environment. *Environ. Microbiol.* 24: 2348-2360.
Tao Y., F. Xun, C. Zhao, Z. Mao, B. Li, P. Xing, *et al.* (2023) Improved assembly of
metagenome-assembled genomes and viruses in Tibetan saline lake sediment
by HiFi metagenomic sequencing. *Microbiol. Spectrum* 11: e03328-03322.
Tian R., D. Ning, Z. He, P. Zhang, S. J. Spencer, S. Gao, *et al.* (2020) Small and
mighty: adaptation of superphylum Patescibacteria to groundwater
environment drives their genome simplicity. *Microbiome* 8: 51.
Tu Q., L. Lin, L. Cheng, Y. Deng, Z. He (2019) NCycDB: a curated integrative
database for fast and accurate metagenomic profiling of nitrogen cycling genes.
Bioinformatics 35: 1040-1048.
Wang D., Y. Meng, F. Meng (2022) Genome-centric metagenomics insights into
functional divergence and horizontal gene transfer of denitrifying bacteria in
anammox consortia. *Water Res.* 224: 119062.
Yang Y., Z. Lu, M. Azari, B. Kartal, H. Du, M. Cai, *et al.* (2022) Discovery of a new
genus of anaerobic ammonium oxidizing bacteria with a mechanism for
oxygen tolerance. *Water Res.* 226: 119165.
Yuan C., J. K. Lei, J. Cole, Y. N. Sun (2015) Reconstructing 16S rRNA genes in
metagenomic data. *Bioinformatics* 31: 35-43.
Zhao Y. P., Y. Feng, J. Q. Li, Y. Z. Guo, L. M. Chen, S. T. Liu (2018a) Insight into the
aggregation capacity of anammox consortia during reactor start-up. *Environ.
Sci. Technol.* 52: 3685-3695.
Zhao Y. P., S. F. Liu, B. Jiang, Y. Feng, T. T. Zhu, H. C. Tao, *et al.* (2018b)
Genome-centered metagenomics analysis reveals the symbiotic organisms
possessing ability to cross-feed with anammox bacteria in anammox consortia.
Environ. Sci. Technol. 52: 11285-11296.
Zheng W., S. Zhao, Y. Yin, H. Zhang, D. M. Needham, E. D. Evans, *et al.* (2022)
High-throughput, single-microbe genomics with strain resolution, applied to a
human gut microbiome. *Science* 376: eabm1483.

Table 1. Characteristics of the 30 cMAGs obtained in this study.

ID	Phylum (Genus)	Genome size (Mb)	GC (%)	CDS	GTDB-Tk marker gene	Types of tRNA	Features (5S/16S/23S)
cMAG.1	Actinobacteriota (Unclassified)	2.2	68.58	2,260	118	23	1/1/1
cMAG.2	Armatimonadota (H1-ARM1)	2.9	60.95	2,651	111	23	1/1/1
cMAG.3	Bacteroidota (OLB6)	3.0	49.64	2,427	119	22	2/2/2
cMAG.4	Bacteroidota (BCD5)	3.3	36.15	2,825	119	22	1/1/1
cMAG.5	Bacteroidota (UTCHB3)	3.4	42.13	2,623	118	23	1/1/1
cMAG.6	Bacteroidota (SURF-28)	3.7	34.47	3,351	118	22	1/1/1
cMAG.7	Bacteroidota (OLB8)	3.9	65.21	3,252	119	22	1/1/1
cMAG.8	Bacteroidota (PHOS-HE28)	4.0	41.08	3,258	119	22	2/2/2
cMAG.9	Bacteroidota (UBA2336)	4.1	43.59	3,752	117	23	2/2/2
cMAG.10	Bdellovibrionota (Unclassified)	2.3	55.04	2,089	93	18	1/1/1
cMAG.11	Chloroflexota (UBA7227)	3.5	59.60	3,200	116	24	1/1/1
cMAG.12	Chloroflexota (UBA7227)	3.6	59.76	3,268	116	24	1/1/1
cMAG.13	Chloroflexota (UBA12294)	3.9	53.51	3,590	111	23	1/1/1
cMAG.14	Chloroflexota (UBA12294)	4.0	53.65	3,738	110	23	1/1/1
cMAG.15	Chloroflexota (OLB14)	4.1	54.13	3,820	113	23	1/1/1
cMAG.16	Chloroflexota (OLB14)	4.4	53.56	4,388	115	24	1/1/1
cMAG.17	Desulfobacterota (UBA5799)	2.8	57.87	2,559	119	23	1/1/1
cMAG.18	Patescibacteria (Unclassified)	1.0	44.78	956	89	20	1/1/1
cMAG.19	Patescibacteria (Unclassified)	1.1	39.76	982	84	21	1/1/1
cMAG.20	Patescibacteria (Unclassified)	1.2	60.26	1,204	95	21	1/1/1

cMAG.21	Patescibacteria (OLB21)	1.2	38.42	1,073	93	21	1/1/1
cMAG.22	Patescibacteria (Unclassified)	0.6	52.17	629	82	21	1/1/1
cMAG.23	Patescibacteria (BOKC01)	0.7	40.37	698	84	22	1/1/2
cMAG.24	Patescibacteria (UBA12026)	0.8	34.42	947	80	20	1/1/1
cMAG.25	Patescibacteria (UBA919)	0.9	36.84	827	90	22	1/2/1
cMAG.26	Patescibacteria (Unclassified)	0.9	41.50	858	87	21	1/1/1
cMAG.27	Planctomycetota (*Ca*. Jettenia)	3.5	41.60	3,145	108	23	2/2/2
cMAG.28	Planctomycetota (*Ca*. Jettenia)	3.8	39.42	3,292	107	24	2/2/2
cMAG.29	Planctomycetota (*Ca*. Jettenia)	4.0	40.01	3,510	107	24	2/2/2
cMAG.30	Proteobacteria (QUBU01)	3.4	66.68	3,021	118	23	1/1/1

Figure legends

Fig. 1. Workflow for filtering the HiFi cMAGs from three assemblies by hifiasm-meta,

Flye, and Canu.

Fig. 2. The phylogenetic placement and abundance of cMAGs recovered from

anammox granular sludge. The phylogenetic tree was inferred based on the multiple

sequence alignments of conserved marker genes from 30 cMAGs using the

GTDB-Tk.

Fig. 3. The MEBS scores for the nitrogen (N), iron (Fe), oxygen (O), carbon (C) and

sulfur (S) metabolic pathways of 30 cMAGs.

Fig. 4. The potential for nitrogen metabolism of 30 cMAGs. (**A**) Nitrogen metabolism

marker genes. (**B**) Proposed pathway for nitrogen conversion in the anammox

community. The number indicates the number of species with genes encoding a given

step. Colored squares indicate gene abundances.

Fig. 5. Mobile genetic elements (MGEs) identified in the cMAGs and reference

genomes. (**A**) MGEs identified in cMAGs. (**B**) The *tnpA* counts identified in two *Ca.*

Jettenia species of this study and other anammox species.

Fig. 6. The predicted 28 gene clusters containing *tnpA* of cMAG.29.

Fig. 7. Network and functions of HGT genes of the anammox community. (**A**) HGT

network of the anammox community; the direction and width of blue arrows indicate

the direction of transfer and gene counts, respectively; the size of the node represents

the count of connected edges. (**B**) and (**C**) Bubble size indicates the count of HGT

genes annotated with COG functional categories. C: Energy production and

conversion; D: Cell cycle control, cell division, chromosome partitioning; E: Amino

acid transport and metabolism; H: Coenzyme transport and metabolism; J: Translation,

ribosomal structure and biogenesis; L: Replication, recombination and repair; N: Cell

motility; O: Posttranslational modification, protein turnover; S: Function unknown; T:

Signal transduction mechanisms.

718   **Fig. 1**

720 **Fig. 2**

722   **Fig. 3**

Fig.

726   **Fig. 5**

728   **Fig. 6**

730   **Fig. 7**