1

The evolution of sexual dimorphism in gene expression

in response to a manipulation of mate competition

Prashastha Mishra[1], Howard D. Rundle[2,3], and Aneil F. Agrawal[1,3]

8   [1]Department of Ecology & Evolutionary Biology, University of Toronto, Toronto, Ontario, Canada

9   [2]Department of Biology, University of Ottawa, Ottawa, Ontario, Canada; E-mail: hrundle@uottawa.ca

10   [3]HDR and AFA contributed equally to this work.

Abstract

Many genes are differentially expressed between males and females and patterns of sex-biased gene expression (SBGE) vary among species. This variation is thought to have evolved in response to differences in mate competition among species that causes varying patterns of sex-specific selection. We used experimental evolution to test this by quantifying SBGE and sex-specific splicing in 15 *Drosophila melanogaster* populations that evolved for 104 generations in mating treatments that removed mate competition via enforced monogamy (MCabs), or allowed mate competition in either small, simple (MCsim) or larger, structurally more complex (MCcom) mating environments. Consistent with SBGE being the product of sex-specific selection, initially sex-biased genes diverged in expression more among treatments than unbiased genes, and there was greater expression divergence for male- than female-biased genes. It has been suggested the transcriptome should be 'feminized' under monogamy because of the removal of sexual selection on males; we did not observe this, likely because selection differs in additional ways between monogamy vs. polygamy. Significant divergence in average expression dimorphism between treatments was observed, and in some treatment comparisons the direction of the divergence differed across different sex-bias categories. There was not a generalized reduction in expression dimorphism under enforced monogamy.

Keywords: sex-biased gene expression, sex-specific splicing, sexual selection, sexual conflict, *Drosophila*,

Introduction

In many animals, substantial differences between the sexes exist in a myriad of phenotypes involving morphology, behaviour, physiology, and life history. Dimorphism in these phenotypes arises, at least in part, from sex differences in expression of the underlying genes, which themselves are another set of phenotypes for which sexual dimorphism can be considered. Studies across many taxa report the existence of sex-biased expression in a large fraction of genes for any given tissue and/or developmental stage (Grath & Parsch, 2016). However, the extent of transcriptional dimorphism can vary considerably between species (Ingleby et al., 2015), echoing patterns of variation in phenotypic dimorphism.

Contrasting selection between the sexes is widely believed to be the major reason for sexual dimorphism. Sex-specific optima exist for many traits, causing intralocus sexual conflict over optimal expression of underlying genes that are shared between males and females (Bonduriansky & Chenoweth 2009). Such conflict can be resolved via the evolution of sex-biased gene expression (Parsch & Ellegren, 2013). The different reproductive strategies of males and females are thought to be a major source of such sex-specific selection. In particular, selection arising from mate competition, including pre- and post-copulatory sexual selection, often differs between the sexes, shaping the traits that mediate intra- and intersexual interactions. We use "mating system" to refer to the set of inter- and intrasexual interactions related to mating and reproduction.

Mating systems vary widely among species and, hence, this variation is presumed to be a major source of variation in sexual dimorphism, though this has received limited direct attention (Fernandes Martins et al., 2017). At the expression level, Harrison et al. (2015) used a comparative approach to examine the effect of mating system on variation in dimorphism among six bird species. The proportion of genes that were male-biased was positively correlated with presumed indices of both pre- and post-copulatory sexual selection (e.g., sexual ornamentation, sperm number, residual testis mass). Further, the rate of turnover of male-biased genes was positively associated with male sexual ornamentation. This suggests that sexual selection drives the evolution of expression dimorphism. Further evidence consistent with the importance of mating system comes from studies showing that sex-biased gene expression differs between alternative reproductive morphs, for example between 'dominant' vs. 'auxiliary' males (Dean et al., 2017; Pointer et al., 2013; Stuglik et al., 2014).

Experimental evolution offers a powerful means to directly test whether a change in mating system drives evolutionary changes in the transcriptome. In one such study, Hollis et al. (2014) subjected

replicate populations of the naturally polygamous *D. melanogaster* to experimental evolution under two

mating systems: polygamy and enforced monogamy (i.e., randomly assigned male-female pairings). The

motivation for imposing monogamy was to eliminate sexual selection on males, which is thought to be

the primary reason why selection differs between the sexes. Hollis et al. predicted that in the absence of

sexual selection on males, a population would no longer be constrained by conflicting selection between

the sexes, and hence phenotypes in males and females would evolve towards female optima. Assuming

existing patterns of sex-biased gene expression represent only a partial resolution of intralocus sexual

conflict, they predicted that evolution under monogamy would result in 'feminisation' of the

transcriptome (i.e., upregulation of female-biased genes and down-regulation of male-biased genes) in

both sexes. These predictions were borne out, with expression being feminised in males and females

under enforced monogamy compared to polygamy. However, a similar evolution experiment using *D.

pseuodoobscura* found results that were very different from these predictions, with expression being

largely masculinised in populations evolving under monogamy compared to polygamy (Veltsos et al.,

2017).

Though Hollis et al. (2014) used enforced monogamy with the goal of eliminating sexual selection on

males, such manipulations of mating systems are likely to have additional consequences on selection

more broadly in both sexes (Rowe & Rundle, 2021). Males are known to inflict harm on females, for

example through persistent courtship and toxic seminal fluid proteins (Fowler and Partridge 1989,

Partridge and Fowler 1990; Chapman et al. 1995, Arnqvist and Rowe 2005). Under enforced monogamy,

there should be strong selection on males to be less harmful to females, and experimental evidence

supports this (Holland and Rice 1999; Yun et al. 2021). Selection on females also likely differs between

these mating systems. Reduced harm by males may subsequently yield selection against costly female

traits involved in avoiding or reducing male harm (Wigby & Chapman, 2004). In addition, under some

polygamous but not monogamous conditions, high-quality females may suffer a 'cost of attractiveness'

as a result of being the targets of preferential male harassment (Long et al., 2009; Yun et al. 2017,

MacPherson et al., 2018), ultimately weakening natural selection through females. A change from

polygamy to enforced monogamy is thus likely to alter selection in both sexes in a variety of ways, such

that contrasts between these mating systems are more nuanced than simply the presence vs. absence

of sexual selection. Expression changes may therefore be more difficult to predict than Hollis et al.

(2014) suggested. Further complicating this contrast is that 'polygamy' can take many forms,

characterized by differences in inter- and intrasexual interactions that occur under different contexts

(e.g., when reproductive interactions and mating happen in different environments).

Here, we analyze sex-biased gene expression in replicate populations of *D. melanogaster* evolved in

three treatments that varied with respect to mating system. One treatment involved the absence of

mate competition (MCabs) via the application of enforced monogamy. The remaining two treatments

allowed for mate competition, but in distinct 'mating environments'. In the first of these treatments

(MCsim), mate competition occurred in a relatively simple environment (e.g., *Drosophila* vials), similar to

the 'polygamy' treatment of earlier studies (i.e., Hollis et al. 2014; Veltsos et al. 2017). In the third

treatment (MCcom) mate competition occurred in larger, less dense containers with multiple food

sources and greater spatial complexity, presumably allowing females to more readily evade males.

Consistent with this, in the 'complex' relative to the 'simple' mating environment, intersexual

interactions and mating are less frequent, males are less harmful to females, and females no longer

suffer a 'cost of attractiveness' that weakens viability selection on them in the simple environment (Yun

et al. 2017, 2019; MacPherson et al. 2018). Populations maintained in the complex mating environment

also evolved males that are less harmful to females compared to their counterparts from the simple

mating environment (Yun et al. 2021), yet these males are highly successful in siring offspring in

competition with other males (Yun et al 2019).

Here we analyze gene expression divergence among these three mating treatments from several

perspectives. We begin by asking whether divergence among mating treatments is random across the

transcriptome with respect to pre-existing sex-bias in expression. If sex-biased gene expression (SBGE)

evolves because of differential selection arising from mating interactions, then one would expect that

sex-biased genes would be more likely to diverge among mating treatments than would unbiased genes.

Second, we contrast female- vs. male-biased genes. Past studies have documented that male-biased

genes diverge in expression more rapidly among populations or species than female-biased genes (e.g.,

Meiklejohn et al. 2003, Zhang et al. 2007, Allen et al. 2018), but this is yet to be considered in relation to

variation in mating system. We ask whether male-biased genes are more likely to diverge than female-

biased genes in response to changes in mating system.

Third, we examine the directionality of expression changes from the perspective of Hollis et al. (2014),

testing their prediction that enforced monogamy should lead to a feminization of the transcriptome

relative to polygamous mating systems in which there is a much greater opportunity for sexual selection

on males. We compare our enforced monogamy treatment with each of two different polygamous

mating treatments in which mate competition occurs in different environments. We also compare the

latter two mate competition treatments with one another, asking whether they differ with respect to

transcriptome feminization or masculinization.

Changes in dimorphism can occur because of changes in just one sex or both. Moreover, examination of

dimorphic traits in each sex separately can provide clues as to whether dimorphism is hindered by

intersexual genetic covariances (Lande 1980; Prasad et al. 2007). For such reasons, we examine gene

expression in each sex separately, as have past studies (Hollis et al. 2014; Veltsos et al. 2017). For

example, Hollis et al. (2014) reported feminization in the transcriptomes of females, and also of males,

under enforced monogamy. However, reporting results in this way alone does not provide a clear view

of how dimorphism itself changed (this was not a goal of past work). For example, feminization in both

sexes under enforced monogamy could mean that dimorphism had increased, decreased, or remained

constant, depending on the relative magnitudes of the changes within each sex. Though we are

interested in the evolutionary divergence of dimorphic traits within each sex, like past studies (Hollis et

al. 2014; Veltsos et al. 2017), we are also interested in whether mating system affects dimorphism per

se, so we also explicitly compare dimorphism across mating treatments.

We examine the above questions with respect to expression dimorphism in whole bodies as well as

heads. In being less sexually dimorphic, heads offer a point of contrast. Finally, in addition to examining

SBGE, we also examine some of these questions with respect to another form of expression

dimorphism, sex-specific splicing (SSS), i.e., quantitative differences between the sexes in the relative

usage of different isoforms.

Methods

Samples for RNAseq

We used 15 populations from a previously described evolution experiment (Yun et al. 2018, 2019, 2021)

that were created in September 2014 by sampling from a single laboratory stock population of *D.*

melanogaster. During experimental evolution, all populations used here were raised under the same

larval conditions consisting of standard cornmeal medium supplemented with 5% NaCl (6% after

generation 6) and a constant exposure to 28 C (rather than the standard 25 C of the ancestor) during

larval development. The 15 populations were divided equally among three treatments that manipulated

the mating system that adults experienced, including the opportunity for mate competition and the

environment in which this occurred. The treatments were: mate competition absent (MCabs) via

randomly assigned single-pair monogamy, and two polygamy treatments in which mate competition

was permitted either in small, structurally simple *Drosophila* vials (MCsim) or in larger, 1.65-L cylindrical

containers with added structural complexity (MCcom).

Populations were maintained via 3-week non-overlapping generations. Within a given population, 140

male and 140 female adults spent 6 days each generation in their respective mating treatment. In

MCabs, this involved 140 male-female pairs separately allocated to individual wide-mouth straws, while

in the polygamy treatments this involved four groups of 35 males and 35 females, each group being held

in either separate vials (MCsim) or separate containers (MCcom). Within each of the MCcom containers

were five small cups of food (with plastic barriers inserted into the food that further subdivided the

surface) and two coiled pipe cleaners, anchored in the lid, that extended into the interior space. After 6

d in these mating treatments, males were discarded and 105 of the surviving females were randomly

allocated among seven vials to lay eggs for ~24 h. Females were subsequently discarded and the

resulting offspring developed under the same larval conditions in all populations (i.e., cornmeal food

with added salt at 28 C). Adults that eclosed were used to create the next generation's mating

treatments.

After 104 generations of selection, samples for RNAseq were obtained as follows. Flies were reared

under a controlled density of 40 larvae per vial on a benign yeast-agar food (no salt). Adults emerged

eight days after hatching and were collected under light CO_2 (< 20 seconds) as virgins (within 8 hours of

emergence) and then held in single-sex vials at low density (10 flies per vial). Two days later, flies were

processed. For whole body samples, flies were transferred under light CO_2 (< 20 sec) to microcentrifuge

tubes (10 flies per tube). A few minutes later (and well after awakening from CO_2) tubes were flash

frozen in liquid nitrogen and then stored at -80 C. For head samples, the same procedure was used

except after flash freezing, tubes were vortex shaken for ca. 10 s to separate heads from bodies. Heads

were transferred to a new microcentrifuge tube (8-10 heads per tube) and then stored at -80 C. RNA

extraction was performed using ThermoFisher PicoPure RNA Isolation Kit. Paired-end (100 bp)

sequencing was performed using Illumina NovaSeq S4.

One of the female replicates from MCsim treatment was found to have elevated expression of Y-linked

genes, indicating possible contamination with male tissue. Consequently, we excluded this replicate

from all further analyses.

Differential Expression between Mating Treatments

Differential expression was analysed between each pair of mating treatments, separately for each sex

and tissue. Each RNAseq file was aligned to the *D. melanogaster* reference genome (dos Santos et al.,

2015) using STAR v2.7 (Dobin et al., 2013) with default parameters. The resulting alignment files were

processed with *htseq-count* to obtain gene-level read counts for each sample (Anders et al., 2015). The R

package *DESeq2* (Love et al., 2014) was used to estimate differential expression between pairs of mating

treatments. For all genes averaging >50 reads across all replicates, we estimated the $\log_2$ fold change in

expression between treatments (henceforth, 'treatment effect'). For any genes thus tested, *DESeq2*

yields an adjusted p-value by applying Benjamini and Hochberg corrections for multiple testing. A gene

was designated to have significant differential expression between treatments if the treatment effect

was accompanied by an adjusted p-value < 0.1. Such genes are called 'Treatment Differentially

Expressed genes' or 'TDE genes'.

Sex-Biased Gene Expression

We used an external dataset (Osada et al., 2017) to characterise genes with respect to sex-biased gene

expression for our analyses. This choice for an external dataset is not critical for our purposes; though

expression in this external dataset undoubtedly differs somewhat from our own, variation among genes

in SBGE tends to be much larger than variation in SBGE for a gene across studies (e.g., genes that are

male- or female-biased in one study are generally male- or female-biased in other studies, though they

may vary quantitively the magnitude of bias). Even across species separated by millions of years, among-

gene SBGE is strongly correlated (Zhang et al. 2007).

The Osada et al. RNAseq dataset consisted of data for whole bodies and heads for a male and female

sample from each of 18 lines. Gene-level read counts were obtained as described above. Sex-biased

gene expression was estimated using the R package *DESeq2* (Love et al., 2014). In estimating differential

expression between sexes, we excluded all genes which averaged <50 reads across all male and female

replicates in the given tissue. *DESeq2* yields $\log_2$ estimates of fold change in male to female expression

("$\log_2$ FC male/female"). For some analyses, we assigned genes into one of three sex bias categories:

female-biased ($-\infty < \log2FC \leq -0.5$), unbiased ($-0.5 < \log2FC \leq 0.5$), and male-biased ($0.5 < \log2FC < \infty$).

Differential Isoform Usage Analysis

In addition to analysing changes to total gene expression, we also considered changes in patterns of

alternative splicing using the R package *JunctionSeq* (Hartley and Mullikin 2016). *JunctionSeq* utilises

read count data from *QoRTs* (Hartley & Mullikin, 2015), which partitions genes into exons and splice

junctions. It then counts the number of reads that overlap an annotated feature (i.e., exon or splice

junction). Then, through *JunctionSeq*, generalised linear models (GLM) are used to test for differential

splicing. At the gene level, corrections for multiple testing are applied through the Benjamini and

Hochberg method, yielding an adjusted p-value for each gene.

We used *QoRTs* to gather read count data from alignment files, previously obtained using *STAR* aligner

on the RNA-seq data. Following this, we utilised *JunctionSeq* to analyse differential isoform usage

between the sexes, as well as between treatments. In both, we restricted our analyses to features which

averaged >50 reads across all replicates. A gene was considered to have significant differential isoform

usage between sexes if the gene-wise p-adjusted was below 0.1. For comparisons between treatments,

a gene was considered to have significant differential isoform usage if p-adjusted < 0.1. The latter set of

genes are referred to as 'TDS genes' (for 'Treatment Differentially Spliced genes').

Identification of Gonad-Specific Genes

Patterns of differential expression or isoform usage in whole bodies might be driven in part by the

gonads. To examine this possibility, we repeated our analyses for the whole bodies without genes that

are expressed specifically in the gonads ('gonad-specific genes', GSGs). We identified GSGs using tissue-

specific estimates of gene expression from the supplementary data of Witt et al. (2017), which in turn is

based on RNAseq data from FlyAtlas2 (Leader et al., 2018). Each gene's expression in a tissue was

expressed in terms of FPKM. We estimated a 'gonad specificity index' (GSI) separately for males and

females, as per the following expression:

$$\mathrm{GSI_j} = \frac{\mathrm{FPKM_{gonads,j}}}{\Sigma_i \mathrm{FPKM_{i,j}}},$$

where *j* = sex and *i* = tissue. For the denominator, we only included FPKMs from non-overlapping tissues.

Any gene that yielded a GSI > 0.95 in either sex was designated as gonad-specific. We then proceeded to

repeat our analyses excluding this set of genes.

Results

Divergence in expression is non-random with respect to SBGE

We first examined the frequency of SBGE within each mating treatment. The frequency of genes with significant sex bias in expression was similar across treatments in whole bodies and in heads (Table 1). Though we found statistically significant variation in the frequency of SBGE among treatments in the whole body, these differences are not drastic. This is not surprising because, on a short microevolutionary time, one would not expect large scale expression divergence that would cause genes to change categories of SBGE (especially given that SBGE is fairly consistent across species; Zhang et al. 2007). However, this does not preclude the possibility of many subtle, quantitative changes in expression.

252 Table 1: Frequency of genes with significant sex-biased gene expression (p-adj < 0.05) and significant

253 sex-specific splicing (SSS; p-adj < 0.1) for whole bodies and heads. P-values are from Fisher's exact tests

254 for non-random association between the percentage of genes with SBGE/SSS and mating treatments.

255 For sex-biased gene expression, the numerator includes only genes that are sex-biased (i.e., $|\log_2FC| >$

256 0.5) with p-adj < 0.05, while the denominator includes all genes.

Mating treatment	Sample	Sex-biased gene expression		Sex-specific splicing	
		Percent of genes with SBGE	p-value	Percent of genes with SSS	p-value
MCabs		88.4% (10357/11717)		49.8% (3614/7250)	
MCsim	Body	87.6% (10266/11717)	**3.4 x 10⁻⁵**	48.1% (3484/7250)	0.097
MCcom		87.2% (10222/11717)		49.0% (3552/7250)	
MCabs		9.8% (913/9356)		4.2% (316/7547)	
MCsim	Head	10.4% (973/9356)	0.56	13.6% (1023/7547)	**<2.2 x 10⁻¹⁶**
MCcom		10.5% (978/9356)		10.4% (785/7547)	

To that end, we performed pairwise contrasts of mating treatments within each sex to identify genes

that were differentially expressed between treatments (hereafter 'Treatment Differentially Expressed',

TDE, genes). Though a relatively small number of genes qualified as TDE (Table S1), we considered

whether these TDE genes are disproportionally represented among different categories of sex-biased

expression. If genes that are sex-biased had previously evolved to be so because of a history sex-

differential selection arising from mate competition, one might expect that sex-biased genes would be

disproportionately likely to diverge among populations evolving in varying mating treatments. However,

we may not expect male- vs. female-biased genes to be equally likely to diverge in expression among

mating treatments, given results of past studies: relative to the female-biased and unbiased genes,

male-biased genes have been reported to evolve higher rates of evolution in sequence (Zhang et al.,

2004) and expression (Meiklejohn et al., 2003).

In female bodies, TDE genes are significantly more common among sex-biased than unbiased genes

(Table 2), with the exception of the MCabs-MCsim comparison for which too few TDE genes exist to

draw any conclusions. No significant difference is observed in the frequency of TDE genes among

unbiased vs. sex-biased genes in comparisons for male bodies, however. Comparing male- vs. female-

biased genes, TDE genes are significantly more frequent among male-biased genes in both male and

female body samples (again ignoring the MCabs - MCsim comparison due to the low total number of

TDE genes). Both of these patterns hold to some degree in heads too: in those comparisons with

statistically significant differences, TDE genes are more common among sex-biased than unbiased genes

and TDE genes are more common among male- than female-biased genes.

A possible proximate mechanism for the body results is a change in the relative sizes of gonads (Mank,

2017). We repeated the analysis presented in Table 2 after excluding genes with highly gonad-specific

expression. The results remain similar (Table S2). While this does not rule out the possibility that

changes in gonads play an important role in expression divergence among treatments, the fact that the

patterns are not weakened after removal of gonad-specific genes—along with the existence of the

patterns in heads—suggests that there is more to the underlying mechanisms behind expression

divergence than solely a change in gonad size.

We analysed the TDE genes for significant enriched biological processes or functions using a gene

ontology enrichment test (Eden et al., 2009), but failed to find any significant associations. However,

among the 574 TDE genes in total, we found 16 genes that putatively code for seminal fluid proteins

(SFPs), identified in a recent study of the seminal fluid proteome (Wigby et al., 2020). One of these 16

putative SFP genes (CG42807) was divergently expressed in two pairwise comparisons (MCabs vs

MCcom, MCsim vs MCcom).

Table 2: Results of Fisher's exact tests for the proportion of treatment differentially expressed (TDE)

genes in each sex bias category. Values in parentheses represent the fraction of TDE genes out of all

genes in the sex bias category.

Sample	Comparison	Percent TDE genes		p-value (UB vs SB)	Percent TDE genes		p-value (MB vs FB)
		UB	SB		FB	MB	
Female body	MCabs vs MCsim	0% (0/1693)	0.03% (2/7025)	1	0.03% (1/3721)	0.03% (1/3304)	1
	MCabs vs MCcom	0.5% (9/1693)	2.0% (137/7025)	**7.0 x 10^{-6}**	0.7% (26/3721)	3.4% (111/3304)	**2.3 x 10^{-16}**
	MCsim vs McCom	1.2% (21/1693)	2.5% (177/7025)	**1.0 x 10^{-3}**	0.8% (29/3721)	4.5% (148/3304)	**<2.2 x 10^{-16}**
Male body	MCabs vs MCsim	0.5% (9/1699)	2.5% (23/9101)	0.082	0.1% (5/3546)	0.3% (18/5555)	0.13
	MCabs vs MCcom	1.6% (27/1699)	1.5% (139/9101)	0.83	0.6% (21/3546)	2.1% (118/5555)	**1.1 x 10^{-9}**
	MCsim vs MCcom	0.7% (12/1699)	0.8% (75/9101)	0.77	0.3% (9/3546)	1.2% (66/5555)	**3.0 x 10^{-7}**
Female head	MCabs vs MCsim	0.7% (54/7767)	3.7% (32/872)	**1.5 x 10^{-11}**	3.4% (24/698)	4.6% (8/174)	0.50
	MCabs vs MCcom	0.3% (27/7767)	0.8% (7/872)	0.076	0.4% (3/698)	2.3% (4/174)	**0.032**
	MCsim vs MCcom	0.3% (26/7767)	1.0% (9/872)	**6.6 x 10^{-3}**	0.9% (6/698)	1.7% (3/174)	0.39
Male head	MCabs vs MCsim	0.5% (35/7774)	1.1% (10/874)	**0.021**	1.2% (8/688)	1.1% (2/186)	1
	MCabs vs MCcom	1.4% (108/7774)	3.0% (26/874)	**1.2 x 10^{-3}**	2.2% (15/688)	5.9% (11/186)	**0.013**
	MCsim vs MCcom	0.7% (56/7774)	1.9% (17/874)	**1.2 x 10^{-3}**	1.7% (12/688)	2.7% (5/186)	0.38

Divergence cannot be summarized as feminization/masculinization of the transcriptome

In the previous section we tested whether different types of genes (with respect to sex-bias) are more

likely to diverge, but we did not test for patterns in the direction and magnitude of divergence. In their

study, Hollis et al. (2014) predicted that, relative to polygamy, expression would be feminized in

populations evolving in monogamy, i.e., female-biased genes evolve to be upregulated in monogamy,

while male-biased genes become downregulated. Similar patterns are predicted in both female and

male tissues due to the presumed shared genetic architecture that constrains dimorphism under any

given mate competition regime (Prasad et al. 2007; Hollis et al. 2014).

We evaluated expression divergence in our populations from this Hollis et al. perspective by examining

divergence separately for female- and male-biased genes, focusing on the contrasts of our monogamy

treatment (MCabs) with each the two polygamy treatments (MCsim, MCcom). To visualise and test for

patterns of transcriptomic feminization/masculinization, we used the 'treatment effect' for each gene as

a measure of expression change, defined as the logarithmic ratio of expression in treatment 1 relative to

treatment 2. A positive treatment effect in a male-biased gene, for instance, would imply that its

expression is relatively masculinized in treatment 1; for female-biased genes, a positive treatment effect

would similarly imply relatively feminized expression in treatment 1. (We included unbiased genes for

completeness but these are not pertinent to the Hollis et al. prediction.) Results (Fig. 1) suggest that the

relationship between treatment effect and sex bias is inconsistent across treatment pairs, sexes and

sample types (head/body). Matching the Hollis et al. (2014) prediction, there is an overall feminization

of the female transcriptome in MCabs relative to the other two treatments, with a significant increase in

expression of female-biased genes and a significant decrease in expression of male-biased genes in

MCabs (Fig. 1A-B). In male bodies (Fig. 1D-E), the net changes are smaller, but more importantly, the

pattern is different. Though female-biased genes have increased expression in MCabs relative to the

other two treatments (similar to the female body result), male-biased genes are not significantly

reduced in expression in MCabs relative to MCsim and, in fact, have significantly increased expression in

MCabs relative to MCcom (i.e., opposite the prediction). A further layer of complication is added when

examining the heads, where the results differ between female and male samples and in neither sex is

there a consistent pattern of the predicted feminization of MCabs (Fig 1G-H, K-L).

Fig. 1 depicts mean treatment effects in discrete, and somewhat arbitrarily-bounded categories of sex

bias. We also performed LOESS regressions of treatment effect against continuously varying sex bias to

discern any patterns missed by treating variation in sex-biased gene expression as discrete. These

regressions (Fig. S2) show that the relationship between treatment effect and sex bias can vary within a

sex bias category, a nuance not captured in the discrete version of the plots. However, the overall sign

of mean treatment effect for a sex bias category is largely consistent in the two analyses.

As another approach to evaluating the Hollis et al. prediction, we looked for evidence of net changes in

feminization/masculinization via a different method using only the TDE genes. In female bodies, there is

a significant net feminization of TDE genes in MCabs relative to MCcom (Fig. S3), similar to the results

depicted in Fig. 1 and matching the Hollis et al. prediction. In male bodies, there is a significant net

masculinization of TDE genes in MCabs relative to MCcom (Fig. S3), opposite to the Hollis et al.

prediction. In most other contrasts, there is no significant net feminization or masculinization.

Figure 1: Average treatment effect (± 1 SE) for three sex bias categories for female body (A-C), male body (D-F), female head (G-I), male head (K-M), in pairwise comparisons between mating treatments. Asterisks denote the mean treatment effect is significantly different from zero, determined using one-sample permutation tests (i.e., by changing the sign of the treatment effect of a randomly chosen 50% of the genes and re-calculating the mean in each permutation). Significance is denoted as follows: * p < 0.05, ** p < 0.01, *** p < 0.001. Panel N summarises the order of expression levels for female- and male-biased genes in panels A-M; tilda (~) denotes no significant difference. An analogous figure from an analysis excluding gonad-specific genes is shown in Fig. S1; most of the patterns are very similar.

Changes in degree of sexual dimorphism

In the previous section we examined expression changes for each sex separately, but these within-sex changes are not directly informative of changes in sexual dimorphism. Here we investigate differences

between treatments in the degree of expression dimorphism across genes with different levels of sex-

bias as determined from an external dataset (see Methods). We used LOESS regressions to evaluate how

treatment differences in 'local average dimorphism' varied with sex-bias (where local average refers to

the difference in dimorphism averaged across a local range of sex-bias). Patterns were complex (Fig. 2)

though treatment differences in local average dimorphism did tend to be larger for sex-biased than

unbiased genes. Among the sex-biased genes, the magnitude and even direction of treatment

differences in local average dimorphism varied between, but also within, sex-bias categories. In bodies,

local average dimorphism is highest in MCsim for both highly female- and male-biased genes, but

patterns differ for moderately-biased genes (summarized in Fig. 2G). In heads, where phenotypic sexual

dimorphism is not as marked, changes in local average dimorphism are detectable despite the relatively

low number of sex-biased genes. Female-biased genes are, on average, most dimorphic in MCcom, but

male-biased genes are least dimorphic in MCcom. Analyses excluding gonad-specific genes yielded

nearly identical results.

Figure 2: Treatment difference in dimorphism as a function of sex-biased gene expression in A-C)

body and D-F) head. The vertical axis is (sign of sex bias)*(difference in dimorphism between

treatments) so that positive (negative) values indicate increased (reduced) dimorphism in treatment

1 relative to treatment 2. In panels A-F, the fitted line is obtained from LOESS regressions (fit using

geom_smooth in the R package *ggplot2* with the width of the sliding-window set at 0.5); 95%

confidence intervals are indicated in green. The plots show a subset of the genes within the vertical

axis range [-1,1] to better display the pattern for the majority of genes. Dashed vertical lines

demarcate the sex bias categories listed in the panel G. Panel G summarises the relative strength of

dimorphism among treatments for each SBGE category. The sex bias categories used in panel G are

designated as follows: highly FB genes = (-∞,-5], moderately FB genes = (-5, -0.5], moderately MB

genes = [0.5, 5), [5, ∞). No highly male- or female-biased genes are depicted for heads due to the

low numbers of such genes.

 Differential splicing amongst mating treatments

We also examined sex-specific splicing (SSS), testing whether alternative splicing differs between mating

treatments. Congruent to our analysis for TDE genes, we tested for genes with significant differential

splicing between treatments, i.e., 'treatment differentially spliced' (TDS) genes. Frequencies of TDS

genes are in the range of 0.2-2.3% of all genes analysed (Table S3). We performed Fisher's exact tests to

ascertain if TDS genes are disproportionately represented among sex-biased genes. In contrast to the

analogous test for TDE genes (Table 2, S2), there was no over-representation of TDS genes among sex-

biased genes, though there was some evidence of TDS genes being more common among male- than

female-biased genes (Tables S4, S5). Considering TDE and TDS together, there was significant overlap

between these in most sample types (Table S6).

We also tested for sex-specific splicing (SSS) within each treatment. The proportion of SSS genes was

similar among treatments, with one exception: the proportion of such genes is noticeably lower in the

heads in MCabs than the other two treatments (Table 1). The lower frequency of SSS in MCabs heads is

unlikely to be due to reduced power to detect SSS: MCabs heads had the highest mean coverage (Table

S7) and there was no comparable in reduction in the frequency of sex-biased expression (Table 1). This

difference in treatment effects on the prevalence SBGE and SSS suggests that these two forms of

expression dimorphism evolve somewhat independently.

Discussion

Differential selection on the sexes is thought to drive the evolution of sexual dimorphism in gene

expression (Parsch & Ellegren, 2013). Selection arising from sexual interactions and mate competition is

hypothesized to be a major cause of this differential selection. Thus, variation in mate competition

among populations or species should be a major source of variation in SBGE across such taxonomic

groups. Here, we tested this idea by measuring shifts in gene expression in response to evolutionary

manipulations of mate competition in *D. melanogaster*. The MCabs treatment eliminates mate

competition through enforced monogamy, while the two other treatments (MCsim, MCcom)

incorporate mate competition in differing mating environments that we know alter intersexual

interactions and the selection these generate (MacPherson et al., 2018; Yun et al., 2017, 2019).

If a history of sex-differential selection arising from mate competition is responsible for pre-existing

SBGE, then we expect that selection on genes that are initially sex-biased is more likely to be sensitive to

a change in mate competition than selection on unbiased genes. Thus, we predicted that sex-biased

genes would be more likely to diverge among mate competition treatments than unbiased genes. The

results largely supported this prediction (Table 2); all seven of the comparisons in which there was a

significant difference were in the predicted direction (though not all comparisons yielded a significant

difference).

Though the patterns are consistent with the prediction, other explanations likely also contribute to this

pattern. Core house-keeping genes will tend to be unbiased and their expression may be under strong

stabilizing selection that changes little across any of a large range of environments. For this reason, we

might predict that unbiased genes are less likely to diverge in expression than sex-biased genes across

any environmental change, not just changes to mate competition. A better test, which may be possible

via meta-analysis in future work, would be to ask whether the preferential divergence of sex-biased

relative to unbiased genes is more pronounced in response to mate competition treatments than it is

with other types of environmental changes.

We also compared the relative frequency of DE genes in male- versus female-biased genes. Past studies

have shown that male-biased genes evolve more rapidly than female-biased genes, both in DNA

sequence (Meisel, 2011; Zhang et al., 2004) and gene expression (Allen et al., 2018; Meiklejohn et al.,

2003), though some studies describe more mixed results (Whittle & Johannesson, 2013; Yang et al.,

2016). Adaptive evolution in response to sexual selection has been invoked as an explanation (Ellegren

& Parsch, 2007), with some studies suggesting that male-biased genes are targeted more often by

strong positive selection (Pröschel et al., 2006; Zhang et al., 2004; but see Singh and Agrawal 2023)

presumably resulting from reproductive interactions. Our manipulation of mate competition resulted in

more divergence in expression of male- than female-biased genes (Table 2): in the six comparisons

where there was a significant difference, male-biased genes had a higher propensity for expression

divergence than female-biased genes. In some sense, this matches with fitness data showing stronger

signatures of local adaptation to the mate competition environment in males than females (Yun et al.

2019), suggesting greater phenotypic divergence in males than females. However, Table 2 contrasts

male- and female-biased genes within each sex, rather than contrasting divergence of males versus that

of females. There are not obvious differences in the proportion of significantly diverged genes between

male and female samples. This apparent discrepancy could be reconciled if the expression divergence of

male-biased genes has greater phenotypic and fitness consequences in males than females.

Hollis et al. (2014) predicted that enforced monogamy would result in a "feminization" of the

transcriptome (i.e., increased expression of female-biased genes and decreased expression of male-

biased genes) in both sexes. This prediction has now been tested in three experiments with highly

heterogeneous results. The data of Hollis et al. (2014) matched their prediction, while the results of

Veltsos et al. (2017) were mixed but largely in the opposite direction. Our results are mixed with respect

to the predicted changes.

The basis of the Hollis et al. prediction is that the primary consequence of enforced monogamy is the

removal of sexual selection on males. However, enforced monogamy not only removes sexual selection,

but it imposes (potentially strong) selection on males to inflict less harm on their mates (Holland and

Rice 1999; Martin and Hosken 2003; Crudginton et al. 2005, 2010; Yun et al. 2021). This can lead to

changes in selection on female resistance to harm (Holland & Rice 1999, Martin and Hosken 2003). Even

in the absence of evolved changes in male harm, there are strong reasons to suspect that enforced

monogamy will change selection on females because males will be unable to bias their attention

towards particular phenotypes (Long et al., 2009; Arbuthnott and Rundle 2012; Chenoweth et al. 2015;

Yun et al. 2017). Considering this multitude of possible changes in selection, it is difficult to predict how

the transcriptome will respond, and it seems unlikely that net change will be consistent or easily

classifiable as 'feminization' or 'masculinization.'

Because selection in enforced monogamy can differ from selection under mate competition in a variety

of ways, other details could become important to how divergence occurs. These three studies differ in

various ways which could, in principle, contribute to the among-study heterogeneity in results (e.g.,

species studied, nature of starting variation, food, temperature, density and population size, and

maintenance procedures (Li Richter & Hollis 2021)). One potentially important aspect is the precise

nature of mate competition, which can take many forms (e.g., there is no one true form of "polygamy"

for a "monogamy vs. polygamy" contrast). Our study directly shows that differences in how mate

competition occurs has major consequences for expression evolution. Within the context of our

experiment, where most other factors are not varied among treatments, the transcriptional differences

between our two different mate competition treatments (i.e., MCsim vs. MCcom) are generally as large

as those between either one of these with the enforced monogamy treatment (MCabs).

Though selection arising from mate competition is almost certainly an important driver of SBGE (Parsch

& Ellegren, 2013), there is little direct evidence (Harrison et al. 2014). It seems intuitive that, in the

absence of mate competition, expression dimorphism would become reduced. For example,

monogamous bird species are often less dimorphic with respect to plumage (and other traits) than non-

monogamous species, and presumably this reflects a reduction in expression dimorphism for genes

affecting such traits. Yet, it is unclear what the prediction for genome-wide expression dimorphism

should be as intralocus conflict on expression can easily exist under monogamy given that males and

females play different roles in reproduction and offspring rearing, even if overt traits such as plumage

do not experience divergent selection.

While the underlying motivation for the studies of Hollis et al. (2014) and Veltsos et al. (2017) was based

around sex-biased gene expression, neither attempted to examine how enforced monogamy changed

expression dimorphism per se. We examined how the average treatment effect on expression

dimorphism varied across local ranges of sex bias as determined in LOESS regressions. We found

differences in 'local average dimorphism' among mating regimes. These changes in local average

dimorphism tend to occur to a greater extent among sex-biased than unbiased genes, and more so in

bodies, which have much more sex-biased gene expression than heads (Fig. 2). However, we did not find

a consistent reduction in local average dimorphism between the enforced monogamy treatment and the

two treatments with mate competition; while some categories of sex-biased genes became less

dimorphic under monogamy, others became more so (Fig. 2). For example, relative to MCsim, MCabs

evolved reduced local average dimorphism of strongly sex-biased genes but increased dimorphism of

moderately male-biased genes in body samples. However, one aspect of expression was consistent with

the intuition of reduced dimorphism under monogamy: there was a notable reduction in the frequency

of genes with sex-specific splicing in heads of MCabs relative to either of the other two treatments,

though not in bodies (Table 1). While our experiment is the result of more than 100 generations of

selection, this is a relatively short period relative to other evolutionary time scales (e.g., sister species);

perhaps a more consistent reduction in expression dimorphism would be evident after enough time.

Nonetheless, this study shows that, though expression dimorphism readily evolves among mate

competition environments, enforced monogamy does not necessarily cause a rapid and consistent

reduction in dimorphism.

The most striking change in local average dimorphism occurred between the two treatments involving

mate competition, with MCsim showing greater dimorphism in bodies than MCcom across all sex bias

categories (Fig. 2C). Relative to MCcom, the MCsim environment is one in which intersexual

interactions, including mating, occur more frequently (Yun et al. 2017, 2019); exposure to males is more

harmful to females in the simple than complex environment and MCsim males have evolved to be more

harmful (Yun et al. 2017, 2021). Thus, it is reasonable to infer that interlocus sexual conflict is relatively

more important in MCsim than MCcom. However, this does not provide an obvious explanation for the

observed difference in expression dimorphism, as dimorphism is thought to evolve in response to intra-,

rather than interlocus, conflict.

In this study, we have treated each gene's expression as a trait (i.e., a measurable property) and

examined patterns in how these 'traits' change in response to an evolutionary manipulation of mate

competition. Just as with traditional traits (i.e., morphological, behavioral, or physiological phenotypes),

multiple possible genetic mechanisms could underly each change and a single genetic change could

affect multiple traits. At one (unlikely) extreme, every gene expression change could be due to a change

in that gene's *cis*-regulatory region. At the other extreme, a single genetic change could affect

expression levels of many genes, possibly by changing the size of different organs or relative abundance

of cell types within organs (Stewart et al. 2010). The truth likely lies somewhere between these two

extremes. In an evolutionary manipulation of mating system in *Drosophila pseudoobscura*, Wiberg et al.

(2021) found that genomically diverged sites were enriched near sites with expression divergence. That

result would not be expected if a very small number of genetic changes underlay most expression

differences, but it also does not imply that most expression changes are due to individual *cis*-regulatory

changes.

It is not our objective here to resolve this issue for our own experiment (and we have little ability to do

so). In the spirit of many past transcriptomic studies, we test for patterns among different 'types' of

traits (e.g., genes with male-biased, female-biased, or unbiased expression). From this perspective, each

gene represents a separate instance of that trait 'type' but it is important to remember that variation in

each of those individual 'traits' need not be governed by independent proximate mechanisms. This

unknown level of independence should temper inferences about genetic changes and the true nature

and number of targets of selection (e.g., could some patterns result from changes in the relative size of

gonads and, if so, why has selection caused such changes?). Our study clearly reveals patterns of

expression evolution across SBGE categories in response to changes in mate competition, though we are

limited in interpreting the reasons for these patterns. The type of work presented here represents only

one step towards understanding the relationship between mate competition and SBGE, which can help

evaluate hypotheses in the literature and provide fodder for new ones.

Acknowledgements

We thank George Sandler for doing the RNA extractions. This work was supported by the Natural Sciences and Engineering Research Council of Canada (AFA & HDR).

Literature Cited

Allen, S. L., R. Bonduriansky, and S. F. Chenoweth. 2018. Genetic constraints on microevolutionary

divergence of sex-biased gene expression. Philos. Trans. R. Soc. Lond. B Biol. Sci. 373:20170427.

Anders, S., P. T. Pyl, and W. Huber. 2015. HTSeq--a Python framework to work with high-throughput

sequencing data. Bioinformatics 31: 166-169.

Arbuthnott, D., and H. D. Rundle. 2012. Sexual selection is ineffectual or inhibits the purging of

deleterious mutations in *Drosophila melanogaster*. Evolution 66:2127–2137.

Arnqvist, G., and L. Rowe. 2005. Sexual conflict. Princeton University Press, Princeton, NJ.

Bonduriansky, R., and S. F. Chenoweth. 2009. Intralocus sexual conflict. Trends Ecol. Evol. 24:280–288.

Chapman, T., L. F. Liddle, J. M. Kalb, M. F. Wolfner, and L. Partridge. 1995. Cost of mating in *Drosophila*

melanogaster females is mediated by male accessory gland products. Nature 373:241–244.

nature.com.

Chenoweth, S. F., N. C. Appleton, S. L. Allen, and H. D. Rundle. 2015. Genomic evidence that sexual

selection impedes adaptation to a novel environment. Curr. Biol. 25:1860–1866.

Crudgington, H. S., A. P. Beckerman, L. Brüstle, K. Green, and R. R. Snook. 2005. Experimental removal

and elevation of sexual selection: does sexual selection generate manipulative males and resistant

females? Am. Nat. 165: S72–87.

Crudgington, H. S., S. Fellows, and R. R. Snook. 2010. Increased opportunity for sexual conflict promotes

harmful males with elevated courtship frequencies. J. Evol. Biol. 23:440–446.

Dean, R., A. E. Wright, S. E. Marsh-Rollo, B. M. Nugent, S. H. Alonzo, and J. E. Mank. 2017. Sperm

competition shapes gene expression and sequence evolution in the ocellated wrasse. Mol. Ecol.

26:505–518.

Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, and T. R.

Gingeras. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15–21.

dos Santos, G., A. J. Schroeder, J. L. Goodman, V. B. Strelets, M. A. Crosby, J. Thurmond, D. B. Emmert,

W. M. Gelbart, and FlyBase Consortium. 2015. FlyBase: introduction of the *Drosophila*

melanogaster Release 6 reference genome assembly and large-scale migration of genome
annotations. Nucleic Acids Res. 43:D690–7.

Eden, E., R. Navon, I. Steinfeld, D. Lipson, and Z. Yakhini. 2009. GOrilla: a tool for discovery and
visualization of enriched GO terms in ranked gene lists. BMC Bioinformatics 10:48.

Ellegren, H., and J. Parsch. 2007. The evolution of sex-biased genes and sex-biased gene expression. Nat.
Rev. Genet. 8:689–698.

Fernandes Martins, M. J., G. Hunt, R. Lockwood, J. P. Swaddle, and D. J. Horne. 2017. Correlation
between investment in sexual traits and valve sexual dimorphism in Cyprideis species (Ostracoda).
PLoS One 12:e0177791.

Fowler, K., and L. Partridge. 1989. A cost of mating in female fruitflies. Nature 338:760–761.

Grath, S., and J. Parsch. 2016. Sex-biased gene expression. Annu. Rev. Genet. 50:29–44.

Harrison, P. W., A. E. Wright, F. Zimmer, R. Dean, S. H. Montgomery, M. A. Pointer, and J. E. Mank. 2015.
Sexual selection drives evolution and rapid turnover of male gene expression. Proc. Natl. Acad.
Sci. U. S. A. 112:4393–4398.

Hartley, S. W., and J. C. Mullikin. 2016. Detection and visualization of differential splicing in RNA-Seq
data with JunctionSeq. Nucleic Acids Res. 44:e127.

Holland, B., and W. R. Rice. 1999. Experimental removal of sexual selection reverses intersexual
antagonistic coevolution and removes a reproductive load. Proc. Natl. Acad. Sci. U. S. A. 96:5083–
5088.

Hollis, B., D. Houle, Z. Yan, T. J. Kawecki, and L. Keller. 2014. Evolution under monogamy feminizes gene
expression in *Drosophila melanogaster*. Nat. Commun. 5:3482.

Ingleby, F. C., I. Flis, and E. H. Morrow. 2014. Sex-biased gene expression and sexual conflict throughout
development. Cold Spring Harb. Perspect. Biol. 7:a017632.

Lande, R. 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic characters. Evolution
34:292–305. JSTOR.

Leader, D. P., S. A. Krause, A. Pandit, S. A. Davies, and J. A. T. Dow. 2018. FlyAtlas 2: a new version of the

Drosophila melanogaster expression atlas with RNA-Seq, miRNA-Seq and sex-specific data. Nucleic

Acids Res. 46:D809–D815.

Li Richter, X.-Y., and B. Hollis. 2021. Softness of selection and mating system interact to shape trait

evolution under sexual conflict. Evolution 75:2335–2347.

Long, T. A. F., A. Pischedda, A. D. Stewart, and W. R. Rice. 2009. A cost of sexual attractiveness to high-

fitness females. PLoS Biol. 7:e1000254.

Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 15:550.

MacPherson, A., L. Yun, T. S. Barrera, A. F. Agrawal, and H. D. Rundle. 2018. The effects of male harm

vary with female quality and environmental complexity in *Drosophila melanogaster*. Biol. Lett.

14:20180443.

Martin, O. Y., and D. J. Hosken. 2003. Costs and benefits of evolving under experimentally enforced

polyandry or monogamy. Evolution 57:2765–2772.

Meiklejohn, C. D., J. Parsch, J. M. Ranz, and D. L. Hartl. 2003. Rapid evolution of male-biased gene

expression in *Drosophila*. PNAS 100:9894 –9899.

Osada, N., R. Miyagi, and A. Takahashi. 2017. Cis- and trans-regulatory effects on gene expression in a

natural population of *Drosophila melanogaster*. Genetics 206:2139–2148.

Parsch, J., and H. Ellegren. 2013. The evolutionary causes and consequences of sex-biased gene

expression. Nat. Rev. Genet. 14:83–87.

Partridge, L., and K. Fowler. 1990. Non-mating costs of exposure to males in female *Drosophila*

melanogaster. J. Insect Physiol. 36:419–425.

Pointer, M. A., P. W. Harrison, A. E. Wright, and J. E. Mank. 2013. Masculinization of gene expression is

associated with exaggeration of male sexual dimorphism. PLoS Genet. 9:e1003697.

Prasad, N. G., S. Bedhomme, T. Day, and A. K. Chippindale. 2007. An evolutionary cost of separate

genders revealed by male-limited evolution. Am. Nat. 169:29–37.

Pröschel, M., Z. Zhang, and J. Parsch. 2006. Widespread adaptive evolution of *Drosophila* genes with sex-
biased expression. Genetics 174:893–900.

Rowe, L., and H. D. Rundle. 2021. The alignment of natural and sexual selection. Annu. Rev. Ecol. Evol.
Syst. 52:499–517.

Singh, A., and A. F. Agrawal. 2023. Two forms of sexual dimorphism in gene expression in *Drosophila*
melanogaster: their coincidence and evolutionary genetics. Mol. Biol. Evol. 40.

Stewart, A. D., A. Pischedda, and W. R. Rice. 2010. Resolving intralocus sexual conflict: genetic
mechanisms and time frame. J. Hered. 101:S94–9.

Stuglik, M. T., W. Babik, Z. Prokop, and J. Radwan. 2014. Alternative reproductive tactics and sex-biased
gene expression: the study of the bulb mite transcriptome. Ecol. Evol. 4:623–632.

Veltsos, P., Y. Fang, A. R. Cossins, R. R. Snook, and M. G. Ritchie. 2017. Mating system manipulation and
the evolution of sex-biased gene expression in *Drosophila*. Nat. Commun. 8:2072.

Whittle, C. A., and H. Johannesson. 2013. Evolutionary dynamics of sex-biased genes in a hermaphrodite
fungus. Mol. Biol. Evol. 30:2435–2446.

Wiberg, R. A. W., P. Veltsos, R. R. Snook, and M. G. Ritchie. 2021. Experimental evolution supports
signatures of sexual selection in genomic divergence. Evol Lett 5:214–229.

Wigby, S., N. C. Brown, S. E. Allen, S. Misra, J. L. Sitnik, I. Sepil, A. G. Clark, and M. F. Wolfner. 2020. The
Drosophila seminal proteome and its role in postcopulatory sexual selection. Philos. Trans. R. Soc.
Lond. B Biol. Sci. 375:20200072.

Wigby, S., and T. Chapman. 2004. Female resistance to male harm evolves in response to manipulation
of sexual conflict. Evolution 58:1028–1037.

Witt, E., N. Svetec, S. Benjamin, and L. Zhao. 2021. Transcription factors drive opposite relationships
between gene age and tissue specificity in male and female *Drosophila* gonads. Mol. Biol. Evol.
38:2104–2115.

Yang, L., Z. Zhang, and S. He. 2016. Both male-biased and female-biased genes evolve faster in fish
genomes. Genome Biol. Evol. 8:3433–3445.

Yun, L., A. F. Agrawal, and H. D. Rundle. 2021. On male harm: How it is measured and how it evolves in
different environments. Am. Nat. 198:219–231.

Yun, L., M. Bayoumi, S. Yang, P. J. Chen, H. D. Rundle, and A. F. Agrawal. 2019. Testing for local
adaptation in adult male and female fitness among populations evolved under different mate
competition regimes. Evolution 73:1604–1616.

Yun, L., P. J. Chen, K. E. Kwok, C. S. Angell, H. D. Rundle, and A. F. Agrawal. 2018. Competition for mates
and the improvement of nonsexual fitness. Proc. Natl. Acad. Sci. U. S. A. 29:201805435.

Yun, L., P. J. Chen, A. Singh, A. F. Agrawal, and H. D. Rundle. 2017. The physical environment mediates
male harm and its effect on selection in females. Proc. Biol. Sci. 284:20170424.

Zhang, Y., D. Sturgill, M. Parisi, S. Kumar, and B. Oliver. 2007. Constraint and turnover in sex-biased gene
expression in the genus *Drosophila*. Nature 450:233–237.

Zhang, Z., T. M. Hambuch, and J. Parsch. 2004. Molecular evolution of sex-biased genes in *Drosophila*.
Mol. Biol. Evol. 21:2130–2139.