Original Article

Novel Insights into the Aortic Mechanical Properties of Mice Modeling

Hereditary Aortic Diseases

Nicolo Dubacher, Kaori Sugiyama, Jeffrey D. Smith, Vanessa Nussbaumer, Máté Csonka, Szilamér

Ferenczi, Krisztina J. Kovács, Sylvan M. Caspar, Lisa Lamberti, Janine Meienberg, Hiromi

Yanagisawa*, Mary B. Sheppard*, Gabor Matyas*

Affiliations

Center for Cardiovascular Genetics and Gene Diagnostics, Swiss Foundation for People with Rare

Diseases, Switzerland (N.D., V.N., M.C., S.M.C., L.L., J.M., G.M.), Translational Cardiovascular

Technologies, Department of Health Sciences, ETH Zurich, Switzerland (N.D.), Institute for

Advanced Research of Biosystem Dynamics, Research Institute for Science and Engineering, Waseda

University, Japan (K.S.), Life Science Center for Survival Dynamics, Tsukuba Advanced Research

Alliance, University of Tsukuba, Japan (K.S., H.Y.), Saha Cardiovascular Research Center, University

of Kentucky, USA (J.D.S.), Heart and Vascular Center, Semmelweis University, Hungary (M.C.),

Laboratory of Molecular Neuroendocrinology, Institute of Experimental Medicine, Hungary (S.F.,

K.J.K.), Department of Family and Community Medicine, University of Kentucky, USA (M.B.S.),

Saha Aortic Center, University of Kentucky, USA (M.B.S.), and Zurich Center for Integrative Human

Physiology, University of Zurich, Switzerland (G.M.).

*H. Yanagisawa, M.B. Sheppard, and G. Matyas are joint senior authors.

Short title: Aortic Rupture Force in Mice Modeling Aortic Diseases

Correspondence to: Gabor Matyas, PhD, Center for Cardiovascular Genetics and Gene Diagnostics,

Swiss Foundation for People with Rare Diseases, 8952 Schlieren-Zurich, Switzerland. Email

matyas@genetikzentrum.ch, Phone +41-43-433-86-86

ABSTRACT

OBJECTIVE: Hereditary aortic diseases (hADs) increase the risk of aortic dissections and ruptures. Recently, we have established an objective approach to measure the rupture force of the murine aorta, thereby explaining the outcomes of clinical studies and assessing the added value of approved drugs in vascular Ehlers-Danlos syndrome (vEDS). Here, we applied our approach to six additional mouse hAD models.

APPROACH AND RESULT: We used two mouse models of Marfan syndrome (MFS) as well as one smooth-muscle-cell-specific knockout (SMKO) of *Efemp2* and three CRISPR/Cas9-engineered knock-in models (*Ltbp1*, *Mfap4*, and *Timp1*). Moreover, one of the two MFS models was subjected to 4-week-long losartan treatment. Per mouse, three rings of the thoracic aorta were prepared, mounted on a tissue puller, and uniaxially stretched until rupture. The aortic rupture force of the SMKO and both MFS models was significantly lower compared with wild-type mice but in both MFS models higher than in mice modeling vEDS. In contrast, the *Ltbp1*, *Mfap4*, and *Timp1* knock-in models presented no impaired aortic integrity. As expected, losartan treatment reduced aneurysm formation but surprisingly had no impact on the aortic rupture force of our MFS mice.

CONCLUSIONS: We demonstrate for the first time that our read-out system is able to characterize the aortic biomechanical integrity of mice modeling not only vEDS but also related hADs, allowing the aortic-rupture-force-focused comparison of mouse models. Furthermore, aneurysm progression alone may not be a sufficient read-out for aortic rupture, as blood-pressure-lowering therapies preventing aortic aneurysms might not strengthen the weakened aortic wall. These results may contribute to better medical therapies of hADs.

Key Words: aortic aneurysm, dissection, rupture, biomechanical integrity, *Efemp2*, *Fbn1*, Marfan syndrome

    Nonstandard Abbreviations and Acronyms

hAD(s)   hereditary aortic disease(s)

AGTR1   angiotensin II receptor type 1

Bgn   biglycan (encoding gene in mice)

ECM   extracellular matrix

Efemp2   fibulin-4 (Fbln4) (encoding gene in mice)

Fbn1   fibrillin-1 (encoding gene in mice)

Ltbp1   latent TGF-β binding protein 1 (encoding gene in mice)

Mfap4   microfibril associated protein 4 (encoding gene in mice)

MFS   Marfan syndrome

MMP   matrix metalloproteinase

SMKO   smooth-muscle-specific knockout (*Efemp2$^{loxP/KO}$;SM22-Cre$^{Tg/Tg}$*)

TAA(s)   thoracic aortic aneurysm(s)

TGF   transforming growth factor

Timp1   tissue inhibitor of matrix metalloproteinase 1 (encoding gene in mice)

vEDS   vascular Ehlers-Danlos syndrome

Hereditary aortic diseases (hADs) with life-threatening aneurysms and dissections are

associated with high morbidity and mortality and are the cause for up to ~2% of all deaths in

the Western world.[1] Syndromic and non-syndromic forms of hADs are rare (i.e., prevalence

<1:2000) but several forms have been identified so far. Despite considerable phenotypic

similarity, disease management can profoundly differ depending on the disease-underlying

gene(s) and, thus, molecular genetic testing is required to identify (likely) pathogenic

sequence variant(s) in hAD-associated genes.[2,3] The advent of high-throughput sequencing

has enabled genetic data generation at hitherto unprecedented scale, allowing for novel

insights into the molecular etiology of hereditary (aortic) diseases but requiring a high level of

expertise for appropriate data analysis and variant interpretation.[4] Currently, at least 50 genes

have been associated with hereditary thoracic aortic aneurysms (TAAs) and classified based on the strength of association.[5] However, these genes cannot explain all the cases of heritable TAAs, suggesting the presence of additional genes which, when mutated, may be causative for hADs.[6]

Changes in the structural components of the extracellular matrix (ECM) as well as in chemical and mechanical stimuli can influence the molecular signaling pathways that regulate aortic wall homeostasis. Accordingly, hAD-causing sequence variants (mutations) can affect the proper ECM organization, thereby directly impacting the mechanical properties of the aorta as well as the mechanoregulation of vascular smooth muscle cells and their ability to sense matrix-mediated mechanical stimuli.[7] This can result in an impaired aortic integrity and mechanical functionality causing progressive weakening of the aortic wall, ultimately leading to arterial dissection and/or rupture.[8] The mechanical properties of the thoracic aorta can be derived from *in vivo* imaging parameters such as aortic wall distensibility and local pulse wave velocity or, especially aortic rupture, studied directly in *ex vivo* aortic samples.[9-11] In humans, however, studies assessing the clinically highly relevant rupture (with or without aneurysm/dissection) of the thoracic aorta in hAD are significantly limited. Thus, non-human models harboring (likely) disease-causing mutations in candidate or already established hAD-associated gene(s) are the (only) alternative in combination with an appropriate read-out at individual level.

Recently, we have established an objective approach to measure the aortic rupture force (maximum tensile force) of uniaxially stretched murine thoracic aortic rings as read-out for the biomechanical integrity of the aorta.[12] We have used this approach to clarify whether or not antihypertensive drugs may have an additional benefit (added value) to strengthen the weakened thoracic aorta (drug repurposing). Indeed, we showed that celiprolol, a 3rd-generation β-blocker, can improve the biomechanical integrity of the aortic wall (i.e., beyond the β-blocker-typical prevention of rise in blood pressure and heart rate),[12] explaining the

outcomes of clinical and retrospective studies in vascular Ehlers-Danlos syndrome (vEDS,

OMIM #130050).[13-15] This added value of celiprolol, however, cannot simply be extrapolated to

other β-blockers, as we have demonstrated for bisoprolol using aortic rupture force as read-out.[16]

Here, we applied our established read-out approach[12] to measure the aortic rupture

force in six additional mouse hAD models and compared their aortic rupture force with that of

vEDS mice: (i) We used the two most widely investigated mouse models of Marfan syndrome

(MFS, OMIM #154700) ($Fbn1^{C1041G}$ and $Fbn1^{mgR}$),[17-19] a rare autosomal-dominant inherited

disease with severe cardiovascular manifestations, such as TAAs predisposing to aortic

dissection and rupture, caused by mutations in the gene encoding fibrillin-1 (*FBN1* in

humans, *Fbn1* in mice).[20] In one of the two mouse MFS models ($Fbn1^{mgR}$), we also aimed to

clarify whether or not the frequently prescribed angiotensin-II-receptor-type-1 (AGTR1)

inhibitor losartan, which has been shown to slow down aneurysm progression in MFS,[21-25]

also strengthens the potentially weakened thoracic aorta. (ii) Furthermore, we assessed the

aortic rupture force in an *Efemp2* (encoding fibulin-4 [Fbln4]) smooth-muscle-specific knock-

out (SMKO) model recapitulating key vascular phenotypic features (aortic aneurysms and

arterial tortuosity)[26] of the autosomal-recessive inherited connective tissue disorder cutis laxa

type 1B (OMIM #614437).[27,28] (iii) Moreover, we used three novel CRISPR/Cas9-engineered

knock-in models ($Ltbp1^{G1132S}$, $Mfap4^{D42A}$, $Timp1^{I120T}$) to assess the causality of human-

equivalent variants in hAD candidate genes detected by whole-genome sequencing (WGS).[6]

METHODS

Mouse strains

In this study, multiple mouse strains were used. An overview can be found in the Major Resources Table in the Supplemental Material. Wild-type mice are referred to as $^{+/+}$.

$Fbn1^{C1041G}$ and $Fbn1^{mgR}$ Mouse Models

For the $Fbn1^{C1041G}$ mouse model (C57BL/6J background), 10-week-old heterozygous ($Fbn1^{+/C1041G}$, n=10) and wild-type ($Fbn1^{+/+}$, n=10) mice were ordered in frozen condition from Jackson Laboratory (JAX stock #012885, Bar Harbor, ME, USA). Tail biopsies were taken to confirm the genotypes of delivered mice, for which DNA extraction, PCR and subsequent Sanger sequencing were performed as previously described.[12,17]

$Fbn1^{mgR}$ mice (C57BL/6J background) were bred and maintained on a 14/10-hour light/dark cycle receiving standard rodent chow and water *ad libitum*. Mice were genotyped as described elsewhere.[18] In addition, 4-week-old homozygous ($Fbn1^{mgR/mgR}$, n=12) and wild-type ($Fbn1^{+/+}$, n=10) mice were treated for 4 weeks with losartan dissolved in drinking water at a concentration of 600 mg/L (LTK Labs, St. Paul, MN, USA, ~180 mg/kg bodyweight per day), whereas age-matched untreated $Fbn1^{mgR/mgR}$ (n=17) and $Fbn1^{+/+}$ (n=10) littermates receiving normal drinking water served as controls. Animals were weighed at the beginning and at the end of the treatment and water consumption was monitored regularly to ensure that control and treated mice drank comparable quantities. At the end of the treatment, mice were euthanized using CO_2 and subsequently frozen. Treatment protocols were approved by the University of Kentucky IACUC (approval reference number: 2016-2437) in line with the local guidelines.

Efemp2 SMKO Model

Efemp2$^{loxP/KO}$;*SM22-Cre*$^{Tg/Tg}$ (SMKO) and wild-type littermates (wild-type for *Efemp2* but containing the SM22α-Cre transgene, *Efemp2*$^{+/+}$;*SM22-Cre*$^{Tg/Tg}$) with a mixed genetic background (C57BL/6J; 129SvEv) were bred and maintained on a 12-hour light/dark cycle receiving standard rodent chow and water *ad libitum*. Genotyping was performed as previously described.[26] Wild-type (n=11) and SMKO (n=11) littermates were euthanized with CO_2 at 8 weeks of age and subsequently frozen. Animal experimentation and maintenance were in accordance with and approved by the Institutional Animal Experiment Committee of the University of Tsukuba, Japan (approval reference number: 21-108).

CRISPR/Cas9-Engineered Mouse Models

Three CRISPR/Cas9 knock-in mouse models (*Ltbp1*^{emG1132S}, *Mfap4*^{emD42A}, and *Timp1*^{emI120T}) were generated on the C57BL/6NJ background by Cyagen (Santa Clara, CA, USA), thereby introducing human-equivalent sequence variants detected in our WGS cohort of hAD patients (GRCm38/mm10 chr17(Ltbp1):g.75,315,062G>A, chr11(Mfap4):g.61,486,071A>C, and chrX(Timp1):g.20,873,395T>C, modeling human variants *LTBP1* NM_206943.4: c.3418G>A p.(Gly1140Ser), *MFAP4* NM_001198695.2: c.191A>C p.(Asp64Ala), and *TIMP1* NM_003254.3: c.356T>C p.(Ile119Thr), respectively). Founder mice of the three strains were bred and maintained on a 12-hour light/dark cycle receiving standard rodent chow and water *ad libitum*. The genotype was determined after weaning at the age of 4 weeks by PCR and subsequent Sanger sequencing using genomic DNA extracted of ear biopsies as previously described.[12] Primers used for PCR and sequencing are listed in Table S1. One-year-old heterozygous (*Ltbp1*$^{+/G1132S}$, n=10; *Mfap4*$^{+/D42A}$, n=10; *Timp1*$^{+/I120T}$, n=7), homozygous (*Ltbp1*$^{G1132S/G1132S}$, n=10; *Mfap4*$^{D42A/D42A}$, n=8; *Timp1*$^{I120T/I120T}$, n=5), and hemizygous (*Timp1*$^{0/I120T}$, n=9) mice as well as age-matched wild-type littermates serving as controls (*Ltbp1*$^{+/+}$, n=10; *Mfap4*$^{+/+}$, n=9; *Timp1*$^{+/+}$, n=8) were euthanized with CO_2 and subsequently

frozen (cf. one-year-old mice were used to be able to detect any late-onset effects of the

variants). Animal experimentation and maintenance were in accordance with institutional and

local guidelines (approval reference number: ZH107/17, Zurich, Switzerland) conforming to

the EU Directive 2010/63/EU for animal experiments.

Aortic Rupture Force Measurements

Euthanized mice kept in a frozen state were thawed immediately before measurement of

aortic rupture force. For all mouse models, the maximum tensile force at rupture (in mN) of

three uniaxially stretched 1.5-mm thoracic aortic rings/segments (segment S1, ascending

aorta, after heart; segment S2, descending aorta, after aortic arch; segment S3, descending

aorta, adjacent to S2) per mouse was measured and relative rupture force was calculated by

setting the arithmetic mean of the respective wild-type S1 to 100% as previously described.[12]

For the SMKO and both MFS models, the stretching positions of the tissue puller until both 5

mN (load-free aortic diameter) and maximum force (rupture of aortic segment) were

determined (in μm) as moved distance of the mounting pins during stretching. Mice were

randomized and all measurements were performed blinded for the investigator.

Statistical Analyses

For arithmetic means, 95% confidence intervals (CIs) were calculated using two-tailed critical

values of *t*-distribution (vassarstats.net/conf_mean.html). For means with non- or slightly

(<2% of means) overlapping 95% CIs compared in the text, two-tailed *P* values were

calculated by two-sample *t*-test (equal sample variances according to *F*-test) or Welch's *t*-test

(unequal sample variances according to *F*-test), assuming normal distribution of independent

samples (vassarstats.net/tu.html). For proportions, 95% CIs were calculated with continuity

correction (vassarstats.net/prop1.html). For additional information, see Supplemental Material.

RESULTS

$Fbn1^{C1041G}$ Mouse Model

Compared to $Fbn1^{+/+}$ mice, the aortic rupture force of $Fbn1^{+/C1041G}$ mice was significantly decreased in all three aortic segments (S1, S2, and S3) (Figure 1A; Table S2). No significant difference between $Fbn1^{+/+}$ and $Fbn1^{+/C1041G}$ mice was observed in the stretch at both 5 mN (aortic diameter) and maximum force (aortic rupture) in all three aortic segments (Figure 1B; Tables S3 and S4). Differences and trends in the relative aortic rupture force and stretch values of $Fbn1^{+/+}$ and $Fbn1^{+/C1041G}$ mice were comparable in both sexes (Figure S1).

$Fbn1^{mgR}$ Mouse Model

During losartan treatment, the mortality rate of treated $Fbn1^{mgR/mgR}$ mice (8.3% [95% CI, 0.4%-40.2%]) was not significantly lower than that of untreated $Fbn1^{mgR/mgR}$ mice (11.8% [95% CI, 2.1%-37.7%]), while no wild-type mouse died (Figure 2A). Although on average males were heavier than females, the body weight within sexes was not significantly different before treatment (at the age of 4 weeks), but after treatment (at the age of 8 weeks) $Fbn1^{mgR/mgR}$ males (especially losartan-treated) were noticeably heavier compared to wild-type males (Figure 2B).

Compared with untreated $Fbn1^{+/+}$ mice, the aortic rupture force was significantly decreased in all three aortic segments of both untreated and losartan-treated $Fbn1^{mgR/mgR}$ mice, i.e., losartan did not strengthen the weakened aortic wall of $Fbn1^{mgR/mgR}$ mice modeling MFS (Figure 3A; Table S5). Rather, the aortic rupture force of losartan-treated $Fbn1^{+/+}$ mice tended to be reduced compared to untreated $Fbn1^{+/+}$ mice in all three aortic segments measured, which requires further investigation (Figure 3A; Table S5).

The stretch of aortic segment S1 at both 5 mN (aortic diameter) and maximum force (rupture) was significantly increased in untreated female and male $Fbn1^{mgR/mgR}$ mice compared to untreated $Fbn1^{+/+}$ females and males, respectively, being the most pronounced in

males (Figure 3B; Tables S6 and S7). While losartan treatment of $Fbn1^{+/+}$ mice had no effect on the stretch of S1 neither in female nor in male animals (Figure 3B; Tables S6 and S7), losartan treatment considerably reduced the stretch of S1 at both 5 mN and maximum force in both female and male $Fbn1^{mgR/mgR}$ mice (Figure 3B; Tables S6 and S7). In S2 and S3, however, there were no considerable differences in the stretch at 5 mN and maximum force (data not shown).

Efemp2 SMKO Mouse Model

Compared with wild-type mice, our measurements showed significantly reduced relative aortic rupture force in the ascending thoracic aorta (S1) but no difference in the descending aorta (S2 and S3) of SMKO mice (Figure 4A; Table S8). Furthermore, in SMKO mice the stretch at both 5 mN and maximum force was significantly higher in the ascending aorta (S1) but significantly lower in the descending aorta (S3) at maximum force (Figure 4B; Tables S9 and S10).

Ltbp1, *Mfap4*, and *Timp1* CRISPR/Cas9 Knock-In Mouse Models

During the one-year period of aging, no *Ltbp1*, *Mfap4* or *Timp1* CRISPR/Cas9 knock-in mice died due to aortic rupture. Moreover, the rupture force of the aortic segments S1, S2, and S3 showed no considerable difference between wild-type controls and hetero-, homo-, or hemizygous mice of *Ltbp1*, *Mfap4*, and *Timp1* CRISPR/Cas9 knock-in mice (Figure 5; Table S11).

DISCUSSION

Inbred mouse hAD models are valuable tools for the investigation of medical therapeutic approaches and the establishment of gene-disease association and have been widely adopted for the identification of cardiovascular abnormalities by assessing aortic diameter, morphology, and wall thickness as well as by considering histological staining, omics data, or survival rate.[29-33] Although these methods provide observational insights, they cannot directly/objectively reveal how medical therapy or sequence variant(s) influences the strength (i.e., rupture force) of the aortic wall and whether or not there is a risk for aortic dissection and rupture. In this study, by measuring the aortic rupture force we provide hitherto unprecedented insights into the aortic biomechanical integrity of three established (*Fbn1^{C1041G}, Fbn1^{mgR}*, and *Efemp2* SMKO) and three novel candidate-gene (*Ltbp1*, *Mfap4*, and *Timp1*) mouse hAD models.

By applying the aortic rupture force measurement to both mouse MFS models (i.e., the phenotypically milder heterozygous *Fbn1^{C1041G}* and the phenotypically more severe homozygous *Fbn1^{mgR}* mice), we showed for the first time a significantly decreased rupture force as indicator for an impaired biomechanical integrity in the thoracic aorta, especially in the ascending aorta (Figures 1A and 3A). Even though the aortic wall of *Fbn1$^{+/C1041G}$* mice is normal in appearance until about 3 months of age and only later on shows medial thickening and/or progressive deterioration within the medial layer, including elastic fiber fragmentation in association with excess expression of matrix metalloproteinase 2 (MMP2) and 9 (MMP9), our read-out system allowed the identification of an impaired aortic wall integrity already at an early stage compared to previously described methods (cf. *Fbn1$^{+/C1041G}$* mice used in this study were 10-week-old).[17,18]

In *Fbn1$^{mgR/mgR}$* mice, treatment with the AGTR1-inhibitor losartan has been shown to improve survival only before aneurysm development but not after.[34,35] In our study, survival of *Fbn1$^{mgR/mgR}$* mice treated with losartan was not improved, despite treatment start already at

the age of 4 weeks (Figure 2A). However, as the average survival of $Fbn1^{mgR/mgR}$ mice is just

below 16 weeks of age,[18] our treatment may be too short to conclusively evaluate the effect of

losartan on average survival. Regarding weight, survived 8-week-old $Fbn1^{mgR/mgR}$ males

(especially those treated with losartan) were heavier than wild-type males (Figure 2B), which

is unexpected considering the human MFS phenotype but has been observed previously in

this mouse model and is attributed to altered adipose tissue homeostasis.[36]

Furthermore, compared to untreated wild-type controls, in $Fbn1^{mgR/mgR}$ mice there was

a significantly increased stretch until both 5 mN (i.e., diameter) and maximum force (i.e.,

rupture) in the ascending aorta (Figure 3B), reflecting aneurysm development and confirming

macroscopic evaluation. These differences were even more pronounced in male than in

female mice, which is in line with previously published data using ultrasound

measurements.[37-38] Indeed, sex is a well-known risk factor for cardiovascular diseases and

several clinical studies have shown that sex also strongly influences the outcome of MFS,

with men having a higher risk for aortic dilatations and arterial events.[39-41]

Treatment of $Fbn1^{mgR/mgR}$ mice with losartan recognizably reduced stretch and, thus,

aneurysm formation (Figure 3B). Despite preventing aneurysm formation by likely reducing

blood pressure (as previously shown for the same dosage of losartan in the $Fbn1^{C1041G}$ mouse

model),[21] losartan treatment did not impact the aortic rupture force and, hence, in case of

blood pressure spikes or continuously elevated blood pressure (e.g., upon discontinuation of

antihypertensive drug therapy), the vessel is still prone to rupture. This suggests that for the

investigation of potential medical therapeutic approaches in mouse hAD models presenting

with aneurysms, the sole prevention of aneurysm formation or the reduction of aneurysm

progression is not an adequate read-out for the strength of aortic wall. Thus, our read-out

system acts as a clinically relevant approach allowing novel insights into the evaluation of

medical therapies for the prevention of arterial events in hADs.

Moreover, the significantly reduced aortic rupture force demonstrated here indicates

that the absence of Fbln4 in the aortic ECM leads to an impaired aortic integrity in the

ascending aorta (Figure 4A). This is likely caused by the previously described profound

impact of the absence of Fbln4 on elastogenesis[42] and collagen biosynthesis,[26] also involved

in the development of ascending aortic aneurysms.[43,44] Accordingly, our measurements of

significantly increased aortic stretch reflected the aneurysm formation of the ascending aorta

of SMKO mice. Considering that the ascending aorta has the highest hemodynamic stress

burden in terms of pulse dampening (consistent with the highest elastin content),[45,46] it is

coherent that knock-out of *Efemp2* in vascular smooth muscle cells has the more pronounced

effect on the ascending aorta (i.e., S1 compared to S2 and S3).

In comparison to a previously reported mouse vEDS model (*Col3a1^{m1Lsmi}*),[12,16] the

relative aortic rupture forces of both *Fbn1^{+/C1041G}* and *Fbn1^{mgR/mgR}* mouse MFS models were

significantly higher in all three aortic segments (Figure 6; Table S12). The weaker thoracic

aorta of *Col3a1^{+/m1Lsmi}* mice is also reflected, for example, in the higher mortality rate

compared with *Fbn1^{mgR/mgR}* mice (Figure S2). In contrast, SMKO mice showed in the

ascending aorta (S1), where the aneurysms occurred, relative aortic rupture force values

comparable to or lower than those of mice modeling vEDS, while in the descending aorta (S2

and S3) with normal diameter the relative rupture force of SMKO mice was comparable to

that of mice modeling MFS (Figure 6; Table S12).

Because our measurements were able to reveal the reduced aortic rupture force of

established hAD mouse models as demonstrated here (*Fbn1^{C1041G}*, *Fbn1^{mgR}*, and *Efemp2*

SMKO) and elsewhere (*Col3a1^{m1Lsmi}*),[12,16] we applied our read-out approach to assess the

hitherto unknown aortic impact of CRISPR/Cas9-engineered human-equivalent mutations in

mouse models of three hAD candidate genes. In humans, these three genes (*LTBP1*, *MFAP4*,

and *TIMP1*) are expressed at higher levels in arterial tissue than in any other tissue studied in

GTEx (gtexportal.org/home). LTBP1 interacts with ECM proteins, such as MFS-associated

FBN1, and regulates transforming growth factor (TGF)-β,[47] which plays an important role in

the pathogenesis of hADs.[48] MFAP4, an ECM protein found in elastic fibers, actively

promotes tropoelastin self-assembly by binding tropoelastin and interacts with FBN1, FBN2,

and type I collagen.[49] TIMP1 is a strong inhibitor of many MMPs except for some of the

membrane type MMPs, thereby preventing excessive ECM degradation.[50] Knock-out mouse

models of *Ltbp1*, *Mfap4*, and *Timp1* showed cardiovascular phenotypes,[51-56] indicating that

mutations in these genes could be causative for hAD.

In contrast to established hAD mouse models (i.e., mice with mutation in the genes

Fbn1, *Efemp2*, or *Col3a1*), however, our read-out system revealed no significantly reduced

aortic rupture force for any genotypes of the three novel knock-in mouse models $Ltbp1^{G1132S}$,

$Mfap4^{D42A}$, and $Timp1^{I120T}$ at the age of 1 year (Figure 5). This indicates and clarifies the lack

of aortic significance of the sequence variants with unknown pathogenicity knocked into our

mouse models. The non-reduced aortic rupture force of the three knock-in mouse models are

consistent with their non-reduced survival rate (data not shown). It is important to note,

however, that survival rate does not reflect aortic rupture force at the individual level, as not

all mice with significantly reduced rupture force have a reduced survival.[16] Further studies are

needed to assess the clinical significance of the genes *LTBP1*, *MFAP4*, and *TIMP1* in hAD,

also considering a disease-modifying (synergistic) effect of these genes as well as mice with a

different genetic background.

Indeed, although mice can well model human diseases (*in vivo*) as demonstrated here

for hAD, mouse models cannot always accurately mimic human disease phenotypes.[57] For

example, the X-linked *BGN* gene encoding biglycan, which is already associated with hAD in

humans,[58] has been shown not to cause an aortic phenotype when *Bgn* is knocked out on the

C57BL/6 mouse background (likely due to the relative deficiency of TGF-β signaling induced

by the C57BL/6 background),[59] whereas biglycan deficiency in male BALB/cA mice leads to

sudden death due to aortic rupture.[60]

Taken together, this study demonstrates that our read-out system of aortic rupture

force can not only explain the aortic phenotype of established mouse hAD models but also

clarify the unknown/uncertain aortic significance of candidate genes, variants, and drugs,

thereby paving the way for better molecular diagnosis and medical therapies in hAD. Thus, it

is obvious that comprehensive studies related to aortic aneurysms, dissections, and ruptures

should consider and assess the biomechanical integrity (rupture force) of the aortic wall.

Acknowledgements

We thank Cyagen (Santa Clara, CA, USA) for help with the generation of the three

CRISPR/Cas9 knock-in mouse models, Bryana M. Levitan for performing ultrasound

measurements of $Fbn1^{mgR}$ mice as well as Daniela Brkan, Edina Jusufi, Marc T. Schönholzer,

Giancarlo Tomio, and Dzenneta Zukovic for help with data analysis and/or figure preparation.

Sources of Funding

This work was supported by the Isaac Dreyfus-Bernheim Stiftung and the Fondation pour la

Recherche et le Traitement Médical (FRTM) as well as by NIH NHLBI (grant number K01

HL149984) and JSPS KAKENHI (grant number JP20H03762).

Disclosures

None.

Supplemental Material

Statistical Analyses

Figures S1 and S2

Tables S1-S12

Major Resources Table

REFERENCES

1. Lindsay ME, Dietz HC. Lessons on the pathogenesis of aneurysm from heritable conditions. *Nature*. 2011;473:308-316. doi: 10.1038/nature10145

2. Verstraeten A, Luyckx I, Loeys B. Aetiology and management of hereditary aortopathy. *Nat Rev Cardiol*. 2017;14:197-208. doi: 10.1038/nrcardio.2016.211

3. Creamer TJ, Bramel EE, MacFarlane EG. Insights on the pathogenesis of aneurysm through the study of hereditary aortopathies. *Genes*. 2021;12:183. doi: 10.3390/genes12020183

4. Caspar SM, Dubacher N, Kopps AM, Meienberg J, Henggeler C, Matyas G. Clinical sequencing: From raw data to diagnosis with lifetime value. *Clin Genet*. 2018;93:508-519. doi: 10.1111/cge.13190

5. Renard M, Francis C, Ghosh R, Scott AF, Witmer PD, Adès LC, Andelfinger GU, Arnaud P, Boileau C, Callewaert BL, Guo D, Hanna N, Lindsay ME, Morisaki H, Morisaki T, Pachter N, Robert L, Van Laer L, Dietz HC, Loeys BL, Milewicz DM, De Backer J. Clinical Validity of Genes for Heritable Thoracic Aortic Aneurysm and Dissection. *J Am Coll Cardiol*. 2018;72:605-615. doi: 10.1016/j.jacc.2018.04.089

6. Caspar SM, Dubacher N, Matyas G. More Genes for Thoracic Aortic Aneurysms and Dissections. *J Am Coll Cardiol*. 2019;73:528-529. doi: 10.1016/j.jacc.2018.09.094

7. Humphrey JD, Milewicz DM, Tellides G, Schwartz MA. Dysfunctional mechanosensing in aneurysms. *Science*. 2014;344:477-479. doi: 10.1126/science.1253026

8. Humphrey JD, Schwartz MA, Tellides G, Milewicz DM. Role of mechanotransduction in vascular biology: focus on thoracic aortic aneurysms and dissections. *Circ Res*. 2015;116:1448-1461. doi: 10.1161/CIRCRESAHA.114.304936

9. Pierce DM, Maier F, Weisbecker H, Viertler C, Verbrugghe P, Famaey N, Fourneau I, Herijgers P, Holzapfel GA. Human thoracic and abdominal aortic aneurysmal tissues:

Damage experiments, statistical analysis and constitutive modeling. *J Mech Behav Biomed Mater*. 2015;41:92-107. doi: 10.1016/j.jmbbm.2014.10.003

10. Chiu P, Lee H-P, Dalal AR, Koyano T, Nguyen M, Connolly AJ, Chaudhuri O, Fischbein MP. Relative strain is a novel predictor of aneurysmal degeneration of the thoracic aorta: An ex vivo mechanical study. *JVS Vasc Sci*. 2021;2:235-246. doi: 10.1016/j.jvssci.2021.08.003

11. Chittajallu SNSH, Richhariya A, Tse KM, Chinthapenta V. A Review on Damage and Rupture Modelling for Soft Tissues. *Bioengineering (Basel)*. 2022;9:26. doi: 10.3390/bioengineering9010026

12. Dubacher N, Münger J, Gorosabel MC, Crabb J, Ksiazek AA, Caspar SM, Bakker ENTP, van Bavel E, Ziegler U, Carrel T, Steinmann B, Zeisberger S, Meienberg J, Matyas G. Celiprolol but not losartan improves the biomechanical integrity of the aorta in a mouse model of vascular Ehlers-Danlos syndrome. *Cardiovasc Res*. 2019;16:457-465. doi: 10.21037/acs.2017.11.10

13. Ong K-T, Perdu J, De Backer J, Bozec E, Collignon P, Emmerich J, Fauret A-L, Fiessinger J-N, Germain DP, Georgesco G, Hulot J-S, De Paepe A, Plauchu H, Jeunemaitre X, Laurent S, Boutouyrie P. Effect of celiprolol on prevention of cardiovascular events in vascular Ehlers-Danlos syndrome: a prospective randomised, open, blinded-endpoints trial. *Lancet* 2010;376:1476-1484. doi: 10.1016/S0140-6736(10)60960-9

14. Frank M, Adham S, Seigle S, Legrand A, Mirault T, Henneton P, Albuisson J, Denarié N, Mazzella J-M, Mousseaux E, Messas E, Boutouyrie P, Jeunemaitre X. Vascular Ehlers-Danlos Syndrome: Long-Term Observational study. *J Am Coll Cardiol*. 2019;73:1948-1957. doi: 10.1016/j.jacc.2019.01.058

15. Baderkhan H, Wanhainen A, Stenborg A, Stattin E-L, Björck M. Celiprolol Treatment in

Patients with Vascular Ehlers-Danlos Syndrome. *Eur J Vasc Endovasc Surg.*

2021;61:326-331. doi: 10.1016/j.ejvs.2020.10.020

16. Gorosabel MC, Dubacher N, Meienberg J, Matyas G. Vascular Ehlers-Danlos syndrome:

can the beneficial effect of celiprolol be extrapolated to bisoprolol? *EHJ - Cardiovasc*

Pharmacother. 2020;6:199-200. doi: 10.1093/ehjcvp/pvz067

17. Judge DP, Biery NJ, Keene DR, Geubtner J, Myers L, Huso DL, Sakai LY, Dietz HC.

Evidence for a critical contribution of haploinsufficiency in the complex pathogenesis of

Marfan syndrome. *J Clin Invest*. 2004;114:172-181. doi: 10.1172/JCI20641

18. Pereira L, Lee SY, Gayraud B, Andrikopoulos K, Shapiro SD, Bunton T, Biery NJ, Dietz

HC, Sakai LY, Ramirez F. Pathogenetic sequence for aneurysm revealed in mice

underexpressing fibrillin-1. *Procl Natl Acad Sci U S A*. 1999;96:3819-3823. doi:

10.1073/pnas.96.7.3819

19. Bunton TE, Biery NJ, Myers L, Gayraud B, Ramirez F, Dietz HC. Phenotypic Alteration

of Vascular Smooth Muscle Cells Precedes Elastolysis in a Mouse Model of Marfan

Syndrome. *Circ Res*. 2001;88:37-43. doi: 10.1161/01.res.88.1.37

20. Saeyeldin A, Zafar MA, Velasquez CA, Ip K, Gryaznov A, Brownstein AJ, Li Y, Rizzo

JA, Erben Y, Ziganshin BA, Elefteriades JA. Natural history of aortic root aneurysms in

Marfan   syndrome.   *Ann   Cardiothorac   Surg*.   2017;6:625-632.   doi:

10.21037/acs.2017.11.10

21. Habashi JP, Judge DP, Holm TM, Cohn RD, Loeys BL, Cooper TK, Myers L, Klein EC,

Liu G, Calvi C, Podowski M, Neptune ER, Halushka MK, Bedja D, Gabrielson K, Rifkin

DB, Carta L, Ramirez F, Huso DL, Dietz HC. Losartan, an AT1 Antagonist, Prevents

Aortic Aneurysm in a Mouse Model of Marfan Syndrome. *Science*. 2006;312:117-121.

doi: 10.1681/ASN.2006050508

22. Habashi JP, Doyle JJ, Holm TM, Aziz H, Schoenhoff F, Bedja D, Chen Y, Modiri AN,

Judge DP, Dietz HC. Angiotensin II Type 2 Receptor Signaling Attenuates Aortic

Aneurysm in Mice Through ERK Antagonism. *Science*. 2011;332:361-365. doi:

10.1126/science.1192152

23. Bhatt AB, Buck JS, Zuflacht JP, Milian J, Kadivar S, Gauvreau K, Singh MN, Creager

MA. Distinct effects of losartan and atenolol on vascular stiffness in Marfan syndrome.

Vasc Med. 2015;20:317-325. doi: 10.1177/1358863X15569868

24. Cook JR, Clayton NP, Carta L, Galatioto J, Chiu E, Smaldone S, Nelson CA, Cheng SH,

Wentworth BM, Ramirez F. Dimorphic effects of transforming growth factor-β signaling

during aortic aneurysm progression in mice suggest a combinatorial therapy for Marfan

syndrome.   *Arterioscler   Thromb   Vasc   Biol*.   2015;35:911-917.   doi:

10.1161/ATVBAHA.114.305150

25. Sellers SL, Milad N, Chan R, Mielnik M, Jermilova U, Huang PL, Crom R de, Hirota JA,

Hogg JC, Sandor GG, Van Breemen C, Esfandiarei M, Seidman MA, Bernatchez P.

Inhibition of Marfan Syndrome Aortic Root Dilation by Losartan: Role of Angiotensin II

Receptor Type 1-Independent Activation of Endothelial Function. *Am J Pathol*.

2018;188:574-585. doi: 10.1016/j.ajpath.2017.11.006

26. Huang J, Davis EC, Chapman SL, Budatha M, Marmorstein LY, Word RA, Yanagisawa

H. Fibulin-4 Deficiency Results in Ascending Aortic Aneurysms: A Potential Link

Between Abnormal Smooth Muscle Cell Phenotype and Aneurysm Progression. *Circ Res*.

2010;106:583-592. doi: 10.1161/CIRCRESAHA.109.207852

27. Hucthagowder V, Sausgruber N, Kim KH, Angle B, Marmorstein LY, Urban Z. Fibulin-

4: A Novel Gene for an Autosomal Recessive Cutis Laxa Syndrome. *Am J Hum Genet*.

2006;78:1075-1080. doi: 10.1086/504304

28. Renard M, Holm T, Veith R, Callewaert BL, Adès LC, Baspinar O, Pickart A, Dasouki

M, Hoyer J, Rauch A, Trapane P, Earing MG, Coucke PJ, Sakai LY, Dietz HC, De Paepe

AM, Loeys BL. Altered TGFβ signaling and cardiovascular manifestations in patients with autosomal recessive cutis laxa type I caused by fibulin-4 deficiency. *Eur J Hum Genet*. 2010;18:895-901. doi: 10.1038/ejhg.2010.45

29. Lee VS, Halabi CM, Hoffman EP, Carmichael N, Leshchiner I, Lian CG, Bierhals AJ, Vuzman D, Brigham Genomic Medicine, Mecham RP, Frank NY, Stitziel NO. Loss of function mutation in *LOX* causes thoracic aortic aneurysm and dissection in humans. *Proc Natl Acad Sci U S A*. 2016;113:8759-8764. doi: 10.1073/pnas.1601442113

30. Halabi CM, Broekelmann TJ, Lin M, Lee VS, Chu M-L, Mecham RP. Fibulin-4 is essential for maintaining arterial wall integrity in conduit but not muscular arteries. *Sci Adv*. 2017;3:e1602532. doi: 10.1126/sciadv.1602532

31. Parker SJ, Stotland A, MacFarlane E, Wilson N, Orosco A, Venkatraman V, Madrid K, Gottlieb R, Dietz HC, Van Eyk JE. Proteomics reveals Rictor as a noncanonical TGF-β signaling target during aneurysm progression in Marfan mice. *Am J Physiol Heart Circ Physiol*. 2018;315:H1112-H1126. doi: 10.1152/ajpheart.00089.2018

32. Bhushan R, Altinbas L, Jäger M, Zaradzki M, Lehmann D, Timmermann B, Clayton NP, Zhu Y, Kallenbach K, Kararigas G, Robinson PN. An integrative systems approach identifies novel candidates in Marfan syndrome-related pathophysiology. *J Cell Mol Med*. 2019;23:2526-2535. doi: 10.1111/jcmm.14137

33. Bowen CJ, Calderón Giadrosic JF, Burger Z, Rykiel G, Davis EC, Helmers MR, Benke K, Gallo MacFarlane E, Dietz HC. Targetable cellular signaling events mediate vascular pathology in vascular Ehlers-Danlos syndrome. *J Clin Invest*. 2020;130:686-698. doi: 10.1172/JCI130730

34. Xiong W, Meisinger T, Knispel R, Worth JM, Baxter BT. MMP-2 Regulates Erk1/2 Phosphorylation and Aortic Dilatation in Marfan Syndrome. *Circ Res*. 2012;110:e92-e101. doi: 10.1161/CIRCRESAHA.112.268268

35. Sheppard MB, Chen JZ, Rateri DL, Moorleghen JJ, Weiland M, Daugherty A. Renin-

Angiotensin System Inhibitors Do Not Improve Survival in Fibrillin-1 Hypomorphic

Mice with Established Aortic Aneurysm. *J Clin Trans Sci*. 2019;3:112-113. doi:

10.1017/cts.2019.258

36. Muthu ML, Tiedemann K, Fradette J, Komarova S, Reinhardt DP. Fibrillin-1 regulates

white   adipose   tissue   development,   homeostasis,   and   function. *Matrix Biol*.

2022;110:106-128. doi: 10.1016/j.matbio.2022.05.002

37. Chen JZ, Rateri D, Sheppard M, Daugherty A. Abstract 20315: Sexual Dimorphism in a

Marfan Syndrome Mouse Model. *Circulation*. 2017;136:A20315.

38. Chen JZ, Sawada H, Moorleghen JJ, Weiland M, Daugherty A, Sheppard MB. Aortic

Strain Correlates With Elastin Fragmentation in Fibrillin-1 Hypomorphic Mice. *Circ Rep*.

2019;1:199-205. doi: 10.1253/circrep.CR-18-0012

39. Détaint D, Faivre L, Collod-Beroud G, Child AH, Loeys BL, Binquet C, Gautier E,

Arbustini E, Mayer K, Arslan-Kirchner M, Stheneur C, Halliday D, Beroud C, Bonithon-

Kopp C, Claustres M, Plauchu H, Robinson PN, Kiotsekoglou A, De Backer J, Adès L,

Francke U, De Paepe A, Boileau C, Jondeau G. Cardiovascular manifestations in men and

women   carrying   a   FBN1   mutation. *Eur Heart J*.   2010;31:2223-2229.   doi:

10.1093/eurheartj/ehq258

40. Holmes KW, Maslen CL, Kindem M, Kroner BL, Song HK, Ravekes W, Dietz HC,

Weinsaft JW, Roman MJ, Devereux RB, Pyeritz RE, Bavaria J, Milewski K, Milewicz D,

LeMaire SA, Hendershot T, Eagle KA, Tolunay HE, Desvigne-Nickens P, Silberbach M,

for the GenTAC Registry Consortium. GenTAC registry report: Gender differences

among individuals with genetically triggered thoracic aortic aneurysm and dissection. *Am*

J Med Genet. 2013;161:779-786. doi: 10.1002/ajmg.a.35836

41. Renard M, Muiño-Mosquera L, Manalo EC, Tufa S, Carlson EJ, Keene DR, De Backer J,

Sakai LY. Sex, pregnancy and aortic disease in Marfan syndrome. *PLoS ONE*.

2017;12:e0181166. doi: 10.1371/journal.pone.0181166

42. Horiguchi M, Inoue T, Ohbayashi T, Hirai M, Noda K, Marmorstein LY, Yabe D, Takagi

K, Akama TO, Kita T, Kimura T, Nakamura T. Fibulin-4 conducts proper elastogenesis

via interaction with cross-linking enzyme lysyl oxidase. *Proc Natl Acad Sci U S A*.

2009;106:19029-19034. doi: 10.1073/pnas.0908268106

43. Papke CL, Yanagisawa H. Fibulin-4 and fibulin-5 in elastogenesis and beyond: Insights

from mouse and human studies. *Matrix Biol*. 2014;37:142-149. doi:

10.1016/j.matbio.2014.02.004

44. Papke CL, Tsunezumi J, Ringuette L-J, Nagaoka H, Terajima M, Yamashiro Y, Urquhart

G, Yamauchi M, Davis EC, Yanagisawa H. Loss of fibulin-4 disrupts collagen synthesis

and maturation: implications for pathology resulting from *EFEMP2* mutations. *Hum Mol*

Genet. 2015;24:5867-5879. doi: 10.1093/hmg/ddv308

45. Ruddy JM, Jones JA, Spinale FG, Ikonomidis JS. Regional heterogeneity within the

aorta: Relevance to aneurysm disease. *J Thorac Cardiovasc Surg*. 2008;136:1123-1130.

doi: 10.1016/j.jtcvs.2008.06.027

46. Jana S, Hu M, Shen M, Kassiri Z. Extracellular matrix, regional heterogeneity of the

aorta, and aortic aneurysm. *Exp Mol Med*. 2019;51:1-15. doi: 10.1038/s12276-019-0286-3

47. Morisaki T, Morisaki H. Genetics of hereditary large vessel diseases. *J Hum Genet*.

2016;61:21-26. doi: 10.1038/jhg.2015.119

48. Neptune ER, Frischmeyer PA, Arking DE, Myers L, Bunton TE, Gayraud B, Ramirez F,

Sakai LY, Dietz HC. Dysregulation of TGF-β activation contributes to pathogenesis in

Marfan syndrome. *Nat Genet*. 2003;33:407-411. doi: 10.1038/ng1116

49. Pilecki B, Holm AT, Schlosser A, Moeller JB, Wohl AP, Zuk AV, Heumüller SE, Wallis

R, Moestrup SK, Sengle G, Holmskov U, Sorensen GL. Characterization of

Microfibrillar-associated Protein 4 (MFAP4) as a Tropoelastin- and Fibrillin-binding

Protein Involved in Elastic Fiber Formation. *J Biol Chem*. 2016;291:1103-1114. doi:

10.1074/jbc.M115.681775

50. Brew K, Nagase H. The tissue inhibitors of metalloproteinases (TIMPs): An ancient

family with structural and functional diversity. *Biochim Biophys Acta*. 2010;1803:55-71.

doi: 10.1016/j.bbamcr.2010.01.003

51. Roten L, Nemoto S, Simsic J, Coker ML, Rao V, Baicu S, Defreyte G, Soloway PJ, Zile

MR, Spinale FG. Effects of Gene Deletion of the Tissue Inhibitor of the Matrix

Metalloproteinase-type 1 (TIMP-1) on Left Ventricular Geometry and Function in Mice.

J Mol Cell Cardiol. 2000;32:109-120. doi: 10.1006/jmcc.1999.1052

52. Eskandari MK, Vijungco JD, Flores A, Borensztajn J, Shively V, Pearce WH. Enhanced

abdominal aortic aneurysm in TIMP-1-deficient mice. *J Surg Res*. 2005;123:289-293. doi:

10.1016/j.jss.2004.07.247

53. Todorovic V, Frendewey D, Gutstein DE, Chen Y, Freyer L, Finnegan E, Liu F, Murphy

A, Valenzuela D, Yancopoulos G, Rifkin DB. Long form of latent TGF-β binding protein

1 (Ltbp1L) is essential for cardiac outflow tract septation and remodeling. *Development*.

2007;134:3723-3732. doi: 10.1242/dev.008599

54. Todorovic V, Finnegan E, Freyer L, Zilberberg L, Ota M, Rifkin DB. Long form of latent

TGF-β binding protein 1 (Ltbp1L) regulates cardiac valve development. *Dev Dyn*.

2011;240:176-187. doi: 10.1002/dvdy.22521

55. Schlosser A, Pilecki B, Hemstra LE, Kejling K, Kristmannsdottir GB, Wulf-Johansson H,

Moeller JB, Füchtbauer E-M, Nielsen O, Kirketerp-Møller K, Dubey LK, Hansen PBL,

Stubbe J, Wrede C, Hegermann J, Ochs M, Rathkolb B, Schrewe A, Bekeredjian R, Wolf

E, Gailus-Durner V, Fuchs H, Hrabě de Angelis M, Lindholt JS, Holmskov U, Sorensen

GL. MFAP4 Promotes Vascular Smooth Muscle Migration, Proliferation and Accelerates

Neointima Formation. *Arterioscler Thromb Vasc Biol*. 2016;36:122-133. doi:

10.1161/ATVBAHA.115.306672

56. Arpino V, Brock M, Gill SE. The role of TIMPs in regulation of extracellular matrix

proteolysis. *Matrix Biol*. 2015;44-46:247-254. doi: 10.1016/j.matbio.2015.03.005

57. Ritskes-Hoitinga M, Leenaars C, Beumer W, Coenen-de Roo T, Stafleu F, Meijboom

FLB. Improving Translation by Identifying Evidence for More Human-Relevant

Preclinical Strategies. *Animals*. 2020;10:1170. doi: 10.3390/ani10071170

58. Meester JAN, Vandeweyer G, Pintelon I, Lammens M, Van Hoorick L, De Belder S,

Waitzman K, Young L, Markham LW, Vogt J, Richer J, Beauchesne LM, Unger S,

Superti-Furga A, Prsa M, Dhillon R, Reyniers E, Dietz HC, Wuyts W, Mortier G,

Verstraeten A, Van Laer L, Loeys BL. Loss-of-function mutations in the X-linked

biglycan gene cause a severe syndromic form of thoracic aortic aneurysms and

dissections. *Genet Med*. 2017;19:386-395. doi: 10.1038/gim.2016.126

59. Tang T, Thompson JC, Wilson PG, Nelson C, Williams KJ, Tannock LR. Decreased

Body Fat, Elevated Plasma Transforming Growth Factor-β Levels, and Impaired BMP4-

Like Signaling in Biglycan-Deficient Mice. *Connect Tissue Res*. 2013;54:5-13. doi:

10.3109/03008207.2012.715700

60. Heegaard A-M, Corsi A, Danielsen CC, Nielsen KL, Jorgensen HL, Riminucci M, Young

MF, Bianco P. Biglycan Deficiency Causes Spontaneous Aortic Dissection and Rupture in

Mice. *Circulation*. 2007;115:2731-2738. doi: 10.1161/CIRCULATIONAHA.106.653980

 Highlights

- Assay to measure the aortic rupture force as read-out for the biomechanical integrity identified weakened murine thoracic aorta prior to micro- and macroscopic changes in $Fbn1^{+/C1041G}$ mice.

- Despite reducing aneurysm growth, losartan treatment did not have any impact on the aortic rupture force of $Fbn1^{mgR/mgR}$ mice.

- Smooth-muscle-cell-knock-out of $Efemp2$ significantly impaired the rupture force of the murine thoracic aorta.

- Aortic rupture force measurements could clarify the causality of novel candidate gene(s)/variant(s) in mouse models of aortic diseases.

Figures and Figure Legends

Figure 1

Figure 1. Aortic parameters of the *Fbn1^{C1041G}* mouse model.
A, Relative aortic rupture force (%) of three aortic segments (S1-S3) of 10-week-old *Fbn1^{+/+}* and *Fbn1^{+/C1041G}* mice. **B**, Stretch (in μm) of aortic segments at 5mN (load-free aortic diameter) and maximum force of 10-week-old *Fbn1^{+/+}* and *Fbn1^{+/C1041G}* mice. Bars represent arithmetic means and error bars indicate 95% confidence intervals. The sample size (n) is displayed (damaged segments and outliers, i.e. >2× standard deviation, were excluded). For mean values with non- or slightly overlapping 95% CIs compared in text, two-tailed *P* values of unpaired t-tests are shown (applies to **A** only).

Figure 2

Figure 2. Survival and bodyweight of the *Fbn1^{mgR}* mouse model treated with the AGTR1-inhibitor losartan.
A, Kaplan-Meier curve for the survival of untreated *Fbn1^{+/+}* mice (n=10) and *Fbn1^{mgR/mgR}* mice (n=17), and *Fbn1^{+/+}* (n=10) and *Fbn1^{mgR/mgR}* (n=12) mice receiving losartan between the age of 4 and 8 weeks. Note that the survival curve of losartan-treated *Fbn1^{+/+}* mice is masked by that of untreated *Fbn1^{+/+}* mice. 95% confidence intervals (shaded areas) are shown for *Fbn1^{mgR/mgR}* mice with survivals differing from 100%. **B**, Bodyweights of mice surviving the treatment period of 4 weeks measured before and after treatment at the age of 4 and 8 weeks, respectively. Bars represent arithmetic means, error bars indicate 95% confidence intervals. Significant two-tailed *P* values of unpaired t-tests are shown (*P*<0.05). For the comparison between female and male mice (B), the respective groups were pooled prior to *P* value calculation.

Figure 3

Figure 3. Aortic parameters of the *Fbn1mgR* mouse model treated with the AGTR1-inhibitor losartan.
A, Relative aortic rupture force (%) of three aortic segments (S1-S3) of 8-week-old *Fbn1+/+* and *Fbn1mgR/mgR* mice after 4-week-long treatment with losartan. Untreated *Fbn1+/+* and *Fbn1mgR/mgR* mice served as control. **B**, Stretch (in μm) of the aortic segment S1 at 5 mN (load-free aortic diameter) as well as at maximum force in losartan-treated and untreated *Fbn1+/+* and *Fbn1mgR/mgR* mice separated according to sex. Bars represent arithmetic means and error bars indicate 95% confidence intervals (95% CI). The sample size (n) is displayed (damaged segments and outliers, i.e. >2× standard deviation, were excluded). For mean values with non- or slightly overlapping 95% CIs compared in text, two-tailed *P* values of unpaired t-tests are shown.

Figure 4

Figure 4. Aortic parameters of the *Efemp2* SMKO mouse model.
A, Relative aortic rupture force (%) of three aortic segments (S1-S3) of 8-week-old wild-type and SMKO mice. **B**, Stretch (in μm) of aortic segments at 5mN (load-free aortic diameter) as well as at maximum force in wild-type and SMKO mice. Bars represent arithmetic means and error bars indicate 95% confidence intervals. The sample size (n) is displayed (damaged segments and outliers, i.e. >2× standard deviation, were excluded). For mean values with non- or slightly overlapping 95% CIs compared in text, two-tailed *P* values of unpaired t-tests are shown.

Figure 5

Figure 5. Aortic rupture force of the *Ltbp1*, *Mfap4*, and *Timp1* CRISPR/Cas9 knock-in mouse models.
A-C, Relative aortic rupture force (%) of three aortic segments (S1-S3) of 1-year-old wild-type and hetero- as well as hemi- and homozygous *Ltbp1* (**A**), *Mfap4* (**B**), and *Timp1* (**C**) CRISPR/Cas9 knock-in mice. Bars represent arithmetic means, error bars indicate 95% confidence intervals. The sample size (n) is displayed and damaged segments and outliers (>2× standard deviation) were excluded. Note the absence of non- or slightly overlapping 95% CIs per aortic segment. Circle, female; triangle, male.

Figure 6

Figure 6. Aortic rupture force of the *Fbn1^C1041G^*, *Fbn1^mgR^*, and *Efemp2* SMKO mouse models compared to *Col3a1^m1Lsmi^* mice modeling vEDS.

Relative aortic rupture force (%) of three aortic segments (S1-S3) of the mouse models *Col3a1^m1Lsmi^* (data as described previously),[12] *Fbn1^C1041G^*, *Fbn1^mgR^*, and *Efemp2* SMKO (cf. wild-type controls of the *Fbn1^C1041G^*, *Fbn1^mgR^*, and *Efemp2* SMKO mouse models were pooled). Bars represent arithmetic means, error bars indicate 95% confidence intervals, The sample size (n) is displayed (damaged segments and outliers, i.e. >2× standard deviation, were excluded). For mean values with non- or slightly overlapping 95% CIs compared in text, two-tailed *P* values of unpaired t-tests are shown. *Col3a1^+/+^* and *Col3a1^+/-^* denotes wild-type and heterozygous mice of the *Col3a1^m1Lsmi^* mouse model, respectively.