Evolution of *E. coli* in a mouse model of inflammatory bowel disease leads to a disease-specific bacterial genotype and trade-offs with clinical relevance

Rahul Unni[1,2,#], Nadia Andrea Andreani[1,2,#], Marie Vallier[1,2,§,#], Silke S. Heinzmann[3], Jan Taubenheim[4], Martina A. Guggeis[5,6], Florian Tran[5,6], Olga Vogler[1], Sven Künzel[1], Jan-Bernd Hövener[7], Philip Rosenstiel[5], Christoph Kaleta[4], Astrid Dempfle[8], Daniel Unterweger[1,2,*], John F. Baines[1,2,*]

[1] *Max Planck Institute for Evolutionary Biology, Plön, Germany.*

[2] *Section of Evolutionary Medicine, Institute for Experimental Medicine, Kiel University, Kiel, Germany.*

[3] *Research Unit Analytical BioGeoChemistry, Helmholtz Munich, Neuherberg, Germany.*

[4] *Research Group Medical Systems Biology, Institute for Experimental Medicine, Kiel University, Kiel, Germany.*

[5] *Institute of Clinical Molecular Biology, Christian-Albrechts-University Kiel and University Medical Center Schleswig-Holstein, Kiel, Germany.*

[6] *Department of Internal Medicine I, University Medical Center Schleswig-Holstein, Kiel, Germany.*

[7] *Section Biomedical Imaging, Molecular Imaging North Competence Center (MOIN CC), Department of Radiology and Neuroradiology, University Medical Center Kiel, Kiel University, Kiel, Germany.*

[8] *Institute of Medical Informatics and Statistics, Kiel University, Kiel, Germany.*

[#] These authors contributed equally.

§ Current affiliation: Valbiotis, R&D Center, Riom, France.

* Corresponding authors: baines@evolbio.mpg.de, unterweger@iem.uni-kiel.de

ORCID:

Rahul Unni: 0000-0002-4656-7542

Nadia Andrea Andreani: 0000-0003-0330-9874

Marie Vallier: 0000-0002-7188-7070

Silke Heinzmann: 0000-0003-0257-8837

Jan Taubenheim: 0000-0001-7283-1768

Martina A. Guggeis: 0000-0003-4505-3731

Florian Tran: 0000-0002-3735-9872

Olga Vogler: 0000-0001-8930-7671

Sven Künzel:

Jan-Bernd Hövener: 0000-0001-7255-7252

Philip Rosenstiel: 0000-0002-9692-8828

Christoph Kaleta: 0000-0001-8004-9514

Astrid Dempfle: 0000-0002-2618-3920

Daniel Unterweger: 0000-0001-7459-6386

John F. Baines: 0000-0002-8132-4909

Evolution of *E. coli* in a mouse model of inflammatory bowel disease leads

to a disease-specific bacterial genotype and trade-offs with clinical

relevance

Abstract

Inflammatory bowel disease (IBD) is a persistent inflammatory condition that affects

the gastrointestinal tract and presents significant challenges in its management and

treatment. Despite the knowledge that within-host bacterial evolution occurs in the

intestine, the disease has rarely been studied from an evolutionary perspective. In this

study, we aimed to investigate the evolution of resident bacteria during intestinal

inflammation and whether- and how disease-related bacterial genetic changes may

present trade-offs with potential therapeutic importance. Here, we perform an *in vivo*

evolution experiment of *E. coli* in a gnotobiotic mouse model of IBD, followed by

multiomic analyses to identify disease-specific genetic and phenotypic changes in

bacteria that evolved in an inflamed versus a non-inflamed control environment. Our

results demonstrate distinct evolutionary changes in *E. coli* specific to inflammation,

including a single nucleotide variant that independently reached high frequency in all

inflamed mice. Using *ex vivo* fitness assays, we find that these changes are associated

with a higher fitness in an inflamed environment compared to isolates derived from

non-inflamed mice. Further, using large-scale phenotypic assays, we show that

bacterial adaptation to inflammation results in clinically relevant phenotypes, which

intriguingly include collateral sensitivity to antibiotics. Bacterial evolution in an

inflamed gut yields specific genetic and phenotypic signatures. These results may serve

as a basis for developing novel evolution-informed treatment approaches for patients

with intestinal inflammation.

Keywords: Inflammatory bowel disease, *E. coli*, experimental evolution, evolutionary trade-

offs.

Introduction

Dysbiosis of the intestinal microbial community, characterized by alterations in bacterial composition and function, is a hallmark of inflammatory bowel disease (IBD). The nature of these microbial imbalances has been described extensively and at numerous complementary levels, including diversity parameters, taxonomic- and functional genomic changes, and at the level of gene expression [1–6]. This important body of work accounts for changes observed at the ecological level, while the latter also considers the potential contribution of phenotypic plasticity of gut microbes to disease susceptibility. Moreover, numerous studies have highlighted the association of dysbiotic microbial signatures with distinct clinical IBD subtypes, including Crohn's disease (CD) and ulcerative colitis (UC), further emphasizing the intricate relationship between the gut microbiota and disease pathogenesis [3,4,7].

While research has focused extensively on the altered ecology of the gut microbiota in IBD, its potential for evolutionary change during disease pathogenesis has received comparatively limited attention. However, numerous recent studies have revealed the capacity of bacteria to undergo adaptive evolution within the host environment, including the gut [reviewed in [8]]. This phenomenon is evident in both mouse models [9–11] and human subjects [12], even in the absence of overt disease. Importantly, recent studies indicate that evolutionary changes also occur in the context of host inflammation. Elhenawy et al. focused on CD-related adherent-invasive *E. coli* (AIEC) using a murine model of chronic colonization, and revealed the evolution of lineages displaying enhanced invasive and metabolic capabilities [13]. Notably, they found that the fitness benefits conferred by increased motility were specific to the host environment, suggesting an evolutionary trade-off. In a second study exploring bacterial evolution in aging mice, significant differences in *E. coli* evolution between old and young mice were revealed [14]. The aged mouse environment

exhibited increased inflammation, leading to the specific targeting of stress-related functions

in *E. coli*.

These findings emphasize the importance of including evolutionary perspectives

when studying dysbiosis of the gut microbiome. In particular, documenting bacterial

evolution within the inflamed gut has the potential to reveal the *timing* of disease-specific

signatures, i.e., it may help disentangle the classical "chicken or egg" dilemma in microbiome

research [15], as well as to shed light on potential trade-offs resulting from adaptive changes.

Evolutionary trade-offs occur when an increase in fitness in one environment is accompanied

by a decrease in fitness in another [16,17]. These trade-offs can have significant therapeutic

implications and are observed in various related fields, including antibiotic resistance [18–21]

and cancer treatment [22–25]. Thus, elucidating trade-offs associated with bacterial adaptations

in the inflamed gut may offer insight into potential collateral effects on bacterial fitness and

prove useful for developing novel treatment strategies based on evolutionary principles.

In this study, we investigated the evolutionary dynamics of resident gut bacteria

during intestinal inflammation using an established gnotobiotic mouse model of IBD [26].

Through the monocolonization of both *Interleukin 10*-knockout and wildtype mice with a

single *E. coli* strain (NC101), this setup allowed us to track genetic and phenotypic changes

over the course of intestinal inflammation and to evaluate their potential clinical relevance.

By employing multiomic analysis and high-throughput phenotypic screening, we identify

genetic changes in bacteria, alterations in the metabolome, and differences in numerous

phenotypic traits among bacterial populations that are specifically associated with the

evolution in inflamed *Il10*-knockout mice. Remarkably, among the phenotypic changes

observed in inflammation-adapted bacteria are sensitivities to antibiotics with a known

therapeutic value in IBD. These results further confirm the importance of understanding

bacterial adaptation to inflammation and suggest its more widespread study of patients as a

means to develop novel treatment approaches.

Results

Gnotobiotic model of intestinal inflammation

In order to capture the evolutionary dynamics of bacteria evolving in the context of

intestinal inflammation, we implemented a previously established gnotobiotic model, for

which inflammation develops upon colonization with the *E. coli* NC101 strain in IL10-

deficient (*Il10*$^{-/-}$; herein "KO"), but not wild type (WT) mice [26]. We performed two

independent *in vivo* experiments (see Methods), in which WT (total N=14) and KO (total

N=11) mice were monocolonized and monitored over a period of 12 weeks, with longitudinal

sampling of feces for downstream multiomic analyses (Fig 1A; Table S1, see Methods).

Inflammation was monitored via levels of lipocalin-2 in feces, which steadily and

significantly increased in KO mice, but not in WT mice, already at one week post-inoculation

(Fig 1B, Tables S2 and S3, Wilcoxon signed-rank test Benjamini-Hochberg-corrected $P <$

0.05). Histopathological assessment of colonic tissue at the endpoint reveals significantly

higher pathology scores in KO mice than in WT mice (Fig 1C, Kruskal-Wallis H test

Benjamini-Hochberg-corrected $P < 0.05$), further confirming that inflammation was specific

to KO mice. The bacterial load in the fecal samples was stable throughout the experiment, as

measured by CFU counts normalized by wet feces weight (Fig S1, Tables S4 and S5).

Bacterial populations show genetic diversification in healthy and inflamed mice

To assess the genetic diversification of the bacterial populations during the evolution

experiment, we performed shotgun sequencing on each single inoculum and all bacterial

populations at weeks 1, 4, 8, and 12 and identified *de-novo* mutations (i.e., mutations that

were not present in an inoculum; see Methods) by comparison with the genome of the

ancestral strain. No statistically significant difference was observed in the number of

mutations in the bacteria from KO and WT mice at any point in the experiment, suggesting

that differences in the intestinal environment in which the populations evolved did not exert a

measurable effect on the number of mutations (Fig S2, Table S6). One week after gavage, the

bacterial populations showed a median number of 87 *de-novo* mutations, with a subsequent

reduction at the following timepoints, although the differences are not significant (median at

week 4=34, at week 8=45 and at week 12=36, Fig S2, Tab S7 Wilcoxon signed rank test,

corrected $P > 0.05$). Comparison of the evolved populations at week 12 with the ancestral

strain reveals a median number of 39 mutations per mouse in WT and 26 in KO mice (Fig

1D). This indicates that the evolved bacterial populations in both groups of mice genetically

differ from the ancestral strain. Although the unphased shotgun data do not enable us to

directly analyze the role of genetic hitchhiking, Table S8 indicates the number of mutations at

week 12 already present at each earlier time point. Interestingly, there remains a substantial

(on average approx. 60% in each mouse genotype) proportion of *de-novo* mutations that arose

between week 8 and 12, which suggests that a large proportion of the mutations cannot be

explained by hitchhiking from standing genetic variation at earlier time points.

To test for overall differences in the *de-novo* mutations that accumulated in the

bacterial populations that evolved in the WT and KO mice, we performed a partial least-

squares discriminant analysis (PLS-DA) on all mutated loci (genes or intergenic regions

differing in their nucleotide sequence compared to the inocula). We find a clear and

significant distinction between the populations that evolved in the two mouse genotypes (Fig

1E, PERMANOVA $P = 0.0022$). The same result was observed when comparing single *de-*

novo mutated genomic positions between evolved bacterial populations from WT and KO

mice (Fig S3, PERMANOVA $P = 0.0016$).

Single nucleotide polymorphism differentiates evolved bacteria of all inflamed mice from

evolved bacteria of healthy mice

To identify candidates for genetic changes that may be specifically selected in an

inflamed environment, we focused on parallel mutations (i.e., genes or intergenic regions

mutated in at least two mice that were gavaged with an independent inoculum; see Methods)

and their associated functions. Analysis of the KEGG pathways encompassed by these loci

reveals seven pathways unique to the bacterial populations from inflamed mice (Fig 2A).

These pathways include those involved in amino acid metabolism, pyruvate metabolismall of

which have been associated with IBD in clinical settings [27,28]. In addition to the pathways

affected only in bacterial populations from the inflamed mice, 18 pathways are affected only

in populations from healthy mice, and 18 other pathways are mutated in populations from

both healthy and inflamed mice (Fig 2A).

Univariate analysis reveals that only one particular locus is mutated at significantly

higher frequencies in evolved bacterial populations from inflamed than healthy mice

(Wilcoxon signed-rank test, corrected $P = 0.001$, Table S9), which is a single nucleotide

polymorphism (SNP) (C>T) nine nucleotides upstream of the gene *mprA* (position 3009211)

(Fig 2B). It is found in 11/11 KO mice at the endpoint with a frequency between 0.373 and 1,

compared to 2/14 WT mice with a frequency between 0.085 and 0.239. Two other mutations

are detected in the same intergenic region, although in a few populations only. The bacteria

from one WT mouse harbors a C>A mutation at the same position (3009211) with a

frequency of 0.052, whereas those from one inflamed mouse harbors a G>T SNP at a nearby

position (3009194) with a frequency of 0.611.

To further investigate the most frequent C>T mutation, we analyzed its frequency

over the course of the experiment (weeks 1, 4, 8, and 12). At week 1 the mutation is not

present or below the detection threshold (minimum frequency of 0.05; see Methods), while at

week 4, it is observed in 7/11 KO mice with a maximum frequency of 0.616, in comparison

to 1/14 WT mice with a frequency of 0.364 (Fig 2C). After a steady increase in frequency at

weeks 8 and 12, the mutation reaches fixation or near fixation in 7/11 KO mice and is also

present at intermediate frequencies in the remaining 4/11 KO mice (Fig 2C). Importantly, the

increase in frequency of the C>T mutation strongly coincides with the increase in

inflammation as measured by lipocalin-2 levels, both in KO mice alone and when including

all mice (Fig 2C, Fig S4; Spearman's rank correlation rho $P = 0.00005$ for KO mice alone and

$P = 3.005\text{e-}11$ for all mice).

Interestingly, closer inspection of the intergenic region between the hypothetical

protein_02879 and the gene *mprA* reveals that this genomic region is a mutational hotspot,

with six others positions in this region having mutated over the course of the *in vivo*

experiment (Fig S5).

Taken together, we identify a C>T change at position 3009211 in the *E. coli* genome,

which is a parallel mutation significantly associated with intestinal inflammation. This

mutation is located upstream of *mprA* (also known as *emrR*), a transcriptional repressor of

several genes, including those encoding the efflux pump EmrAB and AcrAB, and a putative

outer membrane porin OmpC (also known as NmpC)[29–31].

Bacterial populations show significantly different expression profiles in healthy and

inflamed mice

To quantify the expression of genes that are part of the *mprA* regulon (*mprA*, *emrAB*,

acrAB, and *ompC*)[29–31], as well as gain an overall view on differences in gene expression in

WT- compared to KO-evolved bacterial populations, we perform metatranscriptomic

sequencing of all week 12 bacterial populations. To test for overall differences in the gene

expression, we perform a partial least-squares discriminant analysis (PLS-DA) on all genes,

using RPKM (Reads per Kilobase per million Mapped Reads) as a proxy for gene expression.

We find a clear and significant difference in overall gene expression between populations

according to mouse genotype (Fig S6A, PERMANOVA P < 0.00005). Differential

expression analysis performed with Deseq2 reveals 2,219 genes that are up- or down-

regulated in one of the two conditions (1,022 upregulated for the populations from the

healthy gut and 1,197 for the population from the inflamed gut; Benjamini-Hochberg

corrected P <0.05, Tab S10). Analysis of the KEGG pathways encompassed by these

upregulated genes reveals 25 pathways unique to the bacterial populations from inflamed

mice (Fig S6B). These pathways include Ribosome, Aminoacyl-tRNA biosynthesis,

Glutathione metabolism, RNA degradation, Folate biosynthesis, Sulfur relay system,

Arginine biosynthesis, Protein export, Pantothenate and CoA biosynthesis, and Homologous

recombination, among others (Tab S11). Interestingly, *mprA* and *ompC,* which can be

putatively regulated by *mprA*[30], are significantly more expressed in the populations that

evolved in the inflamed gut (Tab S10; Fig S6C).

Increase in fitness of evolved bacteria in an inflammatory environment

As parallel mutations are a strong indication of adaptation by natural selection, we

hypothesized that the evolved bacterial populations may have a fitness advantage in the

environment in which they evolved, and thus next focused on the phenotypes present in the

evolved bacterial populations. To test whether evolution in the inflamed mouse gut confers a

fitness advantage to *E. coli*, we performed "reciprocal transplant" experiments based on *ex*

vivo assays using filtered cecal content from inflamed versus non-inflamed mice, collected

after sacrifice (referred to subsequently as "*ex vivo* media"). Populations of *E. coli* from fecal

samples collected at week 12 from a subset of mice (N = 5 KO mice and N = 5 WT mice)

were cultured in *ex vivo* media derived from both inflamed and non-inflamed mice. While no

significant difference is observed between bacteria from WT and KO mice when grown in

non-inflamed *ex vivo* media (Kruskal-Wallis test $P = 0.9168$; Fig 3A), bacteria from KO mice

grow significantly better than those from WT mice in the inflamed *ex vivo* media (Kruskal-

Wallis test $P = 0.009023$). Notably, bacteria from KO mice grow significantly better in the

inflamed *ex vivo* media than in the non-inflamed *ex vivo* media (Kruskal-Wallis test $P =$

0.0472), while bacteria from WT mice show no difference in growth in the two media

(Kruskal-Wallis test $P = 0.4647$). Thus, bacteria that evolved in the inflamed gut possess a

fitness advantage in the inflamed gut environment, suggesting that evolution in the inflamed

gut results in specific adaptation to that environment.

Inflamed gut environment displays an altered metabolomic profile

To gain a better understanding of the environment to which these bacteria are adapted,

we performed metabolomic analysis of fecal samples spanning the experimental time course

(weeks 1, 2, 5, 6, 10, and 12) using 1H-NMR in mice from one of the two experiments. While

WT and KO mice initially display very similar fecal metabolomes (PERMANOVA $P =$

0.754), with time, the fecal metabolic composition begins to vary concurrently with the onset

of inflammation and the frequency of the C>T SNP upstream of *mprA* (Fig 3B, 1B, 2C). By

week 12 of the evolution experiment, WT and KO mice exhibit notable differences in their

fecal metabolomes, as revealed by PLS-DA (PERMANOVA $P = 0.001$; Fig 3B, Fig S7).

Among the 77 metabolite features initially found in the samples from week 1, none show

significantly different abundances between the WT and KO mice (Table S12). In contrast, in

the week 12 samples, 59 of the 77 detected features exhibit significant differences in

abundance between WT and KO mice (Benjamini-Hochberg corrected $P < 0.05$; Table S13).

Notably, 42 are significantly more abundant in KO mice than in WT mice, including all

detected amino acids that were significantly enriched in KO mice based on KEGG pathway

analysis of parallelly mutated loci. Among the 17 enriched features in the WT mice, most are

putatively annotated as (poly-)saccharides (Fig S8). Notably, variations in fecal metabolic

composition at week 12 could arise from changes in both the host, linked to inflammation-

related changes, and the bacterial population, associated with bacterial adaptation to

inflammation.

Phenotypic profiling of evolved bacteria compared to ancestor

Having established that the environment in which the bacteria evolved significantly

differs according to the mouse genotype, we next focused on specific bacterial traits that may

have changed as a consequence of adaptation. Accordingly, we tested the metabolic activity

of the *E. coli* populations using the Biolog GENIII test panel, which comprises 94 unique

biochemical tests, including a range of compounds (e.g., sugars, amino acids, and short-chain

fatty acids), conditions (e.g., different pH and salt concentrations), and chemical sensitivities

(e.g., antibiotics; see Methods). Similar to the pattern observed for the fecal metabolome, the

overall bacterial metabolic activity across all tested conditions is similar among populations

derived from the WT and KO mice at week 1 of the experiment (PERMANOVA, $P = 0.519$;

Fig S9), as also evidenced by the overlapping of the groups in the PLS-DA based on the

overall metabolic activity observed in the BIOLOG GENIII tests (Fig 3C). However, by

week 12, the overall pattern of metabolic activity across all tested conditions significantly

differs between the bacteria from KO and WT mice (PERMANOVA, $P = 0.005$, Fig S9).

These results suggest that adaptation to the inflamed intestine results in significantly altered

bacterial metabolism and the ability to grow in the presence of different inhibitors. Among

the compounds included in this screen, bacteria from WT and KO mice from week 1 showed

significantly different metabolic activity in only one compound (Table S14). In contrast,

bacteria from WT and KO mice at week 12 display significantly different metabolic activities

in the presence of 31 compounds (Wilcoxon signed-rank test, corrected $P < 0.05$; Table S15,

Fig S10).

To determine whether the differences in growth observed among the 31 compounds at

week 12 may be related to the altered metabolomic environment of inflamed mice, we

compared our BIOLOG GENIII results to the compounds found to be enriched in the fecal

metabolome of the KO mice. Interestingly, metabolomics analysis revealed a significant

enrichment of the amino acid histidine in the feces of KO mice compared to that of WT mice

(Fig S8). In our *in vitro* metabolic assay, we found that bacteria adapted to the inflamed

intestine had a significantly higher ability to metabolize histidine than bacteria adapted to the

healthy intestine (Fig S10). Thus, adaptation to an environment enriched in histidine may

have conferred a higher metabolic activity in the presence of this amino acid.

To further explore how the results of our *in vitro* phenotypic analyses may relate to

the fecal metabolomic profile observed *in vivo*, we performed a shortest path analysis on the

metabolic network of the genome-scale metabolic model of *E. coli* NC101 [32] using

metabolomic data and data from the *in vitro* metabolic activity screen (BIOLOG plates; see

Methods). Pairs of compounds from the two datasets that are converted easily to each other

by means of pathway length include glutamate and L-arginine (Fig 3D, Table S16).

Glutamate is significantly more abundant in the feces of the inflamed mice (Fig S8), whereas

bacteria adapted to the inflamed intestine are capable of significantly higher metabolism in

the presence of L-arginine (Fig S10). This result suggests that the bacteria may be converting

arginine to glutamate, resulting in the enrichment of glutamate in KO mice (Fig 3D, Table

S16). A second pair of compounds showing the same pattern are taurine and L-arginine.

However, analysis of the potential pathways between these compounds through flux

variability analysis revealed that taurine cannot be produced by *E. coli* NC101 under any of

the tested conditions (Fig S11). Additionally, taurine abundance increased steadily over time

only in KO mice (Fig S12). Thus, taurine is most likely to be produced by the host.

Finally, we examined 31 significant differences in metabolic activity observed

between bacteria from WT and KO mice at week 12, in light of their potential clinical value.

A first compound of interest is N-acetyl beta-D mannosamine (NADM), which is a precursor

to sialic acid and has been shown to promote *E. coli* colonization of inflamed gut [33]. Our *E.*

coli populations show higher levels of metabolic activity in the presence of NADM after

adaptation to the inflamed mouse gut. In contrast, *E. coli* evolved in the non-inflamed mouse

gut show no such change (Fig 4). These findings suggest that increased virulence may occur

in the context of adaptation to an inflamed environment.

Another interesting candidate is lithium chloride, which is known to have anti-

inflammatory effects via the inhibition of a key host regulator of inflammation, glycogen

synthase kinase-3 beta [34]. In our experiment, bacterial populations evolved in WT mice

display improved metabolic capacity in the presence of lithium chloride compared to the

ancestor, while no such change is observed among those evolved in the KO mice. Thus, our

results suggest that an inflamed intestinal environment may prevent the acquisition of this

phenotype in *E. coli*.

A final promising category of candidates are antibiotics, including fusidic acid,

vancomycin, and lincomycin. When considering the loading plot of the PLS-DA of our

Biolog GENIII data (Fig S13; reporting which compounds contribute the most to the

separation between groups), all three of these antibiotics contributed to the overall significant

difference in the metabolic activity of the bacteria from WT and KO mice (Fig 3C, Fig S13).

Furthermore, fusidic acid is also significantly different between bacteria from the two mouse

genotypes in the univariate analysis (Wilcoxon signed-rank test, corrected $P < 0.05$). Fusidic

acid is an antibiotic that targets gram-positive bacteria, and is commonly used to treat skin

infections [35]. Interestingly, it was also shown to be effective in reducing disease activity in a

small number of patients with Crohn's disease, which is thought to be due to its

immunosuppressive properties [36,37]. We observe *E. coli* to display decreased metabolic

activity in the presence of fusidic acid after adaptation to the inflamed intestine but increased

metabolic activity after adaptation to the healthy intestine. These results suggest a potential

trade-off between adaptation to inflammation and resistance to fusidic acid, where *E. coli*

adapted to inflammation also show lower resistance to fusidic acid.

In summary, we observe widespread phenotypic differences among the bacterial

populations that evolved in the inflamed intestines of KO mice compared to non-inflamed

WT mice. These results relate to differences in the metabolome between these two intestinal

environments and include phenotypes that carry the potential for exploitation in a clinical

setting.

Discussion

In this study, we describe the findings of an *in-vivo* bacterial evolution experiment

using a gnotobiotic mouse model of IBD. We find that *E. coli* evolved in the inflamed mouse

gut accumulated specific genetic changes and that these changes confer a fitness advantage in

the inflamed intestinal environment, which significantly differs in its metabolome.

Furthermore, we show that *E.coli* populations in the gut of healthy and inflamed mice have a

distinct metatranscriptomic profile. Finally, we showed that adaptation to the inflamed

intestine resulted in several phenotypic differences that may be clinically relevant, such as

differential tolerance to antibiotics.

Recent studies have highlighted how strain-level changes in members of the gut

microbiome can play a crucial role in the adaptation of the gut microbiome to novel

conditions [14,38,39]. However, the effects of inflammation on the evolution of gut commensals

remain largely unexplored. Barreto et al. (2020) followed the adaptation of *E. coli* in mice of

different ages *in vivo* [14]. They showed that the aged mouse gut, which also showed high

levels of inflammation, was a more stressful environment for *E. coli*, resulting in a higher

number of mutations and more severe selective pressure on commensals, particularly in

bacterial loci associated with stress-related functions. Notably, we find no difference in the

number of mutations acquired by *E. coli* in inflamed mice, suggesting that other factors

associated with aging may contribute to the increased bacterial mutation rate observed by

Barreto et al. (2020) [14].

More recently, Tawk et al. (2023) conducted a study of mice monocolonized with

Bacteroides thetaiotaomicron that were subsequently infected with *Citrobacter rodentium*

and developed inflammation. In this setting, a single-nucleotide variant of *B.*

thetaiotaomicron, which had a higher tolerance to oxidative stress than the ancestral variant,

underwent selective sweeps and dominated the intestinal community [39]. Although the

experimental setup and duration of inflammation differ between this study and the current

study, our candidate mutation is also known to be associated with oxidative stress [40]. This is

consistent with the observations of Barreto et al. (2020) and Tawk et al. (2023) [14,39],

suggesting that oxidative stress is a key selective pressure in an inflamed environment.

Moreover, oxidative stress plays a major role in the pathophysiology of IBD [41]. Interestingly,

this is also confirmed by our metatranscriptomic analysis. In particular, 14 of the genes

upregulated in KO mice belong to the pathway of glutathione metabolism (eco00480), and

previous studies demonstrated that glutathione metabolism is involved in protection against

oxidative stress [42,43]. In addition, Sakamoto et al. reported *mprA* to be involved in resistance

to oxidative stress, together with the activity of *gshA* (glutathione synthase), which is also

upregulated in our populations[40]. Additionally, our metatranscriptomic data confirm the

findings of Tawk et al., suggesting a role for vitamin B6 metabolism [39]. We observe not only

the upregulation of nine genes belonging to the vitamin B6 metabolism (eco00750), but also

a significantly higher concentration of aspartate, a substrate of vitamin B6-dependent

enzymes, in the feces of the inflamed mice[39].

Importantly, many of our findings are consistent with the observations made in

clinical IBD settings. First, the KEGG pathways affected by parallel mutations specific to the

inflamed mice (Fig 2A) include D-amino acid-, pyruvate-, and thiamine metabolism. This is

in line with previous reports that the basic metabolism is reduced in IBD [27,44].

395   Second, the metabolomic profiling of the fecal samples reveals that KO mice have

significantly higher levels of many amino acids, which was also reported in IBD patients [45].

Furthermore, the results of our shortest path analysis suggest that bacterial production of

glutamate from arginine may underlie the enrichment of glutamate in the inflamed gut.

Although taurine-arginine was also one of the shortest paths we found, taurine could not be

produced by *E. coli* according to the flux variability analysis. Thus, taurine is most likely to

be produced by the host. Taurine is known to be a promoter of colonization resistance, and

infection has been shown to prime the microbiota against subsequent infections by inducing

host production of taurine [46]. Indeed, taurine was also reported to ameliorate inflammation in

rat models of inflammation [47]. Furthermore, taurine is a substrate for the microbiota-driven

production of hydrogen sulfide [48], and arginine is a precursor to polyamines that can protect

against reactive species such as hydrogen sulfide [49]. Thus, bacterial adaptation for improved

metabolic activity in the presence of arginine may be a response to the host-induced

inflammatory changes in the intestine.

409   Third, *E. coli* adapted to the inflamed mouse gut also showed some clinically relevant

phenotypes (Fig 4). N-acetyl beta-D mannosamine (NADM) is a precursor of polysialic acid,

a pathogenic determinant, and studies have shown that it can promote colonization of the

inflamed intestine by pathogenic *E. coli* [33,50]. As NADM is a crucial player in mediating

pathogenic activity in the inflamed intestine, the improved metabolic activity of

inflammation-adapted *E. coli* observed in the presence of NADM may be a phenotype

associated with virulence. This is consistent with reports of virulence-associated phenotypes,

such as hypermotility, selected during the evolution of an adherent-invasive *E. coli* in the

inflamed mouse gut [13]. Furthermore, lithium chloride is known to inhibit the activity of

glycogen synthase kinase 3-β, a master regulator of host chronic intestinal inflammation

mediated by toll-like receptors, and has been used as a treatment in a mouse model of IBD

[34,51]. However, the effect of lithium on the microbiome has not yet been elucidated. *E. coli*

tolerance to lithium ions is regulated through antiporters, and proline has been shown to

induce the uptake of lithium ions by *E. coli* [52,53]. Our finding that adaptation to the healthy

mouse intestine confers improved metabolic capacity in the presence of lithium chloride,

while adaptation to the inflamed intestine does not (Fig 4), implies that lithium may affect

both the host and microbiome. This may also imply that adaptation to the healthy intestine

affects the antiporter-dependent detoxification system of *E. coli*, resulting in improved ion

tolerance.

Lastly, we found that bacterial adaptation to the inflamed gut decreases metabolic

activity in the presence of fusidic acid, while adaptation to the healthy mouse gut improved

metabolic activity in the presence of fusidic acid (Fig 4). Fusidic acid is an antibiotic with T

cell-specific immunosuppressive effects that also stimulate gastric mucus secretion.

Furthermore, it was successfully applied to alleviate inflammation in rats, as well as to treat

selected patients with Crohn's disease for whom conventional treatment was ineffective [36,37].

However, the effect of fusidic acid on the gut microbiome in IBD has not yet been studied.

Our results imply that bacterial adaptation to the inflamed gut can result in a trade-off for

resistance to fusidic acid. Importantly, such trade-offs are a fundamental concept in

evolutionary medicine, where increased fitness in one context results in a consequent

decrease in fitness in another context, and are proposed as possible potent therapeutic

strategies against antibiotic resistance and cancer [18,54]. Thus, the therapeutic effect of fusidic

acid observed in patients with CD is due to trade-offs in microbiome adaptation to the

inflamed gut and/or its immunosuppressive effects on the host. Interestingly, it is known that

lipocalin-2 plays a role in iron-sequestration, which has been reported to have a role in

antibiotic resistance and sensitivity[55]. It is thus worth noting that there is no enrichment of

genes involved in iron metabolism among the mutated genes in the populations evolved in the

inflamed gut, as well as among the upregulated genes in these populations.

Current treatment options for IBD are largely limited to the treatment of inflammation

with a chronic risk of relapse. Corticosteroids, aminosalicylates, and immunosuppressive

agents are the conventional drugs of choice, but the safety and efficacy of novel emerging

strategies remain unclear (reviewed in [56]). Given that disease-specific aspects of the

microbiome have been shown to remain stable over long periods in IBD patients despite

treatment [57], a persistent microbial influence on the intestinal environment may be a key risk

factor for relapse. Thus, our results highlight new avenues of research involving evolution-

informed therapeutic strategies that exploit trade-offs to either prevent adaptation to

inflammation and/or help restore desirable ancestral traits in the microbiome.

While our results provide valuable insights into the potential role of evolution-

informed therapeutic strategies, it is important to acknowledge that it remains unknown

whether the same or similar genotypic and phenotypic changes would be observed in the

context of a complex microbial community as present in the human gut microbiome. Future

work should therefore include similar *in vivo* evolution experiments using a complex or

synthetic microbial community.

Methods

Bacterial strains

E. coli NC101 strain was obtained from Balfour Sartor, University of North Carolina,

Chapel Hill, NC, USA [58]. *Escherichia coli* NC101 is a mouse strain isolated from the

intestine of a WT 129S6/SvEv mouse raised under specific-pathogen-free (SPF) conditions

[58]. The strain was cultured at 37 °C in Luria Bertani (LB) medium with continuous shaking.

The ancestor strain for the *in vivo* experiment was obtained by plating the overnight culture

on LB agar and picking single colonies to be used as inocula for the *in vivo* evolution

experiment.

Mouse model

Two independent experiments were conducted to study *E. coli* adaptation to the

inflamed gut (Table S1). In the first experiment, "Exp1", a single inoculum was used for all

mice. In order to ensure that mutations present in a common inoculum could not be falsely

identified as "parallel" mutations, the second experiment, "Exp2" was performed with

independent inocula for each mouse. Accordingly, our definition of a parallel mutated gene

requires it to be mutated in at least two mice that were gavaged with a different inoculum.

Germ-free (GF) C57BL/6NTac (WT) and C57BL/6NTac-Il10em8Tac (KO) male mice were

purchased from Taconic Biosciences (Silkeborg, Denmark) and housed in the Germ-Free

Animal Facility at the Max Planck Institute for Evolutionary Biology (Ploen, Germany). GF

mice were maintained in sterile isolators (MB-10, Quip Laboratories, Delaware, USA) and

fed sterilized 50 kGy V1124-927 Sniff (Soest, Deutschland). The animals were allocated to

independent cages with a maximum of four mice per isolator until they reached an age of 12

weeks. Detailed information regarding the mice used in this study is presented in Table S1.

Initially, 19 WT and 15 KO mice were mono-colonized with the *E. coli* NC101 strain. Only

mice that survived the duration of the experiment (12 weeks; N = 14 WT mice and N = 11

KO mice) were included in the study. The study was performed in accordance with the

approved animal protocols and institutional guidelines of the Max Planck Institute for

Evolutionary Biology, Plön. Mice were maintained, and experiments were performed in

accordance with FELASA guidelines and German animal welfare law (Tierschutzgesetz § 11;

permits from Veterinäramt Kreis Plön: 1401-144/PLÖ-004697 and Veterinäramt Kreis Kiel:

244-509017/2018(107-11/18)).

The strains used as inocula were diluted in sterile PBS, and 200 µl (equivalent to

1x10^8 bacteria) were gavaged using a sterile gavage needle (Reusable Feeding Needles

18G, Fine Science Tools, Heidelberg, Deutschland). Fecal pellets were collected in a sterile

manner twice a week to study the evolution of *E. coli* for a total of 24 samples collected per

mouse and once a week for metabolomic investigation. An overview of the sampling plan is

presented in Fig 1A.

Histopathological evaluation

Mice were sacrificed at week 12, and colon tissue was collected and arranged to form

a Swiss-roll [59]. Hematoxylin and eosin-stained sections of colonic tissue Swiss rolls were

scored by two independent researchers in a blinded manner using the scoring system

described by Adolph et al. (2013) [60]. The score is composed of five sub-scores: mononuclear

cell infiltrate, crypt hyperplasia, epithelial injury or erosion, polymorphonuclear cell

infiltrates, and transmural inflammation. Each of the first four sub scores was awarded a

score from 0 to 3, whereas transmural inflammation was scored from 0 to 4, with a higher

score indicating a more severe level of inflammatory activity. The sum of the sub scores was

then multiplied by a factor based on the percentage of affected bowel length (1= < 10%; 2=

10-25%; 3= 25-50%; 4= >50%).

Feces processing

Feces weight and consistency were recorded. Feces were homogenized in 1.5 ml

sterile PBS on a horizontal vortexer (Vortex Mixer Modell Vortex-Genie® 2, Scientific

Industries, Bohemia, NY, USA) at maximum speed for 30 min, then separated in three

aliquots of 500 µl. Each aliquot was centrifuged at 10,000 rpm for 5 min. The supernatants

were transferred to a new tube and stored at -20 °C for subsequent lipocalin-2 concentration

measurement. One pellet aliquot was resuspended in 500 µl of PBS and used to prepare a 12-point 1/10 dilution series in a 96-well plate. Ten µl were plated onto an LB agar plate. Plates were incubated overnight at 37°C and colonies were counted to estimate the bacterial load. The bacterial load in the fecal samples was calculated by normalizing CFU counts by wet feces weight. The second pellet aliquot was resuspended in 1 ml RNAlater and stored at +4 °C for 24 h, after which the tube was centrifuged at 10,000 rpm for 5 min to remove the RNAlater, and the pellet was stored at -20 °C for subsequent nucleic acid extraction and sequencing. The final pellet aliquot was resuspended in 500 µl of LB containing 20% glycerol and stored at -70 °C for further phenotypic investigation.

Lipocalin-2 quantification

Lipocalin-2 concentration in the supernatants was measured using the commercial kit Mouse Lipocalin-2/NGAL DuoSet ELISA (R&D Systems, Minneapolis, MN, USA) for mouse Lipocalin-2 (DY1857). Testing was performed according to the manufacturer's instructions. The samples were diluted 1:10 and added to the plate. The optical density of each well was determined using a plate reader (SPARK, Tecan, Tecan, Männedorf, Switzerland) at 450 and 540 nm. Lipocalin-2 concentrations were normalized to the weight of feces for each sample.

Shotgun sequencing

Total DNA and RNA were extracted using the ZymoBIOMICS DNA/RNA Mini Kit (Zymo Research, Freiburg, Germany), following the manufacturer's instructions from fecal pellets collected at weeks 1, 4, 8, and 12, as well as from the *E. coli* NC101 cultures that served as inocula. Shotgun metagenomic sequencing was performed on four Illumina

Nextseq (HighOutput 300 cycles) sequencing runs at the Max Planck Institute for

Evolutionary Biology (Plön, Germany).

Raw reads were filtered and trimmed to ensure good quality using Cutadapt (version

3.2, [61]). First, any pair containing Ns, homopolymers (10 nucleotides or more), or those

longer than 151 bp were discarded. Sequences were trimmed with a quality cutoff of 25 at

both ends for both reads, Illumina adapters were removed, and sequences shorter than 50 bp

were discarded. Good quality sequences were then filtered to exclude mouse sequences

(mouse_C57BL) using KneadData [62]. A final filter was applied to remove any adapter

leftovers using Trimmomatic [63] and sequences shorter than 105 bp were discarded. Mutations

were identified using Breseq [64] by comparing the obtained metagenomes with the reference

genome of *E.coli* NC101 (PRJNA596436) using default options (i.e. mutations identified by

a threshold frequency > 0.05) and by subtracting the mutations that were detected in the

respective inoculum. Gdtools was used to compare mutations in the samples.

Metatranscriptomics

Depletion of rRNA from extracted RNA was performed using QIAseq FastSelect -

5S/16S/23S (Qiagen). Samples were incubated at 89°C for 8 minutes. Libraries for

metatranscriptomics were prepared using the Illumina TruSeq® Stranded mRNA Library

Prep Kit (Illumina) according to the manufacturer's instructions. Shotgun metatranscriptomic

sequencing of week 12 fecal samples was performed on an Illumina NextSeq 500/550 using

the HighOutput Kit v2.5 (75 cycles).

Quality control and trimming were performed as described above for shotgun

sequencing. Gene expression in the evolved population was calculated in Geneious (v

2022.2.1; https://www.geneious.com) by mapping reads to the reference genome of *E.coli*

NC101 (PRJNA596436). Differential expression was calculated using DeSeq2[65] as

implemented in Geneious, using the parametric model and by assigning the conditions to

"WT-" or "KO-evolved".

Measurement of the metabolic activity of bacteria

Biolog GENIII MicroPlates (Biolog, Hayward, CA, USA) was used to investigate the

metabolism of evolved populations. Evolved populations stored in glycerol were inoculated

onto the Inoculation Fluid-A (IF-A) provided by the manufacturer such that the OD_{600} was

between 0.02-0.05. After adjusting the OD_{600}, the inoculated IF-A was shaken well and

distributed into a Biolog GENIII MicroPlate (100 μl in each well). The plate was then

covered with a sterile Breathe-Easy membrane (Sigma-Aldrich, St. Louis, MO) and placed in

a plate reader (SPARK, Tecan, Männedorf, Switzerland) at a preset temperature of 37 °C.

The plate reader was then run with the following parameters: temperature: 37 °C, shaking:

250 rpm, OD_{590} measurement: every 15 min, total running time: 36 h. From the OD_{590}

measures, area under the curve (AUC) was used as a proxy for metabolic activity. The Biolog

assay was repeated for populations from fecal samples collected at weeks 1 and 12 from all

25 mice, with two replicates each.

Ex vivo assay

The cecal content was collected from each mouse during dissection at the end of the

second experiment. The amount of cecal content from each mouse was different based on the

amount in the cecum at the time of dissection. Sterile PBS was added to the cecal content,

and the volume was adjusted based on the amount of cecal content collected from each

mouse (20 mg/ml). The protocol was adapted from Kitamoto et al. (2020) [66]. The PBS-cecal

content mixture was centrifuged twice (once at 500*g* for 5 min and again at 10000*g* for 5

min) and then filter-sterilized through a 0.2 μm filter. The *ex vivo* medium from each mouse

was mixed well and 100 μl was spread onto an LB agar plate to ensure that they were sterile.

Ex vivo media from all healthy mice were pooled, as were the *ex vivo* media from all

inflamed mice, resulting in two *ex vivo* media. Populations from fecal samples collected at

week 12 from a subset of mice (n = 5 KO and n = 5 WT mice) were inoculated into the *ex*

vivo media and incubated in the TECAN Spark plate reader at a preset temperature of 37 °C.

The plate reader was then run with the following parameters: temperature: 37 °C, shaking:

250 rpm, OD_{600} measurement: every 15 min, total running time: 36 h. From the OD_{600}

measures, area under the curve (AUC) was used as a proxy for growth.

Metabolomic evaluation of the fecal pellets

During the second experiment, samples for metabolomic investigation were collected

once a week and snap-frozen in liquid nitrogen. Samples were stored at -70 °C and delivered

at Helmholtz Center Munich for metabolomic investigation. A non-targeted metabolomics

approach of mouse fecal samples was undertaken using NMR spectroscopy. To extract the

aqueous metabolites, we homogenized 2-3 fecal pellets in 1 mL H_2O using ceramic beads

(NucleoSpin, Macherey–Nagel, Dueren, Germany) and a TissueLyser (Qiagen, Hilden,

Germany), mixing the sample for 3×30 s at 4,500 rpm with a 10 s cooling break (< 0°C).

Subsequently, the homogenate was centrifuged (13,000 rpm for 10 min at 4 °C), the

supernatant evaporated using a SpeedVac, and the dried extract reconstituted in 200 μL NMR

buffer (10% D_2O, 100 mM phosphate buffer with 0.1% trimethylsilyl-tetradeuteropropionic

acid (TSP), pH 7.4). Samples were transferred to 3 mm NMR tubes, and immediate NMR

analysis was performed in a randomized order with a Bruker 800 MHz spectrometer

operating at 800.35 MHz equipped with a Bruker SampleJet for sample cooling (283 K) and a

QCI-cryogenic probe. A standard one-dimensional pulse sequence (noesygppr1d) provides an

overview of all molecules. The acquisition parameters were as follows: water suppression

irradiation during recycle delay (2 s), mixing time of 200 ms, 90 °pulse of 12.5 μs. We

collected 512 scans of 64 K data points with a spectral width of 12 ppm. The software

TopSpin 3.6 (Bruker BioSpin, Ettlingen, Germany) was used for processing, i.e., Fourier

transformation, manual phasing, baseline correction, and calibration to TSP (δ 0.00). Data

were imported into Matlab software R2011b (Mathworks, Natick, MA, USA) and further

processed, i.e., the water region was removed, baseline adjusted [67] and spectra normalized [68].

Relative quantification of metabolites was performed using the peak heights of selected peaks

and compounds identified as described in our published workflow [69].

Shortest path analysis

For the shortest path analysis, we used the metabolic model reconstructed using the

AGORA2 resource for the human microbiome [32]. The model was translated into a graph with

metabolites as nodes, and the reactions converting metabolites into one another as edges. The

shortest path analysis was performed using the Dijkstra algorithm implemented in the igraph

package (1.3.5) of R (4.2.2) [70,71]. To avoid shortcuts through the network using cofactors as

intermediate compounds, the edges were weighted by the sum of degree of the nodes it

connected, as suggested by Faust et al. [72]. For each compound from the *in vitro* screen, we

calculated the set of shortest paths to all other metabolites in the model. Each value in the set

was the average shortest path of the five shortest paths between two metabolites, which was

performed to account for uncertainties in pathway calculation based only on the network

properties. For our analysis, we only considered pathways between compounds from the two

datasets, which were among the shortest 5% of pathways in a pathway set.

Prediction of possible conversions between metabolites

We wanted to understand whether the altered metabolic capabilities observed in the Biolog

plates or the changed environmental conditions observed in the metabolomic data, together

with the metabolism of *E. coli* could contribute to the observed changes in the metabolomic

data between KO and WT mice. To this end, we performed flux variable analysis (FVA) on

the metabolic model for E. coli NC101 (retrieved from AGORA2) [32]. We defined every

significantly detected metabolite in the Biolog experiment or the metabolomic data set as a

possible source metabolites, while the metabolites from the metabolomic data set were

defined as target metabolites. Metabolites found in the metabolomics are those known to be

present *in vivo*. Compounds found to be significantly differentially metabolized in the Biolog

assay are those that we know the evolved bacteria are capable of metabolizing. By using the

sum of these as the source and the detected metabolites as the target, we aimed to capture as

many of the possible conversions as possible to help explain the composition of the *in vivo*

environment and the role of bacterial metabolism in shaping it. For the simulation

background, we employed a minimal medium (Table S13), removed D-Glucose, and adjusted

the oxygen level according to the conditions tested (anoxic = 0 mmol h^{-1} gDW^{-1},

microaerobic = 1 mmol h^{-1} gDW^{-1}, aerobic = 10 mmol h^{-1} gDW^{-1}). For the simulation, we

added each source metabolite individually to the growth medium (100 mmol h^{-1} gDW^{-1}) and

calculated the maximum production rates of the target metabolites, assuming that at least

50% of maximum growth rates were achieved. We considered a maximum flux of $>1e^{-6}$ as

the possible production of the target metabolite from the source metabolite. FVA was

performed in cobrapy [73] and analyses were performed using the data.table and ggplot2

packages for R.

Statistical investigation

Statistical analyses were performed using packages in RStudio 2023.03.0+386. For

PERMANOVA analyses, adonis was used [74] on Bray-Curtis distances of the data [75] and

mixomics was used for PLS-DA [76]. AUCs were calculated using desctools [77].

Acknowledgments

We are grateful to Balfour Sartor for providing the NC101 *E. coli* strain, the MPI-Plön mouse

team for support with animal experiments, Britt Marie Hermes for support in manuscript

preparation, and Wael Elhenawy and Isabel Gordo for their helpful and constructive

discussions. The authors acknowledge the use of ChatGPT for the purpose of supporting the

drafting of material for the introduction section.

Funding

This study was supported by the Deutsche Forschungsgemeinschaft (DFG) Research Unit

FOR5042 "miTarget – The Microbiome as a Target in Inflammatory Bowel Disease"

(subprojects P10, Z, and INF) and Cluster of Excellence 2167 "Precision Medicine in

Chronic Inflammation" (grant no. EXC2167). RU was funded by the International Max-

Planck Research School for Evolutionary Biology (IMPRS EvolBio). Work in the

Unterweger group is funded by the German Federal Ministry for Education and Research

(grant 01KI2020).

Disclosure statement

The authors report there are no competing interests to declare.

Data availability statement

Escherichia coli NC101 reference genome is available under Bioproject number

PRJNA596436. Shotgun metagenomic sequencing of the inocula and evolved

populations and metatranscriptomic of the populations at week 12 are available with

Bioproject number PRJNA1012288.

Author contributions

MV, DU, and JB conceptualized the study; RU, NAA, MV, SH, MAG, OV, SK and DU

performed experiments; RU, NAA, MV, SH, FT, JH, and PR analyzed the data; JT and CK

performed the metabolic modeling; RU, NAA, and AD performed the statistical analyses.

RU, NAA, DU, and JB wrote the first draft of the manuscript. All coauthors revised the

manuscript and agreed to its publication.

Figures captions

Figure 1. Application of a mouse model of IBD to test the effect of gut inflammation on the evolution of NC 101

A Schematic of the experimental setup. Germ-free WT and *Il10*$^{-/-}$ mice were monocolonized with *E. coli* NC101. Fecal samples were collected from each mouse at the indicated time points and analyzed as described. **B** Fecal lipocalin-2 levels from the WT and *Il10*-/- mice. Lipocalin-2 was measured using the Mouse Lipocalin-2/NGAL DuoSet ELISA and normalized to feces weight. Each dot represents one mouse at each sampling point. Results of Wilcoxon signed-rank test of lipocalin-2 concentrations at each time point are reported in Table S3. Differences were considered statistically significant at Benjamini-Hochberg-corrected $P < 0.05$. **C** Boxplots of the histopathology scores of the colon tissue of mice after sacrifice at the end of the experiment. Each dot represents a single mouse. Differences were considered statistically significant using the Kruskal-Wallis H test at Benjamini-Hochberg-corrected $P < 0.05$. **D** Number of *de-novo* mutations in the bacterial populations at week 12 in

WT and KO mice compared with the reference genome. Each dot represents a bacterial

population from a single mouse. Differences were considered statistically significant using

the Kruskal-Wallis H test at Benjamini-Hochberg-corrected P <0.05. **E** Partial least squares-

discriminant analysis of the *de-novo* mutated genes in the evolved populations of *E. coli*

NC101 at week 12 in WT and KO mice compared to the reference genome. Differences were

considered statistically significant at a PERMANOVA $P < 0.05$.

Figure 2. Parallel evolution of *E. coli* in inflamed mice revealed a disease-specific genetic

signature A Parallelly mutated genes were classified based on KEGG pathways; **B** An

intergenic C>T mutation 10-nt upstream of *mprA* was the unique parallel mutation

significantly more abundant in populations evolved in the inflamed gut (Wilcoxon signed-

rank test Benjamini-Hocheberg corrected *P* < 0.005). The heatmap shows the frequency of

the mutation in bacterial populations from each mouse. **C** Mutation frequency (left axis) of

the intergenic C>T mutation 10-nt upstream of *mprA* and mean of lipocalin-2 concentrations

across KO mice at different timepoints during the *in-vivo* evolution experiment. Each line

represents a population from a single mouse. Each bar represents the mean value of the

normalized concentration of lipocalin-2 in fecal samples from KO mice.

Figure 3. Phenotypic characterization of evolved *E. coli* and the inflammatory environment in which they were selected.

A Evolved populations from a subset of mice (n = 5 KO mice and n = 5 WT mice) were cultured in media derived from the cecal content collected from healthy (WT) and inflamed (KO) mice after sacrifice. The area under the curve (AUC) calculated from growth curves measured during the growth of evolved populations from each mouse in inflamed and healthy cecal contents is shown. Each dot represents the mean of three independent replicates. * *P* < 0.05 (Kruskal-Wallis H test); ns, not significant. **B** Partial least squares discriminant analysis (PLS-DA) of the fecal metabolite abundances at different time points during the evolution experiment. Differences were considered statistically significant at PERMANOVA *P* < 0.05 **C** Bacterial metabolism was measured *in- vitro* in a range of compounds. Measurements were performed on bacterial populations derived from fecal samples collected at weeks 1 and 12 of the *in-vivo* evolution experiments. Partial least squares discriminant analysis (PLS-DA) was performed on the area under the curves (AUCs) of the measured metabolic activity of the

evolved bacterial populations from all mice in the range of compounds. Differences were

considered statistically significant at PERMANOVA $P < 0.05$ **D** Compounds from the fecal

metabolome and the *in vitro* screen with the shortest path were inspected. One of the pairs

with the shortest path was glutamate-arginine. Glutamate was enriched in the fecal

metabolome of KO mice, whereas KO-adapted bacteria showed higher metabolic activity in

the presence of arginine. Each dot represents the abundance of the compound in the fecal

sample from a mouse (for glutamate) or the mean metabolic activity of the bacterial

population from a mouse in the presence of L-arginine from two independent measurements

(for arginine). * $P < 0.05$, (Kruskal-Wallis test at Benjamini-Hochberg-corrected), ns, not

significant.

Figure 4. *E. coli* adapted to the inflamed gut show phenotypes with clinical relevance.

Metabolic activity of bacteria isolated at weeks 1 and 12 of the evolution experiment in the presence of three selected compounds. Area under the curve was calculated from growth curves measured during the growth of evolved populations in the presence of each compound. Each dot represents the mean of two independent replicates of the metabolic activities of the *E. coli* population from a mouse measured in two independent experiments. *, $P < 0.05$ (Kruskal-Wallis test at Benjamini-Hochberg-corrected). The clinical relevance of the three compounds is indicated.

References

References

1.   Ott, S. J. *et al.* Reduction in diversity of the colonic mucosa associated bacterial

microflora in patients with active inflammatory bowel disease. *Gut* **53**, 685–693 (2004).

2.   Gevers, D. *et al.* The treatment-naïve microbiome in new-onset Crohn's disease. *Cell*

Host Microbe **15**, 382–392 (2014).

3.   Rehman, A. *et al.* Geographical patterns of the standing and active human gut

microbiome in health and IBD. *Gut* **65**, 238–248 (2016).

4.   Halfvarson, J. *et al.* Dynamics of the human gut microbiome in inflammatory bowel

disease. *Nat. Microbiol.* **2**, 17004 (2017).

5.   Schirmer, M. *et al.* Dynamics of metatranscription in the inflammatory bowel disease

gut microbiome. *Nat. Microbiol.* **3**, 337–346 (2018).

6.   Schirmer, M., Garner, A., Vlamakis, H. & Xavier, R. J. Microbial genes and pathways

in inflammatory bowel disease. *Nat. Rev. Microbiol.* **17**, 497–511 (2019).

7.   Willing, B. P. *et al.* A pyrosequencing study in twins shows that gastrointestinal microbial

profiles vary with inflammatory bowel disease phenotypes. *Gastroenterology* **139**, 1844-

1854.e1 (2010).

8.   Barreto, H. C. & Gordo, I. Intrahost evolution of the gut microbiota. *Nat. Rev. Microbiol.*

(2023) doi:10.1038/s41579-023-00890-6.

9.   Giraud, A. *et al.* Costs and benefits of high mutation rates: adaptive evolution of bacteria

in the mouse gut. *Science* **291**, 2606–2608 (2001).

10.   Paepe, M. D. *et al.* Trade-Off between Bile Resistance and Nutritional Competence

Drives Escherichia coli Diversification in the Mouse Gut. *PLOS Genet.* **7**, e1002107

(2011).

11. Barroso-Batista, J. *et al.* The First Steps of Adaptation of Escherichia coli to the Gut Are

Dominated by Soft Sweeps. *PLOS Genet.* **10**, e1004182 (2014).

12. Zhao, S. *et al.* Adaptive Evolution within Gut Microbiomes of Healthy People. *Cell Host*

Microbe **25**, 656-667.e8 (2019).

13. Elhenawy, W., Tsai, C. N. & Coombes, B. K. Host-Specific Adaptive Diversification of

Crohn's Disease-Associated Adherent-Invasive Escherichia coli. *Cell Host Microbe* **25**,

301-312.e5 (2019).

14. Barreto, H. C., Sousa, A. & Gordo, I. The Landscape of Adaptive Evolution of a Gut

Commensal Bacteria in Aging Mice. *Curr. Biol.* **30**, 1102-1109.e5 (2020).

15. Daniel, H. Diet and Gut Microbiome and the "Chicken or Egg" Problem. *Front. Nutr.* **8**,

(2022).

16. Garland, T., Downs, C. J. & Ives, A. R. Trade-Offs (and Constraints) in Organismal Biology.

Physiol. Biochem. Zool. PBZ **95**, 82–112 (2022).

17. Nesse, R. M., Ganten, D., Gregory, T. R. & Omenn, G. S. Evolutionary molecular

medicine. *J. Mol. Med. Berl. Ger.* **90**, 509–522 (2012).

18. Merker, M. *et al.* Evolutionary Approaches to Combat Antibiotic Resistance:

Opportunities and Challenges for Precision Medicine. *Front. Immunol.* **11**, (2020).

19. Roemhild, R. & Schulenburg, H. Evolutionary ecology meets the antibiotic crisis: Can we

control pathogen adaptation through sequential therapy? *Evol. Med. Public Health*

2019, 37–45 (2019).

20. Andersson, D. I. *et al.* Antibiotic resistance: turning evolutionary principles into clinical

reality. *FEMS Microbiol. Rev.* **44**, 171–188 (2020).

21. Melnikov, S. V. *et al.* Exploiting evolutionary trade-offs for posttreatment management

of drug-resistant populations. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 17924–17931 (2020).

22. Hausser, J. *et al.* Tumor diversity and the trade-off between universal cancer tasks. *Nat.*

Commun. **10**, 5423 (2019).

23. Zhang, J., Cunningham, J. J., Brown, J. S. & Gatenby, R. A. Integrating evolutionary

dynamics into treatment of metastatic castrate-resistant prostate cancer. *Nat. Commun.*

8, 1816 (2017).

24. Jacqueline, C. *et al.* Cancer: A disease at the crossroads of trade-offs. *Evol. Appl.* **10**,

215–225 (2016).

25. Cunningham, J. J., Gatenby, R. A. & Brown, J. S. Evolutionary dynamics in cancer therapy.

Mol. Pharm. **8**, 2094–2100 (2011).

26. Kim, S. C., Tonkonogy, S. L., Karrasch, T., Jobin, C. & Sartor, R. B. Dual-association of

gnotobiotic IL-10-/- mice with 2 nonpathogenic commensal bacteria induces aggressive

pancolitis. *Inflamm. Bowel Dis.* **13**, 1457–1466 (2007).

27. Morgan, X. C. *et al.* Dysfunction of the intestinal microbiome in inflammatory bowel

disease and treatment. *Genome Biol.* **13**, R79 (2012).

28. Davenport, M. *et al.* Metabolic alterations to the mucosal microbiota in inflammatory

bowel disease. *Inflamm. Bowel Dis.* **20**, 723–731 (2014).

29. Brooun, A., Tomashek, J. J. & Lewis, K. Purification and Ligand Binding of EmrR, a

Regulator of a Multidrug Transporter. *J. Bacteriol.* **181**, 5131–5133 (1999).

30. Coll, J. L., Heyde, M. & Portalier, R. Expression of the *nmpC* gene of *Escherichia coli* K-12

is modulated by external pH. Identification of *cis* -acting regulatory sequences involved

in this regulation. *Mol. Microbiol.* **12**, 83–93 (1994).

31. Hirakawa, H. *et al.* AcrS/EnvR Represses Expression of the *acrAB* Multidrug Efflux Genes

in *Escherichia coli*. *J. Bacteriol.* **190**, 6276–6279 (2008).

32. Heinken, A. *et al.* Genome-scale metabolic reconstruction of 7,302 human

microorganisms for personalized medicine. *Nat. Biotechnol.* 1–12 (2023)

doi:10.1038/s41587-022-01628-0.

33. Huang, Y.-L., Chassard, C., Hausmann, M., von Itzstein, M. & Hennet, T. Sialic acid

catabolism drives intestinal inflammation and microbial dysbiosis in mice. *Nat. Commun.*

6, 8141 (2015).

34. Hofmann, C., Dunger, N., Schölmerich, J., Falk, W. & Obermeier, F. Glycogen synthase

kinase 3-β: A master regulator of toll-like receptor-mediated chronic intestinal

inflammation. *Inflamm. Bowel Dis.* **16**, 1850–1858 (2010).

35. Fernandes, P. Fusidic Acid: A Bacterial Elongation Factor Inhibitor for the Oral Treatment

of Acute and Chronic Staphylococcal Infections. *Cold Spring Harb. Perspect. Med.* **6**,

a025437 (2016).

36. Langholz, E., Brynskov, J., Bendtzen, K., Vilien, M. & Binder, V. Treatment of Crohn's

disease with fusidic acid: an antibiotic with immunosuppressive properties similar to

cyclosporin. *Aliment. Pharmacol. Ther.* **6**, 495–502 (2007).

37. Kilic, F. S., Erol, K., Batu, O., Yildirim, E. & Usluer, G. The effects of fusidic acid on the

inflammatory response in rats. *Pharmacol. Res.* **45**, 265–267 (2002).

38. Dapa, T., Ramiro, R. S., Pedro, M. F., Gordo, I. & Xavier, K. B. Diet leaves a genetic

signature in a keystone member of the gut microbiota. *Cell Host Microbe* **30**, 183-

199.e10 (2022).

39. Tawk, C. *et al.* Infection leaves a genetic and functional mark on the gut population of a

commensal bacterium. *Cell Host Microbe* **31**, 811-826.e6 (2023).

40. Sakamoto, A. *et al.* Three Members of Polyamine Modulon under Oxidative Stress

Conditions: Two Transcription Factors (SoxR and EmrR) and a Glutathione Synthetic

Enzyme (GshA). *PLoS ONE* **10**, e0124883 (2015).

41. Tian, T., Wang, Z. & Zhang, J. Pathomechanisms of Oxidative Stress in Inflammatory

Bowel Disease and Potential Antioxidant Therapies. *Oxid. Med. Cell. Longev.* **2017**,

4535194 (2017).

42. Carmel-Harel, O. & Storz, G. Roles of the Glutathione- and Thioredoxin-Dependent

Reduction Systems in the *Escherichia Coli* and *Saccharomyces Cerevisiae* Responses to

Oxidative Stress. *Annu. Rev. Microbiol.* **54**, 439–461 (2000).

43. Łyżeń, R., Gawron, G., Kadziński, L. & Banecki, B. GSH Protects the Escherichia coli Cells

from High Concentrations of Thymoquinone. *Molecules* **27**, 2546 (2022).

44. Pasternak, B. A. *et al.* Lipopolysaccharide exposure is linked to activation of the acute

phase response and growth failure in pediatric Crohn's disease and murine colitis.

Inflamm. Bowel Dis. **16**, 856–869 (2010).

45. Jagt, J. Z. *et al.* Fecal Amino Acid Analysis in Newly Diagnosed Pediatric Inflammatory

Bowel Disease: A Multicenter Case-Control Study. *Inflamm. Bowel Dis.* **28**, 755–763

(2022).

46. Stacy, A. *et al.* Infection trains the host for microbiota-enhanced resistance to

pathogens. *Cell* **184**, 615-627.e17 (2021).

47. Son, M., Ko, J. I., Kim, W. B., Kang, H. K. & Kim, B. K. Taurine can ameliorate

inflammatory bowel disease in rats. *Adv. Exp. Med. Biol.* **442**, 291–298 (1998).

48. Buret, A. G., Allain, T., Motta, J.-P. & Wallace, J. L. Effects of Hydrogen Sulfide on the

Microbiome: From Toxicity to Therapy. *Antioxid. Redox Signal.* **36**, 211–219 (2022).

49. Maeda, T. *et al.* Role of polyamines derived from arginine in differentiation and

proliferation of human blood cells. *Biol. Pharm. Bull.* **29**, 234–239 (2006).

50. Revilla-Nuin, B. *et al.* Transport of N-acetyl-D-mannosamine and N-acetyl-D-glucosamine

in Escherichia coli K1: effect on capsular polysialic acid production. *FEBS Lett.* **511**, 97–

101 (2002).

51. Freland, L. & Beaulieu, J.-M. Inhibition of GSK3 by lithium, from single molecules to

signaling networks. *Front. Mol. Neurosci.* **5**, 14 (2012).

52. Inaba, K. *et al.* Lithium toxicity and Na+(Li+)/H+ antiporter in Escherichia coli. *Biol.*

Pharm. Bull. **17**, 395–398 (1994).

53. Chen, C. C., Tsuchiya, T., Yamane, Y., Wood, J. M. & Wilson, T. H. Na+ (Li+)-proline

cotransport in Escherichia coli. *J. Membr. Biol.* **84**, 157–164 (1985).

54. Gatenby, R. A. & Brown, J. S. Integrating evolutionary dynamics into cancer therapy.

Nat. Rev. Clin. Oncol. **17**, 675–686 (2020).

55. Ezraty, B. & Barras, F. The 'liaisons dangereuses' between iron and antibiotics. *FEMS*

Microbiol. Rev. **40**, 418–435 (2016).

56. Cai, Z., Wang, S. & Li, J. Treatment of Inflammatory Bowel Disease: A Comprehensive

Review. *Front. Med.* **8**, 765474 (2021).

57. Öhman, L. *et al.* Fecal microbiota dynamics during disease activity and remission in

newly diagnosed and established ulcerative colitis. *Sci. Rep.* **11**, 8641 (2021).

58. Kim, S. C. *et al.* Variable phenotypes of enterocolitis in interleukin 10-deficient mice

monoassociated with two different commensal bacteria. *Gastroenterology* **128**, 891–

906 (2005).

59. Moolenbeek, C. & Ruitenberg, E. J. The 'Swiss roll': a simple technique for histological

studies of the rodent intestine. *Lab. Anim.* **15**, 57–59 (1981).

60. Adolph, T. E. *et al.* Paneth cells as a site of origin for intestinal inflammation. *Nature* **503**,

272–276 (2013).

61. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.journal* **17**, 10 (2011).

62. KneadData – The Huttenhower Lab. https://huttenhower.sph.harvard.edu/kneaddata/.

63. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

64. Deatherage, D. E. & Barrick, J. E. Identification of Mutations in Laboratory-Evolved

Microbes from Next-Generation Sequencing Data Using breseq. in *Engineering and*

Analyzing Multicellular Systems (eds. Sun, L. & Shou, W.) vol. 1151 165–188 (Springer

New York, 2014).

65. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

66. Kitamoto, S. *et al.* Dietary L-serine confers a competitive fitness advantage to

Enterobacteriaceae in the inflamed gut. *Nat. Microbiol.* **5**, 116–125 (2020).

67. Haslauer, K. E., Schmitt-Kopplin, P. & Heinzmann, S. S. Data Processing Optimization in

Untargeted Metabolomics of Urine Using Voigt Lineshape Model Non-Linear Regression

Analysis. *Metabolites* **11**, 285 (2021).

68. Dieterle, F., Ross, A., Schlotterbeck, G. & Senn, H. Probabilistic Quotient Normalization

as Robust Method to Account for Dilution of Complex Biological Mixtures. Application in

1H NMR Metabonomics. *Anal. Chem.* **78**, 4281–4290 (2006).

69. Heinzmann, S. S., Waldenberger, M., Peters, A. & Schmitt-Kopplin, P. Cluster Analysis

Statistical Spectroscopy for the Identification of Metabolites in 1H NMR Metabolomics.

Metabolites **12**, 992 (2022).

70. Csárdi, G. & Nepusz, T. The igraph software package for complex network research. in

(2006).

71. R: A Language and Environment for Statistical Computing | BibSonomy.

https://www.bibsonomy.org/bibtex/7469ffee3b07f9167cf47e7555041ee7.

72. Faust, K., Croes, D. & van Helden, J. Metabolic pathfinding using RPAIR annotation. *J.*

Mol. Biol. **388**, 390–414 (2009).

73. Ebrahim, A., Lerman, J. A., Palsson, B. O. & Hyduke, D. R. COBRApy: COnstraints-Based

Reconstruction and Analysis for Python. *BMC Syst. Biol.* **7**, 74 (2013).

74. Anderson, M. J. A new method for non-parametric multivariate analysis of variance.

Austral Ecol. **26**, 32–46 (2001).

75. Bray, J. R. & Curtis, J. T. An Ordination of the Upland Forest Communities of Southern

Wisconsin. *Ecol. Monogr.* **27**, 325–349 (1957).

76. Rohart, F., Gautier, B., Singh, A. & Cao, K.-A. L. mixOmics: an R package for 'omics

feature selection and multiple data integration. 108597 Preprint at

https://doi.org/10.1101/108597 (2017).

77. Signorell, A. *et al.* DescTools: Tools for Descriptive Statistics. (2023).

937