Inferring genetic structure of European beech from observations of spectral phenotypes.

Ewa A. Czyż[1*], Bernhard Schmid[1], Maarten B. Eppinga[1], Marylaure de La Harpe[1,a], Aboubakr Moradi[1,2,b], Cheng Li[1], Michael E. Schaepman[1], Meredith C. Schuman[1,2]

[1] Department of Geography, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland

[2] Department of Chemistry, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland

[a] Current address: Amt für Natur und Umwelt, Graubünden, Ringstrasse 10, 7000 Chur, Switzerland

[b] Current address: Syngenta Crop Protection AG, Schaffhauserstrasse, 4334 Münchwilen, Switzerland

[*] To whom correspondence should be addressed; E-mail: ewa.czyz@uzh.ch.

Abstract

Biodiversity loss presents a growing threat to the global environment and requires systematic and spatially contiguous monitoring. Monitoring of within-species genetic variation, a key factor when assessing biodiversity loss, is laborious and could be complemented by observations of phenotypes allowing inferences about genetic variation. We studied genetic and phenotypic variations in the common European beech (*Fagus sylvatica* L.) derived from whole-genome sequences and spectral phenotypes of more than 200 individuals at 22 sites across the species' natural range. The spectral phenotypes were collected under standardized illumination/observation conditions from the same top-of-canopy leaves used for nuclear DNA extraction. We found that spectral and environmental information explains 77.7% of the variance along the first two Principal Coordinates representing genetic structure among sampled individuals. Information from spectral phenotypes contributed 12.1% to predictions of between-site genetic structure. We therefore conclude that future remote observations having sufficient spectral and spatial resolution could improve our ability to rapidly and continuously monitor genetic diversity in trees.

Highlights

- Genetic diversity underpins biodiversity which loss requires systematic monitoring
- Spectral phenotypes acquired *in situ* are indicative of genetic variation
- Intraspecific spectral and genomic variation correlate across a species range
- Spectral phenotypes improve prediction of the between-site genetic structure
- Remote observations may contribute to monitoring genetic diversity within species

Keywords

Spectral phenotypes, intraspecific genetic variation, remote sensing, *Fagus sylvatica*, biodiversity monitoring

1. Introduction

Biodiversity loss presents a large and growing risk for the global environment and its ability to support life. Plant diversity loss can negatively affect ecosystem functioning (Purvis et al. 2000, Hector et al. 2007, Isbell et al. 2011, Cardinale et al. 2012, Reich et al. 2012), impact ecosystem stability (Tilman et al. 1994, Hector et al. 2010, Isbell et al. 2015, van Moorsel et al. 2021), and reduce the potential of populations and communities to adapt to environmental change (Des Roches et al. 2018). Biodiversity is under pressure from human activity (Johnson et al. 2009) and global change (Exposito-Alonso et al. 2022). To reduce human impact on biodiversity decline, targets have been set several times aiming to mitigate biodiversity loss worldwide (IPBES 2019), but many further steps need to be taken (Mora et al. 2006, Jenkins and Joppa 2009). One important step towards biodiversity support is its continuous and standardized monitoring (Scholes et al. 2008, Turner 2014). Monitoring indicators of biodiversity should help

51 to set conservation priorities (Scott et al. 1989), evaluate actions taken, and develop adaptive

52 management (Sutherland et al. 2004) and policy adjustments (Perrings et al. 2011) to preserve

53 and restore biodiversity. Monitoring schemes should span levels of biological organization from

54 ecosystems to species and genes. Genetic diversity is especially challenging to monitor, but can

55 provide valuable information about a species' contribution to ecosystem functioning (Fridley and

56 Grim 2010, Tang et al. 2022) and its evolutionary potential, and thus is a key component when

57 accounting for biodiversity loss (Wilson 1988, Exposit-Alonso et al. 2022).

58 The assessment of genetic diversity is generally based on field sample collection, with spatial

59 coverage restricted to a small part of species natural range and few individuals within sites.

60 Furthermore, genotyping is still commonly limited to small portions of whole genomes and may

61 not account for all functionally important genetic variation (Lou et al. 2021). These scale-induced

62 limitations result in sparse and biased representations of genetic diversity, which lead to

63 inaccurate representation of its current status, omission of unique but difficult-to-sample sites,

64 and partially misses components of the variation that reflects the total adaptation potential of a

65 species to environmental change (Hughes et al. 2008, Des Roches et al. 2018). Furthermore,

66 incomplete and biased spatial data inhibits understanding of species phylogeographic history

67 that constrains current genetic patterns. Whole-genome sequencing may provide a means to

68 assess intraspecific genetic diversity without the bias of more restricted sets of molecular

69 markers. However, the spatiotemporal limitations still apply and thus indicators of genetic

70 diversity in the absence of DNA sequence should be assessed (Hoban et al. 2022). Phenotypic

71 variation acquired over space may contribute to indicating and understanding species' genetic

72 diversity if a link between the phenotypic and genetic variation can be established across

73 different sites and environmental conditions.

74 To collect these phenotypic data, imaging spectrometers can provide repeated and spatially

75 contiguous acquisitions of spectral phenotypes comprising integrated signatures of plant

biochemical and structural traits (Guyot et al. 1992, Jacquemont and Ustin 2009, Schaepman et

al. 2009, Schaepman et al. 2015). These plant traits impact light interaction with plant tissues

and thus can affect the spectral phenotypes of individuals from leaf to canopy levels (Meireles et

al. 2020). Leaf spectral phenotypes are commonly estimated from spectroradiometer

measurements under standardized illumination/observations conditions, while the spectral

phenotypes of entire canopies at landscape-scale are derived from remote acquisitions of

reflected radiation by airborne and spaceborne imaging spectrometers (Schaepman et al. 2009).

Spectral phenotypes representative of plant traits result from the interaction of plant genes with

the environment, and thus spectral acquisitions might facilitate monitoring of genetic diversity

under natural conditions, if a link between spectral and genetic variation can be established

(Cavender-Bares et al. 2016, Yamasaki et al. 2017). Until now, however, this potential remains

largely untested. Previous studies associating genetic and spectroscopy data focused on

experimental settings (Matsuda et al. 2012, Cavender-Bares et al. 2016, Li et al. 2023), or when

under natural conditions, on single-site, between-species assessments with few within-species

assessments not covering the studied species' across its full natural range (Madricht et al. 2014,

Czyż et al. 2020, Czyż et al. 2023); furthermore previous studies were mostly based on genetic

markers only covering a small fraction of the species genome rather than integrate whole-

genome information.

Here, we demonstrate the potential of spectral phenotypes to contribute to assessing within-

species genetic structure across 22 sites using whole-genome sequences from more than 200

individual trees of *Fagus sylvatica* L. (European beech). We chose this test species because it is

a dominant, ecologically and economically important tree species in Europe that has gained

interest as a model for ecological genomics of tree species under global change (Capblancq et

al. 2020). We present the largest combined genomic and spectral sampling to date of the *F.*

sylvatica species range using short-read whole-genome data to derive genetic structure, and

top-of-canopy leaf reflectance acquired by contact probe to derive spectral phenotypes. We then

use spectral phenotypes, in combination with environmental variables, to predict the genetic structure with focuse on between-site assessments. We thereby (1) contribute to current knowledge about genetic structure in *F. sylvatica* across its natural range; and (2) indicate how spectral phenotypes acquired in natural environments may help to predict it. The results suggest that remote observations by aerial and upcoming satellite-based imaging spectroscopy could be leveraged to contribute to genetic diversity monitoring and conservation over larger areas than typically attainable when solely relying on field sampling.

2. Materials and Methods

2.1. Study areas and sample collection

We selected 22 sites dominated by *Fagus sylvatica* L. (common beech) across its natural range in Europe (Fig.**1a**). We assigned three-letter site names with the first two letters referring to the country that the site is located in, and the third letter referring to the particular location in the country. Most of the sites are covered by protected, old-growth beech forests. The variation among the sites accounts for climatic (mean 50% of the maximum span of environmental values across the range and 95% of the distribution of environmental values across the range, Supporting information Fig. **S5**), edaphic and phylogeographic variation (Caudullo et al. 2017). Data were collected *in situ* during August–September 2020. At each site we sampled 10 *F. sylvatica* trees that occupied canopy space along two ca. 10 x 200 m transects separated by 50 m. Each transect comprised 5 dominant trees separated by at least 30 m. At some sites we sampled fewer than 10 trees due to a lack of *F. sylvatica* trees in the canopy within the transect (4 trees at RSB, 8 trees at ESP) and in some cases the sampled material was of poor quality for the DNA extraction (1 tree at PLM, 1 tree at ITP). When possible, we selected transects on south-west to south-east-facing slopes to normalize for local adaptation to exposition to solar radiation. From each tree, we collected a top-of-canopy branch of 20 to 100 cm length with the help of tree climbers or, when tree height did not exceed 15 m, with a telescoping carbon fiber

pruner (Takeni Trading Co., Osaka, Japan). From each branch, we randomly selected three

undamaged leaves separated by at least 5 cm, from which we acquired spectral data after

harvesting (see Materials and Methods, Spectral data) and which we subsequently stored on

silica gel for later genetic analysis (see Materials and Methods, Genetic data).

2.2.Genetic data

The collected leaves were taken back to the laboratory where we took 50 mg of undamaged leaf

lamina and then stored these samples in tubes at –20 °C until grinding and DNA extraction. The

DNA was purified using a Norgen Plant and Fungi genomic DNA extraction kit (Norgen Biotek,

Thorold, ON, Canada), modified to reduce RNA contamination and to improve lysis, yield, and

purity. Specifically, samples were flash-frozen in liquid nitrogen immediately prior to grinding and

ground using 3 mm glass beads at 30 Hz for 3 min in a TissueLyser II (Qiagen) to a fine powder.

The powder was briefly centrifuged before adding lysis buffer and vortexed to mix with the

buffer. One µl of RNAse A (DNAse-free, 100,000 units/ml in 50% glycerol, 10 mM Tris-HCl, pH

8.0) and 3 µl Proteinase K (>600 µl ~ 20 mg/ml in 10 mM Tris-HCl pH 7.5, containing calcium

acetate and 50% v/v glycerol) were added to the lysis buffer prior to 10 min incubation at 56°C

with 1000 rpm agitation. An additional 1 µl of RNAse A was subsequently added to the reaction,

followed by a 10 min incubation at 65°C with 2000 rpm agitation. The samples were then

processed according to the manufacturer's protocol with the modification that 70% ethanol was

mixed with lysate by pipetting ca. 10x/sample instead of vortexing. The DNA from clarified lysate

was captured on spin columns (centrifugation until all lysate passed) and columns were washed

three times with Solution A prior to elution with Elution Buffer B (100 µl, 5 min RT incubation,

spinning until all elution buffer was recovered).

The purified DNA was used for sequencing library preparation (Novogene, Cambridge, UK):

genomic DNA was randomly sheared into short fragments, which were end repaired, A-tailed,

and ligated with Illumina adapters. The fragments with adapters were PCR amplified, size

selected, and purified. Libraries were sequenced by Illumina NovaSeq PE150 (Novogene,

Cambridge, UK) to produce ca. 11 Gbp/sample, resulting in a mean ca. 20x coverage of the 541

Mbp nuclear genome (Mishra et al. 2022). The Illumina adapters were trimmed from the raw

reads, and reads with phred scores below 3 and windows of 4 base pair sliding window with

average phred scores below 15 were also removed using *trimmomatic* v. 0.40. We aligned reads

against the chromosome-level reference genome (Mishra et al. 2022) in *bowtie2* v. 2.4.5 and

removed alignment duplicates and unaligned reads in *picard* v. 2.0.1. Sequentially, we derived

sets of alleles per each individual tree by calling variants against the chromosome-level

reference genome (Mishra et al. 2022) using the function *HaplotypeCaller* in *GATK4*. We

consolidated the derived alleles and obtained the joint SNPs among all individuals using the

function *GenotypeGVCFs* in *GATK4*. From the derived SNPs, we selected those with phred

scores above 30 (i.e., accuracy higher than 99.9%), minor allele frequencies of more than 0.1,

and minimum and maximum allele depths of 10 and 50, respectively. We pruned the derived

SNPs for linkage disequilibrium by removing SNPs that had a correlation coefficient greater than

0.1 with each other using a sliding window of 50kb in *plink* v. 1.9 software. This resulted in

1,380,310 SNPs across all sampled individuals used in further analyses. To date, patterns of *F.*

sylvatica genetic diversity and structure have been studied with marker-based approaches

including allozymes (Gallois et al. 1998), AFLP (Pluess and Weber 2012, Guevara et al. 2022),

microsatellites and, more recently, SNPs in candidate genes (Csilléry et al. 2014, Pfenninger et

al. 2020, Meger et al. 2021, Cuervo-Alarcon et al. 2021) and whole-genome sequences from

limited area of the specie range (Stefanini et al. 2023). We are not aware of another study so far

that used whole-genome sequences from across the range of *F. sylvatica*.

2.3. Spectral data

Spectral data comprised leaf reflectance spectra collected in the wavelength range from 350–

2500 nm using an ASD FieldSpec 4 spectroradiometer (serial nr 18739, ASD Inc., Boulder, CO,

USA), coupled to a plant probe with leaf clip attachment, and equipped with a standardized

halogen light source and a set of standard black and white reference backgrounds (serial nr 445,

ASD Inc., Boulder, CO, USA). Measurements of three leaves harvested from branches (1

branch/tree) were acquired within minutes to up to 3 h (for ITE, ITM, 5 trees at ITV, and 5 trees

at PLB) from the time of branch harvest using the contact probe positioned in the middle of the

leaf and to the right of the midrib as viewed from the adaxial side (i.e. the side of the leaf facing

the incident radiation (e.g. sun and light source). Measurements comprised five successive

readings per sample for four types of acquisitions: a white reference R_w, a leaf with white

reference as background T_w, a black reference R_b, and the leaf with black reference as

background T_b (cf., eq. 1). For each reading, five scans, each with an integration time of 8.5 ms,

were recorded and averaged per leaf. The averaged reflectance spectra per tree was calculated

from the mean of leaf reflectance R using a formula for background-differentiated measurements

(Miller et al. 1992):

$$R = \frac{T_b * R_w - T_w * R_b}{R_w - R_b} * 100 \ [\%]$$ [eq. 1]

The ASD spectroradiometer provides convolved sampling intervals of 1 nm spectral sampling,

although the spectral resolution of the instrument ranges from 3–10 nm sampling interval. To

minimize collinearity effects of neighboring spectral bands, we resampled the spectral signal

from the original 2151 spectral bands to 216 spectral bands by linearly interpolating the original

signal using the *interp1* function in *Matlab* (v. R2017b) followed by multiplication with a Spectral

Response Function (SRF) of Gaussian shape with Full Width at Half Maximum (FWHM) of 10

nm.

2.4. Climatic data

Climatic data comprised 19 bioclimatic variables derived from WorldClim Version2 (Fick et al.

2017) at a resolution of 1 km with global coverage. The correlated variables represent climatic

conditions and their annual trends that affect ecological processes across large gradients within

the range of *F. sylvatica* and are from the years 1970-2000, so represent conditions at the time

in which the study trees may have been establishing or recently established, rather than the

most recent growing seasons. We extracted the value of each environmental variable per site

based on the geographic locations of sampled sites and applied min-max normalization of

extracted values for each variable.

2.5. Statistical analysis

First, we used the filtered SNPs (see Materials and Methods, Genetic data) to estimate genetic

structure and relatedness of individuals. Genetic structure was calculated in *ADMIXTURE* v. 1.3

(Alexander et al. 2015) and the number of genetic clusters was set to 4 based on the smallest

cross-validation error (Supporting information Fig. **S6**). To derive the relationships among trees,

we calculated the SNPs-based pairwise distance between individuals using *plink* v. 1.9, and

applied the *cmdscale* function of *Matlab* (v. R2017b) to perform Principal coordinate analysis

(PCoA) and to derive the genetic Principal Coordinates (PCOs). Based on pairwise distances

between individuals, we additionally calculated within-site standard deviations of genetic

relatedness.

Furthermore, we estimated independent and common contributions of climatic and spectral

information to model genetic structure among sampled *F. sylvatica* sites. We used redundancy

analysis (RDA) as a multivariate ordination method to simultaneously account for multiple

response variables (Capblancq and Forester 2021). We used Akaike Information Criterion (AIC)

values for forward variable selection and a threshold of 2.5 for Variance Inflation Factor (VIF)

scores to reduce the impact of collinearity of variables (Johnston et al. 2018). We limited the

response variables to *F. sylvatica* distribution along the first and the second genetic PCO, which

explained together 7.7% of total variation in derived genetic structure (Extended Data Fig. **S7**).

Further genetic PCOs (3-6) each explained less then 2% in the derived genetic structure and did

not correlate strongly with environmental and spectral variation in our dataset (Extended Data

Fig. **S8**).

We constructed RDA models from environmental and spectral information separately, which

resulted in the construction of environment-based (including 4 linearly uncorrelated out of 19

explanatory environmental variables) and spectral-based (including 5 linearly uncorrelated out of

216 explanatory spectral variables) models. For the full model, we considered all 9 variables

selected for the environment- and spectral-based models to construct RDA models including

both environmental and spectral information as explanatory variables. Following the variable

selection procedures described above yielded a full model that included three environmental and

five spectral linearly uncorrelated explanatory variables. We derived the independent

contributions of each environmental and spectral variable to the full model and we partitioned the

genetic variance (as variance along genetic PCO1 and PCO2) to within- and between-site

variation using analysis of variance (ANOVA) with explanatory terms for environment and

spectral information and for site identity. Additionally, we conducted an analysis based on the

five first spectral Principal Components (PCs) derived from a PCA on the resampled spectral

data to compare how well these corresponded to variation along genetic PCOs, and found the

explanatory power to be lower than found in our main analysis using RDA (Supporting

information Tab. S1). We used the *rda* function of the *vegan* package (v. 2.6-4) in *R* (v. 4.1.3) to

construct the models and to find the model fits, and the *pca* function of *Matlab* (v. R2017b) to

derive the spectral PCs.

To determine the contribution of spectral information to genetic structure predictions, we

conducted leave-one-out validation. Accordingly, we created a model of 21 populations to

predict the distribution of individuals of the 22^{nd} population along genetic PCOs from (1)

environmental information only, and (2) environmental information merged with spectral

information. We present the contribution of spectral information to genetic structure prediction as

 the percentage change in predicted versus actual distribution along genetic PCO1 and PCO2

from the model based on (1) environmental and (2) environmental with spectral information,

respectively.

3. Results

3.1. Derived genetic structure of *F. sylvatica* across the species natural range

Fig. 1. *Genetic structure among 22* F. sylvatica *sites presented as kinship probabilities of*

belonging to K=4 genetic clusters with the smallest cross-validation error in ADMIXTURE

inferred from 1,380,310 SNPs across the nuclear genome (see Materials and Methods, Genetic

*data). (**a**) Averaged kinship probability and the standard deviation of genetic variation among the*

individuals from each sampled site (N=10 except for RSB, N=4, ESP, N=8, PLM and ITP, N=9).

*(**b**) The kinship probability among 209 sampled* F. sylvatica *individuals. (**c**) Relationship between*

F. sylvatica *individuals plotted along the first and second Principal Coordinates (PCOs) derived*

from genetic distances among individuals inferred from 1,380,310 SNPs across the nuclear

genome (see Materials and Methods, Genetic data). The horseshoe-like distribution of the

individuals along the first and second PCOs does not affect the conclusions of the presented

analysis (Podani and Miklós 2002). Individuals are colored by combined kinship probabilities to

four genetic clusters.

We present, to our knowledge, the first assessment of genetic structure of a broad-leaf tree

species across its natural range derived from whole-genome sequences of nuclear DNA. Using

ADMIXTURE cross-validation, we identified four genetic clusters representing the current *Fagus*

sylvatica L. genetic structure (Fig. **1**). We identified a high abundance of one genetic cluster in

southern Italy (ITE, ITM, ITP, ITV; cluster indicated in yellow), one genetic cluster in the southern

Pyrenees (ESP, ESM, FRM; cluster indicated in maroon), and one genetic cluster centered in

Corsica (FRV, FRC; cluster indicated in green). The genetic cluster centered in Corsica

(indicated in green) intermixes with the cluster along the French Mediterranean basin (FRB,

FRM) (indicated in maroon), with the genetic cluster along the Appenninan Peninsula (ITB, ITV,

ITF, ITP) (indicated in yellow), and with the fourth genetic cluster (indicated in blue) along the

Balkans (HRP, RSB, RSF, SIK) and the Carpathian mountain range (PLB, PLW). This fourth

cluster (indicated in blue) dominates our sampling across Europe and is the most widespread in

the central and northern parts of the continent (Fig. **1a**).

The sampled sites in central and northern Europe are also characterized by a relatively small

standard deviation of genetic distances within sites (Fig. **1b**). In contrast, per-individual kinship

probabilities to different genetic clusters, and pairwise genetic dissimilarities based on allele

frequencies between individuals, demonstrate relatively higher within and between-population

variation in southern as compared to northern samples (Fig. **1b,c**, Supporting information Fig.

S1).

3.2. Observed genetic structure explained by environmental and spectral

information

Together, environmental and spectral information explained 77.7% of the variance in the

distribution of sampled *F. sylvatica* along the first and second genetic PCOs (Fig. **2**).

Environmental variables explained 39.8% of the variation independently of spectral information.

The environmental variables contributing most were Mean Temperature of Driest Quarter (BIO

09), Isothermality (BIO 03), Precipitation Seasonality (BIO 15) and Precipitation of Driest Quarter

(BIO 17). In contrast, variables from leaf spectra explained 12.2% of variation in genetic

structure among trees (genetic PCO1 and PCO2) independently of environmental information.

The spectral variation explained 12.1% of genetic variation between sites and only 0.1% of

genetic variation within sites (Supporting information Fig. **S2**). Portions of spectra which were

most informative about the genetic structure correspond to variations of spectral phenotypes

centered at: 350 nm in the Ultraviolet (UV) part of spectra, 410 nm and 530 nm in the visible

(VIS) part of spectra, 790 nm in the near infrared (NIR) part of spectra, and 1500 nm in the

Shortwave Infrared (SWIR) part of spectra. Environmental and spectral variation combined

contributed an additional 25.6% to predicting genetic variation along genetic PCO1 and PCO2.

Fig. 2. *RDA models for prediction of* F. sylvatica *genetic variation as captured by the first*

(PCO1) and second (PCO2) genetic Principal Coordinate across the species range derived from

environmental information, spectral information, and environmental information combined with

spectral information. The variables were chosen based on AIC and VIF <2.5. Out of 19

environmental variables, four were identified to be indicative of the genetic structure between

sites, and out of 216 spectral bands, five were indicative of the genetic structure between and

within sites. The full model combined three out of the four selected environmental variables and

five out of the five spectral variables selected for the spectral-based model. The incorporation of

spectral information added 12.2% explanatory power to the model created based on

environmental information (with contribution of 39.8%) and spectral information, which together

explained 77.7% of the variation in genetic structure among sampled F. sylvatica *individuals*

(mean R^2 for PCO1 and PCO2). The F. sylvatica *individuals in RDA space are colored by*

combined kinship probabilities to four genetic clusters.

Spectral information contributed most to the prediction of the first genetic PCO (Fig. **3a**). The

inclusion of predictors derived from leaf spectral phenotypes improved the genetic structure

prediction by 59.7% along genetic PCO1 in comparison to the prediction based on

environmental variables only (R^2 of 0.79 versus 0.63) (Supporting information Fig. **S3**). The

spectral indicators contributing most to prediction of the distribution of individuals along this

genetic PCO1 relate to leaf interaction with spectral bands centered at 530 nm and 410 nm in

the visible (VIS) part of spectra. The contribution of spectral information to prediction of the

second genetic PCO produced a mean 7.9% improvement in comparison to the prediction based

on environmental variables only (R^2 of 0.76 versus 0.74) (Fig. **3b**, Supporting information Fig.

S3). The smaller contribution of spectral information to prediction of the second genetic PCO

might correspond to the shared explanatory power of spectral phenotypes centered at 790 nm

and 350 nm, which contribute most to the second genetic PCO axis of the full RDA model (Fig.

2).

331
332

Fig. 3. *Per site contribution of spectral information to predict* F. sylvatica *genetic variation as a*

*change (%) in predicted versus actual distribution along (**a**) the first genetic Principal Coordinate*

*(mean change of + 59.7 %) and (**b**) second genetic Principal Coordinate (mean change of + 7.9*

%). The change compares the prediction accuracy of RDA models based on (1) environmental

information only and (2) environmental and spectral information combined. The improvement (in

red) or decrease (in blue) in the genetic variation prediction represents the mean per-site change

derived from leave-one-out cross validation of a model based on 21 sites to predict the genetic

distribution of individuals of the 22nd site. The within-site variation in prediction accuracy is

*presented in Supporting information Fig. **S4**.*

4. Discussion

We present contributions of spectral phenotype data in combination with environmental data to

improve the prediction of intraspecific genetic structure derived from whole-genome sequences

of a forest tree species across its natural range. Spectral predictions of spatial patterns in

genetic variation may be further improved by incorporation of the species phylogeographic

history, and hold potential to be further guided by remotely acquired spectral information at a

landscape scale, repeatedly over time (Czyż et al. 2023).

4.1. Species phylogeographic history

While spectra carry information about current genetic composition, further incorporation of

information about the species phylogeographic history explaining development of genetic

patterns would constrain model predictions to the available gene pool of particular areas.

The genetic clusters indicated in yellow and maroon in Fig. **1** may correspond to previously

reported glacial refugia (Magri et al. 2006), while further genetic and phylogeographic studies

would be valuable in Corsica to determine whether the cluster indicated in green may represent

intermixing between refugia or an additional refugium (Petit et al. 2003). The origin of the most

widespread genetic cluster (indicated in blue) in central and northern Europe remains uncertain

based on our sampling and might be assigned to the eastern range of the Alps (Gomory et al.

2003) or southern Moravia (Magri et al. 2006). The phylogenetic structure across the range of *F.*

sylvatica is thought to result from post-glacial expansion: from the west across the central

European plains (cluster indicated in maroon), from the south across the Carpathian mountain

range (cluster indicated in yellow), and from Corsica and nearby sites at the French

Mediterranean basin (FRB, FRM) across the Appenninan Peninsula (ITB, ITV, ITF, ITP),

Balkans (HRP, RSB, RSF, SIK) and the Carpathian mountain range (PLB, PLW), with the last

likely permitted by terrestrial connections between the Adriatic and Mediterranean coasts during

the last glaciation (cluster indicated in green) (Hazler et al. 1997). These genetic patterns

correlate with environmental niches defined by water availability (BIO 09, BIO 15, BIO 17) and

isothermality (BIO 03), which may determine adaptive patterns in European beech (Fang and

Lechowicz 2006, Bolte et al. 2007), and correlate with gene flow. In contrast to lower mountain

ranges, the Alps likely act as a barrier for gene flow between southern and northern populations

(Paule 1995). Additionally, the continuous distribution of the species in the northern part of its

range likely allows greater gene flow across the landscape, while the patchier distribution in the

south limits gene flow and maintains genetic distance between populations. This may result in

higher genetic differences between southern sites, which may also be enhanced by post-glacial

climatic changes causing migration to higher altitudes in the southern part of the species range

and environmental pressures that drive the diversification of *F. sylvatica* populations at their

southern limit (Thiébaut et al. 1982). Merging the current genetic structure and the

phylogeographic history of the species with regularly monitored spectral (phenotypic) variation

could allow more accurate prediction of genetic structure within tree species across their range.

4.2.   Contribution of spectral phenotypes to predict genetic structure

We report spectral-genetic associations based on spectral phenotyping of leaves sampled at

selected sites across the natural range of *F. sylvatica*. These associations showed that the

spectral variation that contributed most to the predicton of genetic structure in this study

corresponds to: pigment absorption features related to electronic transitions induced by light at

410 nm and 530 nm (Gitelson et al. 2006), leaf water content related to vibrational absorption of

photons at 790 nm and 1500 nm, leaf surface morphological properties affecting the reflection of

UV light at 350 nm, and leaf inner structure affecting the scattering of electromagnetic radiation

at 790 nm (Gausman et al. 1971). Spectrally inferred variation in leaf inner structure with leaf

water content (790 nm) and variation in leaf surface micromorphology (350 nm) contributed most

to predicting the distribution of individuals across the first and the second genetic PCOs,

respectively.

Conducting genotype-phenotype association studies could be a next step towards identifying

genetic pathways and the mechanistic foundation of these predictions. Associations with genes

of known function may also be supported by growing seeds from these sites in a common

garden, which would help to separate genetic and environmental influences on spectral variation

including short-term microscale environmental conditions which we could not account for in our

analyses and which have a more direct influence on currently observed phenotypes, as opposed to shaping past survival, development and establishment, and composition of genotypes at a site. Such associations could facilitate research on selection processes that drive genetic adaptations across broad spatial and temporal scales.

4.3. Potential of remote observations to contribute to monitoring genetic variation

We have investigated the largest sampling to date of genomic and spectral variation among *F. sylvatica* individuals from across the species natural range, and by combining genetic structure derived from short-read whole-genome sequencing with variation in spectral phenotypes of leaves, we have shown the positive contribution of spectral information to predict genetic structure. This represents an advance on foundational studies on linking genetic with spectral variations conducted at small portions of a species range, comparing species, or comparing evolutionary trends across taxa (Czyż et al. 2020, Czyż et al. 2023, Li et al. 2023, Cavender-Bares et al. 2016, Seeley et al. 2023, Meireles et al. 2020). Further, our work indicates inferences of genetic variation which may become possible when upscaling the approach to remote observations of spectral phenotypes at a landscape scale, and presents potential to associate spectral variation within species with functional genes.

While we used whole-genome sequences representing the genetic composition of sampled individuals, the *in situ* leaf-level sampling necessarily was spatially sparse. Spectral phenotypes at canopy level acquired from remote observations of airborne and spaceborne imaging spectrometers will allow for spatially dense sampling over a large geographical area. Data from existing spaceborne imaging spectrometers, and the wealth of planned and upcoming imaging spectrometer missions (including NASA's SBG (Cawse-Nicholson et al. 2021) and ESA's CHIME (Celesti et al. 2022)), will bring spatially contiguous and temporally dense information about spectral properties of the Earth surface resolved at spectral intervals which are sufficiently

narrow to be interpreted in terms of molecular absorption features. The potential correlation between genomic and spectral variation acquired over large areas and at coarser spatial resolution with pixels each representing a plant community rather then single individual (Zheng et al. 2022, Helfenstein et al. 2022) still needs to be evaluated to confirm its potential contribution to infer intraspecific genetic structure across species ranges and predict genetic composition. The spectral-based modelling of genetic variation might be further applied across different phylogenetic groups and might facilitate spatial mapping of gene pools, beyond existing trait-based approaches.

Spectral support of predictions for genetic variation across space and time will help to target environmentally, phenotypically and genetically underrepresented sites or hotspots for measurement of genetic resources, which will further contribute to calibration of predictive models. Improved predictions of genetic variation using spectral phenotypes could indicate changes in genetic diversity and adaptation within tree species over time when combined with repeated monitoring and could deliver information for the adjustment of current efforts to safeguard biodiversity at the global scale, at unprecedented speed and spatial completeness.

Acknowledgments

This study is supported the University of Zurich Research Priority Program on Global Change and Biodiversity (URPP-GCB), and funded by the NOMIS grant on "Remotely Sensing Ecological Genomics". We express our gratitude to Emmanuelle Argenti, Ventura Gennaro, Hugot Laetitia, Elodie Magnanou, Joseph Garrigue, Kris Vandekerkhove, Arne Verstraeten, Nilsson Jorgen, Piotr Małż, Katarzyna Gierad, Dejan Stojanović and the Institute of Lowland Forestry and Environment; University of Novi Sad, Gordan Lukač, Katja Konečnik, Janez Šubic, Damen Kocjan, Eustaquio Gil Pelegrín, Juan Pedro Ferrio, Piotr Poloczek and Stanisław

446 Kucharzyk for allowing access to their sampling sites and support on investigation and to Romeo

447 Galiano, Joseph Garrigue, Zbigniew Stebel, Helgessons TrädTjänst AB company and Michał

448 Szydłowski for assistance in sample collection. We express our gratitude to Reinhard Furrer,

449 Anna K. Schweiger, Benjamin Dauphin, Katalin Csilléry and Ingmar Staude for helpful

450 discussions.

Declaration of Competing Interest

453 The authors declare that they have no known competing financial interests or personal

454 relationships that could have appeared to influence the work reported in this paper.

Author contributions

Conceptualization: EAC (lead), MES (lead), BS, MBE, MH, MCS. Methodology: EAC (lead),

MES, MCS, BS, MBE, AM, MH. Software: EAC (lead), BS, CL. Validation: EAC, BS, MBE.

Formal analysis: EAC (lead), BS, MBE. Investigation: EAC (lead), BS, MBE, MH, CL, MCS,

MES. Resources: MES (lead), MCS. Data Curation: EAC. Writing - Original Draft: EAC. Writing -

Review and Editing: EAC (lead), BS, MBE, MH, MCS, CL, MES. Visualization: EAC.

Supervision: MES, MCS, BS. Project administration: MES, MCS. Funding acquisition: MES.

Data availability

463 The raw short-read whole genome sequences genomic data that support the findings in this

464 study are deposited at NCBI Sequence Read Archive (SRA) under the project with submission

465 number SUB13301963. The leaf spectroscopic data are archived in the Specchio Spectral

466 Information System (https://specchio.ch/). All data will be publicly available upon publication.

Code availability

468   The bash codes developed for deriving genetic structure, and *R* and *Matlab* codes developed for

469   the statistical analysis are deposited at GitHub (*to be established upon the publication*) and will

470   be publicly available upon publication.

References

Alexander, D. H., Shringarpure, S. S., Novembre, J., & Lange, K. (2015). Admixture 1.3 software
manual. *Los Angeles: UCLA Human Genetics Software Distribution*.

Bolte, A., Czajkowski, T., & Kompa, T. (2007). The north-eastern distribution range of European
beech—a review. *Forestry*, *80*(4), 413-429.

Capblancq, T., & Forester, B. R. (2021). Redundancy analysis: A Swiss Army Knife for
landscape genomics. *Methods in Ecology and Evolution*, *12*(12), 2298-2309.

Capblancq, T., Morin, X., Gueguen, M., Renaud, J., Lobreaux, S., & Bazin, E. (2020).
Climate associated genetic variation in Fagus sylvatica and potential responses to climate
change in the French Alps. *Journal of Evolutionary Biology*, *33*(6), 783-796.

Cardinale, B. J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., ... & Naeem,
S. (2012). Biodiversity loss and its impact on humanity. *Nature*, *486*(7401), 59-67.

Caudullo, G., Welk, E., & San-Miguel-Ayanz, J. (2017). Chorological maps for the main
European woody species. *Data in brief*, *12*, 662-666.

Cavender-Bares, J., Meireles, J. E., Couture, J. J., Kaproth, M. A., Kingdon, C. C., Singh, A., ...
& Townsend, P. A. (2016). Associations of leaf spectra with genetic and phylogenetic variation in
oaks: prospects for remote detection of biodiversity. *Remote Sensing*, *8*(3), 221.

Cawse-Nicholson, K., Townsend, P. A., Schimel, D., Assiri, A. M., Blake, P. L., Buongiorno, M.
F., ... & SBG Algorithms Working Group. (2021). NASA's surface biology and geology
designated observable: A perspective on surface imaging algorithms. *Remote Sensing of*
Environment, *257*, 112349.

Celesti, M., Rast, M., Adams, J., Boccia, V., Gascon, F., Isola, C., & Nieke, J. (2022, July). The
Copernicus Hyperspectral Imaging Mission for the Environment (Chime): Status and Planning. In
IGARSS 2022-2022 IEEE International Geoscience and Remote Sensing Symposium (pp. 5011-
5014). IEEE.

Csilléry, K., Lalagüe, H., Vendramin, G. G., González Martínez, S. C., Fady, B., &
Oddou Muratorio, S. (2014). Detecting short spatial scale local adaptation and epistatic
selection in climate related candidate genes in E uropean beech (F agus sylvatica) populations.
Molecular Ecology, *23*(19), 4696-4708.

Cuervo-Alarcon, L., Arend, M., Müller, M., Sperisen, C., Finkeldey, R., & Krutovsky, K. V. (2021).
A candidate gene association analysis identifies SNPs potentially involved in drought tolerance
in European beech (Fagus sylvatica L.). *Scientific Reports*, *11*(1), 2386.

Czyż, E. A., Guillén Escribà, C., Wulf, H., Tedder, A., Schuman, M. C., Schneider, F. D., &
Schaepman, M. E. (2020). Intraspecific genetic variation of a Fagus sylvatica population in a
temperate forest derived from airborne imaging spectroscopy time series. *Ecology and evolution*,
10(14), 7419-7430.

Czyż, E. A., Schmid, B., Hueni, A., Eppinga, M. B., Schuman, M. C., Schneider, F. D., ... &
Schaepman, M. E. (2023). Genetic constraints on temporal variation of airborne reflectance
spectra and their uncertainties over a temperate forest. *Remote Sensing of Environment*, *284*,
113338.

Des Roches, S., Post, D. M., Turley, N. E., Bailey, J. K., Hendry, A. P., Kinnison, M. T., ... &
Palkovacs, E. P. (2018). The ecological importance of intraspecific variation. *Nature ecology &*
evolution, *2*(1), 57-64.

Exposito-Alonso, M., Booker, T. R., Czech, L., Gillespie, L., Hateley, S., Kyriazis, C. C., ... &
Zess, E. (2022). Genetic diversity loss in the Anthropocene. *Science*, *377*(6613), 1431-1435.

Fang, J., & Lechowicz, M. J. (2006). Climatic limits for the present distribution of beech (Fagus
L.) species in the world. *Journal of Biogeography*, *33*(10), 1804-1819.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1□km spatial resolution climate surfaces
for global land areas. *International journal of climatology*, *37*(12), 4302-4315.

Fridley, J. D., & Grime, J. P. (2010). Community and ecosystem effects of intraspecific genetic
diversity in grassland microcosms of varying species diversity. *Ecology*, *91*(8), 2272-2283.

Gallois, A., Audran, J. C., & Burrus, M. (1998). Assessment of genetic relationships and
population discrimination among Fagus sylvatica L. by RAPD. *Theoretical and applied genetics*,
97, 211-219.

Gausman, H. W., Allen, W. A., Escobar, D. E., Rodriguez, R. R., & Cardenas, R. (1971). Age
Effects of Cotton Leaves on Light Reflectance, Transmittance, and Absorptance and on Water
Content and Thickness 1. *Agronomy Journal*, *63*(3), 465-469.

Gitelson, A. A., Keydan, G. P., & Merzlyak, M. N. (2006). Three□band model for noninvasive
estimation of chlorophyll, carotenoids, and anthocyanin contents in higher plant leaves.
Geophysical research letters, *33*(11).

Gomory, D., Paule, L., Shvadchak, I. M., Popescu, F., Sulkowska, M., Hynek, V., & Longauer, R.
(2003). Spatial patterns of the genetic differentiation in European beech (Fagus sylvatica L.) at
allozyme loci in the Carpathians and the adjacent regions. *Silvae Genetica*, *52*(2), 78-83.

Guevara, M. Á., Sánchez-Gómez, D., Vélez, M. D., de María, N., Díaz, L. M., Ramírez-Valiente,
J. A., ... & Cervera, M. T. (2022). Epigenetic and Genetic Variability in Contrasting Latitudinal
Fagus sylvatica L. Provenances. *Forests*, *13*(12), 1971.

Guyot, G., Baret, F., & Jacquemoud, S. (1992). *Imaging spectroscopy for vegetation studies*
(Vol. 2, pp. 145-165). Kluwer Academic Publishers: Norwell, MA, USA.

Hazler, K., Comps, B., Sugar, I., Melovski, L., Tashev, A., & Gracan, J. (1997). Genetic structure
of Fagus sylvatica L. populations in Southeastern Europe. *Silvae Genetica*, *46*(4), 229-235.

Hector, A., & Bagchi, R. (2007). Biodiversity and ecosystem multifunctionality. *Nature*,
448(7150), 188-190.

Hector, A., Hautier, Y., Saner, P., Wacker, L., Bagchi, R., Joshi, J., ... & Loreau, M. (2010).
General stabilizing effects of plant diversity on grassland productivity through population
asynchrony and overyielding. *Ecology*, *91*(8), 2213-2220.

Helfenstein, I. S., Schneider, F. D., Schaepman, M. E., & Morsdorf, F. (2022). Assessing
biodiversity from space: Impact of spatial and spectral resolution on trait-based functional
diversity. *Remote Sensing of Environment*, *275*, 113024.

Hoban, S., Archer, F. I., Bertola, L. D., Bragg, J. G., Breed, M. F., Bruford, M. W., ... & Hunter,
M. E. (2022). Global genetic diversity status and trends: towards a suite of Essential Biodiversity
Variables (EBVs) for genetic composition. *Biological Reviews*, *97*(4), 1511-1538.

Hughes, A. R., Inouye, B. D., Johnson, M. T., Underwood, N., & Vellend, M. (2008). Ecological
consequences of genetic diversity. *Ecology letters*, *11*(6), 609-623.

IPBES Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services
(IPBES). (2019). Global assessment report on biodiversity and ecosystem services of the
Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.

Isbell, F., Calcagno, V., Hector, A., Connolly, J., Harpole, W. S., Reich, P. B., ... & Loreau, M.
(2011). High plant diversity is needed to maintain ecosystem services. *Nature*, *477*(7363), 199-
202.

Isbell, F., Craven, D., Connolly, J., Loreau, M., Schmid, B., Beierkuhnlein, C., ... & Eisenhauer,
N. (2015). Biodiversity increases the resistance of ecosystem productivity to climate extremes.
Nature, *526*(7574), 574-577.

Jacquemoud, S., & Ustin, S. (2019). *Leaf optical properties*. Cambridge University Press.

Jenkins, C. N., & Joppa, L. (2009). Expansion of the global terrestrial protected area system.
Biological conservation, *142*(10), 2166-2174.

Johnson, C. N., Balmford, A., Brook, B. W., Buettel, J. C., Galetti, M., Guangchun, L., &
Wilmshurst, J. M. (2017). Biodiversity losses and conservation responses in the Anthropocene.
Science, *356*(6335), 270-275.

Johnston, R., Jones, K., & Manley, D. (2018). Confounding and collinearity in regression
analysis: a cautionary tale and an alternative procedure, illustrated by studies of British voting
behaviour. *Quality & quantity*, *52*, 1957-1976.

Li, C., Czyż, E. A., Halitschke, R., Baldwin, I. T., Schaepman, M. E., & Schuman, M. C. (2023).
Evaluating potential of leaf reflectance spectra to monitor plant genetic variation. *Plant Methods*,
19(1), 108.

Lou, R. N., Jacobs, A., Wilder, A. P., & Therkildsen, N. O. (2021). A beginner's guide to
low☐coverage whole genome sequencing for population genomics. *Molecular Ecology*, *30*(23),
5966-5993.

Madritch, M. D., Kingdon, C. C., Singh, A., Mock, K. E., Lindroth, R. L., & Townsend, P. A.
(2014). Imaging spectroscopy links aspen genotype with below-ground processes at landscape
scales. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *369*(1643),
20130194.

Magri, D., Vendramin, G. G., Comps, B., Dupanloup, I., Geburek, T., Gömöry, D., ... & De
Beaulieu, J. L. (2006). A new scenario for the Quaternary history of European beech
populations: palaeobotanical evidence and genetic consequences. *New phytologist*, *171*(1), 199-
221.

Meger, J., Ulaszewski, B., & Burczyk, J. (2021). Genomic signatures of natural selection at
phenology-related genes in a widely distributed tree species Fagus sylvatica L. *BMC genomics*,
22(1), 1-20.

Meireles, J. E., Cavender☐Bares, J., Townsend, P. A., Ustin, S., Gamon, J. A., Schweiger, A.
K., ... & O'Meara, B. C. (2020). Leaf reflectance spectra capture the evolutionary history of seed
plants. *New Phytologist*, *228*(2), 485-493.

Miller, J. R., Steven, M. D., & Demetriades-Shah, T. H. (1992). Reflection of layered bean leaves
over different soil backgrounds: measured and simulated spectra. *International Journal of
Remote Sensing*, *13*(17), 3273-3286.

Mishra, B., Ulaszewski, B., Meger, J., Aury, J. M., Bodénès, C., Lesur-Kupin, I., ... & Thines, M.
(2022). A Chromosome-level genome assembly of the European Beech (Fagus sylvatica)
reveals anomalies for organelle DNA integration, repeat Content and distribution of SNPs.
Frontiers in genetics, *12*, 691058.

Mora, C., Andréfouët, S., Costello, M. J., Kranenburg, C., Rollo, A., Veron, J., ... & Myers, R. A.
(2006). Coral reefs and the global network of marine protected areas. *Science*, *312*(5781), 1750-
1751.

Perrings, C., Duraiappah, A., Larigauderie, A., & Mooney, H. (2011). The biodiversity and
ecosystem services science-policy interface. *Science*, *331*(6021), 1139-1140.

Petit, R. J., Aguinagalde, I., de Beaulieu, J. L., Bittkau, C., Brewer, S., Cheddadi, R., ... &
Vendramin, G. G. (2003). Glacial refugia: hotspots but not melting pots of genetic diversity.
science, *300*(5625), 1563-1565.

Pfenninger, M., Reuss, F., Kiebler, A., Schönnenbeck, P., Caliendo, C., Gerber, S., ... &
Feldmeyer, B. (2020). Genomic basis of drought resistance in Fagus sylvatica. *bioRxiv*, 2020-12.

Pluess, A. R., & Weber, P. (2012). Drought-adaptation potential in Fagus sylvatica: linking
moisture availability with genetic diversity and dendrochronology. *PloS one*, *7*(3), e33636.

Podani, J., & Miklós, I. (2002). Resemblance coefficients and the horseshoe effect in principal
coordinates analysis. *Ecology*, *83*(12), 3331-3343.

Purvis, A., & Hector, A. (2000). Getting the measure of biodiversity. *Nature*, *405*(6783), 212-219.

Reich, P. B., Tilman, D., Isbell, F., Mueller, K., Hobbie, S. E., Flynn, D. F., & Eisenhauer, N.
(2012). Impacts of biodiversity loss escalate through time as redundancy fades. *Science*,
336(6081), 589-592.

Schaepman, M. E., Jehle, M., Hueni, A., D'Odorico, P., Damm, A., Weyermann, J., ... & Itten, K.
I. (2015). Advanced radiometry measurements and Earth science applications with the Airborne
Prism Experiment (APEX). *Remote Sensing of Environment*, *158*, 207-219.

Schaepman, M. E., Ustin, S. L., Plaza, A. J., Painter, T. H., Verrelst, J., & Liang, S. (2009). Earth
system science related imaging spectroscopy—An assessment. *Remote Sensing of*
Environment, *113*, S123-S137.

Scholes, R. J., Mace, G. M., Turner, W., Geller, G. N., Jürgens, N., Larigauderie, A., ... &
Mooney, H. A. (2008). Toward a global biodiversity observing system. *Science*, *321*(5892),
1044-1045.

Scott, J. M., Csuti, B., Estes, J. E., & Anderson, H. (1989). Status assessment of biodiversity
protection. *Conservation Biology*, *3*(1), 85-87.

Seeley, M. M., Martin, R. E., Vaughn, N. R., Thompson, D. R., Dai, J., & Asner, G. P. (2023).
Quantifying the variation in reflectance spectra of Metrosideros polymorpha canopies across
environmental gradients. *Remote Sensing*, *15*(6), 1614.

Stefanini, C., Csilléry, K., Ulaszewski, B., Burczyk, J., Schaepman, M. E., & Schuman, M. C.
(2023). A novel synthesis of two decades of microsatellite studies on European beech reveals
decreasing genetic diversity from glacial refugia. *Tree Genetics & Genomes*, *19*(1), 3.

Sutherland, W. J., Pullin, A. S., Dolman, P. M., & Knight, T. M. (2004). The need for evidence-
based conservation. *Trends in ecology & evolution*, *19*(6), 305-308.

Tang, T., Zhang, N., Bongers, F. J., Staab, M., Schuldt, A., Fornoff, F., ... & Liu, X. (2022). Tree
species and genetic diversity increase productivity via functional diversity and trophic feedbacks.
Elife, *11*, e78703.

Thiébaut, B., Lumaret, R., & Vernet, P. H. (1982). The bud enzymes of beech (Fagus sylvatica
L.) genetic distinction and analysis of polymorphism in several French populations.

Tilman, D., & Downing, J. A. (1994). Biodiversity and stability in grasslands. *Nature*, *367*(6461),
363-365.

Turner, W. (2014). Sensing biodiversity. *Science*, *346*(6207), 301-302.

van Moorsel, S. J., Hahl, T., Petchey, O. L., Ebeling, A., Eisenhauer, N., Schmid, B., & Wagg, C.
(2021). Co□occurrence history increases ecosystem stability and resilience in experimental
plant communities. *Ecology*, *102*(1), e03205.

Wilson, E. O. (1988). Biodiversity.

Yamasaki, E., Altermatt, F., Cavender-Bares, J., Schuman, M. C., Zuppinger-Dingley, D.,
Garonna, I., ... & Shimizu, K. K. (2017). Genomics meets remote sensing in global change
studies: monitoring and predicting phenology, evolution and biodiversity. *Current opinion in*
environmental sustainability, *29*, 177-186.

653 Zheng, Z., Zeng, Y., Schuman, M. C., Jiang, H., Schmid, B., Schaepman, M. E., & Morsdorf, F.
654 (2022). Individual tree-based vs pixel-based approaches to mapping forest functional traits and
655 diversity by remote sensing. *International Journal of Applied Earth Observation and*
656 *Geoinformation*, *114*, 103074.