# High-throughput identification of functional regulatory SNPs in systemic lupus erythematosus

Qiang Wang[1]*, Taehyeung Kim[1]*, Marta Martínez-Bonet[2,3], Sangwan Sim[1], Vitor Aguiar[1,4], Jing Cui[2], Jeffrey A. Sparks[2], Xiaoting Chen[4], Marc Todd[1], Brian Wauford[1], Matthew T. Weirauch[5,6,7], Maria Gutierrez-Arcelus[1,4], Peter A. Nigrovic[1,2]

1. Division of Immunology, Boston Children's Hospital, Harvard Medical School, Boston MA, USA

2. Division of Rheumatology, Inflammation, and Immunity, Brigham and Women's Hospital, Harvard Medical School, Boston MA, USA

3. Laboratory of Immune-regulation, Instituto de Investigación Sanitaria Gregorio Marañón, Madrid Spain

4. Broad Institute of MIT and Harvard, Cambridge, MA, USA

5. Center of Autoimmune Genomics and Etiology, Division of Human Genetics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA

6. Divisions of Human Genetics, Biomedical Informatics, and Developmental Biology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

7. Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, Ohio, USA

* Co-first authors.

**Correspondence:**
Peter A. Nigrovic, MD
Division of Immunology
Boston Children's Hospital
Karp Family Research Building 10211
One Blackfan Circle
Boston, MA 02115
peter.nigrovic@childrens.harvard.edu

**Conflict of interest:** The authors declare no relevant competing interests.

## Abstract

Genome-wide association studies implicate multiple loci in risk for systemic lupus erythematosus (SLE), but few contain exonic variants, rendering systematic identification of non-coding variants essential to decoding SLE genetics. We utilized SNP-seq and bioinformatic enrichment to interrogate 2180 single-nucleotide polymorphisms (SNPs) from 87 SLE risk loci for potential binding of transcription factors and related proteins from B cells. 52 SNPs that passed initial screening were tested by electrophoretic mobility shift and luciferase reporter assays. To validate the approach, we studied rs2297550 in detail, finding that the risk allele enhanced binding to the transcription factor Ikaros (IKZF1), thereby modulating expression of *IKBKE*. Correspondingly, primary cells from genotyped healthy donors bearing the risk allele expressed higher levels of the interferon / NF-κB regulator IKKε. Together, these findings define a set of likely functional non-coding lupus risk variants and identify a new regulatory pathway involving rs2297550, Ikaros, and IKKε implicated by human genetics in risk for SLE.

## Introduction

Systemic lupus erythematosus (SLE) is a chronic autoimmune disease predominantly affecting women. Identical twin concordance rates range between 30–50%, and first-degree relatives exhibit an approximately 20-fold increased risk of SLE compared with the general population [1-3]. While monogenic SLE has proven extremely informative with respect to causative pathways, most SLE is polygenic, arising through a complex interplay of genetic and environmental factors [4, 5]. Correspondingly, genome-wide association studies (GWAS) have defined dozens of loci that modulate disease risk [6-10]. Deciphering these different genetic contributors, each individually minor but collectively of major importance, will be central to defining SLE heterogeneity at a genetic level.

Unfortunately, translating GWAS data into pathogenic understanding has proven difficult. Few common disease-associated variants affect coding regions, indicating that the associated variants – predominantly single-nucleotide polymorphisms (SNPs) – likely exercise a regulatory role. Distinguishing functional non-coding SNPs from the many incidental SNPs that reside in close linkage disequilibrium (LD) remains a "needle in the haystack" problem of great complexity. Strategies to address this problem in polygenic diseases include fine-mapping [11], proteome-wide profiling of DNA-binding proteins over SNPs [12], integration of expression quantitative trait loci (eQTL) with epigenetic and/or transcriptomic data [13-15], massively parallel reporter assays [16-18], and SNP-seq [19].

Here, we pursue a combined experimental and bioinformatic approach to identify functional non-coding variants in SLE that act by regulating the binding of transcription factors and other DNA-binding proteins. We focus on B cells as a lineage implicated genetically as critical to SLE pathogenesis [20]. We used SNP-seq to screen for alleles that confer differential binding to B cell nuclear proteins, enriching targets further through epigenetic markers of open chromatin and predictors of regulatory function. Beginning with 2180 candidates, we identify 52 plausible regulatory SNPs, each interrogated experimentally to prioritize 5 high-likelihood functional variants. To test our approach, we examined

rs2297550 in detail, finding that this variant modulates the binding of the transcription factor Ikaros to regulate expression of the noncanonical IκB kinase IKKε. Thus, our strategy bridges the gap between GWAS and mechanism while defining a novel regulatory pathway implicated by population genetics in risk for SLE.

## Results

### Candidate functional non-coding variant identification via SNP-seq and bioinformatic screens

To identify functional non-coding variants associated with SLE, we began from GWAS hits with genome-wide significance (P value < 5 x 10[-8]) from five studies [7, 9, 10, 21, 22], finding 87 loci of interest, each defined by a tagging SNP (**Supplementary Table 1**). We identified all common variants in LD with these SNPs from 1000 Genomes Phase 3 (European population), filtering on minor allele frequency [MAF] >1%, LD R$^2$>0.8, and location within 1 Mb of the tagging SNP (**Figure 1A**). This strategy yielded 2180 candidate SNPs for downstream screening and validation (**Supplementary Table 2**).

To narrow this pool, we applied SNP-seq [23]. In this method, a library of double-stranded DNA constructs is generated that includes each allele of each candidate SNP. A 31bp sequence centered on each SNP is flanked by binding sites for the type IIS restriction enzyme BpmI that cleaves DNA in a sequence-independent manner 16bp (5')/14bp (3') adjacent to its binding site. Binding of transcription factors or other nuclear proteins to the SNP site protects the construct from enzymatic digestion (**Figure 1B**), as recognized by subsequent amplification and DNA sequencing [23, 24]. Here, our library contained 4363 dsDNA constructs reflecting each allele of the 2180 SNPs, including 3 with 3 common alleles (**Supplementary Table 2**). The library was then incubated with nuclear extract from either human peripheral blood mononuclear cells (PBMCs) from healthy individuals or BL2 B cells, washed, and then digested with BpmI. As a control, the construct pool was incubated without nuclear extract. The remaining sequences were amplified by PCR and used as the library for the next round of selection. The whole procedure was repeated for 5 cycles, with barcoding at cycles 3 and 5 to identify enriched SNPs by next-generation sequencing (NGS) (**Figure 1C**). The entire procedure was performed in three independent biological replicates, finding high correlation (Spearman ρ>0.8) between sample pairs (**Supplementary Figure 1A, 1B**) and between PBMC and BL2 samples (**Figure 1D**).

Two analytical approaches were employed to identify SNPs offering allele-specific protection, as established previously [23] . First, we chose SNPs that exhibited at least 20% difference in protection between alleles in cycle 5 in both samples treated with nuclear extract from both PBMC and BL2 cells; 452 SNPs exhibited such an allelic differential. Second, we sought SNPs demonstrating progressive allele-specific protection across cycles (cycle 0, 3, and 5) (**Supplementary Figure 1C**); 496 SNPs exhibited this pattern. In the end, 248 SNPs met both criteria (**Figure 1E, Supplementary Table 3**). A diagram of our analytical strategy is shown in **Supplementary Figure 2**.

Next, we examined the same 2180 SNPs using two bioinformatic approaches. First, we employed informative chromatin marks to identify areas of open chromatin, calculating H3K4me3 chromatin scores from 5 chromatin immunoprecipitation (ChIP)-seq studies in human B cells in the Encyclopedia of

DNA Elements (ENCODE) database, as per a described method [25] (**Supplementary Figure 3, Supplementary Table 4**). We selected for further study those SNPs that ranked in the top 100 from at least half of studies, yielding 91 variants (**Supplementary Table 4**). Second, we used Combined Annotation-Dependent Depletion (CADD), a composite measure that employs 60 genomic features to estimate the probability that a variant is functional [26]. We ranked all 2180 SNPs as per the CADD database (https://cadd.gs.washington.edu/), retaining the 25 SNPs that scored in the top 0.5% of all ~8.6 billion single nucleotide variants (**Supplementary Table 5**).

To reduce the number of SNPs to a tractable number for experimental validation, we chose the top 20 SNPs identified uniquely by SNP-seq, the top 8 SNPs identified uniquely by epigenetic and CADD analysis, and 18 SNPs identified in at least two of three analyses (6 SNP-seq+H3K4me3, 4 SNP-seq+CADD, 8 H3K4me3+CADD), for a total of 52 plausible functional SNPs (**Figure 2**).

## Experimental validation of plausible functional SNPs in SLE

As experimental validation of the 52-SNP enriched variant pool, we first employed the electrophoretic mobility shift assay (EMSA). Biotinylated 31bp DNA fragments centered on each allele of each the 52 candidate SNPs (111 alleles) were interrogated using nuclear extract from BL2 cells (triplicate biological repeats) followed by a validation EMSA using nuclear extract from human PBMCs. Overall, 7 biallelic and 2 triallelic (rs2297550 and rs936394) SNPs exhibited allele-imbalanced binding for all nuclear extracts: rs2297550, rs906868, rs936394, rs7302634, rs276461, rs13213604, rs378056, rs2814955 and rs9907966 (**Figure 3A**, shift binding bands marked as red circles). All EMSA blots for every experiment and a summary table can be found in **Supplementary Figure 4 and 5.**

Next, we assessed the allele-specific gene regulatory capacity of these 9 SNPs using a luciferase reporter assay; for rs2297550 and rs936394, we included only two alleles, since the third for each has a MAF <0.1% and is thus unlikely to mediate GWAS signal. We cloned the same risk or non-risk allele DNA fragments as for EMSA (but without biotinylation) into the pGL3 vector, and then transfected each construct into Daudi B cells, simpler to transfect than BL2, together with control vector pRL to normalize transfection efficiency. We found 5 SNPs to retain a significant difference between alleles and thus represent high-probability functional non-coding variants: rs2297550, rs906868, rs936394, rs9907966 and rs13213604 (**Figure 3B**). These SNPs are diagrammed in **Figure 3C**.

## Identification of potential transcription factor binding

To provide independent validation of the identified SNPs, we sought to confirm binding of transcription factors and/or other nuclear proteins in an allele-determined manner. Again, we approached this problem both experimentally and bioinformatically.

First, we employed pulldown and silver staining in SNPs that had exhibited allele-specific differential binding by EMSA, excluding rs2814955 because of its low luciferase activity, again with 31bp oligonucleotides for each allele but now using nuclear extract from Daudi cells [27]. In two biological repeats, four high-probability functional variants again showed differential binding (red circles) (**Supplementary Figure 6**). Each pulldown product for each allele of the five high-probability SNPs was

evaluated by mass spectrometry (MS) to identify binding nuclear proteins, yielding a set of candidate proteins for each (**Supplementary Table 6**). Second, we employed a bioinformatic algorithm based on published ChIP-seq data to identify candidate transcription factors likely to recognize each of the 5 high-probability variants [28, 29]. Bioinformatic predictions are shown in **Supplementary Table 7**. Evaluating these data together, we observed that the transcription factor Ikaros (IKZF1) was identified in the pulldown product for rs2297550 and was predicted by the algorithm, including preferential binding to the SLE-risk G allele as observed by EMSA. We therefore chose rs2297550 as a test case for our selection process.

### rs2297550 regulates *IKBKE*/IKKε expression

As depicted in **Figure 3C**, rs2297550 resides in a promoter/enhancer region, 47bp upstream of *IKBKE*, encoding the kinase IKKε [30-32]. This SNP emerged from Chinese and European GWAS data as a leading candidate in SLE, with *IKBKE* as its likely target gene [7, 33]. To assess the role rs2297550 in regulation of *IKBKE*, we used CRISPR homology directed repair to generate base-edited Daudi cell clones with CC, CG, and GG genotypes. GG clones exhibited lower mRNA for *IKBKE* (**Figure 4A**) and lower expression of IKKε protein (**Figure 4B**), confirming a regulatory relationship between SNP and gene.

Next, we recruited healthy donors bearing different genotype of rs2297550 to test the impact of genotype in primary PBMCs, using the Joint Biology Consortium (JBC) patient recruitment infrastructure and the Mass General Brigham Biobank (www.jbcwebportal.org). In resting cells, IKKε protein levels were comparable across genotypes in PBMCs collectively and in all immune subtypes tested (CD19+ B cells, CD3+CD4+ T cells, CD3+CD8+ T cells, CD3-CD56+ NK cells, CD14+ monocytes) (**Figure 5A-F**). However, after 4h of LPS stimulation, cells from GG donors exhibited differential protein levels in PBMCs generally as well as in B cells, CD8+ T cells, and NK cells; a heterozygote (allele dose) effect was observed in B cells and CD8+ T cells (**Figure 5G-L**). Interestingly, unlike in Daudi cells, the effect of the risk allele was to increase rather than decrease IKKε expression.

### Ikaros binds to rs2297550 to regulate *IKBKE*

Given concordant findings from MS and bioinformatic prediction, we pursued the possibility that Ikaros binds to rs2297550 to regulate *IKBKE*. We therefore performed an oligonucleotide pulldown assay using nuclear extract from Daudi cells, with or without excess non-biotinylated competitor, followed by western blot using anti-Ikaros antibody. Ikaros bound both C and G alleles but exhibited a preference for the G allele, as predicted, reflected in the disappearance signal with excess non-biotinylated G competitor (**Figure 6A**). To confirm this result, we employed EMSA-supershift, finding that anti-Ikaros weakened binding to rs2297550-G compared with isotype control or antibody to another transcription factor candidate from the bioinformatic prediction, E2F transcription factor 4 (E2F4) (**Figure 6B**). These studies establish that Ikaros binds rs2297550 and that this interaction is allele specific.

To assess rs2297550-Ikaros binding in living cells, we performed chromatin immunoprecipitation (ChIP)-qPCR. In Daudi cells, anti-Ikaros antibody precipitated rs2297550 significantly more than did isotype

control antibody (**Figure 6C**). Further, in CRISPR-HDR edited CG Daudi cells, we observed preferential pulldown of the G allele over the C allele (**Figure 6D**), again confirming its allelic preference.

To test whether the interaction of Ikaros with rs2297550 modulated IKKε expression, we generated three Ikaros knockout clones using CRISPR-Cas9. All clones showed Ikaros deficiency (**Figure 7A**) and corresponding loss of the ability of anti-Ikaros to pull down rs2297550 (**Figure 7B**). Consistent with the inhibitory effect of G allele carriage in these cells, loss of Ikaros increased *IKBKE* mRNA (**Figure 7C**) and IKKε protein (**Figure 7D**). These findings establish a new rs2297550-Ikaros-IKKε regulatory axis implicated by human population genetics in risk for SLE.

## Discussion

Genetic risk for SLE is borne by many susceptibility loci, most of which reflect common non-coding variants [34]. Despite the modest effect of particular variants on disease risk, each unambiguously implicates its associated pathway in the pathogenesis of human disease. Further, as the number of risk variants accumulates, incidence of SLE increases and age of onset declines, suggesting a "cumulative hit" model whereby variants operate synergistically to generate disease susceptibility [9]. Understanding which specific variants drive risk and how they do so offers a unique window into disease biology as well as into general mechanisms of immune control [35, 36].

Here, we conducted a systematic experimental and bioinformatic screen across 87 loci implicated by GWAS in SLE, identifying 52 plausible casual non-coding SNPs, of which 5 emerged through EMSA and luciferase as high-likelihood variants. Extrapolation from surrounding genes suggests that these 5 variants likely interface with functional pathways relevant for SLE biology, including the interferon pathway (*IKBKE*), pyroptosis (*IKBKE*), B cell differentiation (*IKZF3*) and WNT signaling (*LBH*) (**Figure 3C**).

To test our approach, we studied one variant in detail: rs2297550, a triallelic SNP (frequency of the three alleles C 0.862, G 0.138, T <0.001 per gnomAD) that resides 4462 bp upstream of the start codon for *IKBKE*, encoding the kinase IKKε. The minor allele G confers an increased risk of SLE with an odds ratio of 1.16 (95% CI: 1.11-1.21, $p=1.31 \times 10^{-11}$) compared with the major C allele [7]. Base editing showed that allelic variation at rs2297550 changes expression of *IKBKE* / IKKε. Variation in IKKε with genotype at rs2297550 was confirmed in healthy volunteer donors. As an orthogonal line of evidence, we showed that Ikaros exhibited allele-dependent affinity for rs2297550 and that deletion of Ikaros altered *IKBKE*/IKKε expression. These results establish a previously undefined rs2297550-Ikaros-IKKε regulatory axis.

Some groups have found that the risk allele G at rs2297550 is associated with lower *IKBKE* mRNA in unstimulated PBMCs [32, 37]. IKKε protein was not tested in these studies, raising the possibility that transcript and protein levels could be discordant. Indeed, we observed a trend toward lower levels of *IKBKE* mRNA in GG PBMC cells, though this effect did not reach significance (**Supplementary Figure 7**). The basis for transcript-protein divergence is not clear but could relate to the relative half-lives of mRNA and protein, among other factors [38, 39].

Interestingly, the relationship between genotype and IKKε varied with context. In Daudi B lymphoblasts, a line derived from a patient with Burkitt lymphoma, introduction of the risk GG genotype decreased IKKε expression and deletion of Ikaros enhanced it, defining Ikaros as an inhibitor of *IKBKE*. By contrast, in healthy donors, circulating B cells showed an *increase* in IKKε with risk allele carriage, including a gene dose effect, but only after LPS stimulation. Similar regulation was observed in CD8+ T cells, NK cells, and monocytes. The dependence of the impact of genetic variants on cell type and developmental/activation state is well recognized [40]. Since Ikaros can toggle between activator and repressor depending on associated cofactors, it may be that the variation observed here reflects differential expression of these proteins, or on other genetic/epigenetic differences between the lineages studied [41]. Under conditions other than those tested here, it is possible that the risk allele at rs2297550 could suppress IKKε in primary cells, as it does in Daudi cells.

An important intracellular intermediate in innate immune responses, IKKε can activate IRF3, IRF7, and NF-κB pathways to drive gene transcription in response to extracellular and intracellular stimuli, resulting in production of inflammatory mediators including type I interferons (**Figure 8**) [42-45]. The known role of interferons in SLE makes this pathway an appealing potential explanation for why rs2297550 is a risk SNP for SLE [46]. IKKε can also protect cells from TNF-induced apoptosis, plausibly thereby attenuating autoantibody production [47, 48]. *IKBKE* mRNA expression has been reported to be either higher [49, 50] or lower [32, 51, 52] in whole-blood and/or PBMC of SLE patients compared with healthy donors. Our primary cell data suggest that the "risk phenotype" attributable to rs2297550 likely reflects enhanced rather than suppressed expression of IKKε, though further work will be required to define the mechanistic connection between IKKε and SLE.

*IKZF1* encodes Ikaros, an essential regulator of lymphocyte development [41, 53]. Ikaros is implicated in SLE by multiple lines of evidence. A region including *IKZF1* carries genetic risk for SLE [33, 54-58]. Individuals bearing mutant *IKZF1* can present with SLE-like autoimmune disease [59, 60]. Clinical efficacy has been established in human SLE for iberdomide, an agent that accelerates degradation of Ikaros as well of the related transcription factor Aiolos [61, 62]. Mechanistically, Ikaros is involved in B cell tolerance and plasmacytoid dendritic cell differentiation [41, 63-65]. In B cells from SLE patients, iberdomide attenuated in vitro differentiation to plasmablasts and plasma cells as well as antibody production [66]. Thus, our findings implicating Ikaros in SLE risk via rs2297550 are consistent with genetic, clinical, and mechanistic observations in SLE.

Our study establishes a new connection between Ikaros, IKKε, and SLE, showing that the regulation of this intracellular signaling molecule is part of the portfolio of this transcription factor, and implicating both proteins in risk of incident SLE in humans. This finding has potential therapeutic relevance, since drug candidates supported by human genetic evidence are much more likely to succeed in clinical trials than those without such support [67, 68]. However, our data illustrate the challenge of extrapolating from a genetic variant to a single mechanism. Depending on cellular context, we find that the same variant has opposite effects on gene expression. Further, while we began our search in B cells, in human primary cells the rs2297550-IKKε effect spans multiple immune lineages, both adaptive and innate. Thus

our data do not establish which lineages, and which pathways downstream of IKKε, contribute to the downstream mechanisms through which rs2297550 drives SLE risk.

Other limitations arise from our methodology. Non-coding functional variants may operate in ways not detectable by SNP-seq or by our bioinformatic screens, for example through chromosome conformation, microRNA, and mRNA splicing. We would not have detected variants that operate via DNA-binding proteins absent from the nuclear extract employed, for example, transcription factors specific for other B cell developmental/activation states or in different immune or non-immune lineages. For this reason, while the high-probability SNPs described here are likely functional, we expect that multiple genuinely functional non-coding SNPs will have been filtered out. Definitive validation of the remaining 4 high-probability SNPs will require dedicated study of the kind devoted here to rs2297550.

Our combined experimental and bioinformatic approach also has clear strengths. The number of SNPs that survived each individual filter was substantial, illustrating the difficulty of separating functional from incidental SNPs. Multiple orthogonal approaches are required to validate any individual SNP and remains highly labor-intensive. The value of such an effort arises from the fact that each functional variant points the way to a pathway relevant in human SLE. It is key to remember that the effect size of a non-coding variant reflects the impact of the polymorphism on disease risk, but not the importance of the gene in disease pathogenesis, since regulatory SNPs often alter gene expression only modestly, exactly as observed here. Further, non-coding variants can be expected to identify pathways that coding variants cannot, either because mutations that disrupt a gene are poorly tolerated and thus eliminated from the population or because they lead to a phenotype that is different from that induced by a more subtle alteration in gene expression. As a result, any effort to understand highly polygenic disease must seek to understand not only highly penetrant coding variants but also each non-coding variant.

In summary, we provide a combined experimental and bioinformatic approach to identify plausible regulatory non-coding variants in SLE, proceeding from our initial screen to establish a previously unrecognized pathway through which Ikaros regulates IKKε, as modified by common allelic variation at rs2297550. Further investigation is required to understand how IKKε participates in SLE, including which lineages and downstream pathways are essential to this role. Our approach is broadly applicable and will help to translate GWAS into the pathogenic understanding that is a prerequisite to the personalized treatment of polygenic diseases such as SLE.

**Acknowledgements.** QW, TK, and M-MB were supported by Joint Biology Consortium microgrants off parent grant NIH/NIAMS P30AR070253. JC was supported by P30AR070253. JAS was supported by NIH/NIAMS R01 AR077607, P30 AR070253, and P30 AR072577, the R. Bruce and Joan M. Mickey Research Scholar Fund, and the Llura Gund Award for Rheumatoid Arthritis Research and Care. MTW was supported by NIH grants R01NS099068, R01HG010730, U01AI130830, R01AI024717, R01AR073228. M-GA was supported by NIH/NIAMS P30AR069625, Gilead Sciences, a Global Team Science Award from the Lupus Research Alliance, and the Arthritis National Research Foundation Vic Braden Family

Fellowship. PAN was supported by a Global Team Science Award from the Lupus Research Alliance and NIH/NIAMS 2R01AR065538, R01AR075906, R01AR073201, and P30AR070253.

**Author Contributions.** QW and PAN conceived and designed the study. QW, TK, MM-B, SS performed experiments and data analysis. VA, JC, and MG-A performed data analysis. JAS led recruitment of genotyped human subjects. XTC, MTW identified candidate transcription factors. MT, BW performed experiments. QW, TK, PAN drafted the manuscript and all authors edited and approved the manuscript. PAN supervised the research.

**Declaration of interests.** The authors declare no relevant competing interests.

## REFERENCES

1. Deapen, D., et al., *A revised estimate of twin concordance in systemic lupus erythematosus.* Arthritis Rheum, 1992. **35**(3): p. 311-8.
2. Block, S.R., et al., *Studies of twins with systemic lupus erythematosus. A review of the literature and presentation of 12 additional sets.* Am J Med, 1975. **59**(4): p. 533-52.
3. Alarcon-Segovia, D., et al., *Familial aggregation of systemic lupus erythematosus, rheumatoid arthritis, and other autoimmune diseases in 1,177 lupus patients from the GLADEL cohort.* Arthritis Rheum, 2005. **52**(4): p. 1138-47.
4. Vinuesa, C.G., N. Shen, and T. Ware, *Genetics of SLE: mechanistic insights from monogenic disease and disease-associated variants.* Nat Rev Nephrol, 2023.
5. Caielli, S., Z. Wan, and V. Pascual, *Systemic Lupus Erythematosus Pathogenesis: Interferon and Beyond.* Annu Rev Immunol, 2023. **41**: p. 533-560.
6. Zhang, H., et al., *Meta-analysis of GWAS on both Chinese and European populations identifies GPR173 as a novel X chromosome susceptibility gene for SLE.* Arthritis Res Ther, 2018. **20**(1): p. 92.
7. Morris, D.L., et al., *Genome-wide association meta-analysis in Chinese and European individuals identifies ten new loci associated with systemic lupus erythematosus.* Nat Genet, 2016. **48**(8): p. 940-6.
8. Bentham, J., et al., *Genetic association analyses implicate aberrant regulation of innate and adaptive immunity genes in the pathogenesis of systemic lupus erythematosus.* Nat Genet, 2015. **47**(12): p. 1457-1464.
9. Langefeld, C.D., et al., *Transancestral mapping and genetic load in systemic lupus erythematosus.* Nat Commun, 2017. **8**: p. 16021.
10. Gateva, V., et al., *A large-scale replication study identifies TNIP1, PRDM1, JAZF1, UHRF1BP1 and IL10 as risk loci for systemic lupus erythematosus.* Nat Genet, 2009. **41**(11): p. 1228-33.
11. Westra, H.J., et al., *Fine-mapping and functional studies highlight potential causal variants for rheumatoid arthritis and type 1 diabetes.* Nat Genet, 2018. **50**(10): p. 1366-1374.
12. Butter, F., et al., *Proteome-wide analysis of disease-associated SNPs that show allele-specific transcription factor binding.* PLoS Genet, 2012. **8**(9): p. e1002982.
13. Farh, K.K., et al., *Genetic and epigenetic fine mapping of causal autoimmune disease variants.* Nature, 2015. **518**(7539): p. 337-43.
14. Nicolae, D.L., et al., *Trait-associated SNPs are more likely to be eQTLs: annotation to enhance discovery from GWAS.* PLoS Genet, 2010. **6**(4): p. e1000888.

15. Perez, R.K., et al., *Single-cell RNA-seq reveals cell type–specific molecular and genetic associations to lupus.* Science, 2022. **376**(6589): p. eabf1970.
16. Ulirsch, J.C., et al., *Systematic Functional Dissection of Common Genetic Variation Affecting Red Blood Cell Traits.* Cell, 2016. **165**(6): p. 1530-1545.
17. Tewhey, R., et al., *Direct Identification of Hundreds of Expression-Modulating Variants using a Multiplexed Reporter Assay.* Cell, 2016. **165**(6): p. 1519-1529.
18. Lu, X., et al., *Global discovery of lupus genetic risk variant allelic enhancer activity.* Nature Communications, 2021. **12**(1): p. 1611.
19. Li, G., et al., *High-throughput identification of noncoding functional SNPs via type IIS enzyme restriction.* Nat Genet, 2018. **50**(8): p. 1180-1188.
20. Karrar, S. and D.S. Cunninghame Graham, *Abnormal B Cell Development in Systemic Lupus Erythematosus: What the Genetics Tell Us.* Arthritis Rheumatol, 2018. **70**(4): p. 496-507.
21. Bentham, J., et al., *Genetic association analyses implicate aberrant regulation of innate and adaptive immunity genes in the pathogenesis of systemic lupus erythematosus.* Nature Genetics, 2015. **47**(12): p. 1457-1464.
22. Armstrong, D.L., et al., *GWAS identifies novel SLE susceptibility genes and explains the association of the HLA region.* Genes Immun, 2014. **15**(6): p. 347-54.
23. Li, G., et al., *High-throughput identification of noncoding functional SNPs via type IIS enzyme restriction.* Nat Genet, 2018. **50**(8): p. 1180-1188.
24. Hardenbol, P. and M.W. Van Dyke, *Sequence specificity of triplex DNA formation: Analysis by a combinatorial approach, restriction endonuclease protection selection and amplification.* Proc Natl Acad Sci U S A, 1996. **93**(7): p. 2811-6.
25. Trynka, G., et al., *Chromatin marks identify critical cell types for fine mapping complex trait variants.* Nat Genet, 2013. **45**(2): p. 124-30.
26. Rentzsch, P., et al., *CADD: predicting the deleteriousness of variants throughout the human genome.* Nucleic Acids Res, 2019. **47**(D1): p. D886-d894.
27. Miller, D.E., et al., *Screening for Functional Non-coding Genetic Variants Using Electrophoretic Mobility Shift Assay (EMSA) and DNA-affinity Precipitation Assay (DAPA).* J Vis Exp, 2016(114).
28. Weirauch, M.T., et al., *Evaluation of methods for modeling transcription factor sequence specificity.* Nat Biotechnol, 2013. **31**(2): p. 126-34.
29. Weirauch, M.T., et al., *Determination and inference of eukaryotic transcription factor sequence specificity.* Cell, 2014. **158**(6): p. 1431-1443.
30. Dvir, S., et al., *Deciphering the rules by which 5′-UTR sequences affect protein expression in yeast.* Proceedings of the National Academy of Sciences, 2013. **110**(30): p. E2792-E2801.
31. Mignone, F., et al., *Untranslated regions of mRNAs.* Genome Biology, 2002. **3**(3): p. reviews0004.1.
32. Zhu, T., et al., *The decreased expression of IKBKE in systemic lupus erythematosus.* Clin Rheumatol, 2020. **39**(9): p. 2611-2617.
33. Wang, Y.F., et al., *Identification of 38 novel loci for systemic lupus erythematosus and genetic heterogeneity between ancestral groups.* Nat Commun, 2021. **12**(1): p. 772.
34. Kwon, Y.C., et al., *Update on the Genetics of Systemic Lupus Erythematosus: Genome-Wide Association Studies and Beyond.* Cells, 2019. **8**(10).
35. Wang, Q., et al., *Identification of a regulatory pathway governing TRAF1 via an arthritis-associated non-coding variant.* bioRxiv, 2022: p. 2022.12.15.520628.
36. Kim, T., et al., *Non-coding autoimmune risk variant accelerates T peripheral helper cell development via ICOS.* bioRxiv, 2022: p. 2022.12.16.520733.

37. Chen, L., D.L. Morris, and T.J. Vyse, *Genetic advances in systemic lupus erythematosus: an update.* Curr Opin Rheumatol, 2017. **29**(5): p. 423-433.

38. Pott, J., et al., *Genetically regulated gene expression and proteins revealed discordant effects.* PLoS One, 2022. **17**(5): p. e0268815.

39. Robinson, J.W., et al., *Evaluating the potential benefits and pitfalls of combining protein and expression quantitative trait loci in evidencing drug targets.* bioRxiv, 2022: p. 2022.03.15.484248.

40. Gutierrez-Arcelus, M., et al., *Allele-specific expression changes dynamically during T cell activation in HLA and other autoimmune loci.* Nat Genet, 2020. **52**(3): p. 247-253.

41. Cytlak, U., et al., *Ikaros family zinc finger 1 regulates dendritic cell development and function in humans.* Nature Communications, 2018. **9**(1): p. 1239.

42. Clément, J.F., S. Meloche, and M.J. Servant, *The IKK-related kinases: from innate immunity to oncogenesis.* Cell Res, 2008. **18**(9): p. 889-99.

43. Fitzgerald, K.A., et al., *IKKε and TBK1 are essential components of the IRF3 signaling pathway.* Nature Immunology, 2003. **4**(5): p. 491-496.

44. Tsukamoto, H., et al., *Lipopolysaccharide (LPS)-binding protein stimulates CD14-dependent Toll-like receptor 4 internalization and LPS-induced TBK1–IKKϵ–IRF3 axis activation.* Journal of Biological Chemistry, 2018. **293**(26): p. 10186-10201.

45. Peters, R.T., S.M. Liao, and T. Maniatis, *IKKepsilon is part of a novel PMA-inducible IkappaB kinase complex.* Mol Cell, 2000. **5**(3): p. 513-22.

46. Psarras, A., M. Wittmann, and E.M. Vital, *Emerging concepts of type I interferons in SLE pathogenesis and therapy.* Nat Rev Rheumatol, 2022. **18**(10): p. 575-590.

47. Lafont, E., et al., *TBK1 and IKKε prevent TNF-induced cell death by RIPK1 phosphorylation.* Nature Cell Biology, 2018. **20**(12): p. 1389-1399.

48. Renner, F., R. Moreno, and M.L. Schmitz, *SUMOylation-dependent localization of IKKepsilon in PML nuclear bodies is essential for protection against DNA-damage-triggered cell death.* Mol Cell, 2010. **37**(4): p. 503-15.

49. Kennedy, W.P., et al., *Association of the interferon signature metric with serological disease manifestations but not global activity scores in multiple cohorts of patients with SLE.* Lupus Sci Med, 2015. **2**(1): p. e000080.

50. Akita, K., et al., *Interferon α Enhances B Cell Activation Associated With FOXM1 Induction: Potential Novel Therapeutic Strategy for Targeting the Plasmablasts of Systemic Lupus Erythematosus.* Frontiers in Immunology, 2021. **11**.

51. Wu, C., et al., *Bioinformatics analysis of differentially expressed gene profiles associated with systemic lupus erythematosus.* Mol Med Rep, 2018. **17**(3): p. 3591-3598.

52. Romano, M., et al., *Expanded Regulatory T Cells Induce Alternatively Activated Monocytes With a Reduced Capacity to Expand T Helper-17 Cells.* Frontiers in Immunology, 2018. **9**.

53. Heizmann, B., P. Kastner, and S. Chan, *The Ikaros family in lymphocyte development.* Curr Opin Immunol, 2018. **51**: p. 14-23.

54. Chen, L., et al., *IKZF1 polymorphisms are associated with susceptibility, cytokine levels, and clinical features in systemic lupus erythematosus.* Medicine (Baltimore), 2020. **99**(41): p. e22607.

55. Han, J.W., et al., *Genome-wide association study in a Chinese Han population identifies nine new susceptibility loci for systemic lupus erythematosus.* Nat Genet, 2009. **41**(11): p. 1234-7.

56. Cunninghame Graham, D.S., et al., *Association of NCF2, IKZF1, IRF8, IFIH1, and TYK2 with systemic lupus erythematosus.* PLoS Genet, 2011. **7**(10): p. e1002341.

57. Wang, C., et al., *Genes identified in Asian SLE GWASs are also associated with SLE in Caucasian populations.* Eur J Hum Genet, 2013. **21**(9): p. 994-9.

58. Vyse, T.J. and D.S. Cunninghame Graham, *Trans-Ancestral Fine-Mapping and Epigenetic Annotation as Tools to Delineate Functionally Relevant Risk Alleles at IKZF1 and IKZF3 in Systemic Lupus Erythematosus.* Int J Mol Sci, 2020. **21**(21).

59. Hoshino, A., et al., *Abnormal hematopoiesis and autoimmunity in human subjects with germline IKZF1 mutations.* J Allergy Clin Immunol, 2017. **140**(1): p. 223-231.

60. Van Nieuwenhove, E., et al., *A kindred with mutant IKAROS and autoimmunity.* J Allergy Clin Immunol, 2018. **142**(2): p. 699-702.e12.

61. Merrill, J.T., et al., *Phase 2 Trial of Iberdomide in Systemic Lupus Erythematosus.* N Engl J Med, 2022. **386**(11): p. 1034-1045.

62. Lipsky, P.E., et al., *Biological impact of iberdomide in patients with active systemic lupus erythematosus.* Ann Rheum Dis, 2022. **81**(8): p. 1136-42.

63. Wojcik, H., et al., *Expression of a non-DNA-binding Ikaros isoform exclusively in B cells leads to autoimmunity but not leukemogenesis.* Eur J Immunol, 2007. **37**(4): p. 1022-32.

64. Schwickert, T.A., et al., *Ikaros prevents autoimmunity by controlling anergy and Toll-like receptor signaling in B cells.* Nature Immunology, 2019. **20**(11): p. 1517-1529.

65. Allman, D., et al., *Ikaros is required for plasmacytoid dendritic cell differentiation.* Blood, 2006. **108**(13): p. 4025-34.

66. Rivellese, F., et al., *Effects of targeting the transcription factors Ikaros and Aiolos on B cell activation and differentiation in systemic lupus erythematosus.* Lupus Sci Med, 2021. **8**(1).

67. Nelson, M.R., et al., *The support of human genetic evidence for approved drug indications.* Nat Genet, 2015. **47**(8): p. 856-60.

68. King, E.A., J.W. Davis, and J.F. Degner, *Are drug targets with genetic support twice as likely to be approved? Revised estimates of the impact of genetic support for drug mechanisms on the probability of drug approval.* PLoS Genet, 2019. **15**(12): p. e1008489.

**Figure 1. Screening candidate regulatory variants for SLE by SNP-seq. (A)** Flow chart illustrating identification of 2,180 candidate SNPs from SLE GWAS data. **(B)** SNP-seq: To generate the SNP-seq construct (top), a 31 bp sequence centered on the SNP is positioned between two type IIS restriction enzyme (IIS RE) binding sites. SNPs that fail to bind regulatory proteins such as transcription factors (TF) are negatively selected (bottom), allowing enrichment of protected constructs by PCR. The whole construct can be amplified using primers as per **Suppl. Table 8**. Bio, biotin. Created with BioRender. **(C)** The experimental procedure for SNP-seq; NE, nuclear extract; Bio, biotin. **(D)** Spearman's correlation analysis of SNP-seq allele counts normalized to control between PBMC and BL2 samples. **(E)** We selected 248 SNPs that passed next generation sequencing quality control (NGS-QC) and demonstrated progressive allele-specific protection (**Suppl. Figure. 1C**).

**Figure 2. Prioritization of candidate SNPs by SNP-seq and bioinformatic enrichment.** SNPs selected for experimental screening were the top 20 of 248 emerging form SNP-seq, the top 8 emerging from H3K4me3 score, the top 8 emerging from CADD, and 18 SNPs identified by at least two analyses: 6 SNP-seq+H3K4me3, 4 SNP-seq+CADD, and 8 H3K4me3+CADD.

**Figure 3. EMSA and luciferase reporter assessment of 52 candidate regulatory SNPs. (A)** 7 biallelic and 2 triallelic (rs2297550 and rs936394) SNPs showed consistent allele-imbalanced binding for nuclear by EMSA. Allele-specific gel shift/binding marked with red circles. PBMC data shown; BL2 data and negative SNPs provided in **Suppl. Fig.4**. **(B)** Luciferase reporter assay between the reference (red) and alternative (blue) alleles of 9 candidate regulatory SNPs from A in Daudi B cells (mean ± s.d, n=8 biological replicates, Mann-Whitney test with two-stage step-up correction for multiple hypothesis testing: rs2297550 p=0.00047, rs906868 p=0.0013, rs936394 p= 0.00047, rs9907966 p=0.0071, rs13213604 p=0.0085). **(C)** Diagram displaying the position of 5 SNPs that showed consistent significant differences between alleles through EMSA and luciferase reporter assay; rs2297550 in promoter of *IKBKE*, rs906868 in intergenic region between *YPEL5* and *LBH*, rs936394 in intron of *WBP2*, rs9907966 in intron of *IKZF3*, rs13213604 in intron of *BLTP3A*. The unit of chromosome position is in kilobases (Kb). Diagram of gene features including exon, intron, and UTR was generated using GSDS 2.0 (http://gsds.gao-lab.org/index.php).

**Figure 4. Characterization of functional SNP rs2297550 by CRISPR-mediated editing in Daudi cells. (A)** mRNA levels of *IKBKE* and $\beta$-actin by RT-qPCR and **(B)** protein levels of IKK$\epsilon$ (80 KDa) and GAPDH (37 KDa) were measured from base-edited 4 C/C, 3 C/G, and 3 G/G clones (mean $\pm$ s.d, *P* values from one-way ANOVA corrected by Tukey's multiple comparisons).

**Figure 5. SNP rs2297550 associated with IKKϵ expression in primary PBMCs.** IKKϵ expression was determined by flow cytometry for 35 healthy subjects with C/C (n=12), C/G (n=11), G/G (n=12) genotype at SNP rs2297550. The mean fluorescence intensity (MFI) of IKKϵ expression in (**A-F**) resting and (**G-L**) 4 hours LPS-stimulated PBMCs, including CD19+ B cells, CD3+CD4+ T cells, CD3+CD8+ T cells, CD3-CD56+ NK cells, and CD14+ monocytes (mean ± s.d, *P* values from one-way ANOVA corrected by Dunnett's multiple comparisons test).

**Figure 6. rs2297550 risk allele G binds Ikaros. (A)** Oligonucleotide pulldown western blot assay testing Ikaros binding to rs2297550-G; binding is eliminated in the presence of a 30-fold excess of non-biotinylated rs2297550-C or G competitor; positive control, nuclear extract. **(B)** EMSA supershift assay showing that anti-Ikaros caused fading of the shifted band (oligo rs2297550-G, Daudi nuclear extract), whereas isotype control did not; anti-E2F4 also exhibited specificity. **(C)** ChIP-qPCR using anti-Ikaros confirms binding to rs2297550 (Daudi cells, mean ± s.d, n=3). **(D)** Allele discrimination ChIP-qPCR shows Ikaros preferentially binds to G allele over C allele of rs2297550 in CRISPR-HDR edited C/G Daudi cells (mean ± s.d, n=3). Statistical analysis for C and D: paired *t* test with two tails, without correction.

**Figure 7. Ikaros binding to rs2297550 regulates IKBKE/IKKϵ expression.** CRISPR-Cas9 Ikaros knockout Daudi clones show **(A)** decreased Ikaros protein expression by western blotting, **(B)** loss of Ikaros binding to rs2297550 by ChIP-qPCR (mean ± s.d, n=3),  and **(C, D)** increased *IKBKE* mRNA by RT-qPCR (mean ± s.d, n=3) and increased IKKϵ protein expression by western blotting. Statistical analysis panels B and C: one-way ANOVA corrected by Dunnett's multiple comparisons test.

**Figure 8.** The lupus risk SNP rs2297550 allelically modulates binding of the transcription factor Ikaros (encoded by *IKZF1*) to regulate expression of IKBKE/IKKε, a protein that modulates multiple processes implicated in SLE pathogenesis. The role of mechanisms downstream of IKKε remains to be defined. Created with BioRender.