Title: In vitro investigation of antibacterial activity of Gum Arabic prepared by

two different processing methods against Enterococcus faecalis

Authors: Nuha Elmubarak[a], Yahyia Ibrahim [a], Abbas Gareeballah[b], Nada Sanhouri[a],

Affiliations:

[a] Restorative Department, Faculty of Dentistry /University of Khartoum, Sudan

[b] Anatomy Department, Faculty of Dentistry /University of Khartoum, Sudan

Corresponding Author:

Full name: Nuha Abdel-Rahman Elmubarak

Department: Restorative Department

Institute/University: Faculty of Dentistry /University of Khartoum

Street Name: Al Qasr Street

City, State, Postal code, Country: Khartoum, Khartoum, 11111, Sudan

Whats App: (+249)912292945

E-mail: naina82000@yahoo.com

Abstract:

Background

E. faecalis is a known cause of endodontic treatment failure. Synthetic drugs have been preferred for over decades but recently many plants have been reported for their antibacterial activity. The aim of this study is to investigate the antibacterial effect of mechanically ground and spray dried Acacia senegal against Enterococcus faecalis.

Method:

Antibacterial susceptibility test against Enterococcus faecalis (ATCC 29212) has been performed for 200mg of ethanolic extracts of spray dried & mechanically ground Gum Arabic. Sodium Hypochlorite (1%), Chlorhexidine (0.2%) and Antibiotic multi disc for gram positive bacteria have been used as positive controls and ethanol as a negative control. The diameter of the inhibition zones have been measured.

Results: spray dried Gum Arabic (14.67 mm) was significantly higher than mechanically ground (8.67mm). (p-value=0.02)

Gum Arabic, with both processing methods, exhibited lower antibacterial activity against E. faecalis than chlorhexidine (0.2%).The result was significant with

mechanically ground variant (P-value=.005), but the spray dried Gum showed no

significance.

Spray dried Gum Arabic displayed significantly higher antibacterial activity

against E. faecalis than (Tetracycline 300mcg) (p-value=.0049). While

mechanically ground showed lower antibacterial activity than Tetracycline with no

statistical significance.

Conclusion: Method of Gum Arabic processing affected its antibacterial potency.

Spray-dried Gum Arabic is considered an active antibacterial agent against E.

faecalis, while mechanically ground type is considered a non-active.

Importance:

Enterococcus faecalis has the ability to colonize and hide within the dentinal tubules in the tooth. Dentinal tubules are not accessible by instrumentation during endodontic treatment, they can only be reached and cleaned by irrigation. The problem faced by dentists is that E. faecalis is resistant to sodium hypochlorite, the standard irrigant for endodontic microbiota.

The possibility of using Gum Arabic as a natural antibacterial agent against E. faecalis will help to improve the oral health with low cost and with no side effects commonly associated with synthetics antibacterial agents.

Main Text:

Introduction:

Enterococcus faecalis is a gram-positive, facultative anaerobic coccus bacteria. It can survive under harsh conditions (1, 2). They are part of the habitual flora in the mouth, human gastrointestinal tract, and female genital tract. E. faecalis is a known cause of endodontic treatment failure as well as some systemic diseases. They have developed high resistance to antimicrobial agents (1).

Isolates from oral infections differ from hospital-derived isolates as they do not present many mobile genetic elements. However, oral isolates have virulence factors related to adhesion and biofilm formation, which help the bacteria to colonize different oral sites. Moreover, oral strains may also carry specific antibiotic resistance determinants that have the potential to be transferred to other pathogenic bacteria in biofilm communities (2).

Synthetic drugs have been preferred for decades, but now herbal medicines are being used. Recently, several plants have been investigated for their ability to inhibit the growth of microorganisms(3). Screening of promising medicinal plants is ongoing now and then.
Gum Arabic "Acacia Senegal" has been considered one of 10 promising medicinal plants in Africa (4). It is a naturally occurring gum that exudes from Acacia trees (5). Sudan is the leading producer of Acacia Gums worldwide, followed by

Nigeria, Chad, Mali, and Senegal (6). Phytochemical screening of Acacia senegal has displayed the presence of tannins, glycosides, and flavonoids, in addition to oxidases, peroxidases, and pectinases, which have antimicrobial properties (7-10).

Many studies have proved the antibacterial efficacy of Gum Arabic against various bacterial species (7, 8, 11-15).

A considerable number of medicinal plants have been investigated for their antimicrobial potency against Enterococcus faecalis (16). No studies consider the effect of Gum Arabic against Enterococcus faecalis.

Gum Arabic "Acacia Senegal" is processed differently according to the required quality. Mechanical milled Gum Arabic, on which the dried Gum nodules are crushed mechanically to produce a granular powder. Spray dried Gum Arabic, on which solution of Gum Arabic is pasteurized and sprayed into a stream of hot air to completely evaporate the water leaving the dry powder of Gum Arabic. Gum Arabic produced by spray drying has an advantage over raw gum as it is free of microbial contamination and dissolves much faster. (10)

The aim of this study is to investigate the antibacterial effect of mechanically ground and spray-dried Gum Arabic "Acacia senegal" against Enterococcus faecalis.

Material and method:

Collection of the Plant:

Gum nodules have been collected and authenticated at the Herbarium of Medicinal and Aromatic Plants & Traditional Medicine research institute with code number G-1983-1- MAPTRI-H. After that, nodules have air-dried at room temperature away from the sun and processed in two different processing methods, spray drying and mechanically grounding with a pestle and mortar.

Preparation of extract:

One hundred and fifty grams of each type of Gum Arabic have soaked in ethanol (70%) for three days with daily filtration. The solvent has been evaporated under reduced pressure using rotary evaporator apparatus. The extracts of both types of Gum Arabic have been inserted in a freeze dryer till complete dryness.

Antibacterial susceptibility test:

Antibacterial susceptibility test has performed by Agar disc diffusion according to the method of Elkhateeb 2017 (17) with the following steps:

1. Preparation of the media: Muller-Hinton Agar (HIMEDIA, REF. M173-500G) has been prepared according to instructions. Each 20 mL of the freshly prepared and autoclaved Muller-Hinton agar has been poured into a sterile petri dish and maintained at room temperature to cool down. Before use, the plates were incubated at 35 °C for 48 hours, and the sterility was checked.

2. Preparation of discs: Filter paper discs of 6 mm diameter have been prepared from Whatman filter paper No. 1, placed in a petri dish, and sterilized in a hot air oven at 160 °C for two hours.

3. Preparation of extracts solutions: 200 mg of each extract has been dissolved in 1ml of solvent (ethanol 70%) and mixed in a vortex to ensure complete dissolution.

4. Reactivation and inoculation of Enterococcus faecalis: E. faecalis (ATCC 29212) has reactivated into 5ml of brain heart infusion (BHI) broth (HIMEDIA, REF. M210-500G) and left under anaerobic conditions for 48 hours. A sterile cotton swab has been used to inoculate E. faecalis from BHI broth into a Muller Hinton Agar plate. The plate was incubated under anaerobic conditions at 37∘ for 24 hours.

5. Preparation and standardization of inoculum suspension: Three to five well-isolated colonies of the same morphological type have been selected from a Muller Hinton Agar plate culture and transferred with a sterile loop into a tube containing 5 ml of BHI broth. The broth was incubated anaerobically at 37 °C for 24 hours. The turbidity of E. faecalis suspension has been adjusted equal to that of 0.5 McFarland standard (1.5×10^8 CFU/ml) by using BHI broth to dilute of E. faecalis suspension.

6. Antibacterial susceptibility test:

Sterile cotton swab was rolled in the adjusted bacterial suspension to streak the five plates' surface of Muller Hinton Agar.

Each three sterile discs have immersed in 10 μL of the following extracts solutions:

- Spray-dried Gum Arabic
- Mechanically ground Gum Arabic

Ethanol (70%) (Reagents Duksan, Ethyl Alcohol, absolute, product No.6923, 2.5L) has been used as a negative control.

Chlorhexidine 0.2% (Clenora mouthwash, BP 0.2%w/v), Sodium hypochlorite 1% (Prevest Denpro Hyposol, 3% Sodium hypochlorite 500 ml), and Readymade multidisc for Gram-positive isolates (Axiom laboratories, Code No. 001, Mantola Pahar Ganj- New Delhi-110055) have used as positive controls.

The antibiotics included in the multidisc for Gram-positive have displayed in (Figure 1).

The saturated discs have transferred to their coded plates which inoculated with E. faecalis. The plates have been incubated at 37°C for 24 hours under anaerobic conditions. The plates have been observed for the formation of a clear zone

154   around each disc (inhibition zones). The diameter of the inhibition zone has been

155   measured.

Results:

Gum Arabic with different processing method inhibited the growth of E. faecalis with variant degrees. The inhibition zone measurements reflect the degree of antibacterial activity (Table 1) (Fig. 2-7). Data has been analyzed using SPSS 20.

Gum Arabic, with both processing methods, displayed antibacterial activity against E. faecalis. However, the inhibition zone of the spray-dried Gum Arabic (14.67 mm) was significantly higher than mechanically ground (8.67mm) (level of significance≤0.05, P-value =0.02).

Gum Arabic, with both processing methods, exhibited lower antibacterial activity against E. faecalis than chlorhexidine (0.2%). The result was significant with the mechanically ground variant, but the spray dried Gum showed no significance (P-Values in Table 2).

 Spray-dried Gum Arabic displayed a greater diameter of the inhibition zone (14.67mm) than 1% Sodium hypochlorite (12mm), but the mechanically ground type showed lower activity (8.67mm). However, both results showed no statistical significance (P-values in Table 2).

Only Ciprofloxacin 5mcg, Levofloxacin 5mcg, Cotrimoxazole 25mcg & Tetracycline 300mcg of antibiotics in the multidisc for Gram-positive bacteria exhibited antibacterial activity against E. faecalis (Table 1).

One sample t-test was used to check the significant difference when comparing

Gum Arabic to antibiotic multidisc (Tables 3, 4, and 5).

Gum Arabic, with both processing methods, displayed significantly lower

antibacterial activity to E. faecalis than (Ciprofloxacin 5mcg) & (Levofloxacin

5mcg) (P-values in Table 3).

Spray-dried Gum Arabic (14.6 mm) showed almost the same antibacterial effect

against E. faecalis as Cotrimoxazole 25 mcg (15mm), but the result was

statistically not significant. However, mechanically ground Gum Arabic was

significantly lower than Cotrimoxazole (P-values in Table 4).

Spray-dried Gum Arabic displayed significantly higher antibacterial activity

against E. faecalis than (Tetracycline 300mcg). In contrast, mechanically ground

showed lower antibacterial activity than Tetracycline with no statistical

significance (P-values in Table 5).

Chlorhexidine (0.2%) showed a greater inhibition zone diameter (16 mm) against

E. faecalis than sodium hypochlorite (1%) (12mm). However, the result was

statistically insignificant (level of significance$\leq$0.05, P-value=0.11).

Discussion:

Spray-dried Gum Arabic was found to be active antibacterial against Enterococcus faecalis. In contrast, mechanically ground Gum was found to be inactive antibacterial. The activity scale used in the study referred to Alves et al.(18), and Mukhtar & Ghori (19), who regarded an inhibition zone less than 9mm (inactive product), 9-12mm (partially active product), 13-18mm (active product) , and more than 18mm(very active product).

The chemical composition of Gum Arabic and its pharmacological properties may not always be consistent or stable. It may vary depending on several climatic factors, edaphic conditions, season, age of the tree from which the gum is collected, and other variables (20).

The current study has displayed better antibacterial activity of spray-dried Gum Arabic against E. faecalis over the mechanically ground type. This result clearly reflects the effect of the processing method on the antibacterial activity of Gum Arabic against E. faecalis. However, variation in the processing method does not necessarily affect the antibacterial activity of Gum Arabic against all bacterial specie. It has been found that variation in the processing method did not affect the antibacterial activity of Gum Arabic against Streptococcus mutans.(N. Elmubarak, Y. Ibrahim, A. Gareeballah, and N. Sanhouri submitted for publication)

Based on the current study, the authors agreed with Ali and his colleagues (20),

who recommended using of a single source of Gum Arabic for research work to

achieve standardization for different studies.

Antibacterial activity of spray dried Gum Arabic and mechanically ground is also

differ in their statistical significance when compare them to chlorhexidine and

sodium hypochlorite

This study has displayed that Chlorhexidine (0.2%) has greater antibacterial

activity against E. faecalis than Sodium hypochlorite (1%). This result come in

corroborates with Ferraz et al. (21) and Echeverri & Alderete (22)

Conclusion:

The method of Gum Arabic processing affected its antibacterial potency. Spray-

dried Gum Arabic is considered an active antibacterial agent against E. faecalis,

while mechanically ground type is regarded as non-active.

Spray-dried Gum Arabic has higher antibacterial activity against E. faecalis than

Tetracycline 300mcg.

References:

1. Rodríguez-Niklitschek C. Clinical implications of Enterococcus faecalis microbial contamination in root canals of devitalized teeth: Literature review. Revista Odontológica Mexicana. 2015;19(3):181-6.

2. Pinheiro E, Mayer M. Enterococcus faecalis in oral infections. J Interdiscipl Med Dent Sci. 2014;3(1):1-5.

3. Gull T, Sultana B, Bhatti IA, Jamil A. Antibacterial potential of Capparis spinosa and Capparis decidua extracts. Int J Agric Biol. 2015;17:727-33.

4. Mahomoodally MF. Traditional Medicines in Africa: An Appraisal of Ten Potent African Medicinal Plants. J Evid Based Complement Altern Med. 2013;2013.

5. Glover DA. The effects of dietary supplementation with Gum Arabic on blood pressure and renal function in subjects with type 2 Diabetes Mellitus: Cardiff University; 2012.

6. Musa HH, Ahmed AA, Musa TH. Chemistry, biological, and pharmacological properties of gum Arabic. Bioactive Molecules in Food; Springer International Publishing AG: Cham, Switzerland. 2018:1-18.

7. Okoro S, Kawo A, Arzai A. Phytochemical screening, antibacterial and toxicological activities of Acacia senegal extracts. Bayero Journal of Pure and Applied Sciences. 2012;5(1):163-70.

8. Kubmarawa D, Ajoku G, Enwerem N, Okorie D. Preliminary phytochemical and antimicrobial screening of 50 medicinal plants from Nigeria. African Journal of Biotechnology. 2007;6(14).

9. Marwah RG, Fatope MO, Al Mahrooqi R, Varma GB, Al Abadi H, Al-Burtamani SKS. Antioxidant capacity of some edible and wound healing plants in Oman. Food chemistry. 2007;101(2):465-70.

10. Montenegro MA, Boiero ML, Valle L, Borsarelli CD. Gum Arabic: More Than an Edible Emulsifier. In: Verbeek DJ, editor. Products and Applications of Biopolymers InTech; 2012.

11. Clark DT, Gazi MI, Cox SW, Eley BM, Tinsley GF. The effects of Acacia arabica gum on the in vitro growth and protease activities of periodontopathic bacteria. J Clin Periodontol. 1993;20(4):238-43.

12. Shehu Z, Lamayi DW, Sabo MA, Shafiu M. Synthesis, Characterization and Antibacterial Activity of Kaolin/Gum Arabic Nanocomposite on Escherichia Coli and Pseudomonas Aeruginosa. Res J Nanosci Eng. 2018;2:23-9.

13. Bnuyan I, Hindi N, Jebur M, Mahdi M. In vitro antimicrobial activity of gum Arabic (Al Manna and Tayebat) prebiotics against infectious pathogens. Ijppr Human. 2015;3(3):77-85.

14. Al Alawi SM, Hossain MA, Abusham AA. Antimicrobial and cytotoxic comparative study of different extracts of Omani and Sudanese Gum acacia. Beni-Suef University Journal of Basic and Applied Sciences. 2018;7(1):22-6.

15. Banjar M, Khafaji A, Maher Y. Antimicrobial activity of hydrogen peroxide, sesame and gum Arabic against Streptococcus. Int J Health Sci Res. 2017;7:97-104.

16. Almadi EM, Almohaimede AA. Natural products in endodontics. Saudi Medical Journal. 2018;39(2):124.

17. Elkhateeb A, Abdel Latif RR, Marzouk MM, Hussein SR, Kassem ME, Khalil WK, et al. Flavonoid constituents of Dobera glabra leaves: amelioration impact against CCl4-induced changes in the genetic materials in male rats. Pharmaceutical Biology. 2017;55(1):139-45.

18. Alves TMdA, Silva AF, Brandão M, Grandi TSM, Smânia EdFA, Smânia Júnior A, et al. Biological screening of Brazilian medicinal plants. Memórias do Instituto Oswaldo Cruz. 2000;95(3):367-73.

19. Mukhtar S, Ghori I. Antibacterial activity of aqueous and ethanolic extracts of garlic, cinnamon and turmeric against Escherichia coli ATCC 25922 and Bacillus subtilis DSM 3256. International Journal of applied biology and pharmaceutical Technology. 2012;3(2):131-6.

20. Ali BH, Beegam S, Al-Lawati I, Waly MI, Al Za'abi M, Nemmar A. Comparative efficacy of three brands of gum acacia on adenine-induced chronic renal failure in rats. Physiol Res. 2013;62(1):47-56.

21.   Ferraz CC, Gomes BP, Zaia AA, Teixeira FB, Souza-Filho FJ. Comparative study of the
antimicrobial efficacy of chlorhexidine gel, chlorhexidine solution and sodium hypochlorite as
endodontic irrigants. Brazilian dental journal. 2007;18(4):294-8.
22.   Echeverri D, Alderete D, ECHEVERRI D, ALDERETE D. In vitro Antibacterial Effect of 2%
Chlorhexidine Against Enterococcus faecalis in Dentin Previously Irrigated with 5% Sodium Hypochlorite.
Int J Odontostomatol. 2015;9(1):25-9.

Figures

Figure 1: Multiple antibiotic disc (Axiom laboratories, multidisc for Gram positive isolates, Code No. 001, Mantola Pahar Ganj- New Delhi-110055)

Figure 2: Inhibition zones around a disc saturated with spray dried Gum Arabic on plates inoculated with E. faecalis

Fig. 3: Inhibition zones around a disc saturated with mechanically ground Gum Arabic on plates inoculated with E. faecalis

Fig. 4: Inhibition zones around a disc saturated with Sodium hypochlorite 1% (Hyposol) wash on plates inoculated with E. faecalis

Fig. 5: Inhibition zones around a disc saturated with Chlorhexidine 0.2% (positive control) on plates inoculated with E. faecalis

Fig. 6: Inhibition zones around a disc multidisc antibiotic for Gram positive bacteria (positive control) on plates inoculated with E. faecalis

Fig. 7: Inhibition zones around a disc saturated with Ethanol 70% (negative control) on plates inoculated with E. faecalis

Table 1: Measurements of the inhibition zones of Gum Arabic against E. faecalis.

Plant extract (200mg/ml)	Inhibition zone diameter (mm)					
	M1	M 2	M3	N	mean	Std. dev.
Spray dried Gum Arabic	14	15	15	3	14.67	0.58
Mechanically Gum Arabic	9	8	9	3	8.67	0.58
Ethanol 70% (-ve)	0	0	0	3	0	0.00
Sodium hypochlorite (1%)	12	12	12	3	12	0.00
Chlorhexidine (0.2%)	15	16	17	3	16	1.00
CP (Ciprofloxacine 5mcg)	20	-	-	1		
LE (Levofloxacine 5mcg)	20	-	-	1		
BA (Co-Trimoxazole 25 mcg)	15			1		
TE (Tetracycline 300mcg)	10			1		

(M1), (M2), and (M3): measurements of inhibition zones around first, second and third discs

and (N): number of measurements for each extract

Table 2: Multiple comparisons between spray dried and mechanically milled Gum Arabic chlorhexidine, Sodium Hypochlorite and antibiotics for their antibacterial activity against E. faecalis using least significant difference (LSD). P-values are presented on the table and the level of significance is 0.05.

(I) Plant2	plant (J)	Mean Difference (I-J)	Std. Error	P-Value	95% Confidence Interval Lower Bound	Upper Bound
Spray dried Gum	Mechanically ground Gum	6.000	2.414	.020	1.037	10.963
	Ethanol 70% (-ve)	14.667	2.414	.000	9.704	19.629
	NaOCL(1%) (+ve)	2.667	2.414	.279	-2.296	7.629
	CHX (0.2%) (+ve)	-1.333	2.414	.585	-6.296	3.629
Mechanically ground Gum	Ethanol 70% (-ve)	8.667	2.414	.001	3.704	13.629
	NaOCL (1%) (+ve)	-3.333	2.414	.179	-8.296	1.629
	CHX (0.2%) (+ve)	-7.333	2.414	.005	-12.296	-2.371

NaOCL: Sodium hypochlorite (Clorox),

CHX: Chlorhexidine

Table 3: One sample t-test statistics to compare the inhibition zone diameter (mm) of Gum Arabic against **E. faecalis** to the inhibition zone of (Ciprofloxacin 5mcg) and (Levofloxacin 5mcg). P-Values are presented on the table and the level of significance is 0.03. **(Test Value = 20)**

No.	Plant extract	Mean of Inhibition zone diameter (mm)	Std. Deviation	t	P-Value	Comment
(1)	Spray dried Gum Arabic	14.67	0.58	-16.000	.004	S
(2)	Mechanically milled Gum Arabic	8.67	0.58	-34.000	.001	S

Table 4: One sample t-test statistics to compare the inhibition zone diameter of Gum Arabic against E. faecalis to the inhibition zone of CoTrimoxazole 25 mcg. P- Values are presented in the table and the Level of significance is 0.03 **(Test Value = 15)**

No.	Plant extract	N	Mean of Inhibition zone diameter (mm)	Std. Deviation	P-Value	Comment
(1)	Spray dried Gum Arabic	3	14.6667	.57735	.423	NS
(2)	Mechanically milled Gum Arabic	3	8.6667	.57735	.003	S

NS: Not significant, **S:** Significant

Table 5: One sample t-test statistics to compare the inhibition zone diameter (mm) of Gum Arabic against E. faecalis to the inhibition zone of Tetracycline 300mcg. P-values are presented on the table and the level of significance is 0.03 **(Test value =10)**

No.	Plant extract	N	Mean	Std. Deviation	P-Value	Comment
(1)	Spray dried Gum Arabic	3	14.667	0.577	.0049	S
(2)	Mechanically milled Gum Arabic	3	8.667	0.577	.057	NS

NS: Not significant**, S:** Significant