Mosaic of local adaptation between white clover and rhizobia along an urbanization

gradient

Authors: David Murray-Stoker[1,2,3,*] and Marc T. J. Johnson[2,3]

Affiliations

[1]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario

M5S 3B2, Canada

[2]Department of Biology, University of Toronto Mississauga, Mississauga, Ontario, Canada L5L

1C6

[3]Centre for Urban Environments, University of Toronto Mississauga, Mississauga, Ontario,

Canada L5L 1C6

David Murray-Stoker, ORCID https://orcid.org/0000-0002-4774-6948

Marc T. J. Johnson, ORCID http://orcid.org/0000-0001-9719-0522

Corresponding author: David Murray-Stoker (dstoker92@gmail.com)

Abstract

1. Urbanization frequently alters the biotic and abiotic environmental context. Variation in

both the biotic and abiotic environment can result in selection mosaics that lead to locally

adapted phenotypes. Ecological changes to urban environments have been found to result

in evolutionary responses across a diverse array of plants and animals, but the effects of

urbanization on local adaptation and coevolution between species remains largely

unstudied.

2. Using a common garden experiment of 30 populations and 1080 plants, we tested for

local adaptation in the mutualism between white clover (*Trifolium repens*) and rhizobia

(*Rhizobium leguminosarum* symbiovar *trifolii*) along an urbanization gradient. We asked:

(Q1) are white clover and rhizobia locally adapted? (Q2) Does nitrogen (N) addition

mediate the strength of local adaptation? (Q3) Is the strength of local adaptation in host

plants related to the strength of local adaptation in rhizobia, as expected if white clover

and rhizobia are co-adapting? And (Q4) How does the degree of local adaptation vary

with urbanization?

3. We found evidence for local adaptation for both white clover and rhizobia, with stronger

signatures of local adaptation for rhizobia than white clover (Q1). While N addition

positively affected plant biomass and negatively affected nodule density, N addition did

not consistently mediate patterns of local adaptation (Q2). The strength of local

adaptation was positively related between white clover and rhizobia under N addition,

suggesting that soil N mediates co-adaptation between white clover and rhizobia (Q3).

We did not find a consistent relationship between measures of urbanization and local

adaptation (Q4). Additionally, fitness responses for white clover and rhizobia were

unrelated to *Rhizobium* abundance and local adaptation was unrelated to the broader

bacterial microbiome in the soil inoculants or root endosphere.

4. *Synthesis*. Our results show a spatial mosaic in local adaptation, with stronger local

adaptation for rhizobia than white clover. While urbanization does influence the ecology

of plant-microbe interactions, our study suggests urbanization does not disrupt local

adaptation and coevolution among mutualists.

Keywords: coevolution, legume-rhizobia interaction, local adaptation, mutualism, microbiome,

plant-microbe interactions, urban evolution

Introduction

Urbanization is a dominant driver of environmental change that frequently alters the ecological

setting of ecosystems (Pickett et al., 2001; Santangelo et al., 2022). Abiotic changes to the

environment commonly include increased impervious surface, habitat fragmentation and

degradation, elevated temperature, higher pollution (e.g., air, water, light, noise), and increased

nutrient deposition (Grimm et al., 2008; Pickett et al., 2011). In addition to urban-driven abiotic

changes, biotic changes frequently include altered species diversity and community composition,

reduced vegetation cover, increased abundance of non-native species, and reduced abundance

and diversity of native species (Faeth et al., 2011; McKinney, 2008). Ecological changes to

urban environments have been found to cause evolution in a diverse array of plants and animals

(Diamond & Martin, 2021; Johnson & Munshi-South, 2017). Despite increasing evidence of

evolutionary consequences in response to urbanization, it remains to be seen if urbanization

affects the local adaptation and (co)evolution of species interactions.

Variation in both the biotic and abiotic environment can result in selection mosaics that

lead to locally adapted phenotypes (Thompson, 2005). Numerous environmental factors can

impose natural selection that lead to local adaptation, whereby local populations have greater

fitness than nonlocal populations (Blanquart et al., 2013). Mutualisms are a useful focal

interaction to study local adaptation because the reciprocal fitness feedbacks among interacting

species should lead to fitness alignment in a given environment but still allow for variation

across environments (Heath & Stinchcombe, 2014; Hoeksema, 2010; Nuismer et al., 1999).

Variation in community composition across space and time can alter mutualistic species

interactions (Thompson, 1988, 2005). Abiotic context can also affect mutualisms by altering the

physical (e.g., temperature; Kiers *et al.* 2010) or resource (e.g., nutrient and water availability;

Bronstein 1994; Shantz *et al.* 2016) environment in which species interact. Moreover, biotic and

abiotic contexts are likely acting in concert to affect the outcome of mutualistic interactions

(Burghardt et al., 2022; Heath et al., 2010; Heath & Tiffin, 2007; Van Cauwenberghe et al.,

2016). Because urbanization alters both the biotic and abiotic environment simultaneously across

spatial and temporal scales (Diamond & Martin, 2021; Johnson & Munshi-South, 2017), urban

environments have the potential to alter selective landscapes, local adaptation, and potentially

coevolution between interacting species.

Mutualistic interactions between legumes and rhizobia are a useful model for evaluating

evolutionary ecology in urban environments. In this mutualism, rhizobia fix atmospheric N_2 into

an accessible form of nitrogen (N) for the plant (NH_4^+) in exchange for photosynthate and

hosting within nodules provided by the plant (Poole et al., 2018). Ecological changes in urban

environments, particularly increased N deposition and enrichment (Grimm et al., 2008; Stevens

et al., 2018), could have consequences for legume-rhizobium local adaptation by disrupting the

economy of the resource exchange: direct acquisition of N is 'cheaper' for the host legume,

which can select for less cooperative rhizobia (Regus et al., 2017; Weese et al., 2015). In a

previous study along an urbanization gradient (Murray☐Stoker & Johnson, 2021), we found that

urbanization altered the ecology of a legume-rhizobia mutualism. Specifically, urban plants

exhibited reduced investment in the mutualism and primarily acquired N through fixation except

at the urban and rural limits of the gradient. Soil N content was the nexus linking the effects of

urbanization to the mutualism (Murray☐Stoker & Johnson, 2021). With this evidence, we

wanted to determine if the ecological changes driven by urbanization affected local adaptation of

white clover and its rhizobia mutualists.

Here we conducted a large, factorial field experiment to test the hypothesis that urban

environmental change drives local adaptation in the mutualism between white clover (*Trifolium*

repens) and rhizobia (*Rhizobium leguminosarum* symbiovar *trifolii*). We asked four specific

questions: **(Q1)** Are white clover and rhizobia locally adapted? **(Q2)** Does N addition mediate

the strength of local adaptation? **(Q3)** Is the strength of local adaptation in host plants related to

the strength of local adaptation in rhizobia, as expected if white clover and rhizobia are

coadapting? **(Q4)** How does the degree of local adaptation vary with urbanization? Results from

this study provide the first test of how urbanization affects the evolution and local adaptation of

interacting species, which provides insight into urban environments and whether they could be

disrupting coevolution.

Materials and Methods

Study System

We used the mutualism between white clover (*Trifolium repens* L., Fabaceae) and its facultative,

rhizobial symbiont (*Rhizobium leguminosarum* symbiovar *trifolii*) to evaluate if and how local

adaptation varied with urbanization. White clover is a perennial, herbaceous legume native to

Eurasia that is now globally distributed. White clover typically reproduces clonally through

stolons as well as through seed via obligately-outcrossed flowers (Burdon, 1983). *Rhizobium*

leguminosarum symbiovar *trifolii* (hereafter rhizobia) is the primary rhizobial symbiont of white

clover (Andrews & Andrews, 2017).

Biological Materials

White clover: The 30 focal white clover populations for this experiment were a subset of those

sampled by Murray-Stoker and Johnson (2021), with focal populations selected to span an

urbanization gradient in the Greater Toronto Area, Ontario, Canada (Figure 1A). To minimize

parental effects, we grew field-collected seeds in the greenhouse to produce outcrossed F_1

populations. All F_1 seeds were derived from a random panmictic cross among 5-10 parent plants

within a source population (detailed methods are provided in Appendix S1).

Soil inoculants: Soil for microbiome inoculants was collected following the methods described

in Murray-Stoker and Johnson (2021). Briefly, 5 soil cores were taken at each site at least 5 m

from the nearest white clover plant and other legumes (e.g., *Medicago lupulina*). Each soil core

was collected to a depth of 5 cm, with the top 2 cm plus any organic material removed.

Individual cores were stored in separate ziploc bags and stored at $-80°C$ until subsequent

processing. Local microbiomes were paired to each of the 30 white clover populations in the

experiment, while the nonlocal$_{Urban}$ and nonlocal$_{Rural}$ microbiome inoculants were pooled from 5

populations at the urban or rural ends of the gradient, respectively (Figure 1A).

We used a local versus nonlocal microbiome inoculation approach (Blanquart et al.,

2013; Heath & Stinchcombe, 2014) to test for local adaptation between white clover and rhizobia

for two principal reasons. First, this approach tests for overall differences in fitness between local

and nonlocal white clover or rhizobia populations, which can be further distinguished based on

the habitat of populations (i.e., urban vs. rural). In particular, increased N deposition in urban

environments can alter the dynamics of the mutualism between white clover and rhizobia by

reducing nodule densities and influencing the source of N used by white clover (Murray☐Stoker

& Johnson, 2021). Additionally, urbanization leads to homogenization of the biotic and abiotic

environment (McKinney, 2006; Santangelo et al., 2022), which might result in homogenized

composition of microbial communities (Delgado-Baquerizo *et al.* 2021; Rosier *et al.* 2021; but

see Epp Schmidt *et al.* 2017). Together, this evidence suggests that the degree of local adaptation

might depend on (1) the nonlocal inoculant representing rural or urban environments and (2)

similarity in microbiome composition between the local and either nonlocal microbiomes (i.e.,

nonlocal$_{Rural}$ and nonlocal$_{Urban}$). Second, by using microbiome inoculants instead of isolated

rhizobia strains, this approach also provided more ecological realism by allowing rhizobia and

other bacteria and fungi to establish in the soil and roots.

Experiment Overview

We tested for local adaptation between populations of white clover and rhizobia using a common

garden experiment (Figure 1B). We used the local versus nonlocal contrast (sensu sympatric vs.

allopatric; Blanquart *et al.* 2013), with local adaptation indicated when mean local fitness was

greater than mean nonlocal fitness. We crossed 30 white clover populations with 3 microbiome

treatments (local, nonlocal$_{Rural}$, and nonlocal$_{Urban}$) and 2 N treatments (no N addition = ambient

N, 10 mM KNO_3 fertilizer = N addition). Each treatment combination had 6 replicates per

population (30 populations $\times$ 3 microbiome treatments $\times$ 2 N treatments $\times$ 6 replicates = 1080

plants). The N addition treatment was representative of the soil N content measured at the focal

populations by Murray-Stoker and Johnson (2021) and was designed to reflect the impacts of

urbanization on soil N. We also had 6 control pots (3 ambient N, 3 N addition) that were not

planted or inoculated but did receive fertilizer additions, and these pots were used to evaluate

dispersal and establishment of bacteria at the experimental site.

Common Garden Experiment

We grew all of the plants outside in a common environment using an incomplete block design at

the University of Toronto Mississauga campus. In total, there were 9 blocks consisting of 120

plants; each block included all treatment combinations but not all population-by-treatment

combinations were represented in each block due to spatial constraints. Prior to planting, seeds

were scarified with sandpaper, surface sterilized (60 s in 70% ethanol, 60 s in 10% household

bleach, 90 s rinse in sterile water) and then germinated in the dark on dampened filter paper for

48 h at 20°C. Germinated seeds were planted in 1 L bleach-sterilized pots in a steam-sterilized (2

h of sterilization) mix of potting soil and sand mixture (2:3 V/V). The soil mixture was selected

to facilitate harvesting of all root biomass at the end of the experiment. We planted two

germinated seeds in each pot, and pots were thinned to one plant between weeks 2-3 of the

experiment.

Microbiome treatments were applied 1 week after planting by applying soil slurries. Soil

slurries for each microbiome treatment were prepared with 7.5 g fresh soil (thawed from –80°C

overnight at 4°C) in 50 mL sterile deionized water; nonlocal treatments were composed of equal

mixtures by mass from 5 populations within the respective habitat type. Slurries were hand

shaken for 60 s after preparation to mix and suspend the soil, with further re-suspension (hand

shaken for 30 s) during inoculation to maintain soil suspension. Each plant received 10 mL of its

respective microbiome inoculant. Weekly N addition treatments began in week 2, and we applied

bi-weekly N-free fertilizer following Batstone *et al.* (2017) starting in week 4 of the experiment.

Both the N-addition and N-free fertilizers were autoclave sterilized (60 min sterilization) prior to

application. Plants were watered daily as needed for the duration of the experiment. We ran the

experiment for 19 weeks (21 May–2 October 2021). Due to germination failure, herbivory, and

other sources of plant loss, our final plant count was 841.

Laboratory Processing

We measured multiple components of fitness for white clover (aboveground biomass and

belowground biomass) and rhizobia (total nodule density and fixing nodule density). We first

washed all plants of soil and debris and then split them into aboveground and belowground

components to facilitate fitness measurements. Aboveground components were placed in

envelopes, dried to constant mass (at least 72 h at 60°C), and aboveground biomass was weighed

to the nearest 1 mg. To measure rhizobia fitness components, a subset of roots was collected

from each plant by cutting the first three roots below the plant base and directly attached to the

stolon (following Murray◻Stoker & Johnson 2021), and we then measured the total root length

of all collected roots (mean ± standard error = 43.5 ± 0.58 cm, range = 2.1-128.7 cm). Nodule

density, an estimate of investment in the mutualism by white clover and rhizobia fitness, was

measured by visually-counting nodules from each measured root and standardizing by the total

root length to generate an estimate of nodule density per cm of root per plant (Murray◻Stoker &

Johnson, 2021). We also calculated the density of fixing nodules, a measure of the effectiveness

of the mutualism, by counting the number of nodules with pink colouration (indicative of N

fixation; Udvardi & Poole 2013) and calculating density as described above. After estimating

nodule density and fixing nodule density, all belowground biomass (including the 3 excised

roots) was placed in an envelope, dried for 72 h at 60°C, and weighed to the nearest 0.1 mg. We

focused on biomass as a measure of plant fitness because white clover undergoes extensive

clonal growth as a perennial. We could not measure sexual fitness accurately because only a

fraction of plants flowered and we could not prevent interpopulation crosses.

DNA Extraction & Sequencing

To determine if white clover and rhizobia fitness responses were influenced by the broader

bacterial microbiome, we sequenced the bacterial microbiome for each of the microbiome

inoculants and a subset of roots from each of the experimental treatments. All 40 microbiome

inoculants (30 focal, 5 nonlocal$_{Rural}$, 5 nonlocal$_{Urban}$) were sequenced to evaluate variation in

bacterial community composition. A total of 60 root microbiomes (10 populations $\times$ 3

microbiome treatments $\times$ 2 N treatments = 60 plants) were sequenced to evaluate: (1) white

clover and rhizobia fitness responses as a function of *Rhizobium* abundance, (2) *Rhizobium*

abundance in response to experimental treatments, and (3) fitness of white clover and rhizobia in

response to soil and root bacterial microbiome composition. We included each treatment

combination for the same 6 populations (i.e., 10 populations $\times$ 6 treatment combinations = 60

samples). Protocols for microbiome DNA extraction are provided in the supplementary material

(Appendix S1).

Statistical Analyses

We provide detailed methods for all statistical analyses in the supplementary material (Appendix

S1). Briefly, we analyzed variation in fitness responses for white clover (aboveground biomass,

belowground biomass) and rhizobia (nodule density, fixing nodule density) in response to the

microbiome treatment (local, nonlocal$_{Rural}$, nonlocal$_{Urban}$), N treatment (ambient N, N addition),

and the two-way interaction using linear mixed-effects models with Type III sums-of-squares.

We also quantified the extent to which population $\times$ environment interactions were due to

changes in population variance and changes in rank order using Cockerhams's method

(Cockerham, 1963; Muir et al., 1992). To test if there was a general signature of local adaptation,

we repeated the same procedure described above but with microbiome recoded as local and

nonlocal$_{Global}$ (i.e., nonlocal$_{Rural}$ and nonlocal$_{Urban}$ combined).

The strength of local adaptation was calculated using three indices: (1) local –

nonlocal$_{Global}$, (2) local – nonlocal$_{Rural}$, and (3) local – nonlocal$_{Urban.}$ These indices were

calculated separately for each of the four fitness response variables (i.e., aboveground biomass,

belowground biomass, nodule density, and fixing nodule density). Positive values were

indicative of local adaptation because they showed greater fitness in the local versus nonlocal

treatment. Local adaptation indices were calculated using the best linear unbiased predictors (i.e.,

BLUPs) from linear mixed-effects models. To test whether the strength of local adaptation was

related between interacting white clover and *Rhizobium* populations (i.e., a signature of

coadaptation), we calculated all pairwise Pearson correlations between the two fitness

components of white clover with the two fitness components of rhizobia. We calculated

correlations separately for each N treatment to examine if N addition mediated coadaptation.

To test how urbanization affects the strength of local adaptation, we regressed local

adaptation indices against measures of urbanization: distance from the city center, mean

impervious surface cover (ISC, 250 m radius around each population), and the Human Influence

Index (HII, 1 km radius around each population). We fitted separate regressions for each local

adaptation index and urbanization metric due to multicollinearity between variables (variance

inflation factors > 5–10). We scaled all predictor variables to a mean of 0 and standard deviation

of 1 prior to analysis to allow for direct comparisons across models.

Microbiome sequences were processed for analysis using the DADA2 pipeline (Callahan

et al., 2016), with taxonomy assigned to amplicon sequence variants using the Ribosomal

Database Project (Cole et al., 2014) and naïve Bayesian classifier (Wang et al., 2007)

implemented in DADA2. We regressed white clover and rhizobia fitness responses against

Rhizobium abundance (number of reads assigned to the genus *Rhizobium*), N treatment, and the

interaction using generalized linear models. We examined differences in *Rhizobium* abundance

in response to microbiome$_{Global}$ (local vs. nonlocal$_{Global}$), N (ambient N, N addition), and the

interaction using ANOVAs with Type III sums-of-squares. Finally, we examined the role of

microbiome composition on local adaptation. We tested if the strength of local adaptation

increased when microbiome inoculant composition was more dissimilar (pairwise Bray-Curtis

dissimilarity) using a linear regression. We also tested if local adaptation indices were related to

root bacterial composition. The effects of the microbiome treatment, N treatment, and their

interaction on root microbiome composition was fit and assessed using PERMANOVA with a

Bray-Curtis dissimilarity matrix (Oksanen et al., 2020). Relationships between root community

composition and local adaptation indices were quantified using the `envfit()` function (Oksanen

et al., 2020).

All statistical analyses were performed using R (version 4.2.2; R Core Team 2022) in the

RStudio environment (version 2022.07.2; RStudio Team 2022). All packages used for data

management, analysis, and figure creation are provided in the detailed statistical analyses in the

supplementary material (Appendix S1). We followed statistical best practices by focusing on

estimates, variation of the estimates, and effect sizes to interpret the results and identify the

strength of evidence (Berner & Amrhein, 2022; Gelman & Hennig, 2017; McShane et al., 2019;

Wasserstein et al., 2019; Wasserstein & Lazar, 2016).

Results

Effects of microbiome and nitrogen treatments on fitness

We found evidence for local adaptation for both white clover and rhizobia, and the effects were

stronger for rhizobia than white clover and varied among populations (Table 1, Figure 2). White

clover demonstrated weak adaptation to local microbiomes for both aboveground biomass (P =

0.047, η^2_P = 0.106, Table 1, Figure 2) and belowground biomass (P = 0.036, η^2_P = 0.008, Table

1, Figure 2). Aboveground biomass was 6.4% and 12.8% greater in local versus nonlocal$_{Rural}$ and

nonlocal$_{Urban}$ microbiomes, respectively (contrasts in Table S1), and 60% of populations

demonstrated patterns consistent with local adaptation (Figure 2). In contrast, belowground

biomass was nearly equivalent (0.8% increase) between local and nonlocal$_{Rural}$ microbiomes,

whereas belowground biomass was 7.4% greater in local than nonlocal$_{Urban}$ microbiomes

(contrasts in Table S1).

Signatures of local adaptation were stronger for rhizobia when estimated as either nodule

density (P < 0.001, η^2_P = 0.247) or fixing nodule density (P < 0.001, η^2_P = 0.267, Table 1, Figure

2). Nodule density was 38.4% and 20.1% greater in local versus nonlocal$_{Rural}$ and nonlocal$_{Urban}$

microbiomes, respectively; however, nodule density depended on the three-way interaction

between microbiome, N, and population (Table 1, Figure 2). There was an increase in the

frequency of populations demonstrating local adaptation under N addition (local vs. nonlocal$_{Rural}$

= 83%, local vs. nonlocal$_{Urban}$ = 62%) compared to ambient N (local vs. nonlocal$_{Rural}$ = 63%,

local vs. nonlocal$_{Urban}$ = 57%; contrasts in Table S1). Fixing nodule density was 53.4% and

33.8% greater in the local microbiome treatment than the nonlocal$_{Rural}$ and nonlocal$_{Urban}$

treatments, respectively (contrasts in Table S1).

As expected, N addition had strong and positive effects on aboveground biomass (81.3%

increase, P < 0.001, η^2_P = 0.871, Table 1, Figure 2) and belowground biomass (75.0% increase, P

< 0.001, η^2_P = 0.324, Table 1, Figure 2;contrasts in Table S1). In contrast, N addition decreased

nodule density by 17.9% for both total nodule fixing density ($P < 0.001$, $\eta^2_P = 0.323$, Table 1,

Figure 2) and fixing nodule density ($P = 0.025$, $\eta2P = 0.167$, Table 1, Figure 2; contrasts in

Table S1). Notably, differences between local and nonlocal microbiomes were greater under N

addition for both nodule density and fixing nodule density (Figure 1). When we further evaluated

microbiome $\times$ population interactions using Cockerham's method, we found that crossing of

reaction norms (i.e., changes in rank order of populations among environments) accounted for

most of the variation in the interaction for aboveground biomass (97.4%), nodule density

(86.9%), and fixing nodule density (97.9%), as opposed to changes in population variance.

Results described above were similar when microbiome treatments were examined as local vs.

nonlocal$_{Global}$: there were stronger effects of local adaptation for both nodule density and fixing

nodule density but no evidence of local adaptation for belowground biomass (Table S2, Figure

S2).

Reciprocal Local Adaptation

White clover and rhizobia exhibited patterns of reciprocal local adaptation consistent with

coevolution, but linkages between white clover and rhizobia depended on the N environment and

the local – nonlocal contrast (Figures 3, S2-S3). A positive correlation between indices of local

adaptation for aboveground biomass and nodule density was apparent under N addition for the

global (local – nonlocal$_{Global}$, r and 95% confidence interval: $r = 0.418$ [0.061, 0.681]; Figure 3),

rural (local – nonlocal$_{Rural}$, $r = 0.396$ [0.034, 0.666]; Figure S2), and urban contrasts (local –

nonlocal$_{Urban}$, $r = 0.301$ [−0.081, 0.606]; Figure S3). There was no clear correlation between

indices of local adaptation for aboveground biomass and nodule density under ambient N (local

– nonlocal$_{Global}$, $r = -0.194$, 95% CI [−0.518, 0.179]; local – nonlocal$_{Rural}$, $r = -0.095$, 95% CI

[−0.440, 0.274]; local – nonlocal$_{Urban}$, $r = -0.087$, 95% CI [−0.434, 0.282]; Figures 3, S2-S3).

Relationships between belowground biomass and nodule density generally switched from

negative under ambient N (local – nonlocal$_{Global}$, r = −0.301 [−0.597, 0.067]; local –

nonlocal$_{Rural}$, r = −0.214 [−0.533, 0.159]; local – nonlocal$_{Urban}$, r = −0.197 [−0.520, 0.176];

Figures 3, S2-S3) to positive under N addition (local – nonlocal$_{Global}$, r = 0.253 [−0.125, 0.567];

local – nonlocal$_{Rural}$, r = 0.305 [−0.069, 0.604]; local – nonlocal$_{Urban}$, r = 0.052 [−0.327, 0.417];

Figures 3, S2-S3). When we examined the potential for reciprocal local adaptation between

fixing nodule density and either aboveground or belowground biomass, we observed positive

correlations in the rural contrast under N addition (aboveground biomass: local – nonlocal$_{Rural}$, r

= 0.424 [0.069, 0.684]; belowground biomass: local – nonlocal$_{Rural}$, r = 0.399 [0.038, 0.668];

Figure S2).

Local Adaptation and Urbanization

We did not identify any strong or consistent relationships between local adaptation and

urbanization, regardless of local adaptation index (Figures 4, S4-S6). There were no consistent

patterns in the strength and direction of relationships between local adaptation indices for our

fitness estimates and urbanization metrics (Figures 4, S4-S6), and all of the relationships

explained a negligible amount of variation (range of $R^2_{Adjusted}$ = 0-0.031).

Fitness and the Broader Microbiome

Variation in the microbiome community did not influence fitness for either white clover or

rhizobia. There was no main effect of *Rhizobium* abundance or *Rhizobium* abundance × N

interaction on any of the four fitness estimates (Table S3, Figure 5); using *Rhizobium* relative

abundance gave similar results (Table S4, Figure S7). Local adaptation was unrelated to

microbiome inoculant dissimilarity (Table S5, Figure 5) or root bacterial composition (Table S6,

Figure S8).

Rhizobium abundance was influenced by the N treatment and an interaction between

microbiome and N treatments (Table S7, Figure S9). Root bacterial community composition did

not vary by microbiome inoculant, N, or the interaction (Table S6, Figure S8). Community

composition was more variable among local microbiome inoculants than among either the

nonlocal$_{Rural}$ or nonlocal$_{Urban}$ inoculants or the experimental controls (Figure S9), and *Rhizobium*

abundances were similar among all microbiome inoculants (Cohen's $d = 0.015$-0.158, Figure

S10).

Discussion

Our experiment documents a mosaic of local adaptation in the mutualism between white clover

and rhizobia along an urbanization gradient, a conclusion supported by four key results. First,

both white clover and rhizobia had greater mean fitness in their local microbiomes relative to the

nonlocal microbiomes despite extensive variation among populations. Second, fitness responses

were unrelated to *Rhizobium* abundance and local adaptation was unrelated to the broader

bacterial microbiome in the soil inoculants or root endosphere. Together, these results suggest

local adaptation is driven by local evolution of *Rhizobium* and not ecological changes in

Rhizobium populations or microbiome composition. Third, N addition did not consistently

mediate the effects of local adaptation in white clover or rhizobia, although N did influence the

presence of reciprocal local adaptation between white clover and rhizobia, whereby the strength

of local adaptation was positively correlated between white clover and rhizobia in the presence

of high N. Finally, we did not find a consistent relationship between measures of urbanization

and local adaptation, suggesting urban environmental change did not alter patterns of local

adaptation. Below, we further discuss these results in the context of local adaptation. legume-

rhizobia coevolution, and urban evolution.

Local adaptation between white clover and rhizobia

Both white clover and rhizobia were locally adapted, although effects of local adaptation were

more frequent and stronger for rhizobia. One explanation for this asymmetry in local adaptation

is stronger selection on rhizobia compared to white clover. *Rhizobium* abundance and fitness is

higher when cooperating with plant hosts rather than living in the soil as a saprophyte (Denison

& Kiers, 2004). In contrast, white clover's mutualism with rhizobia is facultative and dependent

on the N environment, and thus selection to maintain the mutualism may be weaker (D. E. Reid

et al., 2011). Another explanation is that the extensive gene flow among white clover populations

within Toronto (Santangelo et al., 2022) and nearby cities (Johnson et al., 2018) could counteract

the effects of local adaptation (Gomulkiewicz et al., 2007; Thompson, 2005). The population

structure for rhizobia has not been characterized in this system, but a separate study found that

populations of *R. leguminosarum* were genetically differentiated at spatial scales from ~4 km to

~350 km (Van Cauwenberghe et al., 2014), which is within the range of our urbanization

gradient (~50 km). Moreover, Van Cauwenberghe *et al.* (2016) observed genetic differentiation

and associated selection mosaics contributed to local adaptation between rhizobia and their host

plants. Evaluating the population structure for rhizobia is an important next step to understand

the balance of selection and gene flow and its role in shaping local adaptation in our system

(Gomulkiewicz et al., 2007; Hoeksema, 2010; Thompson, 2005; Wadgymar et al., 2022).

379   While there was an overall pattern of local adaptation, there was large variation among

populations in the strength and direction of local adaptation and even maladaptation. This

population $\times$ microbiome treatment interaction was driven by changes in the rank order of

populations between treatments for aboveground biomass, nodule density, and fixing nodule

density. Changes in rank order indicate that the relative-fitness of populations differ among

environments (Wade, 2007), which can maintain genetic variation among populations when a

single genotype is not the most fit in all environments (Bell et al., 2021; Christiansen, 1975;

Dempster, 1955; Gomulkiewicz & Kirkpatrick, 1992). These tradeoffs could have contributed to

the observed locally-adapted and locally-maladapted populations in our study.

Microbe-mediated adaptation

By sequentially examining how the microbiome inoculants could contribute to local adaptation,

we determined that *Rhizobium* evolution - not changes in *Rhizobium* abundance or microbiome

community composition - likely explains local adaptation in white clover and rhizobia. Fitness

responses of white clover and rhizobia were unrelated to *Rhizobium* abundance in the root

endosphere. As only *R. leguminosarum* can initiate nodulation on white clover (Andrews &

Andrews, 2017), our results suggest that nodulation depended on having the 'right' *Rhizobium*

partner (i.e., genotype), which is consistent with the findings from other studies (Burghardt et

al., 2022; Heath, 2010; Heath & Tiffin, 2007; Van Cauwenberghe et al., 2016). We found no

relationships between either microbiome inoculant dissimilarity or root endosphere composition

and local adaptation, indicating that the broader bacterial microbiome did not affect local

adaptation. These findings contrast with recent studies that demonstrated adaptation to local

microbiomes increases fitness (O'Brien et al., 2020; O'Brien, Yu, et al., 2022), and that

dissimilarity between local and nonlocal inoculants modulates fitness responses (O'Brien,

Laurich, et al., 2022). Importantly, rhizobia density in inoculants could also confound nodulation

(Batstone et al., 2020; Burghardt et al., 2022), but our microbiome inoculants had similar

densities of *Rhizobium* (Figure S10).

Coevolution or correlated local adaptation?

We detected a positive correlation in the strength of local adaptation between white clover and

rhizobia, which is consistent with interspecific coevolution between mutualists. While our results

are consistent with a history of coadaptation resulting from coevolution, we note several

qualifications with this interpretation. First, this correlation was only detected between

aboveground biomass and nodule density under N addition, so whether or not a signature of

coevolution is detected depends on the traits measured and the environment. Second, the

presence or absence of a correlation does not necessarily implicate coevolution (Janzen, 1980;

Nuismer et al., 2010). For example, spurious correlations can evolve in the absence of reciprocal

biotic selection if only one of the interacting species is evolving or abiotic factors favour

matching trait optima (Gomulkiewicz et al., 2007; Nuismer et al., 2010; Thompson, 2005).

Alternatively, the absence of a correlation can arise despite ongoing coevolution because traits

could be mismatched (Gomulkiewicz et al., 2007; Nuismer et al., 2010; Thompson, 2005).

Further investigation of potential coevolution between white clover and rhizobia in urban

environments requires characterizing selection acting on both white clover and rhizobia, and

ideally genomic signatures of selection on genes associated with the mutualism (Batstone,

Burghardt, et al., 2022; Batstone, Lindgren, et al., 2022; Epstein et al., 2023; Gomulkiewicz et

al., 2007; Hoeksema, 2010).

Conclusion

Our experiment contributes to the expanding literature on urban evolutionary ecology. Previous

studies of mutualisms have predominantly focused on plant-pollinator interactions and

adaptation for only one of the interacting species (Irwin *et al.* 2018; Theodorou *et al.* 2018;

Rivkin *et al.* 2020; but see Brans *et al.* 2022). Moreover, studies evaluating local adaptation in

urban environments have focused primarily on adaptation to abiotic factors, such as temperature

(Brans et al., 2017; Martin et al., 2021) and pollutants (N. M. Reid et al., 2016; Weston et al.,

2021). Anthropogenic impacts on legume-rhizobia symbioses have been evaluated in both

agricultural (Lau et al., 2022; Weese et al., 2015) and more recently in urban (Forrester &

Ashman, 2018; Regus et al., 2017; Wendlandt et al., 2022) environments. Nitrogen addition can

lead to the evolution of less-cooperative rhizobia (Lau et al., 2022; Regus et al., 2017; Weese et

al., 2015) but legumes can still maintain beneficial associations with rhizobia despite N addition

and deposition (Forrester & Ashman, 2018; Wendlandt et al., 2022). Our results show that

species can be locally adapted in urban environments, and that the mutualism between white

clover and rhizobia can persist despite increased N (Murray☐Stoker & Johnson, 2021). As plant-

microbe interactions are essential for ecosystem functioning (Bardgett & van der Putten, 2014;

van der Heijden et al., 2008), it is necessary to understand how microbiomes contribute to

adaptation and ecosystem processes in anthropogenically-disturbed environments (Nugent &

Allison, 2022).

Acknowledgements

We thank Elana Maria, Oliva Toth, Samer El-Galmady, Wendy Van Drunen, Hind Emad, Kelly

Murray-Stoker, Zain Nassrullah, and Narindra Persaud for assistance with fieldwork, Elana

Maria and Alanah Joyce for help with lab work, and Joan Lee, Brenda Pitton, and the UTM

Facilities staff for experiment setup. Comments from Megan Frederickson, John Stinchcombe,

Sophie Breitbart, Lucas Albano, and Aude Caizergues greatly improved the manuscript. This

work was funded by an NSERC Discovery Grant, Canada Research Chair, and E.W.R. Steacie

Fellowship to M. T. J. Johnson and an American Society of Naturalists Student Research Award

to D. Murray-Stoker.

Author Contributions

DMS and MTJJ designed the experiment. DMS collected and analyzed the data, with input from

MTJJ. MTJJ and DMS secured the funding. DMS led the writing of the manuscript, with support

from MTJJ.

Conflict of Interest Statement

The authors declare no conflict of interest.

Data Availability Statement

Data and code are available on Zenodo (10.5281/zenodo.8253951). Sequences are available at

NCBI under the BioProject accession PRJNA1005557.

References

Andrews, M., & Andrews, M. E. (2017). Specificity in legume-rhizobia symbioses. *International*

Journal of Molecular Sciences, *18*(4), 705. https://doi.org/10.3390/ijms18040705

Bardgett, R. D., & van der Putten, W. H. (2014). Belowground biodiversity and ecosystem

functioning. *Nature*, *515*(7528), Article 7528. https://doi.org/10.1038/nature13855

Batstone, R. T., Burghardt, L. T., & Heath, K. D. (2022). Phenotypic and genomic signatures of

interspecies cooperation and conflict in naturally occurring isolates of a model plant

symbiont. *Proceedings of the Royal Society B: Biological Sciences*, *289*(1978),

20220477. https://doi.org/10.1098/rspb.2022.0477

Batstone, R. T., Dutton, E. M., Wang, D., Yang, M., & Frederickson, M. E. (2017). The

evolution of symbiont preference traits in the model legume *Medicago truncatula*. *New*

Phytologist, *213*(4), Article 4. https://doi.org/10.1111/nph.14308

Batstone, R. T., Lindgren, H., Allsup, C. M., Goralka, L. A., Riley, A. B., Grillo, M. A.,

Marshall-Colon, A., & Heath, K. D. (2022). Genome-wide association studies across

environmental and genetic contexts reveal complex genetic architecture of symbiotic

extended phenotypes. *mBio*, *0*(0), e01823-22. https://doi.org/10.1128/mbio.01823-22

Batstone, R. T., Peters, M. A. E., Simonsen, A. K., Stinchcombe, J. R., & Frederickson, M. E.

(2020). Environmental variation impacts trait expression and selection in the legume–

rhizobium symbiosis. *American Journal of Botany*, *107*(2), Article 2.

https://doi.org/10.1002/ajb2.1432

Bell, D. A., Kovach, R. P., Robinson, Z. L., Whiteley, A. R., & Reed, T. E. (2021). The

ecological causes and consequences of hard and soft selection. *Ecology Letters*, *24*(7),

1505–1521. https://doi.org/10.1111/ele.13754

Berner, D., & Amrhein, V. (2022). Why and how we should join the shift from significance

testing to estimation. *Journal of Evolutionary Biology*, *35*(6), 777–787.

https://doi.org/10.1111/jeb.14009

Blanquart, F., Kaltz, O., Nuismer, S. L., & Gandon, S. (2013). A practical guide to measuring

local adaptation. *Ecology Letters*, *16*(9), Article 9. https://doi.org/10.1111/ele.12150

Brans, K. I., Jansen, M., Vanoverbeke, J., Tüzün, N., Stoks, R., & De Meester, L. (2017). The

heat is on: Genetic adaptation to urbanization mediated by thermal tolerance and body

size. *Global Change Biology*, *23*(12), 5218–5227. https://doi.org/10.1111/gcb.13784

Brans, K. I., Tüzün, N., Sentis, A., De Meester, L., & Stoks, R. (2022). Cryptic eco-evolutionary

feedback in the city: Urban evolution of prey dampens the effect of urban evolution of

the predator. *Journal of Animal Ecology*, *91*(3), 514–526. https://doi.org/10.1111/1365-

2656.13601

Bronstein, J. L. (1994). Conditional outcomes in mutualistic interactions. *Trends in Ecology and*

Evolution, *9*, 214–217.

Burdon, J. J. (1983). *Trifolium repens* L. *Journal of Ecology*, *71*(1), Article 1.

https://doi.org/10.2307/2259979

Burghardt, L. T., Epstein, B., Hoge, M., Trujillo, D. I., & Tiffin, P. (2022). Host-associated

rhizobial fitness: Dependence on nitrogen, density, community complexity, and legume

genotype. *Applied and Environmental Microbiology*, *88*(15), e00526-22.

https://doi.org/10.1128/aem.00526-22

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.

(2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature*

Methods, *13*(7), Article 7. https://doi.org/10.1038/nmeth.3869

Christiansen, F. B. (1975). Hard and soft selection in a subdivided population. *The American*

Naturalist, *109*(965), Article 965. https://doi.org/10.1086/282970

Cockerham, C. C. (1963). Estimation of genetic variances. *Statistical Genetics and Plant*

Breeding, *982*, 53–94.

Cole, J. R., Wang, Q., Fish, J. A., Chai, B., McGarrell, D. M., Sun, Y., Brown, C. T., Porras-

Alfaro, A., Kuske, C. R., & Tiedje, J. M. (2014). Ribosomal Database Project: Data and

tools for high throughput rRNA analysis. *Nucleic Acids Research*, *42*(D1), D633–D642.

https://doi.org/10.1093/nar/gkt1244

Delgado-Baquerizo, M., Eldridge, D. J., Liu, Y.-R., Sokoya, B., Wang, J.-T., Hu, H.-W., He, J.-

Z., Bastida, F., Moreno, J. L., Bamigboye, A. R., Blanco-Pastor, J. L., Cano-Díaz, C.,

Illán, J. G., Makhalanyane, T. P., Siebe, C., Trivedi, P., Zaady, E., Verma, J. P., Wang,

L., … Fierer, N. (2021). Global homogenization of the structure and function in the soil

microbiome of urban greenspaces. *Science Advances*, *7*(28), Article 28.

https://doi.org/10.1126/sciadv.abg5809

Dempster, E. R. (1955). Maintenance of genetic heterogeneity. *Cold Spring Harbor Symposia on*

Quantitative Biology, *20*, 25–32. https://doi.org/10.1101/SQB.1955.020.01.005

Denison, R. F., & Kiers, E. T. (2004). Lifestyle alternatives for rhizobia: Mutualism, parasitism,

and forgoing symbiosis. *FEMS Microbiology Letters*, *237*(2), Article 2.

https://doi.org/10.1111/j.1574-6968.2004.tb09695.x

Diamond, S. E., & Martin, R. A. (2021). Evolution in cities. *Annual Review of Ecology,*

Evolution, and Systematics, *52*(1), 519–540. https://doi.org/10.1146/annurev-ecolsys-

012021-021402

Epp Schmidt, D. J., Pouyat, R., Szlavecz, K., Setälä, H., Kotze, D. J., Yesilonis, I, Cilliers, S.,

Hornung, E., Dombos, M., & Yarwood, S. A. (2017). Urbanization erodes

ectomycorrhizal fungal diversity and may cause microbial communities to converge.

Nature Ecology and Evolution, *1*(5), Article 5. https://doi.org/10.1038/s41559-017-0123

Epstein, B., Burghardt, L. T., Heath, K. D., Grillo, M. A., Kostanecki, A., Hämälä, T., Young, N.

D., & Tiffin, P. (2023). Combining GWAS and population genomic analyses to

characterize coevolution in a legume-rhizobia symbiosis. *Molecular Ecology*, *32*(14),

3798–3811. https://doi.org/10.1111/mec.16602

Faeth, S. H., Bang, C., & Saari, S. (2011). Urban biodiversity: Patterns and mechanisms. *Annals*

of the New York Academy of Sciences, *1223*(1), 69–81. https://doi.org/10.1111/j.1749-

6632.2010.05925.x

Forrester, N. J., & Ashman, T.-L. (2018). Nitrogen fertilization differentially enhances

nodulation and host growth of two invasive legume species in an urban environment.

Journal of Urban Ecology, *4*(1), Article 1. https://doi.org/10.1093/jue/juy021

Gelman, A., & Hennig, C. (2017). Beyond subjective and objective in statistics. *Journal of the*

Royal Statistical Society. Series A (Statistics in Society), *180*(4), 967–1033.

Gomulkiewicz, R., Drown, D. M., Dybdahl, M. F., Godsoe, W., Nuismer, S. L., Pepin, K. M.,

Ridenhour, B. J., Smith, C. I., & Yoder, J. B. (2007). Dos and don'ts of testing the

geographic mosaic theory of coevolution. *Heredity*, *98*(5), Article 5.

https://doi.org/10.1038/sj.hdy.6800949

Gomulkiewicz, R., & Kirkpatrick, M. (1992). Quantitative genetics and the evolution of reaction

norms. *Evolution*, *46*(2), 390–411. https://doi.org/10.1111/j.1558-5646.1992.tb02047.x

Grimm, N. B., Faeth, S. H., Golubiewski, N. E., Redman, C. L., Wu, J., Bai, X., & Briggs, J. M.

(2008). Global change and the ecology of cities. *Science*, *319*(5864), 756–760.

https://doi.org/10.1126/science.1150195

Heath, K. D. (2010). Intergenomic epistasis and coevolutionary constraint in plants and rhizobia.

Evolution, *64*, 1446–1458. https://doi.org/10.1111/j.1558-5646.2009.00913.x

Heath, K. D., & Stinchcombe, J. R. (2014). Explaining mutualism variation: A new evolutionary

paradox? *Evolution*, *68*(2), Article 2. https://doi.org/10.1111/evo.12292

Heath, K. D., Stock, A. J., & Stinchcombe, J. R. (2010). Mutualism variation in the nodulation

response to nitrate. *Journal of Evolutionary Biology*, *23*(11), Article 11.

https://doi.org/10.1111/j.1420-9101.2010.02092.x

Heath, K. D., & Tiffin, P. (2007). Context dependence in the coevolution of plant and rhizobial

mutualists. *Proceedings of the Royal Society B: Biological Sciences*, *274*(1620), Article

1620. https://doi.org/10.1098/rspb.2007.0495

Hoeksema, J. D. (2010). Ongoing coevolution in mycorrhizal interactions. *New Phytologist*,

187(2), Article 2. https://doi.org/10.1111/j.1469-8137.2010.03305.x

Irwin, R. E., Warren, P. S., & Adler, L. S. (2018). Phenotypic selection on floral traits in an

urban landscape. *Proceedings of the Royal Society B: Biological Sciences*, *285*(1884),

Article 1884. https://doi.org/10.1098/rspb.2018.1239

Janzen, D. H. (1980). When is it coevolution? *Evolution*, *34*(3), Article 3.

https://doi.org/10.1111/j.1558-5646.1980.tb04849.x

Johnson, M. T. J., & Munshi-South, J. (2017). Evolution of life in urban environments. *Science*,

358(6363), Article 6363. https://doi.org/10.1126/science.aam8327

Johnson, M. T. J., Prashad, C. M., Lavoignat, M., & Saini, H. S. (2018). Contrasting the effects

of natural selection, genetic drift and gene flow on urban evolution in white clover

(*Trifolium repens*). *Proceedings of the Royal Society B: Biological Sciences*, *285*(1883),

20181019. https://doi.org/10.1098/rspb.2018.1019

Kiers, E. T., Palmer, T. M., Ives, A. R., Bruno, J. F., & Bronstein, J. L. (2010). Mutualisms in a

changing world: An evolutionary perspective. *Ecology Letters*, *13*(12), Article 12.

https://doi.org/10.1111/j.1461-0248.2010.01538.x

Lau, J. A., Hammond, M. D., Schmidt, J. E., Weese, D. J., Yang, W. H., & Heath, K. D. (2022).

Contemporary evolution rivals the effects of rhizobium presence on community and

ecosystem properties in experimental mesocosms. *Oecologia*, *200*(1), 133–143.

https://doi.org/10.1007/s00442-022-05253-1

Martin, R. A., Chick, L. D., Garvin, M. L., & Diamond, S. E. (2021). In a nutshell, a reciprocal

transplant experiment reveals local adaptation and fitness trade-offs in response to urban

evolution in an acorn-dwelling ant. *Evolution*, *75*(4), 876–887.

https://doi.org/10.1111/evo.14191

McKinney, M. L. (2006). Urbanization as a major cause of biotic homogenization. *Biological*

Conservation, *127*(3), 247–260. https://doi.org/10.1016/j.biocon.2005.09.005

McKinney, M. L. (2008). Effects of urbanization on species richness: A review of plants and

animals. *Urban Ecosystems*, *11*(2), 161–176. https://doi.org/10.1007/s11252-007-0045-4

McShane, B. B., Gal, D., Gelman, A., Robert, C., & Tackett, J. L. (2019). Abandon statistical

significance. *The American Statistician*, *73*(sup1), Article sup1.

Muir, W., Nyquist, W. E., & Xu, S. (1992). Alternative partitioning of the genotype-by-

environment interaction. *Theoretical and Applied Genetics*, *84*(1–2), 193–200.

https://doi.org/10.1007/BF00224000

Murray☐Stoker, D., & Johnson, M. T. J. (2021). Ecological consequences of urbanization on a

legume–rhizobia mutualism. *Oikos*, *130*(10), 1750–1761.

https://doi.org/10.1111/oik.08341

Nugent, A., & Allison, S. D. (2022). A framework for soil microbial ecology in urban

ecosystems. *Ecosphere*, *13*(3), e3968. https://doi.org/10.1002/ecs2.3968

Nuismer, S. L., Gomulkiewicz, R., & Ridenhour, B. J. (2010). When is correlation coevolution?

The American Naturalist, *175*(5), Article 5. https://doi.org/10.1086/651591

Nuismer, S. L., Thompson, J. N., & Gomulkiewicz, R. (1999). Gene flow and geographically

structured coevolution. *Proceedings of the Royal Society B: Biological Sciences*,

266(1419), Article 1419. https://doi.org/10.1098/rspb.1999.0679

O'Brien, A., Laurich, J., & Frederickson, M. E. (2022). *Having the "right" microbiome matters*

for host trait expression and the strength of mutualism between duckweeds and microbes.

bioRxiv preprint. https://doi.org/10.1101/2022.02.10.479958

O'Brien, A. M., Laurich, J., Lash, E., & Frederickson, M. E. (2020). Mutualistic outcomes across

plant populations, microbes, and environments in the duckweed *Lemna minor*. *Microbial*

Ecology. https://doi.org/10.1007/s00248-019-01452-1

O'Brien, A. M., Yu, Z. H., Pencer, C., Frederickson, M. E., LeFevre, G. H., & Passeport, E.

(2022). Harnessing plant-microbiome interactions for bioremediation across a freshwater

urbanization gradient. *Water Research*, *223*, 118926.

https://doi.org/10.1016/j.watres.2022.118926

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,

O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H.

(2020). *vegan: Community Ecology Package* (2.5-7) [Computer software].

https://CRAN.R-project.org/package=vegan

Pickett, S. T. A., Cadenasso, M. L., Grove, J. M., Boone, C. G., Groffman, P. M., Irwin, E.,

Kaushal, S. S., Marshall, V., McGrath, B. P., Nilon, C. H., Pouyat, R. V., Szlavecz, K.,

Troy, A., & Warren, P. (2011). Urban ecological systems: Scientific foundations and a

decade of progress. *Journal of Environmental Management*, *92*(3), 331–362.

https://doi.org/10.1016/j.jenvman.2010.08.022

Pickett, S. T. A., Cadenasso, M. L., Grove, J. M., Nilon, C. H., Pouyat, R. V., Zipperer, W. C., &

Costanza, R. (2001). Urban ecological systems: Linking terrestrial ecological, physical,

and socioeconomic components of metropolitan areas. *Annual Review of Ecology and*

Systematics, *32*, 127–157.

Poole, P., Ramachandran, V., & Terpolilli, J. (2018). Rhizobia: From saprophytes to

endosymbionts. *Nature Reviews Microbiology*, *16*(5), Article 5.

https://doi.org/10.1038/nrmicro.2017.171

R Core Team. (2022). *R: a language and environment for statistical computing* (4.2.2)

[Computer software]. . R Foundation for Statistical Computing. https://www.R-

project.org/

Regus, J. U., Wendlandt, C. E., Bantay, R. M., Gano-Cohen, K. A., Gleason, N. J., Hollowell, A.

C., O'Neill, M. R., Shahin, K. K., & Sachs, J. L. (2017). Nitrogen deposition decreases

the benefits of symbiosis in a native legume. *Plant and Soil*, *414*(1–2), Article 1–2.

https://doi.org/10.1007/s11104-016-3114-8

Reid, D. E., Ferguson, B. J., Hayashi, S., Lin, Y.-H., & Gresshoff, P. M. (2011). Molecular

mechanisms controlling legume autoregulation of nodulation. *Annals of Botany*, *108*(5),

789–795. https://doi.org/10.1093/aob/mcr205

Reid, N. M., Proestou, D. A., Clark, B. W., Warren, W. C., Colbourne, J. K., Shaw, J. R.,

Karchner, S. I., Hahn, M. E., Nacci, D., Oleksiak, M. F., Crawford, D. L., & Whitehead,

A. (2016). The genomic landscape of rapid repeated evolutionary adaptation to toxic

pollution in wild fish. *Science*, *354*(6317), Article 6317.

https://doi.org/10.1126/science.aah4993

Rivkin, L. R., Nhan, V. J., Weis, A. E., & Johnson, M. T. J. (2020). Variation in pollinator-

mediated plant reproduction across an urbanization gradient. *Oecologia*, *192*(4), 1073–

1083. https://doi.org/10.1007/s00442-020-04621-z

Rosier, C. L., Polson, S. W., D'Amico, V., Kan, J., & Trammell, T. L. E. (2021). Urbanization

pressures alter tree rhizosphere microbiomes. *Scientific Reports*, *11*(1), 9447.

https://doi.org/10.1038/s41598-021-88839-8

RStudio Team. (2022). *RStudio: Integrated development for R* (2021.09.0) [Computer software].

RStudio, Inc. https://rstudio.com/

Santangelo, J. S., Ness, R. W., Cohan, B., Fitzpatrick, C. R., Innes, S. G., Koch, S., Miles, L. S.,

Munim, S., Peres-Neto, P. R., Prashad, C., Tong, A. T., Aguirre, W. E., Akinwole, P. O.,

Alberti, M., Álvarez, J., Anderson, J. T., Anderson, J. J., Ando, Y., Andrew, N. R., …

Johnson, M. T. J. (2022). Global urban environmental change drives adaptation in white

clover. *Science*, *375*(6586), 1275–1281. https://doi.org/10.1126/science.abk0989

Shantz, A. A., Lemoine, N. P., & Burkepile, D. E. (2016). Nutrient loading alters the

performance of key nutrient exchange mutualisms. *Ecology Letters*, *19*(1), Article 1.

https://doi.org/10.1111/ele.12538

Stevens, C. J., David, T. I., & Storkey, J. (2018). Atmospheric nitrogen deposition in terrestrial

ecosystems: Its impact on plant communities and consequences across trophic levels.

Functional Ecology, *32*(7), Article 7. https://doi.org/10.1111/1365-2435.13063

Theodorou, P., Radzevičiūtė, R., Kahnt, B., Soro, A., Grosse, I., & Paxton, R. J. (2018).

Genome-wide single nucleotide polymorphism scan suggests adaptation to urbanization

in an important pollinator, the red-tailed bumblebee (*Bombus lapidarius* L.). *Proceedings*

of the Royal Society B: Biological Sciences, *285*(1877), Article 1877.

https://doi.org/10.1098/rspb.2017.2806

Thompson, J. N. (1988). Variation in interspecific interactions. *Annual Review of Ecology and*

Systematics, *19*(1), Article 1. https://doi.org/10.1146/annurev.es.19.110188.000433

Thompson, J. N. (2005). *The Geographic Mosaic of Coevolution*. University of Chicago Press.

Udvardi, M., & Poole, P. S. (2013). Transport and metabolism in legume-rhizobia symbioses.

Annual Review of Plant Biology, *64*(1), Article 1. https://doi.org/10.1146/annurev-

arplant-050312-120235

Van Cauwenberghe, J., Verstraete, B., Lemaire, B., Lievens, B., Michiels, J., & Honnay, O.

(2014). Population structure of root nodulating *Rhizobium leguminosarum* in *Vicia*

cracca populations at local to regional geographic scales. *Systematic and Applied*

Microbiology, *37*(8), 613–621. https://doi.org/10.1016/j.syapm.2014.08.002

Van Cauwenberghe, J., Visch, W., Michiels, J., & Honnay, O. (2016). Selection mosaics

differentiate *Rhizobium*-host plant interactions across different nitrogen environments.

Oikos, *125*(12), Article 12. https://doi.org/10.1111/oik.02952

van der Heijden, M. G. A., Bardgett, R. D., & van Straalen, N. M. (2008). The unseen majority:

Soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems.

Ecology Letters, *11*(3), Article 3. https://doi.org/10.1111/j.1461-0248.2007.01139.x

Wade, M. J. (2007). The co-evolutionary genetics of ecological communities. *Nature Reviews*

Genetics, *8*(3), Article 3. https://doi.org/10.1038/nrg2031

Wadgymar, S. M., DeMarche, M. L., Josephs, E. B., Sheth, S. N., & Anderson, J. T. (2022).

Local adaptation: Causal agents of selection and adaptive trait divergence. *Annual Review*

of Ecology, Evolution, and Systematics, *53*(1), 87–111. https://doi.org/10.1146/annurev-

ecolsys-012722-035231

Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R. (2007). Naïve bayesian classifier for rapid

assignment of rRNA sequences into the new bacterial taxonomy. *Applied and*

Environmental Microbiology, *73*(16), Article 16. https://doi.org/10.1128/AEM.00062-07

Wasserstein, R. L., & Lazar, N. A. (2016). The ASA statement on p-values: Context, process,

and purpose. *The American Statistician*, *70*(2), Article 2.

https://doi.org/10.1080/00031305.2016.1154108

Wasserstein, R. L., Schirm, A. L., & Lazar, N. A. (2019). Moving to a world beyond "p < 0.05."

The American Statistician, *73*(sup1), Article sup1.

https://doi.org/10.1080/00031305.2019.1583913

Weese, D. J., Heath, K. D., Dentinger, B. T. M., & Lau, J. A. (2015). Long-term nitrogen

addition causes the evolution of less-cooperative mutualists. *Evolution*, *69*(3), Article 3.

https://doi.org/10.1111/evo.12594

Wendlandt, C. E., Gano-Cohen, K. A., Stokes, P. J. N., Jonnala, B. N. R., Zomorrodian, A. J.,

Al-Moussawi, K., & Sachs, J. L. (2022). Wild legumes maintain beneficial soil rhizobia

populations despite decades of nitrogen deposition. *Oecologia*, *198*(2), 419–430.

https://doi.org/10.1007/s00442-022-05116-9

Weston, L. M., Mattingly, K. Z., Day, C. T. C., & Hovick, S. M. (2021). Potential local

adaptation in populations of invasive reed canary grass (*Phalaris arundinacea*) across an

urbanization gradient. *Ecology and Evolution*, *11*(16), 11457–11476.

https://doi.org/10.1002/ece3.7938

Table 1: Summary of the linear mixed-effects models testing the effects of microbiome (local, nonlocal$_{Rural}$, nonlocal$_{Urban}$), nitrogen (ambient N, N addition), and the two-way interaction on measure of plant and rhizobia fitness. We provide the numerator degrees-of-freedom (NumDF), denominator degrees-of-freedom (estimated using the Kenward-Roger method (DenDF; Kenward & Roger 1997), F-statistics calculated from type III sums-of-squares, P-values, and the effect size (partial eta-squared, η^2_P) for each fixed effect term in the ANOVAs. We also provide the χ^2 statistic, P-values from likelihood ratio tests, and the effect size (intra-class correlation coefficient, ICC) for each random effect term, with random effect terms and statistics in italics. The variance explained by the fixed effects ($R^2_{Marginal}$) and combined fixed and random effects ($R^2_{Conditional}$) is provided for each model.

Term	NumDF	DenDF	F/χ^2	P-value	η^2_P/ICC
Aboveground Biomass					
($R^2_{Marginal}$ = 0.233, $R^2_{Conditional}$ = 0.397)					
Microbiome	2	54.911	3.243	0.047	0.106
Nitrogen	1	27.790	188.399	< 0.001	0.871
Microbiome × Nitrogen	2	742.517	0.470	0.625	0.001
Microbiome × Population			*4.951*	*0.026*	*0.040*
Nitrogen × Population			*3.901*	*0.048*	*0.035*
Population			*10.311*	*0.001*	*0.126*
Block			*3.575*	*0.059*	*0.011*
Belowground Biomass					
($R^2_{Marginal}$ = 0.292 $R^2_{Conditional}$ = 0.376)					
Microbiome	2	802.275	3.338	0.036	0.008
Nitrogen	1	801.735	384.570	< 0.001	0.324
Microbiome × Nitrogen	2	805.884	0.292	0.747	0.001
Population			*46.239*	*< 0.001*	*0.104*
Block			*4.791*	*0.029*	*0.015*
Nodule Density					
($R^2_{Marginal}$ = 0.061, $R^2_{Conditional}$ = 0.261)					
Microbiome	2	56.305	9.257	< 0.001	0.247
Nitrogen	1	27.932	13.328	0.001	0.323
Microbiome × Nitrogen	2	55.584	1.057	0.354	0.037
Microbiome × Population			*1.614*	*0.204*	*0.053*
Nitrogen × Population			*0.019*	*0.889*	*0.004*
Microbiome × Nitrogen × Population			*6.376*	*0.012*	*0.105*
Population			*1.899*	*0.168*	*0.047*
Block			*0.390*	*0.533*	*0.003*
Fixing Nodule Density					
($R^2_{Marginal}$ = 0.043, $R^2_{Conditional}$ = 0.198)					
Microbiome	2	55.390	10.067	< 0.001	0.267
Nitrogen	1	27.929	5.595	0.025	0.167
Microbiome × Nitrogen	2	741.058	0.442	0.643	0.001
Microbiome × Population			*8.159*	*0.004*	*0.062*
Nitrogen × Population			*5.201*	*0.023*	*0.045*
Population			*0.787*	*0.375*	*0.026*
Block			*13.511*	*< 0.001*	*0.029*

725 **Figures**

726
Figure 1: Map of the urbanization gradient in the Greater Toronto Area, Ontario, Canada (A),
with the location of the 30 focal populations indicated by yellow circles. We also show the
location of the 5 rural (blue circles) and 5 urban (pink circles) sites from which nonlocal
inoculants were sourced. Urban and suburban areas in gray, nonurban agricultural and forested
areas in green, and Lake Ontario in light blue. Map tiles by Stamen Design, under CC BY 3.0.
Data by OpenStreetMap, under ODbL. We also show a diagram of the experimental design (B).
Each of the 30 focal populations (yellow circles on the map) received 3 microbiome treatments
(local, nonlocal$_{Urban}$, and nonlocal$_{Rural}$). Additionally, each microbiome treatment received 2
nitrogen treatments (no addition = ambient N, 10 mM KNO$_3$ fertilizer = N addition). The
factorial experiment was conducted in an outdoor common garden at the University of Toronto
Mississauga campus.

739
Figure 2: Treatment means ± 1 SE (left column) and reaction norms of population means (right
column) by microbiome and nitrogen treatments for aboveground biomass (A-B), belowground
biomass (C-D), nodule density (E-F), and fixing nodule density (G-H). Microbiome treatments
are indicated as nonlocal$_{Rural}$ (blue), nonlocal$_{Urban}$ (pink), and local (yellow); individual plots are
faceted by nitrogen treatment (ambient N and N addition). Inset text indicates the factors of
interest in the optimized model structure for the respective variable, with fixed effects in regular
font and random effects in italics. Influential factors and interactions were identified based on
effect sizes (η^2_P range = 0.106-0.871, ICC range = 0.035-0.126) and are in bold, with detailed
test statistics provided in Table 1.

749
750 **Figure 3:** Pairwise correlations between the local – nonlocal_Global local adaptation indices for
751 aboveground biomass, belowground biomass, nodule density, and fixing nodule density.
752 Separate correlations were calculated for each nitrogen treatment (ambient N = purple, N
753 addition = pink). Lines represent lines-of-best-fit (± 95% confidence interval). Inset text provides
754 the correlation coefficient (r) and P-value for each correlation.

755
Figure 4: Relationship between the global local adaptation index (local – nonlocal$_{Global}$, LA
Index$_{Global}$) and urbanization (standardized distance from the city centre; mean = 0, standard
deviation = 1) for each of the fitness estimates [aboveground biomass (A) belowground biomass
(B), nodule density (C), and fixing nodule density, (D)]. Lines represent the lines-of-best-fit (±
95% confidence interval). Inset text provides the slope parameter (β), standard error (SE), and P-
value for each linear regression. These conclusions are unchanged when other metrics of
urbanization (i.e., Human Influence Index and imperious surface cover) and indices of local
adaptation (i.e., local – nonlocal$_{Rural}$ and local – nonlocal$_{Urban}$) are used (Figures S4-S6).

Figure 5: The relationship between fitness estimates and *Rhizobium* abundance (left column) and local adaptation by pairwise microbiome dissimilarity (right column) for aboveground biomass (A and B), belowground biomass (C and D), nodule density (E and F), and fixing nodule density (G and H). Lines represent lines-of-best-fit (± 95% confidence interval), with separate lines for each nitrogen treatment (ambient N = purple, N addition = pink) for the *Rhizobium* abundance models. Inset text for the left column indicates the variables of interest in the model. Influential factors and interactions were identified based on effect sizes and are in bold (η^2_P range = 0.081-0.175). Inset text for the right column provides the slope parameter (β), standard error (SE), and P-value for each linear regression. Detailed test statistics are provided in Table S3 (panels A, C, E, G) and Table S5 (panels B, D, F, H).