Title: Mosaic of local adaptation between white clover and rhizobia along an urbanization gradient

Running Title: Mosaic of local adaptation

Authors: David Murray-Stoker[1,2,3,*] and Marc T. J. Johnson[2,3]

Affiliations
[1]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario M5S 3B2, Canada
[2]Department of Biology, University of Toronto Mississauga, Mississauga, Ontario, Canada L5L 1C6
[3]Centre for Urban Environments, University of Toronto Mississauga, Mississauga, Ontario, Canada L5L 1C6

David Murray-Stoker, ORCID https://orcid.org/0000-0002-4774-6948
Marc T. J. Johnson, ORCID http://orcid.org/0000-0001-9719-0522

Corresponding author: David Murray-Stoker (dstoker92@gmail.com)

Keywords: coevolution, legume-rhizobia interaction, local adaptation, mutualism, microbiome, urban evolution
Type: Letter
Word Count: Abstract = 150, Main Text = 4974
Number of References: 82
Number of Figures: 5
Number of Tables: 1

Author Contributions
DMS and MTJJ designed the experiment. DMS collected and analyzed the data, with input from MTJJ. MTJJ and DMS secured the funding. DMS led the writing of the manuscript, with support from MTJJ.

Data Availability Statement
Data and code are available on Zenodo (10.5281/zenodo.8253951). Sequences are available at NCBI under the BioProject accession PRJNA1005557.

Abstract

Urbanization alters many biotic and abiotic environmental factors, but the effects of urbanization on local adaptation and coevolution between species remains largely unstudied. Using a common garden experiment of 30 populations and 1080 plants, we tested for local adaptation in the mutualism between white clover (*Trifolium repens*) and rhizobia (*Rhizobium leguminosarum* symbiovar *trifolii*) along an urbanization gradient. Our results show a spatial mosaic in local adaptation, with stronger local adaptation for rhizobia than white clover. While nitrogen addition (N) strongly influenced plant biomass and nodule density, it did not consistently mediate patterns of local adaptation. The strength of local adaptation was positively related between white clover and rhizobia under N addition, suggesting that soil N mediates coadaptation between white clover and rhizobia. Local adaptation was not influenced by urbanization, indicating that while urbanization does influence the ecology of plant-microbe interactions, it does not disrupt local adaptation and coevolution among mutualists.

Keywords: coevolution, legume-rhizobia interaction, local adaptation, mutualism, microbiome, urban evolution

Introduction

Urbanization is a dominant driver of environmental change that frequently alters the ecological setting of ecosystems (Pickett *et al.* 2001; Santangelo *et al.* 2022). Abiotic changes to the environment commonly include increased impervious surface, habitat fragmentation and degradation, elevated temperature, higher pollution (e.g., air, water, light, noise), and increased nutrient deposition (Grimm *et al.* 2008; Pickett *et al.* 2011). In addition to urban-driven abiotic changes, biotic changes frequently include altered species diversity and community composition, reduced vegetation cover, increased abundance of non-native species, and reduced abundance

and diversity of native species (McKinney 2008; Faeth *et al.* 2011). Ecological changes to urban

environments have been found to cause evolution in a diverse array of plants and animals

(Johnson & Munshi-South 2017; Diamond & Martin 2021). Despite increasing evidence of

evolutionary consequences in response to urbanization, it remains to be seen if urbanization

affects the local adaptation and (co)evolution of species interactions.

Variation in both the biotic and abiotic environment can result in selection mosaics that

lead to locally adapted phenotypes (Thompson 2005). Numerous environmental factors can

impose natural selection that lead to local adaptation, whereby local populations have greater

fitness than nonlocal populations (Blanquart *et al.* 2013). Mutualisms are a useful focal

interaction to study local adaptation because the reciprocal fitness feedbacks among interacting

species should lead to fitness alignment in a given environment but still allow for variation

across environments (Nuismer *et al.* 1999; Hoeksema 2010; Heath & Stinchcombe 2014).

Variation in community composition across space and time can alter mutualistic species

interactions (Thompson 1988, 2005). Abiotic context can also affect mutualisms by altering the

physical (e.g., temperature; Kiers *et al.* 2010) or resource (e.g., nutrient and water availability;

Bronstein 1994; Shantz *et al.* 2016) environment in which species interact. Moreover, biotic and

abiotic contexts are likely acting in concert to affect the outcome of mutualistic interactions

(Heath & Tiffin 2007; Heath *et al.* 2010; Van Cauwenberghe *et al.* 2016; Burghardt *et al.* 2022).

Because urbanization alters both the biotic and abiotic environment simultaneously across spatial

and temporal scales (Johnson & Munshi-South 2017; Diamond & Martin 2021), urban

environments have the potential to alter selective landscapes, local adaptation, and potentially

coevolution between interacting species.

Mutualistic interactions between legumes and rhizobia are a useful model for evaluating

evolutionary ecology in urban environments. In this mutualism, rhizobia fix atmospheric N_2 into

an accessible form of nitrogen (N) for the plant (NH_4^+) in exchange for photosynthate and

hosting within nodules provided by the plant (Poole *et al.* 2018). Ecological changes in urban

environments, particularly increased N deposition and enrichment (Grimm *et al.* 2008; Stevens *et*

al. 2018), could have consequences for legume-rhizobium local adaptation by disrupting the

economy of the resource exchange: direct acquisition of N is 'cheaper' for the host legume,

which can select for less cooperative rhizobia (Weese *et al.* 2015; Regus *et al.* 2017). In a

previous study along an urbanization gradient (Murray☐Stoker & Johnson 2021), we found that

urbanization altered the ecology of a legume-rhizobia mutualism. Specifically, urban plants

exhibited reduced investment in the mutualism and primarily acquired N through fixation except

at the urban and rural limits of the gradient. Soil N content was the nexus linking the effects of

urbanization to the mutualism (Murray☐Stoker & Johnson 2021). With this evidence, we wanted

to determine if the ecological changes driven by urbanization affected local adaptation of white

clover and its rhizobia mutualists.

Here we conducted a large, factorial field experiment to test the hypothesis that urban

environmental change drives local adaptation in the mutualism between white clover (*Trifolium*

repens) and rhizobia (*Rhizobium leguminosarum* symbiovar *trifolii*). We asked four specific

questions: **(Q1)** Are white clover and rhizobia locally adapted? **(Q2)** Does N addition mediate

the strength of local adaptation? **(Q3)** Is the strength of local adaptation in host plants related to

the strength of local adaptation in rhizobia, as expected if white clover and rhizobia are

coadapting? **(Q4)** How does the degree of local adaptation vary with urbanization? Results from

this study provide the first test of how urbanization affects the evolution and local adaptation of

interacting species, which provides insight into urban environments and whether they could be

disrupting coevolution.

Materials and Methods

Study System

We used the mutualism between white clover (*Trifolium repens* L., Fabaceae) and its facultative,

rhizobial symbiont (*Rhizobium leguminosarum* symbiovar *trifolii*) to evaluate if and how local

adaptation varied with urbanization. White clover is a perennial, herbaceous legume native to

Eurasia that is now globally distributed. White clover typically reproduces clonally through

stolons as well as through seed via obligately-outcrossed flowers (Burdon 1983). *Rhizobium*

leguminosarum symbiovar *trifolii* (hereafter rhizobia) is the primary rhizobial symbiont of white

clover (Andrews & Andrews 2017).

Biological Materials

White clover: The 30 focal white clover populations for this experiment were a subset of those

sampled by Murray-Stoker and Johnson (2021), with focal populations selected to span an

urbanization gradient in the Greater Toronto Area, Ontario, Canada (Figure 1A). To minimize

parental effects, we grew field-collected seeds in the greenhouse to produce outcrossed F_1

populations. All F_1 seeds were derived from a random panmictic cross among 5-10 parent plants

within a source population (detailed methods are provided in Appendix S1).

Soil inoculants: Soil for microbiome inoculants was collected following the methods described

in Murray-Stoker and Johnson (2021). Briefly, 5 soil cores were taken at each site at least 5 m

from the nearest white clover plant and other legumes (e.g., *Medicago lupulina*). Each soil core

was collected to a depth of 5 cm, with the top 2 cm plus any organic material removed.

Individual cores were stored in separate ziploc bags and stored at −80°C until subsequent

processing. Local microbiomes were paired to each of the 30 white clover populations in the

experiment, while the nonlocal$_{Urban}$ and nonlocal$_{Rural}$ microbiome inoculants were pooled from 5

populations at the urban or rural ends of the gradient, respectively (Figure 1A).

We used a local versus nonlocal microbiome inoculation approach (Blanquart *et al.* 2013;

Heath & Stinchcombe 2014) to test for local adaptation between white clover and rhizobia for

two principal reasons. First, this approach tests for overall differences in fitness between local

and nonlocal white clover or rhizobia populations, which can be further distinguished based on

the habitat of populations (i.e., urban vs. rural). In particular, increased N deposition in urban

environments can alter the dynamics of the mutualism between white clover and rhizobia by

reducing nodule densities and influencing the source of N used by white clover (Murray☐Stoker

& Johnson 2021). Additionally, urbanization leads to homogenization of the biotic and abiotic

environment (McKinney 2006; Santangelo *et al.* 2022), which might result in homogenized

composition of microbial communities (Delgado-Baquerizo *et al.* 2021; Rosier *et al.* 2021; but

see Epp Schmidt *et al.* 2017). Together, this evidence suggests that the degree of local adaptation

might depend on (1) the nonlocal inoculant representing rural or urban environments and (2)

similarity in microbiome composition between the local and either nonlocal microbiomes (i.e.,

nonlocal$_{Rural}$ and nonlocal$_{Urban}$). Second, by using microbiome inoculants instead of isolated

rhizobia strains, this approach also provided more ecological realism by allowing rhizobia and

other bacteria and fungi to establish in the soil and roots.

Experiment Overview

We tested for local adaptation between populations of white clover and rhizobia using a common

garden experiment (Figure 1B). We used the local versus nonlocal contrast (sensu sympatric vs.

allopatric; Blanquart *et al.* 2013), with local adaptation indicated when mean local fitness was

greater than mean nonlocal fitness. We crossed 30 white clover populations with 3 microbiome

treatments (local, nonlocal$_{Rural}$, and nonlocal$_{Urban}$) and 2 N treatments (no N addition = ambient

N, 10 mM KNO$_3$ fertilizer = N addition). Each treatment combination had 6 replicates per

population (30 populations × 3 microbiome treatments × 2 N treatments × 6 replicates = 1080

plants). The N addition treatment was representative of the soil N content measured at the focal

populations by Murray-Stoker and Johnson (2021) and was designed to reflect the impacts of

urbanization on soil N. We also had 6 control pots (3 ambient N, 3 N addition) that were not

planted or inoculated but did receive fertilizer additions, and these pots were used to evaluate

dispersal and establishment of bacteria at the experimental site.

Common Garden Experiment

We grew all of the plants outside in a common environment using an incomplete block design at

the University of Toronto Mississauga campus. In total, there were 9 blocks consisting of 120

plants; each block included all treatment combinations but not all population-by-treatment

combinations were represented in each block due to spatial constraints. Prior to planting, seeds

were scarified with sandpaper, surface sterilized (60 s in 70% ethanol, 60 s in 10% household

bleach, 90 s rinse in sterile water) and then germinated in the dark on dampened filter paper for

48 h at 20°C. Germinated seeds were planted in 1 L bleach-sterilized pots in a steam-sterilized (2

h of sterilization) mix of potting soil and sand mixture (2:3 V/V). The soil mixture was selected

to facilitate harvesting of all root biomass at the end of the experiment. We planted two

germinated seeds in each pot, and pots were thinned to one plant between weeks 2-3 of the

experiment.

Microbiome treatments were applied 1 week after planting by applying soil slurries. Soil

slurries for each microbiome treatment were prepared with 7.5 g fresh soil (thawed from –80°C

overnight at 4°C) in 50 mL sterile deionized water; nonlocal treatments were composed of equal

mixtures by mass from 5 populations within the respective habitat type. Slurries were hand

shaken for 60 s after preparation to mix and suspend the soil, with further re-suspension (hand

shaken for 30 s) during inoculation to maintain soil suspension. Each plant received 10 mL of its

respective microbiome inoculant. Weekly N addition treatments began in week 2, and we applied

bi-weekly N-free fertilizer following Batstone *et al.* (2017) starting in week 4 of the experiment.

Both the N-addition and N-free fertilizers were autoclave sterilized (60 min sterilization) prior to

application. Plants were watered daily as needed for the duration of the experiment. We ran the

experiment for 19 weeks (21 May–2 October 2021). Due to germination failure, herbivory, and

other sources of plant loss, our final plant count was 841.

Laboratory Processing

We measured multiple components of fitness for white clover (aboveground biomass and

belowground biomass) and rhizobia (total nodule density and fixing nodule density). We first

washed all plants of soil and debris and then split them into aboveground and belowground

components to facilitate fitness measurements. Aboveground components were placed in

envelopes, dried to constant mass (at least 72 h at 60°C), and aboveground biomass was weighed

to the nearest 1 mg. To measure rhizobia fitness components, a subset of roots was collected

from each plant by cutting the first three roots below the plant base and directly attached to the

stolon (following Murray☐Stoker & Johnson 2021), and we then measured the total root length

of all collected roots (mean ± standard error = 43.5 ± 0.58 cm, range = 2.1-128.7 cm). Nodule

density, an estimate of investment in the mutualism by white clover and rhizobia fitness, was

measured by visually-counting nodules from each measured root and standardizing by the total

root length to generate an estimate of nodule density per cm of root per plant (Murray☐Stoker &

Johnson 2021). We also calculated the density of fixing nodules, a measure of the effectiveness

of the mutualism, by counting the number of nodules with pink colouration (indicative of N

fixation; Udvardi & Poole 2013) and calculating density as described above. After estimating

nodule density and fixing nodule density, all belowground biomass (including the 3 excised

roots) was placed in an envelope, dried for 72 h at 60°C, and weighed to the nearest 0.1 mg. We

focused on biomass as a measure of plant fitness because white clover undergoes extensive

clonal growth as a perennial. We could not measure sexual fitness accurately because only a

fraction of plants flowered and we could not prevent interpopulation crosses.

DNA Extraction & Sequencing

To determine if white clover and rhizobia fitness responses were influenced by the broader

bacterial microbiome, we sequenced the bacterial microbiome for each of the microbiome

inoculants and a subset of roots from each of the experimental treatments. All 40 microbiome

inoculants (30 focal, 5 nonlocal$_{Rural}$, 5 nonlocal$_{Urban}$) were sequenced to evaluate variation in

bacterial community composition. A total of 60 root microbiomes (10 populations $\times$ 3

microbiome treatments $\times$ 2 N treatments = 60 plants) were sequenced to evaluate: (1) white

clover and rhizobia fitness responses as a function of *Rhizobium* abundance, (2) *Rhizobium*

abundance in response to experimental treatments, and (3) fitness of white clover and rhizobia in

response to soil and root bacterial microbiome composition. We included each treatment

combination for the same 6 populations (i.e., 10 populations $\times$ 6 treatment combinations = 60

samples). Protocols for microbiome DNA extraction are provided in the supplementary material

(Appendix S1).

Statistical Analyses

We provide detailed methods for all statistical analyses in the supplementary material (Appendix

S1). Briefly, we analyzed variation in fitness responses for white clover (aboveground biomass,

belowground biomass) and rhizobia (nodule density, fixing nodule density) in response to the

microbiome treatment (local, nonlocal$_{Rural}$, nonlocal$_{Urban}$), N treatment (ambient N, N addition),

and the two-way interaction using linear mixed-effects models with Type III sums-of-squares.

We also quantified the extent to which population $\times$ environment interactions were due to

changes in population variance and changes in rank order using Cockerhams's method

(Cockerham 1963; Muir *et al.* 1992). To test if there was a general signature of local adaptation,

we repeated the same procedure described above but with microbiome recoded as local and

nonlocal$_{Global}$ (i.e., nonlocal$_{Rural}$ and nonlocal$_{Urban}$ combined).

The strength of local adaptation was calculated using three indices: (1) local –

nonlocal$_{Global}$, (2) local – nonlocal$_{Rural}$, and (3) local – nonlocal$_{Urban.}$ These indices were

calculated separately for each of the four fitness response variables (i.e., aboveground biomass,

belowground biomass, nodule density, and fixing nodule density). Positive values were

indicative of local adaptation because they showed greater fitness in the local versus nonlocal

treatment. Local adaptation indices were calculated using the best linear unbiased predictors (i.e.,

BLUPs) from linear mixed-effects models. To test whether the strength of local adaptation was

related between interacting white clover and *Rhizobium* populations (i.e., a signature of

coadaptation), we calculated all pairwise Pearson correlations between the two fitness

components of white clover with the two fitness components of rhizobia. We calculated

correlations separately for each N treatment to examine if N addition mediated coadaptation.

To test how urbanization affects the strength of local adaptation, we regressed local

adaptation indices against measures of urbanization: distance from the city center, mean

impervious surface cover (ISC, 250 m radius around each population), and the Human Influence

Index (HII, 1 km radius around each population). We fitted separate regressions for each local

adaptation index and urbanization metric due to multicollinearity between variables (variance

inflation factors > 5–10). We scaled all predictor variables to a mean of 0 and standard deviation

of 1 prior to analysis to allow for direct comparisons across models.

Microbiome sequences were processed for analysis using the DADA2 pipeline (Callahan

et al. 2016), with taxonomy assigned to amplicon sequence variants using the Ribosomal

Database Project (Cole *et al.* 2014) and naïve Bayesian classifier (Wang *et al.* 2007)

implemented in DADA2. We regressed white clover and rhizobia fitness responses against

Rhizobium abundance (number of reads assigned to the genus *Rhizobium*), N treatment, and the

interaction using generalized linear models. We examined differences in *Rhizobium* abundance

in response to microbiome$_{Global}$ (local vs. nonlocal$_{Global}$), N (ambient N, N addition), and the

interaction using ANOVAs with Type III sums-of-squares. Finally, we examined the role of

microbiome composition on local adaptation. We tested if the strength of local adaptation

increased when microbiome inoculant composition was more dissimilar (pairwise Bray-Curtis

dissimilarity) using a linear regression. We also tested if local adaptation indices were related to

root bacterial composition. The effects of the microbiome treatment, N treatment, and their

interaction on root microbiome composition was fit and assessed using PERMANOVA with a

Bray-Curtis dissimilarity matrix (Oksanen *et al.* 2020). Relationships between root community

composition and local adaptation indices were quantified using the `envfit()` function (Oksanen

et al. 2020).

All statistical analyses were performed using R (version 4.2.2; R Core Team 2022) in the

RStudio environment (version 2022.07.2; RStudio Team 2022). All packages used for data

management, analysis, and figure creation are provided in the detailed statistical analyses in the

supplementary material (Appendix S1). We followed statistical best practices by focusing on

estimates, variation of the estimates, and effect sizes to interpret the results and identify the

strength of evidence (Wasserstein & Lazar 2016; Gelman & Hennig 2017; McShane *et al.* 2019;

Wasserstein *et al.* 2019; Berner & Amrhein 2022).

Results

Effects of microbiome and nitrogen treatments on fitness

We found evidence for local adaptation for both white clover and rhizobia, and the effects were

stronger for rhizobia than white clover and varied among populations (Table 1, Figure 2). White

clover demonstrated weak adaptation to local microbiomes for both aboveground biomass (P =

0.047, η^2_P = 0.106, Table 1, Figure 2) and belowground biomass (P = 0.036, η^2_P = 0.008, Table

1, Figure 2). Aboveground biomass was 6.4% and 12.8% greater in local versus nonlocal$_{Rural}$ and

nonlocal$_{Urban}$ microbiomes, respectively (contrasts in Table S1), and 60% of populations

demonstrated patterns consistent with local adaptation (Figure 2). In contrast, belowground

biomass was nearly equivalent (0.8% increase) between local and nonlocal$_{Rural}$ microbiomes,

whereas belowground biomass was 7.4% greater in local than nonlocal$_{Urban}$ microbiomes

(contrasts in Table S1).

283   Signatures of local adaptation were stronger for rhizobia when estimated as either nodule

density (P < 0.001, η^2_P = 0.247) or fixing nodule density (P < 0.001, η^2_P = 0.267, Table 1, Figure

2). Nodule density was 38.4% and 20.1% greater in local versus nonlocal$_{Rural}$ and nonlocal$_{Urban}$

microbiomes, respectively; however, nodule density depended on the three-way interaction

between microbiome, N, and population (Table 1, Figure 2). There was an increase in the

frequency of populations demonstrating local adaptation under N addition (local vs. nonlocal$_{Rural}$

= 83%, local vs. nonlocal$_{Urban}$ = 62%) compared to ambient N (local vs. nonlocal$_{Rural}$ = 63%,

local vs. nonlocal$_{Urban}$ = 57%; contrasts in Table S1). Fixing nodule density was 53.4% and

33.8% greater in the local microbiome treatment than the nonlocal$_{Rural}$ and nonlocal$_{Urban}$

treatments, respectively (contrasts in Table S1).

As expected, N addition had strong and positive effects on aboveground biomass (81.3%

increase, $P < 0.001$, $\eta^2_P = 0.871$, Table 1, Figure 2) and belowground biomass (75.0% increase, P

< 0.001, $\eta^2_P = 0.324$, Table 1, Figure 2;contrasts in Table S1). In contrast, N addition decreased

nodule density by 17.9% for both total nodule fixing density ($P < 0.001$, $\eta^2_P = 0.323$, Table 1,

Figure 2) and fixing nodule density ($P = 0.025$, $\eta2P = 0.167$, Table 1, Figure 2; contrasts in

Table S1). Notably, differences between local and nonlocal microbiomes were greater under N

addition for both nodule density and fixing nodule density (Figure 1). When we further evaluated

microbiome × population interactions using Cockerham's method, we found that crossing of

reaction norms (i.e., changes in rank order of populations among environments) accounted for

most of the variation in the interaction for aboveground biomass (97.4%), nodule density

(86.9%), and fixing nodule density (97.9%), as opposed to changes in population variance.

Results described above were similar when microbiome treatments were examined as local vs.

nonlocal$_{Global}$: there were stronger effects of local adaptation for both nodule density and fixing

nodule density but no evidence of local adaptation for belowground biomass (Table S2, Figure

S2).

Reciprocal Local Adaptation

White clover and rhizobia exhibited patterns of reciprocal local adaptation consistent with

coevolution, but linkages between white clover and rhizobia depended on the N environment and

the local – nonlocal contrast (Figures 3, S2-S3). A positive correlation between indices of local

adaptation for aboveground biomass and nodule density was apparent under N addition for the

global (local − nonlocal$_{Global}$, r and 95% confidence interval: r = 0.418 [0.061, 0.681]; Figure 3),

rural (local − nonlocal$_{Rural}$, r = 0.396 [0.034, 0.666]; Figure S2), and urban contrasts (local −

nonlocal$_{Urban}$, r = 0.301 [−0.081, 0.606]; Figure S3). There was no clear correlation between

indices of local adaptation for aboveground biomass and nodule density under ambient N (local

− nonlocal$_{Global}$, r = −0.194, 95% CI [−0.518, 0.179]; local − nonlocal$_{Rural}$, r = −0.095, 95% CI

[−0.440, 0.274]; local − nonlocal$_{Urban}$, r = −0.087, 95% CI [−0.434, 0.282]; Figures 3, S2-S3).

Relationships between belowground biomass and nodule density generally switched from

negative under ambient N (local − nonlocal$_{Global}$, r = −0.301 [−0.597, 0.067]; local −

nonlocal$_{Rural}$, r = −0.214 [−0.533, 0.159]; local − nonlocal$_{Urban}$, r = −0.197 [−0.520, 0.176];

Figures 3, S2-S3) to positive under N addition (local − nonlocal$_{Global}$, r = 0.253 [−0.125, 0.567];

local − nonlocal$_{Rural}$, r = 0.305 [−0.069, 0.604]; local − nonlocal$_{Urban}$, r = 0.052 [−0.327, 0.417];

Figures 3, S2-S3). When we examined the potential for reciprocal local adaptation between

fixing nodule density and either aboveground or belowground biomass, we observed positive

correlations in the rural contrast under N addition (aboveground biomass: local − nonlocal$_{Rural}$, r

= 0.424 [0.069, 0.684]; belowground biomass: local − nonlocal$_{Rural}$, r = 0.399 [0.038, 0.668];

Figure S2).

Local Adaptation and Urbanization

We did not identify any strong or consistent relationships between local adaptation and

urbanization, regardless of local adaptation index (Figures 4, S4-S6). There were no consistent

patterns in the strength and direction of relationships between local adaptation indices for our

fitness estimates and urbanization metrics (Figures 4, S4-S6), and all of the relationships

explained a negligible amount of variation (range of $R^2_{Adjusted}$ = 0-0.031).

Fitness and the Broader Microbiome

Variation in the microbiome community did not influence fitness for either white clover or

rhizobia. There was no main effect of *Rhizobium* abundance or *Rhizobium* abundance × N

interaction on any of the four fitness estimates (Table S3, Figure 5); using *Rhizobium* relative

abundance gave similar results (Table S4, Figure S7). Local adaptation was unrelated to

microbiome inoculant dissimilarity (Table S5, Figure 5) or root bacterial composition (Table S6,

Figure S8).

Rhizobium abundance was influenced by the N treatment and an interaction between

microbiome and N treatments (Table S7, Figure S9). Root bacterial community composition did

not vary by microbiome inoculant, N, or the interaction (Table S6, Figure S8). Community

composition was more variable among local microbiome inoculants than among either the

nonlocal$_{Rural}$ or nonlocal$_{Urban}$ inoculants or the experimental controls (Figure S9), and *Rhizobium*

abundances were similar among all microbiome inoculants (Cohen's $d = 0.015$-0.158, Figure

S10).

Discussion

Our experiment documents a mosaic of local adaptation in the mutualism between white clover

and rhizobia along an urbanization gradient, a conclusion supported by four key results. First,

both white clover and rhizobia had overall greater mean fitness in their local microbiomes

relative to the nonlocal microbiomes despite extensive variation among populations. Second,

fitness responses were unrelated to *Rhizobium* abundance and local adaptation was unrelated to

the broader bacterial microbiome in the soil inoculants or root endosphere. Together, these

results suggest local adaptation is driven by local evolution of *Rhizobium* and not ecological

changes in *Rhizobium* populations or microbiome composition. Third, N addition did not

consistently mediate the effects of local adaptation in white clover or rhizobia, although N did

influence the presence of reciprocal local adaptation between white clover and rhizobia, whereby

the strength of local adaptation was positively correlated between white clover and rhizobia in

the presence of high N. Finally, we did not find a consistent relationship between measures of

urbanization and local adaptation, suggesting urban environmental change did not alter patterns

of local adaptation. Below, we further discuss these results in the context of local adaptation.

legume-rhizobia coevolution, and urban evolution.

Local adaptation between white clover and rhizobia

Both white clover and rhizobia were locally adapted, although effects of local adaptation were

more frequent and stronger for rhizobia. One explanation for this asymmetry in local adaptation

is stronger selection on rhizobia compared to white clover. *Rhizobium* abundance and fitness is

higher when cooperating with plant hosts rather than living in the soil as a saprophyte (Denison

& Kiers 2004). In contrast, white clover's mutualism with rhizobia is facultative and dependent

on the N environment, and thus selection to maintain the mutualism may be weaker (Reid *et al.*

2011). Another explanation is that the extensive gene flow among white clover populations

within Toronto (Santangelo *et al.* 2022) and nearby cities (Johnson *et al.* 2018) could counteract

the effects of local adaptation (Thompson 2005; Gomulkiewicz *et al.* 2007). The population

structure for rhizobia has not been characterized in this system, but a separate study found that

populations of *R. leguminosarum* were genetically differentiated at spatial scales from ~4 km to

~350 km (Van Cauwenberghe *et al.* 2014), which is within the range of our urbanization gradient

(~50 km). Moreover, Van Cauwenberghe *et al.* (2016) observed genetic differentiation and

associated selection mosaics contributed to local adaptation between rhizobia and their host

plants. Evaluating the population structure for rhizobia is an important next step to understand

the balance of selection and gene flow and its role in shaping local adaptation in our system

(Thompson 2005; Gomulkiewicz *et al.* 2007; Hoeksema 2010; Wadgymar *et al.* 2022).

While there was an overall pattern of local adaptation, there was large variation among

populations in the strength and direction of local adaptation and even maladaptation. This

population × microbiome treatment interaction was driven by changes in the rank order of

populations between treatments for aboveground biomass, nodule density, and fixing nodule

density. Changes in rank order indicate that the relative-fitness of populations differ among

environments (Wade 2007), which can maintain genetic variation among populations when a

single genotype is not the most fit in all environments (Dempster 1955; Christiansen 1975;

Gomulkiewicz & Kirkpatrick 1992; Bell *et al.* 2021). These tradeoffs could have contributed to

the observed locally-adapted and locally-maladapted populations in our study.

Microbe-mediated adaptation

By sequentially examining how the microbiome inoculants could contribute to local adaptation,

we determined that *Rhizobium* evolution - not changes in *Rhizobium* abundance or microbiome

community composition - likely explains local adaptation in white clover and rhizobia. Fitness

responses of white clover and rhizobia were unrelated to *Rhizobium* abundance in the root

endosphere. As only *R. leguminosarum* can initiate nodulation on white clover (Andrews &

Andrews 2017), our results suggest that nodulation depended on having the 'right' *Rhizobium*

partner (i.e., genotype), which is consistent with the findings from other studies (Heath & Tiffin

2007; Heath 2010; Van Cauwenberghe *et al.* 2016; Burghardt *et al.* 2022). We found no

relationships between either microbiome inoculant dissimilarity or root endosphere composition

and local adaptation, indicating that the broader bacterial microbiome did not affect local

adaptation. These findings contrast with recent studies that demonstrated adaptation to local

microbiomes increases fitness (O'Brien *et al.* 2020, 2022b), and that dissimilarity between local

and nonlocal inoculants modulates fitness responses (O'Brien *et al.* 2022a). Importantly, rhizobia

density in inoculants could also confound nodulation (Batstone *et al.* 2020; Burghardt *et al.*

2022), but our microbiome inoculants had similar densities of *Rhizobium* (Figure S10).

Coevolution or correlated local adaptation?

We detected a positive correlation in the strength of local adaptation between white clover and

rhizobia, which is consistent with interspecific coevolution between mutualists. While our results

are consistent with a history of coadaptation resulting from coevolution, we note several

qualifications with this interpretation. First, this correlation was only detected between

aboveground biomass and nodule density under N addition, so whether or not a signature of

coevolution is detected depends on the traits measured and the environment. Second, the

presence or absence of a correlation does not necessarily implicate coevolution (Janzen 1980;

Nuismer *et al.* 2010). For example, spurious correlations can evolve in the absence of reciprocal

biotic selection if only one of the interacting species is evolving or abiotic factors favour

matching trait optima (Thompson 2005; Gomulkiewicz *et al.* 2007; Nuismer *et al.* 2010).

Alternatively, the absence of a correlation can arise despite ongoing coevolution because traits

could be mismatched (Thompson 2005; Gomulkiewicz *et al.* 2007; Nuismer *et al.* 2010). Further

investigation of potential coevolution between white clover and rhizobia in urban environments

requires characterizing selection acting on both white clover and rhizobia, and ideally genomic

signatures of selection on genes associated with the mutualism (Gomulkiewicz *et al.* 2007;

Hoeksema 2010; Batstone *et al.* 2022b, a; Epstein *et al.* 2023).

Conclusion

Our experiment contributes to the expanding literature on urban evolutionary ecology. Previous

studies of mutualisms have predominantly focused on plant-pollinator interactions and

adaptation for only one of the interacting species (Irwin *et al.* 2018; Theodorou *et al.* 2018;

Rivkin *et al.* 2020; but see Brans *et al.* 2022). Moreover, studies evaluating local adaptation in

urban environments have focused primarily on adaptation to abiotic factors, such as temperature

(Brans *et al.* 2017; Martin *et al.* 2021) and pollutants (Reid *et al.* 2016; Weston *et al.* 2021).

Anthropogenic impacts on legume-rhizobia symbioses have been evaluated in both agricultural

(Weese *et al.* 2015; Lau *et al.* 2022) and more recently in urban (Regus *et al.* 2017; Forrester &

Ashman 2018; Wendlandt *et al.* 2022) environments. Nitrogen addition can lead to the evolution

of less-cooperative rhizobia (Weese *et al.* 2015; Regus *et al.* 2017; Lau *et al.* 2022) but legumes

can still maintain beneficial associations with rhizobia despite N addition and deposition

(Forrester & Ashman 2018; Wendlandt *et al.* 2022). Our results show that species can be locally

adapted in urban environments, and that the mutualism between white clover and rhizobia can

persist despite increased N (Murray☐Stoker & Johnson 2021). As plant-microbe interactions are

essential for ecosystem functioning (van der Heijden *et al.* 2008; Bardgett & van der Putten

2014), it is necessary to understand how microbiomes contribute to adaptation and ecosystem

processes in anthropogenically-disturbed environments (Nugent & Allison 2022).

Acknowledgements

We thank Elana Maria, Oliva Toth, Samer El-Galmady, Wendy Van Drunen, Hind Emad, Kelly

Murray-Stoker, Zain Nassrullah, and Narindra Persaud for assistance with fieldwork, Elana

Maria and Alanah Joyce for help with lab work, and Joan Lee, Brenda Pitton, and the UTM

Facilities staff for experiment setup. Comments from Megan Frederickson, John Stinchcombe,

Sophie Breitbart, Lucas Albano, and Aude Caizergues greatly improved the manuscript.

Funding Information

This work was funded by an NSERC Discovery Grant, Canada Research Chair, and E.W.R.

Steacie Fellowship to M. T. J. Johnson and an American Society of Naturalists Student Research

Award to D. Murray-Stoker.

References

Andrews, M. & Andrews, M.E. (2017). Specificity in legume-rhizobia symbioses. *Int. J. Mol.*

Sci., 18, 705.

Bardgett, R.D. & van der Putten, W.H. (2014). Belowground biodiversity and ecosystem

functioning. *Nature*, 515, 505–511.

Batstone, R.T., Burghardt, L.T. & Heath, K.D. (2022a). Phenotypic and genomic signatures of

interspecies cooperation and conflict in naturally occurring isolates of a model plant

symbiont. *Proc. R. Soc. B Biol. Sci.*, 289, 20220477.

Batstone, R.T., Dutton, E.M., Wang, D., Yang, M. & Frederickson, M.E. (2017). The evolution

of symbiont preference traits in the model legume Medicago truncatula. *New Phytol.*,

213, 1850–1861.

Batstone, R.T., Lindgren, H., Allsup, C.M., Goralka, L.A., Riley, A.B., Grillo, M.A., *et al.*

(2022b). Genome-wide association studies across environmental and genetic contexts

reveal complex genetic architecture of symbiotic extended phenotypes. *mBio*, 0, e01823-

22.

Batstone, R.T., Peters, M.A.E., Simonsen, A.K., Stinchcombe, J.R. & Frederickson, M.E.

(2020). Environmental variation impacts trait expression and selection in the legume–

rhizobium symbiosis. *Am. J. Bot.*, 107, 195–208.

Bell, D.A., Kovach, R.P., Robinson, Z.L., Whiteley, A.R. & Reed, T.E. (2021). The ecological

causes and consequences of hard and soft selection. *Ecol. Lett.*, 24, 1505–1521.

Berner, D. & Amrhein, V. (2022). Why and how we should join the shift from significance

testing to estimation. *J. Evol. Biol.*, 35, 777–787.

Blanquart, F., Kaltz, O., Nuismer, S.L. & Gandon, S. (2013). A practical guide to measuring

local adaptation. *Ecol. Lett.*, 16, 1195–1205.

Brans, K.I., Jansen, M., Vanoverbeke, J., Tüzün, N., Stoks, R. & De Meester, L. (2017). The heat

is on: Genetic adaptation to urbanization mediated by thermal tolerance and body size.

Glob. Change Biol., 23, 5218–5227.

Brans, K.I., Tüzün, N., Sentis, A., De Meester, L. & Stoks, R. (2022). Cryptic eco-evolutionary

feedback in the city: Urban evolution of prey dampens the effect of urban evolution of

the predator. *J. Anim. Ecol.*, 91, 514–526.

Bronstein, J.L. (1994). Conditional outcomes in mutualistic interactions. *Trends Ecol. Evol.*, 9,

214–217.

Burdon, J.J. (1983). Trifolium repens L. *J. Ecol.*, 71, 307.

Burghardt, L.T., Epstein, B., Hoge, M., Trujillo, D.I. & Tiffin, P. (2022). Host-associated

rhizobial fitness: dependence on nitrogen, density, community complexity, and legume

genotype. *Appl. Environ. Microbiol.*, 88, e00526-22.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A. & Holmes, S.P. (2016).

DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods*,

13, 581–583.

Christiansen, F.B. (1975). Hard and soft selection in a subdivided population. *Am. Nat.*, 109, 11–

16.

Cockerham, C.C. (1963). Estimation of genetic variances. *Stat. Genet. Plant Breed.*, 982, 53–94.

Cole, J.R., Wang, Q., Fish, J.A., Chai, B., McGarrell, D.M., Sun, Y., *et al.* (2014). Ribosomal

Database Project: data and tools for high throughput rRNA analysis. *Nucleic Acids Res.*,

42, D633–D642.

Delgado-Baquerizo, M., Eldridge, D.J., Liu, Y.-R., Sokoya, B., Wang, J.-T., Hu, H.-W., *et al.*

(2021). Global homogenization of the structure and function in the soil microbiome of

urban greenspaces. *Sci. Adv.*, 7, eabg5809.

Dempster, E.R. (1955). Maintenance of genetic heterogeneity. *Cold Spring Harb. Symp. Quant.*

Biol., 20, 25–32.

Denison, R.F. & Kiers, E.T. (2004). Lifestyle alternatives for rhizobia: mutualism, parasitism,

and forgoing symbiosis. *FEMS Microbiol. Lett.*, 237, 187–193.

Diamond, S.E. & Martin, R.A. (2021). Evolution in cities. *Annu. Rev. Ecol. Evol. Syst.*, 52, 519–

540.

Epp Schmidt, D.J., Pouyat, R., Szlavecz, K., Setälä, H., Kotze, D.J., Yesilonis, I., *et al.* (2017).

Urbanization erodes ectomycorrhizal fungal diversity and may cause microbial

communities to converge. *Nat. Ecol. Evol.*, 1, 1–9.

Epstein, B., Burghardt, L.T., Heath, K.D., Grillo, M.A., Kostanecki, A., Hämälä, T., *et al.*

(2023). Combining GWAS and population genomic analyses to characterize coevolution

in a legume-rhizobia symbiosis. *Mol. Ecol.*, 32, 3798–3811.

Faeth, S.H., Bang, C. & Saari, S. (2011). Urban biodiversity: patterns and mechanisms. *Ann. N.*

Y. Acad. Sci., 1223, 69–81.

Forrester, N.J. & Ashman, T.-L. (2018). Nitrogen fertilization differentially enhances nodulation

and host growth of two invasive legume species in an urban environment. *J. Urban Ecol.*,

4, juy021.

Gelman, A. & Hennig, C. (2017). Beyond subjective and objective in statistics. *J. R. Stat. Soc.*

Ser. A Stat. Soc., 180, 967–1033.

Gomulkiewicz, R., Drown, D.M., Dybdahl, M.F., Godsoe, W., Nuismer, S.L., Pepin, K.M., *et al.*

(2007). Dos and don'ts of testing the geographic mosaic theory of coevolution. *Heredity*,

98, 249–258.

Gomulkiewicz, R. & Kirkpatrick, M. (1992). Quantitative genetics and the evolution of reaction

norms. *Evolution*, 46, 390–411.

Grimm, N.B., Faeth, S.H., Golubiewski, N.E., Redman, C.L., Wu, J., Bai, X., *et al.* (2008).

Global change and the ecology of cities. *Science*, 319, 756–760.

Heath, K.D. (2010). Intergenomic epistasis and coevolutionary constraint in plants and rhizobia.

Evolution, 64, 1446–1458.

Heath, K.D. & Stinchcombe, J.R. (2014). Explaining mutualism variation: a new evolutionary

paradox? *Evolution*, 68, 309–317.

Heath, K.D., Stock, A.J. & Stinchcombe, J.R. (2010). Mutualism variation in the nodulation

response to nitrate. *J. Evol. Biol.*, 23, 2494–2500.

Heath, K.D. & Tiffin, P. (2007). Context dependence in the coevolution of plant and rhizobial

mutualists. *Proc. R. Soc. B Biol. Sci.*, 274, 1905–1912.

van der Heijden, M.G.A., Bardgett, R.D. & van Straalen, N.M. (2008). The unseen majority: soil

microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol.*

Lett., 11, 296–310.

Hoeksema, J.D. (2010). Ongoing coevolution in mycorrhizal interactions. *New Phytol.*, 187,

286–300.

Irwin, R.E., Warren, P.S. & Adler, L.S. (2018). Phenotypic selection on floral traits in an urban

landscape. *Proc. R. Soc. B Biol. Sci.*, 285, 20181239.

Janzen, D.H. (1980). When is it coevolution? *Evolution*, 34, 611–612.

Johnson, M.T.J. & Munshi-South, J. (2017). Evolution of life in urban environments. *Science*,

358, eaam8327.

Johnson, M.T.J., Prashad, C.M., Lavoignat, M. & Saini, H.S. (2018). Contrasting the effects of

natural selection, genetic drift and gene flow on urban evolution in white clover

(Trifolium repens). *Proc. R. Soc. B Biol. Sci.*, 285, 20181019.

Kiers, E.T., Palmer, T.M., Ives, A.R., Bruno, J.F. & Bronstein, J.L. (2010). Mutualisms in a

changing world: an evolutionary perspective. *Ecol. Lett.*, 13, 1459–1474.

Lau, J.A., Hammond, M.D., Schmidt, J.E., Weese, D.J., Yang, W.H. & Heath, K.D. (2022).

Contemporary evolution rivals the effects of rhizobium presence on community and

ecosystem properties in experimental mesocosms. *Oecologia*, 200, 133–143.

Martin, R.A., Chick, L.D., Garvin, M.L. & Diamond, S.E. (2021). In a nutshell, a reciprocal

transplant experiment reveals local adaptation and fitness trade-offs in response to urban

evolution in an acorn-dwelling ant. *Evolution*, 75, 876–887.

McKinney, M.L. (2006). Urbanization as a major cause of biotic homogenization. *Biol.*

Conserv., Urbanization, 127, 247–260.

McKinney, M.L. (2008). Effects of urbanization on species richness: A review of plants and

animals. *Urban Ecosyst.*, 11, 161–176.

McShane, B.B., Gal, D., Gelman, A., Robert, C. & Tackett, J.L. (2019). Abandon statistical

significance. *Am. Stat.*, 73, 235–245.

Muir, W., Nyquist, W.E. & Xu, S. (1992). Alternative partitioning of the genotype-by-

environment interaction. *Theor. Appl. Genet.*, 84, 193–200.

Murray☐Stoker, D. & Johnson, M.T.J. (2021). Ecological consequences of urbanization on a

legume–rhizobia mutualism. *Oikos*, 130, 1750–1761.

Nugent, A. & Allison, S.D. (2022). A framework for soil microbial ecology in urban ecosystems.

Ecosphere, 13, e3968.

Nuismer, S.L., Gomulkiewicz, R. & Ridenhour, B.J. (2010). When is correlation coevolution?

Am. Nat., 175, 525–537.

Nuismer, S.L., Thompson, J.N. & Gomulkiewicz, R. (1999). Gene flow and geographically

structured coevolution. *Proc. R. Soc. B Biol. Sci.*, 266, 605–609.

O'Brien, A., Laurich, J. & Frederickson, M.E. (2022a). Having the "right" microbiome matters

for host trait expression and the strength of mutualism between duckweeds and microbes.

O'Brien, A.M., Laurich, J., Lash, E. & Frederickson, M.E. (2020). Mutualistic outcomes across

plant populations, microbes, and environments in the duckweed Lemna minor. *Microb.*

Ecol.

O'Brien, A.M., Yu, Z.H., Pencer, C., Frederickson, M.E., LeFevre, G.H. & Passeport, E.

(2022b). Harnessing plant-microbiome interactions for bioremediation across a

freshwater urbanization gradient. *Water Res.*, 223, 118926.

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., *et al.* (2020).

vegan: Community Ecology Package.

Pickett, S.T.A., Cadenasso, M.L., Grove, J.M., Boone, C.G., Groffman, P.M., Irwin, E., *et al.*

(2011). Urban ecological systems: scientific foundations and a decade of progress. *J.*

Environ. Manage., 92, 331–362.

Pickett, S.T.A., Cadenasso, M.L., Grove, J.M., Nilon, C.H., Pouyat, R.V., Zipperer, W.C., *et al.*

(2001). Urban ecological systems: linking terrestrial ecological, physical, and

socioeconomic components of metropolitan areas. *Annu. Rev. Ecol. Syst.*, 32, 127–157.

Poole, P., Ramachandran, V. & Terpolilli, J. (2018). Rhizobia: from saprophytes to

endosymbionts. *Nat. Rev. Microbiol.*, 16, 291–303.

R Core Team. (2022). R: a language and environment for statistical computing.

Regus, J.U., Wendlandt, C.E., Bantay, R.M., Gano-Cohen, K.A., Gleason, N.J., Hollowell, A.C.,

et al. (2017). Nitrogen deposition decreases the benefits of symbiosis in a native legume.

Plant Soil, 414, 159–170.

Reid, D.E., Ferguson, B.J., Hayashi, S., Lin, Y.-H. & Gresshoff, P.M. (2011). Molecular

mechanisms controlling legume autoregulation of nodulation. *Ann. Bot.*, 108, 789–795.

Reid, N.M., Proestou, D.A., Clark, B.W., Warren, W.C., Colbourne, J.K., Shaw, J.R., *et al.*

(2016). The genomic landscape of rapid repeated evolutionary adaptation to toxic

pollution in wild fish. *Science*, 354, 1305–1308.

Rivkin, L.R., Nhan, V.J., Weis, A.E. & Johnson, M.T.J. (2020). Variation in pollinator-mediated

plant reproduction across an urbanization gradient. *Oecologia*, 192, 1073–1083.

Rosier, C.L., Polson, S.W., D'Amico, V., Kan, J. & Trammell, T.L.E. (2021). Urbanization

pressures alter tree rhizosphere microbiomes. *Sci. Rep.*, 11, 9447.

RStudio Team. (2022). RStudio: Integrated development for R.

Santangelo, J.S., Ness, R.W., Cohan, B., Fitzpatrick, C.R., Innes, S.G., Koch, S., *et al.* (2022).

Global urban environmental change drives adaptation in white clover. *Science*, 375,

1275–1281.

Shantz, A.A., Lemoine, N.P. & Burkepile, D.E. (2016). Nutrient loading alters the performance

of key nutrient exchange mutualisms. *Ecol. Lett.*, 19, 20–28.

Stevens, C.J., David, T.I. & Storkey, J. (2018). Atmospheric nitrogen deposition in terrestrial

ecosystems: Its impact on plant communities and consequences across trophic levels.

Funct. Ecol., 32, 1757–1769.

Theodorou, P., Radzevičiūtė, R., Kahnt, B., Soro, A., Grosse, I. & Paxton, R.J. (2018). Genome-

wide single nucleotide polymorphism scan suggests adaptation to urbanization in an

important pollinator, the red-tailed bumblebee (Bombus lapidarius L.). *Proc. R. Soc. B*

Biol. Sci., 285, 20172806.

Thompson, J.N. (1988). Variation in interspecific interactions. *Annu. Rev. Ecol. Syst.*, 19, 65–87.

Thompson, J.N. (2005). *The Geographic Mosaic of Coevolution*. University of Chicago Press.

Udvardi, M. & Poole, P.S. (2013). Transport and metabolism in legume-rhizobia symbioses.

Annu. Rev. Plant Biol., 64, 781–805.

Van Cauwenberghe, J., Verstraete, B., Lemaire, B., Lievens, B., Michiels, J. & Honnay, O.

(2014). Population structure of root nodulating Rhizobium leguminosarum in Vicia

cracca populations at local to regional geographic scales. *Syst. Appl. Microbiol.*, 37, 613–

621.

Van Cauwenberghe, J., Visch, W., Michiels, J. & Honnay, O. (2016). Selection mosaics

differentiate Rhizobium-host plant interactions across different nitrogen environments.

Oikos, 125, 1755–1761.

Wade, M.J. (2007). The co-evolutionary genetics of ecological communities. *Nat. Rev. Genet.*, 8,

185–195.

Wadgymar, S.M., DeMarche, M.L., Josephs, E.B., Sheth, S.N. & Anderson, J.T. (2022). Local

adaptation: causal agents of selection and adaptive trait divergence. *Annu. Rev. Ecol.*

Evol. Syst., 53, 87–111.

Wang, Q., Garrity, G.M., Tiedje, J.M. & Cole, J.R. (2007). Naïve bayesian classifier for rapid

assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ.*

Microbiol., 73, 5261–5267.

Wasserstein, R.L. & Lazar, N.A. (2016). The ASA statement on p-values: context, process, and

purpose. *Am. Stat.*, 70, 129–133.

Wasserstein, R.L., Schirm, A.L. & Lazar, N.A. (2019). Moving to a world beyond "p < 0.05."

Am. Stat., 73, 1–19.

Weese, D.J., Heath, K.D., Dentinger, B.T.M. & Lau, J.A. (2015). Long-term nitrogen addition

causes the evolution of less-cooperative mutualists. *Evolution*, 69, 631–642.

Wendlandt, C.E., Gano-Cohen, K.A., Stokes, P.J.N., Jonnala, B.N.R., Zomorrodian, A.J., Al-

Moussawi, K., *et al.* (2022). Wild legumes maintain beneficial soil rhizobia populations

despite decades of nitrogen deposition. *Oecologia*, 198, 419–430.

Weston, L.M., Mattingly, K.Z., Day, C.T.C. & Hovick, S.M. (2021). Potential local adaptation in

populations of invasive reed canary grass (Phalaris arundinacea) across an urbanization

gradient. *Ecol. Evol.*, 11, 11457–11476.

Table 1: Summary of the linear mixed-effects models testing the effects of microbiome (local, nonlocal$_{Rural}$, nonlocal$_{Urban}$), nitrogen (ambient N, N addition), and the two-way interaction on measure of plant and rhizobia fitness. We provide the numerator degrees-of-freedom (NumDF), denominator degrees-of-freedom (estimated using the Kenward-Roger method (DenDF; Kenward & Roger 1997), F-statistics calculated from type III sums-of-squares, P-values, and the effect size (partial eta-squared, η^2_P) for each fixed effect term in the ANOVAs. We also provide the χ^2 statistic, P-values from likelihood ratio tests, and the effect size (intra-class correlation coefficient, ICC) for each random effect term, with random effect terms and statistics in italics. The variance explained by the fixed effects ($R^2_{Marginal}$) and combined fixed and random effects ($R^2_{Conditional}$) is provided for each model.

Term	NumDF	DenDF	F/χ^2	P-value	η^2_P/ICC
Aboveground Biomass					
($R^2_{Marginal}$ = 0.233, $R^2_{Conditional}$ = 0.397)					
Microbiome	2	54.911	3.243	0.047	0.106
Nitrogen	1	27.790	188.399	< 0.001	0.871
Microbiome × Nitrogen	2	742.517	0.470	0.625	0.001
Microbiome × Population			4.951	0.026	0.040
Nitrogen × Population			3.901	0.048	0.035
Population			10.311	0.001	0.126
Block			3.575	0.059	0.011
Belowground Biomass					
($R^2_{Marginal}$ = 0.292 $R^2_{Conditional}$ = 0.376)					
Microbiome	2	802.275	3.338	0.036	0.008
Nitrogen	1	801.735	384.570	< 0.001	0.324
Microbiome × Nitrogen	2	805.884	0.292	0.747	0.001
Population			46.239	< 0.001	0.104
Block			4.791	0.029	0.015
Nodule Density					
($R^2_{Marginal}$ = 0.061, $R^2_{Conditional}$ = 0.261)					
Microbiome	2	56.305	9.257	< 0.001	0.247
Nitrogen	1	27.932	13.328	0.001	0.323
Microbiome × Nitrogen	2	55.584	1.057	0.354	0.037
Microbiome × Population			1.614	0.204	0.053
Nitrogen × Population			0.019	0.889	0.004
Microbiome × Nitrogen × Population			6.376	0.012	0.105
Population			1.899	0.168	0.047
Block			0.390	0.533	0.003
Fixing Nodule Density					
($R^2_{Marginal}$ = 0.043, $R^2_{Conditional}$ = 0.198)					
Microbiome	2	55.390	10.067	< 0.001	0.267
Nitrogen	1	27.929	5.595	0.025	0.167
Microbiome × Nitrogen	2	741.058	0.442	0.643	0.001
Microbiome × Population			8.159	0.004	0.062
Nitrogen × Population			5.201	0.023	0.045
Population			0.787	0.375	0.026
Block			13.511	< 0.001	0.029

659 **Figures**

660
Figure 1: Map of the urbanization gradient in the Greater Toronto Area, Ontario, Canada (A),
with the location of the 30 focal populations indicated by yellow circles. We also show the
location of the 5 rural (blue circles) and 5 urban (pink circles) sites from which nonlocal
inoculants were sourced. Urban and suburban areas in gray, nonurban agricultural and forested
areas in green, and Lake Ontario in light blue. Map tiles by Stamen Design, under CC BY 3.0.
Data by OpenStreetMap, under ODbL. We also show a diagram of the experimental design (B).
Each of the 30 focal populations (yellow circles on the map) received 3 microbiome treatments
(local, nonlocal$_{Urban}$, and nonlocal$_{Rural}$). Additionally, each microbiome treatment received 2
nitrogen treatments (no addition = ambient N, 10 mM KNO$_3$ fertilizer = N addition). The
factorial experiment was conducted in an outdoor common garden at the University of Toronto
Mississauga campus.

673
Figure 2: Treatment means ± 1 SE (left column) and reaction norms of population means (right
column) by microbiome and nitrogen treatments for aboveground biomass (A-B), belowground
biomass (C-D), nodule density (E-F), and fixing nodule density (G-H). Microbiome treatments
are indicated as nonlocal$_{Rural}$ (blue), nonlocal$_{Urban}$ (pink), and local (yellow); individual plots are
faceted by nitrogen treatment (ambient N and N addition). Detailed test statistics are provided in
Table 1.

680
681 **Figure 3:** Pairwise correlations between the local – nonlocal$_{Global}$ local adaptation indices for
682 aboveground biomass, belowground biomass, nodule density, and fixing nodule density.
683 Separate correlations were calculated for each nitrogen treatment (ambient N = purple, N
684 addition = pink). Lines represent lines-of-best-fit (± 95% confidence interval). Inset text provides
685 the correlation coefficient (r) and P-value for each correlation.

686
Figure 4: Relationship between the global local adaptation index (local – nonlocal$_{Global}$, LA
Index$_{Global}$) and urbanization (standardized distance from the city centre; mean = 0, standard
deviation = 1) for each of the fitness estimates [aboveground biomass (A) belowground biomass
(B), nodule density (C), and fixing nodule density, (D)]. Lines represent the lines-of-best-fit ($\pm$
95% confidence interval). Inset text provides the slope parameter (β), standard error (SE), and P-
value for each linear regression. These conclusions are unchanged when other metrics of
urbanization (i.e., Human Influence Index and imperious surface cover) and indices of local
adaptation (i.e., local – nonlocal$_{Rural}$ and local – nonlocal$_{Urban}$) are used (Figures S4-S6).

Figure 5: The relationship between fitness estimates and *Rhizobium* abundance (left column) and local adaptation by pairwise microbiome dissimilarity (right column) for aboveground biomass (A and B), belowground biomass (C and D), nodule density (E and F), and fixing nodule density (G and H). Lines represent lines-of-best-fit (± 95% confidence interval), with separate lines for each nitrogen treatment (ambient N = purple, N addition = pink) for the *Rhizobium* abundance models. Inset text for the right column provides the slope parameter (β), standard error (SE), and P-value for each linear regression. Detailed test statistics are provided in Table S3 (panels A, C, E, G) and Table S5 (panels B, D, F, H).