Matrix-associated extracellular vesicles modulate smooth muscle cell adhesion and directionality by presenting collagen VI.

Alexander Kapustin[1*], Sofia Serena Tsakali[1], Meredith Whitehead[1], George Chennell[2], Meng-Ying Wu[1, 3], Chris Molenaar[1], Anton Kutikhin[4], Yimeng Chen[1], Sadia Ahmad[1], Leo Bogdanov[4], Maxim Sinitsky[3], Kseniya Rubina[5], Aled Clayton[6], Frederik J Verweij[7], Dirk Michiel Pegtel[8], Simona Zingaro[9], Arseniy Lobov[10], Bozhana Zainullina[11], Dylan Owen[12], Maddy Parsons[9], Richard E. Cheney[13], Derek Warren[14], Martin James Humphries[15], Thomas Iskratsch[16], Mark Holt[8], and Catherine M Shanahan[1*]

[1] School of Cardiovascular and Metabolic Medicine & Sciences, James Black Centre, King's College London, 125 Coldharbour Lane, London, SE5 9NU, UK, Tel. 020 7848 5221, FAX 020 7848 5193,

[2] Wohl Cellular Imaging Centre, King's College London, 5 Cutcombe Road, London, SE5 9NU

[3] Department of Urology, The Third Affiliated Hospital of Soochow University, Changzhou, Jiangsu, China, 213003

[4] Laboratory for Molecular, Translational and Digital Medicine, Research Institute for Complex Issues of Cardiovascular Diseases, 6 Sosnovy Boulevard, Kemerovo, 650002, Russian Federation

[5] Laboratory of Morphogenesis and Tissue Reparation, Faculty of Medicine, Lomonosov Moscow State University, Lomonosovsky av. 27-1, Moscow, 119991, Russia, tel/fax +74959329904

[6] Tissue Microenvironment Research Group, Division of Cancer & Genetics, School of Medicine, Cardiff University, Tenovus Building, Cardiff, UK, CF14 2XN

[7] Division of Cell Biology, Neurobiology & Biophysics, Utrecht University, Padualaan 8, 3584 CH, Utrecht, The Netherlands

[8] Amsterdam UMC, Location Vrije Universiteit Amsterdam, Department of Pathology, Cancer Center Amsterdam, De Boelelaan 1117, Amsterdam, The Netherlands

[9] Randall Centre for Cell and Molecular Biophysics, School of Basic and Medical Biosciences, King's College London, New Hunt's House, Guy's Campus, London, SE1 1UL UK.

[10] Laboratory of Regenerative Biomedicine, Institute of Cytology of the Russian Academy of Sciences, 4 Tikhoretskiy Prospekt, 194064, St. Petersburg, Russia

[11] Centre for Molecular and Cell Technologies, Research Park, St. Petersburg State University, 7/9 Universitetskaya Embankment, 199034, St. Petersburg, Russia

[12] Institute of Immunology and Immunotherapy, School of Mathematics and Centre of Membrane Proteins and Receptors (COMPARE), University of Birmingham, Birmingham, B15 2TT, UK.

[13] Department of Cell Biology and Physiology, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA

[14] School of Pharmacy, University of East Anglia, Norwich Research Park, Norwich, Norfolk, UK, NR4 7TJ

[15] Wellcome Centre for Cell-Matrix Research, Faculty of Biology, Medicine & Health, Manchester Academic Health Science Centre, University of Manchester, Michael Smith Building, Oxford Road, Manchester, M13 9PT, United Kingdom

1   [16]School of Engineering and Materials Science, Faculty of Science and Engineering, Queen Mary

2   University of London, Engineering Building, Mile End Road, E1 4NS

4   * Corresponding authors

5   Email: Alexander.kapustin@kcl.ac.uk

6   Words: small extracellular vesicles, cell adhesion, vascular remodelling

1 Abstract

The extracellular matrix (ECM) supports blood vessel architecture and functionality and undergoes
active remodelling during vascular repair and atherogenesis. Vascular smooth muscle cells (VSMCs)
are essential for vessel repair and, via their secretome, can invade from the vessel media into the
intima to mediate ECM remodelling. Accumulation of fibronectin (FN) is a hallmark of early vascular
repair and atherosclerosis. Here we show that FN stimulates VSMCs to secrete small extracellular
vesicles (sEVs) by activating the β1 integrin/FAK/Src pathway as well as Arp2/3-dependent branching
of the actin cytoskeleton. We found that sEVs are trapped by the ECM *in vitro* and colocalise with FN
in symptomatic atherosclerotic plaques *in vivo*. Functionally, ECM-trapped sEVs induced the
formation of focal adhesions (FA) with enhanced pulling forces at the cellular periphery preventing
cellular spreading and adhesion. Proteomic and GO pathway analysis revealed that VSMC-derived
sEVs display a cell adhesion signature and are specifically enriched with collagen VI on the sEV
surface. *In vitro* assays identified collagen VI as playing a key role in cell adhesion and invasion
directionality. Taken together our data suggests that the accumulation of FN is a key early event in
vessel repair acting to promote secretion of collage VI enriched sEVs by VSMCs. These sEVs stimulate
directional invasion, most likely by triggering peripheral focal adhesion formation and actomyosin
contraction to exert sufficient traction force to enable VSMC movement within the complex vascular
ECM network.

1 Introduction.

The healthy arterial vasculature is dominated by a highly organised medial extracellular matrix (ECM)
containing organised layers of contractile vascular smooth muscle cells (VSMCs). Vascular
pathologies such as atherosclerosis are associated with dramatic remodelling of the ECM ultimately
leading to plaque rupture and myocardial infarction or stroke[1]. Progenitor medial VSMCs are
essential for vessel repair and must invade through the ECM to form the protective intimal fibrous
cap in the plaque. This process of VSMC invasion actively contributes to ECM remodelling[2-4]. The
accumulation of liver-derived fibronectin (FN) in the vasculature is an early biomarker of
atherosclerotic plaque formation and conditional FN knockout in the liver blocked VSMC invasion
and fibrous cap formation in the ApoE mouse model. This suggests that FN is an essential signal for
VSMC recruitment and invasion during vascular repair, yet its exact role remains unknown[5-7].

FN has been shown to play key signalling roles by modulating cellular spreading, adhesion, invasion,
differentiation and viability during both developmental and pathological processes[8]. Cell adhesion to
FN is primarily mediated by $\alpha5\beta1$ integrins and proteoglycans acting together to activate small GTP-
binding proteins, Cdc42, Rac1, and Rho which in turn induce branched actin cytoskeletal
rearrangements to form cellular membrane protrusions, filopodia and lamellipodia, which are
attached to the ECM via transient peripheral focal complexes[8-11]. In turn, maturation of focal
complexes into focal adhesions (FAs) anchors cytoplasmic actin stress fibers to the ECM and actin
polymerisation and actomyosin-mediated contractility generate the traction forces required for cell
body locomotion[12, 13].

Exosome-like small extracellular vesicles (sEVs) are novel, cell-matrix crosstalk entities that decorate
the ECM and form "migration paths" for tumour cells by enhancing cell adhesion, motility and
directionality[14-17]. Mechanistically, sEVs are secreted at the cellular leading edge and promote FA
formation by presenting fibronectin (FN)[14, 15, 18, 19]. In addition, sEVs can stimulate motility of
immune cells and *Dictyostelium discoideum* by presenting cytokines and/or generating
chemoattractants[20-22]. We recently showed that FN is also enriched in VSMC-derived sEVs but the
exact role of sEVs in VSMC migration and invasion within the ECM meshwork environment of the
vasculature remains unexplored[23].

Here we report that FN in the ECM induces sEV secretion by activating $\beta1$ integrin/FAK/Src and
Arp2/3-dependent actin cytoskeleton assembly. In turn, sEVs are trapped to ECM and these ECM-
associated sEVs regulate VSMC invasion directionality in a 3D model. Mechanistically ECM-
associated sEVs induced earlier formation of focal adhesions (FA) with enhanced pulling forces at the
cell periphery thus switching the leading edge from protrusion activity into contractile mode to
enable cell body propulsion. Importantly, we found that that the sEV cargo, collagen VI, is
indispensable for triggering FA formation and directed cell invasion. We hypothesize that sEV-
triggered peripheral FA formation anchors the cellular leading edge and activates cell contraction to
exert sufficient force to allow VSMC invasion into the complex ECM fibre meshwork. This novel
mechanism opens a unique therapeutic opportunity to target specifically VSMC invasion activity
during vascular repair and restenosis.

Results.

1. ECM components involved in vessel injury repair stimulate sEV release via b1 integrins.

FN is detectable in the vasculature as early as the fatty streak stage and preceding major atherogenic
alterations such as neointima formation and it is also a novel marker for symptomatic carotid
plaques[5-7]. Given its presence during early stages of vessel injury we hypothesized that FN may
modulate sEV secretion to enable vessel repair. FN is secreted as a monomeric protein forming
fibrils upon cellular binding so we compared the effects of monomeric FN with the effects of FN
immobilised on tissue culture plates to mimic FN fibrils[8]. Plating VSMCs on immobilised FN
increased the release of CD63+/CD81+ sEVs 3.5±0.6 fold whilst addition of soluble FN had no effect
on EV secretion as detected by CD63-bead capture assay (Fig 1A). Fibrillar collagen I but not a non-
fibrillar laminin also stimulated secretion of CD63+/CD81+ sEVs by VSMCs to the same extent as FN
(Fig S1A).

We next tested if the native VSMC-derived 3D matrix that contains FN and collagen could modulate
sEV secretion[24]. VSMCs were induced to produce ECM, cells were removed and fresh VSMCs plated
onto these 3D matrices. VSMCs acquired an elongated shape (Figs 1B and 1C) which is typical for
mesenchymal cells in a 3D environment[24] and increased secretion of CD63+/CD81+ sEVs compared
to VSMCs plated onto the non-coated plastic (Fig 1D). We observed no changes in the size
distribution of sEVs secreted by VSMCs plated either on plastic or FN matrix as detected using
Nanoparticle Tracking Analysis (NTA) (Fig 1E). To further characterise the EV populations released
western blotting confirmed that sEVs isolated in both conditions were loaded with similar levels of
FN and the sEV-specific markers CD63 and Syntenin-1 and lacked the EV-specific marker, α-actinin-4
(Fig 1F)[25]. In addition, treatment of the cells plated onto the matrix with sphingomyelin
phosphodiesterase 3 inhibitor (3-O-Methyl-Sphingomyelin, 3-OMS) revealed that CD63+/CD81+ sEV
secretion in response to collagen and FN was regulated by sphingomyelin phosphodiesterase 3 (Fig
S1B) which regulates sEV generation in multivesicular bodies[26]. Hence, FN triggers secretion of
CD63+/CD81+ sEVs, most likely with a late endosomal origin.

Both collagen I and FN are ECM ligands that bind and transduce intracellular signalling via β1
integrin. Therefore, we explored the effect of β1 integrin activating (12G10) or inhibiting antibodies
(4B4) on FN-stimulated secretion of CD63+/CD81+ sEVs. Activation of β1 integrin using 12G10
antibody in VSMCs enhanced the effect of FN on sEV secretion (Fig 1G). Inhibition of β1 integrin
using the 4B4 antibody blocked CD63+/CD81+ sEV secretion by VSMCs plated on FN (Fig 1G). Next,
we tested the role of the β1 integrin downstream signalling mediators, FAK and Src[27]. Blocking these
pathways with small inhibitors (FAM14 and PP2, respectively) reduced sEV secretion by cells plated
on FN but not on plastic (Figs 1H and 1I). Taken together these data suggest that FN matrices
stimulate secretion of CD63+/CD81+ sEVs via the β1 integrin signalling pathway.

2. FN-induced sEV secretion is modulated by Arp2/3 and formin-dependent actin cytoskeleton
remodelling

ECM-induced focal FAs initiate dynamic remodelling of the actin cytoskeleton, a process regulated
by the Arp2/3 complex and formins, which have recently been implicated in exosome secretion in
lymphocytes and tumor cells[28-30] . So next we tested the contribution of Arp2/3 and formins by using
the small molecule inhibitors, CK666 and SMIFH2, respectively[31, 32]. CK666 reduced sEV secretion
both in VSMCs plated on plastic and FN matrix while inhibition of formins using SMIFH2 was only

effective on the FN matrix (Figs 2A and 2B) indicating that formins may be involved in FN-dependent
sEV secretion.

As regulators of branched actin assembly, the Arp2/3 complex and cortactin are thought to
contribute to sEV secretion in tumour cells by mediating MVB intracellular transport and plasma
membrane docking[28, 33]. Therefore, we overexpressed the Arp2/3 subunit, ARPC2-GFP and the F-
actin marker, F-tractin-RFP in VSMCs and performed live-cell imaging. As expected, Arp2/3 and F-
actin bundles formed a distinct lamellipodia scaffold in the cellular cortex (Fig S2A and
Supplementary Video S2). Unexpectedly, we also observed numerous Arp2/3/F-actin positive spots
moving through the VSMC cytoplasm that resembled previously described endosome actin tails
observed in Xenopus eggs[33] and parasite infected cells where actin comet tails propel parasites via
filopodia to neighbouring cells[34, 35] (Fig S2A, arrow, and Supplementary Video S2). Analysis of the
intracellular distribution of Arp2/3 and CD63-positive endosomes in VSMCs showed CD63-MVB
propulsion by the F-actin tail in live cells (Fig 2C and Supplementary Video S4). Although these were
rare events, possibly due to the short observational time in a single microscopic plane (Fig 2C) we
observed numerous F-actin spots in fixed VSMCs that were positive both for F-actin and cortactin
indicating that these are branched-actin tails (Fig 2D). Moreover, cortactin/F-actin spots colocalised
with CD63+ endosomes and addition of the SMPD3 inhibitor, 3-OMS, induced the appearance of
enlarged doughnut-like cortactin/F-actin/CD63 complexes resembling invadopodia-like structures
similar to those observed in tumour cells (Fig 2D, arrowheads)[18]. To quantify CD63 overlap with the
actin tail-like structures, we extracted round-shaped actin structures and calculated the thresholded
Manders colocalization coefficient (Fig S2B). We observed overlap between F-actin tails and CD63 as
well as close proximity of these markers in fixed VSMCs (Fig S2B). Approximately 50% of the F-actin
tails were associated with ~13% of all endosomes (tM1=0.44±0.23 and tM2= 0.13±0.06,
respectively, N=3). Addition of 3-OMS enhanced this overlap further (tM1=0.75±0.18 and
tM2=0.25±0.09) suggesting that Arp2/3-driven branched F-actin tails are involved in CD63+ MVB
intracellular transport in VSMCs.

Formins and the Arp2/3 complex play a crucial role in the formation of filopodia, a cellular
protrusion required for sensing the extracellular environment and cell-ECM interactions[36]. To test
whether MVBs can be delivered to filopodia, we stained VSMCs for Myosin-10 (Myo10)[37]. We
observed no difference between total filopodia number per cell on plastic or FN matrices (n=18±8
and n=14±3, respectively) however the presence of endogenous CD63+ MVBs along the Myo10-
positive filopodia were observed in both conditions (Fig 2E, arrows). Filopodia have been implicated
in sEV capture and delivery to endocytosis "hot-spots"[38], so next we examined the directionality of
CD63+ MVB movement in filopodia by overexpressing Myo10-GFP and CD63-RFP in live VSMCs.
Importantly, we observed anterograde MVB transport toward the filopodia tip (Fig 2F and
Supplementary Video S2) indicative of MVB secretion.

We also attempted to visualise sEV release in filopodia using CD63-pHluorin where fluorescence is
only observed upon the fusion of MVBs with the plasma membrane[39]. Using total internal reflection
fluorescence microscopy (TIRF) we observed the typical "burst"-like appearance of sEV secretion at
the cell-ECM interface in full agreement with an earlier report showing MVB recruitment to
invadopodia-like structures in tumor cells[18] (Fig S2B and Supplementary Video S1). Although we also
observed an intense CD63-pHluorin staining along filopodia-like structures we were not able to
detect typical "burst"-like events to confirm sEV secretion in filopodia. (Fig S2C and Supplemental
Video S1).

Taken together these data show that sEV secretion by VSMCs is regulated via Arp2/3 and formin-
dependent actin cytoskeleton remodelling. Branched actin tails can potentially contribute to MVB
intracellular transport and secretion at the VSMC-ECM interface. Interestingly, CD63+ MVBs can be
observed in filopodia-like structures suggesting that sEV secretion can also occur spatially via cellular
protrusion-like filopodia but more studies are needed to confirm this hypothesis.

3. sEVs are trapped in the ECM in vitro and in atherosclerotic plaque.

EV trapping in the ECM is a prominent feature of the blood vessel media and intima and EVs become
more abundant in the ECM with ageing and disease[40, 41]. Therefore, we next set out to determine if
sEVs that are secreted toward the ECM can be trapped in the native matrix _in vitro_. VSMCs were
plated on plastic or FN for 24 h and sEVs were visualised by CD63 immunofluorescence. We
observed CD63 puncta in proximity to protrusions at the cell periphery (Fig 3A). These CD63+ sEVs
were observed both on the plastic and FN-coated plates suggesting that sEVs can bind to the ECM
secreted by VSMCs over the 24h incubation. In agreement with our previous data we also observed
CD63+ sEV trapping in the native ECM produced by VSMCs over a 7 day period and their
reduction/absence from the matrix when VSMCs were treated with SMPD3 siRNA (Fig 3B)[41].

To quantify ECM-trapped sEVs we applied a modified protocol for the sequential extraction of
extracellular proteins using salt buffer (0.5M NaCl) to release sEVs which are loosely-attached to
ECM via ionic interactions, followed by 4M guanidine HCl buffer (GuHCl) treatment to solubilize
strongly-bound sEVs (Fig S3A) [42]. We quantified total sEV and characterised the sEV tetraspanin
profile in conditioned media, and the 0.5M NaCl and GuHCl fractions using ExoView. The total
particle count showed that EVs are both loosely bound and strongly trapped within the ECM. sEV
tetraspanin profiling showed differences between these 3 EV populations. While there was close
similarity between the conditioned media and the 0.5M NaCl fraction with high abundance of
CD63+/CD81+ sEVs as well as CD63+/CD81+/CD9+ in both fractions (Fig S3A). In contrast, the GuHCl
fraction was particularly enriched with CD63+ and CD63+/CD81+ sEVs with very low abundance of
CD9+ EVs (Fig S3A). The abundance of CD63+/CD81+ sEVs was confirmed independently by a CD63+
bead capture assay in the media and loosely bound fractions (Fig S3B).

We previously found that the serum protein prothrombin binds to the sEV surface both in the media
and MVB lumen showing it is recycled in sEVs and catalyses thrombogenesis being on the sEV
surface[43]. So we investigated whether FN can also be associated with sEV surface where it can be
directly involved in sEV-cell cross-talk[43]. We treated serum-deprived primary human aortic VSMCs
with FN-Alexa568 and found that it was endocytosed and subsequently delivered to early and late
endosomes together with fetuin A, another abundant serum protein that is a recycled sEV cargo and
elevated in plaques (Figs S3C and S3D). CD63 visualisation with a different fluorophore (Alexa488)
confirmed FN colocalization with CD63+ MVBs (Fig S3E). Next, we stained non-serum deprived VSMC
cultured in normal growth media (RPMI supplemented with 20% FBS) with an anti-FN antibody and
observed colocalization of CD63 and serum-derived FN. Co-localisation was reduced likely due to
competitive bulk protein uptake by non-deprived cells (Fig S3F). Notably, when we compared FN
distribution in sparsely growing VSMCs versus confluent cells we found that FN intracellular spots, as
well as colocalization with CD63, completely disappeared in the confluent state (Fig S3F and S3G).
This correlated with nearly complete loss of CD63+/CD81+ sEV secretion by the confluent cells
indicating that confluence abrogates intracellular FN trafficking as well as sEV secretion by VSMCs
(Fig S3H). Finally, FN could be co-purified with sEVs from VSMC conditioned media (Fig S3I) and
detected on the surface of sEVs by flow cytometry confirming its loading and secretion via sEVs (Fig
3C).

To understand how VSMCs interact with sEVs in the ECM versus in the media we isolated and
fluorescently-labelled sEVs and added them either to the ECM or the media and tracked sEV
distribution. Super-resolution microscopy (iSIM) revealed that the addition of sEVs to the cell media
resulted in fast uptake by VSMCs and delivery to the nuclear periphery (Fig 3D and Fig S3J). This was
particularly obvious in 3D projections (Figs S3J). In contrast, purified sEVs added to FN-coated plates
resulting in their even distribution across the ECM were not internalised even 24h after cell addition
with no perinuclear localisation of ECM-associated sEVs observed (Fig 3E and Fig S3K). No red
fluorescence was observed in the absence of added sEVs (Fig 3F).

To spatially map the accumulation of sEV markers in the atherosclerotic plaque *in vivo* we collected
12 carotid atherosclerotic plaques and adjacent intact vascular segments excised during carotid
endarterectomy. Unbiased proteomics analysis confirmed considerable differences between these 2
sample groups, revealing 213 plaque-specific and 111 intact arterial-specific proteins (Fig 4A and Fig
4B, Tables S1-S5). Differential expression analysis identified 46 proteins significantly overexpressed
(fold change ≥ 2 and FDR-adjusted P value ≤ 0.05) in plaques and 13 proteins significantly
upregulated in the intact arterial segments (Fig 4C, Table S3). Among the top proteins differentially
expressed in plaques were catalytic lipid-associated bioscavenger paraoxonase 1, atherogenic
apolipoprotein B-100, HDL-associated apolipoprotein D, iron-associated protein haptoglobin, and
inflammation-related matrix metalloproteinase-9 (Table S3). These proteins have previously been
implicated in lipid metabolism alterations as well as inflammation in the plaque hence indicating the
advanced atherosclerotic plaque signature of the analysed samples. Comparison of the differentially
expressed proteins also revealed an accumulation in atherosclerotic regions of fetuin-A (Alpha-2-HS-
glycoprotein, P02765, 2.7 fold increase, p=0.003869659), an abundant sEV cargo protein which is
recycled by VSMCs[23]. Likewise, the level of another sEV cargo, Apolipoprotein B-100 (P04114) was
also significantly elevated (2.8 fold increase, p= 0.000124446) in atherosclerotic plaque (Table S2)[23].
Notably, a negative regulator of sEV secretion, Coronin 1C (Q9ULV4) was downregulated in the
plaque (0.6 fold decrease, p=0.000805372, Table S2) consistent with previous studies that suggested
that sEV secretion is upregulated during plaque progression[40, 44].

To test if FN associates with sEV markers in atherosclerosis, we investigated the spatial association
of FN with sEV markers using the sEV-specific marker CD81. Staining of atherosclerotic plaques with
haematoxylin and eosin revealed well-defined regions with the neointima as well as tunica media
layers formed by phenotypically transitioned or contractile VSMCs, respectively (Fig S4A). Masson's
trichrome staining of atherosclerotic plaques showed abundant haemorrhages in the neointima, and
sporadic haemorrhages in the tunica media (Fig S4B). Staining of atherosclerotic plaques with orcein
indicated weak connective tissue staining in the atheroma with a confluent extracellular lipid core,
and strong specific staining at the tunica media containing elastic fibres which correlated well with
the intact elastin fibrils in the tunica media (Figs S4C and S4D). Using this clear morphological
demarcation, we found that FN accumulated both in the neointima and the tunica media where it
was significantly colocalised with the sEV marker, CD81 (Fig. 4D, 4E and 4F). Notably CD81 and FN
colocalization was particularly prominent in cell-free, matrix-rich plaque regions (Figs. 4E and 4F).
Interestingly, colocalization analysis of CD81 and FN distribution showed that there was a trend of
higher colocalization levels of CD81 with FN in the neointima as compared to tunica media which
was not significant (Thresholded Mander's split colocalisation coefficient for CD81 colocalization
with FN 0.58±0.21 and 0.49±0.14, respectively, N=13, p=0.052, paired T test). An enhanced
expression of CD81 by endothelial cells in early atheroma has been previously reported so to study
the contribution of CD81+ sEVs derived from endothelial cells we investigated the localisation of
CD31 and CD81[45]. In agreement with a previous study, we found that the majority of CD31
colocalises with CD81 (Thresholded Mander's split colocalization coefficient 0.54±0.11, N=6)

indicating that endothelial cells express CD81 (Fig 4G)[45]. However, only a minor fraction of total
CD81 colocalised with CD31 (Thresholded Mander's split colocalization coefficient 0.24±0.06, N=6)
confirming that the majority of CD81 in the neointima is originating from the most abundant VSMCs.
Finally, western blot analysis confirmed FN accumulation in the plaque region (Figs 4H, 4I and S4E).
To understand the origin of plaque FN we performed RT-PCR. FN expression could neither be
detected in plaques nor in intact vessels, although it was abundant in the liver (*data not shown*),
suggesting that the circulation is a key source of FN in atherosclerotic plaques.

4. sEVs stimulate VSMC directional invasion and induce early peripheral focal adhesion assembly
with an enhanced pulling force.

FN as a cargo in sEVs promotes FA formation in tumour cells and increases cell speed[14, 15]. As we
found that FN is loaded into VSMC-derived sEVs we hypothesized that ECM-entrapped sEVs can
enhance cell migration by increasing cell adhesion and FA formation in the context of a FN-rich ECM.
Therefore, we tested the effect of sEV deposition onto the FN matrix on VSMC migration in 2D and
3D models. We found that FN coating promoted VSMC velocity and inhibition of bulk sEV secretion
with 3-OMS reduced VSMC speed in a 2D single-cell migration model (Figs. 5A, 5B) in agreement
with previous studies using tumour cells[14, 15]. However, addition of sEVs to the ECM had no effect on
VSMC speed at baseline but rescued cell speed and distance in the presence of the sEV secretion
inhibitor, 3-OMS suggesting the EVs are not primarily regulating cell speed (Figs 5A and 5B).

To assess the effect of sEVs on cell directionality we exploited a 3D model developed by Sung et al[15]
and examined VSMC invasion using a μ-Slide Chemotaxis assay where cells were embedded into FN-
enriched 3D Matrigel matrices in the absence or presence of exogenously-added sEVs (Fig. 5C). We
developed a script to automatically measure cell invasion parameters (track length, cell speed,
straightness, accumulated distance, lateral and vertical displacement and parallel forward motion
index (FMI)). The addition of sEVs had no effect on cell speed, accumulated distance or straightness
(Figs. 5D, 5E, 5F). However, VSMCs invading in the presence of embedded sEVs migrated more
aligned to the FBS gradient compared to cells plated in the absence of sEVs. To test whether this was
a feature of all EV populations we also performed the experiment in the presence of larger EVs
pelleted at the 10,000x*g* centrifugation step and observed no effect suggesting this mechanism was
specific to sEVs (Figs. 5G and 5H). Hence, ECM-associated sEVs have modest influence on VSMC
speed but influence VSMC invasion directionality.

We wondered whether VSMC migration and invasion were affected by the FN cargo in sEVs that was
shown previously to promote tumour cell migration by enhancing FA assembly and cellular adhesion
to FN [14]. Therefore, we measured VSMC adhesion using an adhesion assay where sEVs were added
to FN-coated plates and cells were plated onto these matrices for 1h and firmly attached cells
counted. Surprisingly we observed a marked reduction of VSMC adhesion to FN in the presence of
sEVs (Fig 6A) suggesting that ECM-associated sEV can impair VSMC adhesion. Mesenchymal cell
attachment to the matrix is mediated by transient focal complexes present in the cellular protrusion
as well as centripetal FAs that link the cellular cytoskeleton to the matrix and mediate the pulling
force to move the cell body[12, 13]. Consistently we observed focal complexes in cellular protrusions as
well as more centripetal FAs which were associated with mature actin stress fibres in VSMCs plated
onto FN matrix (Fig S5A). Therefore, we used a live cell spreading assay to monitor cellular adhesion
over time using ACEA's xCELLigence. As expected, FN alone stimulated VSMC spreading and
adhesion (Figs. 6B and 6C). Addition of sEVs did not impact cellular adhesion over the first ~15 min
but then the adhesion was stalled in the presence of sEVs (Fig 6B and 6C) suggesting a defect in
further cellular spreading. Again, to test whether this was specific for the sEV subset, we also tested

the effect of larger EVs pelleted at 10,000x *g* and found that these EVs had no effect on VSMC
adhesion and spreading onto the FN matrix (Fig 6B and 6C). To interrogate the stalling event induced
by sEVs in VSMCs we plated cells and counted the number of FAs, average size as well as distance
from the cell periphery after 30 mins using TIRF microscopy. Once again plating VSMCs onto FN
increased the cell size, number of FAs as well as the number of centripetal FAs as compared to cells
spreading over non-coated plastic (Figs 6E-I). Importantly, addition of sEVs to FN significantly
reduced the number of FAs and the newly formed FAs were formed in close proximity to the cell
periphery (Figs. 6E and 6F). Interestingly, the average FA size was similar between the various
conditions (Fig 5I).

Cellular traction force is generated by the FA gradient upon FA turnover - formation at the leading
edge and disassembly at the tail[12, 13, 46]. To test if sEVs can influence FA turnover in migrating VSMCs
we examined FA turnover in the presence of sEVs. Adhesion sites were visualised by paxillin-RFP
reporter expression and the FA turnover index was calculated by counting FA overlap over time
using a previously developed algorithm to track individual FAs (Fig S5B)[47]. Interestingly, the FA
turnover index remains the same in the presence of sEVs indicating that FA stability was not altered
(6J). These data correlate well with no changes observed in FA average size across various conditions
(Fig 6I) and indicate that sEVs are not influencing FA assembly or "life-cycle" directly. Rho-dependent
activation of actomyosin contractility stabilises FAs by mechanical forces halting fibroblast spreading
onto FN[48, 49]. We hypothesized that sEVs trigger the appearance of premature, peripheral FAs by
activating Rho-dependent cellular contractility. To test whether sEVs can modulate mature FA
contractility we measured individual traction force which is generated by mature adhesion sites in
the absence or presence of sEVs (Fig. S5C). VSMCs transfected with paxillin-RFP were plated on FN
and sEV-covered PDMS pillars and pillar displacements were calculated as previously described[50-52].
Importantly, the traction force was increased in the presence of sEVs (Fig 6K). Altogether these data
suggest that ECM-associated sEVs can trigger the formation of FAs with enhanced pulling force at
the cell periphery.

5. The sEV cargo Collagen VI regulates focal adhesion formation in VSMCs and cell invasion

To identify the components triggering peripheral FA formation we compared the proteomic
composition of sEVs with the larger EV fraction, which had no effect on FA formation. We identified
257 proteins in sEVs and 168 proteins in EVs with 142 proteins common between both datasets (Fig
7A, Tables S6-S8, Fig S6A). Functional enrichment analysis revealed that the top 5 clusters were
related to cell migration - ECM organization, movement of cell or subcellular component, cell
adhesion, leukocyte migration and cell-cell adhesion (Fig 7B, Fig S6B, Table S8). The cell adhesion
cluster included collagen VI (chains COL6A1, COL6A2 and COL6A3), extracellular matrix glycoproteins
(FN and thrombospondin, THBS2, THBS1) as well as EGF-like repeat and discoidin I-like domain-
containing protein (EDIL3) and transforming growth factor-beta-induced protein ig-h3 (TGFBI) (Fig
7B, Fig S6B, Table S8). Notably, these sEV proteins are predominantly involved in cell-matrix
interactions and cellular adhesion. The lectin galactoside-binding soluble 3 binding protein
(LGALS3BP) which regulates cell adhesion and motility was also presented exclusively in sEVs (Table
S6). These cell adhesion modulating proteins including Collagen VI, TGFBI, EDIL and LGALS3BP were
either uniquely or highly enriched in sEVs compared to larger EVs as detected by western blotting
(Fig 7C).

Collagen VI was the most abundant protein in VSMC-derived sEVs (Fig 7B, Table S7) and was
previously implicated in the interaction with the proteoglycan NG2[53] and suppression of cell
spreading on FN[54]. To confirm the presence of collagen VI in ECM-associated sEVs we analysed sEVs
extracted from the 3D matrix using 0.5M NaCl treatment and showed that both collagen VI and FN

are present (Fig 7D). Next, we analysed the distribution of collagen VI using dot-blot. Alix staining
was bright only upon permeabilization of sEV indicating that it is preferentially a luminal protein (Fig
7E). On the contrary, CD63 staining was similar in both conditions showing that it is surface protein
(Fig 7E). Interestingly, collagen VI staining revealed that 40% of the protein is located on the outside
surface with 60% in the sEV lumen (Fig 7E). To test the role of collagen VI in sEV-induced changes in
VSMC spreading we incubated FN-deposited sEVs with an anti-collagen VI antibody to block its
interaction with the proteoglycan NG2[55]. Addition of the anti-collagen VI antibody restored VSMC
spreading on the FN matrix as compared to non-specific IgG treatment (Fig. 7F). Moreover, sEVs
isolated from VSMCs after collagen VI knockdown using siRNA had no effect in the 3D invasion
model on cell speed (Fig 7G) or direction (Fig. 7H) but reduced cell alignment to the gradient as
compared to VSMCs treated with a scrambled siRNA control (Fig. 7I). Finally, we compared the
expression of collagen VI, EDIL3 and TGFBI between the plaque and control aorta tissues (Table S1).
Importantly, collagen VI and TGFBI expression were significantly elevated in the plaque (Fig 7J)
consistent with increased deposition of sEVs during disease progression. Altogether these data
indicate that sEVs can induce early formation of mature peripheral FAs by presenting collagen VI
thus acting to modulate cell directionality upon the invasion (Graphical Abstract).

Discussion.

VSMC migration and invasion to the site of vascular injury is crucial for vessel repair as well as the
pathogenic development of atherosclerotic plaque, however the mechanisms of effective invasion
through the complex vascular ECM meshwork have been poorly studied. Here we show that FN, a
novel marker of unstable plaques[6], enables VSMC migration and invasion by stimulating secretion of
collagen VI-loaded sEVs which decorate the ECM and stimulate FA maturation. Notably, our data are
consistent and extend previous reports showing that EV secretion by tumour cells enhances nascent
adhesion formation hence facilitating tumor cell migration and invasion[14-16]. In particular, we found
that sEVs induce formation of FAs with an enhanced pulling force at the cell periphery thus favouring
actomyosin-mediated contractility which is essential for cell body propulsion and locomotion.

Mesenchymal cell migration begins with protrusive activity at the leading edge followed by the
forward movement of the cell body[13, 49, 56]. Cell directionality is guided via the integrin β1 "probe" on
the tip of cellular protrusions, filopodia and lamellipodia[57]. Cellular protrusions are attached to the
ECM via focal complexes and extended by the physical force generated by the branched network of
actin filaments beneath the plasma membrane[58]. Focal complexes either disassemble or mature into
the elongated centripetally located FAs[48]. In turn, these mature FAs anchor the ECM to actin stress
fibres and the traction force generated by actomyosin-mediated contractility pulls the FAs rearward
and the cell body forward[12, 13]. Here we report that β1 integrin activation triggers sEV release
followed by sEV entrapment by the ECM. Curiously we observed CD63+ MVB transport toward the
filopodia tips as well as inhibition of sEV-secretion with filopodia formation inhibitors suggesting that
sEV secretion can be directly linked to filopodia but further studies are needed to define the
contribution of this pathway to the overall sEV secretion by cells. To gain insight into how sEVs can
stimulate cell invasiveness we explored VSMC protrusive activity using live cell spreading assays and
TIRF imaging. FN stimulated VSMC spreading by increasing the number of FAs, which were formed
centripetally from the cell plasma membrane. FN-associated sEV dramatically ceased VSMC
spreading by inducing early formation of mature FAs at the cell periphery. Cell spreading and
adhesion are orchestrated by the Rho family of small GTPases, Cdc42, Rac1 and Rho with Cdc42 and
Rac1 modulating filopodia and lamellipodia and focal complex formation and Rho controlling FA
maturation and actomyosin-mediated contractility[11, 59, 60]. Interestingly, in fibroblasts Rho is
specifically degraded in cellular protrusions and cellular spreading on FN is accompanied by transient
Rho suppression during the fast cell spreading phase[49, 61, 62]. This phase is followed by Rho activation
leading to formation of actin stress fibres, tension increase on the focal complexes and their
maturation into elongated centripetally located FAs[48, 49, 61, 63]. Excessive Rho activation leads to
premature FA assembly and stress fiber formation resulting in reduced cellular protrusions,
inhibition of cellular spreading and motility on the FN matrix[49, 61, 64-66] Altogether these data indicate
that Rho activity, in mesenchymal cells with extensive FA networks such as fibroblasts or VSMCs, can
slow down cell migration by immobilising cells[13]. However, we found that sEVs were not influencing
mature FA stability. On the contrary, sEV-induced FAs were spatially restricted to the cell periphery
near cellular protrusions and were characterised by an enhanced pulling force activity. Interestingly,
fibroblast polarization is driven by Smurf1-dependent RhoA ubiquitinylation and localised
degradation in cellular protrusions[62]. A recent study indicated that Rho-dependent contractility is
essential for cell invasion in a 3D model and it is tempting to speculate that local Rho activity can
enable VSMC invasion by stabilizing FAs at the leading edge and stimulating actomyosin-mediated
contractility and cell body movement[67]. On the other hand Rho–ROCK activity is critical for the
protease-independent rounded motility of tumour cells and cell types with few adhesion contacts
(ameboid) when Rho-dependent contractile forces generate hydrostatic pressure forming multiple
membrane blebs to invade the ECM[68, 69]. In fact, we observed that an extensive secretion of sEVs

effectively ceased protrusion activity; also VSMCs acquired a rounded morphology when "hovering"
over the FN matrix decorated with sEVs (data not shown). Hence, it will be interesting in future
studies to investigate whether sEVs can stimulate Rho activity by presenting adhesion modulators—
particularly collagen VI—on their surface, thereby guiding cell directionality during invasion.

Collagen VI is a nonfibrillar collagen that assembles into beaded microfilaments upon secretion and
it plays both structural and signalling roles[70]. Interestingly, type VI collagen deposition by interstitial
fibroblasts is increased in the infarcted myocardium and collagen VI knockout in mice improves
cardiac function, structure and remodelling after myocardial infarction via unknown pathways[71, 72].
Immunohistochemical analysis showed that in the healthy vasculature collagen VI is detected in the
endothelial basement membrane in the intima and it also forms fibrillar structures between smooth
muscle cells in the media. In the fibrous plaque collagen VI is diffusely distributed throughout both
the fibrous cap and atheroma[73]. Curiously, treatment of ApoE-/- mice with antibodies to collagen VI
reduced atherosclerosis but its exact role has remained unknown[74]. We and others detected
Collagen VI in sEVs but its functional significance remained unknown[23, 75]. Here we showed that
collagen VI is essential for early FA formation at the VSMC periphery as well as VSMC invasion in a
3D model. We propose that collagen VI modulates FA formation either by activating cellular
signalling or acting as a structural component changing local ECM stiffness[12]. Identified collagen VI
receptors include $\alpha 3\beta 1$ integrin, the cell surface proteoglycan chondroitin sulfate proteoglycan-4
(CSPG4; also known as NG2), and the anthrax toxin receptors 1 and 2[76-79]. Interestingly, a novel,
recently identified Collagen VI signalling receptor (CMG2/ANTXR2) mediates localised Rho activation
upon collagen VI binding[79, 80]. NG2 also activates the Rho/ROCK pathway leading to effective
amoeboid invasiveness of tumour cells which is characterised by excessive blebbing and enhanced
actomyosin contractility[68, 81]. Hence, binding of collagen VI to these receptors can potentially locally
activate Rho and trigger FA formation and actomyosin contractility thus increasing VSMC motility
and directional invasiveness. Of note, it was also reported that NG2 expression is elevated in VSMCs
in the atherosclerotic plaque and NG2 knockout prevents plaque formation[53, 82]. Curiously,
endotrophin, a collagen VI $\alpha 3$ chain-derived profibrotic fragment has recently been associated with
a higher risk of arterial stiffness and cardiovascular and all-cause death in patients with diabetes
type 1 and atherosclerosis, respectively[83, 84].

In addition to collagen VI the unique adhesion cluster in VSMC-derived sEVS also includes EGF-like
repeat and discoidin I-like domain-containing protein (EDIL3), transforming growth factor-beta-
induced protein ig-h3 (TGFBI) and the lectin galactoside-binding soluble 3 binding protein
(LGALS3BP) and these proteins are also directly implicated in activation of integrin signalling and
cellular invasiveness[85-87]. Although we found that collagen VI plays the key role in sEV-induced early
formation of FAs in VSMCs, it is tempting to speculate that the high sEV efficacy in stimulating FA
formation is driven by cooperative action of this unique adhesion complex on the sEVs surface and
targeting this novel sEV-dependent mechanism of VSMC invasion may open-up new therapeutic
opportunities to modulate atherosclerotic plaque development or even to prevent undesired VSMC
motility in restenosis.

We also identified a positive feedback loop and showed that FN stimulates sEV secretion by VSMCs.
sEV secretion is a tightly regulated process which is governed by activation of signalling pathways
including activation of G-protein coupled receptors[39] and alterations in cytosolic calcium[88], small
GTPases Rab7, Rab27a, Rab27b and Rab35[89, 90], vesicular trafficking scaffold proteins including
syntenin[91], sortilin[92] and CD63[93] or accumulation of ceramide[26]. We found that the fibrillar ECM
proteins, FN and collagen I induce sEV secretion by activating the $\beta 1$ integrin/FAK/Src and Arp2/3-
dependent pathways. Our *in vivo* FN staining data are in good agreement with previous studies

showing accumulation of FN in late-stage plaques[6, 7]. Moreover, we observed close co-distribution of
FN and CD81 in the plaque suggesting that accumulating FN matrices can stimulate sEV secretion *in*
vivo. Notably, expression of the major FN receptor, α5β1 integrin is re-activated upon VSMC de-
differentiation following vascular injury[94]. It is thought the α5β1 integrin mediates FN matrix
assembly, whilst β3 integrins are important for cell-matrix interactions[94, 95]. Our data sheds new light
on the functional role of α5β1 as an ECM sensor inducing production of collagen VI-loaded sEVs
which in turn enhance cell invasion in the complex ECM meshwork. Although our small inhibitors
and integrin modulating antibody data clearly indicate that β1 activation triggers sEV secretion via
activation of actin assembly we cannot fully rule out that FN may also be modulating growth factor
activity which in turn contributes to sEV secretion by VSMCs[23]. Excessive collagen and elastin matrix
breakdown in atheroma has been tightly linked to acute coronary events hence it will be interesting
to study the possible link between sEV secretion and plaque stability as sEV-dependent invasion is
also likely to influence the necessary ECM degradation induced by invading cells [96].

In summary, cooperative activation of integrin signalling and F-actin cytoskeleton pathways results in
the secretion of sEVs which associate with the ECM and play a signalling role by controlling FA
formation and cell-ECM crosstalk. Further studies are needed to test these mechanisms across
various cell types and ECM matrices.

1 Materials and Methods

2 *Materials*

Proteins were fibronectin (Cell Guidance Systems, AP-23), collagen I (Gibco, #A1048301), laminin
(Roche, 11243217001), Matrigel (Corning, #356237), Phalloidin-rhodamin (ThermoFisherScientific,
R415). Peptides were Gly-Arg-Gly-Asp-Ser-Pro (GRGDSP, Merck, SCP0157) and scramble control Gly-
Arg-Ala-Asp-Ser-Pro (GRADSP, Merck, SCP0156). All chemical inhibitors were diluted in DMSO and
were: 3-O-Methyl-Sphingomyelin (SMPD3 inhibitor, Enzo Life technologies, BML-SL225-0005),
FAM14 inhibitor (FAK inhibitor, Abcam, ab146739), PP2 (Src inhibitor, Life technologies, PHZ1223),
CK666 (Arp2/3 inhibitor, Abcam ab141231), SMIFH2 (Formin inhibitor, Sigma, S4826). Control siRNA
pool (ON-TARGETplus siRNA, Dharmacon, D-001810-10-05), collagen VI siRNA ON-TARGETplus siRNA
(*COL6A3*, Human), SMARTPool, Horizon, L-003646-00-0005). Antibodies were CD9 (SA35-08 clone,
NBP2-67310, Novus Biologicals), CD63 (BD Pharmingen, 556019), CD81 (BD Pharmingen™, 555676,
B-11, SantaCruz, sc-166029 and M38 clone, NBP1-44861, Novus Biologicals), Syntenin-1 (Abcam,
ab133267), Syndecan-4 (Abcam, ab24511), α-Actinin-4 (Abcam, ab108198), fibronectin (Abcam,
ab2413, ab6328 [IST-9] (3D matrix staining) and F14 clone, ab45688 (clinical samples analysis)), β1
activating (12G10) antibody was previously described[97], 4B4 integrin inhibiting antibody (Beckman
Coulter, 41116015), vinculin (Sigma, V9264), α5 integrin (P1D6, Abcam, ab78614), Myo10 (Sigma,
HPA024223), gelatin-3BP/MAC-2BP (R&D systems, AF2226), EDIL3 antibody (R&D systems,
MAB6046), TGFBI (Sigma, SAB2501486), IgG mouse (Sigma PP54), Anti-collagen Type VI antibody,
clone 3C4 (Sigma, MAB1944) (functional blocking), Anti-collagen Type VI antibody (Abcam, AB6588)
(western blot) Anti-collagen Type VI antibody (Abcam, AB182744) (dot blot) p34-Arc/ARPC2
antibody (Millipore, #07-227), Cortactin, LGALS3BP (R&D, AF2226), GAPDH (ab139416, Abcam), Alix
(Clone 3A9, ThermoFisher Scientific, MA1-83977) Donkey Anti-mouse Secondary IRDye@680 LT (Li-
COR, 926-68022) and Donkey Anti-Rabbit Secondary IRDye@800 CW (Li-COR, 926-32213) DNA
plasmids were: CD63-GFP was kindly provided by Dr Aviva Tolkovsky[98], CD63-pHluorin was
previously described[39], CD63-RFP was previously published[99], Paxillin-RFP was previously
described[97], ARPC2−GFP DNA vector was kind gift from Professor Michael Way laboratory[34], F-
tractin-RFP was kindly provided by Dr Thomas S. Randall (King's College London, UK), Myo10-GFP
(Addgene Plasmid#135403) was previously described[100].

30 *Cell culture*

31   Human VSMCs were isolated as previously described[101] and were cultured in Dulbecco's modified
32   Eagle medium (Sigma) supplemented with 20% fetal calf serum, 100 U/ml penicillin, 100 μg/ml
33   streptomycin and 2 mmol/L L-glutamine (Gibco) and used between passages 4 and 12.

34 *VSMC adhesion*

24 well plate (Corning Costar) was incubated with 5μg/ml fibronectin in PBS overnight at +4°C. Next,
sEVs (10μg/ml) diluted in PBS were added to the wells and incubated overnight at +4°C. Wells were
washed with PBS, blocked with PBS-1% BSA for 30min at 37°C and washed 3 times with PBS again.
VSMCs were incubated in serum free media, M199 supplemented with 0.5% BSA, 100 U/ml penicillin
and 100μg/ml streptomycin overnight and removed by brief trypsin treatment. Next, 20,000 cells
were added to each well and incubated for 30min at +37°C. Unattached cells were washed away
with PBS and attached cells were fixed with 3.7% PFA for 15 min at 37°C. Cells were washed with
H_2O 3 times and stained with 0.1% crystal violet for 30min at 10% CH_3COOH for 5min at room
temperature. Samples were transferred to 96 well plate and absorbance was measured at 570nm
using the spectrophotometer (Tecan GENios Pro).

iCELLigence adhesion protocol

E-Plate L8 (Acea) was coated with FN (5µg/mL) in PBS at 4°C overnight. Next, E-Plate L8 was gently

washed once with cold PBS followed by the incubation with EV or sEV (10µg/ml) at 4°C overnight.

The following day, E-Plate L8 was incubated with 0.1% BSA in PBS for 30min at 37°C to block non-

specific binding sites and washed with PBS before adding 250µL of M199 media supplemented with

20% exosome-free FBS. VSMCs were serum-deprived by incubation in M199 media supplemented

with 0.5% BSA overnight, were passaged with trypsin and resuspended in media M199

supplemented with 20% exosome-free FBS to a final concentration 80,000 cells/mL. Cell aliquot

(250µL) was added to E8-plate L8 which was then transferred to iCELLigence device for the adhesion

assay. Cell adhesion was measured at 37°C with time intervals 20sec for 1h.

Focal adhesion turnover assay

iBIDI 35mm dishes were incubated with FN (5µg/ml) in PBS overnight at +4⬚C and treated with

exosomes (10µg/ml) diluted in PBS overnight at +4⬚C. 500,000 cells were transfected with Paxillin-

RFP/Myo10-GFP by using electroporation (see below) and plated on iBIDI 35mm dishes coated with

FN and sEVs in DMEM supplemented with 20% exosome-free FBS, 100 U/ml penicillin, and 100

µg/ml streptomycin. Next, cells were transferred to DMEM supplemented with 20% exosome-free

FBS, 10mM HEPES, 100 U/ml penicillin, and 100 µg/ml streptomycin and images were acquired

(Nikon Spinning Disk) every 1min for 30min at 37°C. FA images were extracted from timelapses and

FA turnover was quantified using Mathematica by producing the adhesion map as previously

described[47].

Confocal spinning disk microscopy and live cell imaging

VSMC were transfected using electroporation (see below) and plated onto FN-coated 35mm iBIDI

glass bottom dish and incubated for 48h. Then cells were transferred to DMEM supplemented with

20% exosome-free FBS, 10mM HEPES, 100 U/ml penicillin, and 100 µg/ml streptomycin and images

were acquired every 1min for 20min at 37°C using a Nikon Ti-E (inverted) microscope equipped with

a Yokogawa spinning disk and a Neo 5.5 sCMOS camera (Andor) and 60x or 100x/1.40 NA Plan Apo λ

oil objectives (Nikon) were used. Images were acquired using NIS Elements AR 4.2 software. Cells

were maintained at 37°C, 5% CO_2 throughout the experiment via a CO_2 chamber and a temperature-

regulated Perspex box which housed the microscope stage and turret.

iSIM

Slides was imaged and super-resolved images collected using a Visitech-iSIM module coupled to a

Nikon Ti-E microscope using a Nikon 100x 1.49NA TIRF oil immersion lens. Blue fluorescence was

excited with a 405nm laser and emission filtered through a 460/50 filter. Green fluorescence was

excited with a 488nm laser and emission filtered through a 525/50 filter. Red Fluorescence was

excited with a 561nm laser and emission filtered through a 630/30 filter. Far Red fluorescence was

excited with a 640nm laser and emission filtered through a 710/60 filter. Images were collected at

focal planes spaced apart by 0.08um. Data was deconvolved using a Richardson-Lucy algorithm

utilizing 20 iterations using the Nikon deconvolution software module (Nikon Elements). 65 stack

images were taken with Z step 0.08µm. ImageJ (153t) or NIS Elements (5.21.00, built1483, 64bit)

software were used for image analysis and assembly.

Cell electroporation

iBidi dishes were coated with FN (5µg/ml) in PBS for 1 hour at 37°C or overnight at 4°C. VSMCs were

grown to 70% confluence, washed twice with EBSS and detached by trypsin treatment for 5min at

+37°C. Cells (500,000) were mixed with plasmids (2.5µg each) in electroporation buffer (100µl,
Lonza) and transfected using either Nucleofector II (program U25) or Nuclefector III (program CM-
137).

Isolation of apoptotic bodies, extracellular vesicles and small extracellular vesicles

Flasks (T150) were incubated with PBS or FN (5µg/ml) in PBS overnight at +4°C and VSMCs ($\approx 10^6$
cells) were plated and incubated for 16h at +37°C. Next, cells were washed with EBSS 3 times and
incubated with DMEM supplemented with 0,1% BSA, 100 U/ml penicillin, and 100 µg/ml
streptomycin. Conditioned media was collected and centrifuged at 2,500 rpm (Thermo Scientific
Heraeus Multifuge 3SR+ centrifuge, rotor Sorvall 75006445) for 5min to remove apoptotic bodies
(AB, 1.2K pellet). Supernatant was transferred to centrifugation tube (Nalgene™ Oak Ridge High-
Speed Polycarbonate Centrifuge Tubes, Thermo Fisher Scientific, 3138-0050) and centrifuged 10,000
x*g* for 30min. 10K pellet (EVs) was collected, washed in PBS once and kept at -80°C until further
analysis. 10K supernatant was transferred to ultracentrifugation tubes (Polycarbonate tubes for
ultracentrifugation, BeckmanCoulter, 355647) and centrifuged at 35,000 rpm (100,000x*g*) for 40min
at 4°C (Fixed-angle rotor, Beckman Coulter Optima Max Unltracentrifuge). 100K pellet (sEVs) was
washed with PBS twice and re-suspended in PBS. EV and sEV pellets were kept at -80°C for the
proteomic analysis and freshly isolated sEVs were used for all other assays.

EV and sEV proteomic analysis

EV and sEV samples were submitted to the CEMS Proteomics Facility in the James Black Centre,
King's College London for mass spectrometric analysis. The data were processed by Proteome
Discover software for protein identification and quantification. Scaffold 5 proteome software was
utilized to visualize differential protein expression. Individual proteins intensities were acquired and
transformed to log2 scale before Hierarchical clustering analysis based on Euclidean distance and k-
means processing. Differential genes defined by multiple groups comparation were applied to
perform the gene ontology functional assay through using the DAVID Gene Ontology website
(https://david.ncifcrf.gov/). Selected genes which showed abundant in each condition were applied
to generate Venn diagram through Interacti website (http://www.interactivenn.net/).

Cell lysis

Cells were washed with PBS and removed by cell-scraper in 1ml of PBS. The cells were pelleted by
centrifugation at 1000x*g* for 5min and cell pellets were kept at -80°C until further analysis. To
prepare cell lysates, pellets were incubated with lysis buffer (0.1M TrisHCl(pH8.1), 150 mM NaCl, 1%
Triton X-100 and protease inhibitors cocktail (Sigma, 1:100)) for 15min on ice. Cell lysates were
subjected to ultrasound (Branson Sonifier 150), centrifuged at 16,363x*g* for 15 minutes (Eppendorf)
at 4°C and supernatants were collected and analysed by western blotting.

CD63-bead capturing assay

CD63-bead capturing assay was conducted as previously described[23] with some modifications. In
brief, CD63 antibody (35µg) was immobilised on 1×10^8 4µm aldehyde-sulfate beads (Invitrogen) in
30mM MES buffer (pH 6.0) overnight at room temperature and spun down by centrifugation
(3,000x*g*, 5min). Next, beads were washed 3 times with PBS containing 4%BSA and kept in PBS
containing 0,1% Glycine and 0,1% NaN$_3$ at +4°C. VSMCs were plated onto 24 well plate (10,000 cells
per well) and incubated in the complete media overnight. Next, cells were washed 3 times with EBSS
and incubated in M199 supplemented with 2.5% exosome-free FBS in the presence of absence of
inhibitors for 2h. Then, conditioned media was replaced once to the fresh aliquot of 2.5% exosome-

free FBS in the presence or absence of inhibitors and cells were incubated for 18-24h. VSMC conditioned media was collected and centrifuged and centrifuged at 2,500xg for 5 min. VSMCs were detached from the plate by trypsin treatment and viable cells were quantified using NC3000 (ChemoMetec A/S). The conditioned media supernatants were mixed with 1µL of anti-CD63-coated beads and incubated on a shaker overnight at +4°C. Beads were washed with PBS-2%BSA and incubated with anti-CD81-PE antibody (1:50 in PBS containing 2% BSA) for 1h at room temperature. Next, beads were washed with PBS-2%BSA and PBS and analysed by flow cytometry (BD Accuri™ C6). sEV secretion (fold change) was calculated as ratio of Arbitrary Units (fluorescence units x percentage of positive beads and normalized to the number of viable VSMCs) in the treatment and control conditions.

Exosome labelling with Alexa Fluor® 568 C5 Maleimide

Exosomes (10µg/ml) were incubated with 200µg/mL Alexa Fluor® 568 C5 Maleimide (ThermoFisherScientific, #A20341) in PBS for 1h at room temperature. An excessive dye was removed by using Exosome Spin Columns (MW 3000, ThermoFisherScientific, #4484449) according to the manufacturer protocol.

Fetuin-A and Fibronectin labelling and uptake studies.

Bovine fetuin-A (Sigma) and Fibronectin were labelled using an Alexa488 (A10235) and Alex568 (A10238) labelling kits in accordance with the manufacturer's protocol (Invitrogen). VSMCs were serum-starved for 16h and then incubated with Alexa488-labeled fetuin-A (10µg/mL) and Alexa555-labeled fibronectin (10µg/mL) from 30 to 180 minutes at 37°C.

VSMC and beads-coupled sEVs flow cytometry.

VSMC were removed from the plate by brief trypsin treatment, washed with M199 media supplemented with 20% FBS and resuspended in HBSS supplemented with 5%FBS. Cells were kept on ice throughout the protocol. Next, cells (200,000) were incubated with primary antibody for 30min on ice, washed with HBSS-5%FBS and incubated with secondary fluorescently-labelled antibody 30min on ice. Then cells were washed twice with HBSS-5%FBS and once with PBS and analysed by flow cytometry (BD FACScalibur, BD Bioscience).

Flow cytometry analysis of beads-coupled sEVs was conducted as described before[102]. In brief, exosomes (10µg) isolated by differential centrifugation were coupled to 4µm surfactant-free aldehyde/sulfate latex beads (Invitrogen) and were incubated with primary antibody and fluorescently labeled secondary antibody and analysed by flow cytometry on BD FACScalibur (BD Bioscience). Data were analysed using the Cell Quest Software (BD). Cells and beads stained with isotype-control antibodies were used as a negative control in all experiments.

Nanoparticle Tracking Analysis

Exosomes were diluted 1:150 and analysed using LM-10 using the light scattering mode of the NanoSight LM10 with sCMOS camera (NanoSight Ltd, Amesbury, United Kingdom). 5 frames (30 s each) were captured for each sample with camera level 10 and background detection level 11. Captured video was analysed using NTA software (NTA 3.2 Dev Build 3.2.16).

Western blotting and dot blotting

For western blotting, cell lysates or apoptotic bodies, microvesicles and exosomes in Laemmli loading buffer were separated by 10% SDS-PAGE. Next, separated proteins were transferred to ECL immobilon-P membranes (Millipore) using semi-dry transfer (BioRad). Membranes were incubated

in the blocking buffer (TBST supplemented with 5% milk and 0.05% tween-20) for 1h at room
temperature, then incubated with primary antibody overnight at +4°C. Next, membranes were
washed with blocking buffer and incubated with secondary HRP-conjugated antibody and washed
again in PBS supplemented with 0.05% tween-20 and PBS. Protein bands were detected using ECL
plus (Pierce ECL Western Blotting Substrate, #32109). Alternately, membranes were incubated with
the secondary fluorescent antibody (Alexa fluor antibody) and washed again with TBST or PBST.
Blots were detected using Odyssey Licor.

Dot blot was conducted as previously described[103]. In brief, an aliquot of isolated sEVs (0.5ug) was
absorbed onto nitrocellulose membrane (Odyssey® P/N 926-31090) for 30min at room temperature
and air dried. Next, membranes were blocked with PBS-3% BSA in PBS and incubated with primary
antibodies (anti-CD63 1:500, anti-COLVI 1:200, anti-Alix, 1:500) in the absence or presence 0.2%
(v/v) Tween-20 for 16h at the cold room (+4C) room temperature. Membranes were washed 3 times
in PBS-3% BSA, and were incubated with Donkey Anti-Rabbit Secondary IRDye@800 CW (Li-COR,
926-32213) and Donkey Anti-Mouse Secondary IRDye@680 LT (Li-COR, 926-68022) (1:10,000),
washed with PBS-0.2% tween20 and PBS and visualized and quantified using Odyssey ® CLx Infrared
Imaging System.

Generation of 3D Extracellular Matrix

Generation of 3D matrices were conducted as previously described[24] with some modifications. In
brief, plates or glass coverslides were covered with 0.2% gelatin in PBS for 1h at 37°C. Wells were
washed with PBS and fixed with 1% glutaraldehyde in PBS for 30min at room temperature. Plates
were washed with PBS and incubated with 1M ethanolamine for 30min at room temperature, then
washed with PBS again. VSMCs (5 ×10^5 per well) were plated and cultured for 24h. Then the medium
was replaced with medium containing 50µg/ml of ascorbic acid and cells were incubated for 9days.
The medium was replaced every 48h. To extract matrix, cells were rinsed with PBS and incubated
with pre-warmed extraction buffer (20mM NH$_4$OH containing 0.5%Triton X-100) for 3min until intact
cells are not seen. Next, equal volume of PBS was added to extraction buffer and plates were
incubated for 24h at +4°C. Plates were washed with PBS twice and kept with PBS containing 100
U/ml penicillin and 100 µg/ml streptomycin up to 2 weeks at +4°C.

sEVs isolation from 3D Extracellular Matrix

VSMCs were cultured on dishes pre-coated with 0.1% gelatin as described above, with minor
modifications. Briefly, cells were maintained in DMEM supplemented with EV-free fetal bovine
serum, 1% penicillin-streptomycin, and 50µg/ml ascorbic acid. Cultures were maintained for 10
days, with media changes and collections performed every 2–3 days.At the end of the culture
period, the medium was gently removed and collected as the "Media" fraction. Conditioned media
was clarified by centrifugation at 2,500 × g for 15 minutes to remove cellular debris; the resulting
supernatant was retained for ExoView analysis and small EV (sEV) isolation. The cell monolayer was
then rinsed with phosphate-buffered saline (PBS) and incubated with extraction buffer (0.5 M NaCl,
20 mM Tris, pH 7.5) for 4 hours at room temperature as previously described[42]. This NaCl-extracted
ECM fraction, termed the "loosely-bound sEV fraction"," was collected and centrifuged at 2,500 × g
for 15 minutes. The supernatant was retained for ExoView analysis and sEV isolation. Following NaCl
extraction, cells were treated with 0.1% sodium dodecyl sulfate and 25 mM
ethylenediaminetetraacetic acid (EDTA), incubating at 37°C for 10 minutes to lyse any remaining
cellular material. To extract the remaining ECM-associated fraction, the matrix was incubated with 4
M guanidine hydrochloride (GuHCl) in 50 mM sodium acetate buffer (pH 5.8) for 48 hours at room
temperature. The GuHCl fraction, referred to as "strongly-bound sEV fraction" was collected and

1. centrifuged at 2,500 × g for 15 minutes at 4°C; the supernatant was retained for ExoView analysis.
2. sEVs from all fractions were isolated by ultracentrifugation as described above.

ExoView analysis of ECM-derived sEV fractions

4. ECM-derived EV fractions (media, loosely bound and strongly trapped sEVs) were prepared as
5. detailed above. For ExoView analysis, each fraction was diluted 1:2 in the manufacturer-supplied
6. incubation solution. Leprechaun Exosome Human Tetraspanin chips (Unchained Labs, REF 251-1044)
7. were used according to the manufacturer's protocol.

8. Briefly, 35–50⬚µL of diluted sample (1:2-1:10) was applied to each chip well. Chips were incubated
9. overnight at room temperature in a sealed humidity chamber to allow EV capture onto surface-
10. immobilized antibodies. The following day, chips were washed per the supplied protocol to remove
11. unbound material.

12. Secondary immunostaining was performed using fluorescently labeled antibodies provided in the
13. Leprechaun Exosome Human Tetraspanin Kit (CD9, CD63, and CD81, Unchained Labs, LOT
14. UL038460001F). After antibody incubation, chips were washed thoroughly as recommended and air-
15. dried briefly.

16. Chips were scanned on an ExoView R200 platform (Unchained Labs). Imaging and data analysis were
17. performed using ExoView Analyzer software, with spot intensity and particle counts determined for
18. each tetraspanin (CD9, CD63, CD81) according to the manufacturer's guidelines. Data were exported
19. for downstream quantitative and statistical analysis.

Immunocytochemistry

21. VSMCs (10,000 cells per well) were plated onto the coverslips and incubated for 48h. Next, cells
22. were treated with inhibitors in M199 supplemented with 2.5% exosome-free FBS and fixed using
23. 3.7% paraformaldehyde for 15 minutes at +37°C. Cells were then washed with PBS, permeabilised
24. with PBS-0,1% triton X-100 for 5 min at room temperature and washed with PBS again. Cells were
25. blocked with PBS-3% BSA for 1 hour at room temperature and incubated with primary antibodies
26. overnight at +4°C. Following washing 3 times with PBS-3% BSA cells were incubated with secondary
27. fluorescently-labelled antibodies in the dark for 1 hour at room temperature, stained with DAPI for
28. 5min and mounted onto 75-Superdex slides using gelvatol mounting media and analysed by spinning
29. disk confocal microscopy as above (Nikon). Primary antibodies were used in the following dilutions:
30. CD63 1:500, Cortactin 1:500, Arp2C 1:500, Myo10 1:250, Vinculin 1:500, fibronectin 1:500, CD81
31. 1:500, Rhodamine phalloidin 1:200 and secondary fluorescently labelled antibodies 1:200.

Focal adhesion turnover/Individual traction force

33. PDMS pillar (500 nm diameter, 1.3 µm height, 1µm centre-to-centre) substrates were prepared as
34. described previously[104]. Briefly, CdEq3: $T_{tilt}(v) = a\frac{(1+v)}{2\pi}\left\{2(1-v) + \left(1 - \frac{1}{4(1-v)}\right)\right\}$SeS/ZnS alloyed
35. quantum dots (490nm, Sigma) were spun on the master 30s at 10,000rpm with a 150i spin processor
36. (SPS), before the addition of PDMS. PDMS (Sylgard 184, Dow Corning) was mixed thoroughly with its
37. curing agent (10:1), degassed, poured over the silicon master, placed upside-down on a plasma-
38. treated coverslip-dish (Mattek), or coverslip 4-well dishes (Ibidi) and cured at 80C for 2h. The mold
39. was then removed and the pillars were incubated with fibronectin for 1h at 37C, after which pillars
40. were incubated with sEVs (10µg/ml in PBS) at 4°C overnight.
41. VSMC previously transfected with Paxillin-RFP by using electroporation as above were plated on the
42. Pillars and imaged on a Nikon Eclipse Ti-E epifluorescent microscope with a 100x 1.4NA objective,
43. Nikon DS-Qi2 Camera and a Solent Scientific chamber with temperature and CO_2 control. For

calculation of pillar displacements, a perfect grid was assumed and deviations from the grid were calculated using reference pillars outside the cell. The traction stress was calculated for a 10 µm wide region at the cell edge that was enriched in paxillin staining, taking into account all pillar displacements above a ~20 nm noise level, which was calculated from pillars outside the cells.

The Stiffness of the pillars was calculated as described by Ghibaudo et al[52] but taking into account substrate warping as described by Schoen et al (Eq1-5)[51]:

$$Eq1:\ k_{bend} = \frac{3}{64}\pi E \frac{D^4}{H^3};$$

$$Eq2:\ corr = \frac{\frac{16}{3}\left(\frac{L}{D}\right)^3}{\left(\frac{16}{3}\left(\frac{L}{D}\right)^3 + \frac{7+6vL}{3\ D} + 8T_{tilt}(v)\left(\frac{L}{D}\right)^2\right)};$$

$$Eq3:\ T_{tilt}(v) = a\frac{(1+v)}{2\pi}\left\{2(1-v) + \left(1 - \frac{1}{4(1-v)}\right)\right\}$$

$$Eq4:\ k = k_{bend} * corr;$$

$$Eq5:\ E_{Eff} = \frac{9k}{4\pi a};$$

Protein concentration

Protein concentration was determined by DC protein assay (BioRad).

Focal adhesion TIRF Imaging

µ-Slides (iBidi) were coated with FN (5µg/mL) in PBS at 4°C overnight. Next, µ-Slides were gently washed once with cold PBS following incubation with EV or sEV at 4°C overnight. The following day µ-Slides were incubated with 0.1% BSA in PBS for 30 minutes at 37°C to block non-specific binding sites and washed with PBS three times and 250µL of M199 media supplemented with 20% sEV-free FBS was added to each well. VSMCs were incubated in M199 media supplemented with 0.5% BSA overnight, detached with trypsin and resuspended to 80,000 cells/mL in M199 supplemented with 20% FBS. Cells (250µL) were added to µ-Slides and were incubated for 30min at 37°C. VSMC were fixed with 4% PFA for 10 minutes at 37°C, washed with PBS three times and were permeabilised with PBS supplemented with 0.2% Triton X-100 for 10 minutes at 37°C. Next, VSMC were washed three times with PBS for 5 minutes and were incubated with 3% BSA in PBS for 1h at room temperature. VSMC were incubated with anti-vinculin antibody (1:400) in blocking buffer at 4°C overnight. Next, VSMC were washed with PBS for 5min and incubated with secondary antibody (1:500, AlexaFluor488) diluted in blocking buffer for 1h at room temperature. Cells were incubated with CellMaskRed (ThermoFisherScientific) and DAPI (0.1 mg/mL) for 5min, washed and analysed by TIRF microscopy.

TIRF Microscopic images were collected on a Nikon Ti2 upright microscope with a TIRF illuminator (Nikon) attached using a 100x oil immersion (1.49 NA) objective lens. In each fluorescent channel the sample was excited with a laser beam at an angle indicated as follows (488nm at 62.3°, 561nm at 62.5° and 640nm at 65.5°). All emission light was filtered through a quad channel filter set (Chroma 89000) in the upper filter turret and a secondary bandpass emission filter in the lower turret position was used as follows for each channel (488nm excited: filter BP 525/50 - Chroma; 561nm excited: filter BP 595/50 - Chroma; 640nm excited: filter BP 700/75 - Chroma). Light was then passed to a

Hamamatsu Orca-Flash 4v3.0 sCMOS camera and exposures for channels was set as follows (488nm:
100ms, 561nm: 100ms, 640nm:50ms) with 16bit channel images collected.

Acquisition was all controlled through NIS-Elements software (Nikon). Analysis of TIRF images was
completed using NIS-Elements General Analysis Module. A general description of the process
follows: Cell Area measurements were determined by use of an intensity threshold on the HCS
CellMask Deep Red channel at an intensity of 100 AU (grey levels or arbitrary units) above the
background to mask the cell. The Cell area, whole cell Vinculin and Far-Red intensity measurements
calculated from this binary region. In addition, an inversion of the cell mask binary was generated to
have a means to measure distance to edge of the cell.

Vinculin derived Foci were masked using an intensity-based threshold (Above 10000 AU) and a size
minimum for objects was set to 0.5μm^2 to eliminate smaller non foci objects from the binary mask.
The foci mask was then limited to the cell mask regions to allow interpretation of the foci in the cell
area. Minimum Distance to the outside of the cell for each foci was calculated by measurement of
the nearest distance to the inverted non-cell mask. The average distance was then recorded for each
image. For each foci intensity measurements were taken for the Vinculin and FarRed channel and
the Mean intensity of the foci were recorded.

2D VSMC migration time lapse assay

24 well plates were incubated with FN (5μg/ml) in PBS overnight at +4°C and treated with sEVs
(10μg/ml) diluted in PBS overnight at +4°C. VSMC (8,000 cells per well) were plated on the plate in
M199 supplemented with 20% exosome-free FBS and incubated for 16h. Cells were washed twice
with DMEM supplemented with 2.5% exosome-free FBS and incubated in this media for 2hrs +/- 3'-
OMS. Media was replaced for a fresh aliquot and cells were imaged using Opera Phenix High Content
Screening System (PerkinElmer) every 6min20sec for 8h in a transmitted light channel.
Quantification of cellular velocity and directionality was performed in Harmony 4.9 (Perkin Elmer)
and the following filters were applied to the raw data: "Number of Timepoints" >= 5, "First
Timepoint" < 5.

3D VSMC Invasion time lapse assay

Acquisition. Migration assay were completed using IBidi Chemotaxis μ-slides (Ibidi μ-slides #80326).
VSMC were deprived overnight in M199 supplemented with 0.5% BSA, counted and mixed with
Matrigel supplemented with/without sEV (5μg total in 100μl volume) to a final cell concentration of
ca. 3 x 10^6 cells/ml and stained with Draq5 according to the manufacturer protocol. Slides were left
for 30min at 37°C for gelation and chemoattractant-free medium (65uL of M199 supplemented with
0.5% BSA) was filled through Filling Port E. Next, the empty reservoir was filled by injecting
chemoattractant medium (65uL of M199 supplemented with 20% FBS exosome-free through Filling
Port C. 3 slides were loaded and cells were imaged with the Opera Phenix (Perkin Elmer) high
content imaging platform (10x Objective NA 0.3). Images were collected for digital phase contrast
and a fluorescence channel for Draq5. Fluorescence was obtained by exciting with a 640nm laser and
emission light was filtered with a 700/75 bandpass filter to a sCMOS camera. Images were recorded
every 10min for 12h in several locations for each slide.

Migration tracking and analysis of tracking. Analysis of cell motion was completed on Draq5 images
using Harmony Software (Perkin Elmer) with measurements of cell area, speed and direction were
obtained for each cell/track and timepoint. Cells were not selected for tracking if they were in
proximity to other cells of a distance of 35μm. The measurements and object data were exported
and further analysis was performed. Data for each track included total length (timepoints),

speed(μm/s), Straightness, accumulated distance (μm), lateral displacement (μm) and vertical
displacement (μm).

Data analysis. This data was then analysed using an in-house script using Python library Pandas.
Tracks were discarded if they were less than 3 time-points in length. Means and Standard Deviations
were calculated for track length (timepoints), Speed (μm/s), Track Straightness and accumulated
distance. For each track the Parallel FMI (forward motion index) was calculated by taking a ratio of
vertical displacement to accumulated distance. The means and standard deviations for these were
calculated and reported for each condition similarly. Track Straightness is calculated as the ratio of
total displacement over total track length. Accumulated distance is the total length of the track from
the first to last time point of the track.

Clinical samples

Patients enrollment. The study was approved by the Local Ethical committee of the Research
Institute for Complex Issues of Cardiovascular Diseases (Kemerovo, Russia, protocol number
20200212, dates of approval: 12 February 2020), and a written informed consent was provided by all
study participants after receiving a full explanation of the study. The investigation was carried out in
accordance with the Good Clinical Practice and the Declaration of Helsinki. Criteria of inclusion were:
1) performance of carotid endarterectomy due to chronic brain ischemia or ischemic stroke; 2) a
signed written informed consent to be enrolled. A criterion of exclusion was incomplete
investigation regardless of the reason; in this case, we enrolled another subject with similar age,
gender and clinicopathological features who met the inclusion criteria. Cerebrovascular disease
(chronic brain ischemia and ischemic stroke) as well as comorbid conditions (arterial hypertension,
chronic heart failure, chronic obstructive pulmonary disease, asthma, chronic kidney disease,
diabetes mellitus, overweight and obesity) were diagnosed and treated according to the respective
guidelines of European Society of Cardiology, Global Initiative for Chronic Obstructive Lung Disease,
Global Initiative for Asthma, Kidney Disease: Improving Global Outcomes, American Diabetes
Association, and European Association for the Study of Obesity. eGFR was calculated according to
the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation. Extracranial artery
stenosis was assessed using the color duplex screening (Vivid 7 Dimension Ultrasound System,
General Electric Healthcare). Data on age, gender, smoking status and pharmacological anamnesis
were collected at the time of admission. In total, we enrolled 20 patients. The detailed
characteristics of the study samples are presented in Table S1.

Sample collection and preparation. Carotid atherosclerotic plaques (n = 14) and adjacent intact
arterial segments (n = 14) were pairwise excised during the carotid endarterectomy and divided into
3 segments each. The first segment was snap-frozen in the optimal cutting temperature compound
(Tissue-Tek, 4583, Sakura) using a liquid nitrogen and was then sectioned on a cryostat (Microm
HM525, 387779, Thermo Scientific). To ensure the proper immunofluorescence examination, we
prepared 8 sections (7μm thickness), evenly distributed across the entire carotid artery segment, per
slide. The second and third segments were homogenised in TRIzol Reagent (15596018, Thermo
Fisher Scientific) for RNA extraction or in T-PER Tissue Protein Extraction Reagent (78510, Thermo
Fisher Scientific) supplied with Halt protease and phosphatase inhibitor cocktail (78444, Thermo
Fisher Scientific) for the total protein extraction according to the manufacturer's protocols.
Quantification and quality control of the isolated RNA was performed employing Qubit 4
fluorometer (Q33238, Thermo Fisher Scientific), Qubit RNA BR assay kit (Q10210, Thermo Fisher
Scientific), Qubit RNA IQ assay kit (Q33222, Thermo Fisher Scientific), Qubit RNA IQ standards for
calibration (Q33235, Thermo Fisher Scientific) and Qubit assay tubes (Q32856, Thermo Fisher
Scientific) according to the manufacturer's protocols. Quantification of total protein was conducted

using BCA Protein Assay Kit (23227, Thermo Fisher Scientific) and Multiskan Sky microplate
spectrophotometer (51119700DP, Thermo Fisher Scientific) in accordance with the manufacturer's
protocol.

Immunofluorescence examination. Upon the sectioning, vascular tissues were dried at room
temperature for 30min, fixed and permeabilised in ice-cold acetone for 10min, incubated in 1%
bovine serum albumin (Cat. No. A2153, Sigma-Aldrich) for 1h to block non-specific protein binding,
stained with unconjugated mouse anti-human CD81 (M38 clone, NBP1-44861, 1:100, Novus
Biologicals) and rabbit anti-human fibronectin (F14 clone, ab45688, 1:250, Abcam) primary
antibodies and incubated at 4°C for 16h. Sections were further treated with pre-adsorbed donkey
anti-mouse Alexa Fluor 488-conjugated (ab150109, 1:500, Abcam) and donkey anti-rabbit Alexa
Fluor 555-conjugated secondary antibodies (ab150062, 1:500, Abcam) and incubated for 1h at room
temperature. Between all steps, washing was performed thrice with PBS (pH 7.4, P4417, Sigma-
Aldrich). Nuclei were counterstained with 4ᵇ,6-diamidino-2-phenylindole (DAPI) for 30min at room
temperature (10μg/mL, D9542, Sigma-Aldrich). Coverslips were mounted with ProLong Gold
Antifade (P36934, Thermo Fisher Scientific). Sections were examined by confocal laser scanning
microscopy (LSM 700, Carl Zeiss). Colocalization analysis (n=12 images) was performed using the
respective ImageJ (National Institutes of Health) plugins (Colocalisation Threshold and Coloc2). To
evaluate the colocalization, we calculated Pearson's r above threshold (zero-zero pixels),
thresholded Mander's split colocalisation coefficient (the proportion of signal in each channel that
colocalizes with the other channel) for both (red and green) channels, percent volume colocalised
with each channel, and intensity volume above threshold colocalised with each channel in both
neointima and media.

Reverse transcription-quantitative polymerase chain reaction (RT-qPCR). Reverse transcription was
carried out utilising High Capacity cDNA Reverse Transcription Kit (4368814, Thermo Fisher
Scientific). Gene expression was measured by RT-qPCR using the customised primers (500nmol/L
each, Evrogen, Moscow, Russian Federation, Table S5), cDNA (20ng) and PowerUp SYBR Green
Master Mix (A25778, Thermo Fisher Scientific) according to the manufacturer's protocol for Tm ≥
60°C (fast cycling mode). Technical replicates (n=3 per each sample collected from one vascular
segment) were performed in all RT-qPCR experiments. The reaction was considered successful if its
efficiency was 90-105% and R^2 was ≥ 0.98. Quantification of the *CD9, CD63, CD81, COL6A3, EDIL3,*
CSPG4, TGFBI, FN1, and *MYADM* mRNA levels in carotid atherosclerotic plaques (n=5) and adjacent
intact arterial segments (n = 5) was performed by using the $2^{-\Delta\Delta Ct}$ method. Relative transcript levels
were expressed as a value relative to the average of 3 housekeeping genes (*ACTB, GAPDH, B2M*).

Western blotting. Equal amounts of protein lysate (15μg per sample) of carotid atherosclerotic
plaques (n=12), adjacent intact arterial segments (n=12), and plaque-derived sEVs (n=6) were mixed
with NuPAGE lithium dodecyl sulfate sample buffer (NP0007, Thermo Fisher Scientific) at a 4:1 ratio
and NuPAGE sample reducing agent (NP0009, Thermo Fisher Scientific) at a 10:1 ratio, denatured at
99°C for 5 minutes, and then loaded on a 1.5mm NuPAGE 4-12% Bis-Tris protein gel (NP0335BOX,
Thermo Fisher Scientific). The 1:1 mixture of Novex Sharp pre-stained protein standard (LC5800,
Thermo Fisher Scientific) and MagicMark XP Western protein standard (LC5602, Thermo Fisher
Scientific) was loaded as a molecular weight marker. Proteins were separated by the sodium dodecyl
sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) at 150V for 2h using NuPAGE 2-(N-
morpholino)ethanesulfonic acid SDS running buffer (NP0002, Thermo Fisher Scientific), NuPAGE
Antioxidant (NP0005, Thermo Fisher Scientific), and XCell SureLock Mini-Cell vertical mini-protein gel
electrophoresis system (EI0001, Thermo Fisher Scientific). Protein transfer was performed using
polyvinylidene difluoride (PVDF) transfer stacks (IB24001, Thermo Fisher Scientific) and iBlot 2 Gel

Transfer Device (IB21001, Thermo Fisher Scientific) according to the manufacturer's protocols using a standard transfer mode for 30-150 kDa proteins (P0 – 20 V for 1min, 23 V for 4min, and 25 V for 2 min). PVDF membranes were then incubated in iBind Flex Solution (SLF2020, Thermo Fisher Scientific) for 1h to prevent non-specific binding.

Blots were probed with rabbit primary antibodies to CD9, fibronectin or CD81, and GAPDH (loading control). Horseradish peroxidase-conjugated goat anti-rabbit (7074, Cell Signaling Technology) or goat anti-mouse (AP130P, Sigma-Aldrich) secondary antibodies were used at 1:200 and 1:1000 dilution, respectively. Incubation with the antibodies was performed using iBind Flex Solution Kit (SLF2020, Thermo Fisher Scientific), iBind Flex Cards (SLF2010, Thermo Fisher Scientific) and iBind Flex Western Device (SLF2000, Thermo Fisher Scientific) during 3h according to the manufacturer's protocols. Chemiluminescent detection was performed using SuperSignal West Pico PLUS chemiluminescent substrate (34580, Thermo Fisher Scientific) and C-DiGit blot scanner (3600-00, LI-COR Biosciences) in a high-sensitivity mode (12min scanning). Densitometry was performed using the ImageJ software (National Institutes of Health) using the standard algorithm (consecutive selection and plotting of the lanes with the measurement of the peak area) and subsequent adjustment to the loading control (GAPDH).

Statistics

Data were analysed using GraphPad Prizm (version 8.4.3). CD63-beads assay, NanoView, adhesion assay, FA quantification were analysed by one-way ANOVA test. Two group CD63-bead assay was analysed by non-paired T-test. 2D migration multiple comparison data and VSMC invasion multiple comparison data were analysed using non-parametric Kruskal-Wallis test. Two group invasion data were analysed using non-parametric Kolmogorov-Smirnov test. Clinical data for two groups was analysed by non-paired T-test and multiple comparison was analysed by Mann-Whitney U-test. All analysis was conducted using PRISM software (GraphPad, San Diego, CA). Values of $P<0.05$ were considered statistically significant.

Short abbreviations

ECM   extracellular matrix

EM   electron microscopy

FN   fibronectin

MVB   multivesicular body

NTA   nanoparticle tracking analysis

SMPD3 sphingomyelin phosphodiesterase 3

VSMC   vascular smooth muscle cell

Acknowledgments

ANK and CS were supported by BHF-PG/17/37/33023. TI was supported by BHF PG/20/6/34835 and
FS/14/30/30917. REC was supported by National Institutes of General Medical Sciences grant
R01GM134531. AK, MS and LB were supported by the Ministry of Science and Higher Education of
the Russian Federation-Complex Program of Basic Research under the Siberian Branch of the Russian
Academy of Sciences within the Basic Research Topic of Research Institute for Complex Issues
of Cardiovascular Diseases № 0419-2021-001. We are thankful to Dr Adam A Walters for the flow
cytometry consultancy and help.

Competing interests

A. Kapustin is currently an employee and shareholder of AstraZeneca.

References

1. Libby, P., Ridker, P.M. & Hansson, G.K. Progress and challenges in translating the biology of
atherosclerosis. *Nature* **473**, 317-325 (2011).
2. Chappell, J. *et al.* Extensive Proliferation of a Subset of Differentiated, yet Plastic, Medial
Vascular Smooth Muscle Cells Contributes to Neointimal Formation in Mouse Injury and
Atherosclerosis Models. *Circ Res* **119**, 1313-1323 (2016).
3. Misra, A. *et al.* Integrin beta3 regulates clonality and fate of smooth muscle-derived
atherosclerotic plaque cells. *Nat Commun* **9**, 2073 (2018).
4. Durham, A.L., Speer, M.Y., Scatena, M., Giachelli, C.M. & Shanahan, C.M. Role of smooth
muscle cells in vascular calcification: implications in atherosclerosis and arterial stiffness.
Cardiovasc Res **114**, 590-600 (2018).
5. Rohwedder, I. *et al.* Plasma fibronectin deficiency impedes atherosclerosis progression and
fibrous cap formation. *EMBO Mol Med* **4**, 564-576 (2012).
6. Langley, S.R. *et al.* Extracellular matrix proteomics identifies molecular signature of
symptomatic carotid plaques. *J Clin Invest* **127**, 1546-1560 (2017).
7. Glukhova, M.A. *et al.* Expression of extra domain A fibronectin sequence in vascular smooth
muscle cells is phenotype dependent. *J Cell Biol* **109**, 357-366 (1989).
8. Pankov, R. & Yamada, K.M. Fibronectin at a glance. *J Cell Sci* **115**, 3861-3863 (2002).
9. Hocking, D.C., Sottile, J. & McKeown-Longo, P.J. Activation of distinct alpha5beta1-mediated
signaling pathways by fibronectin's cell adhesion and matrix assembly domains. *J Cell Biol*
141, 241-253 (1998).
10. Bass, M.D. *et al.* Syndecan-4-dependent Rac1 regulation determines directional migration in
response to the extracellular matrix. *J Cell Biol* **177**, 527-538 (2007).
11. Nobes, C.D. & Hall, A. Rho, rac, and cdc42 GTPases regulate the assembly of multimolecular
focal complexes associated with actin stress fibers, lamellipodia, and filopodia. *Cell* **81**, 53-62
(1995).
12. Sheetz, M.P., Felsenfeld, D.P. & Galbraith, C.G. Cell migration: regulation of force on
extracellular-matrix-integrin complexes. *Trends Cell Biol* **8**, 51-54 (1998).
13. Ridley, A.J. Rho GTPases and cell migration. *J Cell Sci* **114**, 2713-2722 (2001).
14. Sung, B.H., Ketova, T., Hoshino, D., Zijlstra, A. & Weaver, A.M. Directional cell movement
through tissues is controlled by exosome secretion. *Nat Commun* **6**, 7164 (2015).
15. Sung, B.H. & Weaver, A.M. Exosome secretion promotes chemotaxis of cancer cells. *Cell Adh*
Migr **11**, 187-195 (2017).
16. Sung, B.H. *et al.* A live cell reporter of exosome secretion and uptake reveals pathfinding
behavior of migrating cells. *Nat Commun* **11**, 2092 (2020).
17. Koumangoye, R.B., Sakwe, A.M., Goodwin, J.S., Patel, T. & Ochieng, J. Detachment of breast
tumor cells induces rapid secretion of exosomes which subsequently mediate cellular
adhesion and spreading. *PLoS One* **6**, e24234 (2011).
18. Hoshino, D. *et al.* Exosome secretion is enhanced by invadopodia and drives invasive
behavior. *Cell Rep* **5**, 1159-1168 (2013).
19. Clayton, A. *et al.* Adhesion and signaling by B cell-derived exosomes: the role of integrins.
FASEB J **18**, 977-979 (2004).
20. Brown, M. *et al.* Lymphatic exosomes promote dendritic cell migration along guidance cues.
J Cell Biol **217**, 2205-2221 (2018).
21. Majumdar, R., Tavakoli Tameh, A. & Parent, C.A. Exosomes Mediate LTB4 Release during
Neutrophil Chemotaxis. *PLoS Biol* **14**, e1002336 (2016).
22. Kriebel, P.W. *et al.* Extracellular vesicles direct migration by synthesizing and releasing
chemotactic signals. *J Cell Biol* **217**, 2891-2910 (2018).
23. Kapustin, A.N. *et al.* Vascular smooth muscle cell calcification is mediated by regulated
exosome secretion. *Circ Res* **116**, 1312-1323 (2015).

24. Beacham, D.A., Amatangelo, M.D. & Cukierman, E. Preparation of extracellular matrices produced by cultured and primary fibroblasts. *Curr Protoc Cell Biol* **Chapter 10**, Unit 10 19 (2007).

25. Kowal, J. *et al.* Proteomic comparison defines novel markers to characterize heterogeneous populations of extracellular vesicle subtypes. *Proc Natl Acad Sci U S A* **113**, E968-977 (2016).

26. Trajkovic, K. *et al.* Ceramide triggers budding of exosome vesicles into multivesicular endosomes. *Science* **319**, 1244-1247 (2008).

27. Klinghoffer, R.A., Sachsenmaier, C., Cooper, J.A. & Soriano, P. Src family kinases are required for integrin but not PDGFR signal transduction. *EMBO J* **18**, 2459-2471 (1999).

28. Sinha, S. *et al.* Cortactin promotes exosome secretion by controlling branched actin dynamics. *J Cell Biol* **214**, 197-213 (2016).

29. Ruiz-Navarro, J. *et al.* Formin-like 1beta phosphorylation at S1086 is necessary for secretory polarized traffic of exosomes at the immune synapse in Jurkat T lymphocytes. *Elife* **13** (2024).

30. Campellone, K.G., Lebek, N.M. & King, V.L. Branching out in different directions: Emerging cellular functions for the Arp2/3 complex and WASP-family actin nucleation factors. *Eur J Cell Biol* **102**, 151301 (2023).

31. Nolen, B.J. *et al.* Characterization of two classes of small molecule inhibitors of Arp2/3 complex. *Nature* **460**, 1031-1034 (2009).

32. Rizvi, S.A. *et al.* Identification and characterization of a small molecule inhibitor of formin-mediated actin assembly. *Chem Biol* **16**, 1158-1168 (2009).

33. Taunton, J. *et al.* Actin-dependent propulsion of endosomes and lysosomes by recruitment of N-WASP. *J Cell Biol* **148**, 519-530 (2000).

34. Abella, J.V. *et al.* Isoform diversity in the Arp2/3 complex determines actin filament dynamics. *Nat Cell Biol* **18**, 76-86 (2016).

35. Thomas, S., Popov, V.L. & Walker, D.H. Exit mechanisms of the intracellular bacterium Ehrlichia. *PLoS One* **5**, e15775 (2010).

36. Mellor, H. The role of formins in filopodia formation. *Biochim Biophys Acta* **1803**, 191-200 (2010).

37. Bohil, A.B., Robertson, B.W. & Cheney, R.E. Myosin-X is a molecular motor that functions in filopodia formation. *Proc Natl Acad Sci U S A* **103**, 12411-12416 (2006).

38. Heusermann, W. *et al.* Exosomes surf on filopodia to enter cells at endocytic hot spots, traffic within endosomes, and are targeted to the ER. *J Cell Biol* **213**, 173-184 (2016).

39. Verweij, F.J. *et al.* Quantifying exosome secretion from single cells reveals a modulatory role for GPCR signaling. *J Cell Biol* **217**, 1129-1142 (2018).

40. Hutcheson, J.D. *et al.* Genesis and growth of extracellular-vesicle-derived microcalcification in atherosclerotic plaques. *Nat Mater* **15**, 335-343 (2016).

41. Whitehead, M. *et al.* Vascular smooth muscle cell senescence accelerates medin aggregation via small extracellular vesicle secretion and extracellular matrix reorganization. *Aging Cell* **22**, e13746 (2023).

42. Didangelos, A. *et al.* Proteomics characterization of extracellular space components in the human aorta. *Mol Cell Proteomics* **9**, 2048-2062 (2010).

43. Kapustin, A.N. *et al.* Prothrombin Loading of Vascular Smooth Muscle Cell-Derived Exosomes Regulates Coagulation and Calcification. *Arterioscler Thromb Vasc Biol* **37**, e22-e32 (2017).

44. Tagliatela, A.C. *et al.* Coronin 1C inhibits melanoma metastasis through regulation of MT1-MMP-containing extracellular vesicle secretion. *Sci Rep* **10**, 11958 (2020).

45. Rohlena, J. *et al.* Endothelial CD81 is a marker of early human atherosclerotic plaques and facilitates monocyte adhesion. *Cardiovasc Res* **81**, 187-196 (2009).

46. Burridge, K. Foot in mouth: do focal adhesions disassemble by endocytosis? *Nat Cell Biol* **7**, 545-547 (2005).

47. Holt, M.R. *et al.* Quantifying cell-matrix adhesion dynamics in living cells using interference reflection microscopy. *J Microsc* **232**, 73-81 (2008).

48. Chrzanowska-Wodnicka, M. & Burridge, K. Rho-stimulated contractility drives the formation of stress fibers and focal adhesions. *J Cell Biol* **133**, 1403-1415 (1996).

49. Arthur, W.T. & Burridge, K. RhoA inactivation by p190RhoGAP regulates cell spreading and migration by promoting membrane protrusion and polarity. *Mol Biol Cell* **12**, 2711-2720 (2001).

50. Tabdanov, E. *et al.* Micropatterning of TCR and LFA-1 ligands reveals complementary effects on cytoskeleton mechanics in T cells. *Integr Biol (Camb)* **7**, 1272-1284 (2015).

51. Schoen, I., Hu, W., Klotzsch, E. & Vogel, V. Probing cellular traction forces by micropillar arrays: contribution of substrate warping to pillar deflection. *Nano Lett* **10**, 1823-1830 (2010).

52. Ghibaudo, M. *et al.* Traction forces and rigidity sensing regulate cell functions *Soft Matter* **4**, 1836-1843 (2008).

53. Tillet, E., Gential, B., Garrone, R. & Stallcup, W.B. NG2 proteoglycan mediates beta1 integrin-independent cell adhesion and spreading on collagen VI. *J Cell Biochem* **86**, 726-736 (2002).

54. Nishida, S. *et al.* Collagen VI suppresses fibronectin-induced enteric neural crest cell migration by downregulation of focal adhesion proteins. *Biochem Biophys Res Commun* **495**, 1461-1467 (2018).

55. Sardone, F. *et al.* Collagen VI-NG2 axis in human tendon fibroblasts under conditions mimicking injury response. *Matrix Biol* **55**, 90-105 (2016).

56. Pankova, K., Rosel, D., Novotny, M. & Brabek, J. The molecular mechanisms of transition between mesenchymal and amoeboid invasiveness in tumor cells. *Cell Mol Life Sci* **67**, 63-71 (2010).

57. Galbraith, C.G., Yamada, K.M. & Galbraith, J.A. Polymerizing actin fibers position integrins primed to probe for adhesion sites. *Science* **315**, 992-995 (2007).

58. Pollard, T.D. & Borisy, G.G. Cellular motility driven by assembly and disassembly of actin filaments. *Cell* **112**, 453-465 (2003).

59. Ridley, A.J. & Hall, A. The small GTP-binding protein rho regulates the assembly of focal adhesions and actin stress fibers in response to growth factors. *Cell* **70**, 389-399 (1992).

60. Price, L.S., Leng, J., Schwartz, M.A. & Bokoch, G.M. Activation of Rac and Cdc42 by integrins mediates cell spreading. *Mol Biol Cell* **9**, 1863-1871 (1998).

61. Ren, X.D., Kiosses, W.B. & Schwartz, M.A. Regulation of the small GTP-binding protein Rho by cell adhesion and the cytoskeleton. *EMBO J* **18**, 578-585 (1999).

62. Wang, H.R. *et al.* Regulation of cell polarity and protrusion formation by targeting RhoA for degradation. *Science* **302**, 1775-1779 (2003).

63. Rottner, K., Hall, A. & Small, J.V. Interplay between Rac and Rho in the control of substrate contact dynamics. *Curr Biol* **9**, 640-648 (1999).

64. Roberts, L.A., Glenn, H., Hahn, C.S. & Jacobson, B.S. Cdc42 and RhoA are differentially regulated during arachidonate-mediated HeLa cell adhesion. *J Cell Physiol* **196**, 196-205 (2003).

65. Nobes, C.D. & Hall, A. Rho GTPases control polarity, protrusion, and adhesion during cell movement. *J Cell Biol* **144**, 1235-1244 (1999).

66. Cox, E.A., Sastry, S.K. & Huttenlocher, A. Integrin-mediated adhesion regulates cell polarity and membrane protrusion through the Rho family of GTPases. *Mol Biol Cell* **12**, 265-277 (2001).

67. Matera, D.L., Lee, A.T., Hiraki, H.L. & Baker, B.M. The Role of Rho GTPases During Fibroblast Spreading, Migration, and Myofibroblast Differentiation in 3D Synthetic Fibrous Matrices. *Cell Mol Bioeng* **14**, 381-396 (2021).

68. Sahai, E. & Marshall, C.J. Differing modes of tumour cell invasion have distinct requirements for Rho/ROCK signalling and extracellular proteolysis. *Nat Cell Biol* **5**, 711-719 (2003).

69. Sanz-Moreno, V. *et al.* Rac activation and inactivation control plasticity of tumor cell movement. *Cell* **135**, 510-523 (2008).

70. Okada, M. *et al.* Primary collagen VI deficiency is the second most common congenital muscular dystrophy in Japan. *Neurology* **69**, 1035-1042 (2007).

71. Bryant, J.E. *et al.* Cardiac myofibroblast differentiation is attenuated by alpha(3) integrin blockade: potential role in post-MI remodeling. *J Mol Cell Cardiol* **46**, 186-192 (2009).

72. Luther, D.J. *et al.* Absence of type VI collagen paradoxically improves cardiac function, structure, and remodeling after myocardial infarction. *Circ Res* **110**, 851-856 (2012).

73. Katsuda, S. *et al.* Collagens in human atherosclerosis. Immunohistochemical analysis using collagen type-specific antibodies. *Arterioscler Thromb* **12**, 494-502 (1992).

74. Liu, X. *et al.* Collagen VI antibody reduces atherosclerosis by activating monocyte/macrophage polarization in ApoE(-/-) mice. *Int Immunopharmacol* **111**, 109100 (2022).

75. He, M. *et al.* Hepatocellular carcinoma-derived exosomes promote motility of immortalized hepatocyte through transfer of oncogenic proteins and RNAs. *Carcinogenesis* **36**, 1008-1018 (2015).

76. Doane, K.J., Howell, S.J. & Birk, D.E. Identification and functional characterization of two type VI collagen receptors, alpha 3 beta 1 integrin and NG2, during avian corneal stromal development. *Invest Ophthalmol Vis Sci* **39**, 263-275 (1998).

77. Nishiyama, A. & Stallcup, W.B. Expression of NG2 proteoglycan causes retention of type VI collagen on the cell surface. *Mol Biol Cell* **4**, 1097-1108 (1993).

78. Nanda, A. *et al.* TEM8 interacts with the cleaved C5 domain of collagen alpha 3(VI). *Cancer Res* **64**, 817-820 (2004).

79. Burgi, J. *et al.* CMG2/ANTXR2 regulates extracellular collagen VI which accumulates in hyaline fibromatosis syndrome. *Nat Commun* **8**, 15861 (2017).

80. Burgi, J. *et al.* Ligand Binding to the Collagen VI Receptor Triggers a Talin-to-RhoA Switch that Regulates Receptor Endocytosis. *Dev Cell* **53**, 418-430 e414 (2020).

81. Pankova, D. *et al.* NG2-mediated Rho activation promotes amoeboid invasiveness of cancer cells. *Eur J Cell Biol* **91**, 969-977 (2012).

82. She, Z.G. *et al.* NG2 Proteoglycan Ablation Reduces Foam Cell Formation and Atherogenesis via Decreased Low-Density Lipoprotein Retention by Synthetic Smooth Muscle Cells. *Arterioscler Thromb Vasc Biol* **36**, 49-59 (2016).

83. Holm Nielsen, S. *et al.* The novel collagen matrikine, endotrophin, is associated with mortality and cardiovascular events in patients with atherosclerosis. *J Intern Med* **290**, 179-189 (2021).

84. Frimodt-Moller, M. *et al.* A marker of type VI collagen formation (PRO-C6) is associated with higher arterial stiffness in type 1 diabetes. *Acta Diabetol* **56**, 711-712 (2019).

85. Lee, J.E. *et al.* Identification of EDIL3 on extracellular vesicles involved in breast cancer cell invasion. *J Proteomics* **131**, 17-28 (2016).

86. Costanza, B. *et al.* Transforming growth factor beta-induced, an extracellular matrix interacting protein, enhances glycolysis and promotes pancreatic cancer cell migration. *Int J Cancer* **145**, 1570-1584 (2019).

87. Stampolidis, P., Ullrich, A. & Iacobelli, S. LGALS3BP, lectin galactoside-binding soluble 3 binding protein, promotes oncogenic cellular events impeded by antibody intervention. *Oncogene* **34**, 39-52 (2015).

88. Savina, A., Furlan, M., Vidal, M. & Colombo, M.I. Exosome release is regulated by a calcium-dependent mechanism in K562 cells. *J Biol Chem* **278**, 20083-20090 (2003).

89. Jae, N., McEwan, D.G., Manavski, Y., Boon, R.A. & Dimmeler, S. Rab7a and Rab27b control secretion of endothelial microRNA through extracellular vesicles. *FEBS Lett* **589**, 3182-3188 (2015).

1   90.   Savina, A., Vidal, M. & Colombo, M.I. The exosome pathway in K562 cells is regulated by
2   Rab11. *J Cell Sci* **115**, 2505-2515 (2002).

3   91.   Baietti, M.F. *et al.* Syndecan-syntenin-ALIX regulates the biogenesis of exosomes. *Nat Cell*
4   *Biol* **14**, 677-685 (2012).

5   92.   Goettsch, C. *et al.* Sortilin mediates vascular calcification via its recruitment into extracellular
6   vesicles. *J Clin Invest* **126**, 1323-1336 (2016).

7   93.   van Niel, G. *et al.* The tetraspanin CD63 regulates ESCRT-independent and -dependent
8   endosomal sorting during melanogenesis. *Dev Cell* **21**, 708-721 (2011).

94.   Pickering, J.G. *et al.* alpha5beta1 integrin expression and luminal edge fibronectin matrix
assembly by smooth muscle cells after arterial injury. *Am J Pathol* **156**, 453-465 (2000).

95.   Slepian, M.J., Massia, S.P., Dehdashti, B., Fritz, A. & Whitesell, L. Beta3-integrins rather than
beta1-integrins dominate integrin-matrix interactions involved in postinjury smooth muscle
cell migration. *Circulation* **97**, 1818-1827 (1998).

96.   Libby, P. & Aikawa, M. Stabilization of atherosclerotic plaques: new mechanisms and clinical
targets. *Nat Med* **8**, 1257-1262 (2002).

97.   Parsons, M., Messent, A.J., Humphries, J.D., Deakin, N.O. & Humphries, M.J. Quantification
of integrin receptor agonism by fluorescence lifetime imaging. *J Cell Sci* **121**, 265-271 (2008).

98.   Bampton, E.T.W., Goemans, C.G., Niranjan, D., Mizushima, N. & Tolkovsky, A.M. The
Dynamics of Autophagy Visualised in Live Cells: from Autophagosome Formation to Fusion
with Endo/lysosomes. *Autophagy* **1**, 23-36 (2005).

99.   Leifer, C.A. *et al.* TLR9 is localized in the endoplasmic reticulum prior to stimulation. *J*
Immunol **173**, 1179-1183 (2004).

100.   Berg, J.S. & Cheney, R.E. Myosin-X is an unconventional myosin that undergoes
intrafilopodial motility. *Nat Cell Biol* **4**, 246-250 (2002).

101.   Reynolds, J.L. *et al.* Human vascular smooth muscle cells undergo vesicle-mediated
calcification in response to changes in extracellular calcium and phosphate concentrations: a
potential mechanism for accelerated vascular calcification in ESRD. *J Am Soc Nephrol* **15**,
2857-2867 (2004).

102.   Kapustin, A.N. *et al.* Calcium regulates key components of vascular smooth muscle cell-
derived matrix vesicles to enhance mineralization. *Circ Res* **109**, e1-12 (2011).

103.   Kapustin, A.N. *et al.* Antisense oligonucleotide activity in tumour cells is influenced by
intracellular LBPA distribution and extracellular vesicle recycling. *Communications Biology* **4**,
1241 (2021).

104.   Pandey, P. *et al.* Cardiomyocytes Sense Matrix Rigidity through a Combination of Muscle and
Non-muscle Myosin Contractions. *Dev Cell* **45**, 661 (2018).

Figure Abstract

Vascular smooth muscle cells sense fibronectin via β1 integrin and secrete small extracellular vesicles loaded with collagen VI. These extracellular vesicles are entrapped in the extracellular matrix and induce formation of peripheral focal adhesions presenting adhesion complex ECM proteins including collagen VI, LGALS3BP, EDIL3 and TGFBI. Focal adhesions anchor the extracellular matrix to actin fibrils in the cell. Contraction of the actin fibrils generates the mechanical force for directional cell invasion through the matrix. This figure was created with BioRender (https://biorender.com/).

Figure 1. FN matrix stimulates sEV secretion by VSMCs *in vitro*. **A,** Immobilised FN but not a soluble FN promotes sEV secretion. Cells were cultured for 24h and sEVs in conditioned media were measured by CD63-beads assay. N=3 biological replicates, n=4 technical replicates, ANOVA, ****p<0.0001 **B,** Micrograph showing VSMC plated onto plastic or 3D matrix. **C,** VSMCs were plated on the 3D matrix for 24h, fixed and labelled for Syndecan-4 (green), F-actin (phalloidin, red) and fibronectin (blue) Size bar, 10μm. **D**, 3D matrix promote sEV secretion. Cells were cultured for 24h and conditioned media was collected and sEV secretion was measured by CD63-beads assay. N=5, biological replicates, n=4 technical replicates, t-test **E,** FN matrix does not affect sEV mode size. VSMCs were plated on non-coated or FN-coated flasks and incubated for 24h. Isolated sEVs were analysed by Nanoparticle Tracking Analysis. Representative from N=3 biological replicates. **F,** sEV and EV markers distribution is not altered by FN matrix. Cells were plated on non-coated or FN-coated flasks and apoptotic bodies (AB, 1.2K pellet), extracellular vesicles (EV, 10K pellet) and small extracellular vesicles (sEVs, 100K pellet) were isolated by differential ultracentrifugation and analysed by western blotting. Representative image from N=3 biological replicates. **G,** FN induces secretion of sEVs by activating β1 integrin. VSMCs were plated on non-coated or FN-coated plates in the absence or presence of integrin activating (12G10) or inhibiting (4B4) antibodies for 24h and conditioned media was analysed by CD63-bead assay. N=3 biological replicates, n=4 technical replicates, ANOVA, ****p<0.0001 **H,** Src is required for the sEV secretion. Cells were plated and sEV secretion was measured as in 2A. N=3, biological replicates, n=4, technical replicates, ANOVA, **p<0.0001 **I,** Inhibition of FAK blocks FN-induced sEV secretion. Cells were plated and sEV secretion was measured as in 1A. N=3 biological replicates, n=4 technical replicates, ANOVA.

Figure 2. sEV secretion is regulated by Arp2/3 and formin-dependent actin cytoskeleton remodelling.
A, Arp2/3 inhibition with CK666 reduces sEV secretion in VSMC. Cells were plated onto non-coated or FN-coated plates for 24h and conditioned media was analyzed by CD63-bead assay. N=4 biological replicates, n=4 technical replicates, ANOVA. **B,** Formin inhibitor, SMIFH2 blocking filopodia formation reduces sEV secretion from FN-plated cells only. Cells were plated onto non-coated or FN-coated plates for 24h and conditioned media was analyzed by CD63-bead assay. N=4-5 biological replicates, n=2-4 technical replicates, ANOVA. **C,** Still images of a time-lapse showing that transported MVBs are attached to F-actin tails. VSMCs were co-transfected with CD63-GFP and F-tractin-RFP and time-lapse was captured using confocal spinning-disk microscopy. Arrow head – position of CD63 MVB across time. Time, min. Size bar, 10μm. **D,** VSMC were plated for 24h in the absence (top, -ve) or presence of sEV secretion inhibitor (3-OMS, 10μM). Cells were fixed and stained for cortactin, F-actin and CD63. Note an overlap between CD63 endosomes and branched actin proteins (F-actin and cortactin), which is enhanced by 3-OMS treatment (arrow). Size bar, 10μm. **E,** CD63 MVBs (arrows) are detected in filopodia-like structures. VSMCs were plated on FN-coated plates for 24h and cells were stained for Myo10 (green), CD63 (blue), F-actin (phalloidin, red). Size bar, 10μm. **C,** Still images of a time-lapse showing that MVBs are transported to the filopodia tip in the live VSMC. Cells were co-transfected by CD63-RFP and Myo10-GFP, cultured for 24h and time-lapse video was captured using confocal spinning disk microscopy. Snapshots were taken at T=3s and T=4s after start of the video.

Figure 3. Endogenous and exogenous sEVs are trapped by ECM *in vitro*. **A,** VSMC were plated onto non-coated (top) or FN-coated (bottom) plastic for 24h and stained for the membrane sEV marker, CD63 and F-actin. Note accumulation of CD63 puncta in close proximity to filopodia-like projections. Size bar, 10μm. **B,** VSMCs were plated onto gelatin-coated plates and treated with control or SMPD3 siRNA for 72h. 3D matrices were generated and stained for CD63 and FN. Images were acquired using Nikon AX inverted confocal microscope. Note the decrease in CD63-positive sEVs associated with the FN fibrils. **C,** FN is presented on the surface of the VSMC-derived sEV along with α5β1 integrin. VSMC sEVs were immobilised on the 4μm beads. sEV-beads and VSMCs were stained with the antibodies (filled graphs) in non-permeabilised conditions and analysed by flow cytometry. **D**, VSMC were plated on FN-coated dishes and Alexa568-labelled sEV were added to the cell media for 3h. Cells were fixed and stained for filopodia marker Myo10 (green) and vinculin (blue). Note perinuclear localisation of internalised sEVs. Size bar, 10μm. Representative image from N=3 biological replicates. **E**, VSMC were plated on FN-coated dishes pre-coated with Alexa568-labelled sEV and incubated for 24h. Cell staining as in Fig 3D. Note even distribution of sEVs across the matrix and cell area. **F**, VSMC were plated on the FN-coated dishes in the absence of Alexa568-labelled sEV and incubated for 24h. Cell staining as in Fig 3D. Note the absence of signal in sEV channel.

Figure 4. Fibronectin deposition in the atherosclerotic plaque is spatially associated with sEV markers.

A, Proteomic profiling of pairwise collected carotid atherosclerotic plaques and adjacent intact arterial segments. Venn diagram shows the number of plaque-specific (n = 213) and intact-specific (n = 111) proteins as well as the number of proteins which are common for both vascular regions (1,368). **B**. Partial least-squares discriminant analysis indicates clear classification pattern between the carotid plaques (indicated by red triangles) and adjacent intact arterial segments (indicated by blue circles). N = 14 patients. **C**. Volcano plot illustrates that 46 proteins are significantly overexpressed in plaques whilst 13 proteins are significantly upregulated in adjacent intact arterial segments. **D**, **E.** Atherosclerotic plaques were co-stained for fibronectin (FN) and sEV marker, CD81. Cell nuclei were counterstained with DAPI. Main figure: x200 magnification, size bar, 50 μm. Box: x400 magnification, size bar, 15μm. Note an accumulation of FN in the neointima. **F,** Spatial distribution of FN and CD81 in the neointima. Note high overlap between FN and CD81 in the extracellular matrix. x200 magnification, size bar, 50 μm. Manders' split colocalization coefficient for the overlap of FN with CD81 (M1) and CD81 with FN (M2). Neointima region as in Fig. 4E. **G,** Co-localisation of endothelial cells (stained for CD31/PECAM1, red colour) and sEVs (stained for CD81, green colour). Nuclei are counterstained with 4',6-diamidino-2-phenylindole (DAPI, blue colour). Magnification: ×200, scale bar: 50 μm. Note endothelialised (i.e., those covered with CD31-positive cells) segments at the luminal side and abundant CD81 expression both in the neointima and in the tunica media. Endothelial cells (CD31, red colour) are largely co-localised with CD81-positive extracellular vesicles (green colour). Note CD31-positive capillaries in the neointima. **H, I,** Quantification of FN content in atherosclerotic plaques. Samples were analysed by western blot and bands intensity was quantified in ImageJ. Fold change was calculated as ratio of band intensity in the atherosclerotic plaque to band intensity in the adjacent intact arterial segments normalised to GAPDH. Note that FN content is elevated in atherosclerotic plaques relative to the adjacent intact arterial segments. Paired t-test.

Figure 5. sEV induces directional VSMC invasion. **A, B,** VSMC migration in 2D assay. VSMC were plated onto FN in the absence or presence of SMPD3 inhibitor (3-OMS) and/or sEV (25ug). Cells were tracked for 8h. Kruskal-Wallis with Dunn's multiple comparison test, N=2-3 biological replicates, n indicates the total number of tracked cells per condition, **, p<0.01, ****, p<0.0001. **C,** Chemotaxis μ-slide diagram. Yellow, cell chamber, blue chemoattractant-free medium chamber, dark blue – chemoattractant medium chamber. **D-H,** sEV promote directional VSMC invasion. Cells were seeded to the FN-enriched Matrigel matrix in μ-Slide Chemotaxis assay and stained with Draq5. Cell tracking was conducted by OperaPhenix microscope for 12h and cell invasion parameters were quantified using Columbus. Kruskal-Wallis with Dunn's multiple comparison test, N=4, biological replicates, n indicates the total number of tracked cells per condition, **, p<0.01, ***, p<0.001, ****, p<0.0001.

Figure 6. sEVs induce formation of peripheral FAs. **A,** VSMC were plated on FN matrix for 30min and adhered cells were counted by crystal violet staining. N=6 biological replicates, n=4-10 technical replicates, ANOVA, ***p<0.001, ****, p<0.0001 **B, C,** VSMC spreading onto FN was tracked by using ACEA's xCELLigence Real-Time Cell Analysis. Note that FN matrix promoted VSMC adhesion but addition of sEVs inhibited cell spreading. N=3 biological replicates, n=2-3 technical replicates, ANOVA. **D,** VSMCs were spread onto FN for 30 min and cells were fixed and stained with CellMask (magenta) and vinculin (green). Size bar, 10μm. **E, F, G, I** Quantification of FA number, distance from plasma membrane, cell size and mean FA size per cell, respectively. FA were stained as in 5D and quantified. Representative data from N=3 biological replicates, n=80-82 total number of tracked cells per condition, ANOVA, *p<0.05, **p<0.01. **J,** Focal adhesion turnover is not affected by sEVs. VSMC were transfected with Paxillin-RFP and plated on the FN in the absence or presence of immobilised sEVs. Images were captured for 30min using confocal spinning disk microscopy and FA turnover was quantified using extracted images analysis, N=4, total n=53-61 cells per condition, Unpaired T-test. **K,** sEV induces formation of strong-pulling FAs. VSMC transfected with Paxillin-RFP were plated on the PDMS pillars which were covered with FN and sEVs and pillar displacements were quantified**. **p<0.01, Unpaired t-test, Data represent 2,170-2322 measurements per condition pooled from two independent biological replicates.

Figure 7. sEVs block focal adhesion formation by presenting collagen VI.

A, Proteomic analysis of VSMC-derived sEVs and EVs. Venn diagram. N=3 biological replicates. **B,** Protein enrichment in the EV and sEV proteome. Heat Map. N=3 biological replicates. **C,** Western blot validation of sEV cargos. EVs and sEVs were isolated from VSMC conditioned media by differential ultracentrifugation and analysed by western blotting. Representative image from N=3 biological replicates. **D,** Collagen VI was presented in sEVs from conditioned media and in ECM-associated sEVs. sEVs were isolated from conditioned media and 0.5M NaCl ECM fraction by differential ultracentrifugation and analysed by western blotting. **E,** Collagen VI is presented on the surface of sEVs. sEVs were analysed by dot-blot in the non-permeabilised or permeabilised (PBS-0.2%-tween20) conditions. Staining intensity was quantified for non-permeabilised (outside) and permeabilised (inside) conditions. Representative from N=2 biological replicates. **F,** VSMC adhesion is regulated by collagen VI loaded on sEV. FN matrices were incubated with sEV and anti-collagen VI antibody (COLVI IgG) or control IgG. Cell adhesion was tracked by using ACEA's xCELLigence Real-Time Cell Analysis. ANOVA, N=3 biological replicates, ***, p<0.001, ****, p<0.0001. **G, H, I,** sEVs promote directional VSMC invasion. VSMCs were treated with control siRNA (Scramble) or collagen VI-specific siRNA pools for 24h and were seeded to the FN-enriched Matrigel matrix in a μ-Slide Chemotaxis assay and stained with Draq5. Cell tracking was conducted by OperaPhenix microscope for 12h and cell invasion parameters were quantified using Columbus, n indicates the total number of tracked cells per condition, Kolmogorov-Smirnov test, *, p<0.05. **I,** Real-time PCR analysis of expression of CD9, CD63, CD81, COL6A3, EDIL3 and TGFBI in atherosclerotic plaque. *, p<0.05, Paired t-test, N=5 patients.

Supplementary Figures

Figure S1. FN matrix stimulates sEV secretion by VSMCs. **A,** FN and collagen I but not laminin stimulate secretion of CD63-enriched sEVs. VSMCs were plated on the various matrices for 24h and sEV secretion was measured by CD63-beads assay. N=3, biological replicates, n=6 technical replicates. ANOVA **p<0.01 **B,** Inhibition of SMPD3 blocks sEV secretion by VSMC plated onto FN matrix. VSMCs were plated on non-coated or FN-coated plates for 24h and conditioned media was analysed by CD63-bead assay. N=2 biological replicates, n=4 technical replicates, ANOVA, ***, p<0.001, ****p<0.0001

Figure S2. A, Still images of a time-lapse showing that Arp2/3 and F-actin form tails in VSMC cytosol. VSMCs were co-transfected by ARPC2–GFP and F-tractin-RFP and cultured for 24h. Time-lapse video was captured using confocal spinning disk microscopy. Note, that Arp2/3 and F-actin are observed in lamellipodia but also detected in the cytosol with the unknown activity (*arrow*). Size bar, 10µm **B**, Analysis of F-actin/CD63 overlap using ImageJ in a fixed VSMCs. F-actin and CD63 images (Fig 2D) were converted to 8 bits and thresholded. Actin stress fiber structures were extracted from the image by using ParticleAnalysis plug-in (Size (pixel^2) 10-infinity; Circularity 0.5-1) and overlap image created (merged). CD63 endosomes (pseudo coloured in green) were overlapped with F-actin spot-like structures (pseudo coloured in red). Note that endosomes partially overlapped with F-actin tail-like structures (arrows) though some F-actin spots and endosomes were not colocalised (arrowheads).
C, sEV secretion detected with CD63-pHluorin. VSMCs were co-transfected with CD63-pHluorin and incubated for 24h. Time-lapse was captured using confocal spinning-disk microscopy. Arrows, typical "burst"-like appearance of sEV secretion at the cell-ECM interface. Arrows, an intense CD63-pHluorin staining along filopodia-like structures indicating that sEV release can occur in filopodia. Size bar, 10µm

Figure S3 SMPD3-dependent sEVs are trapped in ECM. A, Characterisation of sEVs extracted from VSMC-derived media and ECMs fractions. sEVs were sequentially extracted and fractions were collected and analysed by using ExoView. X-axis – capturing spot antibodies (CD63, CD81, CD9 and control MigG1). **B,** CD63-bead capturing assay shows similar enrichment in CD63+/CD81+ sEVs in the conditioned media and ECM-associated sEVs extracted using 0.5M NaCl. VSMCs were plated onto gelatin-covered plates for 10 days and media was collected every 3d. sEVs were extracted in a mild (0.5M NaCl) conditions and media and 0.5M NaCl fractions were analysed by using CD63-beads capturing assay. **C, D,** Exogenously added FN-Alexa568 can be detected in the early endosomes and MVBs. VSMC were serum-deprived for 24h and then incubated with FN-Alexa568 for 30min (C) or 3h (D) in RPMI-0.5% BSA, thoroughly washed and fixed and stained for EEA-1 (C, green) or CD63 (D, blue). Size bar, 10um, **E,** Exogenously added FN-Alexa568 can be detected in MVBs. VSMC were incubated with FN-Alexa568 for 3h and stained for CD63 which was visualised with anti-mouse-Alexa488 antibody. Size bar, 10um, Note the partial colocalization of FN and CD63. **F, G,** FN can be detected in the CD63+ MVBs in a sparsely growing VSMCs. VSMC were plated at low (5,000 cells per well, **F**) or high density (20,000 cells per well, **G**) and were cultured for 24h in RMPI supplemented with 2.5% EV-free FBS, fixed and stained for F-actin, FN and CD63. **H,** VSMC plating density influences secretion of CD63+/CD81+ sEVs. Cells were plated at the different density and cultured for 24h and sEVs in conditioned media were measured by CD63-beads assay. N=1 biological replicates, n=8 technical replicates. **I,** FN is presented in sEV along with β1 integrin. **J,** VSMC were plated on the FN-coated dishes and Alexa568-labelled sEV were added to the cell media for 3h. Cells were fixed and stained for filopodia marker Myo10 (green) and vinculin (blue). Note perinuclear localisation of internalised sEVs. Size bar, 10μm. 3D projection. Myo10 staining channel is removed. **K,** VSMC were plated on the FN-coated dishes pre-coated with Alexa568-labelled sEV and incubated for 24h. Cell staining as in Fig S3J. Note even distribution of sEVs across the extracellular matrix and cell area.

Figure S4. Fibronectin deposition in the atherosclerotic plaque is spatially associated with sEV marker CD81.
A. Staining of atherosclerotic plaques with haematoxylin and eosin. Unstable atheroma with an extracellular lipid or necrotic core and calcification (type IV according to the American Heart Association). Below are representative fields of view focusing on the lumen (blue contour), neointima (green contour), and tunica media (yellow contour). Red contours demarcate representative fields of view within these tissue compartments. Note multiple cells at the luminal side, numerous leukocytes in the neointima, and abundant cells in the tunica media. Overview images: magnification: ×1.8, scale bar: 500 µm; close-ups demarcated by blue, green, and gold contour: magnification: ×200, scale bar, 150 µm; close-ups demarcated by red contour: magnification: ×400, scale bar: 50 µm. **B.** Staining of atherosclerotic plaques with Masson's trichrome. Note the superficial erosions at the luminal side, abundant haemorrhages in the neointima, and sporadic haemorrhages in the tunica media. Overview images: magnification: ×1.8, scale bar: 500 µm; close-ups demarcated by blue, green, and gold contour: magnification: ×200, scale bar: 150 µm; close-ups demarcated by red contour: magnification: ×400, scale bar, 50 µm. **C.** Staining of atherosclerotic plaques with orcein. Note the moderate connective tissue staining at the luminal side, weak connective tissue staining in the atheroma with a confluent extracellular lipid core, and strong specific staining at the tunica media containing elastic fibres. Overview images: magnification: ×1.8, scale bar: 500 µm; close-ups demarcated by blue, green, and gold contour: magnification: ×200, scale bar: 150 µm; close-ups demarcated by red contour: magnification: ×400, scale bar, 50 µm. **D.** Demarcation of tunica media from the neointima using specific antibodies to elastin (red colour) and non-specific autofluorescence of elastic fibers (green colour). Nuclei are counterstained with 4',6-diamidino-2-phenylindole (DAPI, blue colour). Two representative images per each vascular compartment (luminal side, neointima, and tunica media) are provided. Negative controls (i.e., sections stained with species-specific pre-adsorbed fluorescent-labeled secondary antibodies and DAPI but without the respective antigen-specific primary antibody) are provided to the right from representative images. Magnification: ×200, scale bar, 50 µm. Note that elastin is expressed exclusively in the tunica media and is associated with elastic fibers. **E.** Expression of FN in atherosclerotic plaques. Atherosclerotic plaques (A) and adjacent intact arterial segments (I) were homogenised and analysed by Western blotting. FN is abundantly presented in both regions and 50 kDa FN fragment (Homandberg et al., 1992) intensity was used to quantify FN amount.

Supplementary Figure S5 sEVs regulate VSMC motility and invasion. A, Centripetal FAs are linked to actin stress fibers. VSMCs were plated on FN-coated plate for 24h and cells were stained for Myo10 (green), CD63 (blue) and F-actin (phalloidin, red). Note the dot-like focal complexes in lamellipodium which are not associated with the contractile actin bundles (arrow) and an appearance of elongated FAs associated with the mature actin bundles (arrowhead). Dotted line, approximate position of the lamellipodium boundaries. Size bar, 10μm. **B,** Mature focal adhesion turnover is not affected by sEVs. VSMC were transfected with Paxillin-RFP and plated on the FN in the absence or presence of immobilised sEVs. Images were captured for 30min using confocal spinning disk microscopy. Note the appearance of the mature FAs in the lamellipodium (Arrowhead). Arrow, direction of the VSMC movement. Representative from N=4 biological replicates. **C,** sEV induces formation of FAs with the enhanced pulling force. VSMC transfected with Paxillin-RFP were plated on the PDMS pillars which were covered with FN and sEVs and pillar displacements were quantified**.** Representative image from N=2 biological replicates.

Supplementary Figure S6. Proteomic analysis of VSMC-derived sEVs and EV. VSMC-derived EVs and sEVs were isolated from cells and analyzed by protein mass-spectrometry. N=3 biological replicates. **A.** Clustered proteomic heatmap for EV and sEV. **B,** The GO term analysis was performed using the differentially expressed genes between the sEV and EV groups.