1 Reward contingency modulates olfactory bulb output via pathway-dependent peri-somatic
2 inhibition
3

4 Sander Lindeman, Xiaochen Fu, Janine K Reinert, Izumi Fukunaga
5
6 Sensory and Behavioural Neuroscience Unit, Okinawa Institute of Science and Technology
7 Graduate University, 1919-1 Tancha, 904-0412, Okinawa, Japan
8
Correspondence: Izumi.fukunaga@oist.jp

Abstract

Associating values to environmental cues is a critical aspect of learning from experiences,
allowing animals to predict and maximise future rewards. Value-related signals in the brain were
once considered a property of higher sensory regions, but its wide distribution across many
brain regions is increasingly recognised. Here, we investigate the nature of reward-related
signals at the earliest stage of olfactory processing, namely, in the olfactory bulb. In head-fixed
mice performing Go/No-Go discrimination of closely related olfactory mixtures, rewarded
odours evoke widespread inhibition in one class of output neurons, that is, in mitral cells but
not tufted cells. The temporal characteristics of this reward-related inhibition suggest it is
odour-driven, but it is also context-dependent since it is absent during pseudo-conditioning and
pharmacological silencing of the piriform cortex. Further, the reward-related modulation is
present in the somata but not in the apical dendritic tuft of mitral cells, suggesting an
involvement of circuit component located deep in the olfactory bulb. Depth-resolved imaging
from granule cell dendritic gemmules suggests that granule cells that target mitral cells receive
a reward-related extrinsic drive. Our results support the notion that value-related modulation
appears at the early stages of sensory processing and provide constraints on long-range and
local circuit mechanisms.

Introduction

Sensory systems of the brain play a crucial role in guiding animals' choices. One of their uses is
in the reward-driven decision-making, where the system is thought to adjust the
representations of sensory cues depending on the past reward encounters, to influence their
future behavioural choices and learning. Decades of studies across brain areas have
demonstrated that reward expectations are, in turn, potent modulators of sensory activity. For
example, stimulus evoked responses in many sensory regions of the brain scale with the

quantity of expected reward [1-5]. This modulation is often interpreted as representing subjective
value [6,7]. Such a system where sensory processing is fined-tuned flexibly may be crucial for
maximising returns in a dynamic and uncertain world [7].
42
Decision and value-related modulations of sensory responses are featured prominently in
higher sensory areas [1,8]. However, recent studies indicate that even early stages of sensory
processing, especially in rodents, participate in value-like representations, showing ample
modulations associated with decision-making [9-11]. The olfactory system is an extreme case in
this regard, where apparent reward-related modulation is readily observed as peripherally as in
the olfactory bulb [12,13], the primary olfactory region situated just one synapse away from the
site of sensory transduction. This peripheral location, along with the saliency of the olfactory
cues for rodents, makes the olfactory bulb an attractive structure to study the mechanisms that
generate value-like signals in the brain [12].
52
The nature of this apparent reward-related modulation in the olfactory bulb remains
unresolved. For example, one study observed that evoked responses to rewarded vs.
unrewarded odours in the principal neurons of the olfactory bulb diverge only transiently as rats
learn to discriminate between the cues [12]. This may reflect the level of animal's engagement [14],
where the learning-related modulation corresponds mainly to the changes in the inputs from
the sensory periphery arising from sniff pattern changes. Rodents indeed adjust the odour
sampling patterns exquisitely according to the behavioural contexts [15,16]. However, given that
the olfactory bulb is a major target of feedback and neuromodulatory projections from many
brain regions, value-related information could affect how the olfactory bulb represents odours.
For example, electrical and optogenetic stimulations and pharmacological manipulations of
neuromodulatory and feedback inputs to the olfactory bulb change the gain of odour responses
in the principal neurons of this region [17-22]. Therefore, whether the reward-related signals in the
primary olfactory area simply reflect changes in the sensory input or internally generated
contextual signals need to be clarified.
67
Here, we show that the olfactory bulb exhibits robust and consistent reward-related signals
during a trace olfactory conditioning paradigm, where mice discriminate between closely
related olfactory mixtures. This phenomenon is characterised by widespread inhibitory
responses following the rewarded odour presentation, in mitral cells but not tufted cells. This
divergence is not explained by the odour identity or sampling strategy and reflects the
congruence of sensory drive and contextual signals. By imaging from specific subcellular
compartments of mitral cells, we demonstrate that the divergent responses first become
evident peri-somatically. Depth-resolved imaging from the dendrites of adult-born granule cells

suggests that the cell-type specific modulation may involve an extrinsic drive to putative mitral
cell-targeting granule cells.
78
79 **Results**
80
The olfactory bulb integrates both feedforward sensory stimuli, as well as long-range
projections from other brain areas (**Fig. 1A**). The latter input is thought to convey behavioural
contextual signals to the olfactory bulb and tune activity patterns flexibly. To study how the
behavioural context modulates the olfactory bulb output in olfactory decision making, we
trained head-fixed mice to perform an olfactory discrimination task (**Fig. 1**). Water restricted
mice were trained to associate a rewarded odour (S+ odour) with a water reward, and an
unrewarded odour (S- odour) with no water delivery (**Fig. 1B**). Note that this paradigm includes
a trace period, as we reasoned that an early cessation in the feedforward signal may maximise
the chance of observing context-related activity patterns. The mice were first trained to
discriminate between easily distinguishable odour mixtures, which comprised ethyl butyrate
(EB) and methyl butyrate (MB), mixed at 80%/20% ratio versus a 20%/80% ratio for the S+ vs. S-
stimuli, respectively (**Fig. 1C**). When the mice reached a criterion of 80% accuracy (3 ± 0.9 days,
n = 6 mice, Fig. 1D), they were trained to discriminate between more similar odour mixtures
("Difficult discrimination task"; 60%/40% mixture of ethyl butyrate and methyl butyrate versus a
40%/60% mixture). This is a task known to engage many components of the olfactory bulb
circuitry [23]. Well-trained mice discriminated between these similar mixtures in 1.63 ± 0.53 s
(**Supplementary fig. 1**), with comparable sniffing patterns for the S+ vs. S- odours
(**Supplementary Fig. 2**), consistent with previous reports where similar odours and reward
timing were used [24,25].
100
In the mice proficiently performing the difficult olfactory discrimination task, we studied the
responses of olfactory bulb output to the rewarded vs. unrewarded odours. The calcium
indicator GCaMP6f was expressed in mitral and tufted cells using *Tbx21-Cre* mice crossed with
Ai95D mice [26,27], and was imaged using a two-photon microscope (n=428 ROIs in 6 mice, and
n=150 ROIs in 3 mice, respectively; **Fig. 1E-G**). Mitral and tufted cells were distinguished by
depth (**Fig. 1E**). Tufted cells responded largely similarly to both odours (mean ΔF/F during odour
= 0.628 ± 0.135 and 0.655 ± 0.148 for S+ and S- respectively; p = 0.777, Wilcoxon rank-sum test;
mean ΔF/F post odour = 0.203 ± 0.264 and 0.237 ± 0.264 for S+ and S- respectively; p = 0.149,
Wilcoxon rank-sum test; **Fig. 1F**). Peculiarly, responses of the mitral cell somata to the rewarded
odour were characterised by widespread inhibitory responses (Mean ΔF/F S+ = -0.048 ± 0.058;
S- = -0.022 ± 0.054; p < 0.001, Wilcoxon rank-sum test; **Fig.1G**) . This dominance of inhibition for
the S+ odour was present soon after the odour onset, but was particularly pronounced during

the post-odour period (Mean ΔF/F S+ = -0.048 ± 0.095; S- = 0.034 ± 0.102; p < 0.001, Wilcoxon
rank-sum test; **Fig. 1G**).

The late onset of the reward-associated inhibition in mitral cells raises the question regarding
the underlying drive: Is the inhibitory component locked to the anticipatory motor output, or to
the odour? To analyse this, we divided the rewarded trials into two sets based on the animals'
reaction times ("early onset" vs. "late onset"), and reverse-correlated the GCaMP6f signals to
the onsets of anticipatory signals ("lick-aligned average"; **Fig. 2**). If the peak of inhibition in the
averages occur at the same time for the early lick sets and late lick sets, it would imply the
inhibition is locked more to the behavioural output (**Fig. 2B**). This analysis revealed, in contrast,
that the time of peak inhibition is shifted depending on the reaction time (Pearson's correlation
coefficient = -0.555, p = 0.026; N = 15 fields of view, 6 mice; **Fig. 2C-E**), indicating that the
inhibition is locked to the odour.

The prevalence of inhibitory responses in mitral cells following the rewarded odour
presentation is striking, but this level of inhibitory dominance has not been reported previously,
even though several studies already studied how mitral cells respond to odours during difficult
odour discrimination paradigms [28-30]. The difference here may be the short duration of odour
pulse used, followed by a two seconds-long trace period. It is possible that, with a longer odour
presentation, the feed-forward component may dominate over any modulatory influences in
the olfactory bulb (**Fig. 3A**). To test this possibility, in well-trained mice, we presented the
odours for a longer period (4 s), making the task a delay task (**Fig. 3B**). In this condition, mitral
cells responded to the rewarded and unrewarded odours similarly (**Fig. 3C-E**). Notably, both
responses were characterised by widespread inhibitory component (% of ROIs showing
significant inhibition = 18.1 for S+ and 11.9 for S-, and 35.2 for S+ and 23.9 for S- in early and
late time windows, respectively, n = 5 mice). The divergent responses may therefore originate
from a conjunction of olfactory and contextual signals.

To test if the behavioural state of the animal is crucial for the response divergence in mitral
cells, we used two pseudo-conditioning paradigms using the same odours (**Fig. 4A,B**). In the first
case ("Disengagement"), the water was delivered every trial, approximately 15 seconds before
the odour presentation (**Fig. 4B**). In the second case ("Random association"), we delivered the
water on randomly selected trials, so that both 60/40 and 40/60 odour mixtures were followed
by water 50% of the time (**Fig. 4B**). These two paradigms decouple the odour-reward
association, while inducing different levels of engagement in the head-fixed mice [31]. Imaging
sessions took place after the mice, previously trained on the fine discrimination task, were
switched to, and experienced at least one session of the new paradigm (**Fig. 4C**). In both control
paradigms, the head-fixed mice showed no preferential licking for the 60/40 mixture (average

anticipatory licks for disengagement paradigm = 1.5 ± 1.6 and 0.7 ± 0.8 on 60/40 and

40/60,respectively; p = 0.999, 1-way ANOVA with *post-hoc* multiple-comparisons; average

anticipatory licks for random association paradigm = 6.9 ± 6.1 and 5.5 ± 4.3 on 60/40 and 40/60,

respectively; p = 0.890, 1-way ANOVA with *post-hoc* multiple-comparisons, **Fig. 4D**).

Importantly, disengagement and randomized paradigms differed in the general levels of

anticipatory licks (average anticipatory licks for all trials = 1.1 ± 1.3 and 6.2 ± 5.2 for

disengagement and random association paradigms, respectively; p = 0.0021, 1-way ANOVA with

post-hoc multiple-comparisons), indicating that different levels of behavioural engagement were

indeed achieved by these paradigms. In both cases, the mitral cell somata responded similarly

to the two odour mixtures (mean ΔF/F for disengagement = -0.02 ± 0.05 and -0.01 ± 0.06

during odour for 60/40 and 40/60, respectively; p = 0.465; for post odour = 0.05 ± 0.09 and 0.05

± 0.09; p=0.553; mean ΔF/F for random association = -0.03 ± 0.09 and -0.04 ± 0.10 during

odour; p = 0.259; post odour = -0.01 ± 0.14 and 7.7×10^{-5} ± 0.14; p=0.617, Wilcoxon rank-sum

test; **Fig. 4E-G**). Note that the inhibition during the post-odour, anticipatory period that is

normally present in discriminating mice was generally reduced in the two control paradigms.

Together, these results indicate that the observed divergent responses in mitral cell somata are

state-dependent, and not explained by the odour identities.

What is the cellular origin of the widespread inhibition associated with the rewarded odour?

Previous studies showed that a variety of feedback and neuromodulatory projections to the

olfactory bulb modulate the physiology of olfactory bulb neurons [18-20,32-34]. Further, several

studies showed that such modulations manifest differently for mitral cells and tufted cells

[17,18,29,35]. Recent works indicate that mitral cells receive more potent feedback modulation from

the piriform cortex [17,18]. Thus, even though it is beyond the scope of the current work to

systematically investigate all sources, the piriform cortex is a reasonable candidate for the

source of the contextual signal resulting in the mitral cell-specific, reward-related inhibition we

observe.

To test the involvement of the piriform cortex, we pharmacologically inactivated the ipsilateral

anterior piriform cortex while the head-fixed mice performed the difficult olfactory

discrimination task (**Fig. 5A**). This was achieved by infusing the $GABA_A$ antagonist, muscimol,

through an implanted canula. Muscimol and control sessions were carried out on alternate

days, but the same fields of view were sampled for the two conditions, so that the responses of

the same ROIs could be compared directly. The infusion of muscimol disrupted the behavioural

performance significantly (behavioural accuracy = 64.0 ± 14.5 % during muscimol sessions; 92.8

± 7.8 % during control sessions; p = 0.004, Wilcoxon rank-sum test; n=6 control sessions and 6

muscimol sessions, 3 mice; **Fig. 5F**). When responses of mitral cells were imaged in this

condition, divergence in the rewarded vs. unrewarded odour responses were significantly

reduced (Mean ΔF/F during odour = -0.007 ± 0.073 and -0.023 ± 0.091 for S+ and S-
respectively; p = 0.174, Wilcoxon rank-sum test; Mean ΔF/F post-odour = 0.067 ± 0.124 and
0.079 ± 0.171 for S+ and S- respectively; p = 0.252, Wilcoxon rank-sum test, **Fig. 5G-I**). This was
characterised by a reduction in the inhibitory responses evoked by the rewarded stimulus
(normalised S+ − S- difference = -0.007 ± 0.156 and 0.043 ± 0.142 in control and muscimol
sessions respectively; p = 0.008, Wilcoxon rank-sum test; **Fig. 5J**), and during the post- odour
phase (normalised S+ − S- difference = -0.263 ± 0.175 and -0.040 ± 0.168 in control and
muscimol sessions respectively; p = 3.53 x 10^{-18}, Wilcoxon rank-sum test; **Fig. 5J**). Together,
these data indicate that an intact piriform cortex and/or accurate behavioural performance, is
required to observe the widespread inhibitory responses associated with the rewarded odour.
199
The results so far indicate that the widespread inhibitory responses associated with the
rewarded odours come from sources extrinsic to the olfactory bulb. One of the major targets of
such long-range projections within the olfactory bulb is the granule cells. These cells contact
mitral cells on their lateral dendrites at a deeper portion of the external plexiform layer. If the
granule cells convey the contextual signals to mitral cells, the divergent responses may be
observable perisomatically, but not in the superficial compartment (Fig. 6A). To test this, we
compared the GCaMP6f signals from the apical dendrites of mitral cells in the glomeruli, vs.
signals from the somata, which reflect signals derived from all subcellular compartments. Since
tufted cells and mitral cells both send their apical dendrites to the glomeruli, to study signals
from mitral cells in isolation, we used Lbhd2CreERT2::Ai95D mice, where GCaMP6f is expressed
predominantly in mitral cells [24] (**Fig. 6A,B**).
211
Imaging from the superficial plane, the apical dendrites showed no significant differences
between responses to S+ and S- odours (mean ΔF/F during odour = 0.307 ± 0.439 and 0.328 ±
0.466 for S+ and S- respectively; p = 0.687; post odour = 0.338 ± 0.512 and 0.382 ± 0.549 for S+
and S- respectively; p = 0.423, Wilcoxon rank-sum test, **Fig. 6C**). As before, signals from the
mitral cell somata imaged in the *Lbhd2-CreERT2::Ai95D* mice were characterised by the
widespread inhibitory component (mean ΔF/F during odour = -0.058 ± 0.077 and -0.034 ± 0.061
for S+ and S- respectively; p = 1.86 x 10^{-4}; post-odour = -0.024 ± 0.130 and 0.038 ± 0.131 for S+
and S- respectively; p = 1.08 x 10^{-5}, Wilcoxon rank-sum test; **Fig. 6D**) . Together, these data
suggest that the reward-related inhibition in mitral cells originates perisomatically.
221
If the inhibition in response to the rewarded cue in mitral cells is mediated via granule cells, we
should observe a greater GCaMP6f signal change to S+ odours specifically in the granule cells
that target mitral cells (**Fig. 7A**). The granule cells whose dendrites ramify in the deeper portion
of the external plexiform layer are thought to synapse with mitral cells [36-38], where mitral cell

lateral dendrites are found. These mitral cell-targeting granule cells are, however, present
intermixed with tufted cell-targeting granule cells.

To distinguish the putative mitral cell-targeting granule cells from those that target tufted cells,
the depth of external plexiform layer needs to be distinguished accurately *in vivo*. Towards this
end, we crossed *Lbhd2-CreERT2* mice with *Ai14* mice to express tdTomato preferentially in
mitral cells. We reasoned that despite tissue curvature or non-uniform thickness of the external
plexiform layer, this method would allow us to accurately separate the deeper from the
superficial portions based on the density and distribution of the tdTomato expression. Indeed,
the deep portion of the external plexiform layer showed higher density of thin red fluorescent
processes (**Fig. 7B,C**), while at more superficial depths, we observed occasional fluorescence
from thick processes, likely corresponding to the primary dendrites of mitral cells.

To study if the divergent odour responses in mitral cells can be explained by the evoked activity
of putative mitral cell-targeting granule cells, we turned to adult-born granule cells that develop
their dendrites in the deep external plexiform layer (**Fig. 7D,E**). Adult-born granule cells are
thought to be critical for refining odour responses in mitral and tufted cells when mice need to
discriminate between similar odours [39-43]. Further, since the mature adult-born granule cells
form dendro-dendritic synapses with mitral cell lateral dendrites, where GABA release can occur
locally [44], we sought to image directly from dendritic gemmules. Due to their small size, we
were cautious to exclude images from sessions that showed motion artefact, which was
determined by correlating the structural fluorescence pattern to the baseline period and
discarded those that showed low correlation (**Supplementary Fig. 3**). As a result, 70%
(1343/1917 trials) of the acquired data was discarded. Deep dendritic gemmules showed more
frequent inhibition compared to superficially located dendritic gemmules (**Fig 7F,**
Supplementary Fig. 4). This may reflect the reduced excitatory drive in the deep granule cells,
due to the prevalent reward-related inhibition in mitral cells. Thus, we analysed how the evoked
amplitude distribution in the deep vs. superficial dendritic gemmules relative to that of the
presynaptic counterparts, that is, against the distribution of evoked responses in mitral cells and
tufted cells, respectively. The S+ vs. S- tuning showed a close overlap between tufted cells and
superficial gemmules of adult-born granule cells. On the other hand, S+ vs. S- tuning distribution
of deep gemmules could not be explained by the mitral cell tuning distribution (**Fig. 7G**). There
was a tendency for these gemmules to respond more positively to the rewarded odours than
would be predicted from mitral cell activity alone. In other words, our data suggests that mitral
cell-targeting granule cells may receive an additional excitatory drive associated with the
rewarded odour.

Discussion

In this study, we observed a cell-type specific reward-associated inhibition in the primary

olfactory area of the mouse. This inhibition is cell-type specific and subcellular specific: first, it

manifests in the somata of mitral cells but not in tufted cells, and second, it appears in the

somata but not in their apical dendritic tuft in the input layer. This subcellular specificity

suggests that the generation of this phenomenon involves circuit components at a deeper layer

of the olfactory bulb. Further, the results of pseudo-conditioning and pharmacological

manipulations suggest that the mitral cell-specific, reward-related inhibition arises from an

acquired congruence of sensory and contextual signals. Thus, our study supports the notion

that value-related modulation of olfactory signals is a characteristic of olfactory processing in

the primary olfactory area and provides constraints on possible underlying mechanisms.

Many questions remain regarding the origin of the reward-related signals to the olfactory bulb.

Many brain regions send long-range projections to the olfactory bulb and are, therefore,

candidate drivers of the reward-related inhibition we observed. One important source of

reward-related signals to OB is the direct or indirect feedback projections from olfactory

cortices. Value-like modulation of olfactory responses occurs in many parts of the brain: It has

been observed in the prefrontal cortex [4,45], orbitofrontal cortex [4,45], hippocampus [46], olfactory

tubercle [47-50], piriform cortex [4,51], and anterior olfactory nucleus [4], although there may be

regional differences, for example, in the long-term stability of expression [45]. Of particular

interest is the piriform cortex, which serves as a gateway for processed signals for modulation of

the mitral cells [17]. While only a subset of pyramidal neurons from the piriform cortex project to

the olfactory bulb [52], a recent imaging study from olfactory bulb-projecting fibres showed value-

like activity when the task depended on olfactory cues [53]. It is unclear why we did not observe

the widespread reward-related modulation in tufted cells, even though some value-like activity

is present in the anterior olfactory nucleus [4], a region known to have modulatory influence over

tufted cells [17]. Since the anterior olfactory nucleus has multiple compartments [54] with each with

distinct long-range connectivity [55], it will be crucial for future studies to resolve how these

subregions contribute to associating values with olfactory stimuli.

Another potential source of reward-related signals are classical neuromodulatory regions. Of

particular interest are those that innervate the deeper layers of the olfactory bulb, including

cholinergic neurons of the nucleus of the horizontal limb of the diagonal band [19,56],

noradrenergic neurons of the locus coeruleus [57,58] and serotonergic neurons of the dorsal raphe

nucleus [59,60]. These regions contain neurons that show reward-related signals, although they

differ in the details. Cholinergic neurons of the basal forebrain are locked to the time of

anticipatory behaviour in reward-driven sensory tasks [61,62]. This timing is relatively late

compared to our phenomenon. In addition, an optogenetic activation of cholinergic fibers in OB

enhances, rather than inhibits, odour-evoked responses in mitral and tufted cells [19]. Normal
adrenergic transmission within the olfactory bulb is required for learning to discriminate similar
odours [63,64]. Neurons of the locus coeruleus, too, show instances of reward-locked activations [65].
As with acetylcholine, however, the timing is locked more to the anticipatory actions and
further, modulation may enhance, rather than inhibit, olfactory responses in the bulb during
delay conditioning [22]. Therefore, cholinergic and adrenergic inputs may not relate directly to the
phenomenon described in this study. On the other hand, among serotonergic neurons of the
dorsal raphe nucleus, some lock to rewarded cues [66-68], making the serotonergic input a
promising candidate for the reward-related inhibition in mitral cells. Precisely what the
serotonergic signal represents remains an active area of research, with possibilities including
reward [67], uncertainty [68], or a motor-sensory variable [69], though this may reflect regional
differences, too [68,70]. Since serotonergic neurons also target the striatum to affect the dopamine
release in the region [71], future experiments must resolve if they pose direct reward-related
effects on OB or via other brain areas.
316
The value-like activity in the higher olfactory areas mentioned above appears as excitatory
responses to the rewarded odour. In contrast, mitral cells of the olfactory bulb showed
enhanced inhibitory responses. This sign reversal suggests an involvement of inhibitory
interneurons like the granule cells. Due to the perisomatic nature of the reward-related
inhibition, the granule cells are a prime candidate. They are the major recipients of long-range
projections in the olfactory bulb and may target mitral cells vs. tufted cells separately by
ramifying dendrites at the deep vs. superficial levels of the external plexiform layer [36-38]. Despite
their numerical dominance [72], how granule cells contribute to signal transformation in the
olfactory bulb remains enigmatic. They are thought to contribute to the temporal precision of
the OB output [73-75], which is perceptible to the animals [76]. Granule cells seem to contribute only
subtly to mitral and tufted cells' spontaneous and evoked firing rates under anaesthesia or
when mice are not engaged in behavioural tasks [75]. However, they can potently silence mitral
and tufted cells when activated in bulk [75]. As a result, it has been hypothesised that an
excitatory drive from long-range projections onto granule cells is critical for potent physiological
effects [77]. Because feedback projections synapse more proximally to the soma, these have more
significant electrical impacts at the soma than inputs from the mitral and tufted lateral
dendrites arriving at the distal dendrites. This drive may be critical to a global activation of these
neurons [77], serving as a potential associative mechanism [44]. Our depth-specific imaging
approach may open new ways to investigate the physiology of these enigmatic interneurons.
336
Feedback signals from higher sensory regions may allow refined or processed information to be
integrated into more peripheral processing, as in the olfactory bulb. This system may be used to
dynamically tune the nature of early sensory processing to the behavioural demands at hand.

But given the timing we observed, where the modulation occurs after decisions have already
been made, feedback signals may be used to fine-tune the process of maturation of adult-born
granule cells and their integration into the existing circuitry within the olfactory bulb in a
behaviourally relevant manner [39-41]. It will be an intriguing future investigation to test if
disruption of feedback signals prevents the proper establishment of acquired connectivity and,
as a result, the expression of task-dependent activity patterns. In summary, our work brings us
closer to a mechanistic understanding of context-dependent modulation involving a congruent
interaction between local and long-range inputs.
348

Acknowledgement
We thank Yu-Pei Huang for technical assistance, Adam Mago and Sourjya Nath Baibhabee for
help with tamoxifen administration, the dedicated staff from the Animal Resources Section,
Imaging section, and mechanical engineering section of OIST Graduate University for the
support, and the members of Sensory and Behavioural Neuroscience Unit for helpful comments
on the project. This work was supported by Grant-in-Aid for Scientific Research (C) 22K06490
and OIST Graduate University.

Materials and Methods
358

Animals
All animal experiments had been approved by the Okinawa Institute of Science and Technology
Graduate University Graduate Animal Care and Use Committee (Protocol 2020–310). *Tbx21-Cre*
(Haddad et al., 2013), B6J.Cg-Gt(ROSA)26Sortm95.1(CAG- GCaMP6f)Hze/MwarJ, also known as
Ai95D (Madisen et al., 2015), and *Ai14* (Madisen et al., 2010) mice were originally obtained
from the Jackson Laboratory (stock numbers: 024507, 028865 and 007914, respectfully).
LBHD2-CreERT2 mice were generated previously and are also available through Jackson
Laboratory (stock number 036054; [24]). *Tbx21-Cre::Ai95D* and *LBHD2-CreERT2::Ai95D* mice were
generated by crossing parents homozygous for each transgene. *LBHD2-CreERT2::Ai14* mice were
generated by crossing LBHD2-CreERT2 mice with Ai14 mice.
369

Tamoxifen administration
To induce Cre-recombinase activity in *LBHD2-CreERT2::Ai95D* and *LBHD2-CreERT2::Ai14* mice,
tamoxifen injections (3 x 80 mg/kg at p21, s.c.), or tamoxifen diet (553.8 ± 362.6 mk/kg at p21)
was used. For injections, each of three consecutive days, tamoxifen solution (8 mg/ml; Sigma-
Aldrich T5648) was first dissolved in absolute ethanol. This was then added to corn oil, resulting
in a 5% ethanol and 95% corn oil mixture, and heated at 65 °C on a shaker for 30 minutes. After
it cooled down to room temperature, the solution was injected intraperitoneally (~ 100 μl).
Mice that were treated with the tamoxifen diet (2 mg/kg) were exposed to this for 2-4 days,
based on their initial weight, after which they were switched back to normal food (threshold to
switch to normal food: 80% initial body weight). Tamoxifen intake was calculated based on the
amount of diet food provided and amount of diet food left after switching back to normal food.

381
382
Surgery
All recovery surgeries were conducted in an aseptic manner. For the cranial window and
headplate implantations, 9-11 week-old male mice were deeply anesthetised with isoflurane (3-
5% for induction, 1-2% for maintenance; IsoFlo, Zoetis Japan). A craniotomy of approximately
1.5 x 1 mm was performed over the left olfactory bulb, and a custom cut glass window
(thickness No. 1; Matsunami, Japan) was implanted. Once the window was sealed with
cyanoacrylate (Histoacryl, B. Braun, Germany), a custom-made metal headplate (26 x 12 mm)
was implanted posterior to the cranial window. Dental acrylic (Kulzer, Hanau, Germany) was
then added to cover the exposed skull and to secure both the headplate and cranial window.
392
393 For experiments involving pharmacological infusion, an additional cannula (10 mm length;
394 C315GS- 4/SPC, Plastics One) was inserted to target the left anterior piriform cortex (coordinate:
395 AP -2.2 mm & ML -2.4 mm from bregma; DV -6.1 mm at 45-degree angle from brain surface),
396 similar to [18].
397
398 All mice were administered carprofen (5 mg/kg, i.p.) post-operatively for three consecutive
399 days. All mice were recovered for at least two weeks before the start of behavioural
400 experiments.
401
402
Virus injection
To express GCaMP6f in adult-born granule cells, during cranial window and headplate
implantations, *Lbhd2-CreERT2::Ai14* mice were injected with AAV1-syn-GCaMP6f-WPRE-SV40
(Addgene 100837-AAV1; titre was 1.84 x 10^{13} GC/mL at the time of synthesis) in the left SVZ
(200 µL; coordinate: AP -1.0 mm & ML -1.0 mm from bregma; DV -2.2 mm vertically) using
Nanoject III (Drummond, 3-000-207).
409
410
Habituation
Male mice were habituated to head fixation on a custom-made running wheel. Thereafter,
water access was restricted by removing water bottles from their home cages. The mice were
habituated to receive water from the port at the experimental setup on the following two to
three days mice, until they learned to drink at least 1 ml at the setup. The body weight was
recorded daily, to ensure that it stayed above 80% of the original weight. Lick responses were
measured using an IR beam sensor (PM- F25, Panasonic, Osaka, Japan).
418
419 During calcium imaging sessions, respiration of the mice was recorded using a flow sensor
420 (AWM3100V, Honeywell, NC) placed closely to the right nostril.
421
422
Discrimination training

After habituation, the mice were trained to associate one odour stimulus with a water reward
(S+ odour) and another odour stimulus with no reward (S- odour). Both S+ and S- odours were a
binary mix of ethyl butyrate (Sigma-Aldrich; W242705) and methyl butyrate (Tokyo Chemical
Industry; B0763), but mixed with different ratios based on the photoionization detector
readings. For the initial, easy, discrimination training, a 80/20 vs 20/80 ratio was used. When
mice reached 80% behavioural accuracy, fine discrimination training started, using 60/40 vs
40/60 odour mixtures. The correct response to S+ odours was to lick within a 3-second window
after stimulus onset, while the correct response to S- odours was to refrain from licking. The
water reward consisted of multiple drops, with a total of ~18 μL per trial. Olfactory stimuli were
presented using a custom flow-dilution olfactometer [78]. On each trial, odour was presented for
1 second and delay to the reward was 3 seconds from the odour onset. Inter-trial interval was
approximately 20 seconds.
436
437
438 *Long odour discrimination*
439 The mice proficient at the difficult discriminaition task were trained to discriminate between the
440 same odour mixtures but with 4 seconds of odour duration. The response window and reward
441 timing on S+ trials was the same between the two paradigms.
442
Disengagement paradigm
In this paradigm, the same odour mixtures as the difficult discrimination were used, but the
water reward was delivered every trial, approximately 15 seconds before the odour onset. The
time window used for measuring the anticipatory licks was identical to that of the fine
discrimination paradigm. The first session was considered a transition session and excluded
from imaging analysis.
449
450 *Random association paradigm*
451 In this paradigm, the water reward was presented in 50% of the trials, regardless of the odour
452 identity. This decoupled the odour identity and reward, but kept mice engaged, indicated by the
453 anticipatory licks. The first session was considered a transition session and excluded from
454 imaging analysis.
455
456
Pharmacological inactivation of anterior piriform cortex
For pharmacological inactivation of the anterior piriform cortex, muscimol (M1523, Sigma-
Aldrich, MO) was infused (2 mM in Ringer; 500 nL at 100 nL/min) through the previously
implanted cannula, using a Hamilton syringe (1 μL Model 7001KH PST-3 80100, Hamilton
Company, Nevada USA), approximately 10 minutes before the start of the imaging sessions. Two
days prior to the first muscimol infusion, Ringer solution (500 nL at 100 nl/min) was infused.
463
Histology
After the conclusion of the behavioural experiments, the mice were perfused transcardially with
phosphate buffer (in mM): NaH2 PO4 (225.7), Na2HPO4 (774.0) with pH adjusted to 7.4,
followed by PFA solution (4% dissolved in phosphate buffer). For mice that were implanted with

a cannula, 500 nL DiI (Invitrogen, V22885) was injected prior to the perfusion to mark the

cannula tip location. Coronal sections of 100 µm thickness were cut on a vibratome (5100 mz-

Plus, Campden Instruments, Leicestershire, UK) and counterstained using DAPI (D9542, Sigma-

Aldrich). Images were acquired using a Leica SP8 confocal microscope using a ×10 (NA 0.40 Plan-

Apochromat, 506407, Leica) objective.

Immunohistochemistry

Free floating olfactory bulb sections from above were first blocked in blocking solution (0.025 M

Tris-HCl, 0.5 M NaCl, 0.2% triton X-100, 7.5% normal goat serum, 2.5% BSA, pH = 7.5) for 60min

at room-temperature. Sliced were subsequently stained with chicken anti-GFP (Abcam,

ab13970; 1:500 in blocking solution) at 4°C over-night. Slices were washed three times in TBS

(0.025 M Tris-HCl, 0.5 M NaCl, pH = 7.5) and incubated in goat anti-chicken Alexa-488 (Abcam,

ab150169; 1:1000 in TBS supplemented with 0.2% triton X-100) for 2 hours at room-

temperature. All slices were counter stained with DAPI (D9542, Sigma- Aldrich). Images were

acquired using a Leica SP8 confocal microscope using a 10X (NA 0.40 Plan-Apochromat) or a 40X

(NA 1.10 Plan-Apochromat, 506357, Leica) objective.

In vivo calcium imaging

All the calcium data presented in this manuscript were obtained from awake mice. Two-photon

fluorescence of GCaMP6f and tdTomato were measured simultaneously with a custom-made

miscroscope (INSS, UK) fitted with a 25x objective (Nikon N25X-APO-MP1300, 1.1 N.A.) or a 16x

objective (Nikon N16XLWD-PF, 0.8 NA), and high-power laser (930 nm; Insight DeepSee, MaiTai

HP, Spectra-Physics, USA) at depths 50–400 µm below the surface of the olfactory bulb. Images

from a single plane were obtained at ~30 Hz with a resonant scanner. In each trial, 400 image

frames were acquired, with 100 frames before odour stimulus to obtain a baseline. Each day,

the stage coordinates were chosen relative to a reference location, which was determined by

the surface blood vessel pattern. Fields of view were 512 µm x 512 µm for apical dendrites, 256

µm x 256 µm for tufted and mitral cell somata, 128 µm x 128 µm and for adult-born granule cell

gemmules. Calcium data during difficult discrimination and disengaged experiments were

obtained from *Tbx21-Cre::Ai95D* mice (Figs. 1 and 4). 6 male mice were used for somata

imaging. All 6 were used to image mitral cell somata, while a subset (3 mice) were used to

image tufted cell somata. To obtain calcium data from different subcelluar compartments of

mitral cells, we used *LBHD2-CreERT2::Ai95D* mice (Fig. 6). For long odour discrimination,

random association, and muscimol infusion experiments, calcium data was obtained from both

Tbx21-Cre::Ai95D and *LBHD2-CreERT2::Ai95D* mice (Figs. 3-5). Finally, *LBHD2-CreERT2::Ai14*

mice were used in to record red (tdTomato, mitral cells) and green (calcium indicator,

gemmules) fluorescent signals during adult-born granule cell imaging experiments (Fig. 7).

Data analysis

All data was analyzed offline using custom MATLAB (MathWorks, USA) routines. To calculate the

behavioural accuracy, the number of licks during a 3 second window from final valve opening

until reward presentation was counted for each trial (anticipatory licks). Correct response to the

rewarded odour was a minimum of 2 anticipatory licks, and correct response to the unrewarded

odour was less then 2 anticipatory licks. Behavioural accuracy was calculated as the percentage
of correct trials from the total number of trials.

To calculate the sniffing frequency and speed of inhalation, the sniffing signal was first filtered (1
Hz high-pass and 30 Hz low-pass) and normalised (z-score). Inhalation peaks were detected
using the *findpeaks* MATLAB function. Sniff onsets were determined by searching back in time
from each detected inhalation peak to the point where the signal crossed a threshold value. The
detected onsets and peaks were then used to calculate the frequency (as 1/inter-onset time)
and speed of inhalation (as onset-to-peak time).

Image analysis
For each field of view, the imaging data was manually curated based on motion artifacts and
drift over time. Data with motion artifacts and/or drift were motion corrected using the
NoRMCorre toolbox (Pnevmatikakis and Giovannucci, 2017) and, when unsuccessfully
corrected, excluded from analysis. Regions of interest (ROIs) were manually drawn using ImageJ
(NIH, Bethesda, USA) based on the average field of view from each imaging session and
exported for usage in MATLAB. Average pixel value from each ROI was offset with a value from
the darkest region in the frame (e.g. a blood vessel). To account for bleaching over the course of
the imaging session, the mean pixel values for all trials were concatenated and detrended using
the MATLAB function *detrend*, then reshaped back into an array (individual trials x frames)
before relative fluorescence change was obtained. For each ROI, the change in fluorescence
(ΔF/F) was calculated by subtracting the mean pixel value from the baseline period (1 second
before odour stimulus onset), and dividing by the baseline value. Odour evoked responses were
calculated as the mean fluorescence change during the odour stimulus presentation, and post-
odour evoked responses as the mean fluorescence change between odour stimulus offset and
reward presentation. For the 'long odour' and 'random association' experiments, the time
windows to calculate the evoked responses were based on the fine odour discrimination
experiments.

Lick-aligned average
Rewarded trials were analysed for each imaging session. Onsets of anticipatory licks were
defined as the average time of first two licks observed after the start of odour presentation.
Rewarded trials were grouped into early vs late lick trials if the anticipatory lick onsets occurred
before or after the median onset time, respectively. Within each group, calcium transients were
aligned to the anticipatory lick onset time for each rewarded trial, and averaged.

Quality assessment abGC imaging
Small abGC gemmules makes them susceptible to motion artifacts in behaving animals. To
objectively assess the quality of the imaged trial, the tdTomato signals from the MC dendrites in
LBHD2-CreERT2::Ai14 mice were analysed. Rolling averages of 5 frames (step size: 1 frame) were
made and compared against the average of 50 frames obtained during the baseline period to
compute the correlation coefficient. If the mean correlation value for the period analysed
(between the odour offset and onset of water reward) was below 95% of the mean value during
the baseline period, the trial was rejected. Separate quality check was conducted for the odour

and post-odour phase. Further, trials where the baseline correlations deviated significantly were
considered outliers and rejected. This was assessed using the Matlab function *isoutlier*. This
quality control method resulted in 16 accepted fields of view from 5 mice for the odour period,
and 13 fields of view from 4 mice for the post-odour period. On average, a given accepted fields
of view yielded 3.2 ± 1.4 ROIs (3.8 ± 1.5 ROIs for deep FOVs, and 2.9 ± 1.4 ROIs for superficial
fields of view for the magnification used (128 μm x 128 μm frame size).
561
562
External plexiform layer depth determination based on red fluorescence
Depth within the external plexiform layer was estimated using the red fluorescence signal from
mitral dendrites in Lbhd2-CreERT2::Ai14 mice, which is dense in the deeper portion. Z-stack
ranging from the superficial layer to mitral cell layer spanned 250 μm (100 frames averaged
every 4 μm) obtained from the same x-y location as the functional imaging was used. Since the
fibre-like structures are the relevant signals, the averaged frame from each depth was passed
through a filter available as a plug-in in ImageJ ("Tubeness"; [79]), with the sigma parameter set to
2 μm.
571
572 *Normalised difference*
573 For each trial, the average value of relative fluorescence change was calculated for the odour
574 period (first 1 second after the odour onset) and the post-odour period (1-3 s after the odour
575 onset). The normalized difference between S+ and S- response amplitudes for the odour period,
576 as well as the post-odour period, is calculated as follows:
577

$$\frac{(\frac{\sum_i^n x_i}{n} - \frac{\sum_i^n y_i}{n})}{(max(\boldsymbol{x}) + max(\boldsymbol{y}))}$$

579
580 where *i* denotes the trial index, *n* is the number of trials, x is the evoked fluorescence change in
581 response to the rewarded odour, and y is the evoked fluorescence change in response to the
582 unrewarded odour.
583
584
Statistics
Divergent responders
To determine if a ROI showed a divergent response, for each ROI odour evoked and post-odour
response amplitudes for S+ and S- trials were tested for statistical significance using the
Wilcoxon rank sum test. Summary transients presented in figures show mean ± SEM, unless
otherwise stated.
591
Discrimination time
The method for determining the discrimination time was modified from [80]. Cumulative
histograms of the detected licks were calculated for all trials using 50 ms time bins. For each
time bin, the histogram values for S+ trials and for S- trials were tested for statistical significance

using a Wilcoxon rank sum test. The first time bin where histograms were significantly different
(p-value less than 0.05) was taken as the discrimination time.

1. Platt, M.L., and Glimcher, P.W. (1999). Neural correlates of decision variables in parietal
cortex. Nature *400*, 233-238. 10.1038/22268.
2. Sugrue, L.P., Corrado, G.S., and Newsome, W.T. (2004). Matching Behavior and the
Representation of Value in the Parietal Cortex. Science *304*, 1782-1787.
10.1126/science.1094765.
3. McCoy, A.N., Crowley, J.C., Haghighian, G., Dean, H.L., and Platt, M.L. (2003).
Saccade Reward Signals in Posterior Cingulate Cortex. Neuron *40*, 1031-1040.
https://doi.org/10.1016/S0896-6273(03)00719-0.
4. Ottenheimer, D.J., Hjort, M.M., Bowen, A.J., Steinmetz, N.A., and Stuber, G.D. (2023).
A stable, distributed code for cue value in mouse cortex during reward learning. Elife *12*.
10.7554/eLife.84604.
5. Schultz, W., Dayan, P., and Montague, P.R. (1997). A neural substrate of prediction and
reward. Science *275*, 1593-1599. 10.1126/science.275.5306.1593.
6. Louie, K., and Glimcher, P.W. (2012). Efficient coding and the neural representation of
value. Annals of the New York Academy of Sciences *1251*, 13-32.
https://doi.org/10.1111/j.1749-6632.2012.06496.x.
7. Schultz, W. (2015). Neuronal Reward and Decision Signals: From Theories to Data.
Physiological Reviews *95*, 853-951. 10.1152/physrev.00023.2014.
8. Wurtz, R.H., and Mohler, C.W. (1976). Enhancement of visual responses in monkey
striate cortex and frontal eye fields. Journal of Neurophysiology *39*, 766-772.
10.1152/jn.1976.39.4.766.
9. Goltstein, P.M., Reinert, S., Bonhoeffer, T., and Hübener, M. (2021). Mouse visual cortex
areas represent perceptual and semantic features of learned visual categories. Nature
Neuroscience *24*, 1441-1451. 10.1038/s41593-021-00914-5.
10. Roelfsema, P.R., Lamme, V.A.F., and Spekreijse, H. (1998). Object-based attention in the
primary visual cortex of the macaque monkey. Nature *395*, 376-381. 10.1038/26475.
11. Shuler, M.G., and Bear, M.F. (2006). Reward Timing in the Primary Visual Cortex.
Science *311*, 1606-1609. doi:10.1126/science.1123513.
12. Doucette, W., and Restrepo, D. (2008). Profound Context-Dependent Plasticity of Mitral
Cell Responses in Olfactory Bulb. PLOS Biology *6*, e258. 10.1371/journal.pbio.0060258.
13. Kay, L.M., and Laurent, G. (1999). Odor- and context-dependent modulation of mitral
cell activity in behaving rats. Nature Neuroscience *2*, 1003-1009. 10.1038/14801.
14. Jordan, R., Fukunaga, I., Kollo, M., and Schaefer, A.T. (2018). Active Sampling State
Dynamically Enhances Olfactory Bulb Odor Representation. Neuron *98*, 1214-
1228.e1215. 10.1016/j.neuron.2018.05.016.
15. Wachowiak, M. (2011). All in a Sniff: Olfaction as a Model for Active Sensing. Neuron
71, 962-973. https://doi.org/10.1016/j.neuron.2011.08.030.

16. Verhagen, J.V., Wesson, D.W., Netoff, T.I., White, J.A., and Wachowiak, M. (2007). Sniffing controls an adaptive filter of sensory input to the olfactory bulb. Nature Neuroscience *10*, 631-639. 10.1038/nn1892.

17. Chae, H., Banerjee, A., Dussauze, M., and Albeanu, D.F. (2022). Long-range functional loops in the mouse olfactory system and their roles in computing odor identity. Neuron *110*, 3970-3985 e3977. 10.1016/j.neuron.2022.09.005.

18. Otazu, G.H., Chae, H., Davis, M.B., and Albeanu, D.F. (2015). Cortical Feedback Decorrelates Olfactory Bulb Output in Awake Mice. Neuron *86*, 1461-1477. 10.1016/j.neuron.2015.05.023.

19. Rothermel, M., Carey, R.M., Puche, A., Shipley, M.T., and Wachowiak, M. (2014). Cholinergic inputs from Basal forebrain add an excitatory bias to odor coding in the olfactory bulb. J Neurosci *34*, 4654-4664. 10.1523/JNEUROSCI.5026-13.2014.

20. Petzold, G.C., Hagiwara, A., and Murthy, V.N. (2009). Serotonergic modulation of odor input to the mammalian olfactory bulb. Nat Neurosci *12*, 784-791. 10.1038/nn.2335.

21. Linster, C., and Fontanini, A. (2014). Functional neuromodulation of chemosensation in vertebrates. Curr Opin Neurobiol *29*, 82-87. 10.1016/j.conb.2014.05.010.

22. Doucette, W., Gire, David H., Whitesell, J., Carmean, V., Lucero, Mary T., and Restrepo, D. (2011). Associative Cortex Features in the First Olfactory Brain Relay Station. Neuron *69*, 1176-1187. https://doi.org/10.1016/j.neuron.2011.02.024.

23. Beshel, J., Kopell, N., and Kay, L.M. (2007). Olfactory bulb gamma oscillations are enhanced with task demands. J Neurosci *27*, 8358-8365. 10.1523/JNEUROSCI.1199-07.2007.

24. Koldaeva, A., Zhang, C., Huang, Y.P., Reinert, J.K., Mizuno, S., Sugiyama, F., Takahashi, S., Soliman, T., Matsunami, H., and Fukunaga, I. (2021). Generation and Characterization of a Cell Type-Specific, Inducible Cre-Driver Line to Study Olfactory Processing. J Neurosci *41*, 6449-6467. 10.1523/JNEUROSCI.3076-20.2021.

25. Reuschenbach, J., Reinert, J.K., Fu, X., and Fukunaga, I. (2023). Effects of Stimulus Timing on the Acquisition of an Olfactory Working Memory Task in Head-Fixed Mice. J Neurosci *43*, 3120-3130. 10.1523/JNEUROSCI.1636-22.2023.

26. Dana, H., Sun, Y., Mohar, B., Hulse, B.K., Kerlin, A.M., Hasseman, J.P., Tsegaye, G., Tsang, A., Wong, A., Patel, R., et al. (2019). High-performance calcium sensors for imaging activity in neuronal populations and microcompartments. Nat Methods *16*, 649-657. 10.1038/s41592-019-0435-6.

27. Haddad, R., Lanjuin, A., Madisen, L., Zeng, H., Murthy, V.N., and Uchida, N. (2013). Olfactory cortical neurons read out a relative time code in the olfactory bulb. Nat Neurosci *16*, 949-957. 10.1038/nn.3407.

28. Chu, Monica W., Li, Wankun L., and Komiyama, T. (2016). Balancing the Robustness and Efficiency of Odor Representations during Learning. Neuron *92*, 174-186. https://doi.org/10.1016/j.neuron.2016.09.004.

29. Yamada, Y., Bhaukaurally, K., Madarasz, T.J., Pouget, A., Rodriguez, I., and Carleton, A. (2017). Context- and Output Layer-Dependent Long-Term Ensemble Plasticity in a Sensory Circuit. Neuron *93*, 1198-1212 e1195. 10.1016/j.neuron.2017.02.006.

30. Kudryavitskaya, E., Marom, E., Shani-Narkiss, H., Pash, D., and Mizrahi, A. (2021). Flexible categorization in the mouse olfactory bulb. Current Biology *31*, 1616-1631.e1614. https://doi.org/10.1016/j.cub.2021.01.063.

31. Adefuin, A.M., Lindeman, S., Reinert, J.K., and Fukunaga, I. (2022). State-dependent
representations of mixtures by the olfactory bulb. Elife *11*. 10.7554/eLife.76882.
32. Markopoulos, F., Rokni, D., Gire, D.H., and Murthy, V.N. (2012). Functional properties
of cortical feedback projections to the olfactory bulb. Neuron *76*, 1175-1188.
10.1016/j.neuron.2012.10.028.
33. Boyd, A.M., Sturgill, J.F., Poo, C., and Isaacson, J.S. (2012). Cortical feedback control of
olfactory bulb circuits. Neuron *76*, 1161-1174. 10.1016/j.neuron.2012.10.020.
34. Chaudhury, D., Escanilla, O., and Linster, C. (2009). Bulbar acetylcholine enhances
neural and perceptual odor discrimination. J Neurosci *29*, 52-60.
10.1523/JNEUROSCI.4036-08.2009.
35. Kapoor, V., Provost, A.C., Agarwal, P., and Murthy, V.N. (2016). Activation of raphe
nuclei triggers rapid and distinct effects on parallel olfactory bulb output channels. Nat
Neurosci *19*, 271-282. 10.1038/nn.4219.
36. Mori, K., Kishi, K., and Ojima, H. (1983). Distribution of dendrites of mitral, displaced
mitral, tufted, and granule cells in the rabbit olfactory bulb. J Comp Neurol *219*, 339-355.
10.1002/cne.902190308.
37. Orona, E., Scott, J.W., and Rainer, E.C. (1983). Different granule cell populations
innervate superficial and deep regions of the external plexiform layer in rat olfactory
bulb. J Comp Neurol *217*, 227-237. 10.1002/cne.902170209.
38. Ezeh, P.I., Wellis, D.P., and Scott, J.W. (1993). Organization of inhibition in the rat
olfactory bulb external plexiform layer. J Neurophysiol *70*, 263-274.
10.1152/jn.1993.70.1.263.
39. Lepousez, G., Nissant, A., Bryant, A.K., Gheusi, G., Greer, C.A., and Lledo, P.M. (2014).
Olfactory learning promotes input-specific synaptic plasticity in adult-born neurons. Proc
Natl Acad Sci U S A *111*, 13984-13989. 10.1073/pnas.1404991111.
40. Livneh, Y., Adam, Y., and Mizrahi, A. (2014). Odor processing by adult-born neurons.
Neuron *81*, 1097-1110. 10.1016/j.neuron.2014.01.007.
41. Wu, A., Yu, B., Chen, Q., Matthews, G.A., Lu, C., Campbell, E., Tye, K.M., and
Komiyama, T. (2020). Context-dependent plasticity of adult-born neurons regulated by
cortical feedback. Sci Adv *6*. 10.1126/sciadv.abc8319.
42. Grelat, A., Benoit, L., Wagner, S., Moigneu, C., Lledo, P.-M., and Alonso, M. (2018).
Adult-born neurons boost odor–reward association. Proceedings of the National Academy
of Sciences *115*, 2514-2519. 10.1073/pnas.1716400115.
43. Shani-Narkiss, H., Vinograd, A., Landau, I.D., Tasaka, G., Yayon, N., Terletsky, S.,
Groysman, M., Maor, I., Sompolinsky, H., and Mizrahi, A. (2020). Young adult-born
neurons improve odor coding by mitral cells. Nature Communications *11*, 5867.
10.1038/s41467-020-19472-8.
44. Egger, V., Svoboda, K., and Mainen, Z.F. (2005). Dendrodendritic synaptic signals in
olfactory bulb granule cells: local spine boost and global low-threshold spike. J Neurosci
25, 3521-3530. 10.1523/JNEUROSCI.4746-04.2005.
45. Wang, P.Y., Boboila, C., Chin, M., Higashi-Howard, A., Shamash, P., Wu, Z., Stein, N.P.,
Abbott, L.F., and Axel, R. (2020). Transient and Persistent Representations of Odor Value
in Prefrontal Cortex. Neuron *108*, 209-224 e206. 10.1016/j.neuron.2020.07.033.
46. Biane, J.S., Ladow, M.A., Stefanini, F., Boddu, S.P., Fan, A., Hassan, S., Dundar, N.,
Apodaca-Montano, D.L., Zhou, L.Z., Fayner, V., et al. (2023). Neural dynamics

underlying associative learning in the dorsal and ventral hippocampus. Nature
Neuroscience *26*, 798-809. 10.1038/s41593-023-01296-6.

47. Martiros, N., Kapoor, V., Kim, S.E., and Murthy, V.N. (2022). Distinct representation of
cue-outcome association by D1 and D2 neurons in the ventral striatum's olfactory
tubercle. Elife *11*. 10.7554/eLife.75463.

48. Millman, D.J., and Murthy, V.N. (2020). Rapid Learning of Odor-Value Association in
the Olfactory Striatum. J Neurosci *40*, 4335-4347. 10.1523/JNEUROSCI.2604-19.2020.

49. Gadziola, M.A., Tylicki, K.A., Christian, D.L., and Wesson, D.W. (2015). The olfactory
tubercle encodes odor valence in behaving mice. J Neurosci *35*, 4515-4527.
10.1523/JNEUROSCI.4750-14.2015.

50. Gadziola, M.A., Stetzik, L.A., Wright, K.N., Milton, A.J., Arakawa, K., del Mar Cortijo,
M., and Wesson, D.W. (2020). A Neural System that Represents the Association of Odors
with Rewarded Outcomes and Promotes Behavioral Engagement. Cell Reports *32*,
107919. https://doi.org/10.1016/j.celrep.2020.107919.

51. Gire, D.H., Whitesell, J.D., Doucette, W., and Restrepo, D. (2013). Information for
decision-making and stimulus identification is multiplexed in sensory cortex. Nat
Neurosci *16*, 991-993. 10.1038/nn.3432.

52. Diodato, A., Ruinart de Brimont, M., Yim, Y.S., Derian, N., Perrin, S., Pouch, J.,
Klatzmann, D., Garel, S., Choi, G.B., and Fleischmann, A. (2016). Molecular signatures
of neural connectivity in the olfactory cortex. Nat Commun *7*, 12238.
10.1038/ncomms12238.

53. Hernandez Trejo, D., Ciuparu, P., Rebouillat, M., Davis, M.B., Gross, M., Muresan, R.C.,
and Albeanu, D.F. (2022). Cortical-bulbar feedback supports behavioral flexibility during
rule-reversal. Society for Nerosience Annual Meeting.

54. Haberly, L.B., and Price, J.L. (1978). Association and commissural fiber systems of the
olfactory cortex of the rat. II. Systems originating in the olfactory peduncle. J Comp
Neurol *181*, 781-807. 10.1002/cne.901810407.

55. Brunert, D., Medinaceli Quintela, R., and Rothermel, M. (2023). The anterior olfactory
nucleus revisited - An emerging role for neuropathological conditions? Prog Neurobiol
228, 102486. 10.1016/j.pneurobio.2023.102486.

56. Zaborszky, L., Carlsen, J., Brashear, H.R., and Heimer, L. (1986). Cholinergic and
GABAergic afferents to the olfactory bulb in the rat with special emphasis on the
projection neurons in the nucleus of the horizontal limb of the diagonal band. J Comp
Neurol *243*, 488-509. 10.1002/cne.902430405.

57. Shipley, M.T., Halloran, F.J., and de la Torre, J. (1985). Surprisingly rich projection from
locus coeruleus to the olfactory bulb in the rat. Brain Res *329*, 294-299. 10.1016/0006-
8993(85)90537-2.

58. McLean, J.H., Shipley, M.T., Nickell, W.T., Aston-Jones, G., and Reyher, C.K. (1989).
Chemoanatomical organization of the noradrenergic input from locus coeruleus to the
olfactory bulb of the adult rat. J Comp Neurol *285*, 339-349. 10.1002/cne.902850305.

59. Steinfeld, R., Herb, J.T., Sprengel, R., Schaefer, A.T., and Fukunaga, I. (2015). Divergent
innervation of the olfactory bulb by distinct raphe nuclei. J Comp Neurol *523*, 805-813.
10.1002/cne.23713.

60. McLean, J.H., and Shipley, M.T. (1987). Serotonergic afferents to the rat olfactory bulb:
I. Origins and laminar specificity of serotonergic inputs in the adult rat. J Neurosci *7*,
3016-3028. 10.1523/JNEUROSCI.07-10-03016.1987.

61. Parikh, V., Kozak, R., Martinez, V., and Sarter, M. (2007). Prefrontal acetylcholine
release controls cue detection on multiple timescales. Neuron *56*, 141-154.
10.1016/j.neuron.2007.08.025.
62. Hangya, B., Ranade, S.P., Lorenc, M., and Kepecs, A. (2015). Central Cholinergic
Neurons Are Rapidly Recruited by Reinforcement Feedback. Cell *162*, 1155-1168.
10.1016/j.cell.2015.07.057.
63. Doucette, W., Milder, J., and Restrepo, D. (2007). Adrenergic modulation of olfactory
bulb circuitry affects odor discrimination. Learn Mem *14*, 539-547. 10.1101/lm.606407.
64. Mandairon, N., Peace, S., Karnow, A., Kim, J., Ennis, M., and Linster, C. (2008).
Noradrenergic modulation in the olfactory bulb influences spontaneous and reward-
motivated discrimination, but not the formation of habituation memory. Eur J Neurosci
27, 1210-1219. 10.1111/j.1460-9568.2008.06101.x.
65. Varazzani, C., San-Galli, A., Gilardeau, S., and Bouret, S. (2015). Noradrenaline and
dopamine neurons in the reward/effort trade-off: a direct electrophysiological comparison
in behaving monkeys. J Neurosci *35*, 7866-7877. 10.1523/JNEUROSCI.0454-15.2015.
66. Ranade, S.P., and Mainen, Z.F. (2009). Transient firing of dorsal raphe neurons encodes
diverse and specific sensory, motor, and reward events. J Neurophysiol *102*, 3026-3037.
10.1152/jn.00507.2009.
67. Liu, Z., Zhou, J., Li, Y., Hu, F., Lu, Y., Ma, M., Feng, Q., Zhang, J.E., Wang, D., Zeng, J.,
et al. (2014). Dorsal raphe neurons signal reward through 5-HT and glutamate. Neuron
81, 1360-1374. 10.1016/j.neuron.2014.02.010.
68. Grossman, C.D., Bari, B.A., and Cohen, J.Y. (2022). Serotonin neurons modulate
learning rate through uncertainty. Curr Biol *32*, 586-599 e587.
10.1016/j.cub.2021.12.006.
69. Kawashima, T., Zwart, M.F., Yang, C.T., Mensh, B.D., and Ahrens, M.B. (2016). The
Serotonergic System Tracks the Outcomes of Actions to Mediate Short-Term Motor
Learning. Cell *167*, 933-946 e920. 10.1016/j.cell.2016.09.055.
70. Ren, J., Friedmann, D., Xiong, J., Liu, C.D., Ferguson, B.R., Weerakkody, T., DeLoach,
K.E., Ran, C., Pun, A., Sun, Y., et al. (2018). Anatomically Defined and Functionally
Distinct Dorsal Raphe Serotonin Sub-systems. Cell *175*, 472-487 e420.
10.1016/j.cell.2018.07.043.
71. Wang, H.L., Zhang, S., Qi, J., Wang, H., Cachope, R., Mejias-Aponte, C.A., Gomez, J.A.,
Mateo-Semidey, G.E., Beaudoin, G.M.J., Paladini, C.A., et al. (2019). Dorsal Raphe Dual
Serotonin-Glutamate Neurons Drive Reward by Establishing Excitatory Synapses on
VTA Mesoaccumbens Dopamine Neurons. Cell Rep *26*, 1128-1142 e1127.
10.1016/j.celrep.2019.01.014.
72. Burton, S.D. (2017). Inhibitory circuits of the mammalian main olfactory bulb. J
Neurophysiol *118*, 2034-2051. 10.1152/jn.00109.2017.
73. Lagier, S., Carleton, A., and Lledo, P.M. (2004). Interplay between local GABAergic
interneurons and relay neurons generates gamma oscillations in the rat olfactory bulb. J
Neurosci *24*, 4382-4392. 10.1523/JNEUROSCI.5570-03.2004.
74. Bathellier, B., Lagier, S., Faure, P., and Lledo, P.M. (2006). Circuit properties generating
gamma oscillations in a network model of the olfactory bulb. J Neurophysiol *95*, 2678-
2691. 10.1152/jn.01141.2005.

75.   Fukunaga, I., Herb, J.T., Kollo, M., Boyden, E.S., and Schaefer, A.T. (2014). Independent
control of gamma and theta activity by distinct interneuron networks in the olfactory
bulb. Nat Neurosci *17*, 1208-1216. 10.1038/nn.3760.
76.   Chong, E., Moroni, M., Wilson, C., Shoham, S., Panzeri, S., and Rinberg, D. (2020).
Manipulating synthetic optogenetic odors reveals the coding logic of olfactory
perception. Science *368*. 10.1126/science.aba2357.
77.   Balu, R., Pressler, R.T., and Strowbridge, B.W. (2007). Multiple modes of synaptic
excitation of olfactory bulb granule cells. J Neurosci *27*, 5621-5632.
10.1523/JNEUROSCI.4630-06.2007.
78.   Koldaeva, A., Schaefer, A.T., and Fukunaga, I. (2019). Rapid task-dependent tuning of
the mouse olfactory bulb. Elife *8*. 10.7554/eLife.43558.
79.   Sato, Y., Nakajima, S., Shiraga, N., Atsumi, H., Yoshida, S., Koller, T., Gerig, G., and
Kikinis, R. (1998). Three-dimensional multi-scale line filter for segmentation and
visualization of curvilinear structures in medical images. Med Image Anal *2*, 143-168.
10.1016/s1361-8415(98)80009-1.
80.   Abraham, N.M., Spors, H., Carleton, A., Margrie, T.W., Kuner, T., and Schaefer, A.T.
(2004). Maintaining Accuracy at the Expense of Speed: Stimulus Similarity Defines Odor
Discrimination Time in Mice. Neuron *44*, 865-876. 10.1016/j.neuron.2004.11.017.
838
839

Figure 1: *Widespread inhibition is observed in mitral cells in response to rewarded odour.*
A, Schematic showing two major sources of inputs to the olfactory bulb. Left arrow represents olfactory nerve inputs. Right arrow represents long-range inputs from other brain areas. **B**, Odour was presented for 1 s and did not overlap with the reward that was delivered 2 s after the odour offset. **C**, Odours used in the Go/NoGo olfactory discrimination tasks. Blue font corresponds to the rewarded (S+) odour. Mice were head-fixed and had a cranial window implanted. **D**, Time course of task acquisition for easy and difficult discriminations defined in **C**. Imaging sessions took place in proficient mice (n = 6 mice). **E**, Left, imaging configuration. Mitral cell (MC) and tufted cell (TC) somata were distinguished by depth. Right, example fields of view for TC somata and MC somata. **Fi-iii**, Responses to S+ and S- odours measured in TC somata. **Fi**, Colormap representation of fluorescence change over time for all ROIs. **Fii**, Average responses to S+ (blue) and S- (black) odours from all ROIs. **Fiii**, Scatter plot comparing S- vs. S+ response amplitudes for the period shown in **Fii**. Dotted line represents unity (S- amplitude = S+ amplitude). Individual points correspond to ROIs. Black dots indicate S+ and S- responses that were significantly different. **Gi-iii**, Same as **Fi-iii**, but for MC somata. N = 150 and 428 ROIs, and 3 and 6 mice for TC somata and MC somata, respectively.

858
859
Figure 2: *Reward-related inhibition is locked to the odour presentation.*
A, Fluorescence change from mitral cell somata aligned to the time of 1st anticipatory lick after odour onset
(S+ trials). **B**, Predictions for two alternative hypotheses for late-lick vs. early-lick trials; if inhibition is odour-
locked, a shift in the peak inhibition is observed in the lick-aligned average. If the reward-related inhibition is
locked to generation of licks, the trough times will be the same for late vs. early lick trials relative to the onset
time of anticipatory lick. **C**, Lick raster plots from two example sessions. Late (black) vs. early (red) lick trials
were defined as trials where first anticipatory lick occurred later or earlier than the median lick onset time for
each session. **D**, Lick-aligned averages for early vs. late lick for all sessions. Each trace is an average for one
session. **E**, Shift in the peak trough time in the lick-aligned average compared against mean difference in the
lick onsets for the late vs. early trials. Each point corresponds to one imaging session. Pearson's correlation
coefficient = -0.555, p = 0.026 (N = 16 sessions, 6 mice).
871
872

Figure 3: *Longer odour presentation masks the appearance of divergent responses.*
A, Schematic showing dominance of sensory drive. **B**, A four-second dour pulse overlapped temporally with reward delivery. **C**, Colour map display showing GCaMP6f fluorescence change from mitral cell somata evoked by S+ and S- odours. **D**, Average fluorescence change from all ROIs in response to S+ (blue) and S- (black) odours. Mean and SEM shown. **E**, Scatter plots of average fluorescence change for S+ vs. S- odours for the time period indicated in **D**. Black dots indicate S+ and S- responses that were significantly divergent. (N = 210 ROIs, 5 mice).

886
887
Figure 4: *Occurrence of reward-related inhibition depends on the behavioural context, not odour identity.*
A, Hypotheses on the source of signals underlying differential S+ and S- responses in mitral cell somata; it
could derive from sensory stimuli (top) or from long-range inputs to the olfactory bulb (bottom). **B**,
Behavioural paradigms to decouple reward association while disengaging mice (middle) or engaging mice
(Random association). In disengagement sessions, reward was delivered every trial, preceding odour
presentations. In random association, reward followed both mixtures of EB and MB 50% of the time. **C**,
Timeline of experiments. Mice first performed fine olfactory discrimination, then went through either
disengagement or random association sessions. Imaging took place from day 2 in both cases. **D**, Number of
anticipatory licks (licks 3 seconds' window from odour onset) for the two odours for three behavioural
paradigms. Individual points correspond to each imaging session analysed. *** corresponds to p < 0.001
(*post-hoc* Tukey-Kramer multiple comparisons after 1-way ANOVA) **E**, Average fluorescence change of all ROIs
(mitral cell somata) for the odours indicated. **F**, Comparison of fluorescence change in response to the the
two odours for the odour period for disengagement sessions (left) and random association sessions (right).
Individual points correspond to ROIs. Darker points represent significantly divergent responses. **G**, Same as F,
but for post-odour period. N = 125 ROIs, 3 mice for disengagement and 301 ROIs, 5 mice for random
association.

910
911

912
913
Figure 5: *Intact piriform cortex is needed to observe the reward-related inhibition*
A, Hypothesis tested; anterior periform cortex is necessary for mitral cell divergence during odour
discrimination. **B**, Muscimol solution (2 mM; 500 nL) was infused via an implanted cannula targeted to
anterior piriform cortex. **C**, Timeline of experiments. After mice were trained on the discrimination task,
control and muscimol sessions alternated. One imaging session occurred per day. **D**, Example of DiI location
infused via an implanted cannula. Scale bar = 1 mm. **E**, Summary of cannula tip locations. **F**, Accuracy in
performance (% of trials with correct lick response) for control vs. muscimol sessions. Individual points
correspond to each imaging session. P = 0.004. **G**, Example fields of view matched across two conditions. **H**,
Timecourse of fluorescence change for control (top) and muscimol (bottom) sessions in matched ROIs. **I**,
Average fluorescence change for each ROI for S+ and S- odours during odour and post-odour periods. Darker
points represent significantly divergent responses. **J**, Cumulative fraction of ROIs for normalized difference in
fluorescence changes evoked by S+ and S-. N = 124 ROIs, 3 mice.
926
927

929
930
931   Figure 6: *Reward-related inhibition originates peri-somatically.*

932
A, Schematic showing possible subcellular compartments where divergent signals may arrive, namely, apical
dendrites in the glomerulus (upper arrows), and deeper, lateral dendrites (lower arrows). **B**, Left, Confocal
image showing GCaMP6f preferentially in mitral cells (MCs) in Lbhd2CreERT2::Ai95D mice. Right, Illustration
of imaging planes to obtain signals from the MC apical dendrites and somata. Scale bar = 100 μm. **C**, Analysis
of GCaMP6f signals from MC apical dendrites. **Ci**, Example field of view from the glomerular layer. **Cii**, Average
fluorescence change from all ROIs (glomeruli) in response to S+ and S- odours. **Ciii**, Comparison of average
GCaMP6f fluorescence change for individual ROIs evoked by S+ vs. S- odours for the periods indicated. Darker
points represent significantly divergent responses. **Di-iii**, Same as **Ci-iii**, but for MC somata. Scale bar = 50 μm.
N = 140 ROIs, 4 mice for apical dendrites and 321 ROIs, 7 mice for somata.

942
943
944

945
946
947 Figure7: *Imaging from deep adult-born granule cells suggests an extrinsic drive for mitral-targeting granule*
948 *cells.*
949

A, Schematic of local circuitry and hypothesis; mitral cells (MCs) synapse with granule cells (GCs) whose
dendrites ramify in the lower portion of EPL. The deep-ramifying granule cells may receive the contextual
signal that leads to the divergent responses in MC somata. **B**, Lower and upper portions of external plexiform
layer can be distinguished by the density of MC dendrites. tdTomato is preferentially expressed in MCs in
Lbhd2CreERT2::Ai14 mice. Scale bars = 30 μm for top and bottom images. **C**, Depth-dependent pixel intensity
histogram from an example z-stack obtained with a two-photon microscope in a Lbhd2CreERT2::Ai14 mouse.
D, Adult-born granule cells (abGCs) are made to express GCaMP6f by injecting AAVs in the subventricular zone
(SVZ). Adult-born granule cells are imaged 4 weeks after injection. RMS = rostral migratory stream. **E**, Example
confocal images showing amplified GCaMP6f signal from deep abGC (green), shown with tdTomato signals
(red) from MCs and DAPI signals (blue). Scale bars = 100 μm and 50 μm for left and right images, respectively.
F, Comparison of S+ vs. S- response difference distributions for deep abGCs (green) and superficially ramifying
abGCs (light blue, dotted line). **G**, Left, Comparison of S+ vs. S- response difference distributions for tufted cell
(TC) somata (black trace) and superficially ramifying abGCs (light blue); right, Comparison of S+ vs. S- response
difference distributions for MC somata (black trace) and deep abGCs (green).

Supplementary figure 1: *Time-course of decision-related behavioural output.*

A, Example raster plots showing lick times relative to the onset of odour (t = 0) and reward delivery (t = 3 s) of a proficient mouse. Trials have been sorted into rewarded (S+) and unrewarded (S-) trials. Whether the mice made the correct or incorrect decision was determined based on the number of licks observed between the odour onset and reward onset. Green ticks on the right indicate correct trials, and red ticks indicate incorrect trials. **B**, Calculation of discrimination time is the earliest time at which licking patterns for S+ and S- trials diverge significantly, at the 0.05 level, shown for the example session in **A**. **C**, Discrimination time for all mice presented in Figure 1. N = 17 sessions, 6 mice.

Supplementary figure 2: *Sniff patterns are not starkly different between rewarded vs. unrewarded trials.*

A, Example sniff signal from a flow sensor placed next to a nostril on the contralateral side to odour presentation. Upward signal (signal above dotted horizontal line) corresponds to inhalation. Inhalation peaks are shown with circles (top) and exhalation peaks are annotated with short vertical ticks (bottom). **B**, Instantaneous sniff frequency is defined as the reciprocal of the sniff interval, measured from one inhalation onset to the next inhalation onset. Speed of inhalation (time to peak) is defined as the time elapsed from the inhalation onset to the inhalation peak. **C**, Time course of instantaneous sniff frequency change relative to the odour period (light brown background) and post-odour period (demarcated with dotted lines). Mean and SEM shown. **D**, Cumulative histogram of instantaneous frequencies observed during post-odour period. Thin lines correspond to individual sessions, and thick lines correspond to average across imaging sessions. **E**, same as **D**, but for inhalation speed. N = 13 sessions, 7 mice.

1007
1008
1009 Supplementary figure 3: *A method for assessing the quality of adult-born granule cell imaging.*
1010
A, Example field of view from the red channel showing mitral cell dendrites. **B**, Image quality was determined
by the frame-by-frame similarity of red fluorescence patterns by calculating correlation in the tdTomato image
between the baseline period and other time points within the trial. Those with a high correlation coefficient
throughout the trial is deemed to have less drift e.g., due to animal's movements. **C**, Time course of red
fluorescence correlation values for the accepted dataset. **D**, Same as **C** but for rejected dataset. Of the 1917
trials imaged in total 574 trials were accepted and 1343 trials were rejected. This amounts to, on average,
27.7% acceptance rate for deep gemmules and 36.2% acceptance rate for superficial gemmules.

1037   Supplementary figure 4: *Statistics of evoked responses in depth-resolved adult-born granule cell dendrites.*
1038
1039   Histograms represent the frequencies of observed fluorescence changes in response to the rewarded odour
1040   (S+, left column) and unrewarded odour (S-, right column) for the odour period (top row) and post-odour
1041   period (bottom row). Green histograms correspond to data from deep gemmules, and light blue histograms
1042   correspond to superficial gemmules.