Spider mites collectively avoid plants with cadmium irrespective of their

frequency or the presence of competitors

Diogo Prino Godinho*[+], Inês Fragata*, Maud Charlery de la Masselière & Sara
Magalhães

cE3c, Centre for Ecology, Evolution and Environmental changes, Faculdade de Ciências da

Universidade de Lisboa, Edifício C2, Campo Grande 1749- 016 Lisbon, Portugal.

+ Corresponding author: dagodinho@fc.ul.pt

*co-first authors

Abstract

1. Plants can accumulate heavy metals from polluted soils on their shoots and use this to

defend themselves against herbivory. One possible strategy for herbivores to cope with the

reduction in performance imposed by heavy metal accumulation in plants is avoidance of

contaminated tissues. Such avoidance, however, may hinge upon the specific conditions faced by

herbivores.

2. Here, we tested whether the spider-mite *Tetranychus urticae* avoids tomato plants

contaminated with cadmium in presence of conspecifics or heterospecifics and depending on the

frequency of contaminated plants.

3. We show that individual spider mite females do not discriminate between discs with or

without cadmium. However, in a set-up where 200 mites were simultaneously given the choice

between four plants with or without cadmium, they collectively avoided plants with cadmium. This

happened when 50% of the plants were uncontaminated, but also when only a single plant

contained no cadmium. In addition, *T. urticae* did not discriminate between plants infested with its

competitor *T. evansi* and other uncontaminated plants but they preferred plants with competitors

when the other plants contained cadmium.

4. Our results show that aggregation may be an important mechanism through which spider

mites avoid contaminated plants. They also indicate that cadmium accumulation in plants is a

stronger selective pressure than interspecific competition with *T. evansi.* This suggests that non-

accumulating plants will suffer more from herbivory than accumulating plants in metal polluted

environments.

5. *Synthesis and applications*. We show that collective avoidance of metal-accumulating plants

by herbivores is robust to environmental conditions and may have important consequences for

species distribution and interactions in metal contaminated sites.

Keywords: metal hyperaccumulation; elemental defences; herbivore performance; host choice;
interspecific competition.

Introduction

Soils often contain heavy metals, either due to geochemical properties or to human activity (Alloway, 2013; Hou et al., 2017). Some plants, known as hyperaccumulators, can store metals translocated from the soil in their upper parts, up to levels that are toxic to most organisms (Baker, 1987). Such hyperaccumulation often results in a significant reduction in the performance of herbivores that feed on those plants (Boyd & Moar, 1999; Freeman et al., 2007; Kazemi-Dinan et al., 2014; Mohiley et al., 2020; Quinn et al., 2010). This observation is at the basis of the elemental defence hypothesis, which posits that plants use such hyperaccumulated metals as a defence against herbivores (Martens & Boyd, 1994).

Because metal accumulation is detrimental to most herbivores, one would expect the latter to evolve means to overcome this effect. Indeed, some herbivores may cope with elemental defences by evolving tolerance mechanisms. Evidence for such adaptation is, however, still scarce(Freeman et al., 2006; Xu et al., 2020), and not universal (Godinho, et al., 2023a).

Alternatively (or in addition), herbivores may avoid plants that have accumulated metals. This seems to be a common strategy (Kazemi-Dinan et al., 2014; Mogren and Trumble, 2010; Vickerman et al., 2003). For example, when given the choice between *Thlaspi caerulascens* plants with high or low zinc concentrations, the desert locust, *Schistocerca gregaria* preferred the latter, a preference pattern that was recapitulated on artificial food supplemented with zinc (Behmer et al., 2005). Another study showed that cadmium modifies the volatiles of *Populus yunnanensis*, and herbivores (the sawfly *Stauronematus compressicornis* and the leaf beetle *Plagiodera versicolora*) use these cues to avoid those plants (Lin et al., 2022).

Herbivores that have been exposed to contaminated food may use this information to ameliorate their choices. Indeed, grass miners (*Chromatomyia milii*) that have fed on grasses (*Holcus lanatus*) with cadmium avoid these plants more than naïve individuals (Pollard & Baker, 1997). However, feeding on metal-contaminated plants may also impair the behavioural decisions of herbivores (Mogren & Trumble, 2010). This, in turn, may affect their ability to avoid contaminated

food. Indeed, the ability of flies (*Drosophila melanogaster*) to avoid substrates with lead vanishes

when they develop on lead-contaminated food (Peterson et al., 2020).

Apart from affecting the choice between contaminated and non-contaminated food,

exposure to heavy metals may also affect other relevant behavioural decisions in herbivores

(Oliveira et al., 2022). For example, in presence of competitors, pyramid ants that had ingested

selenium had impaired foraging behaviour (De La Riva & Trumble, 2016). Similarly, exposure to zinc

compromised the foraging behaviour and predator avoidance in damselfly larvae (Janssens et al.,

2014). Because herbivores are often simultaneously exposed to abiotic and biotic pressures, it is key

to understand how their interactive effect on herbivore decisions affect their distribution (Tran et

al., 2019).

Herbivores are rarely alone. The presence of conspecifics may affect their distribution by

means of aggregative behaviour (Hunter, 2000; McLellan & Montgomery, 2023). Herbivores may

benefit from being with conspecifics, for example to better overcome plant defences (Wertheim et

al., 2005). Moreover, behavioural decisions may be affected by the presence of others (Prokopy &

Roitberg, 2001). For example, fruit flies (*Bactrocera tryoni*) avoid low-quality food patches more

efficiently when in groups (Morimoto et al., 2018).

Knowledge that metals affect collective behaviour has only been gathered in zebrafish

(Shelton et al., 2023). Whether organisms collectively avoid metal-contaminated food is expected to

hinge upon the relative role of acquisition of information and behavioural impairment that such

environments represent.

In this study, we test whether herbivores avoid plants with cadmium and whether such

avoidance hinges on the presence of conspecifics or heterospecifics, or on the frequency of

hyperaccumulating hosts. To this aim, we use the system composed of the spider mites *Tetranychus*

urticae and *T. evansi* feeding on tomato plants (*Solanum lycopersicum*), which hyperaccumulate

cadmium in their upper parts, resulting in decreased performance of the spider mites at high

concentrations (Godinho et al., 2018, 2022). *T. urticae* relies on several environmental cues to avoid

predators (Pallini et al., 1999), pathogens (Zélé et al., 2019) but see (Rodrigues et al., 2022), conspecific

(Pallini et al., 1997) and heterospecific competitors (Godinho, et al., 2020a). However, whether they

avoid metal-accumulating tomato plants and, if so, whether this hinges upon the environmental

context, remains an open question.

Material and Methods

Plant and mite rearing conditions

Tomato plants (*Solanum lycopersicum*, *var* MoneyMaker) were sown in a climatic room (25ºC, 61% of

humidity, light:dark = 16:8), in pots with soil (pH 5.0-6.0; SIRO, Portugal) and watered twice a week

with 100 mL of tap water for the two first weeks. Subsequently, plants were transplanted to a mixture

of 1:4 vermiculite, gardening soil (SIRO, Portugal), where they grew for three more weeks and were

watered twice a week with 100 mL of either distilled water or a cadmium solution (3mM or 2mM

according to the experiment, see below). All plants were also watered once a week with tap water to

avoid the lack of nutrients. Just before the installation of the setup, each plant was watered with 70

mL of tap water. This concentration of cadmium added to the soil ensures that the amount

accumulated by tomato plants in their leaves is higher than the hyperaccumulating threshold

(Godinho et al., 2018) and falls within values encountered in natural populations of hyperaccumulating

plants (Meyer et al., 2015).

The populations of spider mites used in this study were created in 2018 by performing

controlled crosses between three populations of *T. urticae* and four populations of *T. evansi* collected

in the field of Portugal in 2017 from tomato plants (Godinho et al., 2020b). The resulting outbred

populations have been maintained in high numbers (> 1000 adult females) in cages (28 x 39 x 28 cm)

each containing two entire cadmium-free tomato plants, under controlled conditions (25 ºC ± 1, 68%

of humidity, light: dark = 16:8). For the experiment on individual choice and performance, populations

were maintained in 5 replicates during 14 discrete generations with 220 adult mites being transferred

every 14 days, as part of a larger experiment. To ensure that females used in the experiments were

approximately of the same age, adult females where isolated on separate tomato leaves in plastic

boxes (14 x 20 x 14 cm) and allowed to lay eggs for 48h. In all experiments, we used adult females

resulting from this cohort, 14 days after egg laying.

The animals used in this research are not vertebrates or invertebrates with ethical and care

limitations.

Experimental set-up for individual choice and performance

To test whether spider mites individually choose between leaf discs with or without cadmium, mated

females were placed at the centre of plastic strips (1x3 cm) connecting two leaf discs (Ø 16 mm) placed

on soaked cotton wool, one made from a tomato plant supplied with cadmium (2 mM) and the other

from a plant without cadmium (Figure 1A). For each replicate, leaf discs were made from leaves of the

same age (either the 3rd or 4th counting from the cotyledons) and the age and position of the disc with

cadmium was randomized across trials. Females (N = 17 to 24 per replicate population; total N = 101)

were allowed to choose between leaves with or without cadmium for 24h, after which the position of

the female was registered. To measure the effects of each environment on spider mite performance,

females (N = 94 for plants with no cadmium and N= 97 for plants with cadmium) were isolated on leaf

discs (Ø 18 mm) made from leaves 3 to 5 (from the cotyledons), from plants supplied with 2mM

cadmium chloride or not. These leaf discs were kept on top of soaked cotton wool in petri dishes. 48

hours later, females were sacrificed and the number of eggs was counted. Assays were divided in

blocks consisting of four consecutive days.

Figure 1. Experimental set-ups to measure individual (A) and collective host choice (B) and performance (C). The white and black shapes represent plants without and with cadmium, respectively. A) Experimental set-up for individual host choice (left panel) and individual performance (right panel). B) Experimental set-up for collective host choice and performance and schematic representation of the 5 treatments. The leaf marked in red streaks (treatments 4 and 5) identifies the presence of competitors.

Experimental set-up for collective choice and performance

To measure how the presence of competitors and the frequency of plants with cadmium affected the

collective choice of spider mites, four entire tomato plants were placed in a plastic box and were

connected by a round plastic arena (15 cm of diameter) in which their 4th leaf was inserted in a hole

of 2 cm long at equal distance from each other and from the centre of the arena (Fig. 1B). We applied

a ring of Vaseline around the petiole of the 4th leaf close to the node to prevent mite dispersal to other

leaves of the plant. We then released 200 females of *T. urticae* in the centre of the plastic arena, from

where they could move to any plant, and counted the number of mites on each plant 24 hours later.

This setup was used to test whether spider mite distribution was random or affected by the choice of

conspecifics (i.e., if there was aggregation), by giving mites the choice between four uninfested

cadmium-free plants (treatment 1; Figure 1B). Additionally, we used this set up to address how the

distribution of spider mites was affected by the frequency of contaminated plants and the presence

of a competitor in contaminated and uncontaminated environments. To address the first question,

we placed mites in set-ups with either two (treatment 2) or three (treatment 3) plants with cadmium

(Figure 1B), with the remaining plants being cadmium-free. To answer the second question, we

created two other treatments, in which the leaf of one plant without cadmium was infested with 100

adult females of the competitor species *T. evansi* (50 females on the first leaflet and 25 on the two

lateral leaflets to avoid overexploitation by the competitor). This plant was surrounded by three plants

either with (treatment 4) or without cadmium (treatment 5; Fig. 1B). The 200 *T. urticae* females were

placed in the centre of the plastic arena 24 hours after the installation of *T. evansi*. The number of *T.*

urticae females on each plant was counted 24 hours later.

To measure the performance of *T. urticae* on plants with or without cadmium, the number of

eggs on each leaf was counted on all plants in treatments 1, 2 and 3 after having counted (and

removed) the females to measure choice. The number of eggs was then divided by the number of

females on the same plant, to obtain the per capita oviposition rate. To measure the performance of

T. urticae in presence of competitors (treatments 4 and 5), we quantified the number of adult female

offspring after fifteen days, because the eggs of the two species are not distinguishable. To this aim,

we removed the females from the infested leaf of the plant with competitors (from treatments 4 and

5) or from one random plant without cadmium in the respective controls (treatments 1 and 3). Each

leaf was detached and placed in a pot with water in an isolated box. Fifteen days later, the number of

adult females was counted. Twenty replicates per treatment were done (except for treatment 5 for

which 19 replicates were done).

Statistical analysis

<u>Individual choice and performance</u>

To test if individual females choose between cadmium or cadmium-free leaf discs, a general linear

mixed model with a Binomial error distribution was used. The dependent variable was the presence

or absence (coded as 1 or 0, respectively) of the female in each environment (using the cbind function

in R). To test the effect of cadmium on the female performance, the number of eggs laid on each patch

was analysed using a general linear model with a Poisson error distribution. The dependent variable

was the number of eggs per patch, the test environment (cadmium-free vs. cadmium leaf discs) was

coded as a fixed factor. For these models leaf age and replicate were used as random factors.

<u>Collective choice and performance</u>

We first tested whether spider mites distribute themselves evenly across cadmium-free plants. To this

aim, we performed a G-test goodness of fit per replicate, assuming an even distribution of mites across

plants (*i.e.*, 25% per plant). To control for multiple tests, a Bonferroni correction was applied.

Moreover, we performed a general linear model comparing the control treatment to a mock

treatment where each plant 'received' 50 individuals of the 200 released.

All statistical analyses to test the collective spider mite behaviour were done using a general

linear model with number of individuals on a plant as the dependent variable. All models included

block and interaction between identity of the plant and replicate as random factors. Models where

initially fit with a Poisson distribution, but due to overdispersion, a negative binomial error distribution

was used instead.

To test whether spider mites collectively avoided plants with cadmium, we compared the

distribution of females in plants with and without cadmium (treatment 2, 3 and 4) using a model with

cadmium and treatment as fixed factors and their interaction. Post hoc contrasts were performed to

test whether spider mites avoided plants with cadmium in each treatment, and to test whether

treatments differed concerning this avoidance. To test if cadmium concentration affected spider mite

choice we compared treatments 2 and 3, and to test if presence of competitors affected choice of

plants with cadmium we compared treatments 3 and 4. Finally, we tested whether the presence of

cadmium in the environment affected the avoidance of competitors by comparing treatments 4 and

5 with a model with competitors and treatment as fixed factors and their interaction. Post hoc

contrasts were performed to test whether spider mites avoided plants with competitors in each

treatment, and whether treatments differed concerning this avoidance.

All statistical analyses to test collective spider mite performance were done using general

linear models with a gamma error distribution, and the number of per capita eggs or female offspring

(for the treatments with competitors) as dependent variables. All models included block and replicate

as random factors.

We first assessed spider mite performance in the control treatment (i.e., treatment 1) to test

whether mite oviposition rate was even across plants, using a generalized mixed model with plant as

fixed factor. The effect of cadmium on spider mite performance was assessed by comparing the per

capita oviposition rate across all plants among treatments containing plants with and without

cadmium (i.e., treatments 2, 3 and 4). For that we applied a model with cadmium and treatment and

their interaction as fixed factors. Next, we performed pairwise comparisons to test whether spider

mite performance differed on plants with or without cadmium within each treatment. We also

performed pairwise comparisons to test whether the performance on plants with cadmium differed

across treatments, only using data from plants with cadmium in each treatment.

To estimate if the effect of cadmium was similar for the individual and collective tests, we

calculated the standardized effect size for each treatment and experiment using the effectsize package

(Ben-Shachar et al., 2020).

To test the effect of the competitor on the number of per capita offspring we used a model

with presence (or absence) of competitors as fixed factor. We used posthoc tests to compare the

treatments with competitors and their respective control (i.e. treatments 1 vs 5 and 3 vs 4).

All statistical analyses were performed using the software R v3.6 (R Core Team, 2022), using

packages glmmTMB and lme4 to perform the general linear models (Bates et al., 2015; Brooks et al.,

2017), car for the ANOVA tests (Fox & Weisberg, 2019), emmeans for post-hoc contrasts (Lenth, 2023)

and ggplot2 for visualization (Wickham, 2009).

Results

<u>Individual choice and performance</u>

Individual spider mite females did not occupy differently leaf discs with and without cadmium

(estimated overall probability = 0.0554, z ratio= 1.064, P = 0.2875, Fig. 2A). They laid significantly fewer

eggs on leaf discs with cadmium than on leaf discs without cadmium (ratio between water and

cadmium= 2.1, z ratio= 10.432, p<0.001, Fig. 2B, Table S1).

Figure 2. Individual host choice and performance

(A) Number of females on leaf discs from plants with cadmium (dark grey) or without cadmium (light grey) after 24 hours. (B) Number of eggs laid on leaf discs from plants with or without cadmium. Error bars represent replicate standard error between replicates.

Collective choice and performance

Spider mites distributed themselves unevenly across plants in our set-up. In 18 out of the 19 replicates, they showed a distribution significantly different from the expected 25%, even after Bonferroni correction (Table S2). Indeed, the difference between the treatment with only cadmium-free plants and no competitors and a mock variable with an even distribution across plants was significant (effect of treatment: X^2_2=7410,5; P < 0.0001, Fig 3, treatment 1).

Cadmium significantly affected the number of females on each plant, but not treatment or their interaction (ANOVA results cadmium: X^2_1 =8.70, p-value =0.0031; treatment: X^2_1 = 1.678, p-value= 0.432; cadmium*treatment: X^2_1 = 1.674, p-value = 0.432; Fig. 3). Spider mites avoided plants with cadmium in all treatments (treatment 2 plants: z ratio=-55.987, p-value <0.0001, treatment 3: z ratio=-49.560, p-value <0.0001; treatment 4: z ratio=- 49.343, p-value <0.0001). This avoidance was

not different between treatments with two or with three plants (contrast: z ratio=-0.390, p value =

0.9196).

The presence of a competitor, and its interaction with treatment significantly affected the

number of females on each plant (competitor: X^2_1 = 4.556, p-value= 0.033; treatment: X^2_1 = 1.666, p-

value=0.197; cadmium*treatment: X^2_1 = 5.556, p-value =0.0184; Fig. 3). The number of mites on plants

with or without competitors when all plants were cadmium-free did not differ significantly (contrast:

z ratio=1.203, p-value = 0.229). However, in the treatment where plants without competitors had

cadmium, females chose more often the plant with the competitor (i.e. they avoided cadmium),

resulting in a significant difference among these treatments (contrast: z ratio= -2.134, p-value = 0.033).

This choice was not different from that documented when the uncontaminated plant did not have

competitors (contrast between treatment 3 and 4: z ratio=-0.457, p value= 0.891).

Figure 3. Collective host choice. Number of females after 24h on each plant for all treatments.

Numbers indicate the treatments: 1) control, 2) two plants with cadmium, 3) three plants with

cadmium, 4) one plant infested with competitors, 5) one plant infested with competitors and three

plants with cadmium. Dark grey bars represent plants with cadmium, light grey colour plants without

cadmium or competitor, white bars represent presence of competitors on the plants. Error bars

represent replicate standard error between replicates.

In the control treatment, spider mite oviposition rate was not significantly different among

plants (X^2_3 = 3.473, p-value = 0.324; Figure 4). The presence of cadmium in plants, in treatments 2, 3

and 4 significantly reduced the oviposition rate of females (X^2_2 = 89.745, p-value <0.0001; Figure 4).

However, the number of plants with cadmium did not affect spider-mite performance (X^2_2 = 0.413, p-

value = 0.813; Figure 4). In fact, the standardized effect size associated with the presence of cadmium

was remarkably similar for treatments 2 and 3 (Table S1). The effect size of cadmium was larger in

individual than in collective performance experiments but showed a clear overlap of the confidence

intervals (Table S1).

Figure 4. Collective performance. A) Number of eggs per female after 24h on each plant for

treatments 1-3. Dark grey bars represent plants with cadmium, light grey colour plants without

cadmium or competitor, white bars represent presence of competitors on the plants. Error bars

represent replicate standard error between replicates.

In environments with cadmium-free plants only, the presence of the competitor significantly

affected the number of offspring per female after one generation (X^2_1 = 15.877, p-value = 0.0004,

Figure 5). Contrasts between plants with or without competitors revealed that the number of female

offspring on uninfested plants (*i.e.*, in the absence of competitors or cadmium) was higher than on

cadmium-free plants with competitors (contrasts between treatment 1 and 5: t ratio= 3.771, p-

value=0.002). However, this difference was not observed when competitor-free plants contained

cadmium (contrasts between treatment 3 and 4: t ratio= -1.713, p-value=0.325).

Figure 5. Collective performance in presence of competitors. Light grey bars represent plants without

cadmium or competitor, white bars represent presence of competitors on the plants. Error bars

represent replicate standard error between replicates. Both types of plants contained no cadmium.

Discussion

Our study shows that *T. urticae* females do not avoid tomato leaves with cadmium when they are

given the choice individually, although feeding on these leaves entails a reduction in their

performance. In contrast, when in group, facing the choice between cadmium and cadmium-free

plants, females moved more often to plants without cadmium. They also showed a non-even

distribution, even when all plants were uninfested and uncontaminated, suggesting aggregation

behaviour. Moreover, the frequency of cadmium-free plants had no effect on spider-mite avoidance.

Finally, when all plants were cadmium-free, spider mites did not discriminate between plants with or

without competitors, but they avoided plants with cadmium when the alternative was moving to a

uncontaminated plant with competitors.

We found here that spider mites feeding tomato plants that have hyperaccumulated cadmium

have a lower oviposition rate than mite feeding on non-contaminated plants. This recapitulates

previous findings (Godinho et al., 2023b; Godinho et al., 2018, 2022), and implies that they are robust

to the set-up used. Curiously, however, *T. urticae* females did not choose between leaf discs with or

without cadmium, a result also previously obtained for *T. evansi* (Godinho et al., 2023a). This is

surprising, not only because of the abovementioned consequences of cadmium for performance, but

also because this choice set-up had been validated previously (Rodrigues et al., 2017; Zélé et al., 2019).

Possibly, cadmium does not affect plant cues that inform spider mites of its content. Alternatively,

such cues may be produced in whole plants but not in cut leaf discs.

In contrast to the individual choice assays, spider mites systematically avoided plants with

cadmium in a set-up that simultaneously tested 200 mites. Our results from the control treatment

unequivocally show that spider mites do aggregate on particular plants, a result also found in the field

(Rijal et al., 2016). Aggregation behaviour has been shown to be beneficial to *T. urticae*, by increasing

fecundity and decreasing the death rate of females living in groups (Le Goff et al., 2010). In the

treatments containing plants with and without cadmium, such aggregation occurred on plants without

cadmium. It is thus tempting to hypothesize that spider mites use the information of their conspecifics

to avoid plants with cadmium. Spider mites often communicate by means of odour cues, for example

when searching for mates (Regev & Cone, 1976; Rodrigues et al., 2017) or avoiding predators (Choh

& Takabayashi, 2010; Janssen et al., 1999). Possibly, aversive chemical information of spider mites

that have been on plants with cadmium is available in our set-up and used by spider mites to make

choices that are beneficial for their performance. Alternatively, cadmium disrupts the production of

plant volatiles induced by spider mites, which are also used by conspecifics to find infested plants (Lin et al., 2022; Pallini et al., 1997). In this case, cadmium-free plants are expected to become more attractive than plants with cadmium.

In environments without cadmium, *T. urticae* spider mites did not avoid plants with competitors h, which is at odds with competitors negatively affecting the number of offspring produced, as found here and in other studies (Fragata et al., 2022; Sarmento, et al., 2011). Possibly, avoidance of *T. evansi* is only observed in later stages of plant infestation, when resources become scarcer. Indeed, *T. evansi* is known to downregulate plant defences, which is beneficial to *T. urticae* in the short term (Godinho et al., 2016; Sarmento, et al., 2011). Moreover, *T. urticae* is able to avoid *T. evansi* within a single plant (Godinho, et al., 2020a), which may release the selection pressure to avoid whole plants.

In contrast, spider mites avoid plants with cadmium in detriment of avoiding competitors. An earlier study had shown that spider mites prefer young over old leaves, as the former result in higher oviposition rate for them, but that this preference vanishes when this leaf is infested with competitors (Godinho, et al., 2020a). Here, instead, we show that an abiotic selection pressure seems to be more relevant to spider mite host choice than the presence of competitors. The complex interaction between responses to abiotic and biotic stressors has been documented in other systems (Murphy & Loewy, 2015; Tran et al., 2019). One possible explanation for our results is that selecting hosts with *T. evansi* might be less risky than selecting plants with cadmium because *T. urticae* could still avoid *T. evansi* once on the plant by finding a leaf free of competitors (e.g.,(Godinho et al., 2020a), while cadmium is accumulated in all leaves and there is no way to avoid it within a plant.

The avoidance of cadmium even in the presence of competitors implies stronger competition on cadmium-free plants. This is in line with studies showing increased intra or interspecific competition in resource patches with higher quality in heterogeneous environments (Shen et al., 2020; Wang et al., 2012). This suggests that spider mites will be exposed to different selection pressures in heterogeneous environments: on plants with cadmium, selection will favour mites that

are able to physiologically cope with the toxic effects of metal, whereas on plants with no cadmium

mites will face stronger intra- and inter-specific competition. If these two selection pressures persist

for long enough, they may lead to high population differentiation within herbivores. Corroborating

this hypothesis would necessitate studying populations occurring on sites with both types of plants

(Freeman et al., 2007).

Alternatively, spider mites may not adapt to plants with cadmium, for example due to lack of

genetic diversity, in which case the observed cadmium avoidance would lead to plants with such metal

suffering few attacks by herbivores in the long run, as seen in other systems (Galeas et al., 2008; Noret

et al., 2007). This should select for the maintenance of metal accumulation in plants, especially

because spider mites do not individually avoid tomato plants, and not all mites avoid plants, even

when collectively choosing. Thus, plants with cadmium will still suffer some herbivore damage. Hence,

as long as the fitness cost of uptaking metal does not surpass the fitness gain of herbivore avoidance,

selection to uptake metals as a defence against herbivores will be maintained.

Conclusions

We demonstrated that spider-mites do not individually avoid tomato plants contaminated with

cadmium, despite having reduced oviposition on those plants. However, when choosing collectively,

they avoid cadmium contaminated plants, even when clean plants are colonized by a superior

competitor. These results suggest that collective choice in herbivores contributes to the maintenance

of plant elemental defences, as it entails less herbivory on those plants. Moreover, they imply that

hyperaccumulation of metals by plants may lead to stronger herbivore competition on nearby non-

accumulating plants, thus having broad effects on metal-polluted ecosystems.

Acknowledgements

We thank Cátia Eira, Inês Santos and Lucie de Sousa for maintaining plant and mite populations and

all the mite squad for fruitful discussions. This work was financed by an ERC (European Research

Council) consolidator grant COMPCON, GA 725419 attributed to SM and by FCT (Fundação para

Ciência e Tecnologia) with a Junior researcher contract (CEECIND/02616/2018) attributed to IF.

390   **Author contribution**

391   SM conceived the ideas and designed the methodology, with help from MCM; DG and MCM collected

392   the data; DG, IF and SM analysed the data and wrote the manuscript. All authors contributed critically

393   to the drafts and gave final approval for publication.

395   **Conflict of interest**

396   Authors declare no conflict of interest.

398   **Data availability**

399   Data is deposited in Figshare, DOI: 10.6084/m9.figshare.24511093.

401   **References**

Alloway, B. J. (2013). Heavy metals in soils: trace metals and metalloids in soils and their bioavailability.
In *Choice Reviews Online* (Vol. 50, Issue 07). Springer Science \& Business Media.
https://doi.org/10.5860/choice.50-3862

Baker, A. J. M. (1987). Metal Tolerance. *New Phytologist*, *106*, 93–111. https://doi.org/10.1111/j.1469-
8137.1987.tb04685.x

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects models using
lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/jss.v067.i01

Behmer, S. T., Lloyd, C. M., Raubenheimer, D., Stewart-Clark, J., Knight, J., Leighton, R. S., Harper, F. A.,
& Smith, J. A. C. (2005). Metal hyperaccumulation in plants: mechanisms of defence against insect
herbivores. *Functional Ecology*, *19*(1), 55–66. https://doi.org/https://doi.org/10.1111/j.0269-
8463.2005.00943.x

Ben-Shachar, M. S., Lüdecke, D., & Makowski, D. (2020). {e}ffectsize: Estimation of Effect Size Indices
and Standardized Parameters. *Journal of Open Source Software*, *5*(56), 2815.
https://doi.org/10.21105/joss.02815

Boyd, R. S., & Moar, W. J. (1999). The defensive function of Ni in plants: Response of the polyphagous
herbivore Spodoptera exigua (Lepidoptera: Noctuidae) to hyperaccumulator and accumulator

species of Streptanthus (Brassicaceae). *Oecologia*, *118*(2), 218–224.
https://doi.org/10.1007/s004420050721

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J.,
Maechler, M., & Bolker, B. M. (2017). {glmmTMB} Balances Speed and Flexibility Among Packages
for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal*, *9*(2), 378–400.
https://doi.org/10.32614/RJ-2017-066

Choh, Y., & Takabayashi, J. (2010). Predator avoidance by phytophagous mites is affected by the
presence of herbivores in a neighboring patch. *Journal of Chemical Ecology*, *36*(6), 614–619.
https://doi.org/10.1007/s10886-010-9792-4

De La Riva, D. G., & Trumble, J. T. (2016). Selenium exposure results in reduced reproduction in an
invasive ant species and altered competitive behavior for a native ant species. *Environmental*
Pollution, *213*, 888–894. https://doi.org/10.1016/j.envpol.2016.03.045

Fox, J., & Weisberg, S. (2019). *An {R} Companion to Applied Regression* (Third). Sage.

Fragata, I., Costa-Pereira, R., Kozak, M., Majer, A., Godoy, O., & Magalhães, S. (2022). Specific sequence
of arrival promotes coexistence via spatial niche pre-emption by the weak competitor. *Ecology*
Letters, *25*(7), 1629–1639. https://doi.org/10.1111/ele.14021

Freeman, J. L., Lindblom, S. D., Quinn, C. F., Fakra, S., Marcus, M. A., & Pilon-Smits, E. A. H. (2007).
Selenium accumulation protects plants from herbivory by Orthoptera via toxicity and deterrence.
New Phytologist, *175*(3), 490–500. https://doi.org/10.1111/j.1469-8137.2007.02119.x

Freeman, J. L., Quinn, C. F., Marcus, M. A., Fakra, S., & Pilon-Smits, E. A. H. (2006). Selenium-Tolerant
Diamondback Moth Disarms Hyperaccumulator Plant Defense. *Current Biology*, *16*(22), 2181–
2192. https://doi.org/10.1016/j.cub.2006.09.015

Galeas, M. L., Klamper, E. M., Bennett, L. E., Freeman, J. L., Kondratieff, B. C., Quinn, C. F., & Pilon-Smits,
E. A. H. (2008). Selenium hyperaccumulation reduces plant arthropod loads in the field. *New*
Phytologist, *177*(3), 715–724. https://doi.org/https://doi.org/10.1111/j.1469-8137.2007.02285.x

Godinho, D. P., Branquinho, C., & Magalhães, S. (2023b). Intraspecific variability in herbivore response
to elemental defences is caused by the metal itself. *Journal of Pest Science*, *96*(2), 797–806.
https://doi.org/10.1007/s10340-022-01525-0

Godinho, D. P., Cruz, M. A., Charlery de la Masselière, M., Teodoro-Paulo, J., Eira, C., Fragata, I.,
Rodrigues, L. R., Zélé, F., & Magalhães, S. (2020b). Creating outbred and inbred populations in
haplodiploids to measure adaptive responses in the laboratory. *Ecology and Evolution*, *10*(14),
7291–7305. https://doi.org/10.1002/ece3.6454

Godinho, D. P., Fragata, I., Majer, A., Rodrigues, L. R., & Magalhães, S. (2023a). Limits to the adaptation
of herbivorous spider mites to metal accumulation in homogeneous and heterogeneous
environments. *BioRxiv*. https://doi.org/10.1101/2023.03.15.532545

Godinho, D. P., Janssen, A., Dias, T., Cruz, C., & Magalhães, S. (2016). Down - regulation of plant defence
in a resident spider mite species and its effect upon con - and heterospecifics. *Oecologia*, *180*(1),
161–167. https://doi.org/10.1007/s00442-015-3434-z

Godinho, D. P., Janssen, A., Li, D., Cruz, C., & Magalhães, S. (2020a). The distribution of herbivores
between leaves matches their performance only in the absence of competitors. *Ecology and*
Evolution, *10*(15), 8405–8415. https://doi.org/10.1002/ece3.6547

Godinho, D. P., Serrano, H. C., Da Silva, A. B., Branquinho, C., & Magalhães, S. (2018). Effect of cadmium
accumulation on the performance of plants and of herbivores that cope differently with organic
defenses. *Frontiers in Plant Science*, *871*(November), 1–14.
https://doi.org/10.3389/fpls.2018.01723

Godinho, D. P., Serrano, H. C., Magalhães, S., & Branquinho, C. (2022). Concurrent herbivory and metal
accumulation: The outcome for plants and herbivores. *Plant-Environment Interactions*, *3*(4), 170–
178. https://doi.org/10.1002/pei3.10088

Hou, D., O'Connor, D., Nathanail, P., Tian, L., & Ma, Y. (2017). Integrated GIS and multivariate statistical
analysis for regional scale assessment of heavy metal soil contamination: A critical review.
Environmental Pollution, *231*, 1188–1200. https://doi.org/10.1016/j.envpol.2017.07.021

Hunter, A. F. (2000). Gregariousness and repellent defences in the survival of phytophagous insects.
Oikos, *91*(2), 213–224. https://doi.org/10.1034/j.1600-0706.2000.910202.x

Janssen, A., Pallini, A., Venzon, M., & Sabelis, M. W. (1999). Absence of odour-mediated avoidance of
heterospecific competitors by the predatory mite Phytoseiulus persimilis. *Entomologia*
Experimentalis et Applicata, *92*(1), 73–82. https://doi.org/https://doi.org/10.1046/j.1570-
7458.1999.00526.x

Janssens, L., Dinh Van, K., Debecker, S., Bervoets, L., & Stoks, R. (2014). Local adaptation and the
potential effects of a contaminant on predator avoidance and antipredator responses under global
warming: A space-for-time substitution approach. *Evolutionary Applications*, *7*(3), 421–430.
https://doi.org/10.1111/eva.12141

Kazemi-Dinan, A., Thomaschky, S., Stein, R. J., Krämer, U., & Müller, C. (2014). Zinc and cadmium
hyperaccumulation act as deterrents towards specialist herbivores and impede the performance
of a generalist herbivore. *New Phytologist*, *202*(2), 628–639. https://doi.org/10.1111/nph.12663

Le Goff, G. J., Mailleux, A. C., Detrain, C., Deneubourg, J. L., Clotuche, G., & Hance, T. (2010). Group
effect on fertility, survival and silk production in the web spinner tetranychus urticae (acari:
Tetranychidae) during colony foundation. *Behaviour*, *147*(9), 1169–1184.
https://doi.org/10.1163/000579510X510980

Lenth, R. V. (2023). *emmeans: Estimated Marginal Means, aka Least-Squares Means*.

Lin, T., Zhu, G., He, W., Xie, J., Li, S., Han, S., Li, S., Yang, C., Liu, Y., & Zhu, T. (2022). Soil cadmium stress
reduced host plant odor selection and oviposition preference of leaf herbivores through the
changes in leaf volatile emissions. *Science of the Total Environment*, *814*, 152728.
https://doi.org/10.1016/j.scitotenv.2021.152728

Martens, S. N., & Boyd, R. S. (1994). The ecological significance of nickel hyperaccumulation: a plant
chemical defense. *Oecologia*, *98*(3–4), 379–384. https://doi.org/10.1007/BF00324227

McLellan, C. F., & Montgomery, S. H. (2023). Towards an integrative approach to understanding
collective behaviour in caterpillars. *Philosophical Transactions of the Royal Society B: Biological*
Sciences, *378*(1874). https://doi.org/10.1098/rstb.2022.0072

Meyer, C.-L., Juraniec, M., Huguet, S., Chaves-Rodriguez, E., Salis, P., Isaure, M.-P., Goormaghtigh, E., &
Verbruggen, N. (2015). Intraspecific variability of cadmium tolerance and accumulation, and
cadmium-induced cell wall modifications in the metal hyperaccumulator Arabidopsis halleri.
Journal of Experimental Botany, *66*(11), 3215–3227. https://doi.org/10.1093/jxb/erv144

Mogren, C. L., & Trumble, J. T. (2010). The impacts of metals and metalloids on insect behavior.
Entomologia Experimentalis et Applicata, *135*(1), 1–17. https://doi.org/10.1111/j.1570-
7458.2010.00967.x

Mohiley, A., Tielbörger, K., Seifan, M., & Gruntman, M. (2020). The role of biotic interactions in
determining metal hyperaccumulation in plants. *Functional Ecology*, *34*(3), 658–668.
https://doi.org/10.1111/1365-2435.13502

Morimoto, J., Nguyen, B., Tabrizi, S. T., Ponton, F., & Taylor, P. (2018). Social and nutritional factors
shape larval aggregation, foraging, and body mass in a polyphagous fly. *Scientific Reports*, *8*(1), 1–
10. https://doi.org/10.1038/s41598-018-32930-0

Murphy, S. M., & Loewy, K. J. (2015). Trade-offs in host choice of an herbivorous insect based on
parasitism and larval performance. *Oecologia*, *179*(3), 741–751. https://doi.org/10.1007/s00442-
015-3373-8

Noret, N., Meerts, P., Vanhaelen, M., Dos Santos, A., & Escarré, J. (2007). Do metal-rich plants deter
herbivores? A field test of the defence hypothesis. *Oecologia*, *152*(1), 92–100.
https://doi.org/10.1007/s00442-006-0635-5

Oliveira, C. S., Nogara, P. A., Lima, L. S., Galiciolli, M. E. A., Souza, J. V, Aschner, M., & Rocha, J. B. T.
(2022). Toxic metals that interact with thiol groups and alteration in insect behavior. *Current
Opinion in Insect Science*, *52*, 100923. https://doi.org/https://doi.org/10.1016/j.cois.2022.100923

Pallini, A., Janssen, A., & Sabelis, M. W. (1997). Odour-mediated responses of phytophagous mites to
conspecific and heterospecific competitors. *Oecologia*, *110*(2), 179–185.
https://doi.org/10.1007/s004420050147

Pallini, A., Janssen, A., & Sabelis, M. W. (1999). Spider mites avoid plants with predators. *Experimental
and Applied Acarology*, *23*, 803–815. https://doi.org/10.1023/A

Peterson, E. K., Stark, A., Varian-Ramos, C. W., Hollocher, K. T., & Possidente, B. (2020). Exposure to
Lead (Pb2+) Eliminates Avoidance of Pb-Treated Oviposition Substrates in a Dose-Dependent
Manner in Female Vinegar Flies. *Bulletin of Environmental Contamination and Toxicology*, *104*(5),
588–594. https://doi.org/10.1007/s00128-020-02825-w

Pollard, A. J., & Baker, A. J. M. (1997). Deterrence of herbivory by zinc hyperaccumulation in Thlaspi
caerulescens (Brassicaceae). *New Phytologist*, *135*(4), 655–658. https://doi.org/10.1046/j.1469-
8137.1997.00689.x

Prokopy, R. J., & Roitberg, B. D. (2001). JOINING AND AVOIDANCE BEHAVIOR IN NONSOCIAL INSECTS.
Annual Review of Entomology, *46*(1), 631–665. https://doi.org/10.1146/annurev.ento.46.1.631

Quinn, C. F., Freeman, J. L., Reynolds, R. J. B., Cappa, J. J., Fakra, S. C., Marcus, M. A., Lindblom, S. D.,
Quinn, E. K., Bennett, L. E., & Pilon-Smits, E. A. H. (2010). Selenium hyperaccumulation offers
protection from cell disruptor herbivores. *BMC Ecology*, *10*. https://doi.org/10.1186/1472-6785-
10-19

R Core Team. (2022). *R: A language and environment for statistical computing.* R Foundation for
Statistical Computing.

Regev, S., & Cone, W. W. (1976). Analyses of pharate female twospotted spider mites for Nerolidol and
Geraniol: Evaluation for Sex Attraction of Males. *Environmental Entomology*, *5*(6), 133–138.
https://doi.org/10.1093/ee/5.1.133

Rijal, J. P., Wilson, R., & Godfrey, L. D. (2016). Characterization of spatial distribution of Tetranychus
urticae in peppermint in California and implication for improving sampling plan. *Experimental and*
Applied Acarology, *68*(2), 155–171. https://doi.org/10.1007/s10493-015-0006-1

Rodrigues, L. R., Figueiredo, A. R. T., Varela, S. A. M., Olivieri, I., & Magalhães, S. (2017). Male spider
mites use chemical cues, but not the female mating interval, to choose between mates.
Experimental and Applied Acarology, *71*(1), 1–13. https://doi.org/10.1007/s10493-016-0103-9

Rodrigues, L. R., Zélé, F., Santos, I., & Magalhães, S. (2022). No evidence for the evolution of mating
behavior in spider mites due to Wolbachia-induced cytoplasmic incompatibility. *Evolution*, *76*(3),
623–635. https://doi.org/10.1111/evo.14429

Sarmento, R. A., Lemos, F., Bleeker, P. M., Schuurink, R. C., Pallini, A., Oliveira, M. G. A., Lima, E. R., Kant,
M., Sabelis, M. W., & Janssen, A. (2011). A herbivore that manipulates plant defence. *Ecology*
Letters, *14*(3), 229–236. https://doi.org/10.1111/j.1461-0248.2010.01575.x

Sarmento, R. A., Lemos, F., Dias, C. R., Kikuchi, W. T., Rodrigues, J. C. P., Pallini, A., Sabelis, M. W., &
Janssen, A. (2011). A herbivorous mite down-regulates plant defence and produces web to
exclude competitors. *PLoS ONE*, *6*(8), 8–14. https://doi.org/10.1371/journal.pone.0023757

Shelton, D. S., Dinges, Z. M., Khemka, A., Sykes, D. J., Suriyampola, P. S., Shelton, D. E. P., Boyd, P., Kelly,
J. R., Bower, M., Amro, H., Glaholt, S. P., Latta, M. B., Perkins, H. L., Shaw, J. R., & Martins, E. P.
(2023). A pair of cadmium-exposed zebrafish affect social behavior of the un-exposed majority.
Environmental Toxicology and Pharmacology, *100*(April), 104119.
https://doi.org/10.1016/j.etap.2023.104119

Shen, N., Liu, C., Yu, H., & Qu, J. (2020). Effects of resource heterogeneity and environmental
disturbance on the growth performance and interspecific competition of wetland clonal plants.
Global Ecology and Conservation, *22*, e00914.
https://doi.org/https://doi.org/10.1016/j.gecco.2020.e00914

Tran, T. T., Janssens, L., Dinh, K. V., & Stoks, R. (2019). An adaptive transgenerational effect of warming
but not of pesticide exposure determines how a pesticide and warming interact for antipredator
behaviour. *Environmental Pollution*, *245*, 307–315. https://doi.org/10.1016/j.envpol.2018.11.022

Vickerman, D. B., Young, J. K., & Trumble, J. T. (2003). Erratum: Effect of selenium-treated alfalfa on
development, survival, feeding, and oviposition preferences of Spodoptera exigua (Lepidoptera:
Noctuidae) (Environmental Entomology 31 (953-959)). *Environmental Entomology*, *32*(1), 953–
959.

Wang, P., Lei, J.-P., Li, M.-H., & Yu, F.-H. (2012). Spatial heterogeneity in light supply affects intraspecific
competition of a stoloniferous clonal plant. *PloS One*, *7*(6), e39105.
https://doi.org/10.1371/journal.pone.0039105

Wertheim, B., Van Baalen, E. J. A., Dicke, M., & Vet, L. E. M. (2005). Pheromone-mediated aggregation
in nonsocial arthropods: An evolutionary ecological perspective. *Annual Review of Entomology*,
50(94), 321–346. https://doi.org/10.1146/annurev.ento.49.061802.123329

Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag.

Xu, L., Van Doorn, S., Hildenbrandt, H., & Etienne, R. S. (2020). Inferring the Effect of Species
Interactions on Trait Evolution. *Systematic Biology*, *0*(0), 1–17.
https://doi.org/10.1093/sysbio/syaa072

Zélé, F., Santos-Matos, G., Figueiredo, A. R. T., Eira, C., Pinto, C., Laurentino, T. G., Sucena, É., &
Magalhães, S. (2019). Spider mites escape bacterial infection by avoiding contaminated food.
Oecologia, *189*(1), 111–122. https://doi.org/10.1007/s00442-018-4316-y.