Small molecule inhibitors of transcriptional Cyclin Dependent Kinases impose HIV-1 latency, presenting "block and lock" treatment strategies

Riley M. Horvath[1], Zabrina L. Brumme[2,3], and Ivan Sadowski[1]*

[1]Department of Biochemistry and Molecular Biology
Molecular Epigenetics Group, LSI
University of British Columbia, Vancouver, B.C., Canada

[2]Faculty of Health Sciences
Simon Fraser University, Burnaby, B.C., Canada

[3]British Columbia Centre for Excellence in HIV/AIDS
Vancouver, B.C., Canada

Running Title: Senexin A inhibition of CDK8/19 robustly enforces proviral latency, a "Block-and-Lock" strategy for HIV-1 treatment

Keywords: CDK7; CDK8; CDK9; CDK19; YKL-5-124; LDC000067; Senexin A; HIV-1; Latency; Transcription; TFIIH; Mediator kinase; P-TEFb; block and lock

*Correspondence to: I. Sadowski, Dept. of Biochemistry and Molecular Biology, UBC, 2350 Health Sciences Mall, Vancouver, B.C., V6T 1Z3, CANADA; Email: ijs.ubc@gmail.com; Phone: (604) 822-4524; FAX: (604) 822-5227.

Abstract

Antiretroviral therapy is not a cure for HIV-1 as viral rebound ensues immediately following discontinuation. The block and lock therapeutic strategy seeks to enforce proviral latency and durably suppress viremic reemergence in the absence of antiretroviral therapy. Transcriptional Cyclin Dependent Kinase activity regulates LTR transcription, however, the effect and therapeutic potential of inhibiting these kinases for enforcing HIV-1 latency remains unrecognized. Using newly developed small molecule inhibitors that are highly selective for either CDK7 (YKL-5-124), CDK9 (LDC000067), or CDK8/19 (Senexin A), we found that targeting any one of these kinases prevented HIV-1 expression at concentrations that showed no toxicity. Furthermore, although CDK7 inhibition induced cell cycle arrest, inhibition of CDK9 and/or CDK8/19 did not. Of particular interest, proviral latency as induced by CDK8/19 inhibition was maintained following drug removal while CDK9 inhibitor induced latency rebounded within 24 hrs of discontinuation. Our results indicate that the Mediator complex kinases, CDK8/CDK19, are attractive block and lock targets while sole disruption of P-TEFb is unlikely to be efficacious.

Introduction

The barrier towards a cure for HIV-1 is the population of extremely long-lived CD4$^+$ T cells that harbor latent provirus. Antiretroviral therapy (ART) is capable of halting disease progression but has no effect on the latent HIV-1 reservoir and viral rebound proceeds immediately following discontinuation of treatment (Davey *et al*, 1999) (Zhang *et al*, 2000). The seemingly antithetical "shock and kill" and "block and lock" HIV-1 therapeutic strategies are currently being pursued to eliminate the current requirement of lifetime ART adherence (Sadowski & Hashemi, 2019). These aforementioned strategies rely on modulating proviral latency with "shock and kill" involving pharmacological reactivation mediated by latency reversal agents (LRAs) allowing for the subsequent destruction of infected cells through viral cytotoxicity and immunological surveillance (Kim *et al*, 2018) (Yeh & Ho, 2021). Unfortunately, formidable barriers to "shock and kill" strategies exist as it may be impossible to achieve reactivation of the majority of latent provirus (Ait-Ammar *et al*, 2020) (McMahon *et al*, 2021) (Siliciano & Siliciano, 2021) and it has become apparent that HIV-1 specific immune-mediated clearance is inefficient (Gunst *et al*, 2020) (Powell *et al*, 2020). Given the barriers to viral reactivation and clearance, the more feasible approach may be the alternative "block and lock" strategy by which durable latency is enforced to suppress viral rebound in the absence of ART (Moranguinho & Valente, 2020) (Vansant *et al*, 2020).

Cyclin Dependent Kinases (CDKs) are a family of proteins that catalyze the phosphorylation of serine/threonine residues, an activity that requires association with a cognate cyclin subunit. CDKs have historically been classified as either cell cycle associated kinases (including CDK1, CDK2, CDK4 and CDK6) or RNA polymerase II transcription associated kinases (CDK7, CDK8, CDK9, CDK12, CDK13 and CDK19). The multiprotein RNAPII

complex mediates the synthesis of mRNA from a DNA template via a series of highly

coordinated events that begins with the assembly of RNAPII and general transcription factors

(GTFs) at a gene promoter. Initiation of RNA synthesis is coupled with Serine 5 (Ser5) and

Serine 7 (Ser7) phosphorylation of the RNAPII Rpb1 subunit within the heptapeptide repeats of

the carboxy-terminal domain (CTD) as mediated by the TFIIH associated kinase, CDK7

(Buratowski, 2009) (Rodríguez-Molina *et al*, 2016). Ser5 phosphorylation (Ser5-P) frees

RNAPII from associated GTFs and the Mediator complex, facilitating promoter release

(Rodríguez-Molina *et al*, 2016) (Wong *et al*, 2014). Additionally, CTD Ser5-P stabilizes the

nascent transcript as the modification is recognized by enzymes that catalyze 7-methylguanosine

(m7GTP) capping of the mRNA 5' terminus (Ho & Shuman, 1999) (Fabrega *et al*, 2003) (Tome

et al, 2018). In contrast, the effect of Serine 7 phosphorylation (Ser7-P) is ill-defined but is

thought to prime subsequent CTD phosphorylation steps (Czudnochowski *et al*, 2012).

CDK7 initiates transcription from the core promoter, however, RNAPII may be

associated with the pause factors DRB sensitivity-inducing factor (DSIF) and negative

elongation factor (NELF) which cause stalling 20-60 nucleotides downstream of the

transcriptional start site (TSS) in an effect known as promoter proximal pausing (Core &

Adelman, 2019). Recruitment of the Cyclin T1/ CDK9 heterodimeric P-TEFb complex catalyzes

RNAPII CTD Serine 2 phosphorylation (Ser2-P), a post-translational mark that is also deposited

by CDK12 and CDK13 (Bartkowiak *et al*, 2010) (Blazek *et al*, 2011). In addition to Ser2-P,

CDK9 phosphorylates DSIF and NELF causing the former to be converted to an elongation

factor and the latter to be released, enabling processive elongation (Yamaguchi *et al*, 2013). As

RNAPII reaches the 3' end of the gene, the CTD is dephosphorylated, elongation transitions to a

slowed pace, the mRNA is cleaved, and transcription is terminated (Cortazar *et al*, 2019) (Huang

et al, 2020).

The 26-subunit Mediator complex facilitates genome-wide transcription by bridging

sequence-specific DNA-binding transcription factors (TFs) with RNAPII. A quaternary complex

of Cyclin C, CDK8 or CDK19, MED12, and MED13 forms the Mediator kinase module that

associates reversibly with Mediator. Unlike the Mediator complex, the kinase module is not a

genome-wide transcriptional co-activator. Rather, the Mediator kinase module selectively

regulates transcription by phosphorylating transcription factors. The outcome of CDK8/19 kinase

activity can be activation or repression depending on the functional effect of the modification.

For instance, CDK8/19 phosphorylation of β-catenin (Firestein *et al*, 2008), NFκB (Chen e*t al*,

2017) (Li e*t al*, 2019), SREBP-1 (Zhao *et al*, 2012), SMAD1/3 (Alarcón e*t al*, 2009), and STAT1

(Bancerek e*t al*, 2013) (Steinparzer e*t al*, 2019) enhances transcriptional activity while

phosphorylation of Notch ICD promotes its degradation leading to target gene repression (Fryer

et al, 2004) (Kuang e*t al*, 2020).

CDK7, CDK9, and CDK8/19 are transcriptional Cyclin Dependent Kinases (tCKDs) that

are involved in HIV-1 expression. Recruitment of CDK7 (TFIIH) to the HIV-1 LTR has been

described as the rate limiting step of latency reversal (Kim e*t al*, 2006) while the viral trans-

activator Tat recruits CDK9 (P-TEFb) through interaction with nascent TAR RNA resulting in

enhanced transcriptional elongation. Interestingly, the host cellular factor TRIM24 has been

observed to stimulate RNAPII HIV-1 promoter clearance, an effect that is associated with

enrichment of CDK9 and Ser2-P at the LTR (Horvath e*t al*, 2023). Moreover, the TRIM24

bromodomain inhibiter IACS-9571 and the BRD4 bromodomain inhibitor JQ1 reverse latency by

promoting association of CDK9 with the LTR (Zhu *et al*, 2012) (Li *et al*, 2013) (Horvath e*t al*,

2023). Given the role of CDK9 for HIV-1 expression, the kinase represents a potential anti-viral

target (Canduri *et al*, 2008) (de la Fuente *et al*, 2003). Inhibitors of CDK9 such as purine

derivatives (Roscovitine) and flavonoids (flavopiridol) have been shown to repress HIV-1

replication (Wang e*t al*, 2001) (Schang *et al*, 2002) (Schonhofer e*t al*, 2021). However, these

drugs lack specificity, targeting numerous CDKs (CDK2, CDK5, CDK7, etc.) in addition to

other host cell enzymes (Meijer *et al*, 1997) (Schonhofer *et al*, 2021) that results in global

inactivation of RNAPII transcription and cell cycle arrest (Blagosklonny, 2004) (Chao & Price,

2001). In addition to the aforementioned tCDKs, a role for CDK8 as a co-activator of HIV-1

transcription has recently been described with inhibitors of CDK8/19 causing repression of

proviral expression both *in vitro* and e*x vivo* (Horvath e*t al*, 2023), though the ability of

CKD8/19 inhibition to enforce robust latency has not been investigated.

Only recently have specific small molecule inhibitors of CDK7 and CDK9 been

developed. Given this technological advancement, we sought to determine the HIV-1 "block and

lock" therapeutic potential of targeting CKD7, CDK9, and/or CDK8/19. Treatment with YKL-5-

124 (CDK7 inhibitor) (Olson *et al*, 2019), LDC000067 (CDK9 inhibitor) (Albert *et al*, 2014), or

Senexin A (CDK8/19 inhibitor) (Porter *et al*, 2012) was able to prevent T cell activation

mediated proviral induction in JLat Jurkat latency models and primary CD4$^+$ peripheral blood

monocytes (PBMCs) isolated from people living with HIV-1. Although each of the small

molecules suppressed proviral expression, we observed synergistic inhibition when applied in

combination with LDC000067 (CDK9i) and Senexin A (CDK8/19i) displaying the greatest

magnitude of synergy. Although inhibition of CDK7 with YKL-5-124 was unable to affect

proviral expression without causing cell cycle defects, CDK9 and CDK8/19 inhibition was

extremely well tolerated over long durations of exposure. Given this tolerance, we were able to

establish drug induced latency through multi-day treatment of HIV-1 infected cells with

LDC000067 and/or Senexin A. Upon drug exclusion, cells treated with the CDK9 inhibitor

LDC000067 displayed immediate viral rebound. In contrast, Senexin A induced latency was

found to be robust, lasting following discontinuation. Collectively, our results suggest that the

therapeutic potential of P-TEFb disruption is severely limited while CDK8/19 inhibitors are

attractive latency promoting agents in the "block and lock" strategy to a functional HIV-1 cure.

Results

tCDK activity is required for HIV-1 induction in response to T cell activation. We sought to

examine the requirement of tCDK activity for HIV-1 expression using the small molecule

inhibitors YKL-5-124 (CDK7i), LDC000067 (CDK9i), and Senexin A (CDK8/19i) (Fig. 1*A-C*).

These recently developed chemical inhibitors display vastly improved target selectivity over

previously developed CDK inhibitors (Olson *et al*, 2019) (Albert *et al*, 2014) (Porter *et al*, 2012).

Exposure of Jurkat human T cells to increasing concentrations of inhibitor determined half lethal

concentration's (LC50) of 93 μM, 499 μM, and 200 μM for YKL-5-124 (CDK7i), LDC000067

(CDK9i), and Senexin A (CDK8/19i), respectively (Fig. 1*D*). Having determined non-toxic

concentration ranges, we investigated the effect of these drugs using the previously characterized

JLat10.6 and JLat9.2 models of HIV-1 latency. JLat Jurkat human T cell lines possess a full-

length HIV-1 provirus where GFP replaces *Nef*, and expression of the fluorescent protein is

driven by the 5' LTR (Jordan *et al*, 2003). Treatment with YKL-5-124, LDC000067, or Senexin

A displayed dose-dependent suppression of HIV-1 induction in response to PMA/ionomycin

mediated T cell activation in the JLat10.6 cell lines as measured by the percent of cells

expressing GFP and the GFP Mean Fluorescent Intensity (MFI) (Fig. 2*A-C*). Here, YKL-5-124

(CDK7i) displayed the most potent activity with a half inhibitory concentration (IC50) for

cellular GFP expression of 0.11 μM while LDC000067 and Senexin A followed with GFP

expression IC50's of 1.42 μM and 15.43 μM, respectively (Fig. 2*B*). Comparable results were

obtained when GFP MFI was assessed (Fig. 2*C*). Furthermore, a similar inhibitory effect was

observed for the JLat9.2 latency model (Fig. 2*D*), with lower concentrations of inhibitor being

required to repress proviral expression as compared to the JLat10.6 cell line (Fig. 2*E*, 2*F*). YKL-

5-124 was again the most potent inhibitor (IC50: 0.02 μM), followed by Senexin A (IC50: 0.43

μM) and LDC000067 (IC50: 0.56 μM). Notably, cellular viability of JLat10.6 and JLat9.2 cellse

were not affected by tCDK inhibition (Fig. S1).

The JLat10.6 and JLat9.2 cell lines display a distinct pattern of proviral activation in

response to T cell activation, whereby cells tend to possess provirus that are either inactive or

fully expressed (Fig. 2*A*, 2*D*, Vehicle). Treatment with LDC000067 (CDK9i) or Senexin A

(CDK8/19i) diminished the proportion of GFP+ cells (Fig. 2*B*, 2*E*), however, the bimodal

expression pattern remains apparent where there is still the tendency for those provirus that

escape latency to be fully active (Fig. 2*A*, 2*D*, LDC000067, Senexin A). In contrast, treatment

with the CDK7 inhibitor YKL-5-124 does not follow bimodal expression. Rather, a gradient of

expression is produced with many proviruses displaying activation levels that verge on the GFP-

/GFP+ border (Fig. 2*A*, 2*D*, YKL-5-124). The viral encoded trans-activator of transcription Tat

protein produces a strong positive feedback loop. CDK7 is involved in initiation of transcription,

ostensibly a Tat independent process, while CDK9 is recruited by Tat to the LTR to facilitate

release of RNAPII promoter proximal pausing. As such, it is probable that the different

expression pattern observed between YKL-5-124 inhibition of CDK7 and either LDC000067

inhibition of CDK9 or Senexin A inhibition of CDK8/19 is the result of Tat independent versus

Tat dependent transcriptional induction.

tCDK inhibitors synergistically inhibit HIV-1 expression. CDK7, CDK9, and CDK8/19

regulate different stages in RNAPII gene expression. As such, it would be expected that

combined treatment with inhibitors specific for particular tCDK's would reveal synergistic

interaction. To examine synergistic inhibition, we treated JLat10.6 cells with YKL-5-124

(CDK7i) and LDC000067 (CDK9i) (Fig. 3*A*), YKL-5-124 (CDK7i) and Senxin A (CDK8/19i)

(Fig. 3*B*), or LDC000067 (CDK9i) and Senexin A (CDK8/19i) (Fig. 3*C*), and induced proviral

expression with the addition of PMA/ionomycin. Co-treatment with any pair of tCDK inhibitor

was found to have a greater repressive effect than mono-treatment as indicated by the percent of

GFP positive cells following reactivation (Fig. 3*A-C*). Additionally, cellular viability was not

affected by any of the applied treatments (Fig. S2). We confirmed synergistic inhibition by

performing Bliss independence modelling (Laird *et al*, 2014) (Liu *et al*, 2018) and found that the

combination of LDC000067 (CDK9i) and Senexin A (CDK8/19) displayed the strongest

interaction (Fig. 3*D-F*). Of note, the apparent synergy dips at higher concentrations of chemical

inhibitor as a result of viral expression approaching zero (Fig. 3*D-F*). These observations are

consistent with the tCDK inhibitors being reported as selective for either CDK7, CDK9, or

CDK8/19 and indicate that these kinases function through independent mechanisms.

Inhibition of tCDK activity antagonizes proviral reactivation in response to mechanistically

diverse latency reversal agents. Having observed that tCDK inhibition suppresses reactivation of

latent provirus in response to T cell activation, we next examined the effect on a variety of

mechanistically diverse latency reversal agents (LRAs). As mentioned above, co-treatment with

the phorbol ester PMA and the ionophore ionomycin triggers T cell activation by stimulating

Ras-MAPK, PKC-NFκB, and calcineurin-NFAT. Ingenol 3-angelate (PEP005) activates PKC-

NFκB, partially mimicking T cell activation (Kedei *et al*, 2004). JQ1 and IACS-9571 are

bromodomain inhibitors that reverse latency as mediated by their respective targets, BRD4 (Li *et*

al, 2013) and TRIM24 (Horvath *et al*, 2023). Finally, suberanilohydroxamic acid (SAHA) is a

well-studied histone deacetylase inhibitor (HDACi) known to reactivate HIV-1 expression *in*

vitro and *in vivo* (Contreras *et al*, 2009) (Desimio *et al*, 2017). Of the LRAs applied, the

bromodomain inhibitors JQ1 and IACS-9571 had the smallest effect on proviral reactivation in

the JLat10.6 latency model (Fig. 4*A*, 4*B*). The NFκB agonist PEP005 and the HDACi SAHA

produced intermediate reactivation while co-treatment with either PEP005/IACS-9571 or

PMA/ionomycin exhibited the greatest levels of HIV-1 expression (Fig. 4*A*, 4*B*).

We applied 100 nM YKL-5-124 (CDK7i), 1 μM LDC000067 (CDK9i), or 10 μM

Senexin A (CDK8/19i), concentrations near the inhibitor's LC50 for PMA/ionomycin activation,

in combination with the LRAs discussed above. CDK7 inhibition by YKL-5-124 limited proviral

reactivation as measured by the population of GFP+ cells in response to every LRA other than

JQ1 where oddly, we observe an increased proportion of GFP expressing cells (Fig. 4*A*).

Interestingly, although CDK7 inhibition with concomitant JQ1 treatment produced a greater

proportion of GFP+ cells (Fig. 4*A*), the GFP MFI was decreased as compared to the vehicle

control (Fig. 4*B*) indicating that more cells expressed GFP but at attenuated levels upon

introduction of YKL-5-124. Treatment with LDC000067 (CDK9i) or Senexin A (CDK8/19i)

was found to stunt HIV-1 transcriptional induction in response to all of the tested LRAs (Fig. 4*A*,

4*B*). In agreement with previous reports (Horvath *et al*, 2023), we observed a vigorous

synergistic interaction between PEP005 activation of NFκB and IACS-9571 inhibition of the

TRIM24 bromodomain (Fig. S3). Moreover, tCDK inhibition was least effective at repressing

the PEP005/IACS-9571 reactivation protocol, further demonstrating its latency reversal potential

(Fig. 4*A*, 4*B*). Finally, although these concentrations of tCDK inhibitors equally suppressed the

emergence of GFP positive cells in response to PMA/ionomycin T cell activation (Fig. 4*A*,

PMA/Ionomycin), we found certain inhibitors to be more effective at combatting particular

LRAs. For instance, Senexin A (CDK8/19i) was most effective at suppressing SAHA mediated

activation (Fig. 4*A*, SAHA) while LDC000067 (CDK9i) was most potent at inhibiting expression

in response to PEP005 (Fig. 4*A*). Taken together, these results show that tCDK inhibition

suppresses proviral activation in response to mechanistically diverse LRAs while suggesting that

the kinases have divergent functions for HIV-1 transcription.

Tat dependence of tCDKs for activation of HIV-1 transcription. Next, we examined the Tat

dependence of CDK7, CDK9, and CDK8/19 for activation of HIV-1 transcription. The Jurkat

derived JLatA72 cell line possesses an integrated HIV-1 LTR that drives expression of GFP but

does not encode Tat. Untreated JLatA72 cells are ~30% GFP positive and this proportion

increases to ~60% upon PMA/ionomycin stimulation (Fig. 5*A*, 5*B*, Untreated, Vehicle). To

examine the Tat dependence of the tCDKs, we treated JLatA72 with a kinase inhibitor and

subsequently induced T cell activation with PMA/ionomycin. Surprisingly, we found that YKL-

5-124 inhibition of CDK7 caused an increase in the proportion of GFP positive cells (Fig. 5*A*,

5*B*) while also lowering overall proviral expression (Fig. 5*C*), an effect that is strikingly similar

to co-treatment with JQ1 (Fig. 4). The CDK9 inhibitor LDC000067 had no effect on the

proportion of GFP positive cells (Fig. 5*A*, 5*B*) and only slightly decreased overall GFP

expression (Fig. 5*C*), consistent with the role of Tat for recruitment of CDK9 to the HIV-1 LTR.

Finally, inhibition of CDK8/19 with Senexin A decreased the emergence of GFP (Fig. 5*A*, 5*B*)

while also suppressing overall GFP expression (Fig. 5*C*), indicating Tat independent regulation.

We confirmed Tat dependence by additional examination of the CEM derived GXR-5 cell line.

As with the JLatA72 cell line, The GXR-5 cells possess a chromosomally integrated HIV-1 LTR

that drives GFP expression but does not express Tat. Unlike JLatA72 cells, GXR-5 cells are

almost entirely GFP positive (Fig. S4*A*, Fig. S4*B*) though an increase in GFP expression

following PMA/ionomycin treatment is observable by assessing GFP Mean Fluorescent Intensity

(MFI) (Fig. S4*C*). Similar to the JLatA72 cell line, GXR-5 cells display severely attenuated GFP

induction upon treatment with YKL-5-124 (CDK7i) or Senexin A (CDK/19i) (Fig. S4*C*). In

contrast, LDC000067 (CDK9) caused small but significant inhibition of GFP expression (Fig.

S4*C*).

Effect of tCDK inhibition on cell growth. Above, we examined the effects of acute tCDK

inhibition on proviral expression. Given the stochastic nature of HIV-1 reactivation, a "block and

lock" treatment strategy will likely necessitate longer periods of exposure to generate latency

that persists subsequent to drug withdrawal. To further examine the cellular impacts of tCDK

inhibition, we assessed the growth and viability of Jurkat human T cells exposed to a range of

drug concentrations for 4-days. We found that inhibition of CDK7 with YKL-5-124 caused cell

cycle arrest (Fig. 6*A*) at concentrations shown to inhibit HIV-1 expression (Fig. 2), a result that is

consistent with CDK7 being the CDK-activating kinase (CAK) which phosphorylates and

activates CDKs that have essential roles for cell-cycle progression (Fisher, 2005). Moreover,

cellular toxicity begins to become apparent following 3- and 4-days of CDK7 inhibition (Fig.

6*B*). In contrast of CDK7 inhibition, inhibition of CDK9 with LDC000067 had no effect on cell

division (Fig. 6*C*) or viability (Fig. 6*D*) at concentrations effective at limiting proviral expression

(Fig. 2). Likewise, Senexin A inhibition of CDK8/19 had minimal impact on cell growth (Fig.

6*E*) and did not affect viability (Fig. 6*F*) within a range of concentrations that suppressed HIV-1

reactivation (Fig. 2). As CDK7 inhibition causes cell-cycle arrest at low concentrations, the

therapeutic potential of YKL-5-124 and other CDK7 inhibitors will likely be limited. However,

our results find that inhibition of CDK9 with LDC000067 or inhibition of CDK8/19 with

Senexin is well tolerated over prolonged application within an optimal concentration range for

HIV-1 repression.

Role of tCDKs for the establishment of immediate latency. Upon infection, nearly 50% of

provirus are transcriptionally inactive, a phenomenon known as immediate latency (Dahabieh *et*

al, 2013) (Dahabieh *et al*, 2014) (Battivelli e*t al*, 2018). Having observed that the tCDKs are

required for reactivation of HIV-1 in response to various LRAs, we examined the effect of kinase

inhibition on the occurrence of immediate latency using the Red-Green-HIV-1 (RGH) dual-

reporter virus (Fig. 7*A*) (Dahabieh *et al*, 2013) (Dahabieh *et al*, 2014). RGH is a full-length HIV-

1 construct that is rendered replication incompetent by mutation to *Env* and whereby eGFP

expression is driven by the 5' LTR and a constitutive PGK-mCherry cassette is inserted in place

of *Nef*, allowing for differentiation between latent and productive proviral infection (Fig. 7*A*). To

assess tCDK activity on immediate latency, we pre-treated Jurkat human $CD4^+$ T cells with a

kinase inhibitor and subsequently infected with RGH at a low multiplicity of infection (MOI).

One day following infection, we either washed (removed) or maintained the drug inhibitor,

assessing proviral latency 4-days post-infection by flow cytometry (Fig. 7*B*) (Fig. 7*C*). At the

point of infection, inhibition of either CDK7 with YKL-5-124 or CDK8/19 with Senexin A was

able to reduce the proportion of productive infections (Fig. 7*D*, Washed). Interestingly, inhibition

of CDK9 with LDC000067 at the point of infection had no impact on immediate latency (Fig.

7*D*, Washed), suggesting that CDK9 and Tat are not involved in viral transcription early upon

infection. In agreement with tCDK inhibition preventing reactivation of latent provirus (Fig. 2,

Fig. 4), latency was enforced by all of the small molecule inhibitors when they were maintained

for the 4 days (Fig. 7*D*, Maintained). Surprisingly, we found that YKL-5-124 inhibition of CDK7

caused the rate of infection to roughly double indicating that loss of CDK7 activity renders T

cells susceptible to infection (Fig. 7*E*). Notably, although CDK7 inhibition causes an increase in

HIV-1 infection (Fig 7*E*), integrants were enriched for latent phenotype (Fig. 7*C*, compare

Untreated to YKL-5-124). These results indicate that CDK9 and CDK8/19 are attractive targets

to suppress proviral expression.

Disruption of CDK8/19 but not CDK9 enforces robust latency. An optimal "block and lock"

therapy would involve generating proviral latency that is maintained subsequent to the removal

of the latency promoting agent (LPA). To examine the capability of tCDK inhibitors to create

robust and lasting latency, we applied LDC000067 (CDK9i) and/or Senexin A (CDK8/19i) to the

previously described mdHIV #77 cell line (Hashemi *et al*, 2016). mdHIV #77 is a Jurkat Tat

derived clonal line that contains a mini-dual reporter HIV-1 provirus (Fig. 8*A*) that is integrated

into an intron of the *PACS1* gene in the (+) orientation (Hashemi *et al*, 2016). Although every

cell of this clonal line contains the mini-virus integrated in the same location, stochastic bursts of

transcription produce a slightly fluctuating population of ~3% productive infections as indicated

by the presence of LTR expressed dsRed and eGFP transcribed by an internal EF1α promoter

(Hashemi *et al*, 2016). We applied the tCDKi to mdHIV #77 cells for 14 days after which the

drug was washed (removed from the culture media) and cells were monitored for 10 more days

by flow cytometry (Fig. 8*B*, 8*C*). YKL-5-124 (CDK7i) was not assessed given the toxicity of this

drug when applied for several days (Fig. 6*A*, 6*B*).

Consistent with our findings that tCDK activity is required for reactivation of latent

provirus (Fig. 2) and stimulates transcription early upon infection (Fig. 7), basal HIV-1

expression was practically outright suppressed by LDC000067 (CDK9i) and/or Senexin A

(CDK8/19i) following 4 days of treatment (Fig. 8*C*, 8*D*). To great surprise, removal of the

CDK9 inhibitor LDC000067 corresponded with immediate rebound of proviral expression as the

population of dsRed+/eGFP+ cells nearly matched that of uninhibited cells within 24 hrs of drug

withdrawal (Fig. 8*D*, compare day 14 to 15). Very importantly however, mdHIV #77 cells that

had been treated with Senexin A (CDK8/19i) alone or in combination with LDC000067 (CDK9i)

displayed maintenance of latency, with the tCDKi treated cells displaying reduced HIV-1

expression for more than 10 days following drug removal (Fig. 8*D*, Drug Washed). Of note,

none of the inhibitor treatments affected cellular viability (Fig. 8*E*). Additionally, we examined

the Jurkat Tat mdHIV #110 clonal line (Fig. S6). Akin to mdHIV #77 cells, the mdHIV #110 cell

line possesses a chromosomally integrated mini-dual reporter provirus (Fig. S6*A*), however, this

virus is located in *BTBD10* in the (-) orientation and has a proportion of ~25% productive cells

(Hashemi *et al*, 2016). Treatment of mdHIV #110 resulted in similar observations as either

inhibitor used alone or in combination reduced the population of productive provirus (Fig. S6*D*,

Drug Treated). Once again, latency caused by LDC000067 (CDK9i) was not robust, with viral

expression rebounding within 24 hrs of withdrawal (Fig. S6*D*, compare day 14 to 15). In

contrast, Senexin A (CDK8/19i) treated cells maintained lower viral expression days following

removal (Fig. S6*D*, Drug Washed). The viability of mdHIV #110 cells was unaffected by the

tCDKi treatments (Fig. S6*E*).

Above, we examined the capability of CDK9 and/or CDK8/19 inhibitors to generate

robust, lasting HIV-1 latency (Fig. 8, Fig. S6). However, these cell lines were generated from

Jurkat T cells that constitutively express Tat and with a mini-HIV-1 reporter virus that lacks

expression of viral proteins (Hashemi *et al*, 2016). As such, we wanted to confirm our findings

using a full-length HIV-1 provirus with auxiliary protein expression, including Tat, coupled to 5'

LTR transcriptional activity. To this end, we generated a clonal RGH Jurkat T cell line (Fig. 9*A*,

RGH#20) and treated it for 7 days with LDC000067 (CDK9i) and/or Senexin A (CDK8/19i)

with analysis continuing for 10 days following drug withdrawal (Fig. 9*B*). Near complete viral

latency was established following 4 days of inhibitor treatment (Fig. 9*C*, 9*D*, Drug Treated).

Again, removal of the CDK9 inhibitor LDC000067 was followed by viral rebound while cells

retained decreased HIV-1 transcription for more than a week following the removal of the

CDK8/19 inhibitor Senexin A (Fig. 9*D*, Drug Washed). Furthermore, no change in RGH #20 cell

viability was observed for any of the administered treatments (Fig. 9*E*). Collectively, these

results indicate that inhibitors of CDK9 will be incapable of producing long-lasting robust

latency and as such, are not ideal "block and lock" LPA candidates. In contrast, inhibition of the

Mediator kinases CDK8/CDK19 generates proviral latency that is maintained following drug

withdrawal making them attractive targets for "block and lock" therapeutic approaches.

tCDK inhibitors suppress proviral expression in people living with HIV-1 ex vivo. We sought

to confirm the requirement of tCDK activity for proviral expression by assessing the effect of

kinase inhibition on primary CD4$^+$ PBMCs isolated from individuals living with HIV-1 (Table

1). In all three of the individuals examined, we observed increased abundance in HIV-1 mRNA

in response to PMA/ionomycin stimulation (Fig. 10, compare Unstimulated to Vehicle). In

agreement with the results presented above, treatment with YKL-5-124 (CDK7i), LDC000067

(CDK9i), or Senexin A (CDK8/19i) consistently resulted in a decrease in proviral induction (Fig.

10). Interestingly, the combination of LDC000067 and Senexin A produced the most robust

silencing of viral expression (Fig. 10, LDC000067/Senexin A), with Patient A displaying even

lower quantities of HIV-1 mRNA than when left unstimulated (Fig. 10*A*). Furthermore, for

355   Patient C we observe cooperative HIV-1 inhibition of combined LDC000067 and Senexin A

356   treatment (Fig. 10*C*). Taken together, these results indicate that inhibition of the tCDKs can

357   suppress reactivation of HIV-1 in patients *ex vivo*.

Discussion

Current therapeutic strategies to eliminate the life-long requirement of ART for people

living with HIV-1 are aimed at modulating proviral latency. The "shock and kill" strategy of

latency reversal has been the subject of intense investigation and although a number of LRAs

have been identified, none have succeeded in reducing the pool of latently infected $CD4^+$ T cells

(Sadowski & Hashemi, 2019). The antithetical "block and lock" approach has been proposed to

be a more realistic strategy to functionally eliminate HIV-1. The aforementioned therapeutic

strategy relies on LPA mediated suppression of 5' LTR transcription which may lead to

sustained epigenetic silencing of integrated provirus (Moranguinho & Valente, 2020) (Vansant *et*

al, 2020). Although several LPAs such as didehydro-cortistatin A (dCA) (Mousseau *et al*, 2015)

(Kessing *et al*, 2017) and spironolactone (Mori *et al*, 2021) have displayed promising results, no

human trial with any latency enforcing compound has been performed. In the present study, we

assessed the transcriptional Cyclin Dependent Kinases (tCDKs) CDK7, CDK9, and CDK8/19 as

potential targets for the robust enforcement of proviral latency in various cell line models and

primary $CD4^+$ T cells from people living with HIV-1.

Recruitment of the general transcription factor TFIIH to the LTR is a rate limiting step

for reactivation of latent HIV-1 (Kim *et al*, 2006). TFIIH is a complex composed of 10 proteins

consisting of a core component (XPB, XPD, p62, p52, p44, p34, p8) and a CDK-activating

kinase (CAK) module (CDK7, cyclin H, MAT1) (Yan *et al*, 2019). In agreement with TFIIH

recruitment being required for viral reactivation, we found that treatment with the selective

CDK7 inhibitor YKL-5-124 prevented transcriptional induction of HIV-1 in response to T cell

activation (Fig. 2). Furthermore, YKL-5-124 was found to limit reactivation in response to

several diverse LRAs including the TRIM24 bromodomain inhibitor IACS-9571 and the HDACi

SAHA (Fig. 4). Surprisingly, CDK7 inhibition with concomitant JQ1 mediated induction

resulted in an increased proportion of cells possessing productive HIV-1 (Fig. 4*A*), although viral

expression was severely dampened (Fig. 4*B*). Interestingly, we observed a similar expression

pattern when we assessed the reactivation of YKL-5-124 treated JLatA72 cells (Fig. 5). The

JLatA72 cell line possesses an integrated HIV-1 reporter that lacks Tat and as such, it would be

reasonable to attribute this expression pattern to a Tat dependence. However, inhibition of CDK9

does not affect the reactivation of JLatA72 cells (-Tat) (Fig. 5*B*) but does impede reactivation of

JLat10.6 cells in response to JQ1 (Fig. 4). As mentioned above, TFIIH is a multi-subunit

complex with both helicase and kinase activities that facilitate promoter opening and

phosphorylate the CTD of RNAPII at Ser-5. Our peculiar finding that CDK7 inhibition increases

the pool of productive infections in certain conditions likely reflects the diverse functions of

TFIIH for transcription circuit procession.

Additionally, we found that the CDK7 inhibitor YKL-5-124 was unable to prevent HIV-1

expression at concentrations that did not cause major disturbances in cell cycle progression (Fig.

6*A*, 6*B*). Perhaps even more intriguing, we observed that CDK7 inhibition was associated with

increased susceptibility of T cells for HIV-1 infection (Fig. 7). Taken together, our assessment

suggests that CDK7 inhibitors such as YKL-5-124 are unlikely to be appropriate for therapeutic

use. Spironolactone causes the proteasomal degradation of the XBP subunit of TFIIH and has

been shown to enforce HIV-1 latency (Mori *et al*, 2021). As such, spironolactone has been

suggested as an LPA suitable for "block and lock" although viral rebound occurs immediately

following drug withdrawal (Mori *et al*, 2021). Interestingly, although spironolactone severely

restricts cellular XPB protein, it also leads to a reduction in CDK7 (Mori *et al*, 2021). Given our

findings, inhibitors of TFIIH should be approached cautiously with regards to application in

clinical settings.

P-TEFb, a heterodimer of Cyclin T1 and CDK9, is recruited to the HIV-1 LTR TAR

element through interaction with Tat and facilitates escape of promoter proximal paused

RNAPII, resulting in a strong positive feedback loop (Karn, 2011). We sought to interrupt the

CDK9-Tat feedback loop and enforce latency by applying the selective inhibitor of CDK9 kinase

activity, LDC000067 (Albert *et al*, 2014). Here, we found that LDC000067 suppressed proviral

induction in various cell models (Fig. 2, Fig. 4) and in people living with HIV-1 *ex vivo* (Fig.

10). We also confirmed the Tat dependence of CDK9 for HIV-1 expression as LDC000067 had

no appreciable effect in the absence of the viral trans-activator of transcription (Fig. 5, Fig. S4).

Given the predominate role of CDK9 for HIV-1 transcription, the enzyme has been proposed as a

potential therapeutic target (Canduri *et al*, 2008) (Ahlenstiel *et al*, 2020) and to this point, the

CDK9 inhibitors Roscovitine, flavopiridol, and seliciclib have been shown to supress HIV-1

replication (Wang e*t al*, 2001) (Schang *et al*, 2002) (Biglione *et al*, 2007) (Schonhofer e*t al*,

2021). However, the aforementioned compounds are properly described as pan-CDK inhibitors

as they are not selective and cause global inactivation of RNAPII transcription and cell cycle

arrest (Blagosklonny, 2004) (Chao & Price, 2001). Of slight surprise, we found that the selective

inhibition of CDK9 with LDC000067 had no effect on cell division and no apparent toxicity at

concentrations effective at restricting proviral expression (Fig. 6*C*, 6*D*).

As LDC000067 administration was well tolerated, we were able to assess the capability

of CDK9 inhibition to generate robust, lasting latency. In all T cell models examined,

LDC000067 treatment generated proviral latency without affecting cell growth (Fig. 8, Fig. 9,

Fig. S6, see Drug Treated). Disappointingly though, removal of the CDK9 inhibitor from media

by washing corresponded with immediate HIV-1 rebound (Fig. 8, Fig. 9, Fig. S6, see Drug

Washed). There is great interest in disrupting the Tat-CDK9 positive feedback loop in order to

initiate "block and lock", and several Tat targeting mechanisms have been identified including

degradation of Tat by the long noncoding RNA NRON (Li *et al*, 2016) and introduction of the

Tat mutant, Nullbasic (Jin *et al*, 2016) (Jin *et al*, 2019). Given that the predominate role of Tat

for HIV-1 replication is the recruitment of P-TEFb (CDK9/Cyclin T1) to the 5' LTR promoter

and that we have observed that CDK9 inhibition is unable to generate durable latency, future

studies should closely assess the ability of Tat inhibition to create long-lasting latency.

Recently, work from our lab has identified that CDK8 is a coactivator of HIV-1

expression and small molecule inhibition of CDK8/19 kinase activity restricts proviral

reactivation (Horvath *et al*, 2023). Similarly, here we show that Senexin A inhibition of

CDK8/19 suppresses viral induction in response to a diverse set of LRAs (Fig. 2, Fig. 4). The

function of CDK8/19 is not dependent on Tat (Fig 5, Fig. S4), consistent with previous findings

that CDK8/19 inhibition limits reactivation of ACH2 and U1 proviral latency cell models

(Horvath *et al*, 2023). Additionally, we display a synergistic interaction between the CDK8/19

inhibitor Senexin A and inhibitors of either CDK7 or CDK9, suggesting distinct mechanism(s) of

action. Intriguingly, CDK8/19 inhibition was effective beyond temporally acute treatment.

Application of Senexin A at the point of infection alone caused a significant decrease in the

proportion of transcriptionally active provirus in the days that followed (Fig. 7). Furthermore,

inhibition of CDK8/19 with Senexin A was able to generate latency that was maintained for days

following removal of the drug (Fig. 8, Fig. 9, Fig. S6). Collectively, our results indicate that the

Mediator complex kinases, CDK8 and CDK19, are attractive targets for the development of a

"block and lock" therapeutic strategy of sustained latency.

465  In the present study, we have evaluated the therapeutic potential of targeting CDK7,

466 CDK9, and/or CDK8/19 for a functional HIV-1 cure. Inhibition of CDK7 presented several

467 deficiencies including cellular toxicity and susceptibility to further HIV-1 infection. Although

468 the CDK9 inhibitor LDC000067 was well tolerated and suppressed proviral expression, durable

469 latency was not manifested. On the other hand, the CDK8/19 inhibitor Senexin A restricted HIV-

470 1 expression at concentrations that did not affect cellular growth while also showing the ability

471 to establish latency that was maintained following drug withdrawal. These results indicate that

472 continued treatment with LDC000067 (CDK9i) and/or Senexin A (CDK8/19i) could be used

473 therapeutically to deplete low-level viral replication in patients on ART. However, inhibition of

474 CDK8/19 has the possibility of maintaining viral suppression following discontinuation of

475 treatment.

Materials and Methods

Cell culture, virus culture, and lentiviral transduction

Jurkat E6-1, Jurkat Tat mHIV-Luciferase, JLat10.6, JLat9.2, JLatA72, and GXR-5 CEM

cells were cultured in Roswell Park Memorial Institute 1640 (RPMI-1640) medium

supplemented with 10% FBS, penicillin [100 units/ml], streptomycin [100 g/ml], and L-

glutamine [2 mM]. HEK293T were cultured in Dulbecco's modified Eagle's (DMEM) medium

supplemented with 10% fetal bovine serum (FBS), penicillin [100 units/ml], streptomycin [100

g/ml], and L-glutamine [2 mM]). All cell lines were incubated in a humidified 37°C and 5% CO_2

atmosphere.

Peripheral Blood Mononuclear Cells (PBMC) from participants with HIV-1 on ART

were isolated from whole blood by density gradient centrifugation using Lymphoprep[TM] and

SepMate[TM] tubes (StemCell Technologies), and cryopreserved. Upon thawing, PBMCs were

cultured in RPMI supplemented with 10% FBS, penicillin [100 units/ml], streptomycin [100

g/ml], and L-glutamine [2 mM]. Samples from participants were collected with written informed

consent under a protocol jointly approved by the research ethics boards at Providence Health

Care/UBC and Simon Fraser University (certificate H16-02474).

Vesicular stomatitis virus G (VSV-G) pseudotyped viral stocks were produced by co-

transfecting HEK293T cells with a combination of viral molecular clone, psPAX, and VSV-G at

a ratio of 8 μg : 4 μg : 2 μg. Transfections were performed with polyethylenimine (PEI) at a ratio

of 6:1 (PEI:DNA) in Gibco Opti-MEM[tm]. Lentiviral infections were performed by plating $1x10^6$

cells in 24-well plates with media containing 8 μg/mL polybrene and the amount of viral

supernatant to give the desired multiplicity of infection (M.O.I.) as indicated. Plates were

subsequently spinoculated for 1.5 hrs at 1500 rpm.

RT-PCR

RNA was extracted from cells using the RNeasy Kit (Qiagen) and analyzed with the

Quant Studio 3 Real-Time PCR system (Applied Biosystems) using *Power* SYBR® Green

RNA-to-CT™ 1-Step Kit (Thermo Fisher) as per the manufacturer's instructions. RT-PCR data

was normalized to GAPDH expression using the ΔΔCt method as previously described (Livak &

Schmittgen, 2001). Cycling parameters were as follows: 48 ^{o}C, 30 min, 1x; 95 ^{o}C, 10 min, 1x; 95

^{o}C, 15 sec, 60 ^{o}C, 1 min, 60x. Primers were as follows: HIV-1 mRNA (multiply spliced Tat-Rev

transcripts), Fwd 5' CTTAGGCATCTCCTATGGCAGGA, Rev 5'

GGATCTGTCTCTGTCTCTCTCTCCACC; GAPDH, Fwd 5'

TGCACCACCAACTGCTTAGC, Rev 5' GGCATGGACTGTGGTCATGAG.

Flow cytometry

Cells were treated as indicated in the figure legends. Human T cell lineages were

suspended in PBS and a BD Biosciences LSRII-561 system was used for flow cytometry with

threshold forward scatter (FSC) and side scatter (SSC) parameters being set so that a

homogenous population of live cells was counted. FlowJo software (TreeStar) was used to

analyze data and determine the indicated Mean Fluorescence Intensity (MFI).

Quantitative analysis of tCDK inhibitor latency promoting agent interaction

Bliss independence modeling was used as previously described (Horvath *et al*, 2023) to

statistically assess the inhibitory activity of combinatorial tCDK inhibitor treatment on HIV-1

reactivation. The equation $Fa_{xy, P} = Fa_x + Fa_y - (Fa_x)(Fa_y)$ defines the Bliss independence

model, in which $Fa_{xy, P}$ is the predicted fraction affected by a combination of drug x and drug y

that is derived from the experimentally observed fraction affected by drug x (Fa_x) and drug y

(Fa_y) individually. Comparison of the predicted combinatorial affect ($Fa_{xy, P}$) with the

experimentally observed impact ($Fa_{xy,O}$) was then performed: $\Delta Fa_{xy} = Fa_{xy,O} - Fa_{xy,P}$. If

ΔFa_{xy} is greater than 0, the combination of drugs x and y exceed that of the predicted affect

indicating that the drugs display synergistic interaction. If $\Delta Fa_{xy} = 0$, the drug combination

follows the Bliss model for independent action. If ΔFa_{xy} is less than 0, the drug interaction is

antagonistic as the observed effect of the drug combination is less than predicted.

Statistics and reproducibility

All replicates are independent biological replicates and are presented as mean values

with $\pm$ standard deviation shown by error bars. The number of times that an experiment was

performed is indicated in the figure legends. P-values were determined by performing unpaired

samples *t*-test with the use of GraphPad Prism 9.0.0. Statistical significance is indicated at

*$*P < 0.05$, **$P < 0.01$, or ***$P < 0.001$, with n.s. denoting non-significant $P \geq 0.05$.

Data availability

All data supporting the findings of this study are available within the article or from the

corresponding author upon reasonable request (I. Sadowski, ijs.ubc@gmail.com).

Acknowledgments

We thank the laboratory staff at the BC Centre for Excellence in HIV/AIDS for processing PBMCs from study participants. This research was supported by program project grant F16-01210, from the Canadian Institutes of Health Research (CIHR), and F19-05392 Discovery Grant from the Natural Sciences and Engineering Research Council of Canada (NSERC) (to I.S.). PBMC collection from participants with HIV-1 was supported by CIHR project grant PJT-159625 (to ZLB). ZLB is supported by a Michael Smith Health Research BC Scholar Award.

Author Contributions

Experimental design was conceived by Horvath R. M who also performed all experiments. Brumme Z.L. provided PBMC samples from individuals on ART. Horvath R. M. and Sadowski I. wrote the manuscript.

Conflicts of Interest

The authors declare no conflicts of interest.

References

1. Ahlenstiel, C. L., Symonds, G., Kent, S. J. & Kelleher, A. D. Block and Lock HIV Cure Strategies to Control the Latent Reservoir. *Front Cell Infect Microbiol* **10,** 424 (2020).

2. Ait-Ammar, A., Kula, A., Darcis, G., Verdikt, R., De Wit, S., Gautier, V., Mallon, P. W. G., Marcello, A., Rohr, O. & Van Lint, C. Current Status of Latency Reversing Agents Facing the Heterogeneity of HIV-1 Cellular and Tissue Reservoirs. *Front Microbiol* **10,** 3060 (2019).

3. Alarcón, C., Zaromytidou, A.-I., Xi, Q., Gao, S., Yu, J., Fujisawa, S., Barlas, A., Miller, A. N., Manova-Todorova, K., Macias, M. J., Sapkota, G., Pan, D. & Massagué, J. Nuclear CDKs drive Smad transcriptional activation and turnover in BMP and TGF-beta pathways. *Cell* **139,** 757–769 (2009).

4. Albert, T. K., Rigault, C., Eickhoff, J., Baumgart, K., Antrecht, C., Klebl, B., Mittler, G. & Meisterernst, M. Characterization of molecular and cellular functions of the cyclin-dependent kinase CDK9 using a novel specific inhibitor. *Br J Pharmacol* **171,** 55–68 (2014).

5. Bancerek, J., Poss, Z. C., Steinparzer, I., Sedlyarov, V., Pfaffenwimmer, T., Mikulic, I., Dölken, L., Strobl, B., Müller, M., Taatjes, D. J. & Kovarik, P. CDK8 kinase phosphorylates transcription factor STAT1 to selectively regulate the interferon response. *Immunity* **38,** 250–262 (2013).

6.

Bartkowiak, B., Liu, P., Phatnani, H. P., Fuda, N. J., Cooper, J. J., Price, D. H., Adelman, K., Lis, J. T. & Greenleaf, A. L. CDK12 is a transcription elongation-associated CTD kinase, the metazoan ortholog of yeast Ctk1. *Genes Dev* **24,** 2303–2316 (2010).

7. Battivelli, E., Dahabieh, M. S., Abdel-Mohsen, M., Svensson, J. P., Tojal Da Silva, I., Cohn, L. B., Gramatica, A., Deeks, S., Greene, W. C., Pillai, S. K. & Verdin, E. Distinct chromatin functional states correlate with HIV latency reactivation in infected primary CD4+ T cells. *Elife* **7,** e34655 (2018).

8. Biglione, S., Byers, S. A., Price, J. P., Nguyen, V. T., Bensaude, O., Price, D. H. & Maury, W. Inhibition of HIV-1 replication by P-TEFb inhibitors DRB, seliciclib and flavopiridol correlates with release of free P-TEFb from the large, inactive form of the complex. *Retrovirology* **4,** 47 (2007).

9. Blagosklonny, M. V. Flavopiridol, an inhibitor of transcription: implications, problems and solutions. *Cell Cycle* **3,** 1537–1542 (2004).

10. Blazek, D., Kohoutek, J., Bartholomeeusen, K., Johansen, E., Hulinkova, P., Luo, Z., Cimermancic, P., Ule, J. & Peterlin, B. M. The Cyclin K/Cdk12 complex maintains genomic stability via regulation of expression of DNA damage response genes. *Genes Dev* **25,** 2158–2172 (2011).

11.

Buratowski, S. Progression through the RNA polymerase II CTD cycle. *Mol Cell* **36,** 541–546 (2009).

12. Canduri, F., Perez, P. C., Caceres, R. A. & de Azevedo, W. F. CDK9 a potential target for drug development. *Med Chem* **4,** 210–218 (2008).

13. Chao, S. H. & Price, D. H. Flavopiridol inactivates P-TEFb and blocks most RNA polymerase II transcription in vivo. *J Biol Chem* **276,** 31793–31799 (2001).

14. Chen, M., Liang, J., Ji, H., Yang, Z., Altilia, S., Hu, B., Schronce, A., McDermott, M. S. J., Schools, G. P., Lim, C.-U., Oliver, D., Shtutman, M. S., Lu, T., Stark, G. R., Porter, D. C., Broude, E. V. & Roninson, I. B. CDK8/19 Mediator kinases potentiate induction of transcription by NFκB. *Proc Natl Acad Sci U S A* **114,** 10208–10213 (2017).

15. Contreras, X., Schweneker, M., Chen, C.-S., McCune, J. M., Deeks, S. G., Martin, J. & Peterlin, B. M. Suberoylanilide hydroxamic acid reactivates HIV from latently infected cells. *J Biol Chem* **284,** 6782–6789 (2009).

16. Core, L. & Adelman, K. Promoter-proximal pausing of RNA polymerase II: a nexus of gene regulation. *Genes Dev* **33,** 960–982 (2019).

17. Cortazar, M. A., Sheridan, R. M., Erickson, B., Fong, N., Glover-Cutter, K., Brannan, K. & Bentley, D. L. Control of RNA Pol II Speed by PNUTS-PP1 and Spt5 Dephosphorylation

Facilitates Termination by a 'Sitting Duck Torpedo' Mechanism. *Mol Cell* **76,** 896-908.e4

(2019).

18.

Czudnochowski, N., Bösken, C. A. & Geyer, M. Serine-7 but not serine-5 phosphorylation

primes RNA polymerase II CTD for P-TEFb recognition. *Nat Commun* **3,** 842 (2012).

19.

Davey, R. T., Bhat, N., Yoder, C., Chun, T. W., Metcalf, J. A., Dewar, R., Natarajan, V.,

Lempicki, R. A., Adelsberger, J. W., Miller, K. D., Kovacs, J. A., Polis, M. A., Walker, R. E.,

Falloon, J., Masur, H., Gee, D., Baseler, M., Dimitrov, D. S., Fauci, A. S. & Lane, H. C. HIV-1

and T cell dynamics after interruption of highly active antiretroviral therapy (HAART) in

patients with a history of sustained viral suppression. *Proc Natl Acad Sci U S A* **96,** 15109–15114

(1999).

20.

de la Fuente, C., Maddukuri, A., Kehn, K., Baylor, S. Y., Deng, L., Pumfery, A. & Kashanchi, F.

Pharmacological cyclin-dependent kinase inhibitors as HIV-1 antiviral therapeutics. *Curr HIV*

Res **1,** 131–152 (2003).

21.

Desimio, M. G., Giuliani, E. & Doria, M. The histone deacetylase inhibitor SAHA

simultaneously reactivates HIV-1 from latency and up-regulates NKG2D ligands sensitizing for

natural killer cell cytotoxicity. *Virology* **510,** 9–21 (2017).

22.

Fabrega, C., Shen, V., Shuman, S. & Lima, C. D. Structure of an mRNA capping enzyme bound

to the phosphorylated carboxy-terminal domain of RNA polymerase II. *Mol Cell* **11,** 1549–1561

(2003).

651   23.

652   Firestein, R., Bass, A. J., Kim, S. Y., Dunn, I. F., Silver, S. J., Guney, I., Freed, E., Ligon, A. H.,

653   Vena, N., Ogino, S., Chheda, M. G., Tamayo, P., Finn, S., Shrestha, Y., Boehm, J. S., Jain, S.,

654   Bojarski, E., Mermel, C., Barretina, J., Chan, J. A., Baselga, J., Tabernero, J., Root, D. E., Fuchs,

655   C. S., Loda, M., Shivdasani, R. A., Meyerson, M. & Hahn, W. C. CDK8 is a colorectal cancer

656   oncogene that regulates beta-catenin activity. *Nature* **455,** 547–551 (2008).

657   24.

658   Fryer, C. J., White, J. B. & Jones, K. A. Mastermind recruits CycC:CDK8 to phosphorylate the

659   Notch ICD and coordinate activation with turnover. *Mol Cell* **16,** 509–520 (2004).

660   25.

661   Gunst, J. D., Højen, J. F. & Søgaard, O. S. Broadly neutralizing antibodies combined with

662   latency-reversing agents or immune modulators as strategy for HIV-1 remission. *Curr Opin HIV*

663   *AIDS* **15,** 309–315 (2020).

664   26.

665   Ho, C. K. & Shuman, S. Distinct roles for CTD Ser-2 and Ser-5 phosphorylation in the

666   recruitment and allosteric activation of mammalian mRNA capping enzyme. *Mol Cell* **3,** 405–

667   411 (1999).

668   27.

Horvath, R., Brumme, Z. L. & Sadowski, I. *CDK8 inhibitors antagonize HIV-1 reactivation and*

promote provirus latency in T cells. (Microbiology, 2023). doi:10.1101/2023.06.07.544021

28.

Horvath, R. M., Brumme, Z. L. & Sadowski, I. Inhibition of the TRIM24 bromodomain

reactivates latent HIV-1. *Sci Rep* **13,** 556 (2023).

29.

Horvath, R. M., Dahabieh, M., Malcolm, T. & Sadowski, I. TRIM24 controls induction of latent

HIV-1 by stimulating transcriptional elongation. *Commun Biol* **6,** 86 (2023).

30.

Huang, K.-L., Jee, D., Stein, C. B., Elrod, N. D., Henriques, T., Mascibroda, L. G., Baillat, D.,

Russell, W. K., Adelman, K. & Wagner, E. J. Integrator Recruits Protein Phosphatase 2A to

Prevent Pause Release and Facilitate Transcription Termination. *Mol Cell* **80,** 345-358.e9 (2020).

31.

Jin, H., Li, D., Sivakumaran, H., Lor, M., Rustanti, L., Cloonan, N., Wani, S. & Harrich, D.

Shutdown of HIV-1 Transcription in T Cells by Nullbasic, a Mutant Tat Protein. *mBio* **7,**

e00518-16 (2016).

32.

Jin, H., Sun, Y., Li, D., Lin, M.-H., Lor, M., Rustanti, L. & Harrich, D. Strong In Vivo Inhibition

of HIV-1 Replication by Nullbasic, a Tat Mutant. *mBio* **10,** e01769-19 (2019).

33.

Jordan, A., Bisgrove, D. & Verdin, E. HIV reproducibly establishes a latent infection after acute

infection of T cells in vitro. *EMBO J* **22,** 1868–1877 (2003).

34.

Karn, J. The molecular biology of HIV latency: breaking and restoring the Tat-dependent

transcriptional circuit. *Curr Opin HIV AIDS* **6,** 4–11 (2011).

35.

Kedei, N., Lundberg, D. J., Toth, A., Welburn, P., Garfield, S. H. & Blumberg, P. M.

Characterization of the interaction of ingenol 3-angelate with protein kinase C. *Cancer Res* **64,**

3243–3255 (2004).

36.

Kessing, C. F., Nixon, C. C., Li, C., Tsai, P., Takata, H., Mousseau, G., Ho, P. T., Honeycutt, J.

B., Fallahi, M., Trautmann, L., Garcia, J. V. & Valente, S. T. In Vivo Suppression of HIV

Rebound by Didehydro-Cortistatin A, a 'Block-and-Lock' Strategy for HIV-1 Treatment. *Cell*

Rep **21,** 600–611 (2017).

37.

Kim, Y. K., Bourgeois, C. F., Pearson, R., Tyagi, M., West, M. J., Wong, J., Wu, S.-Y., Chiang,

C.-M. & Karn, J. Recruitment of TFIIH to the HIV LTR is a rate-limiting step in the emergence

of HIV from latency. *EMBO J* **25,** 3596–3604 (2006).

38.

Kim, Y., Anderson, J. L. & Lewin, S. R. Getting the 'Kill' into 'Shock and Kill': Strategies to

Eliminate Latent HIV. *Cell Host Microbe* **23,** 14–26 (2018).

39.

Kuang, Y., Golan, O., Preusse, K., Cain, B., Christensen, C. J., Salomone, J., Campbell, I.,

Okwubido-Williams, F. V., Hass, M. R., Yuan, Z., Eafergan, N., Moberg, K. H., Kovall, R. A.,

Kopan, R., Sprinzak, D. & Gebelein, B. Enhancer architecture sensitizes cell specific responses

to Notch gene dose via a bind and discard mechanism. *Elife* **9,** e53659 (2020).

40.

Laird, G. M., Bullen, C. K., Rosenbloom, D. I. S., Martin, A. R., Hill, A. L., Durand, C. M.,

Siliciano, J. D. & Siliciano, R. F. Ex vivo analysis identifies effective HIV-1 latency-reversing

drug combinations. *J Clin Invest* **125,** 1901–1912 (2015).

41.

Li, J., Ji, H., Porter, D. C., Broude, E. V., Roninson, I. B. & Chen, M. Characterizing CDK8/19

Inhibitors through a NFκB-Dependent Cell-Based Assay. *Cells* **8,** 1208 (2019).

42.

Li, J., Chen, C., Ma, X., Geng, G., Liu, B., Zhang, Y., Zhang, S., Zhong, F., Liu, C., Yin, Y.,

Cai, W. & Zhang, H. Long noncoding RNA NRON contributes to HIV-1 latency by specifically

inducing tat protein degradation. *Nat Commun* **7,** 11730 (2016).

43.

Li, Z., Guo, J., Wu, Y. & Zhou, Q. The BET bromodomain inhibitor JQ1 activates HIV latency

through antagonizing Brd4 inhibition of Tat-transactivation. *Nucleic Acids Res* **41,** 277–287

(2013).

44.

Liu, Q., Yin, X., Languino, L. R. & Altieri, D. C. Evaluation of drug combination effect using a

Bliss independence dose-response surface model. *Stat Biopharm Res* **10,** 112–122 (2018).

45.

Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time

quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25,** 402–408 (2001).

46.

McMahon, D. K., Zheng, L., Cyktor, J. C., Aga, E., Macatangay, B. J., Godfrey, C., Para, M.,

Mitsuyasu, R. T., Hesselgesser, J., Dragavon, J., Dobrowolski, C., Karn, J., Acosta, E. P.,

739   Gandhi, R. T. & Mellors, J. W. A Phase 1/2 Randomized, Placebo-Controlled Trial of

740   Romidespin in Persons With HIV-1 on Suppressive Antiretroviral Therapy. *J Infect Dis* **224,**

741   648–656 (2021).

742   47.

743   Meijer, L., Borgne, A., Mulner, O., Chong, J. P., Blow, J. J., Inagaki, N., Inagaki, M., Delcros, J.

744   G. & Moulinoux, J. P. Biochemical and cellular effects of roscovitine, a potent and selective

745   inhibitor of the cyclin-dependent kinases cdc2, cdk2 and cdk5. *Eur J Biochem* **243,** 527–536

746   (1997).

747   48.

748   Moranguinho, I. & Valente, S. T. Block-And-Lock: New Horizons for a Cure for HIV-1. *Viruses*

749   **12,** 1443 (2020).

750   49.

751   Mori, L., Jenike, K., Yeh, Y.-H. J., Lacombe, B., Li, C., Getzler, A., Mediouni, S., Cameron, M.,

752   Pipkin, M., Ho, Y.-C., Ramirez, B. C. & Valente, S. The XPB Subunit of the TFIIH Complex

753   Plays a Critical Role in HIV-1 Transcription and XPB Inhibition by Spironolactone Prevents

754   HIV-1 Reactivation from Latency. *J Virol* **95,** e01247-20, JVI.01247-20 (2021).

755   50.

756   Mousseau, G., Kessing, C. F., Fromentin, R., Trautmann, L., Chomont, N. & Valente, S. T. The

757   Tat Inhibitor Didehydro-Cortistatin A Prevents HIV-1 Reactivation from Latency. *mBio* **6,**

758   e00465 (2015).

759   51.

760   Olson, C. M., Liang, Y., Leggett, A., Park, W. D., Li, L., Mills, C. E., Elsarrag, S. Z., Ficarro, S.

761   B., Zhang, T., Düster, R., Geyer, M., Sim, T., Marto, J. A., Sorger, P. K., Westover, K. D., Lin,

C. Y., Kwiatkowski, N. & Gray, N. S. Development of a Selective CDK7 Covalent Inhibitor Reveals Predominant Cell-Cycle Phenotype. *Cell Chem Biol* **26,** 792-803.e10 (2019).

52. Porter, D. C., Farmaki, E., Altilia, S., Schools, G. P., West, D. K., Chen, M., Chang, B.-D., Puzyrev, A. T., Lim, C., Rokow-Kittell, R., Friedhoff, L. T., Papavassiliou, A. G., Kalurupalle, S., Hurteau, G., Shi, J., Baran, P. S., Gyorffy, B., Wentland, M. P., Broude, E. V., Kiaris, H. & Roninson, I. B. Cyclin-dependent kinase 8 mediates chemotherapy-induced tumor-promoting paracrine activities. *Proc Natl Acad Sci U S A* **109,** 13799–13804 (2012).

53. Powell, A. B., Ren, Y., Korom, M., Saunders, D., Hanley, P. J., Goldstein, H., Nixon, D. F., Bollard, C. M., Lynch, R. M., Jones, R. B. & Cruz, C. R. Y. Engineered Antigen-Specific T Cells Secreting Broadly Neutralizing Antibodies: Combining Innate and Adaptive Immune Response against HIV. *Mol Ther Methods Clin Dev* **19,** 78–88 (2020).

54. Rodríguez-Molina, J. B., Tseng, S. C., Simonett, S. P., Taunton, J. & Ansari, A. Z. Engineered Covalent Inactivation of TFIIH-Kinase Reveals an Elongation Checkpoint and Results in Widespread mRNA Stabilization. *Mol Cell* **63,** 433–444 (2016).

55. Sadowski, I. & Hashemi, F. B. Strategies to eradicate HIV from infected patients: elimination of latent provirus reservoirs. *Cell Mol Life Sci* **76,** 3583–3600 (2019).

56. Schang, L. M., Bantly, A., Knockaert, M., Shaheen, F., Meijer, L., Malim, M. H., Gray, N. S. & Schaffer, P. A. Pharmacological cyclin-dependent kinase inhibitors inhibit replication of wild-

type and drug-resistant strains of herpes simplex virus and human immunodeficiency virus type 1 by targeting cellular, not viral, proteins. *J Virol* **76,** 7874–7882 (2002).

57. Schonhofer, C., Yi, J., Sciorillo, A., Andrae-Marobela, K., Cochrane, A., Harris, M., Brumme, Z. L., Brockman, M. A., Mounzer, K., Hart, C., Gyampoh, K., Yuan, Z., Montaner, L. J. & Tietjen, I. Flavonoid-based inhibition of cyclin-dependent kinase 9 without concomitant inhibition of histone deacetylases durably reinforces HIV latency. *Biochem Pharmacol* **186,** 114462 (2021).

58. Siliciano, J. D. & Siliciano, R. F. Low Inducibility of Latent Human Immunodeficiency Virus Type 1 Proviruses as a Major Barrier to Cure. *J Infect Dis* **223,** 13–21 (2021).

59. Steinparzer, I., Sedlyarov, V., Rubin, J. D., Eislmayr, K., Galbraith, M. D., Levandowski, C. B., Vcelkova, T., Sneezum, L., Wascher, F., Amman, F., Kleinova, R., Bender, H., Andrysik, Z., Espinosa, J. M., Superti-Furga, G., Dowell, R. D., Taatjes, D. J. & Kovarik, P. Transcriptional Responses to IFN-γ Require Mediator Kinase-Dependent Pause Release and Mechanistically Distinct CDK8 and CDK19 Functions. *Mol Cell* **76,** 485-499.e8 (2019).

60. Tome, J. M., Tippens, N. D. & Lis, J. T. Single-molecule nascent RNA sequencing identifies regulatory domain architecture at promoters and enhancers. *Nat Genet* **50,** 1533–1541 (2018).

61. Vansant, G., Bruggemans, A., Janssens, J. & Debyser, Z. Block-And-Lock Strategies to Cure HIV Infection. *Viruses* **12,** 84 (2020).

62.

808 Wang, D., de la Fuente, C., Deng, L., Wang, L., Zilberman, I., Eadie, C., Healey, M., Stein, D.,

809 Denny, T., Harrison, L. E., Meijer, L. & Kashanchi, F. Inhibition of human immunodeficiency

810 virus type 1 transcription by chemical cyclin-dependent kinase inhibitors. *J Virol* **75,** 7266–7279

811 (2001).

812   63.

813 Wong, K. H., Jin, Y. & Struhl, K. TFIIH phosphorylation of the Pol II CTD stimulates mediator

814 dissociation from the preinitiation complex and promoter escape. *Mol Cell* **54,** 601–612 (2014).

815   64.

816 Yamaguchi, Y., Shibata, H. & Handa, H. Transcription elongation factors DSIF and NELF:

817 promoter-proximal pausing and beyond. *Biochim Biophys Acta* **1829,** 98–104 (2013).

818   65.

819 Yan, C., Dodd, T., He, Y., Tainer, J. A., Tsutakawa, S. E. & Ivanov, I. Transcription preinitiation

820 complex structure and dynamics provide insight into genetic diseases. *Nat Struct Mol Biol* **26,**

821 397–406 (2019).

822   66.

823 Yeh, Y.-H. J. & Ho, Y.-C. Shock-and-kill versus block-and-lock: Targeting the fluctuating and

824 heterogeneous HIV-1 gene expression. *Proc Natl Acad Sci U S A* **118,** e2103692118 (2021).

825   67.

826 Zhang, L., Chung, C., Hu, B. S., He, T., Guo, Y., Kim, A. J., Skulsky, E., Jin, X., Hurley, A.,

827 Ramratnam, B., Markowitz, M. & Ho, D. D. Genetic characterization of rebounding HIV-1 after

828 cessation of highly active antiretroviral therapy. *J Clin Invest* **106,** 839–845 (2000).

829   68.

830   Zhao, X., Feng, D., Wang, Q., Abdulla, A., Xie, X.-J., Zhou, J., Sun, Y., Yang, E. S., Liu, L.-P.,

831   Vaitheesvaran, B., Bridges, L., Kurland, I. J., Strich, R., Ni, J.-Q., Wang, C., Ericsson, J., Pessin,

832   J. E., Ji, J.-Y. & Yang, F. Regulation of lipogenesis by cyclin-dependent kinase 8-mediated

833   control of SREBP-1. *J Clin Invest* **122,** 2417–2427 (2012).

834   69.

835   Zhu, J., Gaiha, G. D., John, S. P., Pertel, T., Chin, C. R., Gao, G., Qu, H., Walker, B. D.,

836   Elledge, S. J. & Brass, A. L. Reactivation of latent HIV-1 by inhibition of BRD4. *Cell Rep* **2,**

837   807–816 (2012).

 Legends to Figures

Figure 1. Chemical inhibition of tCDKs is well tolerated in T cells. A-C: Molecular structure

of YKL-5-124 (CDK7 inhibitor) (**A**), LDC000067 (CDK9 inhibitor) (**B**), and Senexin A (CDK8

and CDK19 inhibitor) (**C**). **D:** Jurkat human T cells were treated with increasing concentrations

of the indicated compound for 24 hrs after which cellular viability was determined. Results are

normalized to untreated control ($n = 2$, mean ±□SD).

Figure 2. HIV-1 latency cell line models require tCDK kinase activity for reactivation. A:

Representative flow cytometry scatter plots of JLat10.6 cells that were pre-treated for 30 min

with DMSO (Vehicle), 100 nM YKL-5-124 (CDK7i), 1 μM LDC000067 (CDK9i), or 10 μM

Senexin A (CDK8/19i) and subsequently incubated with 5 nM PMA and 1 μM ionomycin for 20

hrs. **B, C:** Following 30 min pre-treatment with the indicated concentration of tCDK inhibitor,

JLat10.6 cells were incubated with 5 nM PMA and 1 μM ionomycin. After 20 hrs, flow

cytometric analysis was performed to determine HIV-1 expression as reported as the proportion

of GFP positive cells (**B**) and the GFP delta (Δ) Mean Fluorescent Intensity (MFI) (MFI of

treatment subtracted by the MFI of untreated) (**C**) ($n = 2$, mean ±□SD). **D:** As in (**A**), but JLat9.2

cells were assessed. **E, F:** As in B, C, but JLat9.2 cells were examined ($n = 2$, mean ±□SD).

Figure 3. tCDK inhibitors synergistically inhibit HIV-1 transcription. A - C: JLat10.6 cells

were pre-treated for 30 min with 100 nM YKL-5-124 (CDK7i) and the indicated concentration

of LDC000067 (CDK9i) (**A**) or Senxexin A (CDK8/19i) (**B**), or were pre-treated 30 min with 1

μM LDC000067 and the indicated concentration of Senexin A (**C**). Subsequently, cells were

incubated with 5 nM PMA/ 1 μM ionomycin for 20 hrs and analyzed by flow cytometry

following 20 hrs (*n* = 2, mean ±□SD). **D - F:** Calculation of synergy between the indicated

treatments in **A - C** using Bliss Independence Modeling. Data are presented as the difference

between the predicted and the observed fractional HIV-1 expression response to the given drug

combination (*n* = 2, mean ±□SD). See Materials and Methods for more details.

Figure 4. tCDK kinase activity mediates proviral expression in response to latency reversal

agents. A, B: JLat10.6 cells were pre-treated for 30 min with 100 nM YKL-5-124 (CDK7i), 1

μM LDC000067 (CDK9i), 10 μM Senexin A (CDK8/19i), or a vehicle control (Ve, DMSO).

Subsequently, cells were incubated with10 μM JQ1, 20 μM IACS-9571, 5 nM PEP005, 5 nM

PEP005/ 20 μM IACS-9571, 5 nM PMA/ 1 μM ionomycin, 5 μM SAHA, or were left untreated.

Following 20 hrs, viral expression was assessed by flow cytometry and reported as the percent of

GFP positive cells (**A**) and the Mean Fluorescent Intensity (MFI) of GFP (**B**) (*n* = 2, mean

±□SD).

Figure 5. Tat dependency of tCDK kinase activity. A: Representative flow cytometry scatter

plots of JLatA72 Jurkat cells (-Tat) that were pre-treated for 30 min with a vehicle control

(DMSO), 100 nM YKL-5-124 (CDK7i), 1 μM LDC000067 (CDK9i), or 50 μM Senexin A

(CDK8/19i) and subsequently incubated with 5 nM PMA and 1 μM ionomycin for 20 hrs. **B, C:**

JLatA72 cells were left untreated or pre-treated for 30 min with a vehicle control (DMSO), 100

nM YKL-5-124, 1 μM LDC00067, or 50 μM Senexin A prior to being incubated with 5 nM

PMA/ 1 μM ionomycin. Following 20 hrs, flow cytometry was performed with viral expression

reported as the percent of GFP positive cells (**B**) and the delta GFP Mean Fluorescent Intensity

(MFI) (**C**) ($n = 2$, mean ±□SD).

Figure 6. Effect of sustained tCDK inhibition on cellular growth and viability. A, B: Jurkat

T cells were incubated with the indicated concentration of YKL-5-124 or a vehicle control (Ve,

DMSO) with cell number (**A**) and viability (**B**) assessed on the indicated day. Media was

replaced daily with inhibitor ($n = 3$, mean ±□SD). **C, D:** As in (**A**) and (**B**), but cells were treated

with LDC000067 ($n = 3$, mean ±□SD). **E, F:** Performed as previously described but Jurkat cells

were incubated with the indicated concentration of Senexin A ($n = 3$, mean ±□SD).

Figure 7. tCDK activity mediates the establishment of latency upon infection. A: Simplified

depiction of the replication incompetent Red-Green-HIV-1 (RGH) dual reporter virus (Dahabieh

et al, 2013). RGH is a full-length virus that is rendered to a single round of infection through

mutation in *Env*, and that expresses GFP as driven by 5' LTR transcription while also expressing

mCherry from a constitutive PGK promoter that replaces *Nef,* allowing for latent and productive

infections to be differentiated. **B:** Schematic representation of the RGH assay. Jurkat CD4[+]

human T cells were left untreated or were pre-treated for 30 min with 100 nM YKL-5-124, 1 μM

LDC000067, or 10 μM Senexin A. Subsequently, cells were infected with RGH using equivalent

amounts of virion. 1-day post-infection, media was refreshed and cells were either washed of the

drug or maintained with the same concentration of inhibitor as prior. 4 days post-infection,

proviral expression was assessed by flow cytometry. **C:** Representative flow cytometry scatter

plots of RGH infected cells, treated as indicated in (**B**) and assessed 4 days post-infection. **D:**

Jurkat T cells were treated as in (**B**), with the percentage of productive infections determined by

flow cytometry 4 days post-infection. The %Productive Infections was determined from the ratio

of GFP+/mCherry+ cells to all mCherry+ cells ($n = 3$, mean $\pm$ SD). **E:** Bar graph summary of

the infection rate for the indicated treatment determined by cells displaying PGK driven mCherry

expression ($n = 3$, mean $\pm$ SD).

Figure 8. Effect of CDK9 and/or CDK8/19 inhibition on mdHIV #77 proviral latency. A:

Simplified depiction of the replication incompetent mini-dual HIV-1 reporter virus that is

chromosomally integrated in the mdHIV #77 Jurkat Tat clonal T cell line. 5' LTR transcription

drives dsRed expression as a fusion with p24 Gag while an internal EF1α promoter expresses

eGFP. All other auxiliary proteins including Tat are not present. **B:** Experimental design. mdHIV

#77 cells were incubated with a vehicle control (DMSO), 1 μM LDC000067, 10 μM Senexin A,

or 1 μM LDC000067 and 10 μM Senexin A for 14 days after which the drug was removed from

the media by washing and viral expression was assessed by flow cytometry for 10 more days. **C:**

Representative flow cytometry scatter plots following 4 days of drug treatment and 4 days

following drug withdrawal. All "dots" are indicative of a single HIV-1 infected cell with Q2

(dsRED+/eGFP+) possessing cells with transcriptionally active provirus. **D:** mdHIV #77 cells

were treated as in (**B**) with viral expression assessed on the indicated day by flow cytometry ($n =$

2, mean $\pm$ SD). **E:** mdHIV #77 cells treated as indicated in (**B**) were examined for viability,

with values normalized to the vehicle control ($n = 2$, mean $\pm$ SD).

Figure 9. Effect of CDK9 and/or CDK8/19 inhibition on RGH #20 proviral latency. A:
Simplified depiction of the full-length, replication incompetent HIV-1 reporter virus that is
chromosomally integrated in the RGH #20 Jurkat clonal T cell line. 5' LTR transcription drives
eGFP expression as a fusion with p24 Gag while mCherry is expressed from an internal PGK
promoter that replaces *Nef*. Other than *Nef*, all auxiliary proteins are expressed while a mutation
in *Env* renders the provirus replication incompetent **B:** Experimental design. RGH #20 cells were
incubated with a vehicle control (DMSO), 1 μM LDC000067, 10 μM Senexin A, or 1 μM
LDC000067 and 10 μM Senexin A for 7 days after which the drug was removed from the media
by washing and viral expression was assessed by flow cytometry for 10 more days. **C:**
Representative flow cytometry scatter plots following 4 days of drug treatment and 4 days
following drug withdrawal. All "dots" are indicative of a single HIV-1 infected cell with Q2
(eGFP+/mCherry+) displaying cells with transcriptionally active provirus. **D:** RGH #20 cells
were treated as in (**B**) with viral expression assessed on the indicated day by flow cytometry ($n =$
2, mean $\pm$ SD). **E:** RGH #20 cells treated as indicated in (**B**) were examined for viability, with
values normalized to the vehicle control ($n = 2$, mean $\pm$ SD).

Figure 10. tCDK activity is required for reactivation of HIV-1 in patient CD4+ PBMCs *ex*
***vivo*. A - C:** CD4+ PBMCs, isolated from HIV-1 infected individuals on ART, were either left
untreated or were pre-treated for 30 min with a Vehicle (DMSO), 100 nM YKL-5-124, 1 μM
LDC000067, 10 μM Senexin A, or 1 μM LDC000067/ 10 μM Senexin A. Following the pre-
treatment, cells were incubated with 5nM PMA/ 1μM Ionomycin for 24 hrs after which
intracellular RNA was extracted and analyzed using RT-PCR with oligos specific for multiply

spliced Tat-Rev HIV-1 mRNA transcripts. HIV-1 mRNA expression is normalized to *GAPDH* (*n*

= 3 - 4, mean ±☐SD).

Legends to Supplementary Figures

Figure S1. Effect of tCDK inhibition on cellular viability. A – C: JLat10.6 cells were pre-

treaterd 30 min with the indicated concentration of YKL-5-124 (CDK7i) (**A**), LDC000067

(CDK9i) (**B**), or Senexin A (CDK8/19i) (**C**) and were subsequently stimulated with 5 nM PMA/

1 μM ionomycin. Following 20 hrs, cell viability was assessed and is normalized to cells absent

tCDK inhibitor (*n* = 2, mean ±☐SD). **D – F:** As in (**A – C**), but JLat9.2 cells were assessed (*n* =

2, mean ±☐SD).

Figure S2. **Effect of combined tCDK inhibitor treatment on cellular viability. A – C:**

JLat10.6 cells were pre-treated for 30 min with 100 nM YKL-5-124 (CDK7i) and the indicated

concentration of LDC000067 (CDK9i) (**A**) or Senxexin A (CDK8/19i) (**B**), or were pre-treated

30 min with 1 μM LDC000067 and the indicated concentration of Senexin A (**C**). Subsequently,

cells were incubated with 5 nM PMA/ 1 μM ionomycin for 20 hrs. Following incubation, cell

viability was determined and normalized to cells absent tCDK inhibitor (*n* = 2, mean ±☐SD).

Figure S3. IACS-9571 strongly synergizes with PEP005 to reactivate latent HIV-1. JLat10.6

cells were left untreated, or were incubated with 5 nM PEP005, 20 μM IACS-9571, or 5 nM

PEP005/ 20 µM IACS-9571. Following 20 hrs, cells were assessed for GFP expression by flow

cytometry as shown in Fig. 4A. Calculation of synergy between the indicated treatments was

performed using Bliss Independence Modeling. Data are presented as the difference between the

predicted and the observed fractional HIV-1 expression response to the given drug combination

($n = 2$, mean ±□SD). See Materials and Methods for more details.

Figure S4. Tat dependence of tCDK for proviral induction A: Representative flow cytometry

scatter plots of GXR-5 CEM cells (-Tat) that were pre-treated for 30 min with a vehicle control

(DMSO), 100 nM YKL-5-124 (CDK7i), 1 µM LDC000067 (CDK9i), or 50 µM Senexin A

(CDK8/19i) and subsequently incubated with 5 nM PMA and 1 µM ionomycin for 20 hrs. **B, C:**

GXR-5 cells were left untreated or pre-treated for 30 min with a vehicle control (DMSO), 100

nM YKL-5-124, 1 µM LDC00067, or 50 µM Senexin A prior to being incubated with 5 nM

PMA/ 1 µM ionomycin. Following 20 hrs, flow cytometry was performed with viral expression

reported as the percent of GFP positive cells (**B**) and the delta GFP Mean Fluorescent Intensity

(MFI) (**C**) ($n = 2$, mean ±□SD).

Figure S5. Effect of tCDK inhibitors on cell viability during HIV-1 infection. Jurkat human

T cells were left untreated or were pre-treated for 30 min with 100 nM YKL-5-124, 1 µM

LDC000067, or 10 µM Senexin A. Cells were subsequently infected with RG and 1-day post-

infection, media was refreshed and cells were either washed of the drug or maintained with the

same concentration of inhibitor as prior. 4 days post-infection, viability was determined and is

normalized to untreated cells ($n = 3$, mean ±□SD).

1004

Figure S6. Effect of CDK9 and/or CDK8/19 inhibition on mdHIV #110 proviral latency. A:

Simplified depiction of the replication incompetent mini-dual HIV-1 reporter virus that is

chromosomally integrated in the mdHIV #110 Jurkat Tat clonal T cell line. 5' LTR transcription

drives dsRed expression as a fusion with p24 Gag while an internal EF1α promoter expresses

eGFP. All other auxiliary proteins including Tat are not present. **B:** Experimental design. mdHIV

#110 cells were incubated with a vehicle control (DMSO), 1 μM LDC000067, 10 μM Senexin A,

or 1 μM LDC000067 and 10 μM Senexin A for 14 days after which the drug was removed from

the media by washing and viral expression was assessed by flow cytometry for 10 more days. **C:**

Representative flow cytometry scatter plots following 4 days of drug treatment and 4 days

following drug withdrawal. All "dots" are indicative of a single HIV-1 infected cell with Q2

(dsRED+/eGFP+) possessing cells with transcriptionally active provirus. **D:** mdHIV #110 cells

were treated as in (**B**) with viral expression assessed on the indicated day by flow cytometry ($n =$

2, mean ±SD). **E:** mdHIV #110 cells treated as indicated in (**B**) were examined for viability,

with values normalized to the vehicle control ($n = 2$, mean ±SD).

Figure 1

A

YKL-5-124 – CDK7 Inhibitor

B

LDC000067 – CDK9 Inhibitor

C

Senexin A – CDK8/19 Inhibitor

D

LC50: 93 µM, 499 µM, 200 µM

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

A

B

C

Table 1

Patient	ID	Sex	Intact HIV Copies/Million CD4$^+$ T cells	Total HIV Copies/Million CD4$^+$ T cells	%Intact
A	BC024	Male	339.88	4941.32	6.88
B	BC030	Male	397.27	3774.98	10.52
C	BC002	Male	197.31	4857.65	4.06