Habitat fragmentation weakens the positive relationship between grassland plant richness and above-ground biomass

Yongzhi Yan[1], Scott Jarvie[2], Qing Zhang[1, 3, 4*]

[1]Ministry of Education Key Laboratory of Ecology and Resource Use of the Mongolian Plateau, School of Ecology and Environment, Inner Mongolia University, Hohhot 010021, China

[2]Otago Regional Council, Dunedin 9016, New Zealand

[3]Collaborative Innovation Center for Grassland Ecological Security (Jointly Supported by the Ministry of Education of China and Inner Mongolia Autonomous Region), Hohhot, 010021, China

[4]Autonomous Region Collaborative Innovation Center for Integrated Management of Water Resources and Water Environment in the Inner Mongolia Reaches of the Yellow River, Hohhot 010018, China

*Correspondence:

Qing Zhang

Email: qzhang82@163.com

Highlights:

1. BEF relationship is moderated by landscape context in fragmented landscapes.

2. Habitat loss and fragmentation per se have inconsistent effects.

3. Habitat loss can weaken the positive BEF relationship.

Abstract:

Anthropogenic habitat fragmentation has been shown to be a major threat to global biodiversity and ecosystem function. However, little is known about how habitat fragmentation alters the relationship between biodiversity and ecosystem function (BEF relationship). Based on 130 landscapes identified using a stratified random sampling in the agro-pastoral ecotone of northern China, we investigated using a structural equation model the effects of habitat fragmentation (including habitat loss and fragmentation per se) on plant richness, above-ground biomass, and the relationship between them in grassland communities. We found that habitat loss decreased plant richness, while fragmentation per se increased plant richness. Fragmentation per se decreased above-ground biomass, while habitat loss had no significant effect on above-ground biomass. Neither habitat loss nor fragmentation per se affected the direction of the positive relationship between plant richness and above-ground biomass. However, habitat loss decreased the magnitude of the positive relationship by reducing the percentage of grassland specialists in the community. These results demonstrate that habitat loss and fragmentation per se have inconsistent effects on biodiversity and ecosystem function, with the BEF relationship being moderated by landscape context. Our findings emphasize that habitat loss rather than fragmentation per se can weaken the positive BEF relationship by decreasing the degree of habitat specialization of the community.

Keywords:

BEF relationship, biodiversity, ecosystem function, fragmentation per se, generalists, habitat loss, moderating effect, specialists.

1 Introduction

Evidence from biodiversity-ecosystem function (BEF hereafter) experiments during the past 30 years generally show positive relationships between biodiversity and productivity, soil carbon storage, decomposition rates, and other ecosystem functions in experimental communities, revealing the importance of biodiversity in maintaining ecosystem functioning (Tilman et al., 2012; van der Plas, 2019). When research expands from experiments to natural systems, however, BEF relationships remain unclear in the natural assembled communities, with significant context dependency (Hagan et al., 2021; van der Plas, 2019). One of the main reasons for these differences

is because the landscape context surrounding natural communities regulates biodiversity and ecosystem function (Gonzalez et al., 2020; Liu et al., 2018). Consideration of the impacts of landscape context on surrounding communities is key to understand complex BEF relationships in natural systems.

Human activities have modified natural ecosystems globally, with habitat fragmentation currently being a typical landscape context (Chase et al., 2020; Maxwell et al., 2016). For example, at least 1.5 billion hectares of natural habitats on Earth have been converted to human-modified land since 2014, breaking apart continuous habitat into smaller and isolated fragments (IPBES, 2019). Habitat fragmentation in natural landscapes consists of two facets: 1) habitat loss, i.e., reducing habitat amount, and 2) fragmentation per se, i.e., breaking apart of habitat such as decreased size of habitat patches, increased number of habitat patches, increased isolation among habitat patches, etc. (Fahrig, 2003; Ibáñez et al., 2014; Wang et al., 2014). Recent studies suggest that habitat loss and fragmentation per se are related to different processes, with inconsistent ecological effects, and should be considered simultaneously to fully understand the ecological consequences of habitat fragmentation (Fahrig et al., 2019; Fletcher et al., 2018; Miller-Rushing et al., 2019).

Habitat fragmentation is often considered the major near-term threat to biodiversity and ecosystem function (Chase et al., 2020; Fletcher Jr et al., 2023; Haddad et al., 2015), while the impact of fragmentation per se remains debated (Miller-Rushing et al., 2019). Habitat fragmentation can also affect BEF relationships. Previous studies have found that the magnitude and direction of BEF relationships vary across fragmented landscapes, including positive, negative, and non-significant relationships (Godbold et al., 2011; Hagan et al., 2021; Rolo et al., 2018; Zirbel et al., 2019). However, few studies have been conducted on how habitat fragmentation regulates BEF relationships (but see, for example, Liu et al., 2018; Wilson et al., 2016). This lack of studies hampers our understanding of complex BEF relationships in fragmented natural ecosystems. In theory, habitat fragmentation can regulate the BEF relationship by altering species composition and interactions regardless of changes in species richness (Liu et al., 2018). However, species in communities are not ecologically equivalent and may respond differently to habitat fragmentation and have different associations with ecosystem function (Devictor et al., 2008;

Wardle and Zackrisson, 2005). Therefore, considering changes in species composition can help to

understand how habitat fragmentation regulates BEF relationships.

The degree of habitat specialization is a key ecological characteristic determining responses

of species to habitat fragmentation and the slope of BEF relationships (Clavel et al., 2011; Gravel

et al., 2011). In fragmented landscapes, species with low levels of habitat specialization

(generalists) can use resources from different land covers, including focal habitat and non-habitat

matrix, and thus are not sensitive to habitat fragmentation (Matthews et al., 2014). In contrast,

species with high levels of habitat specialization (specialists) depend highly on resources in

specific habitats, and thus are vulnerable to adverse effects from habitat fragmentation (Matthews

et al., 2014). Meanwhile, specialists in communities are often highly productive and competitive,

which are closely associated with ecosystem function, whereas generalists are less productive and

competitive, which are redundant for ecosystem function (Dehling et al., 2021; Gravel et al., 2011;

Mello et al., 2015). Therefore, habitat fragmentation is often considered to decrease the degree of

habitat specialization (the replacement of specialists in communities by generalists), possibly

resulting in functional homogenization of communities and reduced BEF relationships (Clavel et

al., 2011; Matthews, 2021). However, few studies have evaluated this process in fragmented

landscapes.

Currently, research on habitat fragmentation focuses primarily on forest ecosystems (Fardila

et al., 2017; Haddad et al., 2015). Grasslands have received considerably less attention, despite

being the largest terrestrial ecosystem type, and suffering severe fragmentation due to human

activities, such as agricultural reclamation and urbanization (Fardila et al., 2017). The

agro-pastoral ecotone of northern China is a typical anthropogenically fragmented grassland

landscape caused by historical agricultural reclamation, especially in the late Qing Dynasty (about

1840–1912). Due to land policy reforms, the region has experienced a rapid expansion of farmland

since the 1960s, converting continuous natural grasslands into smaller and isolated fragments (Wu

et al., 2015; Yang et al., 2020). Habitat fragmentation has reduced plant diversity and ecosystem

function in these grassland fragments (Yan et al., 2022; Yan et al., 2023; Yang et al., 2020).

However, whether habitat fragmentation affects the relationship between plant diversity and

ecosystem function is still unclear.

Based on 130 landscapes with different fragmentation levels in the agro-pastoral ecotone of northern China, we investigated how habitat fragmentation (including habitat loss and fragmentation per se) regulates the relationship between grassland plant diversity and above-ground productivity in the community. Specifically, we aimed to evaluate whether habitat fragmentation would weaken the positive relationship between grassland plant diversity and above-ground productivity by reducing the habitat specialization of the community.

2 Materials and methods

2.1 Study area

Our study area is in the agro-pastoral ecotone of northern China, the Tabu River Basin in Siziwang Banner, Inner Mongolia Autonomous Region. The mean annual temperature ranges from 1.5 to 5.0 □, and the mean annual precipitation ranges from 225 to 322 mm. The type of soil is light chestnut soil. This area is a typical fragmented grassland landscape caused by agricultural intensification. Grassland is the dominant natural habitat type in this area, accounting for about 40.8 % of the total area, with the dominant plant species being *Stipa krylovii* and S. *breviflora*. Farmland is the dominant matrix type in this area, accounting for about 30.6 % of the total area, with the main crops grown are potatoes and maize. Further background information about the study area is described in our previous papers (Yan et al., 2022; Yan et al., 2023; Yan et al., 2021; Zhang et al., 2021).

2.2 Sampling landscape selection

We quantified landscape-scale habitat fragmentation in the study area to determine the spatial gradient of habitat fragmentation (including habitat loss and fragmentation per se), then established the sampling landscapes. Grassland was defined as the focal habitat. Habitat amount was represented by the percentage of grassland cover in the landscape. Habitat loss was represented by non-grassland cover in the landscape. Fragmentation per se was estimated using four landscape indices, reflecting different landscape-scale fragmentation processes (Fahrig, 2003;

Fahrig, 2017): (1) mean patch area metric, representing a decrease in grassland patches area in the

landscape; (2) mean nearest-neighbor distance metric, representing an increase in the isolation of

grassland patches in the landscape; (3) patch density metric, representing an increase in the

number of grassland patches in the landscape; and (4) edge density metric, representing an

increase in the grassland edges in the landscape. Given that habitat amount (or habitat loss) and

fragmentation per se are typically highly correlated in natural landscapes, it is hard to disentangle

their relative effects (Fahrig, 2017; Smith et al., 2009). We therefore used stratified sampling to

select sampling landscapes across the relative independent spatial gradient of habitat amount and

fragmentation per se, which can reduce collinearity between them (Pasher et al., 2013; Reynolds

et al., 2018).

To do so, we first used the moving window method (window size: 500-m radius buffer) to

quantify grassland amount and the four landscape indices surrounding all grassland cells. The

500-m radius buffer was used because our previous studies showed this buffer includes the

optimal scale of spatial processes influencing grassland plant diversity in this region (Yan et al.,

2022; Yan et al., 2023; Zhang et al., 2021). Second, we derived the first principal component (PC1)

of the four landscape indices as a single fragmentation per se index (Hertzog et al., 2019; Rolo et

al., 2018). This index was positively correlated with patch density, edge density, mean

nearest-neighbor distance metric, and negatively with mean patch area (Appendix Figure A1 and

Table A1). Based on the quartiles of grassland amount and PC1 of the four landscape indices, we

ranked these grassland cells into nine types: high-high, high-moderate, high-low, moderate-high,

moderate-moderate, moderate-low, low-high, low-moderate, low-low grassland amount and

fragmentation per se (Appendix Figure A2). Given landscapes with a high grassland amount and

high fragmentation are scarce in this region (Appendix Figure A2), we did not consider this type

of landscape. Finally, we selected at least 20 landscapes using stratified sampling from each of the

remaining eight grassland types as sites for field surveys.

The land-cover data used to quantify grassland fragmentation was obtained via supervised

classification on a cloud-free Landsat 8 TOA composite image (30-m resolution) from 2019 (Yan

et al., 2022). We used the random forest classifier in the Google Earth Engine platform (Gorelick

et al., 2017) for the supervised classification. The moving window analysis and all landscape

metric calculations were performed in FRAGSTATS v4.2.1 based on the eight-cells neighborhood

rule (McGarigal et al., 2012). The principal component analysis was performed in the R

programming language v. 4.0.3 (R Development Core Team, 2020). Stratified sampling was

conducted in ArcGIS v10.3.

2.3 Biodiversity and ecosystem function surveys

Based on the landscapes selected above, we established 130 sites (30 m * 30 m) between late

July to mid-August 2020 in the Tabu River Basin in Siziwang Banner, Inner Mongolia

Autonomous Region (Figure 1). To prevent overlapping landscapes and potential spatial

autocorrelation, the shortest distance between each site was at least 1000 m. The type of habitat of

the selected sites was grasslands with regional vegetation characteristics. Each site showed no

signs of having been reclaimed. In the center of each site, we surveyed grassland vascular plant

diversity and above-ground productivity in 1 m * 1 m plots set within 10 m * 10 m

topographically flat areas. Plant diversity was obtained by recording the number of vascular plant

species in each plot. The above-ground productivity was obtained by harvesting the above-ground

biomass of the plants in each plot and drying biomass at 65 °C to a constant weight.

As grassland is the dominant habitat type in the fragmented landscape and farmland is the

dominant matrix type, the specialists and generalists in this study were grouped as grassland

specialists, i.e., species that occur only in grassland, and weeds, i.e., species that occur in both

grassland and farmland. The classification of grassland specialists and weeds in this study was

based on our experience with plant surveys in this region, the List of Main Crop Weeds in China,

and available information in the Flora of China (http://www.iplant.cn/frps).

Figure 1. Map of study area in the Tabu River Basin, Inner Mongolia Autonomous Region,

northern China. (a) Location of the 130 survey sites in the study area. (b) Examples of four survey

sites with varying levels of habitat loss and fragmentation per se shown with a 500-m radius

buffer.

2.4 Data analysis

For each site, we used data from three 1 m * 1 m plots to calculate the mean richness of

vascular plant, representing the vascular plant diversity, and mean above-ground biomass of

vascular plants, representing the above-ground productivity. The mean vascular plant richness and

the mean above-ground biomass of vascular plants were assessed to be normally distributed by a

Shapiro-Wilk normality test. The degree of habitat specialization was represented the percentage

of grassland specialists in the community. Habitat loss was represented by the percentage of

non-grassland cover in the landscape and fragmentation per se was represented by the first

principal component of the four landscape indices (mean grassland patch area, mean

nearest-neighbor distance among grassland patch, grassland patch density, grassland patch edge).

First, to investigate the impact of habitat fragmentation on grassland vascular plant diversity,

we used a linear regression model to estimate the effects of habitat loss and fragmentation per se

on grassland plant richness (including grassland specialist richness and weed richness).

Second, to investigate the relative importance of habitat fragmentation and grassland vascular

plant diversity on above-ground productivity, we used the multi-model averaging method based on

the Akaike information criterion corrected for a small sample size (AICc) (Harrison et al., 2018).

We first constructed linear regression models, including the response variable (above-ground biomass) and all combinations of the predictor variables (habitat loss, fragmentation per se, grassland specialist richness, and weed richness). The models with the lowest AICc value and a difference of less than two ΔAICc from the lowest AICc value were selected as optimal models. We then calculated the model-averaged standardized parameter estimate based on the optimal models as the relative effect of each influencing factor on above-ground biomass (Harrison et al., 2018). Before the analysis, we calculated the variance inflation factors (VIF) for each predictor variable to assess multicollinearity. The VIF of all explanatory variables was less than three (Appendix Table A2), suggesting no significant multicollinearity in the analysis (Carrara et al., 2015; Dormann et al., 2013).

Finally, we used the piecewise structural equation model to investigate how habitat fragmentation regulates the relationship between grassland plant diversity and above-ground productivity. We first constructed a hypothetical conceptual model (Figure 2). Based on previous studies, we hypothesized that habitat loss and fragmentation per se could decrease above-ground biomass directly and decrease plant richness indirectly (Allan et al., 2015; Haddad et al., 2015; Rolo et al., 2018). We also hypothesized that habitat loss and fragmentation per se could both decrease the impact of plant richness on above-ground biomass by decreasing the percentage of grassland specialists in the community (Clavel et al., 2011; Gravel et al., 2011; Liu et al., 2018). We included the interaction between the percentage of grassland specialists and the plant richness as predictor variables to quantify the moderating effect of the percentage of grassland specialists for the impact of plant richness on above-ground biomass. We tested the hypothetical conceptual model using a piecewise structural equation model (Lefcheck, 2016). The global fit of the model was evaluated using the Fisher's C statistic, and the hypothetical pathways were evaluated by standardized path coefficients. In addition, we used linear regression models to evaluate the relationship between plant richness and above-ground biomass at low, moderate, and high levels of habitat loss and fragmentation per se, respectively.

All data analyses were performed in the R programming language v. 4.0.3 (R Development Core Team, 2020) with the following functions and packages. The linear regression model and the Shapiro-Wilk normality test were conducted with 'lm' and 'shapiro.test' functions of the stats

package (R Development Core Team, 2020). The multi-model averaging were conducted with the

'dredge' and 'model.avg' functions of the MuMIn package (Bartoń, 2020). The VIF was

calculated with the 'vif' function of the car package (Fox and Weisberg, 2019). The piecewise

structural equation model was conducted and tested with the 'psem' function of the piecewiseSEM

package (Lefcheck, 2016).

Figure 2. Hypothetical model of habitat fragmentation that alters the relationship between

grassland plant diversity and above-ground biomass by affecting the percentage of grassland

specialists in the community in the Tabu River Basin, a typical agro-pastoral ecotone of northern

China. Arrows represent the hypothesized paths among variables. See the main text for a

description of arrow directions.

3 Results

3.1 Effect of habitat fragmentation on grassland plant richness

Habitat loss had a significant negative effect on overall species richness ($p < 0.05$) and

grassland specialist richness ($p < 0.01$), but a significant positive effect on weed richness ($p < 0.01$,

Figure 3a). Fragmentation per se had no significant effect on overall species richness and

grassland specialist richness, but a significant positive effect on weed richness ($p < 0.01$, Figure

3b).

Figure 3. Relationship between habitat fragmentation and grassland plant richness in the Tabu

River Basin, a typical agro-pastoral ecotone of northern China. The R^2 values in the figure are

from linear regression models. The n in the figure is the number of surveying sites used in the

linear regression models. * and ** represent significance at the 0.05 and 0.01 levels, respectively.

3.2 The relative impacts of habitat fragmentation and grassland plant richness on above-ground biomass

Habitat loss had no significant effect on above-ground biomass, and fragmentation per se had

a significant negative effect on above-ground biomass ($p < 0.05$). Grassland specialist richness

and weed richness had a significant positive effect on above-ground biomass ($p < 0.01$). The

effects of grassland specialist richness and weed richness on above-ground biomass were stronger

than fragmentation per se (Figure 4).

Figure 4. Standardized parameter estimates and 95% confidence intervals for habitat

fragmentation and plant richness affecting above-ground biomass in the Tabu River Basin, a

typical agro-pastoral ecotone of northern China. Standardized estimates and 95% confidence

intervals are calculated by the multi-model averaging method. * and ** represent significance at

267   the 0.05 and 0.01 levels, respectively.

3.3 The impact of habitat fragmentation on the relationship between

grassland plant richness and above-ground biomass

The Fisher's C statistic indicated that the piecewise structural equation model fitted the data

well (*Fisher's C*=5.5, *P*-value > 0.05, Figure 5). Habitat loss was shown by the piecewise

structural equation model to have no direct effect on above-ground biomass, and an indirect

negative effect on above-ground biomass through decreasing overall species richness.

Fragmentation per se had a direct negative effect on above-ground biomass and an indirect

positive effect on above-ground biomass through increasing overall species richness.

Meanwhile, the piecewise structural equation model showed that the percentage of grassland

specialists increased the effect of overall species richness on above-ground biomass (Figure 5).

Habitat loss decreased the effect of overall species richness on above-ground biomass by

decreasing the percentage of grassland specialists. Fragmentation per se had no significant effect

on the percentage of grassland specialists.

Figure 5. Results from piecewise structural equation model with hypothesized paths showing

283   how habitat fragmentation alters the relationship between grassland plant richness and

284   above-ground biomass in the Tabu River Basin, a typical agro-pastoral ecotone of northern China.

285   The widths of arrows are proportional to the standardized path coefficients. Black and red

286   solid arrows represent the positive and negative effects, respectively. Black solid and grey dashed

287   arrows represent the significant and nonsignificant effects at the 0.05 level, respectively. The

288   Fisher's C and P-values are from the piecewise structural equation model.

Results of the linear regression showed that there was a positive relationship between plant

richness and above-ground biomass at low, moderate, and high levels of habitat loss (Figure 6).

The positive relationship was strongest at low levels of habitat loss and weakest at high levels of

habitat loss.

Figure 6. Relationships between grassland plant richness and above-ground biomass at low,

moderate, and high levels of habitat loss from 130 landscapes in the Tabu River Basin, a typical

agro-pastoral ecotone of northern China. The R^2 values in the figure are from linear regression

298   models. ** represent significance at the 0.01 level.

4 Discussion

4.1 Habitat loss and fragmentation per se had inconsistent effects on grassland plant diversity and ecosystem function

302   Previous studies investigating the influence of habitat fragmentation on biodiversity and

303 ecosystem function mainly conducted at the patch level generally indicate the negative effects of

habitat fragmentation on multitrophic biodiversity and multiple ecosystem function (Haddad et al., 2015; Hertzog et al., 2019; Rolo et al., 2018). However, habitat fragmentation is often considered a landscape-wide phenomenon ideally to be understood at the landscape level (Fahrig, 2017; McGarigal and Cushman, 2002). We evaluated the effects of two facets of habitat fragmentation at the landscape scale (habitat loss and fragmentation per se) on grassland plant diversity and above-ground productivity in the agro-pastoral ecotone of northern China, with our results showing the effects of these two facets to not be consistent.

Consistent with previous studies, we found habitat loss significantly reduced grassland plant diversity, suggesting that habitat loss is a major threat to the biodiversity conservation of fragmented landscapes in this region (Chase et al., 2020; Haddad et al., 2015; Ibáñez et al., 2014). While for fragmentation per se, we found a positive effect on grassland plant diversity, in accordance with some recent evidence the effects of fragmentation per se on biodiversity are more likely positive than negative (Gestich et al., 2022; Palmeirim et al., 2019; Riva and Fahrig, 2023). This is mainly because when the amount of habitat in the landscape is not reduced fragmentation per se may promote biodiversity through increasing habitat diversity, increasing landscape connectivity, reducing competition, and other mechanisms (Fahrig, 2017). Our study suggests that biodiversity conservation strategies for fragmented landscapes should consider optimizing the habitat configuration in the landscape in addition to preventing habitat loss, such as increasing the number of habitat patches (Arroyo-Rodríguez et al., 2020).

Our results showed for a metric of ecosystem function that fragmentation per se significantly decreased above-ground productivity, whereas habitat loss did not. A possible reason for this finding is fragmentation per se could directly alter environmental factors affecting ecosystem function through enhanced edge effects (Laurance et al., 2011; Smith et al., 2018). The negative effect of fragmentation per se on above-ground productivity in our study may be due to increased drought severity caused by edge effects (Brookshire and Weaver, 2015). Previous studies showed edge effects could lead to ecosystem drought by causing greater desiccation and evapotranspiration rate, which is a major factor limiting productivity in grassland ecosystems (Smith et al., 2018; Tuff et al., 2016). Our study, generally, suggests habitat loss and fragmentation per se have inconsistent effects on plant diversity and ecosystem function in fragmented

333 landscapes. Future studies should simultaneously consider habitat loss and fragmentation per se to

334 understand the consequences of habitat fragmentation.

4.2 Habitat fragmentation did not change the direction of the positive relationship between plant diversity and ecosystem function, but weakened the magnitude of the relationship

Understanding the direction and magnitude of BEF relationships in fragmented landscapes is

essential to understand the importance of biodiversity for ecosystem function in the changing

world (Gonzalez et al., 2020; van der Plas, 2019). In naturally assembled communities, ecosystem

functions may be dominated by complex environmental factors and landscape context, showing a

weak or even negative correlation with biodiversity (Grace et al., 2007; Hagan et al., 2021; Zirbel

et al., 2019). Our study found grassland plant diversity showed a significant positive correlation

with aboveground productivity and its relative importance was stronger than that of habitat

fragmentation. This result is consistent with findings from BEF experiments (Fanin et al., 2018;

Hong et al., 2022), indicating habitat fragmentation did not change the direction of the positive

BEF relationship in this region. Thus, our study shows grassland plant diversity has an important

role in maintaining grassland ecosystem functions in the fragmented landscapes of the

agro-pastoral ecotone of northern China.

Although habitat fragmentation did not change the direction of the positive BEF relationship,

we found it could weaken the magnitude of the relationship through habitat loss. This is consistent

with predictions of theoretical models of habitat fragmentation affecting the relationship between

biodiversity and ecosystem function (Clavel et al., 2011; Liu et al., 2018). Our finding suggests

habitat loss could reduce the importance of plant diversity to ecosystem function in fragmented

landscapes of the agro-pastoral ecotone of northern China. However, a recent study by Hertzog et

al. (2019) on temperate forest ecosystems found that habitat fragmentation could strengthen the

positive relationship between plant diversity and ecosystem multifunctionality. This inconsistency

can be explained by trade-offs between different ecosystem functions that may differ in their

response to habitat fragmentation (Banks-Leite et al., 2020). Therefore, future studies are needed

to focus on multiple ecosystem functions and consider their potential trade-offs.

We found habitat loss weakened the positive BEF relationship by decreasing the percentage

of specialists in grassland communities, as specialists were more vulnerable to the negative effect

of habitat loss and more associated with ecosystem function than generalists. These findings

indicate reducing the degree of habitat specialization may be the mechanism of habitat

fragmentation altering the BEF relationship in the fragmented landscapes of this region (Clavel et

al., 2011; Gravel et al., 2011). Thus, our study provides evidence that habitat fragmentation could

decrease the degree of habitat specialization by leading to the replacement of specialists by

generalists in the community, thus weakening the BEF relationship (Clavel et al., 2011). Our

findings suggest further expansion of farmland in the agro-pastoral ecotone of northern China

would decrease the grassland plant diversity and its importance to ecosystem function.

371 5 Conclusions

Our study investigated the effects of two aspects of habitat fragmentation (habitat loss and

fragmentation per se) on grassland plant diversity, above-ground productivity, and their

relationships in the agro-pastoral ecotone of northern China. We found habitat loss and

fragmentation per se had inconsistent effects. Habitat loss decreased grassland plant diversity,

while fragmentation per se increased grassland plant diversity. Habitat loss did not affect

above-ground biomass, while fragmentation per se decreased above-ground biomass. We also

found habitat fragmentation did not change the direction of the positive relationship between

grassland plant diversity and above-ground biomass, but weakened the magnitude of the

relationship by decreasing the habitat specialization of the community. Our study suggests

biodiversity conservation in fragmented landscapes should focus on preventing habitat loss, which

decreases plant diversity and its importance to ecosystem function.

383 ACKNOWLEDGMENTS

384   We are very grateful to Peng Han, Shuangshuang Zhang, Xiaoqian Gong, Fengshi Li, Nier Su,

Luyao Liu, Linjie Yao, and Guofeng Shi for assistance with field work. This study was supported by the National Natural Science Foundation of China (32071582) and the Inner Mongolia Autonomous Region Science and Technology Plan Project (2021GG0392). No permits were required for fieldwork in this research.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

DATA ACCESSIBILITY STATEMENT

We have unloaded the data as supplementary materials. If the manuscript is accepted, we will upload the data to a publicly accessible digital repository.

AUTHORS' CONTRIBUTIONS

Q.Z. and Y.Z. conceived the research; Y.Z., S.J., and Q.Z. contributed to the writing of the manuscript; Y.Z. and S.J. contributed to data compilation and analysis; Y.Z. and Q.Z. contributed to the fieldwork.

References

Allan, E., Manning, P., Alt, F., Binkenstein, J., Blaser, S., Blüthgen, N., et al., 2015. Land use intensification alters ecosystem multifunctionality via loss of biodiversity and changes to functional composition. Ecology Letters 18, 834-843. https://doi.org/10.1111/ele.12469

Arroyo-Rodríguez, V., Fahrig, L., Tabarelli, M., Watling, J.I., Tischendorf, L., Benchimol, M., et al., 2020. Designing optimal human-modified landscapes for forest biodiversity conservation. Ecology Letters 23, 1404-1420. https://doi.org/10.1111/ele.13535

Banks-Leite, C., Ewers, R.M., Folkard-Tapp, H., Fraser, A., 2020. Countering the effects of habitat loss,
fragmentation, and degradation through habitat restoration. One Earth 3, 672-676.
https://doi.org/10.1016/j.oneear.2020.11.016
Bartoń, K., 2020. MuMIn: multi-model inference. R package version 1.43.17.
https://CRAN.R-project.org/package=MuMIn
Brookshire, E.N.J., Weaver, T., 2015. Long-term decline in grassland productivity driven by increasing
dryness. Nature Communications 6, 7148. https://doi.org/10.1038/ncomms8148
Carrara, E., Arroyo-Rodríguez, V., Vega-Rivera, J.H., Schondube, J.E., de Freitas, S.M., Fahrig, L.,
2015. Impact of landscape composition and configuration on forest specialist and generalist
bird species in the fragmented Lacandona rainforest, Mexico. Biological Conservation 184,
117-126. https://doi.org/10.1016/j.biocon.2015.01.014
Chase, J.M., Blowes, S.A., Knight, T.M., Gerstner, K., May, F., 2020. Ecosystem decay exacerbates
biodiversity loss with habitat loss. Nature 584, 238-243.
https://doi.org/10.1038/s41586-020-2531-2
Clavel, J., Julliard, R., Devictor, V., 2011. Worldwide decline of specialist species: toward a global
functional homogenization? Frontiers in Ecology and the Environment 9, 222-228.
https://doi.org/10.1890/080216
Dehling, D.M., Bender, I.M.A., Blendinger, P.G., Böhning-Gaese, K., Muñoz, M.C., Neuschulz, E.L.,
et al., 2021. Specialists and generalists fulfil important and complementary functional roles in
ecological processes. Functional Ecology 35, 1810-1821.
https://doi.org/10.1111/1365-2435.13815
Devictor, V., Julliard, R., Jiguet, F., 2008. Distribution of specialist and generalist species along spatial
gradients of habitat disturbance and fragmentation. Oikos 117, 507-514.
https://doi.org/10.1111/j.0030-1299.2008.16215.x
Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., et al., 2013. Collinearity: a
review of methods to deal with it and a simulation study evaluating their performance.
Ecography 36, 27-46. https://doi.org/10.1111/j.1600-0587.2012.07348.x
Fahrig, L., 2003. Effects of habitat fragmentation on biodiversity. Annual Review of Ecology Evolution
and Systematics 34, 487-515. https://doi.org/10.1146/annurev.ecolsys.34.011802.132419
Fahrig, L., 2017. Ecological responses to habitat fragmentation per se. Annual Review of Ecology,
Evolution, and Systematics 48, 1-23. https://doi.org/10.1146/annurev-ecolsys-110316-022612
Fahrig, L., Arroyo-Rodríguez, V., Bennett, J.R., Boucher-Lalonde, V., Cazetta, E., Currie, D.J., et al.,
2019. Is habitat fragmentation bad for biodiversity? Biological Conservation 230, 179-186.
https://doi.org/10.1016/j.biocon.2018.12.026
Fanin, N., Gundale, M.J., Farrell, M., Ciobanu, M., Baldock, J.A., Nilsson, M.-C., et al., 2018.
Consistent effects of biodiversity loss on multifunctionality across contrasting ecosystems.
Nature Ecology & Evolution 2, 269-278. https://doi.org/10.1038/s41559-017-0415-0
Fardila, D., Kelly, L.T., Moore, J.L., McCarthy, M.A., 2017. A systematic review reveals changes in
where and how we have studied habitat loss and fragmentation over 20years. Biological
Conservation 212, 130-138. https://doi.org/10.1016/j.biocon.2017.04.031
Fletcher Jr, R.J., Smith, T.A.H., Kortessis, N., Bruna, E.M., Holt, R.D., 2023. Landscape experiments
unlock relationships among habitat loss, fragmentation, and patch-size effects. Ecology 104,
e4037. https://doi.org/10.1002/ecy.4037
Fletcher, R.J., Didham, R.K., Banks-Leite, C., Barlow, J., Ewers, R.M., Rosindell, J., et al., 2018. Is

habitat fragmentation good for biodiversity? Biological Conservation 226, 9-15.

https://doi.org/10.1016/j.biocon.2018.07.022

Fox, J., Weisberg, S., 2019. An R companion to Applied Regression. Sage.

https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Gestich, C.C., Arroyo-Rodríguez, V., Saranholi, B.H., da Cunha, R.G.T., Setz, E.Z.F., Ribeiro, M.C.,

2022. Forest loss and fragmentation can promote the crowding effect in a forest-specialist

primate. Landscape Ecology 37, 147-157. https://doi.org/10.1007/s10980-021-01336-1

Godbold, J.A., Bulling, M.T., Solan, M., 2011. Habitat structure mediates biodiversity effects on

ecosystem properties. Proceedings of the Royal Society B: Biological Sciences 278,

2510-2518. https://doi.org/10.1098/rspb.2010.2414

Gonzalez, A., Germain, R.M., Srivastava, D.S., Filotas, E., Dee, L.E., Gravel, D., et al., 2020.

Scaling-up biodiversity-ecosystem functioning research. Ecology Letters 23, 757-776.

https://doi.org/10.1111/ele.13456

Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., Moore, R., 2017. Google Earth

Engine: planetary-scale geospatial analysis for everyone. Remote Sensing of Environment 202,

18-27. https://doi.org/10.1016/j.rse.2017.06.031

Grace, J.B., Michael Anderson, T., Smith, M.D., Seabloom, E., Andelman, S.J., Meche, G., et al., 2007.

Does species diversity limit productivity in natural grassland communities? Ecology Letters

10, 680-689. https://doi.org/10.1111/j.1461-0248.2007.01058.x

Gravel, D., Bell, T., Barbera, C., Bouvier, T., Pommier, T., Venail, P., et al., 2011. Experimental niche

evolution alters the strength of the diversity–productivity relationship. Nature 469, 89-92.

https://doi.org/10.1038/nature09592

Haddad, N.M., Brudvig, L.A., Clobert, J., Davies, K.F., Gonzalez, A., Holt, R.D., et al., 2015. Habitat

fragmentation and its lasting impact on Earth's ecosystems. Science Advances 1, e1500052.

https://doi.org/10.1126/sciadv.1500052

Hagan, J.G., Vanschoenwinkel, B., Gamfeldt, L., 2021. We should not necessarily expect positive

relationships between biodiversity and ecosystem functioning in observational field data.

Ecology Letters 24, 2537-2548. https://doi.org/10.1111/ele.13874

Harrison, X.A., Donaldson, L., Correa-Cano, M.E., Evans, J., Fisher, D.N., Goodwin, C.E., et al., 2018.

A brief introduction to mixed effects modelling and multi-model inference in ecology. PeerJ 6,

e4794. https://doi.org/10.7717/peerj.4794

Hertzog, L.R., Boonyarittichaikij, R., Dekeukeleire, D., de Groote, S.R.E., van Schrojenstein Lantman,

I.M., Sercu, B.K., et al., 2019. Forest fragmentation modulates effects of tree species richness

and composition on ecosystem multifunctionality. Ecology 100, e02653.

https://doi.org/10.1002/ecy.2653

Hong, P., Schmid, B., De Laender, F., Eisenhauer, N., Zhang, X., Chen, H., et al., 2022. Biodiversity

promotes ecosystem functioning despite environmental change. Ecology Letters 25, 555-569.

https://doi.org/10.1111/ele.13936

Ibáñez, I., Katz, D.S.W., Peltier, D., Wolf, S.M., Connor Barrie, B.T., 2014. Assessing the integrated

effects of landscape fragmentation on plants and plant communities: the challenge of

multiprocess–multiresponse dynamics. Journal of Ecology 102, 882-895.

https://doi.org/10.1111/1365-2745.12223

IPBES, 2019. Global assessment report on biodiversity and ecosystem services of the

intergovernmental science-policy platform on biodiversity and ecosystem services. Bonn:

IPBES. https://doi.org/10.5281/zenodo.5657041

Laurance, W.F., Camargo, J.L.C., Luizão, R.C.C., Laurance, S.G., Pimm, S.L., Bruna, E.M., et al., 2011.
The fate of Amazonian forest fragments: a 32-year investigation. Biological Conservation 144,
56-67. https://doi.org/10.1016/j.biocon.2010.09.021

Lefcheck, J.S., 2016. PIECEWISESEM: piecewise structural equation modelling in R for ecology,
evolution, and systematics. Methods in Ecology and Evolution 7, 573-579.
https://doi.org/10.1111/2041-210x.12512

Liu, J., Wilson, M., Hu, G., Liu, J., Wu, J., Yu, M., 2018. How does habitat fragmentation affect the
biodiversity and ecosystem functioning relationship? Landscape Ecology 33, 341-352.
https://doi.org/10.1007/s10980-018-0620-5

Matthews, T.J., 2021. On the biogeography of habitat islands: the importance of matrix effects, noncore
species, and source-sink dynamics. The Quarterly Review of Biology 96, 73-104.
https://doi.org/10.1086/714482

Matthews, T.J., Cottee-Jones, H.E., Whittaker, R.J., 2014. Habitat fragmentation and the species–area
relationship: a focus on total species richness obscures the impact of habitat loss on habitat
specialists. Diversity and Distributions 20, 1136-1146. https://doi.org/10.1111/ddi.12227

Maxwell, S.L., Fuller, R.A., Brooks, T.M., Watson, J.E.M., 2016. Biodiversity: the ravages of guns,
nets and bulldozers. Nature 536, 143-145. https://doi.org/10.1038/536143a

McGarigal, K., Cushman, S.A., 2002. Comparative evaluation of experimental approaches to the study
of habitat fragmentation effects. Ecological Applications 12, 335-345.
https://doi.org/10.1890/1051-0761(2002)012[0335:CEOEAT]2.0.CO;2

McGarigal, K., Cushman, S.A., Ene, E., 2012. FRAGSTATS v4: spatial pattern analysis program for
categorical and continuous maps. Computer software program produced by the authors at the
University of Massachusetts, Amherst. Available at the following web site:
http://www.umass.edu/landeco/research/fragstats/fragstats.html.

Mello, M.A.R., Rodrigues, F.A., Costa, L.d.F., Kissling, W.D., Şekercioğlu, Ç.H., Marquitti, F.M.D., et
al., 2015. Keystone species in seed dispersal networks are mainly determined by dietary
specialization. Oikos 124, 1031-1039. https://doi.org/10.1111/oik.01613

Miller-Rushing, A.J., Primackh, R.B., Devictor, V., Corlett, R.T., Cumming, G.S., Loyola, R., et al.,
2019. How does habitat fragmentation affect biodiversity? a controversial question at the core
of conservation biology. Biological Conservation 232, 271-273.
https://doi.org/10.1016/j.biocon.2018.12.029

Palmeirim, A.F., Figueiredo, M.S.L., Grelle, C.E.V., Carbone, C., Vieira, M.V., 2019. When does
habitat fragmentation matter? a biome-wide analysis of small mammals in the Atlantic Forest.
Journal of Biogeography 46, 2811-2825. https://doi.org/10.1111/jbi.13730

Pasher, J., Mitchell, S.W., King, D.J., Fahrig, L., Smith, A.C., Lindsay, K.E., 2013. Optimizing
landscape selection for estimating relative effects of landscape variables on ecological
responses. Landscape Ecology 28, 371-383. https://doi.org/10.1007/s10980-013-9852-6

R Development Core Team, 2020. R: a language and environment for statistical computing. R
Foundation for Statistical Computing.

Reynolds, C., Fletcher, R.J., Carneiro, C.M., Jennings, N., Ke, A., LaScaleia, M.C., et al., 2018.
Inconsistent effects of landscape heterogeneity and land-use on animal diversity in an
agricultural mosaic: a multi-scale and multi-taxon investigation. Landscape Ecology 33,
241-255. https://doi.org/10.1007/s10980-017-0595-7

Riva, F., Fahrig, L., 2023. Landscape-scale habitat fragmentation is positively related to biodiversity, despite patch-scale ecosystem decay. Ecology Letters 26, 268-277. https://doi.org/10.1111/ele.14145

Rolo, V., Olivier, P.I., Pfeifer, M., van Aarde, R.J., 2018. Functional diversity mediates contrasting direct and indirect effects of fragmentation on below- and above-ground carbon stocks of coastal dune forests. Forest Ecology and Management 407, 174-183. https://doi.org/10.1016/j.foreco.2017.10.059

Smith, A.C., Koper, N., Francis, C.M., Fahrig, L., 2009. Confronting collinearity: comparing methods for disentangling the effects of habitat loss and fragmentation. Landscape Ecology 24, 1271-1285. https://doi.org/10.1007/s10980-009-9383-3

Smith, I.A., Hutyra, L.R., Reinmann, A.B., Marrs, J.K., Thompson, J.R., 2018. Piecing together the fragments: elucidating edge effects on forest carbon dynamics. Frontiers in Ecology and the Environment 16, 213-221. https://doi.org/10.1002/fee.1793

Tilman, D., Reich, P.B., Isbell, F., 2012. Biodiversity impacts ecosystem productivity as much as resources, disturbance, or herbivory. Proceedings of the National Academy of Sciences 109, 10394-10397. https://doi.org/10.1073/pnas.1208240109

Tuff, K.T., Tuff, T., Davies, K.F., 2016. A framework for integrating thermal biology into fragmentation research. Ecology Letters 19, 361-374. https://doi.org/10.1111/ele.12579

van der Plas, F., 2019. Biodiversity and ecosystem functioning in naturally assembled communities. Biological Reviews 94, 1220-1245. https://doi.org/10.1111/brv.12499

Wang, X., Blanchet, F.G., Koper, N., 2014. Measuring habitat fragmentation: an evaluation of landscape pattern metrics. Methods in Ecology and Evolution 5, 634-646. https://doi.org/10.1111/2041-210X.12198

Wardle, D.A., Zackrisson, O., 2005. Effects of species and functional group loss on island ecosystem properties. Nature 435, 806-810. https://doi.org/10.1038/nature03611

Wilson, M.C., Chen, X.-Y., Corlett, R.T., Didham, R.K., Ding, P., Holt, R.D., et al., 2016. Habitat fragmentation and biodiversity conservation: key findings and future challenges. Landscape Ecology 31, 219-227. https://doi.org/10.1007/s10980-015-0312-3

Yan, Y., Jarvie, S., Liu, Q., Zhang, Q., 2022. Effects of fragmentation on grassland plant diversity depend on the habitat specialization of species. Biological Conservation 275, 109773. https://doi.org/10.1016/j.biocon.2022.109773

Yan, Y., Jarvie, S., Zhang, Q., Han, P., Liu, Q., Zhang, S., et al., 2023. Habitat heterogeneity determines species richness on small habitat islands in a fragmented landscape. Journal of Biogeography 50, 976-986. https://doi.org/10.1111/jbi.14594

Yan, Y., Jarvie, S., Zhang, Q., Zhang, S., Han, P., Liu, Q., et al., 2021. Small patches are hotspots for biodiversity conservation in fragmented landscapes. Ecological Indicators 130, 108086. https://doi.org/10.1016/j.ecolind.2021.108086

Yang, Y., Wang, K., Liu, D., Zhao, X., Fan, J., 2020. Effects of land-use conversions on the ecosystem services in the agro-pastoral ecotone of northern China. Journal of Cleaner Production 249, 119360. https://doi.org/10.1016/j.jclepro.2019.119360

Zhang, S., Zhang, Q., Yan, Y., Han, P., Liu, Q., 2021. Island biogeography theory predicts plant species richness of remnant grassland patches in the agro-pastoral ecotone of northern China. Basic and Applied Ecology 54, 14-22. https://doi.org/10.1016/j.baae.2021.04.010

Zirbel, C.R., Grman, E., Bassett, T., Brudvig, L.A., 2019. Landscape context explains ecosystem

582   multifunctionality in restored grasslands better than plant diversity. Ecology 100, e02634.

583   https://doi.org/10.1002/ecy.2634

Overall species · Grassland specialists · Weeds

(a)
$R^2 = 0.05^*$, n = 130
$R^2 = 0.17^{**}$, n = 130
$R^2 = 0.10^{**}$, n = 130

Species richness

Habitat loss (%)

(b)
$R^2 = 0.00$, n = 130
$R^2 = 0.03$, n = 130
$R^2 = 0.07^{**}$, n = 130

Species richness

Fragmentation per se (PC1)

Fisher's C = 5.5, *P*-value = 0.48