# Habitat loss weakens the positive relationship between grassland plant richness and above-ground biomass

Yongzhi Yan[1], Scott Jarvie[2], Qing Zhang[1, 3, 4*]

[1]Ministry of Education Key Laboratory of Ecology and Resource Use of the Mongolian Plateau, School of Ecology and Environment, Inner Mongolia University, Hohhot 010021, China

[2]Otago Regional Council, Dunedin 9016, New Zealand

[3]Collaborative Innovation Center for Grassland Ecological Security (Jointly Supported by the Ministry of Education of China and Inner Mongolia Autonomous Region), Hohhot, 010021, China

[4]Autonomous Region Collaborative Innovation Center for Integrated Management of Water Resources and Water Environment in the Inner Mongolia Reaches of the Yellow River, Hohhot 010018, China

*Correspondence:

Qing Zhang

Email: qzhang82@163.com

# Highlights:

1. BEF relationship is moderated by landscape context in fragmented landscapes.

2. Habitat loss and fragmentation per se have inconsistent effects.

3. Habitat loss can weaken the positive BEF relationship via reducing the percentage of grassland specialists in the community.

**Abstract:**

Habitat loss and fragmentation per se have been shown to be a major threat to global biodiversity and ecosystem function. However, little is known about how habitat loss and fragmentation per se alters the relationship between biodiversity and ecosystem function (BEF relationship) in the natural landscape context. Based on 130 landscapes identified by a stratified random sampling in the agro-pastoral ecotone of northern China, we investigated the effects of landscape context (habitat loss and fragmentation per se) on plant richness, above-ground biomass, and the relationship between them in grassland communities using a structural equation model. We found that habitat loss directly decreased plant richness and hence decreased above-ground biomass, while fragmentation per se directly increased plant richness and hence increased above-ground biomass. Fragmentation per se also directly decreased soil water content and hence decreased above-ground biomass. Meanwhile, habitat loss decreased the magnitude of the positive relationship between plant richness and above-ground biomass by reducing the percentage of grassland specialists in the community, while fragmentation per se had no significant modulating effect on this relationship. These results demonstrate that habitat loss and fragmentation per se have inconsistent effects on biodiversity and ecosystem function, with the BEF relationship being modulated by landscape context. Our findings emphasise that habitat loss rather than fragmentation per se can weaken the positive BEF relationship by decreasing the degree of habitat specialisation of the community.

Keywords:

BEF relationship, biodiversity, ecosystem function, fragmentation per se, generalists, habitat loss, modulating effect, specialists.

# 1 Introduction

Evidence from biodiversity-ecosystem function (BEF hereafter) experiments during the past 30 years generally show positive relationships between biodiversity and productivity, soil carbon storage, decomposition rates, and other ecosystem functions in experimental communities, revealing the importance of biodiversity in maintaining ecosystem functioning (*Tilman et al., 2012; van der Plas, 2019*). When research expands from experiments to natural systems, however, BEF

relationships remain unclear in the natural assembled communities, with significant context

dependency (*Hagan et al., 2021; van der Plas, 2019; but see Duffy et al., 2017*). One of the main

reasons for these differences is because the landscape context surrounding natural communities

regulates biodiversity and ecosystem function (*Gonzalez et al., 2020; Liu et al., 2018*).

Consideration of the impacts of landscape context on surrounding communities is key to

understand complex BEF relationships in natural systems.

Human activities have modified natural ecosystems globally, with fragmented landscape

contexts becoming increasingly widespread around the world (*Chase et al., 2020; Maxwell et al.,*

*2016*). For example, at least 1.5 billion hectares of natural habitats on Earth have been converted

to human-modified land since 2014, breaking apart continuous habitat into smaller and isolated

fragments (*IPBES, 2019*). Fragmented landscape context typically involves different processes,

such as habitat loss, i.e., reducing habitat amount in the landscape, and fragmentation per se, i.e.,

breaking apart of habitat for a given habitat amount in the landscape, including decreased mean

size of habitat patches, increased number of habitat patches, increased isolation among habitat

patches, etc. (*Fahrig, 2003; Ibáñez et al., 2014; Wang et al., 2014*). Habitat loss is often

considered the major near-term threat to the biodiversity of terrestrial ecosystems (*Chase et al.,*

*2020; Haddad et al., 2015*), while the impact of fragmentation per se remains debated (*Fletcher Jr*

*et al., 2023; Miller-Rushing et al., 2019*). Thus, habitat loss and fragmentation per se may have

inconsistent ecological consequences and should be considered simultaneously to establish

effective conservation strategies in fragmented landscapes (*Fahrig et al., 2019; Fletcher et al.,*

*2018; Miller-Rushing et al., 2019*).

Fragmented landscape context can also affect BEF relationships. Previous studies have found

that the magnitude and direction of BEF relationships vary across fragmented landscapes,

including positive, negative, and non-significant relationships (*Godbold et al., 2011; Hagan et al.,*

*2021; Rolo et al., 2018; Zirbel et al., 2019*). However, few studies have been conducted on how

habitat loss and fragmentation per se regulate BEF relationships (*Liu et al., 2018; Wilson et al.,*

*2016*). This lack of studies hampers our understanding of complex BEF relationships in

fragmented natural ecosystems. In theory, habitat loss and fragmentation per se can regulate

ecosystem function and the BEF relationship by altering species composition, interactions, and

spatial asynchrony regardless of changes in species richness (*Liu et al., 2018; Thompson and Gonzalez, 2016; Tscharntke et al., 2012*). This is because species in communities are not ecologically equivalent and may respond differently to habitat loss and fragmentation per se, and contribute unequally to ecosystem function (*Devictor et al., 2008; Wardle and Zackrisson, 2005*). Therefore, considering changes in species composition can help to understand how habitat loss and fragmentation per se regulate BEF relationships.

The degree of habitat specialisation is a key ecological characteristic determining responses of species to habitat loss and fragmentation per se and the slope of BEF relationships (*Clavel et al., 2011; Gravel et al., 2011*). In fragmented landscapes, species with low levels of habitat specialisation (generalists) can use resources from different land covers, including focal habitat and non-habitat matrix, and thus are not sensitive to habitat loss and fragmentation per se (*Matthews et al., 2014*). Conversely, species with high levels of habitat specialisation (specialists) depend highly on resources in specific habitats, and thus are vulnerable to adverse effects from habitat loss and fragmentation per se (*Matthews et al., 2014*). In communities, specialists with specialised niches in resource use may contribute complementary roles to ecosystem functioning, whereas generalists with unspecialised niches in resource use may contribute redundant roles to ecosystem functioning due to overlapping niches (*Dehling et al., 2021; Denelle et al., 2020; Gravel et al., 2011; Wilsey et al., 2023*). Therefore, communities composed of specialists should have a higher niche complementarity effect in maintaining ecosystem functions and a more significant BEF relationship than communities composed of generalists. Habitat loss and fragmentation per se are often predicted to decrease the degree of habitat specialisation (the replacement of specialists in communities by generalists), possibly resulting in functional homogenisation of communities and reduced BEF relationships (*Clavel et al., 2011; Matthews, 2021*). However, few studies have evaluated this process in fragmented landscapes.

Currently, research on habitat loss and fragmentation per se focuses primarily on forest ecosystems (*Fardila et al., 2017; Haddad et al., 2015; Ma et al., 2023*). Grasslands have received considerably less attention, despite being one of the largest terrestrial ecosystems, and suffering severe fragmentation due to human activities, such as agricultural reclamation and urbanisation (*Fardila et al., 2017*). The agro-pastoral ecotone of northern China is a typical anthropogenically

fragmented grassland landscape caused by historical agricultural reclamation, especially in the late

Qing Dynasty (about 1840–1912). Due to land policy reforms, the region has experienced a rapid

expansion of farmland since the 1960s, converting continuous natural grasslands into smaller and

isolated fragments, seriously threatening the conservation of biodiversity and ecosystem functions

(*Yan et al., 2022; Yan et al., 2023; Yang et al., 2020*). Based on 130 landscapes with different

fragmentation levels in the agro-pastoral ecotone of northern China, we investigated how

fragmented landscape context (habitat loss and fragmentation per se) impact the relationship

between grassland plant diversity and above-ground productivity in the community. Specifically,

we aimed to evaluate whether habitat loss and fragmentation per se would weaken the positive

relationship between grassland plant diversity and above-ground productivity by reducing the

habitat specialisation of the community.

# 2 Materials and methods

## 2.1 Study area

Our study area is in the agro-pastoral ecotone of northern China, the Tabu River Basin in

Siziwang Banner, Inner Mongolia Autonomous Region. The mean annual temperature ranges from

1.5 to 5.0 □, and the mean annual precipitation ranges from 225 to 322 mm. The type of soil is

light chestnut soil. This area is a typical fragmented grassland landscape caused by agricultural

intensification. Grassland is the dominant natural habitat type in this area, accounting for about

40.8 % of the total area, with the dominant plant species being *Stipa krylovii* and S. *breviflora*.

Farmlandd is the dominant matrix type in this area, accounting for about 30.6 % of the total area,

with the main crops grown being potatoes and maize. Further background information about the

study area is described in our previous papers (*Yan et al., 2022; Yan et al., 2023; Yan et al., 2021;*

*Zhang et al., 2021*).

## 2.2 Sampling landscape selection

We quantified landscape-scale habitat loss and fragmentation per se in the study area to

determine the spatial gradient of landscape context, then established the sampling landscapes.

Grassland was defined as the focal habitat. Habitat amount was represented by the percentage of

grassland cover in the landscape. Habitat loss was represented by the loss of grassland amount in

the landscape. As the remaining grassland fragments in this region were mainly caused by

grassland loss due to human activities such as cropland expansion (*Chen et al., 2019; Yang et al.,*

*2020*), the percentage of non-grassland cover in the landscape was used in our study to represent

habitat loss. Fragmentation per se was estimated by calculating four landscape indices that reflect

the different fragmentation processes for a given amount of habitat in the landscape (*Fahrig, 2003;*

*Fahrig, 2017*): (1) patch density metric, representing an increase in the number of grassland

patches in the landscape; (2) edge density metric, representing an increase in the grassland edges

in the landscape; (3) mean patch area metric, representing a decrease in the mean size of grassland

patches in the landscape; and (4) mean nearest-neighbor distance metric, representing an increase

in the isolation among grassland patches in the landscape. The patch density metric reflects the

breaking apart of habitat in the landscape, which is a direct reflection of the definition of

fragmentation per se (*Fahrig et al., 2019*). The edge density metric reflects the magnitude of the

edge effect caused by fragmentation (*Fahrig, 2017*). The mean patch area metric and the mean

nearest-neighbor distance metric are associated with the area and distance effects of island

biogeography, respectively, reflecting the processes of local extinction and dispersal of species in

the landscape (*Fletcher et al., 2018*).

Given that habitat amount (or habitat loss) and fragmentation per se are typically highly

correlated in natural landscapes, it is hard to disentangle their relative effects (*Fahrig, 2017; Smith*

*et al., 2009*). We therefore used a quasi-experimental method to select sampling landscapes across

the relative independent spatial gradient of habitat amount and fragmentation per se, which can

reduce collinearity between them (*Butsic et al., 2017; Pasher et al., 2013; Reynolds et al., 2018*).

To do so, we first used the moving window method (window size: 500-m radius buffer) to

quantify grassland amount and the four landscape indices (patch density, edge density, mean patch

area, and mean nearest-neighbor distance metric) surrounding all grassland cells. The 500-m

radius buffer was used because our previous studies showed this buffer includes the optimal scale

of spatial processes influencing grassland plant diversity in this region (*Yan et al., 2022; Yan et al.,*

*2023; Zhang et al., 2021).* Second, to quantify the fragmentation per se level, we derived the first

principal component (PC1) of the four landscape indices (*Hertzog et al., 2019; Rolo et al., 2018*).

We took the inverse of the PC1 as a single fragmentation per se index, which was positively

correlated with patch density, edge density, mean nearest-neighbor distance metric, and negatively

with mean patch area (Appendix Figure A1 and Table A1).

Based on the quartiles of grassland amount and single fragmentation per se index, we ranked

these grassland cells into nine types: high-high, high-moderate, high-low, moderate-high,

moderate-moderate, moderate-low, low-high, low-moderate, low-low grassland amount and

fragmentation per se (Appendix Figure A2). Given landscapes with a high grassland amount and

high fragmentation are scarce in this region (Appendix Figure A2), we did not consider this type

of landscape. Finally, we selected at least 20 grassland landscapes with a minimum distance

condition using stratified sampling from each of the remaining eight grassland types as alternative

sites for field surveys. The minimum distance between each landscape was at least 1000 m to

prevent overlapping landscapes and potential spatial autocorrelation.

The land-cover data used to quantify grassland fragmentation was obtained via supervised

classification on a cloud-free Landsat 8 TOA composite image (30-m resolution) from 2019 (*Yan*

*et al., 2022*). We used the random forest classifier in the Google Earth Engine platform (*Gorelick*

*et al., 2017*) for the supervised classification. The overall classification accuracy was 84.3 %, and

the kappa coefficient was 0.81. The moving window analysis and all landscape metric calculations

were performed in FRAGSTATS v4.2.1 based on the eight-cell neighborhood rule (*McGarigal et*

*al., 2012*). The principal component analysis was performed in the R programming language v.

4.0.3 (*R Development Core Team, 2020*), and stratified sampling was conducted in ArcGIS v10.3.

### 183  2.3 Biodiversity and ecosystem function surveys

Based on the alternative sites selected above, we established 130 sites (30 m * 30 m) between

late July to mid-August 2020 in the Tabu River Basin in Siziwang Banner, Inner Mongolia

Autonomous Region (Figure 1). The types of the 130 sites were: 19 high-moderate, 14 high-low,

19 moderate-high, 16 moderate-moderate, 18 moderate-low, 16 low-high, 17 low-moderate, 11

low-low habitat amount and fragmentation per se. In order to exclude the impact of historical

agricultural activities, the habitat type of the established sites was natural grasslands with regional

vegetation characteristics. Each site was not abandoned agricultural land, and there was no sign of

agricultural reclamation.

At the 10 m * 10 m center of each site, we randomly set up three 1 m * 1 m plots in a flat

topographic area to investigate grassland vascular plant diversity and above-ground productivity.

Plant diversity was obtained by recording the number of vascular plant species in each plot. The

above-ground productivity was obtained by harvesting the above-ground biomass of the plants in

each plot and drying biomass at 65 °C to a constant weight. We also investigated two

environmental factors related to water and temperature (soil water content and land surface

temperature) in each site to consider their potential impact on plant diversity and ecosystem

function. The soil water content (SWC, %) was obtained by measuring the weight of wet soil

samples before and after oven-drying at 105 □ to a constant weight. The wet soil samples were

collected from three 30 cm deep cores using a 5 cm diameter soil auger within three plots of each

site. The land surface temperature (LST, □) was extracted from MODIS Land Surface

Temperature/Emissivity daily product (MOD11A1) using the Google Earth Engine platform. For

each site, the average daily LST of late July to mid-August 2020 was used in the following data

analysis.

As grassland is the dominant habitat type in the fragmented landscape and farmland is the

dominant matrix type, the specialists and generalists in this study were grouped as grassland

specialists, i.e., species that occur only in grassland, and weeds, i.e., species that occur in both

grassland and farmland. The classification of grassland specialists and weeds in this study was

based on our experience with plant surveys in this region, the List of Main Crop Weeds in China,

and available information in the Flora of China (http://www.iplant.cn/frps).

**Figure 1.** Map of study area in the Tabu River Basin, Inner Mongolia Autonomous Region,

northern China. (a) Location of the 130 survey sites in the study area. (b) Examples of four survey

sites with varying levels of habitat loss and fragmentation per se shown with a 500-m radius

buffer.

## 2.4 Data analysis

For each site, we calculated the mean vascular plant richness of the three 1 m * 1 m plots,

representing the vascular plant diversity, and the mean above-ground biomass of the three 1 m * 1

m plots, representing the above-ground productivity. The mean vascular plant richness and the

mean above-ground biomass were assessed to be normally distributed by a Shapiro-Wilk

normality test. The degree of habitat specialisation was represented by the percentage of grassland

specialists in the community. Landscape context includes habitat loss and fragmentation per se.

Habitat loss was represented by the percentage of non-grassland cover in the landscape and

fragmentation per se was represented by the inverse of the first principal component of the four

landscape indices (mean grassland patch area, mean nearest-neighbor distance among grassland

patch, grassland patch density, grassland patch edge). Environmental factors were LST and SWC.

First, to investigate the overall pattern between landscape context and grassland vascular

plant diversity and above-ground productivity in this region, we used scatter plots and Pearson

correlation analyses to present the pairwise relationship of habitat loss and fragmentation per se

with plant richness (including grassland specialist richness and weed richness) and above-ground

biomass.

Second, to investigate the relative importance of landscape context, grassland vascular plant

diversity, and environmental factors on above-ground productivity, we used the multi-model

averaging method based on the Akaike information criterion corrected for a small sample size

(AICc) (*Harrison et al., 2018*). We first constructed linear regression models, including the

response variable (above-ground biomass) and all combinations of the predictor variables (habitat

loss, fragmentation per se, grassland specialist richness, weed richness, LST, and SWC). The

models with the lowest AICc value and a difference of less than two ΔAICc from the lowest AICc

value were selected as optimal models. We then calculated the model-averaged standardized

parameter estimate based on the optimal models as the relative effect of each influencing factor on

above-ground biomass (*Harrison et al., 2018*). Before the analysis, we calculated the variance

inflation factors (VIF) for each predictor variable to assess multicollinearity. The VIF of all

explanatory variables was less than four (Appendix Table A2), suggesting no significant

multicollinearity in the analysis (*Carrara et al., 2015; Dormann et al., 2013*).

Finally, to investigate how landscape context impacts the relationship between grassland

plant diversity and above-ground productivity, we first used linear regression models to evaluate

relationship between plant richness and above-ground biomass at low, moderate, and high levels

of habitat loss and fragmentation per se, respectively. We then assessed the significance of

interaction terms between habitat loss and fragmentation per se and plant richness in the linear

regression models to evaluate whether they modulate the relationship between plant richness and

above-ground biomass. Further, we used a piecewise structural equation model to investigate the

specific pathways in which habitat loss and fragmentation per se modulate the relationship

between plant richness and above-ground biomass.

To do so, we first constructed a hypothetical conceptual model (Figure 2). Based on previous

studies, we hypothesised that habitat loss and fragmentation per se could decrease above-ground

biomass directly and indirectly by affecting plant richness and environmental factors (*Allan et al.,*

*2015; Haddad et al., 2015; Rolo et al., 2018*). As habitat loss is often the main cause of

fragmentation per se in natural landscapes, we hypothesised a causal relationship between habitat

loss and fragmentation per se (*Didham et al., 2012*). We also hypothesised that habitat loss and

fragmentation per se could both decrease the impact of plant richness on above-ground biomass by

decreasing the percentage of grassland specialists in the community (*Clavel et al., 2011; Gravel et*

*al., 2011; Liu et al., 2018*). We included the interaction between the percentage of grassland

specialists and the plant richness as predictor variables to quantify the modulating effect of the

percentage of grassland specialists for the impact of plant richness on above-ground biomass. We

tested the hypothetical conceptual model using a piecewise structural equation model (*Lefcheck,*

*2016*). The global fit of the model was evaluated using the Fisher's C statistic, and the

hypothetical pathways were evaluated by standardized path coefficients.

All data analyses were performed in the R programming language v. 4.0.3 (*R Development*

*Core Team, 2020*), with the following functions and packages. The Pearson correlation analysis,

linear regression model, and the Shapiro-Wilk normality test were conducted with 'cor', 'lm' and

'shapiro.test' functions of the stats package (*R Development Core Team, 2020*). The multi-model

averaging was conducted with the 'dredge' and 'model.avg' functions of the MuMIn package

(*Bartoń, 2020*). The VIF was calculated with the 'vif' function of the car package (*Fox and*

*Weisberg, 2019*). The piecewise structural equation model was conducted and tested with the

'psem' function of the piecewiseSEM package (*Lefcheck, 2016*).

**Figure 2.** Hypothetical model of habitat loss and fragmentation per se for the relationship between

grassland plant richness and above-ground biomass affecting the percentage of grassland

specialists in the community from 130 landscapes in the Tabu River Basin, a typical agro-pastoral

ecotone of northern China. Arrows represent the hypothesised paths among variables. See the

282                  main text for a description of arrow directions.

# 3 Results

## 3.1 Relationship of habitat loss and fragmentation per se with grassland plant richness and above-ground biomass

A total of 130 vascular plant species were identified in our study sites, including 91 grassland specialists and 39 weeds (Appendix Table A4). Habitat loss was significantly negatively correlated with overall species richness ($R$ = -0.21, $p < 0.05$, Figure 3a) and grassland specialist richness ($R$ = -0.41, $p < 0.01$, Figure 3a), but positively correlated with weed richness ($R$ = 0.31, $p < 0.01$, Figure 3a). Fragmentation per se was not significantly correlated with overall species richness and grassland specialist richness, but was significantly positively correlated with weed richness ($R$ = 0.26, $p < 0.01$, Figure 3b). Habitat loss ($R$ = -0.39, $p < 0.01$, Figure 3c) and fragmentation per se ($R$ = -0.26, $p < 0.01$, Figure 3d) were both significantly negatively correlated with above-ground biomass.

**Figure 3.** Correlation of habitat loss and fragmentation per se with grassland plant richness and

above-ground biomass from 130 landscapes in the Tabu River Basin, a typical agro-pastoral

ecotone of northern China: (a) habitat loss and plant richness, (b) fragmentation per se and plant

richness, (c) habitat loss and above-ground biomass, (d) fragmentation per se and above-ground

biomass. The $R$ value in each panel is from the Pearson correlation coefficient analyses. The n in

each panel is the number of surveying sites used in the Pearson correlation analyses. The trend

lines in the figure are from linear regression models. The shaded area around the trend line

represents the 95% confidence interval. * and ** represent significance at the 0.05 and 0.01 levels,

respectively.

## 3.2 The relative effects of landscape context, plant diversity, and environmental factors on above-ground biomass

Results of the multi-model averaging for the four optimal models affecting above-ground

biomass (Appendix Table A3) showed that plant diversity had the strongest relative effects on

above-ground biomass than landscape context and environmental factors. Grassland specialist

richness (estimate: 0.61, $p < 0.01$, Figure 4) and weed richness (estimate: 0.24, $p < 0.01$, Figure 4)

had significant positive relative effects on above-ground biomass. Habitat loss (estimate: -0.16, $p >$

0.05, Figure 4) and fragmentation per se (estimate: -0.05, $p > 0.05$, Figure 4) had insignificant

negative relative effects on above-ground biomass. Soil water content had a significant positive

relative effect on above-ground biomass (estimate: 0.19, $p < 0.01$, Figure 4), and land surface

temperature had an insignificant negative relative effect on above-ground biomass (estimate: -0.01,

$p > 0.05$, Figure 4).

**Figure 4.** Standardised parameter estimates and 95% confidence intervals for landscape context,

plant diversity, and environmental factors affecting above-ground biomass from 130 landscapes in

the Tabu River Basin, a typical agro-pastoral ecotone of northern China. Standardised estimates

and 95% confidence intervals are calculated by the multi-model averaging method based on the

four optimal models affecting above-ground biomass (Appendix Table A3). ** represent

significance at the 0.01 level.

## 3.3 The impact of habitat loss and fragmentation per se on the relationship between grassland plant richness and above-ground biomass

327    The linear regression models showed that habitat loss had a significant positive modulating

328 effect on the positive relationship between plant richness and above-ground biomass, and

329 fragmentation per se had no significant modulating effect (Figure 5). The positive relationship

330 between plant richness and above-ground biomass weakened with increasing levels of habitat loss,

331 strengthened and then weakened with increasing levels of fragmentation per se.

**Figure 5.** Relationships between grassland plant richness and above-ground biomass at different

levels of habitat loss and fragmentation per se from 130 landscapes in the Tabu River Basin, a

typical agro-pastoral ecotone of northern China: (a) high habitat loss and low fragmentation per se,

(b) high habitat loss and moderate fragmentation per se, (c) high habitat loss and high

fragmentation per se, (d) moderate habitat loss and low fragmentation per se, (e) moderate habitat

loss and moderate fragmentation per se, (f) moderate habitat loss and high fragmentation per se, (g)

low habitat loss and low fragmentation per se, (h) low habitat loss and moderate fragmentation per

se. The $R^2$ values in each panel are from linear regression models. The n in each panel is the

number of surveying sites used in the linear regression models. The blue solid and dashed trend

lines represent the significant and not significant effects, respectively. The shaded area around the

trend line represents the 95% confidence interval. * represent significance at the 0.05 level. **

represent significance at the 0.01 level.

The Fisher's C statistic indicated that the piecewise structural equation model fitted the data

well (*Fisher's C* = 19.3, *P*-value > 0.05, Figure 6). The piecewise structural equation model

showed that the percentage of grassland specialists increased the positive effect of plant richness

on above-ground biomass (path coefficient: 0.34, Figure 6). Habitat loss decreased the positive

effect of plant richness on above-ground biomass by decreasing the percentage of grassland

specialists (path coefficient: -0.46, Figure 6). Fragmentation per se had no significant effect on the

percentage of grassland specialists.

Meanwhile, the piecewise structural equation model showed habitat loss to have an indirect

negative effect on above-ground biomass through decreasing plant richness (path coefficient:

-0.39, Figure 6) and through increasing fragmentation per se (path coefficient: 0.69, Figure 6) and

in turn decreasing soil water content (path coefficient: -0.35, Figure 6). Fragmentation per se had

an indirect negative effect on above-ground biomass through decreasing soil water content (path

coefficient: -0.35, Figure 6) and an indirect positive effect on above-ground biomass through

increasing plant richness (path coefficient: 0.26, Figure 6).

**Figure 6.** Results from the piecewise structural equation model with hypothesised paths

showing how habitat loss and fragmentation per se alters the relationship between grassland plant

richness and above-ground biomass from 130 landscapes in the Tabu River Basin, a typical

agro-pastoral ecotone of northern China. HL: habitat loss. FPS: fragmentation per se. LST: land

surface temperature. SWT: soil water content. PS: percentage of grassland specialists. SR: plant

richness. PS×SR: interaction term between percentage of grassland specialists and plant richness.

AGB: above-ground biomass. Black and grey solid arrows represent the significant and not

significant effects at the 0.05 level, respectively. Black and red numbers on the solid arrows

represent the significant positive and negative path coefficients, respectively. The Fisher's C and

369                  *P*-values are from the piecewise structural equation model.

## 4 Discussion

### 4.1 Habitat loss and fragmentation per se had inconsistent effects on grassland plant diversity and ecosystem function

Although habitat loss and fragmentation per se are generally highly associated in natural landscapes, they are distinct ecological processes that determine decisions on effective conservation strategies (*Fahrig, 2017; Valente et al., 2023*). Our study evaluated the effects of habitat loss and fragmentation per se on grassland plant diversity and above-ground productivity in the context of fragmented landscapes in the agro-pastoral ecotone of northern China, with our results showing the effects of these two facets to not be consistent.

Consistent with previous studies, we found habitat loss significantly reduced grassland plant diversity, suggesting that habitat loss is a major threat to the biodiversity conservation of fragmented landscapes in this region (*Chase et al., 2020; Haddad et al., 2015; Ibáñez et al., 2014*). While for fragmentation per se, we found a positive effect on grassland plant diversity, in accordance with some recent evidence, the effects of fragmentation per se on biodiversity are more likely positive than negative for a given amount of habitat (*Gestich et al., 2022; Palmeirim et al., 2019; Riva and Fahrig, 2023*). Our study suggests that biodiversity conservation strategies for fragmented landscapes should consider optimising the habitat configuration in the landscape in addition to preventing habitat loss, such as increasing the number of habitat patches (*Arroyo-Rodríguez et al., 2020; Fahrig, 2017*). However, it is important to stress that the observed positive effect of fragmentation per se does not imply that increasing the isolation of grassland patches would promote biodiversity, as the metric of fragmentation per se used in our study was more related to patch density, edge density and mean patch area while relatively less related to patch isolation (Appendix Table A1). The potential threats from isolation still needs to be carefully considered in the conservation of biodiversity in fragmented landscapes (*Haddad et al., 2015*).

Our results showed for a metric of ecosystem function that landscape context affected

above-ground productivity indirectly by influencing plant diversity and environment factors,

which is consistent with the findings of previous studies in fragmented landscapes (*Allan et al.,*

*2015; Rippel et al., 2020*). And we found fragmentation per se mediated the effect of habitat loss

on environmental factors. A possible reason for this finding is fragmentation per se could directly

alter environmental factors through enhanced edge effects (*Laurance et al., 2011; Smith et al.,*

*2018*). In our study, fragmentation per se could lead to ecosystem drought and thus limit

above-ground productivity in grassland ecosystems. This is mainly because enhanced edge effects

due to fragmentation per se could lead to greater desiccation and evapotranspiration rates in

remaining habitats (*Smith et al., 2018; Tuff et al., 2016*). Therefore, our study suggests that habitat

loss and fragmentation per se have inconsistent consequences on plant diversity and ecosystem

function, which should be considered explicitly in developing conservation strategies in

fragmented landscapes.

In our study, a possible mechanism for the positive impacts of fragmentation per se on plant

diversity and above-ground productivity (indirect positive impact via plant diversity) is that

fragmentation per se increases the habitat heterogeneity in the landscape, which can promote

biodiversity through spatial asynchrony and spatial insurance effects (*Tscharntke et al., 2012*).

Previous studies indicated that heterogeneity typically has nonlinear effects on biodiversity and

ecosystem function, as moderate heterogeneity can maximise spatial asynchrony (*Redon et al.,*

*2014; Wilcox et al., 2017*). However, our study did not observe nonlinear patterns between

fragmentation per se and plant diversity and above-ground productivity. This may be due to the

low spatial heterogeneity of this area as a result of agricultural intensification (*Benton et al., 2003;*

*Chen et al., 2019*). The gradient of fragmentation per se in our study may not cover the optimal

heterogeneity levels for maximising plant diversity and above-ground productivity (*Thompson and*

*Gonzalez, 2016*).

**4.2 Habitat loss rather than fragmentation per se weakened the magnitude of the positive relationship between plant diversity and ecosystem function**

Understanding the direction and magnitude of BEF relationships in fragmented landscapes is essential to understanding the importance of biodiversity for ecosystem function in the changing world (*Gonzalez et al., 2020; van der Plas, 2019*). In naturally assembled communities, ecosystem functions may be dominated by complex environmental factors and landscape context, showing a weak or even negative correlation with biodiversity (*Grace et al., 2007; Hagan et al., 2021; Zirbel et al., 2019*). Our study found grassland plant diversity showed a stronger positive impact on above-ground productivity than landscape context and environmental factors. This result is consistent with findings by *Duffy et al.* (*2017*) in natural ecosystems, indicating grassland plant diversity has an important role in maintaining grassland ecosystem functions in the fragmented landscapes of the agro-pastoral ecotone of northern China.

Consistent with theoretical model predictions (*Liu et al., 2018*), our study found that fragmented landscape context significantly modulated the BEF relationship. We further found that habitat loss weakened the magnitude of the BEF relationship through decreasing the percentage of specialists in grassland communities, as specialists were more vulnerable to the negative effect of habitat loss and more associated with above-ground productivity than generalists. These findings indicate reducing the degree of habitat specialisation may be the mechanism of fragmented landscape context weakening the BEF relationship in this region (*Clavel et al., 2011; Gravel et al., 2011*). Meanwhile, our study demonstrates that habitat loss, rather than fragmentation per se, can decrease the degree of habitat specialisation by leading to the replacement of specialists by generalists in the community, thus weakening the BEF relationship. This is mainly because fragmentation per se did not decrease the grassland specialist richness in this region, whereas habitat loss decreased the grassland specialist richness and led to the invasion of more weeds from the surrounding farmland into the grassland community (*Yan et al., 2022; Yan et al., 2023*). Our findings suggest further expansion of farmland in the agro-pastoral ecotone of northern China would decrease the grassland plant diversity and its importance to above-ground productivity.

In addition, our study found that the BEF relationship showed a nonlinear pattern with

increasing levels of fragmentation per se. For a given level of habitat loss, the positive BEF

relationship was strongest at moderate fragmentation per se level and became neutral at high

fragmentation per se level. This can be explained by the increased spatial asynchrony at moderate

fragmentation per se level, which can promote niche complementary among species in the

community and thus strengthen the BEF relationship (*Gonzalez et al., 2020; Thompson and*

*Gonzalez, 2016; Tscharntke et al., 2012*). The neutral BEF relationship at high fragmentation per

se level may be due to edge effects enhancing environmental filtering, thereby leading to

functional redundancy among species and decoupling the BEF relationship (*Fetzer et al., 2015;*

*Hu et al., 2016; Zambrano et al., 2019*). However, a recent study by *Hertzog et al.* (*2019*) on

temperate forest ecosystems found that the positive relationship between plant diversity and

ecosystem multifunctionality strengthened with increasing forest fragmentation and peaked at the

high level of forest fragmentation. This inconsistency can be explained by trade-offs between

different ecosystem functions that may differ in their response to fragmentation per se

(*Banks-Leite et al., 2020*). Therefore, future studies are needed to focus on multiple ecosystem

functions, such as below-ground productivity, litter decomposition, soil carbon stocks, etc.

## 5 Conclusions

Our study investigated the effects of habitat loss and fragmentation per se on grassland plant

diversity, above-ground productivity, and their relationships in the agro-pastoral ecotone of

northern China. We found habitat loss and fragmentation per se had inconsistent effects. Habitat

loss decreased grassland plant diversity, while fragmentation per se increased grassland plant

diversity. Habitat loss did not affect above-ground biomass, while fragmentation per se decreased

above-ground biomass. We also found that habitat loss rather than fragmentation per se weakened

the magnitude of the BEF relationship by decreasing the habitat specialisation of the community.

Our study suggests biodiversity conservation in fragmented landscapes should focus on preventing

habitat loss, which decreases plant diversity and its importance to ecosystem function.

# ACKNOWLEDGMENTS

We are very grateful to Peng Han, Shuangshuang Zhang, Xiaoqian Gong, Fengshi Li, Nier Su, Luyao Liu, Linjie Yao, and Guofeng Shi for assistance with field work. This study was supported by the National Natural Science Foundation of China (32071582) and the Inner Mongolia Autonomous Region Science and Technology Plan Project (2021GG0392). No permits were required for fieldwork in this research.

# CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

# DATA ACCESSIBILITY STATEMENT

If the manuscript is accepted, we will upload the data to a publicly accessible digital repository.

# AUTHORS' CONTRIBUTIONS

Q.Z. and Y.Z. conceived the research; Y.Z., S.J., and Q.Z. contributed to the writing of the manuscript; Y.Z. led the data compilation and analysis; Y.Z. and Q.Z. contributed to the fieldwork.

# References

Allan, E., Manning, P., Alt, F., Binkenstein, J., Blaser, S., Blüthgen, N., Böhm, S., Grassein, F., Hölzel, N., Klaus, V.H., Kleinebecker, T., Morris, E.K., Oelmann, Y., Prati, D., Renner, S.C., Rillig, M.C., Schaefer, M., Schloter, M., Schmitt, B., Schöning, I., et al. 2015. Land use intensification alters ecosystem multifunctionality via loss of biodiversity and changes to functional composition. *Ecology Letters* **18**: 834-843. https://doi.org/10.1111/ele.12469

Arroyo-Rodríguez, V., Fahrig, L., Tabarelli, M., Watling, J.I., Tischendorf, L., Benchimol, M., Cazetta,
E., Faria, D., Leal, I.R., Melo, F.P.L., Morante-Filho, J.C., Santos, B.A., Arasa-Gisbert, R.,
Arce-Peña, N., Cervantes-López, M.J., Cudney-Valenzuela, S., Galán-Acedo, C., San-José, M.,
Vieira, I.C.G., Slik, J.W.F., et al. 2020. Designing optimal human-modified landscapes for
forest biodiversity conservation. *Ecology Letters* **23**: 1404-1420.
https://doi.org/10.1111/ele.13535

Banks-Leite, C., Ewers, R.M., Folkard-Tapp, H., Fraser, A. 2020. Countering the effects of habitat loss,
fragmentation, and degradation through habitat restoration. *One Earth* **3**: 672-676.
https://doi.org/10.1016/j.oneear.2020.11.016

Bartoń, K. 2020. MuMIn: multi-model inference. *R package version 1.43.17*.
https://CRAN.R-project.org/package=MuMIn

Benton, T.G., Vickery, J.A., Wilson, J.D. 2003. Farmland biodiversity: is habitat heterogeneity the key?
*Trends in Ecology & Evolution* **18**: 182-188. https://doi.org/10.1016/S0169-5347(03)00011-9

Butsic, V., Lewis, D.J., Radeloff, V.C., Baumann, M., Kuemmerle, T. 2017. Quasi-experimental
methods enable stronger inferences from observational data in ecology. *Basic and Applied
Ecology* **19**: 1-10. https://doi.org/10.1016/j.baae.2017.01.005

Carrara, E., Arroyo-Rodríguez, V., Vega-Rivera, J.H., Schondube, J.E., de Freitas, S.M., Fahrig, L.
2015. Impact of landscape composition and configuration on forest specialist and generalist
bird species in the fragmented Lacandona rainforest, Mexico. *Biological Conservation* **184**:
117-126. https://doi.org/10.1016/j.biocon.2015.01.014

Chase, J.M., Blowes, S.A., Knight, T.M., Gerstner, K., May, F. 2020. Ecosystem decay exacerbates
biodiversity loss with habitat loss. *Nature* **584**: 238-243.
https://doi.org/10.1038/s41586-020-2531-2

Chen, X., Wang, F., Jiang, L., Huang, C., An, P., Pan, Z. 2019. Impact of center pivot irrigation on
vegetation dynamics in a farming-pastoral ecotone of Northern China: A case study in
Ulanqab, Inner Mongolia. *Ecological Indicators* **101**: 274-284.
https://doi.org/10.1016/j.ecolind.2019.01.027

Clavel, J., Julliard, R., Devictor, V. 2011. Worldwide decline of specialist species: toward a global
functional homogenization? *Frontiers in Ecology and the Environment* **9**: 222-228.
https://doi.org/10.1890/080216

Dehling, D.M., Bender, I.M.A., Blendinger, P.G., Böhning-Gaese, K., Muñoz, M.C., Neuschulz, E.L.,
Quitián, M., Saavedra, F., Santillán, V., Schleuning, M., Stouffer, D.B. 2021. Specialists and
generalists fulfil important and complementary functional roles in ecological processes.
*Functional Ecology* **35**: 1810-1821. https://doi.org/10.1111/1365-2435.13815

Denelle, P., Violle, C., DivGrass, C., Munoz, F. 2020. Generalist plants are more competitive and more
functionally similar to each other than specialist plants: insights from network analyses.
*Journal of Biogeography* **47**: 1922-1933. https://doi.org/10.1111/jbi.13848

Devictor, V., Julliard, R., Jiguet, F. 2008. Distribution of specialist and generalist species along spatial
gradients of habitat disturbance and fragmentation. *Oikos* **117**: 507-514.
https://doi.org/10.1111/j.0030-1299.2008.16215.x

Didham, R.K., Kapos, V., Ewers, R.M. 2012. Rethinking the conceptual foundations of habitat
fragmentation research. *Oikos* **121**: 161-170.
https://doi.org/10.1111/j.1600-0706.2011.20273.x

Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J.R.G., Gruber, B.,

Lafourcade, B., Leitão, P.J., Münkemüller, T., McClean, C., Osborne, P.E., Reineking, B., Schröder, B., Skidmore, A.K., Zurell, D., Lautenbach, S. 2013. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* **36**: 27-46. https://doi.org/10.1111/j.1600-0587.2012.07348.x

Duffy, J.E., Godwin, C.M., Cardinale, B.J. 2017. Biodiversity effects in the wild are common and as strong as key drivers of productivity. *Nature* **549**: 261-264. https://doi.org/10.1038/nature23886

Fahrig, L. 2003. Effects of habitat fragmentation on biodiversity. *Annual Review of Ecology, Evolution, and Systematics* **34**: 487-515. https://doi.org/10.1146/annurev.ecolsys.34.011802.132419

Fahrig, L. 2017. Ecological responses to habitat fragmentation per se. *Annual Review of Ecology, Evolution, and Systematics* **48**: 1-23. https://doi.org/10.1146/annurev-ecolsys-110316-022612

Fahrig, L., Arroyo-Rodríguez, V., Bennett, J.R., Boucher-Lalonde, V., Cazetta, E., Currie, D.J., Eigenbrod, F., Ford, A.T., Harrison, S.P., Jaeger, J.A.G., Koper, N., Martin, A.E., Martin, J.-L., Metzger, J.P., Morrison, P., Rhodes, J.R., Saunders, D.A., Simberloff, D., Smith, A.C., Tischendorf, L., et al. 2019. Is habitat fragmentation bad for biodiversity? *Biological Conservation* **230**: 179-186. https://doi.org/10.1016/j.biocon.2018.12.026

Fardila, D., Kelly, L.T., Moore, J.L., McCarthy, M.A. 2017. A systematic review reveals changes in where and how we have studied habitat loss and fragmentation over 20years. *Biological Conservation* **212**: 130-138. https://doi.org/10.1016/j.biocon.2017.04.031

Fetzer, I., Johst, K., Schäwe, R., Banitz, T., Harms, H., Chatzinotas, A. 2015. The extent of functional redundancy changes as species' roles shift in different environments. *Proceedings of the National Academy of Sciences* **112**: 14888-14893. https://doi.org/10.1073/pnas.1505587112

Fletcher Jr, R.J., Smith, T.A.H., Kortessis, N., Bruna, E.M., Holt, R.D. 2023. Landscape experiments unlock relationships among habitat loss, fragmentation, and patch-size effects. *Ecology* **104**: e4037. https://doi.org/10.1002/ecy.4037

Fletcher, R.J., Didham, R.K., Banks-Leite, C., Barlow, J., Ewers, R.M., Rosindell, J., Holt, R.D., Gonzalez, A., Pardini, R., Damschen, E.I., Melo, F.P.L., Ries, L., Prevedello, J.A., Tscharntke, T., Laurance, W.F., Lovejoy, T., Haddad, N.M. 2018. Is habitat fragmentation good for biodiversity? *Biological Conservation* **226**: 9-15. https://doi.org/10.1016/j.biocon.2018.07.022

Fox, J., Weisberg, S. 2019. An R companion to Applied Regression. *Sage*. https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Gestich, C.C., Arroyo-Rodríguez, V., Saranholi, B.H., da Cunha, R.G.T., Setz, E.Z.F., Ribeiro, M.C. 2022. Forest loss and fragmentation can promote the crowding effect in a forest-specialist primate. *Landscape Ecology* **37**: 147-157. https://doi.org/10.1007/s10980-021-01336-1

Godbold, J.A., Bulling, M.T., Solan, M. 2011. Habitat structure mediates biodiversity effects on ecosystem properties. *Proceedings of the Royal Society B: Biological Sciences* **278**: 2510-2518. https://doi.org/10.1098/rspb.2010.2414

Gonzalez, A., Germain, R.M., Srivastava, D.S., Filotas, E., Dee, L.E., Gravel, D., Thompson, P.L., Isbell, F., Wang, S., Kéfi, S., Montoya, J., Zelnik, Y.R., Loreau, M. 2020. Scaling-up biodiversity-ecosystem functioning research. *Ecology Letters* **23**: 757-776. https://doi.org/10.1111/ele.13456

Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., Moore, R. 2017. Google Earth Engine: planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment* **202**: 18-27. https://doi.org/10.1016/j.rse.2017.06.031

Grace, J.B., Michael Anderson, T., Smith, M.D., Seabloom, E., Andelman, S.J., Meche, G., Weiher, E., Allain, L.K., Jutila, H., Sankaran, M., Knops, J., Ritchie, M., Willig, M.R. 2007. Does species diversity limit productivity in natural grassland communities? *Ecology Letters* **10**: 680-689. https://doi.org/10.1111/j.1461-0248.2007.01058.x

Gravel, D., Bell, T., Barbera, C., Bouvier, T., Pommier, T., Venail, P., Mouquet, N. 2011. Experimental niche evolution alters the strength of the diversity–productivity relationship. *Nature* **469**: 89-92. https://doi.org/10.1038/nature09592

Haddad, N.M., Brudvig, L.A., Clobert, J., Davies, K.F., Gonzalez, A., Holt, R.D., Lovejoy, T.E., Sexton, J.O., Austin, M.P., Collins, C.D., Cook, W.M., Damschen, E.I., Ewers, R.M., Foster, B.L., Jenkins, C.N., King, A.J., Laurance, W.F., Levey, D.J., Margules, C.R., Melbourne, B.A., et al. 2015. Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances* **1**: e1500052. https://doi.org/10.1126/sciadv.1500052

Hagan, J.G., Vanschoenwinkel, B., Gamfeldt, L. 2021. We should not necessarily expect positive relationships between biodiversity and ecosystem functioning in observational field data. *Ecology Letters* **24**: 2537-2548. https://doi.org/10.1111/ele.13874

Harrison, X.A., Donaldson, L., Correa-Cano, M.E., Evans, J., Fisher, D.N., Goodwin, C.E., Robinson, B.S., Hodgson, D.J., Inger, R. 2018. A brief introduction to mixed effects modelling and multi-model inference in ecology. *PeerJ* **6**: e4794. https://doi.org/10.7717/peerj.4794

Hertzog, L.R., Boonyarittichaikij, R., Dekeukeleire, D., de Groote, S.R.E., van Schrojenstein Lantman, I.M., Sercu, B.K., Smith, H.K., de la Peña, E., Vandegehuchte, M.L., Bonte, D., Martel, A., Verheyen, K., Lens, L., Baeten, L. 2019. Forest fragmentation modulates effects of tree species richness and composition on ecosystem multifunctionality. *Ecology* **100**: e02653. https://doi.org/10.1002/ecy.2653

Hu, G., Feeley, K.J., Yu, M. 2016. Habitat fragmentation drives plant community assembly processes across life stages. *PLoS ONE* **11**: e0159572. https://doi.org/10.1371/journal.pone.0159572

Ibáñez, I., Katz, D.S.W., Peltier, D., Wolf, S.M., Connor Barrie, B.T. 2014. Assessing the integrated effects of landscape fragmentation on plants and plant communities: the challenge of multiprocess–multiresponse dynamics. *Journal of Ecology* **102**: 882-895. https://doi.org/10.1111/1365-2745.12223

IPBES 2019. Global assessment report on biodiversity and ecosystem services of the intergovernmental science-policy platform on biodiversity and ecosystem services. *Bonn: IPBES*. https://doi.org/10.5281/zenodo.5657041

Laurance, W.F., Camargo, J.L.C., Luizão, R.C.C., Laurance, S.G., Pimm, S.L., Bruna, E.M., Stouffer, P.C., Bruce Williamson, G., Benítez-Malvido, J., Vasconcelos, H.L., Van Houtan, K.S., Zartman, C.E., Boyle, S.A., Didham, R.K., Andrade, A., Lovejoy, T.E. 2011. The fate of Amazonian forest fragments: a 32-year investigation. *Biological Conservation* **144**: 56-67. https://doi.org/10.1016/j.biocon.2010.09.021

Lefcheck, J.S. 2016. PIECEWISESEM: piecewise structural equation modelling in R for ecology, evolution, and systematics. *Methods in Ecology and Evolution* **7**: 573-579. https://doi.org/10.1111/2041-210x.12512

Liu, J., Wilson, M., Hu, G., Liu, J., Wu, J., Yu, M. 2018. How does habitat fragmentation affect the biodiversity and ecosystem functioning relationship? *Landscape Ecology* **33**: 341-352. https://doi.org/10.1007/s10980-018-0620-5

Ma, J., Li, J., Wu, W., Liu, J. 2023. Global forest fragmentation change from 2000 to 2020. *Nature*

*Communications* **14**: 3752. https://doi.org/10.1038/s41467-023-39221-x

Matthews, T.J. 2021. On the biogeography of habitat islands: the importance of matrix effects, noncore

species, and source-sink dynamics. *The Quarterly Review of Biology* **96**: 73-104.

https://doi.org/10.1086/714482

Matthews, T.J., Cottee-Jones, H.E., Whittaker, R.J. 2014. Habitat fragmentation and the species–area

relationship: a focus on total species richness obscures the impact of habitat loss on habitat

specialists. *Diversity and Distributions* **20**: 1136-1146. https://doi.org/10.1111/ddi.12227

Maxwell, S.L., Fuller, R.A., Brooks, T.M., Watson, J.E.M. 2016. Biodiversity: the ravages of guns, nets

and bulldozers. *Nature* **536**: 143-145. https://doi.org/10.1038/536143a

McGarigal, K., Cushman, S.A., Ene, E. 2012. FRAGSTATS v4: spatial pattern analysis program for

categorical and continuous maps. *Computer software program produced by the authors at the*

*University of Massachusetts, Amherst. Available at the following web site:*

http://www.umass.edu/landeco/research/fragstats/fragstats.html.

Miller-Rushing, A.J., Primackh, R.B., Devictor, V., Corlett, R.T., Cumming, G.S., Loyola, R., Maas, B.,

Pejchar, L. 2019. How does habitat fragmentation affect biodiversity? A controversial question

at the core of conservation biology. *Biological Conservation* **232**: 271-273.

https://doi.org/10.1016/j.biocon.2018.12.029

Palmeirim, A.F., Figueiredo, M.S.L., Grelle, C.E.V., Carbone, C., Vieira, M.V. 2019. When does habitat

fragmentation matter? a biome-wide analysis of small mammals in the Atlantic Forest.

*Journal of Biogeography* **46**: 2811-2825. https://doi.org/10.1111/jbi.13730

Pasher, J., Mitchell, S.W., King, D.J., Fahrig, L., Smith, A.C., Lindsay, K.E. 2013. Optimizing

landscape selection for estimating relative effects of landscape variables on ecological

responses. *Landscape Ecology* **28**: 371-383. https://doi.org/10.1007/s10980-013-9852-6

R Development Core Team 2020. R: a language and environment for statistical computing. *R*

*Foundation for Statistical Computing*. https://www.R-project.org

Redon, M., Bergès, L., Cordonnier, T., Luque, S. 2014. Effects of increasing landscape heterogeneity

on local plant species richness: how much is enough? *Landscape Ecology* **29**: 773-787.

https://doi.org/10.1007/s10980-014-0027-x

Reynolds, C., Fletcher, R.J., Carneiro, C.M., Jennings, N., Ke, A., LaScaleia, M.C., Lukhele, M.B.,

Mamba, M.L., Sibiya, M.D., Austin, J.D., Magagula, C.N., Mahlaba, T.a., Monadjem, A.,

Wisely, S.M., McCleery, R.A. 2018. Inconsistent effects of landscape heterogeneity and

land-use on animal diversity in an agricultural mosaic: a multi-scale and multi-taxon

investigation. *Landscape Ecology* **33**: 241-255. https://doi.org/10.1007/s10980-017-0595-7

Rippel, T.M., Mooring, E.Q., Tomasula, J., Wimp, G.M. 2020. Habitat edge effects decrease litter

accumulation and increase litter decomposition in coastal salt marshes. *Landscape Ecology* **35**:

2179-2190. https://doi.org/10.1007/s10980-020-01108-3

Riva, F., Fahrig, L. 2023. Landscape-scale habitat fragmentation is positively related to biodiversity,

despite patch-scale ecosystem decay. *Ecology Letters* **26**: 268-277.

https://doi.org/10.1111/ele.14145

Rolo, V., Olivier, P.I., Pfeifer, M., van Aarde, R.J. 2018. Functional diversity mediates contrasting

direct and indirect effects of fragmentation on below- and above-ground carbon stocks of

coastal dune forests. *Forest Ecology and Management* **407**: 174-183.

https://doi.org/10.1016/j.foreco.2017.10.059

Smith, A.C., Koper, N., Francis, C.M., Fahrig, L. 2009. Confronting collinearity: comparing methods

for disentangling the effects of habitat loss and fragmentation. *Landscape Ecology* **24**: 1271-1285. https://doi.org/10.1007/s10980-009-9383-3

Smith, I.A., Hutyra, L.R., Reinmann, A.B., Marrs, J.K., Thompson, J.R. 2018. Piecing together the fragments: elucidating edge effects on forest carbon dynamics. *Frontiers in Ecology and the Environment* **16**: 213-221. https://doi.org/10.1002/fee.1793

Thompson, P.L., Gonzalez, A. 2016. Ecosystem multifunctionality in metacommunities. *Ecology* **97**: 2867-2879. https://doi.org/10.1002/ecy.1502

Tilman, D., Reich, P.B., Isbell, F. 2012. Biodiversity impacts ecosystem productivity as much as resources, disturbance, or herbivory. *Proceedings of the National Academy of Sciences* **109**: 10394-10397. https://doi.org/10.1073/pnas.1208240109

Tscharntke, T., Tylianakis, J.M., Rand, T.A., Didham, R.K., Fahrig, L., Batáry, P., Bengtsson, J., Clough, Y., Crist, T.O., Dormann, C.F., Ewers, R.M., Fründ, J., Holt, R.D., Holzschuh, A., Klein, A.M., Kleijn, D., Kremen, C., Landis, D.A., Laurance, W., Lindenmayer, D., et al. 2012. Landscape moderation of biodiversity patterns and processes - eight hypotheses. *Biological Reviews* **87**: 661-685. https://doi.org/10.1111/j.1469-185X.2011.00216.x

Tuff, K.T., Tuff, T., Davies, K.F. 2016. A framework for integrating thermal biology into fragmentation research. *Ecology Letters* **19**: 361-374. https://doi.org/10.1111/ele.12579

Valente, J.J., Gannon, D.G., Hightower, J., Kim, H., Leimberger, K.G., Macedo, R., Rousseau, J.S., Weldy, M.J., Zitomer, R.A., Fahrig, L., Fletcher, R.J., Wu, J., Betts, M.G. 2023. Toward conciliation in the habitat fragmentation and biodiversity debate. *Landscape Ecology* **38**: 2717-2730. https://doi.org/10.1007/s10980-023-01708-9

van der Plas, F. 2019. Biodiversity and ecosystem functioning in naturally assembled communities. *Biological Reviews* **94**: 1220-1245. https://doi.org/10.1111/brv.12499

Wang, X., Blanchet, F.G., Koper, N. 2014. Measuring habitat fragmentation: an evaluation of landscape pattern metrics. *Methods in Ecology and Evolution* **5**: 634-646. https://doi.org/10.1111/2041-210X.12198

Wardle, D.A., Zackrisson, O. 2005. Effects of species and functional group loss on island ecosystem properties. *Nature* **435**: 806-810. https://doi.org/10.1038/nature03611

Wilcox, K.R., Tredennick, A.T., Koerner, S.E., Grman, E., Hallett, L.M., Avolio, M.L., La Pierre, K.J., Houseman, G.R., Isbell, F., Johnson, D.S., Alatalo, J.M., Baldwin, A.H., Bork, E.W., Boughton, E.H., Bowman, W.D., Britton, A.J., Cahill Jr, J.F., Collins, S.L., Du, G., Eskelinen, A., et al. 2017. Asynchrony among local communities stabilises ecosystem function of metacommunities. *Ecology Letters* **20**: 1534-1545. https://doi.org/10.1111/ele.12861

Wilsey, B., Martin, L., Xu, X., Isbell, F., Polley, H.W. 2023. Biodiversity: Net primary productivity relationships are eliminated by invasive species dominance. *Ecology Letters*. https://doi.org/10.1111/ele.14342

Wilson, M.C., Chen, X.-Y., Corlett, R.T., Didham, R.K., Ding, P., Holt, R.D., Holyoak, M., Hu, G., Hughes, A.C., Jiang, L., Laurance, W.F., Liu, J., Pimm, S.L., Robinson, S.K., Russo, S.E., Si, X., Wilcove, D.S., Wu, J., Yu, M. 2016. Habitat fragmentation and biodiversity conservation: key findings and future challenges. *Landscape Ecology* **31**: 219-227. https://doi.org/10.1007/s10980-015-0312-3

Yan, Y., Jarvie, S., Liu, Q., Zhang, Q. 2022. Effects of fragmentation on grassland plant diversity depend on the habitat specialization of species. *Biological Conservation* **275**: 109773.

https://doi.org/10.1016/j.biocon.2022.109773

Yan, Y., Jarvie, S., Zhang, Q., Han, P., Liu, Q., Zhang, S., Liu, P. 2023. Habitat heterogeneity determines species richness on small habitat islands in a fragmented landscape. *Journal of Biogeography* **50**: 976-986. https://doi.org/10.1111/jbi.14594

Yan, Y., Jarvie, S., Zhang, Q., Zhang, S., Han, P., Liu, Q., Liu, P. 2021. Small patches are hotspots for biodiversity conservation in fragmented landscapes. *Ecological Indicators* **130**: 108086. https://doi.org/10.1016/j.ecolind.2021.108086

Yang, Y., Wang, K., Liu, D., Zhao, X., Fan, J. 2020. Effects of land-use conversions on the ecosystem services in the agro-pastoral ecotone of northern China. *Journal of Cleaner Production* **249**: 119360. https://doi.org/10.1016/j.jclepro.2019.119360

Zambrano, J., Garzon-Lopez, C.X., Yeager, L., Fortunel, C., Cordeiro, N.J., Beckman, N.G. 2019. The effects of habitat loss and fragmentation on plant functional traits and functional diversity: what do we know so far? *Oecologia* **191**: 505-518. https://doi.org/10.1007/s00442-019-04505-x

Zhang, S., Zhang, Q., Yan, Y., Han, P., Liu, Q. 2021. Island biogeography theory predicts plant species richness of remnant grassland patches in the agro-pastoral ecotone of northern China. *Basic and Applied Ecology* **54**: 14-22. https://doi.org/10.1016/j.baae.2021.04.010

Zirbel, C.R., Grman, E., Bassett, T., Brudvig, L.A. 2019. Landscape context explains ecosystem multifunctionality in restored grasslands better than plant diversity. *Ecology* **100**: e02634. https://doi.org/10.1002/ecy.2634