Latitudinal patterns of flowering phenology of three widespread tropical species from

public media and data repositories

Authors: Bhavya Kriti*[1], Sidharth Sankaran [2], Geetha Ramaswami [3]

Affiliations:

1. *Corresponding author, St. Xavier's College, Ranchi, Jharkhand,

India, email: kriti.bhavya22@gmail.com

Orcid ID: bhavyak@orcid

2. Poorna Learning Centre, Bengaluru, India, email:

sidharth.san2005@gmail.com

3. Nature Conservation Foundation, Mysuru, India, email:

geetha@ncfindia.org

Abstract

Plant phenology encompasses the study of periodic events in biological life cycles of plants

and the influence of seasonal changes in weather and other environmental factors. In order to

understand the role of the environment and the changing climate on the life cycle of plants,

long-term records are required to establish baselines against which significant future

deviations can be discerned. This study aims to analyze the spatial flowering pattern of the

plant species -*Bombax ceiba, Butea monosperma* and *Cassia fistula* across India, through

images uploaded on public media repositories like citizen science portals and social media.

Citizen science is rapidly emerging as a reliable source of supplementary scientific

information, and can be utilized to understand broad patterns of tree behavior. We found that

for all three species, flowering period is longer, and begins earlier in southern latitudes as

compared to the northern latitudes. We conclude that publicly available images of flowering

trees and citizen science data can be used to supplement existing information on tree

phenology across India.

Keywords: citizen science, latitudinal patterns, media repositories, tree phenology, tree

seasonality,   SeasonWatch

1. Introduction:

Phenology is the study of the timing of the specific life history events of a species, particularly in relation to cyclic seasonal changes (Raju *et al* . 2005; Jain *et al* . 2009; Sindhi and Bairwa 2010). This timing is an important factor in plant reproductive cycles and hence important to the population dynamics and persistence of a species (Jain *et al*. 2009; Sindhi and Bairwa 2010). Phenological events in mature plants include leaf-emergence, flowering, vegetative growth, fruiting, and leaf fall (Rathcke *et al*. 1985). Environmental variables such as temperature,precipitation, photoperiod, soil moisture, humidity, and soil nutrient content have been shown to influence plant phenology (Abu- Asab *et al* . 2001; Kudo and Ida 2013; Renner and Zohner 2018), and the same species may also show variable phenology depending on spatial factors such as latitude and altitude (Chaudhary and Khadabadi 2012). Of these factors, temperature and precipitation are likely to become highly variable due to global- warming induced climate change, in turn affecting plant phenology, typically observed as changes in the onset of a phenophase from a historical baseline (Walther *et al*. 2002; Cleland 2007; Gordo and Sanz 2010; Numata *et al*. 2022). Variability in environmental factors is likely to be exacerbated by anthropogenic changes such as urbanization, contributing to localized 'urban heat islands' or altered soil moisture due to artificial irrigation (Neil and Wu 2006). Climate change impacts on plant phenology are better-studied in temperate regions - where plants respond to lengthening growing seasons through advancing vegetative and reproductive phenology such as budburst and flowering (Badeck *et al*. 2004; Gordo and Sanz 2010). In recent years, due to climate change, seasonal climatic variations have begun to differ from their normal patterns (Rathcke *et al*. 1985). This could cause variations in the flowering dates (Renner and Zohner 2018; Numata *et al*. 2022) and reproductive cycles of many tree species which could in turn affect a whole host of animal species that depend on these trees to provide food and habitats. Plant phenology is linked to the life history patterns of consumers that feed on these species as well as their interactions

with each other (Rathcke *et al*. 1985; Raju *et al*. 2005; Burli at al. 2007; Jain *et al*. 2009;

Sindhi and Bairwa 2010). Consequently, changes in plant-pollinator relationships – where

change in the timing of cyclic life events could lead to phenological mismatch- can

negatively impact the interacting species (Renner and Zohner 2018). For example, early onset

of flowering in the pan-Asian tuberous plant species *Corydalis ambigua* is associated with a

lowered rate of pollination by bumblebees and therefore in lower seed production (Kudo and

Ida 2013). Thus knowledge of the phenology of a species is important as it can be used to

predict how the species may respond to climate change and the larger ecological implications

at an ecosystem level.

India has a very wide variation in latitudes, extending between 8°4'N and 37°6'S and the

climate varies accordingly – with very little temperature seasonality in the south and high

temperature seasonality in the north, west and eastern regions of the country (Attri and Tyagi

2010). A large part of the Indian subcontinent is thus dominated by widespread tropical

species. We hypothesise that species with ranges spread across latitudes, will show variable

phenological behaviour as one can expect that the optimal temperature, and other

environmental conditions such as day-length and precipitation, affecting phenology in these

species would also vary considerably with latitude. The change in the timing of vegetative

and reproductive phenology in tropical tree species is poorly understood due to lack of

historical documentation.

Information on species-level phenology is often difficult to find outside of regional floras

and other technical sources. Ideally, population-level replicates of species should be

considered for such analyses. However, in the absence of intensively sampled data, one can

derive information from large online public repositories of species images (Barve 2014) or

from citizen science projects such as SeasonWatch (Ramaswami *et al*. 2021). SeasonWatch

is a pan-india project monitoring the phenology of common tropical tree species wherein,

citizen scientists make quantitative assessment of tree phenophases on a weekly basis,

observing registered trees again and again with the aim of understanding long-term changes

in tree seasonality (Ramaswami *et al.* 2021).

In this study, we explore the effect of latitude on the flowering phenology of three wide-

ranging tree species *Bombax ceiba L., Butea monosperma (Lam.) Taub,* and *Cassia fistula*

L. Bombax ceiba (Red Silk Cotton tree), belonging to the family Malvaceae, is a large

deciduous tree, easily identifiable by its spiny bark and large pentamerous red flowers. The

tree typically blooms before the onset of summer and attracts a large diversity of pollinators

such as birds, bats and bees (Raju A. J. S., 2005). *Butea monosperma* (Flame-of-the-Forest

tree), belonging to the family Fabaceae, is found in dry-deciduous, seasonal forest habitats

of India. The bright red flowers of *Butea monosperma* are frequented by as many as seven

species of birds, belonging to six families and one species in particular, the purple sunbird

(*Nectarinia asiatica*), is an effective pollinator (Tandon *et al*. 2003). *Cassia fistula* (Indian

Laburnum tree)*,* also a Fabaceae member, has a wide geographical range, is native to the

Indian subcontinent, and produces bright yellow flowers in large, pendulous inflorescences.

Xylocopa bees (Carpenter bees) are known to be the most effective pollinator of *Cassia*

fistula as they show high specificity towards this particular plant species (Kallur, 1993). All

three species find use in local cuisine, medicine, and cultural practices, and because of their

bright, showy flowers are likely to be identified accurately, observed frequently, and

documented widely in media repositories and citizen science.

2. Methods:

We used two data sources for understanding phenological patterns of these three species: 1)

images of flowering trees available publicly on social media and other image repositories,

and 2) phenology information submitted by citizen scientists to SeasonWatch. We expected

latitudinal patterns in flowering to be discernible in both datasets, with the date of

observation showing correlation with the latitude at which the phenophase was observed.

We expected patterns observed through one-time phenological observations (images) to be

supplemented long-term repeated observations from SeasonWatch to infer advantages of

either method in assessing tree phenology.

In the first data source (dataset 1 hereon) - we collated images uploaded for *Bombax ceiba*,

Butea monosperma and *Cassia fistula* on social media platforms (Facebook, Twitter,

Instagram); image-based citizen science portals (Indian Biodiversity Portal, iNaturalist), and

digital photo repositories (Flickr) between 1 January 2018 and 30 November 2021. All these

platforms allow contributors to provide metadata on species ID, location and date of

observation of a photograph. Images were searched through keywords such as scientific and

common names of the species in English and other Indian languages, and through hashtags

where appropriate. Images which included accurately identified plants with recognisable

phenophases, and had reliable metadata such as the date of observation/image, and location

were considered appropriate for the assessment of the relationship between tree phenology

and latitude. We noted the presence of 4 phenophases - young leaves, mature leaves, unripe

fruits and ripe fruits, in addition to the presence of 2 stages of flowers - buds and open

flowers. Only images with the two flowering phenophases were considered for further

assessment. We found 276 images of *Bombax ceiba*, 350 images of *Butea monosperma*, and

247 images of *Cassia fistula* in which the flowering phenology was clearly identifiable. We

assessed flowering patterns visually by plotting the day of the year on which an image was

taken, against the latitude of observation.

The second data source was the SeasonWatch database[19] which was queried for the three

species of interest. Citizen scientists register trees with the project and record phenophases

(same as in dataset 1) as 'none', 'few', or 'many' based on the proportion of the tree canopy

covered by the phenophase. We assessed flowering visually by plotting the day of the year on

which an observation of 'many' flowers was made (indicating peak flowering), against the

latitude of the observation. While both datasets allow for looking at latitudinal patterns in

flowering, SeasonWatch data additionally allows for observing flowering patterns across the

same individuals over time. We used repeated observations from 762 *Bombax ceiba,* 371

Butea monosperma, and 3329 *Cassia fistula* trees between 1 January 2018 and 30 November

2021 in the SeasonWatch database to assess the date of observation of flowering against the

latitude of observation.

3. Result and Discussion:

We did not combine information from dataset 1 and SeasonWatch as the raw information

was collected in a non-comparable way (i.e. image and quantity of phenophase). In dataset

1, images of flowering *Bombax ceiba* trees appear between days 330 of the previous year

and day 90 of the subsequent year, indicating December - March as the flowering season

for this species (Figure 1a). In more southern latitudes (for example: 8.62°N-16.7°N)

flowering trees were photographed from December - February while in more northern

latitudes flowering trees were photographed in February-March, sometimes extending till

May (Figure 1a). More continuous data from SeasonWatch indicates that at lower latitudes

a few trees may be flowering throughout the year, while at latitudes above 9 degrees, most

trees flowered between the 300th day of one year and the 150th day of the next (Figure 2a).

A slightly more pronounced effect of latitude is seen for flowering in dataset 1 *Butea*

monosperma, with most images appearing across latitudes between December (day 350)

and March (day 90), sometimes extending till April or May (Figure 1b). SeasonWatch data

indicate that trees have shown 'many' flowers as early as mid-year in latitudes below 13

degrees, but in more northern latitudes peak flowering occurred within the first 100 days of

the year (Figure 2b). Images of flowering *Cassia fistula* trees occurred in dataset 1 almost

throughout the year. Most images of flowering *C. fistula* trees across the 3.5 years were

taken between the 50th and 200th day of the year (i.e. between end-February and July,

Figure 1c). In SeasonWatch dataset, *C. fistula* trees were observed to flower almost

throughout the year between 8- and 13-degrees N latitude, while flowering peaked more

seasonally at higher latitudes (Figure 2c).

Across both datasets, the probabiltity of finding flowering phenology dates were

consistently more in the first quarter of the year for *Butea monosperma* and *Bombax ceiba*,

but staggered by latitude for *C. fistula* , when data was pooled across all years

(Supplementary figure 1 and Supplementary figure 2). In the SeasonWatch dataset, C.

Fistula appears to be more variable than the other two species, whereas in public media

repositories, all three species show similar amount of variability in dates of flowering. On

comparing the patterns from the two datasets, we infer that continued monitoring of

individual trees allows for a better understanding of the flowering season of trees across

latitudes - as individual-level variation is better captured through repeated observations. We

do not have adequate understanding of what environmental factors could be driving

phenological variation in these species across latitudes from this study alone, and speculate

that as seen in other species, temperature and precipitation may be key drivers. If these

species have a temperature threshold that triggers flowering then one could expect the

flowering season to increase in duration across latitudes over the next few years as global

temperatures increase (Rathcke *et al*. 1985). Citizen science data on flowering and leaf-out

patterns elsewhere have demonstrated early onset of vegetative and reproductive

phenophases with increasing mean spring temperatures (Fucillo Battle *et al* 2022).

However, the wide variation in existing phenological strategies of tropical species such as

the ones examined in this study, are likely to make temperature-phenology relationships

less discernible than temperate species.

Images in dataset 1 are, at best, one-time snapshot samples of single individuals and may not

represent the entire variation in the phenological behaviour of the study species. Furthermore,

contributor behaviour is likely to affect what images are posted publicly, and which May in-

turn bias the data available on these platforms. For instance, contributors may be more likely

to post photos of clearly visible flowers and of trees in full bloom, than any given instance of

flowering, particularly in the species examined. This may create a bias towards a more

visually identifiable stage of the flowering phenopahse, than the entire duration of the

phenology. Contributor behaviour may also be affected by other factors such as access to

smart phones or cameras to take pictures, or access to public groups and repositories, because

of technological or linguistic constraints. Therefore, dataset 1, may have lower utility in

understanding future changes in tree phenology with changing climate as compared to

SeasonWatch data that has multiple observations of individuals in the same locations over

time. However, in order to utilize citizen science data effectively, high data quality needs to

be ensured. Phenology citizen science assessed elsewhere has found disparities between

citizen contributed data and actual observations (MacEnzie *et al.* 2017). Although a similar

assessment, we conclude that both approaches to observing tree phenology can be used to

supplement existing phenological information on the species, or to arrive at the baseline

patterns of phenology to aid in designing sampling strategies for future studies.

Acknowledgements:

Bhavya Kriti would like to thank the Indian Academy of Science for supporting this study.

SeasonWatch is supported by the Wipro Foundation. We would like to thank the hundreds of

citizen scientists who contributed the data to the SeasonWatch database and facilitated this

comparative study.

Conflict of Interest:

The   authors   do   not   have   any   conflict   of   interest   to   report.

References:

Abu-Asab M S, Peterson P M, Shetler S G, *et al* 2001 Earlier plant flowering in spring as a response

to global warming in the Washington, DC, area. *Biodivers. Conserv.* **10** 597–612

Attri S D and Tyagi A 2010 Climate profile of India. *IMD*

Badeck F W, Bondeau A, Böttcher K, *et al*. 2004 Responses of spring phenology to climate change.

New Phytol. **162** 295–309

Barve V 2014 Discovering and developing primary biodiversity data from social networking sites: A

novel approach. *Ecol. Inform.* **24** 194–199

Burli D and Khade A B 2007 A comprehensive review on *Butea monosperma* (Lam.) Kuntze. *Phcog.*

Rev. **1** 333–337

Chaudhary P H and Khadabadi S S 2012 *Bombax ceiba* Linn.: Pharmacognosy, ethnobotany and

phyto-pharmacology. *Phcog. Commn.* **2** 2–9

Cleland E E 2007 Shifting plant phenology in response to global change. *Trends Ecol. Evol.* **22** 357–

365

Gordo O and Sanz J J 2010 Impact of climate change on plant phenology in Mediterranean

ecosystems. *Glob. Change Biol.* **16** 1082–1106

Jain V, Verma S, Katewa S S 2009 Myths, traditions and fate of multipurpose *Bombax ceiba* L. – An

appraisal. *Indian J. Tradit. Knowl*. **8** 638–644

Kallur M 1993 Differential reproductive success in *Cassia fistula* in different habitats – a case of

pollinator limitation? *Curr. Sci.* **65**

Kudo G and Ida T Y 2013 Early onset of spring increases the phenological mismatch between plants

and pollinators. *Ecol.* **94** 2311–2320

Neil K and Wu J 2006 Effects of urbanization on plant flowering phenology: A review. *Urban*

Ecosyst. **9** 243–257

Numata S, Yamaguchi K, Shimizu M., *et al*. 2022 Impacts of climate change on reproductive

phenology in tropical rainforests of Southeast Asia. *Commun. Biol.* **5** 1–10

Raju A J S, Rao S P, and Rangaiah K 2005 Pollination by bats and birds in the obligate outcrosser

Bombax ceiba L. (Bombacaceae), a tropical dry season flowering tree species in the Eastern Ghats

forests of India. *Jorn.* **4** 81–87

Ramaswami G, Sidhu S, and Quader S 2021 Using citizen science to build baseline data on tropical

tree phenology. *Curr. Sci.* **121** 1409–1416

Rathcke B and Lacey E P 1985 Phenological patterns of terrestrial plants. *Annu. Rev. Ecol. Evol. Syst.*

16 179–214

Renner S S and Zohner C M 2018 Climate change and phenological mismatch in trophic interactions

among plants, insects, and vertebrates. *Annu. Rev. Ecol. Evol. Syst.* **49** 165–182

SeasonWatch Citizen Scientist Network 2023 Data on tree phenology. Data Type: 01/01/2018–

30/11/2021, All India, *Bombax ceiba*, *Butea monosperma*, *Cassia fistula*. Date accessed: 27/02/3023.

Sindhia V and Bairwa R 2010 Plant review: *Butea monosperma*. *Int. J. Pharm. Clin. Res.* **2** 90–94

Tandon R, Shivanna K R, Mohan Ram H Y 2003 Reproductive biology of *Butea monosperma*

(Fabaceae). *Ann. Bot.* **92** 715–723

Walther G R, Post E, Convey P, *et al* . 2002 Ecological responses to recent climate change. *Nature*

416 389–395

 FIGURE LEGENDS

Figure1: Phenology patterns from dataset 1 - date of upload of images of a flowering tree of

A) *Bombax ceiba*, B) *Butea monosperma*, and C) *Cassia fistula*, against latitude at which the

image was taken.

Figure2: Phenology patterns from SeasonWatch - date of observation of a tree in peak

flowering of A) *Bombax ceiba*, B) *Butea monosperma*, and C) *Cassia fistula*, against latitude

at which the observation was made.

SUPPLEMENTARY MATERIALS

Supplementary figure 1: This figure shows the distribution of dates on which images of flowering

were taken across latitudes for three different species on public media repositories over a span of three

years (2018-2021). A) *Bombax ceiba* was consistently photographed across latitudes in the first 100

days of the year with southern latitudes showing an additional peak in December, B) *Butea*

monosperma, showed a similar pattern to *B. ceiba* with most amount of photos were observed across

latitudes in the first 100 days of the year, C) *Cassia fistula* showed a slightly more prominent

latitudinal pattern, with the peak in flowering photos shifting between the 100th day at southern

latitudes to the 150th day at more northern latitudes.

Supplementary figure 2: *This figure shows the distribution of dates on which species were*

reported to have "many" flowers across different latitudes in SeasonWatch between 2018

and 2021. A) Bombax ceiba flowering is represented by a sharp peak within the first 100 days

of the year across latitudes, B) Butea monosperma flowering occurs mostly in Jan-March

with a sharper peak discernible in northern latitudes, C) Cassia fistula is markedly less

seasonal in this dataset, especially in southern latitudes, as compared to the public repository

dataset, but shows a similar shift in peak flowering.