Spatial single-cell isotope tracing reveals heterogeneity of *de novo* fatty acid synthesis in cancer cells

Elena Buglakova[1], Marteinn Snaebjornsson[2], Maans Ekeloef[1], Lisa Schlicker[2], Martijn R. Molenaar[1], Shahraz Mohammed[1], Lachlan Stuart[1], Andreas Eisenbarth[1], Volker Hilsenstein[1], Almut Schulze[2,+], Theodore Alexandrov[1,3,4,+]

1 Structural and Computational Biology Unit, European Molecular Biology Laboratory (EMBL), Heidelberg, Germany

2 German Cancer Research Center, Division of Tumor Metabolism and Microenvironment, Im Neuenheimer Feld 581, 69120 Heidelberg, Germany

3 Metabolomics Core Facility, EMBL, Heidelberg, Germany

4 BioStudio, BioInnovation Institute, Copenhagen, Denmark

+ corresponding authors

Email: theodore.alexandrov@embl.de, almut.schulze@dkfz-heidelberg.de

Keywords: spatial single-cell metabolomics, stable isotope tracing, single-cell analysis, fatty acid synthesis, acetyl-CoA production, ACLY, cancer

Abstract

Metabolism has emerged as a key factor in homeostasis and disease including cancer. Yet, little is known about the heterogeneity of metabolic activity of cancer cells due to the lack of tools to directly probe it. Here, we present a novel method, ^{13}C-SpaceM for spatial single-cell isotope tracing of glucose-dependent *de novo* lipogenesis. The method combines imaging mass spectrometry for spatially-resolved detection of ^{13}C$_6$-glucose-derived ^{13}C label incorporated into fatty acids composing cellular lipids with microscopy and computational methods for data integration and analysis. We validated ^{13}C-SpaceM on a spatially-heterogeneous normoxia-hypoxia model of liver cancer cells. We demonstrated the single-cell heterogeneity of acetyl-CoA pool labelling degree upon ACLY knockdown that is hidden in the bulk analysis and its effect on synthesis of individual fatty acids. Overall, ^{13}C-SpaceM provides a capacity to probe metabolic activity on the single-cell level.

Introduction

Lipids are a highly complex class of hydrophobic biomolecules that are involved in multiple cellular functions. For example, glycerophospholipids and sphingolipids are major structural components of cellular membranes [1] while triglycerides stored in lipid droplets provide cells with an additional energy source [2]. Lipids can also contribute to cellular signaling pathways by functioning as second messengers or lipid mediators [3-5]. Altered lipid metabolism is a hallmark of many cancers, as it is a prerequisite for rapid growth and proliferation and promotes survival under nutrient limiting conditions [6-8]. Altered lipid metabolism in cancer also contributes to oxidative stress resistance and activates oncogenic signaling pathways through the production of lipid mediators [9]. Lipid metabolism is also affected by different stresses imposed by the tumor microenvironment, such as limited nutrient and oxygen supply [10]. Therefore, cancer cells need to adapt to these stresses by adopting different routes for the provision of lipids, thus inducing selective metabolic dependencies that could be targeted for cancer therapy.

Most lipids are synthesized from fatty acids, a group of small linear molecules consisting of a carboxylic acid head and an aliphatic tail (Fig. 1a). Fatty acids differ in their length and degree of saturation, i.e. the number of double bonds located in different positions of the hydrocarbon chain [11]. Cells in the adult mammalian organism generally take up fatty

acids from the blood stream. However, cells of adipose and hepatic tissues as well as
various cancers can also generate fatty acids through *de novo* synthesis using acetyl-
coenzyme A (acetyl-CoA) as a substrate [7]. The first step in fatty acid biosynthesis consists
of the conversion of acetyl-CoA into malonyl-CoA by acetyl-CoA synthetases (ACC1/2).
Malonyl-CoA is then used by fatty acid synthase (FASN), a large enzyme with multiple
catalytic centers, for the repeated condensation reactions of two-carbon units, derived
from acetyl-CoA, to form palmitate (16:0), a 16-carbon saturated fatty acid. Palmitate can
then be further elongated and/or desaturated thus generating a variety of saturated and
monounsaturated fatty acids that can subsequently be incorporated into the lipidome. The
fatty acid composition (or saturation state) of the lipidome is a critical factor for cell survival
under metabolically challenging conditions. An example of this is the mono-unsaturated
fatty acid oleate which protects cancer cells from lipotoxicity and ER stress [12-15], inhibits
apoptosis [16] and promotes ferroptosis resistance [17]. Indeed, is has been shown that
hypoxic and Ras-transformed cells preferentially take up mono-unsaturated fatty acid to
support proliferation and survival [18].

The cytoplasmic (lipogenic) acetyl-CoA pool used as a substrate for fatty acid synthesis
can be derived from different precursors through specific metabolic routes. Glucose-
derived carbons provide fuel for the production of citrate by the TCA cycle in the
mitochondrial matrix. This citrate is then exported from the mitochondria and used in the
cytoplasm by ATP citrate lyase (ACLY) to generate acetyl-CoA. However, the use of
glucose as a precursor for fatty acid synthesis is limited under certain conditions, such as
hypoxia, in which glucose-derived pyruvate is diverted away from mitochondrial
metabolism and converted to lactate. Under these conditions, glutamine can substitute
as a precursor for fatty acid synthesis through the reductive carboxylation of glutamate to
citrate [19, 20]. Furthermore, hypoxic cancer cells also use acetate as a precursor for fatty
acid biosynthesis through acetyl-CoA synthetase 2 (ACSS2) [21, 22].

Besides its function as a substrate for fatty acid synthesis, acetyl-CoA is also needed for
the post-translational modification of proteins, most importantly histones for chromatin
regulation [23]. Indeed, both ACLY and ACSS2 have been found to localize to nucleus and

modulate histone acetylation and thus affect gene expression [24, 25]. The modulation of

acetyl-CoA metabolism by oncogenic signaling pathways and environmental factors can

thereby affect complex gene expression programs in cancer cells. Monitoring the interplay

between fatty acid synthesis and acetyl-CoA metabolism therefore provides vital

information on the state of cancer cells.

The advent of single-cell tools has revolutionized the analysis of individual cells within

heterogenic populations [26]. While metabolomics provides much more dynamic

information about cellular states, its application at single-cell level has somewhat lagged

behind single-cell transcriptomic and proteomic techniques, due to the immense technical

challenges [27]. The high metabolic heterogeneity of cancer cells makes it essential to

develop analysis tools that are capable of interrogating fatty acid synthesis and acetyl-

CoA metabolism in cancer cells on the single-cell level. Different experimental methods

have been applied to evaluate the heterogeneity of lipid metabolism in cancer cells. For

example, a metabolic sensor based on the malonyl-CoA responsive Bacillus subtilis

transcriptional repressor, FapR, was developed to determine the activity of FASN in live

cells [28]. Another approach used surface-immobilized dendrimers to evaluate fatty acid

uptake on a single cell level [29]. Furthermore, a recent study applied dielectric barrier

discharge ionization (DBDI) on single cells to determine alterations in lipid metabolism

upon ACLY inhibition in pancreatic cancer cells [30]. However, these methods fail to provide

information about metabolic precursors of fatty acid synthesis and do not afford sufficient

molecular resolution to interrogate acetyl-CoA metabolism.

We previously developed a spatially resolved mass-spectrometry pipeline for the

quantification of intracellular metabolites at single-cell resolution [31]. Stable-isotope tracing

has been used to determine *de novo* fatty acid biosynthesis and the contribution of

different metabolic precursors to the cytoplasmic acetyl-CoA pool [21, 32]. We now describe

^{13}C-SpaceM, a method that combines stable isotope tracing with the detection of fatty

acids present as acyl-chains in selected lipid classes in a spatially resolved manner, using

MALDI-MS2. We applied this method to interrogate the effect of hypoxia on *de novo* fatty

acid synthesis in a population of murine liver cancer cells. Moreover, we also developed

a method to evaluate the effect of genetic depletion of ACLY on the contribution of glucose

to the cytoplasmic acetyl-CoA pool on a single cell level. Our results show that ^{13}C-

SpaceM can reveal substantial heterogeneity in fatty acid synthesis and acetyl-CoA

metabolism in cancer.

Material and Methods

Cell culture for the normoxia-hypoxia model

Primary murine liver cancer cells derived from a Myc- and Akt-driven tumor model (Myc^{OE};

Akt^{Myr}; $Tp53^{-/-}$) [15] were a gift from Daniel Dauch (University of Tübingen). Cells were

cultured in DMEM supplemented with 1 mM acetate, 2 mM glutamine and 10% FCS in a

37°C incubator with 5% CO_2 either under normoxia (20% O_2) or hypoxia (0.5% O_2) in the

presence of $^{13}C_6$-glucose for 72 hours. Hypoxic conditions (0.5% O_2) were induced in a

hypoxia workstation (H35, Don Whitley). The cells cultured under hypoxic conditions were

modified to express Green Fluorescent Protein (GFP) as a marker. For ^{13}C-SpaceM, the

normoxic and hypoxic cells were mixed using 10,000 cells of each condition, plated on

the same glass slide and allowed to attach for 3 hours before fixation.

ACLY knockdown

For the ACLY knockdown, we used the same primary murine liver cancer cells as in the

hypoxia model. shRNA sequences targeting murine Acly or non-targeting controls were

cloned into LT3-GEPIR (Addgene), the shRNA sequences used were

(TGCTGTTGACAGTGAGCGACCGCAGCAAAGATGTTCAGTATAGTGAAGCCACAGATGTATACTGAACATCTTTGCTGCGGCTGCCTACTGCCTCGGA),

(TGCTGTTGACAGTGAGCGAACCAGTGTCTACTTATGTCAATAGTGAAGCCACAGATGTATTGACATAAGTAGACACTGGTCTGCCTACTGCCTCGGA),

(TGCTGTTGACAGTGAGCGCAGGAATTATAATGCTTATCTATAGTGAAGCCACAGATGTATAGATAAGCATTATAATTCCTATGCCTACTGCCTCGGA) for

ACLYkd oligo 1, oligo 2 and nontargeting control, respectively. Lentiviral particles were

produced in HEK293 cells after transient transfection of the packaging vectors psPAX.2

and pMD.G2 (Addgene #12260 and #12259). After viral transduction, liver cancer cells

were selected with puromycin and used at low passage. Induction of shRNA expression

was achieved by treating cells with 1µg/ml doxycycline (Sigma) for 72 hours. For stable

isotope tracing, cells were cultured in glucose-free DMEM supplemented with U$^{13}C_6$-

glucose (Cambridge Isotope laboratory, Tewksbury, Massachusetts) and 1mM acetate
for 72hrs.

Liquid-chromatography coupled with mass spectrometry (LC-MS)

For bulk LC-MS, cells were washed with cold 154 mM ammonium acetate, snap frozen

in liquid nitrogen and harvested in methanol/H_2O (80/20, v/v) with added standards (10

µL of 100 µM Palmitate-2,2-D2 (Eurisotop: DLM-1153-0)/ 1 x 10E6 cells). 30 µL 0.2 M

HCl, 2 x 100 µL $CHCl_3$, 2 x 100 µL H_2O was then added with vortexing in between. The

suspension was centrifuged at 16,000 g for 5 min, the lower lipid phase was then washed

with synthetic polar phase (CH_3Cl/Methanol/H_2O, 58/33/8, v/v/v) and evaporated to

dryness under N_2 at +45°C. For saponification, lipids were resuspended in Methanol/H_2O

(80/20, v/v) containing 0.3 M KOH, heated at +80°C for 1 h and washed twice with 0.5

mL hexane. After addition of 50 µL formic acid, fatty acids were subsequently extracted

twice with 0.5 mL hexane and evaporated to dryness under N_2 at +45°C. For LC/MS

analysis the fatty acids, were dissolved in 100 µL isopropanol and 5 µL of each sample

was applied to a C8 column (Acclaim RSLC C8 column, 2.2 µm particles, 50 ✕ 2.1 mm,

Thermo Fisher Scientific) at +40°C, with mobile phase buffer A consisting of 0.1% formic

acid in CH_3CN/H_2O (10/90, v/v), and solvent B consisting of 0.1% formic acid in

CH_3CN/H_2O (90/10, v/v). The flow rate was maintained at 350 µL/min and eluent was

directed to the ESI source of the QE-MS from 3 min to 27 min after sample injection. MS

analysis was performed on a Q Exactive Plus Orbitrap mass spectrometer (Thermo

Fisher Scientific) applying the following settings: Sheath gas, 30; spray voltage, 2.6 kV.

Capillary temperature +320°C, aux gas heater temperature: +120°C and S-lens voltage

was 55. A full scan range from 150 to 460 (m/z) in negative ion mode was used. The

resolution was set at 70,000. The maximum injection time was 100 ms with an AGC-target

of 1E6 ions. Peaks corresponding to the calculated fatty acid masses ($\pm$5 ppm) were

integrated using El-Maven (https://resources.elucidata.io/elmaven) and correction for

natural ^{13}C isotopic abundance was done using IsoCorrectoR [33].

SpaceM Analysis

The SpaceM [31] method for fusion of microscopy and MALDI-MS spatial metabolomics data was applied to analyze the de-novo synthesis of fatty acids on a single-cell level.

Microscopy

After cell culturing, the cells were washed with PBS, fixed using Histofix (Roth, 87.3) for 10 minutes, stained using DAPI, washed 3x in PBS and then desiccated in a Lab Companion Cabinet Vacuum Desiccator for 30 min at room temperature and −0.08 MPa. Pre-MALDI brightfield and fluorescent microscopy (620 and 460 nm) images were obtained with a DS-Qi2 camera (Nikon Instruments) with a Plan Fluor 10x (numerical aperture (NA) of 0.30) objective (Nikon Instruments) mounted on a Ti-E inverted microscope (Nikon Instruments). The pixel size was 0.64 µm. For the hypoxia experiment, additional pre-desiccation microscopy images were collected. Rigid registration of pre-desiccation and post-desiccation microscopy images was performed using the Affinder plugin in Napari (https://www.napari-hub.org/plugins/affinder). The cells were imaged in bright-field microscopy after MALDI-imaging, using the same microscopy setup and parameters.

Microscopy image analysis

Cells in the bright-field channel of pre-MALDI microscopy were segmented using a custom Cellpose model [34] trained on manually annotated fragments of images. Since the primary cell culture contains cells of varying size, segmentation for the largest cells had to be manually corrected. Ablation marks were detected in the post- MALDI microscopy by manually fitting a grid of circular shapes. Registration of the pre- and post-MALDI images was done using the SpaceM software tool as previously published [31].

MALDI imaging

The MALDI matrix 1,5-diaminonaphthalene (DAN) was applied to a surface density of 3 µg/µm^2 immediately before MALDI analysis using a TM sprayer (HTX Technologies, Chapel Hill, NC, USA). Atmospheric pressure MALDI imaging was performed using an AP-SMALDI5 ion source (TransMIT GmbH, Gießen, Germany) coupled to a Q Exactive Plus Orbitrap mass spectrometer (Thermo Fisher Scientific). For all experiments the

201 raster pitch was 10x10 μm, with the laser attenuator angle set to 33°. The mass
202 spectrometer method used was set up as all-ion fragmentation (AIF) in negative ion
203 mode, with an isolation range of 600-1000 and a scan range of 100-400 at 140,000
204 nominal resolving power with 500 ms ion injection time. Fragmentation was performed
205 using higher-energy collisional dissociation (HCD) at a normalized collision energy of 25.

Constructing single-cell isotopologue profiles

MALDI images for unlabeled samples were annotated with the cloud software
METASPACE (http://metaspace2020.eu) at 5% FDR to determine the most abundant
detected fatty acids. For each fatty acid, intensities corresponding to theoretical
isotopologue peak masses were extracted. Raw intensities were normalized for natural
isotope abundance using IsoCor [35], after which every isotopologue distribution was
normalized by its sum. Ablation marks with a total raw intensity of less than 200 for a
given fatty acid were removed from the analysis. Ablation marks which had at least 30%
overlap with the cell mask were considered intracellular. After normalizing spectra in each
pixel, the median of values in the intracellular ablation marks was assigned as a single-
cell readout for each isotopologue peak. The resulting single-cell isotopologue distribution
for every fatty acid was normalized by its sum again. Scanpy v.1.8.1 [36] was used for all
single-cell data analysis.

Calculation of single-cell features

For each well of a multi-well slide, the median intensity of the GFP channel outside of the
cell masks was subtracted from the image, and the logarithm of the maximum GFP
intensity inside the cell mask was used to characterize single-cell GFP signal. Both in
normoxia/hypoxia and in ACLYkd experiments, the 95th percentile of the GFP signal in
the GFP-negative wells was used as a threshold to assign cells to either GFP-positive or
GFP-negative state.

In addition to the normalized isotopologue peak intensities, a binomial model for the fatty
acid synthesis described in [32] was used. Fatty acids are synthesized by randomly taking
2-carbon monomers from the cytosolic AcCoA pool. Therefore, if the fraction of labeled
monomers equals p, the probability that a given fatty acid has 2i labeled carbons, can be
described with the following model:

$$P_{binom}(k) = \binom{n}{k} p^k (1-p)^{n-k}, k = 0,1, \dots n$$

$$I_0 = uptake + (1-uptake) * P_{binom}(0)$$

$$I_i = (1-uptake) * P_{binom}(i), i = 1,2, \dots n$$

where uptake is the fraction of the unlabeled FA directly taken from the media, *p* is the

labeling degree of cytosolic AcCoA pool, and n is the number of AcCoA molecules used

for the synthesis (number of carbons in the fatty acid / 2). Longer fatty acids can be

synthesized both using labeled palmitate made by the cell *de novo* and unlabeled

palmitate, therefore we used a modified model for C18:

$$I_0 = uptake + (1-uptake) * P_{binom}(0)$$

$$I_1 = (1-uptake) * \left(uptake_{C16} + (1-uptake_{C16}) * P_{binom}(1)\right)$$

$$I_i = (1-uptake) * (1-uptake_{C16}) * P_{binom}(i), i = 2,3, \dots n$$

where *uptake_{C16}* is the fraction of palmitate taken from the medium.

Fitting a binomial model to the single-cell isotopologue distributions allows summarizing

them as two numbers: uptake, which characterizes the fraction of the fatty acid which was

not synthesized *de novo* but directly taken from the medium, and AcCoA pool labeling

degree *p*, which describes relative contribution of the labeled substrate to the fatty acid

synthesis compared to other carbon substrates consumed by the cell. The limitation of

this method is that if overall substrate usage is very low, such as in the hypoxia/normoxia

experiment, it becomes impossible to reliably fit a binomial distribution and estimate the

uptake fraction, therefore it was only used for the ACLYkd experiment data analysis.

Results

¹³C-SpaceM, a method for single-cell isotope tracing

The key development presented in this paper is ¹³C-SpaceM, a method for single-cell isotope tracing. ¹³C-SpaceM builds upon SpaceM, a method for single-cell metabolomics which integrates imaging mass spectrometry (imaging MS) for the detection of stable isotope-labeled metabolites and lipids, and microscopy for assigning imaging MS pixels to individual cells and for quantifying fluorescence and morphometric properties of single cells [31]. In contrast to the conventional SpaceM method using MS1, in ¹³C-SpaceM we exploited a particular type of mass spectrometry, namely Higher-Energy Collision-induced Dissociation All Ion Fragmentation (HCD AIF), where lipids in the range of 600-1000 m/z in the negative mode were simultaneously fragmented and then the fragments in the range of 200-400 m/z were detected (**Fig. 1b**). This allowed us to interrogate fatty acid composition of the cellular lipids. The selected mass range includes most glycerophospholipids (PC, PE, PS, PG, PA, PI), sphingolipids (SM, Cer, HexCer, CerP), neutral lipids such as but does not include smaller lipids such as lysophospholipids or monoradylglycerolipids nor larger lipids such as triglycerides, cardiolipins or PIP-trisphosphates [37]. The exploited negative ion polarity favors detection of phospholipids and sphingolipids. The construction of single-cell profiles was done as in the conventional SpaceM (**Fig. 1c**); see Methods for details.

Spatially-heterogeneous normoxia-hypoxia model for evaluating single-cell isotope tracing

In this work, we focused on applying 13C-SpaceM to interrogate lipid biogenesis through *de novo* fatty acid synthesis, a highly dynamic metabolic process that integrates multiple substrates and contributes to the complexity of cellular lipids (**Fig. 1a**). In order to validate ¹³C-SpaceM in this context, we developed a controlled spatially-heterogeneous model containing co-plated cells of two distinct phenotypes: cells cultured under either normoxic or hypoxic conditions. We chose this model as hypoxia causes global reprogramming of cellular metabolism, including lipid metabolism. Specifically, hypoxic cells undergo a switch in the use of substrates for the production of cytoplasmic acetyl-CoA used by fatty

acid synthesis, by promoting the reductive carboxylation of glutamine to form citrate [19, 20,

38] or by the direct use of acetate [22]. Moreover, hypoxia blocks fatty acid degradation and

induces fatty acid uptake and lipid storage [39]. We therefore anticipated that cells cultured

under hypoxia should show decreased incorporation of glucose-derived stable-isotope-

labeled carbon atoms into cellular lipids compared to cells cultured under normoxic

conditions. As fatty acid synthesis utilizes the cytoplasmic acetyl-CoA pool, analysis of

the isotopologues distribution of fatty acids incorporated into cellular lipids allows the

assessment of changes in the relative contribution of different substrates to the synthesis

of cytoplasmic acetyl-CoA [21]. The workflow employed by ^{13}C-Space allows the

quantification of all isotopologues detected for fatty acids released from total cell lipids

isolated at a mass range of m/z 600-1000 in the negative mode using all-ion fragmentation

(**Fig. 1b**). Registration of pre- and post-MALDI microscopic images to the MALDI-imaging

spectra allows the assignment of ion intensities to individual cells (**Fig. 1c**).

^{13}C-SpaceM provides isotopologues profiles for single cells, integrated with

microscopy

Pre-MALDI microscopy images overlaying the brightfield and GFP channels for the co-

plated normoxic (GFP-negative) and hypoxic (GFP-positive) cells demonstrate a high

degree of spatial heterogeneity of these two conditions (**Fig. 2a, first plot**). Cell

segmentation and quantification of cellular GFP provides single-cell fluorescence

intensities (**Fig. 2a**, **second plot**). The single-cell isotopologues profiles can be

constructed by ^{13}C-SpaceM for individual fatty acids incorporated into cellular lipids, for

example for palmitate, which is the key intermediate metabolite of the *de novo* fatty acid

synthesis. An important parameter which can be calculated based on an isotopologues

profile is the fraction of the unlabeled fatty acid, namely the fraction of M+0 compared to

other isotopologues. A high fraction of unlabeled palmitate corresponds to low *de novo*

fatty synthesis and lipid biogenesis from glucose. Figure 2a (third and fourth plots)

illustrate the fraction of unlabeled palmitate as measured by imaging MS in pixelated

manner and as calculated for single cells using ^{13}C-SpaceM.

Figure 2b shows the underlying mass spectra collected at two representative pixels, one
corresponding to a cell cultured under normoxia (spectral intensities shown in grey, cell
outline shown in white) and another corresponding to a cell cultured under hypoxia
(spectral intensities and cell outline shown in green); the intensities for the latter are
shown in the negative scale for better visualization. For each spectrum, the highest peak
corresponds to the unlabeled palmitate (M+0), with additional peaks for higher
isotopologues e.g. M+2, M+4, M+6 are marked by circles. The peaks not marked by
circles are unrelated to palmitate isotopologues based on the accurate m/z matching
within ±3 ppm. Notably, most palmitate isotopologues are present only in the spectrum of
the pixel derived from a normoxic cell. The mass spectrum from the hypoxic cell shows
an increased peak for the M+0 mass and only two minor additional isotopologues (M+2
and M+10).

^{13}C-SpaceM reveals reduced *de novo* fatty acid synthesis under hypoxia

Before we applied ^{13}C-SpaceM to the spatially-heterogeneous mix of normoxic and
hypoxic cells, we interrogated each condition separately. This was performed to validate
that the GFP readout is informative of the condition (GFP+ for hypoxia vs GFP- for
normoxia) and to evaluate whether isotopologue distribution at the single-cell level can
reveal reduced glucose contribution to *de novo* fatty acid synthesis and lipid biogenesis
under hypoxia.

Figure 2c shows a single-cell scatterplot showing for each cell its log(GFP intensity)
which is the ground-truth readout characteristic of normoxia (x axis) vs unlabeled
palmitate fraction calculated using the single-cell isotopologue profiles as the normalized
palmitate M+0 intensity (y axis). Palmitate was chosen for this analysis as it is the
immediate product of *de novo* fatty acid synthesis by the enzyme fatty acid synthase
(FASN) and is readily incorporated into cellular lipids, thus representing the most direct
readout of the pathway [7]. Fig. 2c shows that the fractions of unlabeled palmitate are
clearly different between normoxic and hypoxic cells plated separately. Cells cultured
under hypoxic conditions (shown in green) show normalized M+0 intensities close to 0.9,
indicating that only a small amount of palmitate was synthesized from glucose. This

reflects either an overall low rate of *de novo* fatty acid synthesis or a switch from glucose
to other substrates, such as glutamine or acetate [21]. This is opposite to cells cultured
under normoxia, where the proportion of unlabeled palmitate is substantially lower,
indicating higher contribution from [13]C-glucose to palmitate synthesis.

Interestingly, cells cultured under normoxia demonstrate a higher heterogeneity (broad
distribution of normalized M+0 intensities ranging from 0.4 to 0.8) compared to the cells
cultured under hypoxia (narrow distribution of normalized M+0 intensities ranging from
0.85 to 0.95). One explanation for this increased heterogeneity could be that cells cultured
under normoxic conditions are more proliferative. Indeed, proliferating cells switch from
fatty acid synthesis to fatty acid uptake during different phases of the cell cycle [40].
Nevertheless, despite this higher heterogeneity, cells cultured under normoxia displayed
ratios of unlabeled palmitate (< 0.8) that were clearly distinct from those of hypoxic cells
(> 0.85). This demonstrates that *de novo* fatty acid synthesis is active in all normoxic cells,
with a detectable contribution from glucose to the cytoplasmic lipogenic acetyl-CoA pool.

[13]C-SpaceM discerns co-plated single normoxic and hypoxic cells

We next examined glucose-dependent palmitate labeling at the single-cell level of cells
cultured under normoxia and hypoxia using the spatially-heterogeneous co-plated model,
using GFP as the ground-truth for assigning each cell to either normoxia (GFP-) or
hypoxia (GFP+) culturing conditions. Cells with log10 GFP intensity of larger than 3 were
considered to be from the hypoxia condition. These cells showed a large fraction of
unlabeled palmitate (normalized M+0 intensity around 0.9, **Fig. 2d**), indicating low rates
of *de novo* fatty acid synthesis form glucose. However, cells identified as low GFP (log10
GFP intensity less than 3) displayed a mixed phenotype with some cells showing a large
fraction of unlabeled palmitate (normalized M+0 intensity >0.8) while others showed
evidence for palmitate labelling from glucose (normalized M+0 intensity between 0.4-0.8)
(**Fig. 2d**). This can also be seen in the mono-plated scenario (**Fig. 2c**) and is most likely
caused by a proportion of the hypoxic cells having low levels of GFP expression,
potentially due to silencing of the promoter in the expression construct.

Overall, we observed a strong similarity in the values measured for the unlabeled palmitate pool in the mono-plated and co-plated scenarios (**Fig. 2c** cf. **Fig. 2d**). Despite this similarity, the separation of the two cell populations based on unlabeled palmitate was less obvious in the co-plated scenario, as a small proportion of cells show intermediate levels of palmitate labelling (normalized M+0 intensities between 0.75-0.88). This potentially indicates an error of estimating the unlabeled palmitate ratio for the hypoxic cells due to co-sampling of neighboring normoxic cells, and the nature of this error (slight increase of intensities of isotopic peaks at M+1 to M+16, thus reducing the ratio of unlabeled palmitate). However, the presence of two distinct clusters (i.e. cells with high levels of M+0 palmitate vs cells with low levels of M+0 palmitate) and the high similarity between the separate and co-plated cells indicates that the effect of this error is small and still allows a clear separation between the two phenotypes.

We therefore quantified the capacity of ^{13}C-SpaceM and specifically of the ratio of unlabeled palmitate to discern the two considered phenotypes (normoxia vs. hypoxia) in an unbiased manner. For this, we assigned the conditions to single cells by using the levels of the GFP marker at the threshold of log10 of 3 (with values above this threshold indicating the hypoxic condition). After assigning the conditions in this manner, we used the levels of unlabeled palmitate to classify the cells into the two phenotypes, achieving a classification accuracy of 87%. This accuracy validates the capacity of ^{13}C-SpaceM to identify the reduced contribution from glucose to palmitate through *de novo* fatty acid synthesis and thus determine the metabolic activity in a spatially-heterogeneous scenario at single-cell resolution.

^{13}C-SpaceM pseudo-bulk isotopologue profiles for palmitate match bulk results

As an additional step of validation, we compared the single-cell isotopologue profile as provided by ^{13}C-SpaceM with the bulk measurements. For bulk analysis, lipids were extracted from pooled cells individually for each condition using chloroform/methanol extraction [41], which allows the isolation of all major phospholipids as well as neutral lipids

and ceramides [37]. Lipid extracts were subjected to alkaline hydrolysis to release fatty

acids and palmitate isotopologue distribution was determined by LC-MS.

This comparison showed an overall level of similarity, with pseudo-bulk intensities for the

palmitate isotopologues detected by ^{13}C-SpaceM being reproduced by the bulk analysis

in each condition (**Fig. 2e**). Importantly, the expected drastic changes induced upon

hypoxia in the palmitate isotopologue distribution were reproduced in the bulk analysis:

high fraction of unlabeled palmitate in hypoxia (> 0.9) as well as disappearance of

isotopologues higher than M+2. However, there is a noticeable difference in the M+0

fraction detected in normoxic cells (average normalized intensity of 0.6 obtained by ^{13}C-

SpaceM compared to 0.4 detected in the bulk analysis). One reason for this difference

could be that the bulk data represent palmitate released through chemical hydrolysis from

cellular lipids including neutral lipids. In contrast, the collision-induced dissociation of

lipids used by ^{13}C-SpaceM is mostly specific to phospholipids and sphingolipids. Thus,

the two methods interrogate palmitate from somewhat different lipid pools within the cell.

In particular, triglycerides, which are included in the bulk method but excluded in the

single-cell analysis due to it being restricted to the negative mode, could contribute a

significant amount of newly synthesized fatty acids [42]. Consequently, our results imply

that different lipid pools can incorporate newly synthesized fatty acids at different rates,

thereby presenting different proportions of unlabeled molecules.

Summarizing the validation results obtained for the normoxia-hypoxia model, we

conclude that ^{13}C-SpaceM can profile the contribution of glucose into *de novo* fatty acid

synthesis and provide isotopologue distribution reflecting the labeling of palmitate as the

key readout of the activity of this pathway, on the single-cell level.

Inducible ACLY knockdown reduces glucose contribution into *de novo* fatty

acid synthesis

ACLY is a key enzyme of *de novo* fatty acid synthesis as it catalyzes the conversion of

citrate into acetyl-CoA (**Fig. 1a**). We applied ^{13}C-SpaceM to investigate the effect of

inducible ACLY knockdown on *de novo* fatty acid synthesis from glucose. We therefore

engineered cells to express short hairpin RNA sequences (shRNA) targeting murine
ACLY under the control of a doxycycline-inducible promoter. The same promoter also
drives expression of a GFP reporter, allowing identification of shRNA expressing cells.
We used two non-overlapping shRNA sequences (ACLYkd oligo 1 and ACLYkd oligo 2)
as well as a non-targeting control. Silencing was achieved by treating cells with 1 µg/ml
of doxycycline for 72 hours, with both shRNA sequences resulting in a comparable level
of ACLY depletion (**Fig. 3a**).

We next induced ACLY silencing in cells cultured in the presence of $^{13}C_6$-glucose and
interrogated them with ^{13}C-SpaceM. Parallel cultures were analyzed using western blot
(**Fig. 3b**) and bulk LC-MS-based stable isotope tracing. **Figure 3a** shows intensities of
the isotopologues for palmitate released from cellular lipids for the cells expressing non-
targeting shRNA and for cells with ACLYkd induced by either oligo 1 or oligo 2. For the
control cells, the isotopologues M+10 and M+12 have maximal intensities in the bulk and
single-cell analyses (**Fig. 3b**, left graphs). Pseudo-bulk analysis of ^{13}C-SpaceM data
showing averages across all single cells indicates a higher M+0 fraction compared to the
bulk analysis (normalized peak intensity of 0.6 compared to 0.3). This is in line with the
results for the normoxia-hypoxia model in **Figure 2e** and, as discussed earlier, can be
explained by differences in the specific lipid pools interrogated by the two methods. After
ACLY silencing using oligo 1, we observed a marked shift in the isotopologue distribution
with the intensities of isotopologues at M+2, M+4, and M+6 substantially increased (**Fig.**
3a, middle graphs). This is in line with our expectation that ACLY silencing reduces the
contribution of glucose to the cytoplasmic acetyl-CoA pool. ACLY depletion using oligo 2
also induced a shift in the isotopologues distribution, with a peak between M+8 and M+10
(**Fig. 3a, right graphs**).

In contrast to the results obtained by comparing cells cultured under normoxic and
hypoxic conditions, we could not discriminate the effect of knockdown (control vs. oligo 1
vs. oligo 2) based on the unlabeled palmitate ratio alone: the M+0 averages are similar
between all three conditions (with the average values equal to 0.58, 0.52, and 0.62 for
the control, oligo 1, and oligo 2 conditions, respectively). This can be explained by the

overall weaker effect of the induced ACLY knockdown compared to culturing under
hypoxia. However, the intensities of isotopologues from M+2 to M+16 have visibly
different distributions, showing higher intensities for isotopologues with lower mass upon
knockdown when compared to control, clearly indicating a reduction in the glucose
contribution to the cytoplasmic acetyl-CoA pool. Moreover, the impact of ACLY silencing
on the overall contribution of *de novo* synthesis to the palmitate pool was more apparent
in the bulk analysis (with M+0 fraction equal to 0.3, 0.37, and 0.43 for the control, oligo1,
and oligo 2 conditions, respectively). This indicates that the lipid pool interrogated by [13]C-
SpaceM is more dependent on fatty acid uptake.
470

Quantifying acetyl-CoA pool labeling degree for single cells

In **Figure 3a** we observed a pronounced shift towards a lower mass in the isotopologue
distribution of palmitate upon the inducible knockdown of ACLY in both single-cell and
bulk data. This shift indicates an increase of the fraction of unlabeled acetyl-CoA in the
acetyl-CoA pool used for palmitate synthesis and is expected as ACLY catalyzes the
production of cytoplasmic acetyl-CoA from citrate produced in its turn from labeled
glucose [21]. We also noticed that ACLY silencing using shRNA oligo 1 resulted in a much
wider distribution of isotopologues (cf. oligo 2) both in the bulk analysis and the pseudo-
bulk display of the single-cell data. This is visible in **Figure 3a, middle graphs,** as higher
intensities of the M+4 and in particular M+2 isotopologues for oligo 1 compared to oligo
2. The distribution of the pseudo-bulk (or bulk) intensities across the isotopologues for the
oligo 1 data suggests the presence of two distinct populations with distinct responses to
ACLY knockdown, with one population exhibiting a stronger effect of ACLY knockdown
(with higher intensities at M+2 and M+4). We therefore used the power of single-cell
analysis to deconvolve the relative palmitate labelling in individual cells from the three
conditions (control, ACLYkd oligo 1 and ACLYkd oligo 2).
487

To analyze the single-cell heterogeneity of contribution from labeled [13]C-glucose into
acetyl-CoA, for every cell we quantified the acetyl-CoA pool labeling degree as described
in the methods. The estimated value *p* quantifies the ratio of labeled acetyl-CoA in the

cytosolic acetyl-CoA pool. This model can be applied to each single-cell profile of isotopic

intensities of a fatty acid by estimating p that leads to best approximation of the data (the

value n is known for a considered fatty acid, e.g. n=17 for palmitate). **Figure 3c** shows

examples of such modeling for two cells, a control cell (grey) and an ACLYkd oligo 2 cell

(blue). As expected, the ACLYkd cell demonstrates a lower acetyl-CoA pool labeling

degree (p equal to 0.43 vs. 0.61).

Single-cell heterogeneity of flux from glucose to acetyl-CoA

Figure 3d shows results of quantifying the acetyl-CoA pool labeling degree (p) for single

cells, plotted against the single-cell intensities of the GFP knockdown reporter. There is

a clear difference between the control cells (grey) and cells with ACLY knockdown

(ACLYkd oligo 1 in blue and ACLYkd oligo 2 in orange): as expected, an ACLY

knockdown with either oligo 1 or oligo 2 leads to lower values of p (histogram peak in

control at 0.6 compared to 0.35 in ACLYkd oligo1 and 0.4 in ACLYkd oligo 2).

Moreover, the differences between the two ACLYkd conditions (ACLYkd oligo 1 and

ACLYkd oligo 2) are clearly visible. ACLY silencing using oligo 1 resulted in a bimodal

distribution of the values of p, with a visibly higher variance (values ranging from 0.2 to

0.65). A bimodal distribution in single-cell results often indicates the presence of two

subpopulations of cells. Notably, one mode of the distribution for the acetyl-CoA pool

labeling degree in the ACLYkd oligo 1 condition displayed values for p in the range of 0.5-

0.6 that overlaps with the acetyl-CoA pool labeling degree exhibited by the control cells

(p of 0.5-0.7). This can be interpreted as the presence of a subpopulation of "weakly-

silenced" cells where the ACLY knockdown was not sufficiently induced. This hypothesis

is confirmed by the similarity of the GFP reporter data, where the GFP intensities for the

ACLYkd oligo 1 cells also show a bimodal distribution. Indeed, ACLYkd oligo 1 cells with

high values of p (above 0.5) also display GFP intensities similar to those of the WT cells

(average log GFP intensity of 2.2 for the group with higher values of p and 2.6 for the

group with lower p, compared to 1.9 for control). A possible explanation of the weak

silencing in this subpopulation is that the silencing of ACLY did not work in these cells

due to loss of the doxycycline-inducible expression cassette.

Interestingly, the second mode of the distribution of the acetyl-CoA pool labeling degree

(p) for ACLYkd oligo 1 cells has a much lower value (0.35 cf. 0.55 for the first mode). This

was clearly below the mode of the distribution of p for ACLYkd oligo 2 cells (equal to

0.42). This indicates the presence of a "strongly-silenced" subpopulation of ACLY oligo 1

cells in which silencing of ACLY was highly efficient, resulting in a strong reduction of

acetyl-CoA labelling. Thus, we can interpret these results as the heterogeneity of ACLY

knockdown using oligo 1 giving rise to two subpopulations of cells: "weakly-silenced" and

"strongly-silenced" cells.

This is contrasted by the results for ACLY silencing using oligo 2. The acetyl-CoA pool

labelling degree (p) as well as GFP reporter for these cells have values between "weakly-

silenced" and "strongly-silenced" cells from the ACLYkd oligo 1 condition. Thus, one can

interpret this data as ACLY knockdown using oligo 2 resulted in a more homogeneous

cell population of "medium-silenced" cells.

Spatial information provided by [13]C-SpaceM offers another view at the cellular

heterogeneity. **Figure 3e** shows ion images for three isotopic peaks of palmitate (M+0 in

blue, M+2 in green, M+8 in red) for all three conditions (control, ACLYkd oligo 1 and

ACLYkd oligo 2), with 3 replicates per condition. The M+2 peak was chosen as the most

discriminative for the phenotype of low acetyl-CoA labelling that is specific to the

"strongly-silenced" population among ACLYkd oligo 1 cells. The M+8 peak was chosen

as a representative isotope for the control condition where fatty acids contain high

numbers of [13]C. Thus, the difference between M+2 and M+8 as visible in **Fig. 3a** can

serve as an indicator of the relative activity of the *de novo* fatty acid synthesis and thus

be used to display heterogeneity. The data was acquired with the pixel size of 10 µm with

cells having an average diameter of 550 µm, corresponding to 12 pixels per average cell.

The ion images for ACLYkd oligo 1 showed a marked heterogeneity in the intensities of

M+2 and M+8 isotopic peaks for palmitate that visualizes the presence of two distinct

subpopulations in this condition: of "weakly-silenced" and "strongly-silenced" cells.

Interestingly, we can see that these two subpopulations are spatially-heterogeneous that

rules out a batch effect or technical artifacts and indicates a single-cell effect. In contrast,

ACLYkd oligo 2 cells show a more homogenous distribution of palmitate M+8 and an

overall lower abundance of the palmitate M+2 isotopic peak to the degree that it's hardly

visible in cells of ACLYkd oligo 2. The single-cell and spatial heterogeneity cannot be

revealed through bulk analysis, demonstrating the unique advantages of the ^{13}C-SpaceM

method.

^{13}C-SpaceM differentiates fatty acids by their relative uptake in single-cells

We next considered that ^{13}C-SpaceM data could be used to determine the relative uptake

of different fatty acids as compared to their *de novo* synthesis (**Fig. 4b**). For each fatty

acid, this was achieved by first performing natural isotope correction, normalizing all

isotopologue peaks to the sum of all peaks, approximating isotopologues intensities with

the binomial model described in the methods section, and taking the value of the "uptake"

from the model. **Figure 4a** (upper plot) shows single-cell values of the relative uptake for

the detected fatty acids: saturated fatty acids (SFAs) myristate, palmitate, stearate as well

as mono-unsaturated fatty acids (MUFAs) palmitoleate and oleate, as determined in the

control liver cancer cells. Interestingly, the levels of relative uptake at 60%-90% (and thus

de novo synthesis of 10%-40%) are close to the reported rate of 30% of *de novo*

lipogenesis in patients with NAFLD [43]. The two MUFAs, palmitoleate and oleate, showed

a substantially higher relative uptake compared to the corresponding saturated fatty acids

palmitate and stearate. This indicates that these cells mostly utilize uptake rather than *de*

novo synthesis to obtain mono-unsaturated fatty acids. The results on the average levels

were confirmed in the bulk LC-MS-based isotope tracing (**Fig. 4a**, lower plot).

We next employed the power of ^{13}C-SpaceM to investigate the relationships between the

relative uptake of different fatty acids in the same cells. This was done by plotting and

analyzing the single-cell values of relative uptake for each fatty acid vs. the others. This

analysis showed no correlation of the relative uptake of myristate with any of the other

fatty acids (**Fig. 4c, first column**). This lack of correlation is expected, as myristate was

reported to be produced by *de novo* FA synthesis only to a small degree. Instead,

myristate is mainly generated by the shortening of palmitate via peroxisomal β-oxidation

or the elongation of lauric acid [44]. In contrast, single-cell relative uptake values for

palmitate, palmitoleate, stearate and oleate showed strong positive correlations (**Fig. 4c,**

columns 2-5).

Overall, ^{13}C-SpaceM allowed to differentiate fatty acids by their sources specifically by

exogenous uptake vs *de novo* synthesis. The lack of correlation for the single-cell uptake

values for myristate likely reflects the specific mechanism of production or uptake of this

fatty acid.

Effect of the induced ACLY knockdown on fatty acid uptake

Next, we investigated the effect of ACLY knockdown on fatty acid uptake. This analysis

was restricted to ACLYkd using oligo 2 only, as the strong shift in the isotopologue

distribution observed in upon ACLYkd using oligo 1 prohibited an accurate quantification

of the fatty acid relative uptake (**see Fig. 3a, c**). Spatial analysis and visualization

demonstrated a strong single-cell heterogeneity in the relative uptake of either palmitate

or oleate in a mixed population of control and ACLYkd oligo 2 cells (**Fig. 4d and e**). More

detailed and quantified analysis by plotting the single-cell relative uptake values

separately for control and ACLYkd oligo 2 showed that ACLY-silenced cells increased the

uptake of palmitate (histogram peak at 0.55 in control compared to 0.61 in ACLYkd), with

each population displaying a unimodal symmetric distribution (**Fig. 4f**). Oleate uptake,

which was already higher compared to palmitate uptake in the control cells, was also

increased after ACLY knockdown (histogram peak at 0.85 in control compared to 0.95 in

ACLYkd). In addition, we observed an asymmetric (skewed) unimodal distribution of the

oleate single-cell relative uptake values for the ACLYkd population, with more cells

showing high relative uptake values. This indicates that some cells of the population

respond to ACLY silencing by selectively inducing the uptake of mono-unsaturated fatty

acids, such as oleate.

Taken together, we have presented ^{13}C-SpaceM, a novel method for spatial single-cell

isotope tracing, and demonstrated how it can help interrogate the activity of *de novo* fatty

acid synthesis. The key novelty of the method compared to the previously published

method SpaceM for spatial single-cell metabolomics is in using All-Ion-Fragmentation combined with fatty acid isotopologues analysis and modeling. [13]C-SpaceM can provide detailed insights into the heterogeneity of *de novo* fatty acid synthesis in cancer cells by quantifying single-cell heterogeneity of acetyl-CoA metabolism and fatty acid uptake in phenotypically distinct cell populations. Concluding, [13]C-SpaceM addresses the need to investigate the activity of well-characterized metabolic pathways on the single-cell level. [13]C-SpaceM can dissect single-cell metabolic heterogeneity as well as reveal subpopulations of cells which differ in their metabolic state yet are hidden from the bulk analyses.

Discussion

The advance of single-cell omics technologies has revolutionized insight into the genetic and phenotypic heterogeneity of cell populations at the levels of tissues, organs and whole organisms [26]. This is of particular importance in cancer research, as tumors consist of multiple subclones of cancer cells as well as a highly diverse population of stromal cells, both of which are exposed to microenvironmental conditions that can change profoundly in a spatial and temporal manner [45]. Complementing the paradigm of genetic heterogeneity of cancer cells [46] [47], there is an emerging acknowledgment of the heterogeneity of metabolic states of cancer cells. Determined by genetic and epigenetic alterations, their metabolic states are influenced by non-genetic factors such as the tissue of origin, nutrients and oxygen availability in the microenvironment as well as local interactions with the tumor stroma and immune cells [48]. Single-cell transcriptomics has provided insights into co-existence of distinct metabolic states in populations of cancer cells, such as a shift towards mitochondrial oxidative phosphorylation observed in micrometastases compared to primary tumor in breast cancer [49]. Recent technological advances in mass spectrometry, including electrospray ionization (ESI), matrix-assisted laser desorption/ionization (MALDI) or dielectric barrier discharge ionization (DBDI) in combination with advanced computational methods enabled detecting profiles of polar metabolites and lipids from single cells [30, 31, 50]. While these methods have provided unprecedented insight into the metabolic states of single cells, they fail to capture the dynamic activity of metabolic processes and the degree of flexibility within the metabolic

network. At the same time, stable-isotope tracing has emerged as the gold-standard to
reveal metabolic activity in cells and tissues in bulk [51] and in spatially-resolved manner [52],
yet was not demonstrated so far in single cells.

We developed [13]C-SpaceM, a method to combine stable-isotope tracing with spatial
single-cell metabolomics to determine *de novo* fatty acid biosynthesis in cancer cells at
single-cell resolution. [13]C-SpaceM builds upon SpaceM [31] with the major changes
including the mass spectrometry method constructed to mimic chemical saponification
and fatty acid detection as in bulk stable isotope tracing as well as the data analysis. In
particular, we demonstrate how using the binomial model applied to the isotopologue
distribution helps estimate such labeling parameters as uptake and pool labeling degree
for individual cells.

[13]C-SpaceM as employed in this study uses all-ion fragmentation to release fatty acids
from cellular lipids at m/z range of 600-1000 and ionized in the negative mode, thereby
focusing on glycerophospholipids and sphingolipids while mostly excluding
lysophospholipids, triglycerides and cardiolipins [37]. This selectivity in mass range can be
used to selectively monitor labelling in fatty acids present in specific lipid classes, in this
case in the lipids most abundant in cell membranes. Indeed, when comparing [13]C-
SpaceM data to bulk stable isotope tracing in fatty acids after saponification of total
cellular lipids from the same cells, we noticed a larger proportion of palmitate derived from
de novo synthesis in the bulk analysis. As the cells used in this study have already
reached isotopic steady state after 72 hours in the presence of [13]C-glucose, this result
suggests that the synthesis of specific lipids utilizes distinct pools of palmitate that differ
in their contribution from *de novo* fatty acid synthesis and uptake [6]. Local fatty acid
channeling into phospholipid synthesis has been observed during the rapid formation of
the autophagic membrane in yeast [53], indicating that lipid synthesis can take place in a
context-specific manner.

We validated the ability of [13]C-SpaceM to detect changes in glucose-dependent fatty acid
synthesis, and its single-cell resolution, by co-plating cells that had previously been
cultured under either normoxia or hypoxia. This system was chosen as hypoxia leads to
a global reprogramming of cellular lipid metabolism [10]. Spatial single-cell isotope tracing
by [13]C-SpaceM was able to clearly separate the two cell populations based on the strong

increase in the M+0 fraction of palmitate observed in hypoxic cells. This is expected, as
hypoxia reduces the conversion of glucose-derived pyruvate into mitochondrial acetyl-
CoA via induction of pyruvate dehydrogenase kinase 1 (PDK1) [54], thereby lowering the
flux of glucose carbons into the TCA cycle and subsequently into the lipogenic
(cytoplasmic) acetyl-CoA pool. At the same time, hypoxia induces the reductive
carboxylation of glutamine thus generating cytoplasmic citrate to support lipid synthesis
[19, 20] and the direct synthesis of acetyl-CoA from acetate [21, 22]. In addition, hypoxia leads
to the uptake of exogenous fatty acids [18, 39], further increasing the proportion of the M+0
fraction.

We also assessed the application of ^{13}C-SpaceM to monitor the effect of perturbations of
the metabolic network in cancer cells. Both bulk and single-cell analysis revealed a
marked shift in palmitate isotopologue distribution following silencing of ACLY, correctly
reflecting a reduced contribution of ^{13}C-glucose to the lipogenic acetyl-CoA pool. Using a
binomial model to quantify fractional labelling from the isotopologue distribution [21], we
estimated the labelling degree of acetyl-CoA in control and ACLY silenced cells at single-
cell level. In agreement with previous studies using bulk analysis of palmitate
isotopologue distribution in ACLY wildtype vs knockout mouse embryo fibroblasts [24],
silencing ACLY caused a reduction in the fractional labelling of acetyl-CoA from glucose
but only a minor increase in the M+0 fraction. This suggests that other substrates, such
as acetate, are used to synthesize acetyl-CoA under these conditions [24]. Our analysis
also revealed a substantial degree of heterogeneity in fractional labelling of lipogenic
acetyl-CoA within the cell population, particularly after silencing ACLY. This heterogeneity
suggested differences in the silencing efficiency of the different oligonucleotides used to
target ACLY, which were completely obscured in the bulk analysis. Acetyl-CoA plays a
central role in the metabolism of cancer cells, as it is not only used as a substrate for fatty
acid and cholesterol synthesis but is also essential for post-translational modification of
proteins, most importantly histones. Methods to quantify acetyl-CoA synthesis from
different substrates at single-cell level can reveal the link between metabolism and
genetic or environmental factors.

We also used ^{13}C-SpaceM to determine the relative proportion of *de novo* synthesis and
uptake of five non-essential fatty acids at single-cell level. While fatty acids can be taken

up by cells through passive diffusion, it is generally concluded that uptake functions

through receptor-mediated processes [55]. Moreover, fatty acid uptake via the scavenging

receptor CD36 has been shown to facilitate metastasis formation by promoting a stem

cell phenotype in cancer [56]. Our single-cell analysis revealed substantial heterogeneity in

relative uptake for the different fatty acid within the population of cancer cells. Moreover,

correlation analysis indicated increased uptake of palmitoleate and oleate, suggesting a

higher demand for mono-unsaturated fatty acids. ACLY silencing resulted in a general

increase in fatty acid uptake, in particular palmitate and oleate, indicating that limiting the

conversion of citrate to acetyl-CoA lowers the ability of cancer cells to fulfil their fatty acid

demand via de novo synthesis [24].

Interestingly, a recent single-cell lipidomics study investigated the effect of chemical

inhibition of ACLY on intracellular lipids in pancreatic cancer cells [30]. Their findings

revealed a reduction of specific PC species containing saturated and mono-unsaturated

fatty acids upon ACLY inhibition. This is consistent with our observations, as a reduction

in *de novo* fatty acid synthesis would selectively reduce the availability of saturated and

mono-unsaturated fatty acids for lipid synthesis und could therefore explain the reduction

in these lipid species.

Together, the results presented here present ¹³C-SpaceM as a method for spatial single-

cell isotope tracing able to monitor *de novo* fatty acid synthesis, production of lipogenic

acetyl-CoA and fatty acid uptake of heterogenous cell populations exposed to genetic

and environmental perturbations. Through additional modifications of this approach, for

example by applying different metabolic tracers or by restricting collision-induced

dissociation to individual lipid classes, ¹³C-SpaceM can provide broader or more specific

information about activity of the *de novo* lipid biogenesis at the single-cell resolution.

Furthermore, by detecting other metabolite classes with the underlying MALDI-imaging

mass spectrometry as already demonstrated [52], ¹³C-SpaceM can be extended for stable

isotope tracing of other metabolic pathways thus opening avenues towards revealing and

understanding single-cell heterogeneity of metabolic activity.

References

1. Harayama, T. & Riezman, H. Understanding the diversity of membrane lipid composition. *Nat Rev Mol Cell Biol* **19**, 281-296 (2018).
2. Farese, R.V., Jr. & Walther, T.C. Lipid droplets finally get a little R-E-S-P-E-C-T. *Cell* **139**, 855-860 (2009).
3. Wang, D. & Dubois, R.N. Eicosanoids and cancer. *Nat Rev Cancer* **10**, 181-193 (2010).
4. Park, J.B. *et al.* Phospholipase signalling networks in cancer. *Nat Rev Cancer* **12**, 782-792 (2012).
5. Vanhaesebroeck, B., Stephens, L. & Hawkins, P. PI3K signalling: the path to discovery and understanding. *Nat Rev Mol Cell Biol* **13**, 195-203 (2012).
6. Currie, E., Schulze, A., Zechner, R., Walther, T.C. & Farese, R.V., Jr. Cellular fatty acid metabolism and cancer. *Cell Metab* **18**, 153-161 (2013).
7. Rohrig, F. & Schulze, A. The multifaceted roles of fatty acid synthesis in cancer. *Nat Rev Cancer* **16**, 732-749 (2016).
8. Snaebjornsson, M.T., Janaki-Raman, S. & Schulze, A. Greasing the Wheels of the Cancer Machine: The Role of Lipid Metabolism in Cancer. *Cell Metab* **31**, 62-76 (2020).
9. Koundouros, N. & Poulogiannis, G. Reprogramming of fatty acid metabolism in cancer. *Br J Cancer* **122**, 4-22 (2020).
10. Munir, R., Lisec, J., Swinnen, J.V. & Zaidi, N. Lipid metabolism in cancer cells under metabolic stress. *Br J Cancer* **120**, 1090-1098 (2019).
11. Peck, B. & Schulze, A. Lipid desaturation - the next step in targeting lipogenesis in cancer? *FEBS J* **283**, 2767-2778 (2016).
12. Griffiths, B. *et al.* Sterol regulatory element binding protein-dependent regulation of lipid synthesis supports cell survival and tumor growth. *Cancer & metabolism* **1**, 3 (2013).
13. Young, R.M. *et al.* Dysregulated mTORC1 renders cells critically dependent on desaturated lipids for survival under tumor-like stress. *Genes & development* **27**, 1115-1131 (2013).
14. Williams, K.J. *et al.* An essential requirement for the SCAP/SREBP signaling axis to protect cancer cells from lipotoxicity. *Cancer research* (2013).
15. Rudalska, R. *et al.* LXRα activation and Raf inhibition trigger lethal lipotoxicity in liver cancer. *Nature Cancer* **2**, 201-217 (2021).
16. Peck, B. *et al.* Inhibition of fatty acid desaturation is detrimental to cancer cell survival in metabolically compromised environments. *Cancer & metabolism* **4**, 6 (2016).
17. Magtanong, L. *et al.* Exogenous Monounsaturated Fatty Acids Promote a Ferroptosis-Resistant Cell State. *Cell Chem Biol* **26**, 420-432 e429 (2019).
18. Kamphorst, J.J. *et al.* Hypoxic and Ras-transformed cells support growth by scavenging unsaturated fatty acids from lysophospholipids. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 8882-8887 (2013).
19. Metallo, C.M. *et al.* Reductive glutamine metabolism by IDH1 mediates lipogenesis under hypoxia. *Nature* **481**, 380-384 (2011).
20. Wise, D.R. *et al.* Hypoxia promotes isocitrate dehydrogenase-dependent carboxylation of alpha-ketoglutarate to citrate to support cell growth and viability. *Proc Natl Acad Sci U S A* **108**, 19611-19616 (2011).
21. Kamphorst, J.J., Chung, M.K., Fan, J. & Rabinowitz, J.D. Quantitative analysis of acetyl-CoA production in hypoxic cancer cells reveals substantial contribution from acetate. *Cancer & metabolism* **2**, 23 (2014).
22. Schug, Z.T. *et al.* Acetyl-CoA synthetase 2 promotes acetate utilization and maintains cancer cell growth under metabolic stress. *Cancer cell* **27**, 57-71 (2015).
23. Sivanand, S., Viney, I. & Wellen, K.E. Spatiotemporal Control of Acetyl-CoA Metabolism in Chromatin Regulation. *Trends Biochem Sci* **43**, 61-74 (2018).

24. Zhao, S. *et al.* ATP-Citrate Lyase Controls a Glucose-to-Acetate Metabolic Switch. *Cell*
reports **17**, 1037-1052 (2016).
25. Bulusu, V. *et al.* Acetate Recapturing by Nuclear Acetyl-CoA Synthetase 2 Prevents Loss
of Histone Acetylation during Oxygen and Serum Limitation. *Cell reports* **18**, 647-658
(2017).
26. Heath, J.R., Ribas, A. & Mischel, P.S. Single-cell analysis tools for drug discovery and
development. *Nat Rev Drug Discov* **15**, 204-216 (2016).
27. Duncan, K.D., Fyrestam, J. & Lanekoff, I. Advances in mass spectrometry based single-
cell metabolomics. *Analyst* **144**, 782-793 (2019).
28. Ellis, J.M. & Wolfgang, M.J. A genetically encoded metabolite sensor for malonyl-CoA.
Chem Biol **19**, 1333-1339 (2012).
29. Guo, Z. *et al.* Single-Cell Profiling of Fatty Acid Uptake Using Surface-Immobilized
Dendrimers. *J Am Chem Soc* **143**, 11191-11198 (2021).
30. Liu, Q. *et al.* High-Throughput Single-Cell Mass Spectrometry Reveals Abnormal Lipid
Metabolism in Pancreatic Ductal Adenocarcinoma. *Angew Chem Int Ed Engl* **60**, 24534-
24542 (2021).
31. Rappez, L. *et al.* SpaceM reveals metabolic states of single cells. *Nat Methods* **18**, 799-
805 (2021).
32. Tumanov, S., Bulusu, V. & Kamphorst, J.J. Analysis of Fatty Acid Metabolism Using Stable
Isotope Tracers and Mass Spectrometry. *Methods Enzymol* **561**, 197-217 (2015).
33. Heinrich, P. *et al.* Correcting for natural isotope abundance and tracer impurity in MS-,
MS/MS- and high-resolution-multiple-tracer-data from stable isotope labeling experiments
with IsoCorrectoR. *Sci Rep* **8**, 17910 (2018).
34. Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for
cellular segmentation. *Nat Methods* **18**, 100-106 (2021).
35. Millard, P. *et al.* IsoCor: isotope correction for high-resolution MS labeling experiments.
Bioinformatics **35**, 4484-4487 (2019).
36. Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression
data analysis. *Genome Biol* **19**, 15 (2018).
37. Tumanov, S. & Kamphorst, J.J. Recent advances in expanding the coverage of the
lipidome. *Curr Opin Biotechnol* **43**, 127-133 (2017).
38. Mullen, A.R. *et al.* Reductive carboxylation supports growth in tumour cells with defective
mitochondria. *Nature* **481**, 385-388 (2012).
39. Bensaad, K. *et al.* Fatty acid uptake and lipid storage induced by HIF-1alpha contribute to
cell growth and survival after hypoxia-reoxygenation. *Cell reports* **9**, 349-365 (2014).
40. Scaglia, N., Tyekucheva, S., Zadra, G., Photopoulos, C. & Loda, M. De novo fatty acid
synthesis at the mitotic exit is required to complete cellular division. *Cell Cycle* **13**, 859-
868 (2014).
41. Bligh, E.G. & Dyer, W.J. A rapid method of total lipid extraction and purification. *Can J*
Biochem Physiol **37**, 911-917 (1959).
42. Ackerman, D. *et al.* Triglycerides Promote Lipid Homeostasis during Hypoxic Stress by
Balancing Fatty Acid Saturation. *Cell reports* **24**, 2596-2605 e2595 (2018).
43. Donnelly, K.L. *et al.* Sources of fatty acids stored in liver and secreted via lipoproteins in
patients with nonalcoholic fatty liver disease. *J Clin Invest* **115**, 1343-1351 (2005).
44. Rioux, V., Catheline, D. & Legrand, P. In rat hepatocytes, myristic acid occurs through
lipogenesis, palmitic acid shortening and lauric acid elongation. *Animal* **1**, 820-826 (2007).
45. Li, Z., Seehawer, M. & Polyak, K. Untangling the web of intratumour heterogeneity. *Nat*
Cell Biol **24**, 1192-1201 (2022).
46. Hausser, J. & Alon, U. Tumour heterogeneity and the evolutionary trade-offs of cancer.
Nat Rev Cancer **20**, 247-257 (2020).

833 47. Kim, C. *et al.* Chemoresistance Evolution in Triple-Negative Breast Cancer Delineated by
834   Single-Cell Sequencing. *Cell* **173**, 879-893 e813 (2018).
835 48. Kim, J. & DeBerardinis, R.J. Mechanisms and Implications of Metabolic Heterogeneity in
836   Cancer. *Cell Metab* **30**, 434-446 (2019).
837 49. Davis, R.T. *et al.* Transcriptional diversity and bioenergetic shift in human breast cancer
838   metastasis revealed by single-cell RNA sequencing. *Nat Cell Biol* **22**, 310-320 (2020).
839 50. Fujii, T. *et al.* Direct metabolomics for plant cells by live single-cell mass spectrometry. *Nat*
840   *Protoc* **10**, 1445-1456 (2015).
841 51. Jang, C., Chen, L. & Rabinowitz, J.D. Metabolomics and Isotope Tracing. *Cell* **173**, 822-
842   837 (2018).
843 52. Wang, L. *et al.* Spatially resolved isotope tracing reveals tissue metabolic activity. *Nat*
844   *Methods* **19**, 223-230 (2022).
845 53. Schutter, M., Giavalisco, P., Brodesser, S. & Graef, M. Local Fatty Acid Channeling into
846   Phospholipid Synthesis Drives Phagophore Expansion during Autophagy. *Cell* **180**, 135-
847   149 e114 (2020).
848 54. Kim, J.W., Tchernyshyov, I., Semenza, G.L. & Dang, C.V. HIF-1-mediated expression of
849   pyruvate dehydrogenase kinase: A metabolic switch required for cellular adaptation to
850   hypoxia. *Cell Metab* **3**, 177-185 (2006).
851 55. Samovski, D., Jacome-Sosa, M. & Abumrad, N.A. Fatty Acid Transport and Signaling:
852   Mechanisms and Physiological Implications. *Annu Rev Physiol* (2022).
853 56. Pascual, G. *et al.* Targeting metastasis-initiating cells through the fatty acid receptor
854   CD36. *Nature* **541**, 41-45 (2017).
855

Figures

Figure 1. ^{13}C-SpaceM workflow as applied to interrogate *de novo* fatty acid synthesis

a) Generation of cytoplasmic acetyl-CoA and subsequent fatty acid and lipid synthesis from stable-isotope labelled ^{13}C$_6$-glucose. Glucose-derived pyruvate is metabolized in the mitochondria and exported to the cytoplasm in the form of citrate. ATP-citrate lyase (ACLY) converts citrate to acetyl-CoA, thereby contributing to the cytoplasmic (lipogenic) acetyl-CoA pool. *De novo* fatty acid synthesis by fatty acid synthase (FASN) results in the formation of palmitate (C16:0), which is either further modified or directly incorporated into cellular lipids. Exogenously taken up fatty acids are also incorporated into cellular lipids but are not labelled.

b) Using all-ion fragmentation imaging mass spectrometry for ^{13}C-SpaceM. Cells are grown in medium supplemented with uniformly labelled ^{13}C$_6$-glucose for 72 hours. The combination of wide-range isolation of parent lipid ions followed by HCD fragmentation and selective isolation of fatty acid fragments allows high-sensitivity measurements of fatty acids incorporated into lipids,

conceptually similar to bulk extractions followed by saponification but with the benefit of retained spatial information and the exclusion of free fatty acids.

c) Integration of microscopy and imaging mass spectrometry in ^{13}C-SpaceM to obtain single-cell profiles. Pre-MALDI microscopy and post-MALDI microscopy images containing information about the cell outlines and areas ablated by MALDI-imaging are registered. Ion intensities from MALDI-imaging are assigned to single cells through a normalization procedure.

Figure 2. Validating ^{13}C-SpaceM by interrogating *de novo* fatty acid synthesis in spatially-heterogeneous normoxia-hypoxia model

a) Illustration of microscopy and imaging mass spectrometry data from the model of co-plated primary murine liver cancer cells originally cultured under normoxia (GFP-negative) and hypoxia (GFP-positive). The GFP signal was used for discerning the culturing conditions (white and green cell outlines shown normoxic and hypoxic cells, respectively). The normalized intensities of the M+0 isotope of palmitate (representing the fraction of unlabelled palmitate) are shown for MALDI-imaging pixels and as assigned to the single cells.

b) Mass spectra for individual pixels mapped to cells from a normoxic cell (white outline) and a hypoxic cell (green outline). The peaks corresponding to palmitate isotopologues are marked by grey or green dots.

c and d) Discerning cells cultured under normoxia vs. hypoxia using ^{13}C-SpaceM for the cells mono-plated for each culturing condition (C) and for spatially-heterogeneous co-plated cells from both conditions (D). Scatterplot shows the values of the GFP-reporter (ground truth for telling the

condition) and the normalized intensity of the M+0 isotope of palmitate representing the fraction of unlabelled palmitate for single cells.

e) Comparison of single-cell vs. bulk intensities for the M+0 isotope of palmitate. Single-cell intensities are from the spatially-heterogeneous co-plated model. Bulk intensities are from cells of each mono-cultured condition subjected to total fatty acid analysis by saponification followed by LC-MS/MS. For single-cell data, black lines show average values. For bulk data, data are displayed as mean ± standard deviation across 3 replicates.

Figure 3. Single-cell quantitative analysis of lipogenic acetyl-CoA production and heterogeneity

a) Comparison of bulk and single-cell analysis of isotopologue distribution for palmitate after 72 hours of ACLY gene silencing and culture in the presence of $^{13}C_6$-glucose. Bulk data are displayed as mean ± standard deviation across 3 replicates. For single-cell data, black lines show average values.

b) Immunoblot of protein lysates showing efficiency of ACLY knockdown using two shRNA sequences (ACLYkd oligo1 and ACLYkd oligo 2) compared to non-targeting control (shNT). Tubulin B (TUBB) is shown as loading control.

c) Normalized single-cell isotopologues distribution for two individual cells, one from the control and the other from ACLYkd oligo 1 (shown as bar plots). Lines show fit of the fatty acid labelling binomial model: horizontal lines for M+0 showing an estimated uptake, and connected lines for

M+2i showing (1 - uptake) * binomial(i). Legend shows parameters of the binomial model fit.

d) Single-cell analysis of the estimated acetyl-CoA pool labelling degree (*p*) as calculated using the fatty acid labelling binominal model for the three conditions: control (grey), ACLYkd oligo 1 (blue), and ACLYkd oligo 2 (blue). Green dashed line shows 95% quantile of the GFP intensity distribution in the control condition, which was used to classify cells as GFP-positive vs. GFP-negative.

e) Spatial metabolic imaging of *de novo* fatty synthesis for the control, ACLYkd oligo 1, and ACLYkd oligo 2 conditions. Abundance of different palmitate isotopic peaks are displayed in different channels: M+0 (blue), M+2 (green) and M+8 (red). Each channel is normalized to the total ion count (TIC).

Figure 4. ¹³C-SpaceM differentiates fatty acids by their relative uptake in single-cells

a) Correlation of uptake of different fatty acids. Dashed line is the diagonal. For each pair of fatty acids, the linear fit is shown as a grey line, with the fit parameters shown in the legend. Schematic of biosynthesis of different fatty acids?

b) Diagram showing mechanisms involved in fatty acid synthesis, desaturation and uptake.

c) Fatty acids uptake in the control cells as determined for myristate, palmitate, palmitoleate, stearate and oleate; saturated fatty acids in green and unsaturated fatty acids in blue. Top: Single-cell data with each point representing a cell, black horizontal lines showing median values. Bottom: Data from the bulk stable isotope tracing.

d and e) Spatial metabolic imaging of the fatty acid uptake in co-plated cells from two conditions (control and ACLYkd oligo 2). Each square corresponds to one MALDI pixel. Pixels with the total intensity of the fatty acid isotopes below the set limit of detection (1000) are shown in black. Cell outlines are green for GFP-positive cells and white for GFP-negative cells. Pixels are colored

according to the normalized M+0 describing the uptake contribution for a given fatty acid. D: Normalized M+0 of palmitate. E: Normalized M+0 of oleate.

f) Single-cell analysis of how fatty acid uptake changes upon ACLY knockdown. Dashed line is the diagonal. For each condition the linear fit is shown in the corresponding color, with the fit parameters shown in the legend.