Title: Targeting host deoxycytidine kinase attenuates *Staphylococcus aureus* virulence

Authors: Volker Winstel[1,2]*, Evan R. Abt[3], Caius G. Radu[3]

Affiliations:

[1] Research Group Pathogenesis of Bacterial Infections; TWINCORE, Centre for Experimental and Clinical
Infection Research, a joint venture between the Hannover Medical School and the Helmholtz Centre for
Infection Research, Hannover, Germany

[2] Institute of Medical Microbiology and Hospital Epidemiology, Hannover Medical School, Hannover, Germany

[3] Department of Molecular and Medical Pharmacology, David Geffen School of Medicine, UCLA, Los Angeles,
United States of America

* Address **correspondence** to Volker Winstel: winstel.volker@mh-hannover.de

Institute of Medical Microbiology and Hospital Epidemiology, Hannover Medical School
Carl-Neuberg-Str. 1, 30625 Hannover, Germany
Phone: +49 511 22002 7220 / Fax: +49 511 22002 7186

Abstract

Host-directed therapy (HDT) is an emerging approach to overcome antimicrobial resistance in pathogenic microorganisms. Specifically, HDT targets host-encoded factors required for pathogen replication and survival without interfering with microbial growth or metabolism, thereby eliminating the risk of resistance development. By applying HDT and a drug repurposing approach, we demonstrate that (*R*)-DI-87, a clinical-stage anti-cancer drug and potent inhibitor of mammalian deoxycytidine kinase (dCK), attenuates the virulence of antibiotic-resistant *Staphylococcus aureus* in a mouse model of bloodstream infection. Mechanistically, (*R*)-DI-87 shields phagocytes from staphylococcal death-effector deoxyribonucleosides that target dCK and the mammalian purine salvage pathway-apoptosis axis. In this manner, (*R*)-DI-87-mediated protection of immune cells amplifies macrophage infiltration into deep-seated abscesses, a phenomenon coupled with enhanced pathogen control, ameliorated immunopathology, and reduced disease severity. Thus, pharmaceutical blockade of dCK represents an advanced anti-infective intervention strategy against which staphylococci cannot develop resistance and may help to fight fatal infectious diseases in hospitalized patients.

Introduction

Hospital- and community-acquired infections caused by antibiotic-resistant bacterial pathogens represent a global public health threat (Wenzel, 2007; GBD 2019 Antimicrobial Resistance Collaborators, 2022a; b). Specifically, infections caused by multidrug-resistant microbes are often associated with increased morbidity and mortality, as compared to infections caused by antibiotic-sensitive strains (Klevens et al., 2008; Yaw et al., 2014; GBD 2019 Antimicrobial Resistance Collaborators, 2022a; b). In addition, the pathology of various bacterial infections includes multifactorial manipulation of host multicellular assemblies that enable pathogen replication, dissemination of disease, and adaptation to new hosts (Foster, 2005; Finlay and McFadden, 2006; Okumura and Nizet, 2014; Thammavongsa et al., 2015). Thus, prevention and therapy of such infections represent a major challenge in clinical practice and frequently require administering last-resort antibiotics such as vancomycin, linezolid, or colistin (Chambers and Deleo, 2009; McKenna, 2013; Tong et al., 2015). However, many clinically relevant pathogens have already developed resistance-mechanisms against many of these drugs thereby rendering antibiotic therapy of appropriate infections often ineffective (Meka and Gold, 2004; Chambers and Deleo, 2009; GBD 2019 Antimicrobial Resistance Collaborators, 2022a). Moreover, the rapid exchange of antibiotic resistance genes via horizontal gene transfer events, along with the large-scale usage of antimicrobials in hospitals and in the animal industry, continuously triggers the emergence of new drug-resistant and globally spreading bacterial clones, highlighting the desperate need for novel treatment options to fight infectious diseases in human or animal hosts (Chambers and Deleo, 2009; Nubel, 2016; Jian et al., 2021).

Antibiotic-resistant bacteria are frequently associated with local and life-threatening infections in both, hospital environments and in the general community. Of particular concern are Mycobacteria, certain Gram-negative as well as Gram-positive bacteria including *Staphylococcus aureus*, a notorious pathogen that colonizes approximately 30% of the human population (Lowy, 1998; Kuehnert et al., 2006; Lee et al., 2018). *S. aureus* causes skin and soft tissue infections and is a widespread cause of endocarditis, septic arthritis, osteomyelitis, bacteremia, and sepsis (Lowy, 1998; Fridkin et al., 2005; Klevens et al., 2007). Moreover, this microbe triggers toxic shock syndrome, food poisoning, ventilator-associated pneumonia, and associated infections of the respiratory tract (Lowy, 1998; Lee et al., 2018; Torres et al., 2021). Nonetheless, a hallmark of staphylococcal disease is the formation of suppurative abscesses, where *S. aureus* provokes inflammatory responses that attract neutrophils, macrophages, and other immune cells to the site of the infection (Lowy, 1998; Cheng et al., 2011; Spaan et al., 2013). Abscess formation is further accompanied by liquefaction necrosis, the release of purulent exudates into circulating body fluids, and several defensive host responses designed to limit staphylococcal replication and spread (Cheng et al., 2011). For example, staphylococcal invasion of host tissues and abscess formation includes deposition of structural fibrin clusters, which shield

healthy tissues from disseminating staphylococci (Cheng et al., 2011). Moreover, neutrophils combat *S. aureus*
by powerful mechanisms which involve phagocytosis, generation of reactive oxygen species, and the
formation of neutrophil extracellular traps (NETs) (Spaan et al., 2013). NETs represent an extracellular matrix
composed of nuclear and mitochondrial DNA spiked with antimicrobial peptides, cell-specific proteases, and
granular proteins that together ensnare and kill microbial invaders (Brinkmann et al., 2004). However, *S.*
aureus rapidly evades NET-mediated killing due to the secretion of a potent nuclease (Nuc) capable of
disrupting these structures (Berends et al., 2010; von Kockritz-Blickwede and Winstel, 2022). Intriguingly, Nuc-
mediated degradation of NETs leads to the release of deoxyribonucleoside monophosphates that can be
converted by the staphylococcal cell surface protein AdsA (adenosine synthase A) into death-effector
deoxyribonucleosides (Thammavongsa et al., 2013; Tantawy et al., 2022). Notably, AdsA-derived death-
effector deoxyribonucleosides such as deoxyadenosine (dAdo) and deoxyguanosine (dGuo) display toxigenic
properties and contribute to immune cell death during acute and persistent *S. aureus* infections
(Thammavongsa et al., 2013; Winstel et al., 2019; Tantawy et al., 2022). More precisely, dAdo and dGuo-
triggered phagocyte cell death involves the uptake of dAdo and dGuo via human equilibrative nucleoside
transporter 1 (hENT1), intracellular phosphorylation of dAdo and dGuo by adenosine kinase (ADK) and
deoxycytidine kinase (dCK), and subsequent imbalanced expansion of intracellular dATP and dGTP pools which
trigger caspase-3-dependent immune cell death (Winstel et al., 2018; Tantawy et al., 2022). Of note, genetic
disruption of this metabolic cascade rendered macrophages resistant to AdsA-derived death-effector
deoxyribonucleosides, leading to the accumulation of phagocytes within the deeper cavity of abscesses and
accelerated clearance of staphylococci *in vivo* (Winstel et al., 2018; Winstel et al., 2019; Tantawy et al., 2022).
Thus, elements of the mammalian purine salvage pathway represent attractive targets for the design of novel
host-directed therapeutics that aim at boosting macrophage survival during acute and persistent *S. aureus*
infections. In particular, pharmacological inhibition of dCK by small molecule inhibitors may terminate the
intracellular conversion of *S. aureus*- and AdsA-derived death-effector deoxyribonucleosides into
deoxyribonucleoside monophosphates and subsequent accumulation of apoptosis-triggering
deoxyribonucleoside triphosphates in host phagocytes (Figure 1A). If so, *S. aureus*- and dAdo/dGuo-mediated
macrophage cell death may efficiently be prevented thereby aiding in the clearance of staphylococci during
abscess formation and persistent infections.
Here we provide proof of this concept by establishing a novel host-directed therapeutic strategy for
therapy of staphylococcal infectious diseases. Specifically, we show that the dCK-specific small molecule
inhibitor   (*R*)-DI-87   (((*R*)-2-((1-(2-(4-methoxy-3-(2-morpholinoethoxy)phenyl)-5-methylthiazol-4-
yl)ethyl)thio)pyrimidine-4,6-diamine), an orally active, well-tolerated clinical-stage anti-cancer drug (Figure
1B) (Poddar et al., 2020), protects host phagocytes from staphylococcal death-effector deoxyribonucleoside-

mediated cytotoxicity and caspase-3-dependent cell death. Our results further suggest that administration of

(*R*)-DI-87 boosts macrophage survival during abscess formation and thus improves clinical outcomes in *S.*

aureus-infected laboratory animals.

Results

(*R*)-DI-87-mediated blockade of mammalian dCK protects host immune cells from staphylococcal death-

effector deoxyribonucleosides. To analyze whether pharmacological inhibition of mammalian dCK may

represent a suitable strategy to protect host phagocytes from staphylococcal death-effector

deoxyribonucleosides, we initially took advantage of various tissue culture model systems and pre-incubated

human U937 monocytes (U937) or U937-derived macrophages (U937 MФ) with (*R*)-DI-87. Controls received

vehicle only. Following pre-incubation, U937 or U937 MФ were exposed to dAdo or dGuo and analyzed for

viability rates 48 hours post-intoxication. Of note, (*R*)-DI-87-mediated inhibition of mammalian dCK in U937

or U937 MФ efficiently prevented dAdo- or dGuo-induced cell death in a dose-dependent manner (Figure 2A-

D; supplementary Figure 1A-D). Moreover, these effects resembled the phenotype of dCK-deficient U937 or

U937 MФ, which were found to be refractory to dAdo- or dGuo-mediated cytotoxicity (Figure 2A-D) (Winstel

et al., 2018; Tantawy et al., 2022). In addition, we observed that pre-treatment of primary human CD14[+]

monocytes or human monocyte-derived macrophages (HMDMs) with (*R*)-DI-87 blocked dAdo- or dGuo-

triggered cytotoxicity suggesting that dCK inhibition may suppress the toxigenic properties of staphylococcal

death-effector deoxyribonucleosides (Figure 2E-H). To test this conjecture, we expressed a soluble and

affinity-tagged recombinant form of *S. aureus* AdsA (hereafter termed rAdsA) in *Escherichia coli*, which was

purified, released of its affinity tag and subsequently used for cytotoxicity assays. Particularly, we incubated

rAdsA with either dAMP or dGMP according to a published protocol (Tantawy et al., 2022) and added the

resulting, dAdo- or dGuo-containing and filter-sterilized reaction products to U937 or U937 MФ that received

(*R*)-DI-87 or were left untreated prior to intoxication. While enzymatic reactions that contained rAdsA and

purine deoxyribonucleoside monophosphates triggered cell death of human monocytes or macrophages in

this approach, (*R*)-DI-87-exposed cells were fully protected against the cytotoxic effect of dAdo and dGuo

further indicating that inhibition of mammalian dCK helps to neutralize *S. aureus* AdsA-derived death-effector

deoxyribonucleosides (Figure 2I-J). Collectively, these initial data demonstrate that (*R*)-DI-87 is a suitable small

molecule dCK inhibitor capable of preventing host immune cell death induced by toxigenic products of the

staphylococcal Nuc/AdsA pathway.

(*R*)-DI-87 prevents death-effector deoxyribonucleoside-induced activation of apoptosis. Earlier work

demonstrated that staphylococcal dAdo and dGuo target the mammalian purine salvage pathway to trigger

an exaggerated biogenesis of deoxyribonucleoside triphosphates, thereby igniting caspase-3-dependent host

immune cell death (Winstel et al., 2018; Winstel et al., 2019; Tantawy et al., 2022). Since the accumulation of

deoxyribonucleoside triphosphates and associated apoptotic signaling via cleavage of caspase-3 exclusively

occurs in dCK-proficient monocytes or macrophages upon intoxication with death-effector

deoxyribonucleosides such as dAdo (Winstel et al., 2018; Tantawy et al., 2022), it seemed plausible to us at

this stage that administration of (*R*)-DI-87 and inhibition of dCK may suppress dAdo- or dGuo-mediated

activation of the programmed cell death machinery. To test this hypothesis, U937 or U937 MΦ were pre-

incubated with (*R*)-DI-87, exposed to dAdo or dGuo, and used to generate cell extracts for evaluation of

caspase-3 activity by measuring the hydrolysis of the caspase-3-specific peptide substrate Ac-DEVD-pNA. As

expected, (*R*)-DI-87-treatment of U937 or U937 MΦ significantly decreased dAdo- or dGuo-mediated

activation of caspase-3 activity (Figure 3A-D). Since caspase-3 represents the key modulator of the apoptosis

signaling pathway, these data suggest that (*R*)-DI-87-mediated dCK inhibition selectively prevented dAdo- or

dGuo-mediated induction of apoptotic cell death in host immune cells (Figure 3A-D). To verify these results

further, cell extracts were also probed with a specific antibody capable of detecting the active (cleaved) form

of human caspase-3. In agreement with the enzymatic activity assay, (*R*)-DI-87-treatment of U937 or U937 MΦ

prevented caspase-3 activation (Figure 3E-F). Moreover, we exposed U937 phagocytes to (*R*)-DI-87 and

assessed dAdo- or dGuo-mediated activation of apoptotic signaling via immunofluorescence microscopy.

Microscopy-based analysis of (*R*)-DI-87- and death-effector deoxyribonucleoside-exposed macrophages

confirmed that (*R*)-DI-87 inhibited activation of programmed cell death and apoptosis as positive signals for

annexin-V/PI were strongly decreased in samples that have been exposed to the dCK inhibitor (Figure 3G-H).

Of note, similar findings were also obtained with primary HMDMs, suggesting that (*R*)-DI-87 is a suitable agent

to block death-effector deoxyribonucleoside-mediated induction of apoptosis in host phagocytes

(supplementary Figure 2). Together, these data indicate that administration of (*R*)-DI-87 and associated

inhibition of mammalian dCK prevents apoptotic cell death in phagocytes caused by *S. aureus* AdsA-derived

death-effector deoxyribonucleosides.

Pharmacological inhibition of host dCK attenuates *S. aureus* virulence in a mouse model of bloodstream

infection. To evaluate the therapeutic value and safety of (*R*)-DI-87 *in vivo*, cohorts of female C57BL/6 mice

received (*R*)-DI-87 (75 mg/kg) or vehicle (40% Captisol) via oral gavage in 12-hour intervals over a time period

of 16 days. On day 16, peripheral blood was collected from all cohorts of mice and subjected to a FACS-based

immuno-phenotyping approach (Figure 4A). Continuous treatment of animals with (*R*)-DI-87 did not alter the

immune cell composition of peripheral blood (Figure 4A). Further, animals treated with (*R*)-DI-87 did neither

lose weight nor displayed any other obvious phenotype during this procedure suggesting that (*R*)-DI-87

represents a safe and highly selective therapeutic agent (Figure 4B). Next, we sought to investigate whether

(*R*)-DI-87-mediated protection of mammalian phagocytes may represent a valuable strategy to cure

staphylococcal diseases and tested the therapeutic efficacy of the compound in a mouse model of *S. aureus*

bloodstream infection. In this model, C57BL/6 mice were treated with (*R*)-DI-87 (75 mg/kg) or vehicle (40%

Captisol) via oral gavage every 12 hours and challenged with a single dose of 1.0 x 10^7 CFU *S. aureus* Newman,

a human clinical isolate (Duthie and Lorenz, 1952). Five days post-infection, mice were euthanized. Livers and

kidneys were dissected and analyzed for visible abscess lesions. Subsequently, tissue homogenates were

prepared and plated on agar plates to measure bacterial loads in infected organs. Notably, abscess numbers

and bacterial loads were significantly reduced in livers and kidneys in (*R*)-DI-87-treated animals as compared

to control mice validating (*R*)-DI-87 as a novel host-directed and anti-infective drug against staphylococcal

diseases (Figure 4C-F). To test whether *S. aureus* exploits the activity of dCK along with the purine salvage

pathway during infection, cohorts of mice were also challenged with the *S. aureus* Newman *adsA* mutant. (*R*)-

DI-87-treated animals no longer displayed increased resistance to *S. aureus* infection (Figure 4C-F). Moreover,

infection with *S. aureus adsA*-deficient bacteria phenocopied (*R*)-DI-87-mediated inhibition of dCK, in line with

the concept that AdsA is required for establishing persistent infections in host tissues (Figure 4C-F). In light of

these findings, infected organs were also fixed with formalin, embedded into paraffin, thin-sectioned, and

analyzed for histopathology. As expected, histopathological analysis of hematoxylin and eosin (H&E)-stained

liver or renal tissues revealed that wild-type *S. aureus* Newman formed structured abscesses in vehicle-treated

mice (Figure 4G-H). However, lesions obtained from (*R*)-DI-87-treated animals appeared smaller in size and

typically did not harbor a discernable organization of staphylococci, further demonstrating that administration

of (*R*)-DI-87 represents a medically valuable strategy to attenuate staphylococcal virulence (Figure 4G-H).

Presumably, (*R*)-DI-87-treatment protects macrophages against AdsA-derived death-effector

deoxyribonucleosides and therefore enhances phagocyte survival during abscess formation as (*R*)-DI-87

neither improved the killing of *S. aureus* in human or mouse blood, nor it displayed antimicrobial activity

(supplementary Figure 3A-B; supplementary Table 1). Collectively, these data indicate that inhibition of

mammalian dCK by (*R*)-DI-87 attenuates *S. aureus* abscess formation and disease pathogenesis *in vivo*.

(*R*)-DI-87-treatment amplifies macrophage infiltration into staphylococcal infectious foci. Previous studies

revealed that macrophages with defects in the purine salvage pathway-apoptosis axis are refractory to

staphylococcal dAdo or dGuo (Winstel et al., 2018; Winstel et al., 2019; Tantawy et al., 2022). As a result,

death-effector deoxyribonucleoside-resistant tissue macrophages were found to accumulate within deep-

seated abscesses of *S. aureus*-challenged mice, a phenomenon that contributed to accelerated clearance of

staphylococci (Winstel et al., 2019). Thus, we hypothesized that pharmacological inhibition of host dCK might

augment phagocyte infiltration into *S. aureus*-derived abscesses, thereby explaining reduced bacterial burdens

in (*R*)-DI-87-treated laboratory animals. To pursue this possibility, we established an immunofluorescence

microscopy-based approach to detect macrophages in liver or renal tissues of *S. aureus* Newman wild type-

infected C57BL/6 mice that were treated with (*R*)-DI-87 or vehicle only. As expected, immunofluorescence

staining of tissue abscesses obtained from vehicle-treated animals revealed that F4/80-positive macrophages

resided at the periphery of infectious foci (Figure 5A-D). On the contrary, lesions derived from mice that

received (*R*)-DI-87 differed as they contained infiltrates of F4/80-positive macrophages within the neutrophil

cuff, suggesting that (*R*)-DI-87-mediated inhibition of host dCK terminates macrophage exclusion from

staphylococcal abscesses (Figure 5E-H). To further delineate whether these results correlate with AdsA and

staphylococcal death-effector deoxyribonucleoside-mediated manipulation of host dCK in phagocytes, tissues

of *S. aureus* Newman *adsA* mutant-infected mice, which received either (*R*)-DI-87 or vehicle, were also

examined via immunofluorescence microscopy and analyzed for the presence of F4/80-positive macrophages.

Of note, *adsA* mutant-derived abscess lesions were also characterized by increased infiltration rates of F4/80-

positive phagocytes, irrespectively of whether animals were treated with (*R*)-DI-87 or vehicle during the course

of the infection (Figure 5I-P). Thus, (*R*)-DI-87-mediated blockade of host dCK phenocopied the *adsA* mutation

in *S. aureus*. Lastly, (*R*)-DI-87- or vehicle-treated murine bone marrow-derived macrophages (BMDMs) mice

were analyzed for survival rates upon exposure to dAdo or dGuo. Compared to vehicle-treated cells, which

were susceptible to death-effector deoxyribonucleosides, BMDMs exposed to (*R*)-DI-87 displayed increased

resistance toward dAdo and dGuo presumably explaining their abundance in infectious foci obtained from

wild-type *S. aureus* Newman-challenged and dCK inhibitor-treated mice (Figure 5Q-R). In summary, these data

suggest that (*R*)-DI-87-treatment protects macrophages from staphylococcal death-effector

deoxyribonucleosides and therefore boosts their infiltration into persistent abscesses in organ tissues (Figure

6).

Discussion

Bacterial infectious diseases are typically treated with specific or broad-spectrum antibiotics. However, many

microbial pathogens have developed resistance mechanisms against these drugs and other antimicrobial

agents either via mutational changes within the drug target or by importing resistance genes from other

bacteria (Klevens et al., 2008; Yaw et al., 2014; Nubel, 2016; Jian et al., 2021; GBD 2019 Antimicrobial

Resistance Collaborators, 2022a). Accordingly, infections by multidrug-resistant microbes are frequently

associated with failure of therapy and increased mortality in both, industrialized and developing countries

around the world (Klevens et al., 2008; Yaw et al., 2014; GBD 2019 Antimicrobial Resistance Collaborators,

2022a; b). Each year, several million people die due to infections caused by drug-resistant bacteria such as

Mycobacterium tuberculosis, *E. coli*, or *S. aureus* underscoring the very urgent need of new therapeutics and

anti-infective compounds against which bacteria cannot develop resistance (Pai et al., 2016; GBD 2019

Antimicrobial Resistance Collaborators, 2022a; b).

Host-directed therapy is an emerging therapeutic concept that eliminates the possibility of drug

resistance development in microbial pathogens (Kaufmann et al., 2018). This innovative approach aims at

targeting specific host determinants and signaling cascades bacterial or other pathogenic microorganisms may

attack or exploit to establish acute and persistent infections (Kaufmann et al., 2018). Overall, this strategy

seeks to ameliorate host immunity, immune cell survival, and immunopathology, as well as pathogen control

without interfering with microbial replication, growth, or metabolism (Kaufmann et al., 2018). By applying this

concept along with a drug-repurposing approach, we here report that administration of (*R*)-DI-87, a soluble,

orally active, and high-affinity inhibitor of host dCK (Poddar et al., 2020), attenuates *S. aureus* virulence in a

mouse model of bloodstream infection. In particular, we show that (*R*)-DI-87-mediated blockade of host dCK

prevents staphylococcal death-effector deoxyribonucleoside-triggered apoptotic cell death in phagocytes of

human and animal origin. Accordingly, oral administration of (*R*)-DI-87 in mice rendered host phagocytes

refractory to AdsA-derived dAdo and dGuo thereby leading to an accumulation of *S. aureus*-eliminating

macrophages within deeper cavities of infectious foci along with accelerated phagocytic clearance of

staphylococci. Since dCK is a rate-limiting enzyme of the nucleoside salvage pathway that catalyzes the

intracellular conversion of deoxyribonucleosides into deoxyribonucleoside monophosphates (Arner and

Eriksson, 1995), (*R*)-DI-87-mediated inhibition of dCK most likely suppresses an uncontrolled biogenesis of

deoxyribonucleoside di- and triphosphates that may trigger replication errors and apoptosis in host cells.

Consistent with this model, earlier work suggested that dCK deficiency in phagocytes or other cells such as

lymphoblasts prevents an overload of intracellular deoxyribonucleoside triphosphates upon exposure to

death-effector deoxyribonucleosides (Gudas et al., 1978; Winstel et al., 2018). Thus, inhibition of host dCK

represents an attractive host-directed therapeutic intervention strategy that terminates a refined immune-

evasive maneuver *S. aureus* has evolved to establish persistent infections in mammalian hosts. The

technological advantage of this therapeutic approach, beyond the clinical importance and long-term economic

benefits for the medical-therapeutic sector, is undoubtedly the exploitation of an already existing, highly

selective, and clinical-stage compound that can easily be administered to attenuate infections caused by a

predominant bacterial pathogen. In particular, (*R*)-DI-87 does not exhibit detrimental side effects as (*R*)-DI-87-

exposed laboratory animals neither lost weight during long-term treatment nor displayed any other adverse

effects that may negatively affect host immune responses. Specifically, the immune cell composition profile

of mice that received (*R*)-DI-87 was not altered when compared to mice that received the vehicle control. This

suggests that patrolling neutrophils known to play a significant role during staphylococcal infections (Spaan et
al., 2013) along with other crucial elements of the host immune system are fully functional in (*R*)-DI-87-treated
hosts, thereby offering tremendous opportunities to improve infection outcomes in hospitalized or critically
ill patients. Further, usage of (*R*)-DI-87 represents a host-directed and anti-infective therapeutic approach to
which *S. aureus* and other multidrug-resistant bacteria cannot develop resistance mechanisms. In this regard,
we note that many other medically highly relevant bacterial pathogens including streptococci, *Bacillus*
anthracis, and multidrug-resistant *Staphylococcus pseudintermedius* synthesize effector-nucleosides in order
to modulate host immune cell responses (Thammavongsa et al., 2009; Liu et al., 2014; Zheng et al., 2015; Ma
et al., 2017; Dai et al., 2018; Bunsow et al., 2021). Thus, (*R*)-DI-87-mediated inhibition of dCK may also help to
protect host cells from apoptotic cell death or other adverse effects during other bacterial diseases in human
or animal hosts. Even polymicrobial infections caused by various microbial pathogens and other infectious
agents that manipulate the purine salvage pathway-apoptosis axis during infection can eventually be
prevented, attenuated, or cured by using (*R*)-DI-87. Ultimately, double-pronged therapies which cover the
usage of (*R*)-DI-87 combined with antibiotics, various anti-infective agents, or monoclonal antibodies, some of
which are known to neutralize predominant staphylococcal virulence determinants, including secreted toxins
(Ragle and Bubeck Wardenburg, 2009; Hua et al., 2014; Thomsen et al., 2017), protein A (Kim et al., 2012;
Chen et al., 2022b), and specific surface antigens (Bennett et al., 2019; de Vor et al., 2022) along with usage
of other host-directed therapeutics that impede *S. aureus* disease pathogenesis (Bravo-Santano et al., 2019;
Alphonse et al., 2021), may essentially improve disease progression during severe bacterial infectious diseases
and should therefore be considered for pre-clinical tests in animal models.

Overall, (*R*)-DI-87-mediated inhibition of dCK is a novel host-directed therapeutic concept to improve
clinical outcomes of *S. aureus* infectious diseases. Since (*R*)-DI-87 is extremely well tolerated in humans
(personal communication; Kenneth A. Schultz, Trethera Corporation, Los Angeles, CA, USA) and currently
being investigated in phase I clinical trials for the treatment of non-communicable diseases, including cancer
and multiple sclerosis (Poddar et al., 2020; Chen et al., 2022a), this compound already meets the requirements
for a prompt launch of clinical studies in individuals that suffer local or invasive staphylococcal infections.
Optimization of (*R*)-DI-87 or any medically acceptable derivatives thereof may further accelerate this process
and concurrently could also aid in the design of pre-exposure prophylactic agents for hospitalized or high-risk
patients, with the primary aim to enhance infection control and public health.

Materials und methods

Bacterial strains and growth conditions. All bacterial strains used in this study are listed in the *Supplementary*
Materials (supplementary Table 2). Bacteria were grown in tryptic soy broth (TSB), Mueller-Hinton broth

(MHB), or lysogeny broth (LB) at permissive temperatures. Media were supplemented with appropriate antibiotics (ampicillin 100 μg/ml).

Cell lines and tissue culture. U937 cells were obtained from American Type Culture Collection (ATCC) and grown in Roswell Park Memorial Institute (RPMI) 1640 medium (Gibco) supplemented with 10% heat-inactivated fetal bovine serum (hi-FBS) according to the manufacturer's instructions. All mammalian cell lines were grown at 37°C under 5% CO_2. Cell lines used in this study are listed in the *Supplementary Materials* (supplementary Table 3).

Isolation of human primary cells. Peripheral blood mononuclear cells (PBMCs) were isolated from heparinized blood by density gradient centrifugation on Pancoll (PAN Biotech) according to standard laboratory protocols. Next, human primary monocytes were isolated from PBMCs by using magnetic nanobeads and the MojoSort™ Human CD14 Selection Kit (BioLegend) as described earlier (Tantawy et al., 2022). Subsequently, purified primary CD14-positive (CD14+) monocytes were resuspended in RPMI 1640 medium containing 10% hi-FBS and 1% penicillin-streptomycin and used for cytotoxicity assays. Alternatively, purified CD14+ monocytes were differentiated into human monocyte-derived macrophages (HMDMs) in the same medium supplemented with 50 ng/ml of human macrophage colony stimulating factor (hM-CSF; Genscript). HMDMs were used at day 7 post-differentiation for cytotoxicity experiments.

Isolation of murine bone marrow-derived macrophages. Bone marrow-derived macrophages (BMDMs) were isolated from C57BL/6 mice as previously described (Tantawy et al., 2022). In brief, mice were euthanized to remove the femur and tibia which were sterilized by using 70% ethanol. Bones were then washed with sterile phosphate-buffered saline (PBS). Next, the bone ends were removed and the bone marrow was flushed out by using RPMI 1640 containing 10% hi-FBS and 1% penicillin-streptomycin. Following a resuspension step, cells were passed through a nylon cell strainer (40 μm) to remove unwanted tissue and cellular debris. Subsequently, cells were centrifuged (10 min, 200 x g, 4°C), resuspended in red blood cell (RBC) lysis buffer (Roche), and incubated for 5 min at room temperature to lyse RBC. Cells were centrifuged once more, resuspended in BMDM medium (RPMI 1640 containing 10% hi-FBS, 1% penicillin-streptomycin, and 50 ng/ml of mouse macrophage colony-stimulating factor (Genscript)), and seeded into tissue culture-treated dishes to deplete bone marrow cells from fibroblasts. At day 1 post-extraction, suspension bone marrow cells were collected via centrifugation, adjusted to 6.0×10^5 cells/ml in BMDM medium, and re-seeded into bacteriological dishes. At day 4 post-extraction, cells were incubated with an additional 10 ml of BMDM medium. BMDMs were used at day 7 post-extraction for cytotoxicity experiments.

Protein purification. Purification of recombinant *S. aureus* AdsA was performed as described elsewhere (Thammavongsa et al., 2009; Tantawy et al., 2022). Briefly, a glutathione S-transferase (GST)-tagged and recombinant version of *S. aureus* AdsA (rAdsA) was expressed in *E. coli* BL21 using the pGEX-2T plasmid system (GE Healthcare) and purified via glutathione S-transferase affinity chromatography (Thammavongsa et al., 2009; Tantawy et al., 2022). Next, the N-terminal GST tag was removed by using thrombin which was immediately cleared from the protein sample via benzamidine sepharose beads according to the manufacturer's instructions (GE Healthcare). Purified rAdsA was analyzed by a Coomassie-stained SDS-PAGE following standard laboratory protocols.

Cytotoxicity assays. To evaluate the protective role of (*R*)-DI-87 during death-effector deoxyribonucleoside-induced cytotoxicity, 2.0×10^5 U937 monocyte-like cells (U937) per well were seeded in a 24-well plate and pre-incubated for 2 h at 37°C under 5% CO_2 in RPMI 1640 medium containing 10% hi-FBS and 1 µM of (*R*)-DI-87. Control wells received vehicle only. Alternatively, 4.0×10^5 U937 cells per well were seeded in a 24-well plate and incubated for 48 h at 37°C under 5% CO_2 in RPMI 1640 growth medium that contained 160 nM phorbol 12-myristate 13-acetate (PMA). Resulting U937-derived macrophages (U937 MΦ) were washed and further incubated in RPMI 1640 growth medium lacking PMA (24 h). Similarly, 3.5×10^5 BMDMs per well were seeded in 24-well plates and incubated for 24 h at 37°C under 5% CO_2 in corresponding growth media. Next, U937 MΦ or BMDMs were washed once and pre-incubated for 2 h at 37°C under 5% CO_2 in appropriate growth media supplemented with 1 µM of (*R*)-DI-87. Control wells received vehicle only. Following pre-incubation with (*R*)-DI-87, U937, U937 MΦ, or BMDMs were exposed to various concentrations of either dAdo or dGuo (100 µM for U937; 200 µM for U937 MΦ or BMDMs) and incubated for 48 h at 37°C under 5% CO_2 in RPMI 1640 growth medium. Cells were collected via centrifugation (U937) or a detachment-centrifugation step using trypsin-EDTA (U937 MΦ) or accutase (BMDMs) solution. Dead cells were stained with trypan blue and counted by using a microscope to calculate killing efficiency. To analyze the protective effect of (*R*)-DI-87 in primary human cells, 2.0×10^5 CD14[+] monocytes or HMDMs per well were seeded in multi-well plates and pre-incubated for 2 h at 37°C under 5% CO_2 in RPMI 1640 medium containing 10% hi-FBS and 1 µM of (*R*)-DI-87. Control wells received vehicle only. Subsequently, cells were intoxicated by using various concentrations of either dAdo or dGuo (100 µM for CD14[+] monocytes; 200 µM for HMDMs) and incubated in appropriate growth media for 48 h at 37°C under 5% CO_2. Cells were collected via centrifugation (CD14[+] monocytes) or a detachment-centrifugation step (HMDMs) using trypsin-EDTA solution. Viability of cells was determined via trypan blue staining and microscopy as described above. To block the cytotoxic effect of AdsA-derived death-effector deoxyribonucleosides by using (*R*)-DI-87, rAdsA (1.25 µg/µl) was incubated for 16 h at 37°C in a reaction buffer (30 mM Tris-HCl, pH 7.5; 1.5 mM $MgCl_2$; 1.5 mM $MnCl_2$) supplemented with either dAMP or

dGMP (1.19 mM each) according to a published protocol (Tantawy et al., 2022). Controls included reactions

that lacked deoxyribonucleoside monophosphates or rAdsA. Following incubation, all reaction products were

filter-sterilized and added to U937 MΦ which were pre-incubated with vehicle or 1 µM (*R*)-DI-87 for 2 h prior

to intoxication. Next, cells were incubated at 37°C under 5% CO_2 for 48 h, collected, and analyzed via trypan

blue staining and microscopy as described above.

Assessment of caspase-3-activity. Caspase-3 activity was analyzed using a colorimetric caspase-3 detection

kit (Sigma) and a published protocol (Winstel et al., 2018; Tantawy et al., 2022). Briefly, U937 cells or U937

MΦ along with appropriate controls were pre-incubated for 2 h at 37°C under 5% CO_2 in RPMI 1640 medium

containing 10% hi-FBS and 1 µM of (*R*)-DI-87. Controls received vehicle only. Subsequently, cells were exposed

to various concentrations of either dAdo or dGuo (100 µM for U937; 200 µM for U937 MΦ) and incubated for

24 h at 37°C under 5% CO_2 in RPMI 1640 growth medium. Cells were collected via centrifugation (U937) or a

detachment-centrifugation step (U937 MΦ) using trypsin-EDTA solution and washed once in PBS. Next, 1.0 x

10^7 cells were lysed in pre-chilled lysis buffer (Sigma kit) for 20 min. This step was performed on ice. Resulting

lysates were centrifuged at 4°C (18,000 × g for 10 min) to obtain cell- and debris-free supernatants which were

incubated with the caspase-3 substrate Ac-DEVD-pNA according to the manufacturer's instructions. Caspase-

3 activity was determined based on the amount of released pNA that can be detected at 405 nm.

Immunoblotting. Immunoblotting was performed as described elsewhere (Tantawy et al., 2022). In short,

U937 or U937 MΦ and appropriate controls were pre-incubated for 2 h at 37°C under 5% CO_2 in RPMI 1640

medium containing 10% hi-FBS and 1 µM of (*R*)-DI-87. Controls received vehicle only. Next, cells were treated

with dAdo or dGuo (100 µM for U937; 200 µM for U937 MΦ), incubated for 24 h at 37°C under 5% CO_2 in RPMI

1640 growth medium, and collected via centrifugation (U937) or a detachment-centrifugation step (U937 MΦ)

using trypsin-EDTA solution. Cells were washed once in PBS. 1.0 x 10^7 cells were lysed on ice in pre-chilled lysis

buffer (50 mM HEPES, pH 7.4; 5 mM CHAPS; 5 mM DTT) for 20 min. Resulting lysates were centrifuged at 4°C

(18,000 × g for 10 min) to obtain cell- and debris-free supernatants which were mixed with sodium dodecyl

sulfate-polyacrylamide gel (SDS-PAGE) loading buffer. Samples were boiled for 10 min at 95°C. Proteins were

separated via SDS-PAGE (12%) and transferred onto PVDF membranes for immunoblot analysis with the

following rabbit primary antibodies: α-Caspase-3 (α-CASP3, 9662, Cell Signaling) and α-GAPDH (ab181602,

Abcam, loading control). Immunoreactive signals were revealed with a secondary antibody conjugated to

horseradish peroxidase (α-rabbit IgG, 7074, Cell Signaling). Horseradish peroxidase activity was detected with

enhanced chemiluminescent (ECL) substrate (Thermo Fisher).

FITC-annexin-V/PI staining. FITC-annexin-V/PI staining of U937 MΦ or HMDMs exposed to 1 µM (*R*)-DI-87 and death-effector deoxyribonucleosides (200 µM dAdo or dGuo; 24 h at 37°C) along with appropriate controls was performed by using FITC-annexin-V Apoptosis Detection Kit I (BD Biosciences) according to the manufacturer's instructions. Stained cells were analyzed via immunofluorescence microscopy according to standard laboratory protocols.

Determination of the minimal inhibitory concentration (MIC). The MIC of (*R*)-DI-87 was determined in 96-well plates by using the microdilution method and Mueller-Hinton broth according to standard laboratory protocols. Wells containing varying concentrations of (*R*)-DI-87 or vehicle were inoculated with 10^5 CFU/ml of wild-type *S. aureus* Newman or its *adsA* mutant and incubated for 24 h at 37°C under continuous shaking. The MIC was defined as the lowest concentration of compound at which no visible growth was detected.

Bacterial survival in blood. Bacterial survival in mouse or human blood was analyzed as described before (Thammavongsa et al., 2009). In brief, fresh overnight cultures of *S. aureus* Newman wildtype or its *adsA* were diluted into fresh TSB medium and grown at 37°C to an OD of 1.0. Bacteria were washed twice in sterile PBS and adjusted in PBS to a final density of 1.0 x 10^8 CFU/ml. Next, freshly drawn mouse or human blood anticoagulated with heparin was incubated in the presence or absence of 1 µM (*R*)-DI-87 for 60 min (37°C) with *S. aureus* Newman wildtype or its *adsA* mutant using a bacterial dose of 1.0 x 10^6 CFU/ml (murine blood) or 1.0 x 10^7 CFU/ml (human blood). Samples without (*R*)-DI-87 treatment received the vehicle. Following incubation, blood samples were mixed in a 1:1 ratio with sterile lysis buffer (PBS containing 1.0% saponin) and incubated for 10 min at 37°C to lyse eukaryotic cells. Subsequently, serial dilutions were prepared and plated onto TSA plates to determine bacterial survival rates.

Animal infection model. C57BL/6 mice were purchased from Janvier Laboratories and kept under specific pathogen-free conditions in our central mouse facility (TWINCORE, Center for Experimental and Clinical Infection Research, Hannover, Germany). Mice received (*R*)-DI-87 (75 mg/kg) or vehicle (40% Captisol) via oral gavage at the day of the infection, separated by 12-hour intervals over a 5-day observation period. For infection experiments, TSB overnight cultures of wild-type *S. aureus* Newman or its *adsA* mutant were diluted 1:100 in TSB and grown to an optical density (600 nm) of 0.5. Bacteria were then centrifuged (10 min, RT, 8,000 × g), washed twice in sterile PBS, and adjusted to 10^8 CFU/ml. One hundred microliters of the bacterial suspension (10^7 CFU) were administered intravenously (lateral tail vein) into 6- to 8-weeks-old female C57BL/6 mice. Five days post-infection, animals were euthanized. Organs were dissected, examined for surface abscesses, and homogenized in sterile PBS supplemented with 0.1% Triton X-100. Serial dilutions were prepared and plated onto TSA plates to determine bacterial loads. For histopathology, dissected organs were

fixed in 10% Formalin (Sigma), embedded into paraffin, thin-sectioned, and stained with hematoxylin and
eosin (H&E). Stained tissues were examined by microscopy according to standard laboratory protocols.

Immunofluorescence staining. To detect macrophages in *S. aureus*-infected tissues, formalin-fixed and
paraffin-embedded organs were thin-sectioned, deparaffinized, and rehydrated. After heat-induced antigen
retrieval in 10 mM sodium citrate buffer (pH 6.0), non-specific antibody binding was blocked by adding 2%
normal goat serum. Immunofluorescence staining was carried out by using an antibody against F4/80-positive
macrophages (α-F4/80, 70076, Cell Signaling), followed by a fluorescently-labelled secondary antibody
(Alexa546, Invitrogen). Stained tissues were examined by using a Zeiss Apotome 2 microscope (Zeiss).

Safety assessment of (*R*)-DI-87. To assess the therapeutic value and safety of (*R*)-DI-87, cohorts of female
C57BL/6 mice were treated with (*R*)-DI-87 (75 mg/kg) or vehicle (40% Captisol) via oral gavage in 12-hour
intervals over the course of 16 days. On day 16, 100 µl peripheral blood was collected in lithium heparin-
coated tubes via retro-orbital bleeding using heparin-coated capillary tubes. Subsequently, blood samples
were incubated with 5 ml of ACK lysis buffer at room temperature for 5 min, quenched with 5 ml of FACS
buffer (5% FBS in PBS), and centrifuged at 4°C (4 min). This process was repeated and cells were subsequently
stained with the following fluorochrome-conjugated anti-mouse antibodies diluted 1:100 in 100 µl of FACS
buffer for 20 min at 4°C: α-B220-PerCP/Cy5.5 (103236, BioLegend), α-CD4-BV711 (100550, BioLegend), α-
CD8α-PE (100708, BioLegend), α-CD11c-PE/Dazzle594 (117347, BioLegend), α-CD11b-FITC (101206,
BioLegend), α-GR1-APC (108412, BioLegend); α-CD16/32 (FC block; 101319; BioLegend). Following incubation,
cells were centrifuged and washed twice using FACS buffer. Cells were resuspended in FACS buffer and
analyzed using a BD LSRII flow cytometer and the FlowJo software package. During the (*R*)-DI-87 or vehicle
treatment procedure, mice were regularly monitored and weighed to assess the overall health status.

Ethics statement. All animal experiments were conducted in accordance with the local animal welfare
regulations reviewed by the institutional review board and the Niedersächsisches Landesamt für
Verbraucherschutz und Lebensmittelsicherheit (LAVES) under the permission number 33.19-42502-04-
21/3696. Scientific use of animals (murine blood samples; isolation of BMDMs) was further carried out under
approved animal care and use protocols which are granted by the state veterinary authorities and overseen
by the internal animal care and use committee (IACUC). Animal studies required to assess the safety of (*R*)-DI-
87 in mice were approved by the UCLA Animal Research Committee (ARC). Human blood samples were
obtained from adult, consenting healthy donors. Informed consent forms were obtained from all participants.
These studies were reviewed and approved by the medical ethics committee of Hannover Medical School
(Hannover, Germany) under the permission number 8831_BO_K_2019.

Synthesis and preparation of (*R*)-DI-87. (*R*)-DI-87 ((((*R*)-2-((1-(2-(4-methoxy-3-(2-morpholinoethoxy)phenyl)-
5-methylthiazol-4-yl)ethyl)thio)pyrimidine-4,6-diamine) was synthesized as described before (Poddar et al.,
2020). For experiments, (*R*)-DI-87 was either dissolved in Dimethyl sulfoxide (*in vitro* experiments) or in an
aqueous solution containing 40% Captisol (CyDex Pharmaceuticals, Inc.) (*in vivo* experiments).

Statistical analysis. Statistical analysis was performed by using GraphPad Prism (GraphPad Software, Inc., La
Jolla, USA). Statistically significant differences were calculated by using statistical methods as indicated. *P*
values < 0.05 were considered significant.

Data availability. All data supporting the findings of this study are available within the article and its
supplementary material. All other data are available from the corresponding author upon reasonable request.
488
489
490

491 **Acknowledgments**
492 We thank all laboratory members for helpful discussion. We are grateful for support from the German
493 Research Foundation (award WI4582/2-1 to V.W., project number 449712894). E.R.A. is supported by UCLA
494 Tumor Immunology Training Grant T32CA009120. This work was supported by National Institutes of Health
495 grants to C.G.R. (1R01CA25052901A1 and 1R01CA260678-01).
496

Author contributions
Volker Winstel: Conceptualization; data curation; investigation; methodology; formal analysis; supervision;
funding acquisition; writing – original draft; writing – review and editing; visualization; project administration.
Evan R. Abt: Conceptualization; data curation; investigation; methodology; formal analysis; writing – review
and editing. Caius G. Radu: Conceptualization; investigation; methodology; formal analysis; resources; funding
acquisition; writing – review and editing.
503

504 **Conflict of interest**
505 C.G.R. is a co-inventor of the dCK inhibitor used in this study. This intellectual property has been patented by
506 the University of California and optioned to Trethera Corporation, a company that C.G.R. owns equity in. V.W.,
507 E.R.A., and C.G.R. are inventors in a patent application covering the use of dCK inhibitors as a treatment for
508 bacterial infectious diseases.
509
510

References

Alphonse MP, Rubens JH, Ortines RV, Orlando NA, Patel AM, Dikeman D, Wang Y, Vuong I, Joyce DP, Zhang J,
Mumtaz M, Liu H, Liu Q, Youn C, Patrick GJ, Ravipati A, Miller RJ, Archer NK, and Miller LS (2021). Pan-
caspase inhibition as a potential host-directed immunotherapy against MRSA and other bacterial skin
infections. *Sci Transl Med* 13(601). doi: 10.1126/scitranslmed.abe9887.

Arner ES, and Eriksson S (1995). Mammalian deoxyribonucleoside kinases. *Pharmacol Ther* 67(2)**,** 155-186. doi:
10.1016/0163-7258(95)00015-9.

Bennett MR, Bombardi RG, Kose N, Parrish EH, Nagel MB, Petit RA, Read TD, Schey KL, Thomsen IP, Skaar EP,
and Crowe JE (2019). Human mAbs to Staphylococcus aureus IsdA Provide Protection Through Both
Heme-Blocking and Fc-Mediated Mechanisms. *J Infect Dis* 219(8)**,** 1264-1273. doi:
10.1093/infdis/jiy635.

Berends ET, Horswill AR, Haste NM, Monestier M, Nizet V, and von Kockritz-Blickwede M (2010). Nuclease
expression by Staphylococcus aureus facilitates escape from neutrophil extracellular traps. *J Innate*
Immun 2(6)**,** 576-586. doi: 10.1159/000319909.

Bravo-Santano N, Stolting H, Cooper F, Bileckaja N, Majstorovic A, Ihle N, Mateos LM, Calle Y, Behrends V, and
Letek M (2019). Host-directed kinase inhibitors act as novel therapies against intracellular
Staphylococcus aureus. *Sci Rep* 9(1)**,** 4876. doi: 10.1038/s41598-019-41260-8.

Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, Weinrauch Y, and Zychlinsky A (2004).
Neutrophil extracellular traps kill bacteria. *Science* 303(5663)**,** 1532-1535. doi:
10.1126/science.1092385.

Bunsow D, Tantawy E, Ostermeier T, Bahre H, Garbe A, Larsen J, and Winstel V (2021). Methicillin-resistant
Staphylococcus pseudintermedius synthesizes deoxyadenosine to cause persistent infection.
Virulence 12(1)**,** 989-1002. doi: 10.1080/21505594.2021.1903691.

Chambers HF, and Deleo FR (2009). Waves of resistance: Staphylococcus aureus in the antibiotic era. *Nat Rev*
Microbiol 7(9)**,** 629-641. doi: 10.1038/nrmicro2200.

Chen BY, Salas JR, Trias AO, Perez Rodriguez A, Tsang JE, Guemes M, Le TM, Galic Z, Shepard HM, Steinman L,
Nathanson DA, Czernin J, Witte ON, Radu CG, Schultz KA, and Clark PM (2022a). Targeting
deoxycytidine kinase improves symptoms in mouse models of multiple sclerosis. *Immunology*. doi:
10.1111/imm.13569.

Chen X, Schneewind O, and Missiakas D (2022b). Engineered human antibodies for the opsonization and killing
of Staphylococcus aureus. *Proc Natl Acad Sci U S A* 119(4). doi: 10.1073/pnas.2114478119.

Cheng AG, DeDent AC, Schneewind O, and Missiakas D (2011). A play in four acts: Staphylococcus aureus
abscess formation. *Trends Microbiol* 19(5)**,** 225-232. doi: 10.1016/j.tim.2011.01.007.

Dai J, Lai L, Tang H, Wang W, Wang S, Lu C, Yao H, Fan H, and Wu Z (2018). Streptococcus suis synthesizes
deoxyadenosine and adenosine by 5'-nucleotidase to dampen host immune responses. *Virulence* 9(1)**,**
1509-1520. doi: 10.1080/21505594.2018.1520544.

de Vor L, van Dijk B, van Kessel K, Kavanaugh JS, de Haas C, Aerts PC, Viveen MC, Boel EC, Fluit AC, Kwiecinski
JM, Krijger GC, Ramakers RM, Beekman FJ, Dadachova E, Lam MG, Vogely HC, van der Wal BC, van
Strijp JA, Horswill AR, Weinans H, et al. (2022). Human monoclonal antibodies against Staphylococcus
aureus surface antigens recognize in vitro and in vivo biofilm. *Elife* 11. doi: 10.7554/eLife.67301.

Duthie ES, and Lorenz LL (1952). Staphylococcal coagulase; mode of action and antigenicity. *J Gen Microbiol*
6(1-2)**,** 95-107. doi: 10.1099/00221287-6-1-2-95.

Finlay BB, and McFadden G (2006). Anti-immunology: evasion of the host immune system by bacterial and
viral pathogens. *Cell* 124(4)**,** 767-782. doi: 10.1016/j.cell.2006.01.034.

Foster TJ (2005). Immune evasion by staphylococci. *Nat Rev Microbiol* 3(12)**,** 948-958. doi:
10.1038/nrmicro1289.

Fridkin SK, Hageman JC, Morrison M, Sanza LT, Como-Sabetti K, Jernigan JA, Harriman K, Harrison LH, Lynfield
R, and Farley MM (2005). Methicillin-resistant *Staphylococcus aureus* disease in three communities.
N. Engl. J. Med. 352**,** 1436-1444.
GBD 2019 Antimicrobial Resistance Collaborators (2022a). Global burden of bacterial antimicrobial resistance
in 2019: a systematic analysis. *Lancet* 399(10325)**,** 629-655. doi: 10.1016/S0140-6736(21)02724-0.
GBD 2019 Antimicrobial Resistance Collaborators (2022b). Global mortality associated with 33 bacterial
pathogens in 2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*
400(10369)**,** 2221-2248. doi: 10.1016/S0140-6736(22)02185-7.
Gudas LJ, Ullman B, Cohen A, and Martin DW, Jr. (1978). Deoxyguanosine toxicity in a mouse T lymphoma:
relationship to purine nucleoside phosphorylase-associated immune dysfunction. *Cell* 14(3)**,** 531-538.
doi: 10.1016/0092-8674(78)90239-8.
Hua L, Hilliard JJ, Shi Y, Tkaczyk C, Cheng LI, Yu X, Datta V, Ren S, Feng H, Zinsou R, Keller A, O'Day T, Du Q,
Cheng L, Damschroder M, Robbie G, Suzich J, Stover CK, and Sellman BR (2014). Assessment of an anti-
alpha-toxin monoclonal antibody for prevention and treatment of Staphylococcus aureus-induced
pneumonia. *Antimicrob Agents Chemother* 58(2)**,** 1108-1117. doi: 10.1128/AAC.02190-13.
Jian Z, Zeng L, Xu T, Sun S, Yan S, Yang L, Huang Y, Jia J, and Dou T (2021). Antibiotic resistance genes in bacteria:
Occurrence, spread, and control. *J Basic Microbiol* 61(12)**,** 1049-1070. doi: 10.1002/jobm.202100201.
Kaufmann SHE, Dorhoi A, Hotchkiss RS, and Bartenschlager R (2018). Host-directed therapies for bacterial and
viral infections. *Nat Rev Drug Discov* 17(1)**,** 35-56. doi: 10.1038/nrd.2017.162.
Kim HK, Emolo C, DeDent AC, Falugi F, Missiakas DM, and Schneewind O (2012). Protein A-specific monoclonal
antibodies and prevention of Staphylococcus aureus disease in mice. *Infect Immun* 80(10)**,** 3460-3470.
doi: 10.1128/IAI.00230-12.
Klevens RM, Edwards JR, Gaynes RP, and National Nosocomial Infections Surveillance S (2008). The impact of
antimicrobial-resistant, health care-associated infections on mortality in the United States. *Clin Infect*
Dis 47(7)**,** 927-930. doi: 10.1086/591698.
Klevens RM, Morrison MA, Nadle J, Petit S, Gershman K, Ray S, Harrison LH, Lynfield R, Dumyati G, Townes JM,
Craig AS, Zell ER, Fosheim GE, McDougal LK, Carey RB, and Fridkin SK (2007). Invasive methicillin-
resistant *Staphylococcus aureus* infections in the United States. *JAMA* 298**,** 1763-1771.
Kuehnert MJ, Kruszon-Moran D, Hill HA, McQuillan G, McAllister SK, Fosheim G, McDougal LK, Chaitram J,
Jensen B, Fridkin SK, Killgore G, and Tenover FC (2006). Prevalence of *Staphylococcus aureus* nasal
colonization in the United States, 2001-2002. *J. Infect. Dis.* 193**,** 172-179.
Lee AS, de Lencastre H, Garau J, Kluytmans J, Malhotra-Kumar S, Peschel A, and Harbarth S (2018). Methicillin-
resistant Staphylococcus aureus. *Nat Rev Dis Primers* 4**,** 18033. doi: 10.1038/nrdp.2018.33.
Liu P, Pian Y, Li X, Liu R, Xie W, Zhang C, Zheng Y, Jiang Y, and Yuan Y (2014). Streptococcus suis adenosine
synthase functions as an effector in evasion of PMN-mediated innate immunit. *J Infect Dis* 210(1)**,** 35-
45. doi: 10.1093/infdis/jiu050.
Lowy   FD   (1998).   Staphylococcus   aureus   infections.   *N   Engl   J   Med*   339(8)**,**   520-532.   doi:
10.1056/NEJM199808203390806.
Ma F, Guo X, and Fan H (2017). Extracellular Nucleases of Streptococcus equi subsp. zooepidemicus Degrade
Neutrophil Extracellular Traps and Impair Macrophage Activity of the Host. *Appl Environ Microbiol*
83(2). doi: 10.1128/AEM.02468-16.
McKenna M (2013). Antibiotic resistance: the last resort. *Nature* 499(7459)**,** 394-396. doi: 10.1038/499394a.
Meka VG, and Gold HS (2004). Antimicrobial resistance to linezolid. *Clin Infect Dis* 39(7)**,** 1010-1015. doi:
10.1086/423841.
Nubel U (2016). Emergence and Spread of Antimicrobial Resistance: Recent Insights from Bacterial Population
Genomics. *Curr Top Microbiol Immunol* 398**,** 35-53. doi: 10.1007/82_2016_505.
Okumura CY, and Nizet V (2014). Subterfuge and sabotage: evasion of host innate defenses by invasive gram-
positive bacterial pathogens. *Annu Rev Microbiol* 68**,** 439-458. doi: 10.1146/annurev-micro-092412-
155711.

Pai M, Behr MA, Dowdy D, Dheda K, Divangahi M, Boehme CC, Ginsberg A, Swaminathan S, Spigelman M,
Getahun H, Menzies D, and Raviglione M (2016). Tuberculosis. *Nat Rev Dis Primers* 2**,** 16076. doi:
10.1038/nrdp.2016.76.
Poddar S, Capparelli EV, Rosser EW, Gipson RM, Wei L, Le T, Jung ME, Radu C, and Nikanjam M (2020).
Development and preclinical pharmacology of a novel dCK inhibitor, DI-87. *Biochem Pharmacol* 172**,**
113742. doi: 10.1016/j.bcp.2019.113742.
Ragle BE, and Bubeck Wardenburg J (2009). Anti-alpha-hemolysin monoclonal antibodies mediate protection
against Staphylococcus aureus pneumonia. *Infect Immun* 77(7)**,** 2712-2718. doi: 10.1128/IAI.00115-
09.
Spaan AN, Surewaard BG, Nijland R, and van Strijp JA (2013). Neutrophils versus Staphylococcus aureus: a
biological tug of war. *Annu Rev Microbiol* 67**,** 629-650. doi: 10.1146/annurev-micro-092412-155746.
Tantawy E, Schwermann N, Ostermeier T, Garbe A, Bahre H, Vital M, and Winstel V (2022). Staphylococcus
aureus Multiplexes Death-Effector Deoxyribonucleosides to Neutralize Phagocytes. *Front Immunol* 13**,**
847171. doi: 10.3389/fimmu.2022.847171.
Thammavongsa V, Kern JW, Missiakas DM, and Schneewind O (2009). Staphylococcus aureus synthesizes
adenosine to escape host immune responses. *J Exp Med* 206(11)**,** 2417-2427. doi:
10.1084/jem.20090097.
Thammavongsa V, Kim HK, Missiakas D, and Schneewind O (2015). Staphylococcal manipulation of host
immune responses. *Nat Rev Microbiol* 13(9)**,** 529-543. doi: 10.1038/nrmicro3521.
Thammavongsa V, Missiakas DM, and Schneewind O (2013). Staphylococcus aureus degrades neutrophil
extracellular traps to promote immune cell death. *Science* 342(6160)**,** 863-866. doi:
10.1126/science.1242255.
Thomsen IP, Sapparapu G, James DBA, Cassat JE, Nagarsheth M, Kose N, Putnam N, Boguslawski KM, Jones LS,
Wood JB, Creech CB, Torres VJ, and Crowe JE, Jr. (2017). Monoclonal Antibodies Against the
Staphylococcus aureus Bicomponent Leukotoxin AB Isolated Following Invasive Human Infection
Reveal Diverse Binding and Modes of Action. *J Infect Dis* 215(7)**,** 1124-1131. doi: 10.1093/infdis/jix071.
Tong SY, Davis JS, Eichenberger E, Holland TL, and Fowler VG, Jr. (2015). Staphylococcus aureus infections:
epidemiology, pathophysiology, clinical manifestations, and management. *Clin Microbiol Rev* 28(3)**,**
603-661. doi: 10.1128/CMR.00134-14.
Torres A, Cilloniz C, Niederman MS, Menendez R, Chalmers JD, Wunderink RG, and van der Poll T (2021).
Pneumonia. *Nat Rev Dis Primers* 7(1)**,** 25. doi: 10.1038/s41572-021-00259-0.
von Kockritz-Blickwede M, and Winstel V (2022). Molecular Prerequisites for Neutrophil Extracellular Trap
Formation and Evasion Mechanisms of Staphylococcus aureus. *Front Immunol* 13**,** 836278. doi:
10.3389/fimmu.2022.836278.
Wenzel RP (2007). Health care-associated infections: major issues in the early years of the 21st century. *Clin
Infect Dis* 45 Suppl 1**,** S85-88. doi: 10.1086/518136.
Winstel V, Missiakas D, and Schneewind O (2018). Staphylococcus aureus targets the purine salvage pathway
to kill phagocytes. *Proc Natl Acad Sci U S A* 115(26)**,** 6846-6851. doi: 10.1073/pnas.1805622115.
Winstel V, Schneewind O, and Missiakas D (2019). Staphylococcus aureus Exploits the Host Apoptotic Pathway
To Persist during Infection. *mBio* 10(6)**,** e02270-02219. doi: 10.1128/mBio.02270-19.
Yaw LK, Robinson JO, and Ho KM (2014). A comparison of long-term outcomes after meticillin-resistant and
meticillin-sensitive Staphylococcus aureus bacteraemia: an observational cohort study. *Lancet Infect
Dis* 14(10)**,** 967-975. doi: 10.1016/S1473-3099(14)70876-X.
Zheng L, Khemlani A, Lorenz N, Loh JM, Langley RJ, and Proft T (2015). Streptococcal 5'-Nucleotidase A (S5nA),
a Novel Streptococcus pyogenes Virulence Factor That Facilitates Immune Evasion. *J Biol Chem*
290(52)**,** 31126-31137. doi: 10.1074/jbc.M115.677443.

Figure 1. Mode of action of the dCK-specific inhibitor (*R*)-DI-87. (A) Scheme illustrating the mode of action of the dCK-specific inhibitor (*R*)-DI-87. *S. aureus*-derived dAdo and dGuo are pumped into phagocytes via human equilibrative transporter 1 (hENT1). Deoxycytidine kinase (dCK) converts dAdo and dGuo into appropriate deoxyribonucleoside monophosphates thereby triggering an accumulation of apoptosis-stimulating deoxyribonucleoside di- and triphosphates. (*R*)-DI-87 interferes with this pathway by inhibiting dCK, thus preventing host cell death. **(B)** Structure of (*R*)-DI-87.

676
677

Figure 2. (*R*)-DI-87 protects phagocytes from death-effector deoxyribonucleoside-mediated cytotoxicity. (A-D) Survival rates of human U937 monocyte-like cells (U937) (A, B) or U937-derived macrophages (U937 MΦ) (C, D) exposed to dAdo or dGuo in the presence (+) or absence (-) of 1 µM (*R*)-DI-87. Cells were also exposed to the inhibitor or vehicle only. U937 *dCK⁻/⁻* were included as a control. **(E-H)** Survival rates of human CD14⁺ monocytes (E, F) or human monocyte-derived macrophages (HMDMs) (G, H) exposed to dAdo or dGuo in the presence (+) or absence (-) of 1 µM (*R*)-DI-87. Cells were also exposed to the inhibitor or vehicle only. **(I-J)** Survival rates of U937 MΦ exposed to rAdsA-derived dAdo (I) or dGuo (J). rAdsA was incubated with dAMP or dGMP and reaction products containing dAdo or dGuo were used to treat phagocytes in the presence (+) or absence (-) of 1 µM (*R*)-DI-87. Controls lacked rAdsA or deoxyribonucleoside monophosphates, or included reaction buffer only as indicated with + and − symbols. 100 µM (A-B; E-F) or 200 µM (C-D; G-H) of dAdo or dGuo were used to treat the cells. Cell survival rates were analyzed 48 h post-treatment. Data are the mean (± standard deviation [SD]) values from at least three independent determinations. Primary cell experiments include at least three independent donors. Statistically significant differences were analyzed with one-way analysis of variance (ANOVA) and Tukey's multiple-comparison test; ns, not significant ($P \geq 0.05$); *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

693
694
695
696

697
698

Figure 3. Selective inhibition of dCK prevents death-effector deoxyribonucleoside-mediated induction of immune cell apoptosis. (A-D) Analysis of caspase-3 activity in human U937 monocyte-like cells (U937) (A, B) or U937-derived macrophages (U937 MΦ) (C, D) exposed to dAdo or dGuo in the presence (+) or absence (-) of 1 μM (*R*)-DI-87. Cells were also exposed to the inhibitor or vehicle only. U937 *dCK*[-/-] were included as a control. Caspase-3 activity was analyzed using a colorimetric assay. **(E-F)** Immunoblotting of lysates obtained from U937 (E) or U937 MΦ (F) exposed to dAdo or dGuo in the presence (+) or absence (-) of 1 μM (*R*)-DI-87. Controls are indicated (+/− symbols). A specific antibody was used that detects the cleaved (active) form of caspase-3 (α-CASP3). GAPDH was used as a loading control (α-GAPDH). Numbers to the right of blots indicate the migration of molecular weight markers in kilodaltons. **(G-H)** Analysis of (*R*)-DI-87-dependent prevention of host cell apoptosis via immunofluorescence microscopy. U937 MΦ were exposed to dAdo (G) or dGuo (H) in the presence or absence of 1 μM (*R*)-DI-87 and stained using FITC-annexin-V/PI. Controls are indicated. White bars depict a length of 100 μm. Representative images are shown. 100 μM (A-B; E) or 200 μM (C-D; F-H) of dAdo or dGuo were used to treat the cells. Apoptosis rates were analyzed 24 h post-treatment. Data are the mean (± standard deviation [SD]) values from three independent determinations. Statistically significant differences were analyzed with one-way analysis of variance (ANOVA) and Tukey's multiple-comparison test; ns, not significant ($P \geq 0.05$); *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Figure 4. (*R*)-DI-87 protects against *S. aureus* invasive disease. (A, B) Safety assessment of (*R*)-DI-87 in mice. Cohorts of female C57BL/6 mice were treated with (*R*)-DI-87 (75 mg/kg) or vehicle (40% Captisol) via oral gavage in 12-hour intervals for 16 days. On day 16, peripheral blood was collected and subjected to a FACS-based immuno-phenotyping approach (A). Panel B indicates the body weight gain during the course of treatment. **(C-F)** Enumeration of visible surface abscesses and staphylococcal loads in organs of *S. aureus*-challenged C57BL/6 mice treated with (*R*)-DI-87 (75 mg/kg) or vehicle (40% Captisol). Mice received (*R*)-DI-87 (75 mg/kg) or vehicle (40% Captisol) via oral gavage every 12 hours and were challenged with 10^7 CFU of wild-type *S. aureus* Newman (WT) or its Δ*adsA* mutant. Data for female C57BL/6 mice are displayed (*n* = 8). Bacterial burden was enumerated as $\log_{10}$ CFU per gram of tissue at 5 days post-infection. Horizontal blue bars represent the mean values of visible abscesses per organ (C-D) or indicate the mean CFU count in each cohort (E-F). **(G, H)** Microscopic images of H&E–stained liver (G) or renal tissues (H) obtained after necropsy of *S. aureus*-challenged C57BL/6 mice treated with (*R*)-DI-87 (75 mg/kg) or vehicle (40% Captisol). Mice received (*R*)-DI-87 (75 mg/kg) or vehicle (40% Captisol) via oral gavage every 12 hours and were challenged with 10^7 CFU of wild-type *S. aureus* Newman (WT) or its Δ*adsA* mutant. Arrows point to immune cell infiltrates (black) or replicating staphylococci (blue). Black bars depict a length of 100 μm. Representative images are shown. Statistically significant differences were analyzed by a two-tailed Student's t-test (A-B) or with the Kruskal–Wallis test corrected with Dunn's multiple comparison (C-F). ns, not significant (*P* > 0.05); *, *P* < 0.05; **, *P* < 0.01; ***, *P* < 0.001.

Figure 5. (*R*)-DI-87-mediated inhibition of dCK enhances macrophage infiltration into staphylococcal abscesses. (A-P) Immunofluorescence microscopy-based detection of macrophages in liver or renal tissues isolated 5 days after intravenous injection of 10^7 CFU of wild-type *S. aureus* Newman (A-H) or its *adsA* mutant (I-P) into wild type C57BL/6 mice treated with (*R*)-DI-87 (75 mg/kg) or vehicle (40% Captisol). Mice received (*R*)-DI-87 (75 mg/kg) or vehicle (40% Captisol) via oral gavage every 12 hours. Magnifications of boxed areas from upper panels are indicated. Thin sections were stained with α-F4/80 antibodies (macrophages; red). Nuclei were labeled with DAPI (blue). White bars shown in the upper panels depict 100 µm length. Representative images are shown. **(Q, R)** Survival rates of murine bone marrow-derived macrophages (BMDMs) exposed to dAdo (Q) or dGuo (R) in the presence (+) or absence (-) of 1 µM (*R*)-DI-87. Cells were also exposed to the inhibitor or vehicle only. 200 µM of dAdo or dGuo were used to treat the cells. Cell survival rates were analyzed 48 h post-treatment. Data are the mean (± standard deviation [SD]) values from three independent determinations. Statistically significant differences were analyzed with one-way analysis of variance (ANOVA) and Tukey's multiple-comparison test; ns, not significant ($P \geq 0.05$); *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Figure 6. Proposed model of (*R*)-DI-87-mediated protection of phagocytes during *S. aureus* abscess formation. Diagram illustrating the (*R*)-DI-87-mediated protection of macrophages during the development of staphylococcal abscesses. While phagocytes get killed by *S. aureus*-derived death-effector deoxyribonucleosides in vehicle-treated animals, (*R*)-DI-87 protects macrophages and boosts their infiltration into the deeper cavity of infectious foci thereby enhancing eradication of staphylococci.