Digging deeper into pain – an ethological behavior assay correlating well-being in mice with human pain experience.

Luke A. Pattison [1*], Alexander Cloake [1], Sampurna Chakrabarti [1, 2], Helen Hilton [1], Rebecca H. Rickman [1], James P. Higham [1], Michelle Y. Meng [1], Luke W. Paine [1], Lanhui Qiu [1], Anne Ritoux [1], David C. Bulmer [1], Gerard Callejo [1, 3] and Ewan St. John Smith [1*]

From 1) Department of Pharmacology, University of Cambridge, Cambridge, United Kingdom; 2) Currently at: Max Delbrück Center for Molecular Medicine in the Helmholtz Association (MDC), Berlin, Germany; 3) Currently at: Department of Biomedicine, Medical School, Institute of Neurosciences, Universitat de Barcelona, Barcelona, Spain

*To whom correspondence should be addressed: Dr Luke A. Pattison: lap48@cam.ac.uk; Prof. Ewan St. John Smith: es336@cam.ac.uk

Keywords: Behavior; Ethology; Nociception; Inflammation; Arthritis; Colitis

Abstract (249 words)

The pressing need for safer, more efficacious analgesics is felt worldwide. Pre-clinical tests in animal models of painful conditions represent one of the earliest checkpoints novel therapeutics must negotiate before consideration for human use. Traditionally, the pain status of laboratory animals has been inferred from evoked nociceptive assays which measure their responses to noxious stimuli. The disconnect between how pain is tested in laboratory animals and how it is experienced by humans may in part explain the shortcomings of current pain medications and highlights a need for refinement. Here, we survey human chronic pain patients who assert that everyday aspects of life, such as cleaning and leaving the house, are affected by their on-going level of pain. Accordingly, we test the impact of painful conditions on an ethological behavior of mice, digging. Stable digging behavior was observed over time in naïve mice of both sexes. By contrast, deficits in digging were seen following acute knee inflammation. In the monosodium iodoacetate knee osteoarthritis model, the decrease in digging observed was not fully ameliorated by a clinically used drug, gabapentin, which reflected the opinion of human patients regarding its effectiveness. Lastly, in a visceral pain model, the decrease in digging behavior correlated with the extent of disease. Ultimately, we make a case for adopting ethological assays, such as digging, in studies of pain in laboratory animals, which we believe to be more representative of the human experience of pain and thus valuable in assessing clinical potential of novel analgesics in animals.

1. Introduction (500 words)

Chronic pain, resulting from inflammation, tissue damage or injury, is a global problem, with patients experiencing reduced quality of life ranging from social isolation, physical incapacity, and mental health disorders [6,26,35]. The issue is exacerbated by the lack of safe and effective medications to provide pain relief. Therefore, development of novel analgesics, particularly those targeted to disease-specific pain mechanisms, has been identified as a priority [31,43]. However, drug discovery efforts for novel analgesics are plagued by a high attrition rate, particularly during transition from pre-clinical testing, using animal (mostly rodent) models, to clinical trials [22]. The disconnect between how pain is assessed in non-verbal rodents and experienced by humans may offer some explanation for this [27,40]. Indeed, several clinical trials of putative painkillers have failed, despite showing promise in preclinical studies [44].

Historically pain in rodents has been inferred from nociception: the neural encoding of noxious stimuli. During such tests, animals are subjected to noxious stimulation, commonly heat or mechanical pressure. The latency to a response (e.g., withdrawal or vocalization), or frequency of nocifensive behaviors (flinching, licking, etc.) are then used to quantify nociception [29]. Animals experiencing pain due to an experimental disease model (e.g., osteoarthritis) are expected to exhibit quicker or more robust reactions. The analgesic potential of new compounds thus relies on animals being able to tolerate noxious stimulation for longer. However, such evoked approaches are not representative of how humans experience, or measure, their pain. Furthermore, these assays are dependent on being able to probe sites affected by pain, limiting their usefulness for certain conditions [25].

Chronic pain patients, living with various conditions, often report an inability to engage with everyday aspects of human life [6,15], which makes a compelling case for studying how ethological, animal behaviors are affected by pain models. Indeed, non-reflexive behavioral assays, such as facial grimacing [28], dynamic weight bearing [37], nest building [20], burrowing [42], automated home-cage analytics [18], and machine learning approaches to classify spontaneous behavior [4] are gaining traction as methods to track pain progression and assess analgesic intervention. Not only do these assessments correlate better with how pain affects humans, animals benefit from behaving freely during testing (often in a home-cage environment), and thus such approaches also represent a substantial refinement of the use of animals in pain research.

Another ethological behavior that can be used to assess animal welfare is digging. Distinct from burrowing, which relies on animals being provided with a substrate-filled apparatus, or marbles [9], digging captures the inquisitive nature of mice as they explore an open, uniform environment comprising of compressed digging substrate [11]. Compared to burrowing, digging is quicker to test and less task-driven, which removes a confounding influence on an animal's motivation to dig. We have already shown digging deficits in acute inflammatory and joint degeneration pain models that correlate with neuronal hyperexcitability [1,7,8], and here sought to further investigate the utility of digging as a readout of animal welfare by focusing on the robustness of naïve animals' behavior over extended periods of time, as well as the impact of different more translationally relevant pain models.

2. Materials and Methods

2.1. The Impact of Chronic Pain on Daily Life survey Ethical approval for human research was granted by The University of Cambridge Psychology Research Ethics Committee (PRE.2023.012). Participants were recruited via social media and with the help of numerous charities based in English-speaking countries with a focus on helping those living with chronic pain conditions (see Acknowledgements) over a period of three months. All participants consented to the use of the anonymous data collected for subsequent analyses and dissemination. Participants were surveyed about their diagnoses of chronic pain, use of analgesics and the impact of their pain on daily activities, some demographic information was also collected on an optional basis. A copy of the survey is included in Appendix B. Where respondents provided optional additional information about medication regimes and non-pharmacological therapies, responses were assigned to categories determined during analysis. Due to international differences in the legality and accessibility of cannabis and cannabidiol (CBD)-containing products as pharmacological treatments for pain, the response of participants that reported sole use of cannabis/CBD to the question about pharmacological interventions was amended to 'Other'. Use of cannabis/CBD by respondents who also satisfied the criteria to be counted as using prescription, over-the-counter, or both types of drug is still accounted for in the breakdown of medication classes.

2.2. Animals C57BL/6J mice (Envigo) were housed in groups of up to 5 per cage with *ad libitum* access to food and water. Holding rooms were maintained at 21 °C and operated 12-hour light/dark cycles. Mice of both sexes were used for experiments when aged 10-12 weeks (specific compositions of experimental cohorts are defined in corresponding figure legends). All animal work was regulated in accordance with the United Kingdom Animal (Scientific Procedures) Act 1986 Amendment Regulations 2012 and was approved by the University of Cambridge Animal Welfare Ethical Review Body. Behavioral studies commenced after at least a week of habituation in the animal facility. Animals were acclimatized in behavioral testing rooms for at least 30 minutes each day of testing. The room used, time of testing and experimenters present were kept constant for each individual study.

2.3. Complete Freund's adjuvant model of inflammatory joint pain Following induction of anesthesia (ketamine, 100 mg/kg and xylazine, 10 mg/kg, i.p.) and complete loss of consciousness, intra-articular injections were made through the patella tendon using a Hamilton syringe and 30G needle. 10 μl complete Freund's adjuvant (CFA, 10 mg/ml, Chondrex) was injected unilaterally (ipsilateral knee randomly determined). Knee joint width was measured with digital Vernier calipers before and after injections at specified time points.

2.4. Monosodium iodoacetate model of osteoarthritis Intra-articular injections of 10 μl monosodium iodoacetate (MIA, 0.1 mg/μl, Sigma-Aldrich) were made under inhalation anesthesia (isoflurane: 4% induction, 2% maintenance) unilaterally to a randomly determined knee. Mice were monitored daily for weight loss and knee inflammation. 10 days after injections, following behavioral assessments, mice were injected with gabapentin (30 mg/kg, i.p., Merck) and behaviors re-tested 90 minutes later.

2.5. Dextran sulfate sodium model of colitis After capturing baseline behaviors and weights, mice were provided with drinking water containing 3% or 1.5% (w/v) dextran sulphate sodium (DSS, Alfa Aesar) for 5 days, after which normal drinking water was provided again

for a further 2 days. A control cohort maintained on normal drinking water was run in parallel. Mice were monitored daily: a disease activity index was scored for each animal based on weight loss (0, none; 1, 1-5%; 2, 5-10%; 3, 10-15%, 4, >15%), stool consistency (0, normal; 2, loose stool; 4, watery diarrhea) and presence of blood in stool (0, none; 2, slight bleeding; 4, gross hemoccult). Daily water consumption was also monitored.

2.6. Digging behavioral paradigm   Following acclimatization to the behavior room and experimenters, mice were individually transferred to testing cages (49 x 10 x 12 cm) set up with ~4 cm of tightly packed fine-grain aspen midi 8/20 wood chip bedding (LBS Biotechnology), up to five cages were tested simultaneously. Mice were allowed 3-minutes to explore testing cages without interference, under video surveillance (iPhone 6S camera, Apple). Training sessions were carried out the day before baseline behaviors were assessed, to allow mice to gain familiarity with the set up. Training consisted of mice being placed in test cages as per a standard test. On the rare occasion that an animal did not dig, the individual was placed into a test cage with another animal (from the same home cage) that did meet this criterion, and a second training dig was conducted after a 30-minute break. Separate cages were used to test female and male mice, although when both sexes were studied, both were present in the behavior room for the duration of testing. Following testing sessions, the number of visible burrows at the end of the three-minute test was recorded. Where the same test cage was used multiple times on the same day, half of the digging substrate was refreshed between testing individual mice. Digging behavior was further analyzed offline via video playback at the conclusion of each study, following blinding of all videos. The latency of mice to begin digging was noted and the total time mice spent digging was scored independently by at least two observers; since scores were well correlated (Average R^2 = 0.7886 ± 0.03 from an average of 26 ± 4 videos scored per experimenter pairing) an average is reported as the time spent digging. We define digging behavior as the vigorous disturbance of digging substrate with all four limbs, which usually begins with a defined planting of the hind legs in a wide stance, followed by rapid movement of substrate under the body using first the fore limbs, and concluding by kicking of the substrate backwards using the hind limbs to produce a burrow.

2.7. Rotarod   A rotarod apparatus (Ugo Basile 7650) was used to assess the motor coordination of mice. Animals were allowed 1 minute to acclimatize to the apparatus on a slow setting (7 revolutions per minute, rpm), before an accelerating program (7-40 rpm over 5-minutes) was started. Mice were first placed on the rotarod after being trained in the digging assay, to gain some familiarity with the apparatus. Tests were video-recorded, mice were removed from the rotarod if they fell, following two consecutive passive rotations or after 6-minutes of the accelerating program, whichever occurred first. Analysis of videos was performed at the conclusion of studies, following blinding, the latency to passive rotation or fall was scored by a single experimenter.

2.8. Colon histology   After the final behavior assessments of the DSS study, mice were humanely euthanized by cervical dislocation. A segment of colon was resected ~1cm from the anus and fixed in 4% (w/v) paraformaldehyde (Sigma-Aldrich) for 6-hours before overnight cryoprotection in 30% (w/v) sucrose (Thermo Fisher Scientific). The following day tissue was embedded in Shandon M-1 Embedding Matrix (Thermo Fisher Scientific) and snap frozen in liquid nitrogen. 20 μm sections of colon were collected on a SuperFrost Plus slides (Thermo Fisher Scientific) using a Leica CM3000 cryostat. Sections were stained with alcian blue (1% w/v in 3% v/v acetic acid; Alfa Aesar; goblet cell stain), hematoxylin (0.2% w/v; Sigma Aldrich;

nuclei stain), and eosin (0.5% w/v; Acros Organics; counterstain). Slides were mounted with glycerol and imaged using a NanoZoomer S360 (Hamamatsu). Following blinding, images were scored based on histopathology features: inflammation (0, none; 1, increased immune nuclei in lamina propria; 2, confluence of immune nuclei extending to submucosa; 3, transmural extend of infiltrate), crypt damage (0, crypts intact; 1, loss of basal 1/3 crypt; 2, loss of basal 2/3 crypt; 3, entire crypt loss; 4, epithelial surface erosion; 5, confluent erosion) and ulceration (0, none; 1, 1-2 ulcerative foci; 2, 3-4 ulcerative foci; 3, extensive ulceration). At least two sections, separated by 150 µm were scored per colon, sections were scored independently by 3 experimenters, an average of their scores is reported.

2.9. Data analysis and statistics   The number of animals and sex composition of studies are detailed in each corresponding figure legend. Data are presented as mean ± standard error of the mean. Appropriate analyses were selected according to the number of factors being compared and whether data met the assumptions for parametric analyses, the statistical tests employed are stated in corresponding figure legends. All analyses were performed in R, and p values less than 0.05 were considered significant.

3. Results

3.1. Chronic pain prevents full engagement with daily life.

To assess the relevance of using ethological behaviors as a means of assaying nociception in laboratory animals, it was important to first confirm that natural human behaviors are affected in those experiencing chronic pain. To address this, we ran an online survey (Appendix B). Over a three-month recruitment period, 341 participants completed the survey. An optional demographic section (completed by 339 respondents) revealed that the majority (83.2%) of responders identified as female (Fig. S1A), the ages of respondents were relatively uniformly distributed (Fig. S1B), the overwhelming majority self-identified as White (85.8%; Fig. S1C) and most responses came from those living in United Kingdom (53.3%), New Zealand (25.3%) and Australia (11.3%; Fig. S1D). Respondents all confirmed they experience chronic pain (regular bouts of pain persisting at least 3 months) due to a medically diagnosed health condition. The underlying health conditions varied, with the largest numbers experiencing chronic pain due to: fibromyalgia, osteoarthritis, migraine and irritable bowel syndrome (IBS, Figure 1A; full diagnosis breakdown available in Table S1); 55.1% of respondents had diagnoses of more than one condition causing them chronic pain (Fig. 1B).

Respondents were asked about the frequency and intensity of their pain, with constant, low intensity pain being most commonly reported (27.0%), followed by low intensity pain on a daily basis (19.9%; Fig. 1C). In addition to their usual experience of pain, 97.9% of patients expressed that they also experience flare ups of more intense pain (Fig. 1D). The frequency of flare ups varied from daily to less frequent than monthly (Fig. 1E). Participants were also asked about their use of drugs to manage pain, with 65.7% of respondents using prescription medication, a further 21.4% relying on over-the-counter analgesics, 4.1% using a combination of both prescription medication and over-the-counter drugs and 1.5% of respondents using cannabis/cannabidiol-containing products only (classified as 'Other'). 7.3% of respondents reported not using any pharmacological intervention for their pain (Fig. 1F). Of the 258 participants that shared details on their use of drugs, the majority (68.6%) use multiple drugs to try and manage their pain (Fig. S1E). A breakdown of medication classes revealed that weak opioids (used by 31.0% of participants that shared details on their drug use) and paracetamol (28.3%) were the most commonly used (Fig. S1F), with one third (33.3%) taking some form of antidepressant (Fig. S1F). Of the 308 respondents using drugs to treat pain, only 5.5% found them to be completely effective at relieving pain, with most (82.4%) finding them somewhat effective and 12.1% not achieving any pain relief with their medications (Fig. 1G). Additionally, 57.5% of respondents reported experiencing unpleasant side effects from taking medication (Fig 1H). Emphasizing the desire of chronic pain patients for effective therapies, 74.2% of patients also make use of non-pharmacological treatments to treat their pain (Fig. 1I), with 62.8% of those using multiple alternative therapies (Fig. S1G). Analysis of the types of non-pharmacological therapies used by participants placed physiotherapy, massage, thermal, and exercise as the most popular non-pharmacological treatments used (Fig. S1H).

Questions probing the affect of pain on daily life, confirmed that pain has a mostly negative impact on many varied aspects of daily life, often taken for granted by those not living in chronic pain. For instance, 63.9% of respondents agreed that tending to their personal hygiene could cause them or increase their pain (Fig. 1J). An overwhelming majority of participants agreed that pain has prevented them from undertaking housework, engaging with physical activity, socializing and leaving the house. Whether pain interfered with non-physical recreational activities (e.g., reading) divided respondents the most, with 56.6% agreeing that pain can prevent them and 29.9% disagreeing. Regarding sleep, 87.7% of respondents reported that their sleep quality is impaired by their pain and 91.8% participants agreed that their pain contributed to low mood. Perhaps most importantly was the finding that 94.1% of participants agreed that they are unable to live life as they would if they did not experience

pain. Overall, these data support the notion that pain reduces an individual's ability and motivation to engage in everyday activities and that it would make sense to examine how equivalent behaviors in animals are affected by models of pain and pharmacological intervention.

Figure 1. Experiences with pain have a negative impact on the daily lives of human patients living with chronic pain. Summary of results from The Impact of Chronic Pain on

Daily Life survey, completed by 341 participants. **(A)** Number of diagnoses of conditions that caused respondents' chronic pain. **(B)** Frequency distribution of the number of diagnoses of chronic pain conditions by respondent. **(C)** Frequency and intensity of the pain experienced by respondents on usual basis. **(D)** Proportion of respondents experiencing flare ups of pain, in addition to their usual experience with pain. **(E)** Frequency distribution of the frequency of additional flare ups. **(F)** Use of medications to manage pain, OTC = over-the-counter. **(G)** Effectiveness of drugs to manage pain. **(H)** Occurrence of side effects from pain medications. **(I)** Use of non-pharmacological therapies to manage pain. **(J)** Opinion scales on how experience with pain affects various aspects of daily life.

3.2. Digging is a robust measurable ethological behavior

To better understand the stability of digging behavior, naïve mice of both sexes were studied over a period of 10 days, a common timeline for many rodent pain models (Fig. 2A). The latency to dig, time mice spent digging and number of burrows produced during each test period were measured. Additionally, on each test day a random subset of mice was re-tested after a 30-minute break to assess the consistency of digging behaviors. The time taken for mice to begin digging after starting testing remained relatively consistent across the study ($F_{(5,45)}$ = 1.649, p = 0.167; Fig. 2B), with an overall average latency to dig of 63 ± 5.5 s. Latencies remained similar across test digs when repeated after a 30-minute break (Test 1, 60.7 ± 17.4 s vs. Test 2, 52.4 ± 15.7 s, t = 0.678, df = 9, p = 0.515; Fig. 2C). Similarly, the time mice spent digging did not differ with each time point tested ($F_{(5, 45)}$ = 2.333, p = 0.058; Fig. 2D), with mice spending an average of 11.9 ± 1.0 s digging each day. A slight increase in the time mice spent digging was observed during the second time tested, although the difference was not significant (Test 1, 9.67 ± 2.5 s vs. Test 2, 14.5 ± 3.3 s, t = -2.164, df = 9, p = 0.058; Fig. 2E). The cohort of naïve mice produced an average of 3.98 ± 0.1 burrows, regardless of when tested during the study ($\chi^2(5)$ = 5.86, p = 0.320; Fig. 2F). The number of burrows produced between repeat tests on the same day remained consistent as well (Test 1, 4.4 ± 0.2 vs. Test 2, 3.9 ± 0.2, W = -25, p = 0.180; Fig. 2G). In the small cohort tested, the average time male mice spent digging was slightly higher than that recorded for females, although no significant difference was seen (Time digging: $F_{(1, 56)}$ = 3.866, p = 0.054; Fig. 2D), importantly, digging durations remained consistent over time when scrutinized by sex. In summary, the digging behavior of naïve mice is stable over timeframes commonly used in behavioral studies assessing pain.

Figure 2. Digging is a robust and measurable ethological behavior. (A) Timeline of behavioral experiments. **(B)** Average latency to dig across experimental timeline. **(C)** Latencies to dig of individual mice when tested twice in one day, 30-minute break between tests. **(D)** Average time spent digging across experimental timeline. **(E)** Digging durations of individual mice when tested twice in one day, 30-minute break between tests. **(F)** Average number of burrows dug by mice at each test point. **(G)** Number of burrows dug by individual animals when tested twice in one day, 30-minute break between tests. **(B, D)** Repeated measures ANOVA. **(C, E)** Paired t-test. **(F)** Friedman repeated measures ANOVA. **(G)** Wilcoxon signed-rank test. n = 8, 4 females (denoted by open symbols), 4 males (denoted by closed symbols).

3.3 Acute inflammation of the knee joint decrease digging behavior

Injection of complete Freund's adjuvant (CFA) into the knee joint of mice is a commonly used model of acute inflammatory pain [17]. To confirm that digging behaviors can be used as a readout of inflammatory pain, mice were injected unilaterally through the patellar tendon with either CFA, or saline serving as a control for intra-articular injection (Fig. 3A). 24-hours post-injection, mice that received CFA exhibited joint inflammation, evident from an increased ratio in the diameter of the ipsilateral joint relative to the contralateral joint (Baseline, 0.99 ± 0.01 vs. 24-hours, 1.29 ± 0.03, t = -9.240, df = 9, $p < 0.0001$; Fig. 3B). No swelling was observed for mice injected with saline (Baseline, 1.00 ± 0.00 vs. 24-hours, 1.01 ± 0.02, t = -0.778, df = 5, $p = 0.472$; Fig. 3B). Consistent with the absence of any swelling, the latency to dig was not affected by intra-articular injection of saline (Baseline, 17.5 ± 7.8 s vs. 24-hours, 12.7 ± 3.9 s, t = 0.520, df = 5, $p = 0.625$; Fig. 3C), whereas mice injected with CFA took longer to start digging 24-hours post-injection (Baseline, 33.1 ± 8.2 s vs. 24-hours, 68.2 ± 14.5 s, t = -2.857, df = 9, $p = 0.019$; Fig 3C). The time mice spent digging and the number of burrows dug during the test period were also unaffected by saline injection (Duration: Baseline, 34.8 ± 8.3 s vs. 24-hours, 32.7 ± 5.8 s, t = 0.348, df = 5, $p = 0.742$; Fig. 3D; Burrows: Baseline, 4.1 ± 0.5 vs. 24-hours, 4.0 ± 0.4, W = -2, $p = 0.773$; Fig. 3E). However, injection of CFA reduced both the amount of time mice spent digging (Baseline, 30.6 ± 5.2 s vs. 24-hours, 7.7 ± 2.0 s, t = 5.061,

df = 9, *p* = 0.0007; Fig. 3D) and the number of burrowing sites recorded (Baseline, 4.7 ± 0.4 vs. 24-hours, 2.3 ± 0.5, W = -55, *p* = 0.002; Fig. 3E). Importantly, no adverse effect was seen when mice were tested on the rotarod following CFA-induced inflammation (Baseline, 296.6 ± 34.8 s vs. 24-hours, 354.0 ± 6 s, t = -1.614, df = 4, *p* = 0.182; Fig. S2), indicating that mice retain normal locomotor function. These results thus suggest that the inflammation resulting from intra-articular injection of CFA reduces the motivation of mice to dig, while they likely remain able to. Possible explanations for the observed digging deficit may arise from mice finding digging painful following induction of inflammation, or, the occurrence of spontaneous pain, resulting from the inflammatory insult, triggering an apathetic withdrawal.

Figure 3. Acute inflammation of the knee joint decreases digging behavior. (A) Schematic representation of the complete Freund's adjuvant (CFA) knee model (created with BioRender.com) and experimental timeline. **(B)** Ratio of ipsilateral knee width to contralateral knee width on the day of injections and 24-hours after. **(C)** Latency of mice to dig. **(D)** Time spent digging before and 24-hours after CFA injection. **(E)** Number of visible burrows at the conclusion of 3-minutes digging test. ∗ p < 0.05, ∗∗ p < 0.01, ∗∗∗ p < 0.001, ∗∗∗∗ p < 0.0001: **(B – D)** Paired t-test. **(E)** Wilcoxon signed-rank test. n = Saline: 6, 4 females (denoted by open symbols), 2 males (denoted by closed symbols); CFA: 10, 5 females (denoted by open symbols), 5 males (denoted by closed symbols).

3.4 Digging deficits are also seen in a more translationally relevant model of osteoarthritis

Having confirmed that the digging activity of mice is reduced following acute inflammation, the suitability of digging as a welfare readout in a more chronic model was assessed. Injection of monosodium iodoacetate (MIA) into the knee joint of mice results in a transient inflammatory response and progressive joint degeneration, recapitulating many of the aspects of osteoarthritis [32]. Following assessment of baseline behaviors, a cohort of 8 mice were injected with MIA unilaterally and studied for 10 days (Fig. 4A). Swelling of the injected knee, relative to the contralateral joint, was observed following injections, peaking 24-hours post-injection, but then steadily declining throughout the study (F (10, 70) = 52.656, *p* < 0.0001;

Fig. 4B). In line with the progressive degeneration of the injected joint following injection of MIA, the latency of mice to dig was observed to increase over time (F (4, 28) = 14.471, $p < 0.0001$; Fig. 4C). The time spent digging declined with time following injections (F_{GG} (1.55, 10.87) = 17.929, $p = 0.0006$; Fig. 4D), as did the number of burrows dug during the test period ($\chi^2(4) = 16.5$, $p = 0.0025$; Fig. 4E). However, the ability of mice to use their joints when required was not affected by the MIA model, as is evident from the lack of an effect on the time the mice could spend on the rotarod (F (4, 28) = 1.891, $p = 0.14$; Fig. S3). Considered together, these results are indicative of mice experiencing reduced motivation to explore and to interact with the environment with the onset of arthritis-like symptoms.

Gabapentinoids, such as gabapentin and pregabalin, are increasingly being prescribed to human osteoarthritis patients, to alleviate the pain associated with their condition [2]. In addition, administration of gabapentin to rodents following progressive degeneration of the knee joint can remedy mechanical hypersensitivity [4,41]. Although an improvement in certain aspects of spontaneous behavior can be seen, comprehensive 3D pose analysis has shown that gabapentin does not return the spontaneous behavior of mice to their pre-injury state [4]. Such reports question the analgesic properties of gabapentin, when focusing on metrics more relevant to human pain experience. The effect of gabapentin on the digging deficits induced by the MIA model of osteoarthritis was thus tested. Following behavior testing on day 10, mice were injected with gabapentin and behaviors re-assessed 1.5-hours later (Fig. 4A). Digging duration, the most severely affected digging parameter at day 10, was found to increase slightly following administration of gabapentin, but remained statistically distinct to the baseline activity of mice (Baseline, 8.5 ± 0.9 s vs. Post-gabapentin, 4.6 ± 1.7 s, t = 3.44, df = 7, $p.adj = 0.033$; Fig. 4F) and no difference was observed when compared to the digging duration before administration of gabapentin (Pre-gabapentin, 2.8 ± 1.1 s vs. Post-gabapentin, 4.6 ± 1.7 s, t = -1.40, df = 7, $p.adj = 0.612$; Fig. 4F). Gabapentin is known to be a sedative [13], which may help explain these findings, indeed, mice struggled more on the rotarod test 1.5-hours after receiving gabapentin (Pre-gabapentin, 313.6 ± 18.6 s vs. post-gabapentin, 201.6 ± 19.3 s, t = 3.47, df = 7, $p.adj = 0.031$; Fig. 4G). These findings are consistent with the opinions of the 50 participants of our survey, who reported use of gabapentinoids to manage their pain (10 participants reported only using gabapentinoids; Fig. 4H). An overwhelming majority of those using gabapentinoids regularly (80%) felt that their medication regime was only partially effective at reducing their pain, with 16.0% reporting a complete lack of effectiveness (Fig. 4Ii). Additionally, 61.2% of gabapentinoid users stated that they experience unpleasant side-effects from their use of pain medications (Fig. 4Iii. The questionable analgesic properties of gabapentin in the digging assay reported here, corroborates the lack of effectiveness reported by human patients who use similar medications, evident from our own survey (Figs 4 I) and the findings of others [12]. Considering gabapentin often reverses evoked pain assay deficits, our data only strengthen the notion for the use of ethological approaches in preclinical tests of novel analgesics.

Figure 4. Digging behavior is decreased in the MIA model of osteoarthritis. (A) Schematic representation of the MIA model (created with BioRender.com) and experimental timeline. **(B)** Ratio of ipsilateral knee width to contralateral knee width. **(C)** Average latency to dig across experimental timeline. **(D)** Average time digging across experimental timeline. **(E)** Average number of burrows dug across experimental timeline. Effect of gabapentin administration on **(F)** time digging and **(G)** rotarod performance. **(H)** Proportion of respondents from Impact of Chronic Pain on Daily Life survey using gabapentinoids (GBP = gabapentin; PGB = pregabalin) to manage pain. **(I)** Opinion of survey respondents that use gabapentinoids on the **(i)** effectiveness of and **(ii)** experience of side-effects from their pain medications. ∗ p < 0.05, ∗∗ p < 0.01, ∗∗∗ p < 0.001, ∗∗∗∗ p < 0.0001: **(B - D)** Repeated measures ANOVA **(D)** with Greenhouse-Geisser sphericity corrections. **(E)** Friedman repeated measures ANOVA. **(F - G)** Repeated measures ANOVA followed by Bonferroni corrected post-hoc. n = 8 males.

3.5 Digging behavior is predictive of the extent of intestinal inflammation in a visceral pain model

Painful conditions affecting visceral organs are highly prevalent but are difficult to study in both human patients and animal models due to their internal nature. A behavioral readout that offers

insight to animal wellbeing in relation to the extent of disease, would thus likely be highly useful. To this end, the effect of an experimental model of ulcerative colitis, a form of inflammatory bowel disease, on digging behaviors was tested. Addition of dextran sulfate sodium (DSS) to the drinking water of rodents triggers colitis-like intestinal inflammation through disruption of the intestinal epithelia [33]. An acute DSS-colitis model, where mice are provided with DSS-containing drinking water for 5 days, before returning to normal water for a further 2 days was used to assess the suitability of digging behavior to infer animal wellbeing in visceral pain (Fig. 5A). The extent of colitis symptoms correlates with the concentration of DSS provided [14], we thus elected to study two doses (3% w/v and 1.5% w/v) with the aim of interrogating whether digging parameters can distinguish between disease severity. A control cohort kept on water for the duration of the study was run in parallel. Upon starting the model, mice were monitored daily to estimate the extent of disease progression, producing a disease activity index (DAI). As expected, a significant difference in the extent of disease activity was seen dependent on the dose of DSS that mice were administered ($F_{(2, 228)} = 181.144$, $p < 0.0001$; Fig. 5B). In line with the work of other groups using the same model [16], a sex-dependent effect was also observed ($F_{(1, 228)} = 32.186$, $p < 0.0001$; Fig. 5B), which was most noticeable for the 1.5% w/v dose of DSS (DAI at Day 7: Females, 2.4 ± 0.3 vs. Males, 8.0 ± 0.3, t = -14.00, df = 4, $p.adj$ = 0.0002). The DAI data suggest that males are more susceptible to DSS colitis, however, this is not due to higher consumption of DSS-containing water, as the average daily water consumption of animals over the first five days of the experiment was consistent across groups and sexes, when corrected to animal body weight (Interaction: Sex:Group $F_{(2, 24)} = 0.03$, $p = 0.971$; Fig. S4A).

In line with the progressive nature of the model, a trend for an increased latency to dig was seen for mice that received DSS-containing water, with the latency of mice on the 3% w/v dose increasing 64% from day 0 to day 7. Although the effect of dose on latency was not found to be significant ($F_{(2, 120)} = 1.609$, $p = 0.204$; Fig. 5C), minimal change (-22%) was seen in the latency of mice kept on regular drinking when comparing day 7 to day 0, suggesting that mice on the highest dose of DSS began to lose in motivation to dig by the end of the study. In agreement with this, a decrease in the time mice spent digging was observed over experimental time ($F_{(4, 120)} = 6.503$, $p < 0.0001$; Fig. 5D), which was affected by both DSS dose ($F_{(2, 120)} = 4.483$, $p = 0.013$; Fig. 5D) and sex ($F_{(1, 120)} = 8.186$, $p = 0.005$; Fig. 5D). These effects were most noticeable at the conclusion of the study, where mice that received the high dose of DSS dug substantially less than the control cohort for both sexes (Female: 0% DSS, 7.6 ± 0.7 s vs. 3% DSS, 2.3 ± 0.6 s, t = 5.43, df = 4, $p.adj$ = 0.017; Male: 0% DSS, 6.6 ± 0.5 s vs. 3% DSS, 2.0 ± 0.8 s, t = 6.14, df = 4, $p.adj$ = 0.011; Fig. 5E). However, only males were seemingly affected when comparing mice that received 1.5% w/v DSS to the control group (Female: 0% DSS, 7.6 ± 0.7 s vs. 1.5% DSS, 8.8 ± 2.2 s, t = 3.22, df = 4, $p.adj$ = 0.097; Male: 0% DSS, 6.6 ± 0.5 s vs. 1.5% DSS, 1.9 ± 0.5 s, t = 6.05, df = 4, $p.adj$ = 0.011; Fig. 5E). Similarly, the number of burrows dug decreased over time ($F_{(4, 120)} = 14.942$, $p < 0.001$; Fig. 5F) and was found to be dependent on the dose of DSS that mice received ($F_{(2, 120)} = 6.416$, $p = 0.002$; Fig. 5F) and sex ($F_{(1, 120)} = 5.461$, $p = 0.021$; Fig 5F). To confirm that the observed deficits in digging behavior were not the result of mice being physically incapacitated, mice were also assessed on the rotarod after each test of digging. No deficit in locomotor function was seen during the study ($F_{(4, 143)} = 1.543$, $p = 0.193$; Fig. S4B).

Disease activity indexes are useful for monitoring animal welfare during experimental colitis, however histopathological assessment of the affected tissue offers a more conclusive assessment of disease extent. After behavioral assessments on day 7, mice were sacrificed, and colons extracted. In line with previous reports [16,33] a reduced colon length was observed for mice that drank DSS-containing water ($F_{(2,18)} = 15.947$, $p = 0.000104$; Fig. S4C). Sections of the distal colon were prepared and stained (Fig. 5G), and histological scores were assigned. The dose of DSS that animals received had a significant influence over the resulting histological score (F = 23.415, $p < 0.0001$; Fig. 5H) and, in line with the DAI and time spent digging, a sex effect was also seen (F = 7.976, $p = 0.0105$; Fig. 5H). Having confirmed

the extent of intestinal inflammation on Day 7 for each mouse, histological scores were next compared to the digging activity (time spent digging) observed on the final day of the experiment, offering the most comprehensive assessment of whether digging activity is predictive of disease severity. A negative correlation was observed (β = -0.7793, p < 0.0001; R^2 = 0.645, F (1, 22) = 39.97, p < 0.0001; Fig. 5I). Although, DAI is less conclusive, the relationship between DAI and amount of time mice spent digging was also explored, also showing a negative correlation (β = -0.7222, p < 0.0001; R^2 = 0.377, F (1, 51) = 30.86, p < 0.0001; Fig. S4D). The findings described here thus strengthen the notion of digging as a valuable read out of disease severity.

Figure 5. Digging behavior is predictive of the extent of intestinal inflammation in a visceral pain model. (A) Schematic representation of the DSS model (created with BioRender.com) and experimental timeline. **(B)** Disease activity index (DAI) over experimental time. **(C)** Average latency to dig across experimental timeline. **(D)** Average time digging across experimental timeline. **(E)** Summary of digging durations of mice as tested on Day 7. **(F)** Average number of burrows dug across experimental timeline. **(G)** Representative staining of colon sections from male mice that received **(i)** water only, **(ii)** 1.5% w/v DSS, **(iii)** 3% w/v DSS. Scale bars: Full cross-sections = 400 µm; inserts = 200 µm. **(H)** Histological scores of

colon sections according to animal sex and treatment group. **(I)** Correlation between histological score and digging duration of mice as tested on Day 7. $*$ $p < 0.05$, $**$ $p < 0.01$, $****$ $p < 0.0001$: **(B – F)** Three-way ANOVA **(E)** followed by Bonferroni corrected post-hoc. **(H)** Two-way ANOVA followed by Bonferroni corrected post-hoc. **(I)** Simple linear regression. **(B - F)** All groups: n = 10, 5 females (denoted by open symbols) and 5 males (denoted by closed symbols). **(G - H)** 0% group: n = 4 females, 3 males; 1.5% group: n = 5 females, 4 males; 3% group: n = 5 females, 3 males.

4. Discussion (1376 words)

Animal models of painful conditions are an incredibly useful aspect of the drug discovery process, not just for pre-clinical testing of potential new pain medications, but also in the identification of putative targets. With the ever pressing need to deliver safer and more efficacious analgesics, ensuring that data obtained from animal models of painful conditions are as reliable and informative as possible is a top priority. Here we add to the mounting evidence that chronic pain patients are prevented from going about their everyday lives by the pain they experience (Fig. 1). We also show that an analogous everyday behavior of mice, digging, is negatively affected by several models of painful conditions (Figs 3 - 5). We thus believe there is strong justification to include digging as an ethological behavioral readout of animal wellbeing in the field of pain research. We believe that the addition of digging to behavioral characterization, will complement data obtained via other evoked and non-evoked assays, to generate a more comprehensive pain profile, which will aid in smoothing the transition from treating pain in animals to helping human patients.

Burrowing is the main alternative ethological rodent behavior to digging. Undeniably useful as a measure of animal wellbeing, deficits in burrowing behavior have been reported in inflammatory, visceral, neuropathic and post-surgical pain models [10,19,21,30,34,42]. However, we believe that digging [11] provides a better ethological readout than burrowing. Our approach to studying digging is reminiscent of the open field test, whereby mice are placed in a completely uniform environment, which they can freely explore: naturally inquisitive, this often involves digging. Experimental set ups for assaying burrowing involve providing animals with apparatus to burrow from, and can thus be considered task based, which raises questions over the motivation of animals to perform the behavior, especially as in many cases burrows are filled with food pellets. This is evident from observations that mice will spend more time exploring a designated burrow (plastic tube) than in the open when a burrow is available [36]. Motivation has a large sway over any animals' behavior, but the digging assay removes the confounder of object exploration from the motivation to perform the measured behavior, but still allows motivation to be inferred. This is evident from our observation that mice retain full locomotive ability when placed on the rotarod, and as such are given no choice but to move, in all the pain models we explored. However, when tested in our digging paradigm, we report a reduction in behavior across all models tested. We therefore assume that mice also retain the ability to dig, but on some level make a conscious decision not to, and thus believe that digging represents a better way to capture both the motivation and ability of mice to engage in a routine behavior.

Visceral pain conditions are among the worst managed by currently approved medications, evident from findings that anti-inflammatories often exacerbate intestinal inflammation [5,24] and opioid medication producing constipation as a common side effect. Assessment of visceral pain in lab animals is also limited, which may contribute to the lack of effective pharmaceutical treatments. Currently, visceromotor responses to colorectal distention, measured by electromyography, afford the most specificity in assessing visceral pain in rodents. However, experiments often involve invasive surgeries, evoked stimulation, anesthesia and restraint [23], a stark contrast to how visceral pain manifests, and is assessed, in humans. Here we have shown a negative effect on the digging activity of mice in an experimental model of colitis (Fig. 5). Our findings that digging deficits were graded in line with the dose of DSS animals received, and a correlation between digging activity on the final day of behavioral testing and histopathological analyses of disease extent suggest that digging behavior is predictive of internal inflammatory state and thus holds high value in assessing animal wellbeing in DSS colitis. This linkage of internal inflammatory state to digging behavior provides further support for digging analysis to be widely used in the behavioral arsenal of pain scientists, particularly those researching visceral conditions. It will be interesting to see how digging behaviors are affected by other models of visceral conditions associated with pain. Bashir and colleagues [3] have recently shown that mice undergoing a model of

endometriosis are slower to burrow, from a tube filled with food pellets, than sham animals, however, both groups successfully clear the burrow within 24-hours. Such findings align with our argument that the burrowing assay is more task driven than digging. Studying how the digging behavior of mice is affected by this endometriosis model may allow for a more objective measure of animal welfare in the absence of an overt challenge.

Here we only report minimal improvement of digging behavior in the MIA model following administration of gabapentin (Fig. 4F). However, we have previously demonstrated that analgesia, evident from an increase in digging activity, can be obtained in the acute CFA-induced knee inflammation model using a blocker of the transient receptor potential vanilloid 1 (TRPV1) ion channel and chemogenetic approaches [7,8]. We have also observed improved digging in the more chronic destabilization of the medial meniscus model of osteoarthritis following intra-articular mesenchymal stem cell derived extracellular vesicle (MSC-EV) administration; MSC-EVs also reducing neuronal hyperexcitability [1]. Thus, digging behavior also holds value in the testing of analgesics, in addition to tracking welfare through pain models. Furthermore, we have shown that the ethological digging assay may lend itself better to assess analgesia than traditional evoked nociception assays. This is demonstrated by two findings of this study: 1) that gabapentin, a drug clinically used to treat osteoarthritis pain, provides limited analgesia when measured in terms of its effect on digging behavior in an experimental model of osteoarthritis (Fig. 4F), and 2) that a resounding majority of human patients who take gabapentinoids to manage their pain are dissatisfied with the pain relief afforded by their medications (Fig. 4I). Of course, the utility of digging as a predictor of analgesic success could be further tested by expanding investigations to other analgesics already approved to treat conditions that can be modelled in mice and comparing how changes in digging align with patient opinion.

Here, we describe the use of digging as a behavioral read out of animal welfare in various pain models. The utility of digging has been demonstrated by the robustness of behavior in naïve mice (Fig. 2) and deficits observed following acute inflammation (Fig. 3) and more translationally relevant models of osteoarthritis (Fig. 4) and colitis (Fig. 5). Sufficient sensitivity of the assay also allowed capture of graded responses, which correlated with the extent of inflammation. It is important to note the parallels between the observed deficits in digging behavior seen in mouse models of painful conditions and the reduced motivation and/or ability of humans experiencing pain to undertake routine behaviors, as well as the limitations of gabapentin on improving arthritis-triggered digging deficits in mice and complaints regarding the effectiveness of analgesia achieved when similar drugs are used by humans. Together, these demonstrate that digging is a suitable behavior for assessing pain and analgesia in mice. Indeed, the similarities in digging deficits and the human experience of pain potentially permit greater insight into the analgesic potential of novel drugs by measuring their ability to improve the likelihood of engaging in natural behavior. Furthermore, digging behavior can be assessed with minimal investment in equipment or personnel training, animals can be tested in batches and analyses performed offline after blinding. The nature of data collection is also highly amenable to automated analysis with constant developments in tracking algorithms, representing an extra level of possible improvement. Although we have previously shown that regulating articular neuron activity improves digging behavior in acute [7] and chronic [1] joint pain models, we cannot say with certainty that observed deficits in digging behavior are a result of animals experiencing pain. However, we have also argued that our approach allows a more objective assessment of the motivation to spontaneously perform an ethological behavior than alternatives. In this way, we believe that digging adeptly captures a critical aspect of the current IASP definition of pain [38], owing to its high reliance on emotional experience. This is different to many other behavioral measures of pain, which have a higher degree of affective reaction to passive overt or evoked stimuli, as opposed to digging being driven, mostly, by free will. That said, we are advocates for using multiple approaches to profile pain, agreeing with the recent recommendation of Sadler, Mogil and Stucky [39] to increase

diversity and resolution of pain behavior measurements, believing that digging has certainly proved its use to become more commonplace.

Acknowledgements

The authors thank the following charities for their assistance in promoting and recruiting pain patients to 'The Impact of Chronic Pain on Daily Life' survey: Arthritis Action, Arthritis Australia, Arthritis New Zealand, Arthritis Society Canada, Bowel Research UK, Crohn's and Colitis Canada, Crohn's and Colitis New Zealand, Crohn's and Colitis UK, Endometriosis New Zealand, Endometriosis UK, Fibromyalgia Action UK and National Arthritis Foundation Singapore. The authors also thank the staff of the University of Cambridge Combined Animal Facility for their work in supporting this research. For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

LAP and ESS acknowledge funding from the MRC (MR/W002426/1). AMC, GC and ESS were funded by Versus Arthritis (RG21973). GC and ESS were funded by the Rosetrees Trust (CM818). SC was supported by a Gates Cambridge Trust scholarship. HH was supported by a BBSRC/GSK iCASE PhD studentship (BB/V509528/1). RHR, MM, LWP and AR were supported by AstraZeneca PhD Studentships (RHR: G104108; MM: G104109; LWP: G113502; AR: G115018). JPH and DCB acknowledge funding from Crohn's and Colitis UK (PC2019/1-Bulmer).

Author contributions: conceptualization: L. A. Pattison, S. Chakrabarti and E. St. J. Smith; investigation: L. A. Pattison, A. Cloake, S. Chakrabarti, H. Hilton, R. H. Rickman , J. P. Higham, M. Y. Meng, L. W. Paine and G. Callejo; data analysis: L. A. Pattison, A. Cloake, S. Chakrabarti, H. Hilton, R. H. Rickman, J. P. Higham, M. Y. Meng, L. W. Paine, L. Qui and A. Rittoux; writing: L. A. Pattison and E. St. J. Smith; supervision: D. C. Bulmer and E. St. J. Smith; funding acquisition: G. Callejo, D. C. Bulmer and E. St. J. Smith.

Conflict of interest statement

The authors have no conflicts of interest to declare.

Availability of data

Behavioral data sets supporting the conclusions of this article are available in University of Cambridge Apollo Repository (https://doi.org/10.17863/CAM.100117). Summary data from The Impact of Chronic Pain on Daily Life survey will be shared on reasonable request from the corresponding authors.

References

[1] Ai M, Hotham WE, Pattison LA, Ma Q, Henson FMD, Smith ESJ. Role of Human Mesenchymal Stem Cells and Derived Extracellular Vesicles in Reducing Sensory Neuron Hyperexcitability and Pain Behaviors in Murine Osteoarthritis. Arthritis Rheumatol 2023;75:352–363.

[2] Appleyard T, Ashworth J, Bedson J, Yu D, Peat G. Trends in gabapentinoid prescribing in patients with osteoarthritis: a United Kingdom national cohort study in primary care. Osteoarthritis Cartilage 2019;27:1437–1444.

[3] Bashir ST, Redden CR, Raj K, Arcanjo RB, Stasiak S, Li Q, Steelman AJ, Nowak RA. Endometriosis leads to central nervous system-wide glial activation in a mouse model of endometriosis. J Neuroinflammation 2023;20:59–18.

[4] Bohic M, Pattison LA, Jhumka ZA, Rossi H, Thackray JK, Ricci M, Mossazghi N, Foster W, Ogundare S, Twomey CR, Hilton H, Arnold J, Tischfield MA, Yttri EA, St John Smith E, Abdus-Saboor I, Abraira VE. Mapping the neuroethological signatures of pain, analgesia, and recovery in mice. Neuron 2023.

[5] Bonner GF. Exacerbation of inflammatory bowel disease associated with use of celecoxib. Am. J. Gastroenterol. 2001;96:1306–1308.

[6] Breivik H, Collett B, Ventafridda V, Cohen R, Gallacher D. Survey of chronic pain in Europe: prevalence, impact on daily life, and treatment. Eur J Pain 2006;10:287–333.

[7] Chakrabarti S, Pattison LA, Doleschall B, Rickman RH, Blake H, Callejo G, Heppenstall PA, Smith ESJ. Intra-articular AAV-PHP.S mediated chemogenetic targeting of knee-innervating dorsal root ganglion neurons alleviates inflammatory pain in mice. Arthritis Rheumatol 2020:art.41314.

[8] Chakrabarti S, Pattison LA, Singhal K, Hockley JRF, Callejo G, Smith ESJ. Acute inflammation sensitizes knee-innervating sensory neurons and decreases mouse digging behavior in a TRPV1-dependent manner. Neuropharmacology 2018;143:49–62.

[9] Deacon RMJ. Burrowing in rodents: a sensitive method for detecting behavioral dysfunction. Nat Protoc 2006;1:118–121.

[10] Deacon RMJ. Burrowing: a sensitive behavioural assay, tested in five species of laboratory rodents. Behav. Brain Res. 2009;200:128–133.

[11] Deacon RMJ. Digging and marble burying in mice: simple methods for in vivo identification of biological impacts. Nat Protoc 2006;1:122–124.

[12] Derry S, Bell RF, Straube S, Wiffen PJ, Aldington D, Moore RA. Pregabalin for neuropathic pain in adults. Cochrane Database Syst Rev 2019;1:CD007076.

[13] Doleman B, Heinink TP, Read DJ, Faleiro RJ, Lund JN, Williams JP. A systematic review and meta-regression analysis of prophylactic gabapentin for postoperative pain. Anaesthesia 2015;70:1186–1204.

[14]   Egger B, Bajaj-Elliott M, MacDonald TT, Inglin R, Eysselein VE, Büchler MW.
        Characterisation of acute murine dextran sodium sulphate colitis: cytokine profile
        and dose dependency. Digestion 2000;62:240–248.

[15]   Fagan G, Osborne H, Schultz M. Physical Activity in Patients with Inflammatory
        Bowel Disease: A Cross-Sectional Study. Inflamm Intest Dis 2021;6:61–69.

[16]   Francis-Malavé AM, Martínez González S, Pichardo C, Wilson TD, Rivera-García
        LG, Brinster LR, Carrasquillo Y. Sex differences in pain-related behaviors and
        clinical progression of disease in mouse models of colonic pain. PAIN
        2023;164:197–215.

[17]   Gauldie SD, McQueen DS, Clarke CJ, Chessell IP. A robust model of adjuvant-
        induced chronic unilateral arthritis in two mouse strains. J. Neurosci. Methods
        2004;139:281–291.

[18]   Hasriadi, Wasana PWD, Vajragupta O, Rojsitthisak P, Towiwat P. Automated home-
        cage for the evaluation of innate non-reflexive pain behaviors in a mouse model of
        inflammatory pain. Sci Rep 2021;11:12240–16.

[19]   Jirkof P, Cesarovic N, Rettich A, Nicholls F, Seifert B, Arras M. Burrowing behavior
        as an indicator of post-laparotomy pain in mice. Front Behav Neurosci 2010;4:165.

[20]   Jirkof P, Fleischmann T, Cesarovic N, Rettich A, Vogel J, Arras M. Assessment of
        postsurgical distress and pain in laboratory mice by nest complexity scoring. Lab.
        Anim. 2013;47:153–161.

[21]   Jirkof P, Leucht K, Cesarovic N, Caj M, Nicholls F, Rogler G, Arras M, Hausmann M.
        Burrowing is a sensitive behavioural assay for monitoring general wellbeing during
        dextran sulfate sodium colitis in laboratory mice. Lab. Anim. 2013;47:274–283.

[22]   Kissin I. The development of new analgesics over the past 50 years: a lack of real
        breakthrough drugs. Anesth Analg 2010;110:780–789.

[23]   Larauche M, Gourcerol G, Million M, Adelson DW, Taché Y. Repeated psychological
        stress-induced alterations of visceral sensitivity and colonic motor functions in mice:
        influence of surgery and postoperative single housing on visceromotor responses.
        Stress 2010;13:343–354.

[24]   Lichtenstein GR, Feagan BG, Cohen RD, Salzberg BA, Diamond RH, Price S,
        Langholff W, Londhe A, Sandborn WJ. Serious infection and mortality in patients
        with Crohn's disease: more than 5 years of follow-up in the TREAT™ registry. Am.
        J. Gastroenterol. 2012;107:1409–1422.

[25]   López-Estévez S, Gris G, la Puente de B, Carceller A, Martínez V. Intestinal
        inflammation-associated hypersensitivity is attenuated in a DSS model of colitis in
        Sigma-1 knockout C57BL/6 mice. Biomed Pharmacother 2021;143:112126.

[26]   Luque-Suarez A, Martinez-Calderon J, Falla D. Role of kinesiophobia on pain,
        disability and quality of life in people suffering from chronic musculoskeletal pain: a
        systematic review. Br J Sports Med 2019;53:554–559.

[27]   Mao J. Current challenges in translational pain research. Trends Pharmacol. Sci.
        2012;33:568–573.

[28]   Matsumiya LC, Sorge RE, Sotocinal SG, Tabaka JM, Wieskopf JS, Zaloum A, King OD, Mogil JS. Using the Mouse Grimace Scale to reevaluate the efficacy of postoperative analgesics in laboratory mice. J Am Assoc Lab Anim Sci 2012;51:42–49.

[29]   Mogil JS. The history of pain measurement in humans and animals. Front Pain Res (Lausanne) 2022;3:1031058.

[30]   Muralidharan A, Kuo A, Jacob M, Lourdesamy JS, Carvalho LMSPD, Nicholson JR, Corradini L, Smith MT. Comparison of Burrowing and Stimuli-Evoked Pain Behaviors as End-Points in Rat Models of Inflammatory Pain and Peripheral Neuropathic Pain. Front Behav Neurosci 2016;10:88.

[31]   Nawai A, Leveille SG, Shmerling RH, van der Leeuw G, Bean JF. Pain severity and pharmacologic pain management among community-living older adults: the MOBILIZE Boston study. Aging Clin Exp Res 2017;29:1139–1147.

[32]   Ogbonna AC, Clark AK, Gentry C, Hobbs C, Malcangio M. Pain-like behaviour and spinal changes in the monosodium iodoacetate model of osteoarthritis in C57Bl/6 mice. Eur J Pain 2013;17:514–526.

[33]   Okayasu I, Hatakeyama S, Yamada M, Ohkusa T, Inagaki Y, Nakaya R. A novel method in the induction of reliable experimental acute and chronic ulcerative colitis in mice. Gastroenterology 1990;98:694–702.

[34]   Palazzo E, Romano R, Luongo L, Boccella S, De Gregorio D, Giordano ME, Rossi F, Marabese I, Scafuro MA, de Novellis V, Maione S. MMPIP, an mGluR7-selective negative allosteric modulator, alleviates pain and normalizes affective and cognitive behavior in neuropathic mice. PAIN 2015;156:1060–1073.

[35]   Pitcher MH, Korff Von M, Bushnell MC, Porter L. Prevalence and Profile of High-Impact Chronic Pain in the United States. J Pain 2019;20:146–160.

[36]   Pond HL, Heller AT, Gural BM, McKissick OP, Wilkinson MK, Manzini MC. Digging behavior discrimination test to probe burrowing and exploratory digging in male and female mice. J Neurosci Res 2021;99:2046–2058.

[37]   Quadros AU, Pinto LG, Fonseca MM, Kusuda R, Cunha FQ, Cunha TM. Dynamic weight bearing is an efficient and predictable method for evaluation of arthritic nociception and its pathophysiological mechanisms in mice. Sci Rep 2015;5:14648–11.

[38]   Raja SN, Carr DB, Cohen M, Finnerup NB, Flor H, Gibson S, Keefe FJ, Mogil JS, Ringkamp M, Sluka KA, Song X-J, Stevens B, Sullivan MD, Tutelman PR, Ushida T, Vader K. The revised International Association for the Study of Pain definition of pain: concepts, challenges, and compromises. PAIN 2020;161:1976–1982.

[39]   Sadler KE, Mogil JS, Stucky CL. Innovations and advances in modelling and measuring pain in animals. Nat. Rev. Neurosci. 2022;23:70–85.

[40]   Vierck CJ, Hansson PT, Yezierski RP. Clinical and pre-clinical pain assessment: are we measuring the same thing? PAIN 2008;135:7–10.

[41]   Vonsy JL, Ghandehari J, Dickenson AH. Differential analgesic effects of morphine and gabapentin on behavioural measures of pain and disability in a model of osteoarthritis pain in rats. Eur J Pain 2009;13:786–793.

[42]   Wodarski R, Delaney A, Ultenius C, Morland R, Andrews N, Baastrup C, Bryden LA, Caspani O, Christoph T, Gardiner NJ, Huang W, Kennedy JD, Koyama S, Li D, Ligocki M, Lindsten A, Machin I, Pekcec A, Robens A, Rotariu SM, VoB S, Segerdahl M, Stenfors C, Svensson CI, Treede R-D, Uto K, Yamamoto K, Rutten K, Rice ASC. Cross-centre replication of suppressed burrowing behaviour as an ethologically relevant pain outcome measure in the rat. PAIN 2016;157:2350–2365.

[43]   Yekkirala AS, Roberson DP, Bean BP, Woolf CJ. Breaking barriers to novel analgesic drug development. Nat Rev Drug Discov 2017;16:545–564.

[44]   Yezierski RP, Hansson P. Inflammatory and Neuropathic Pain From Bench to Bedside: What Went Wrong? J Pain 2018;19:571–588.