Lineage-based scaling of germline intercellular bridges during oogenesis

Umayr Shaikh[1], Kathleen Sherlock[1], Julia Wilson[1], William Gilliland[2], and Lindsay

Lewellyn[1,3]

[1]Department of Biological Sciences, Butler University, Indianapolis, IN

[2]Department of Biological Sciences, DePaul University, Chicago, IL

[3]Author for correspondence (llewelly@butler.edu)

Key words: oogenesis, *Drosophila,* intercellular bridge, ring canal

<u>Summary Statement</u>

Using the fruit fly egg chamber as a model, this study demonstrates that the size

and scaling of germline intercellular bridges vary based on lineage.

<u>Abstract</u>

The size of subcellular structures must be tightly controlled to maintain normal

cell function. Despite its importance, few studies have determined how the size of

organelles or other structures is maintained during development, when cells are

growing, dividing, and rearranging. The developing egg chamber is a powerful model in

which to study the relative growth rates of subcellular structures. The egg chamber

contains a cluster of sixteen germline cells, which are connected through intercellular

bridges called ring canals. As the egg chamber grows, the germline cells and the ring

canals that connect them increase in size. Here, we demonstrate that ring canal size

scaling is related to lineage; the largest, "first born" ring canals increase in size at a

relatively slower rate than ring canals derived from subsequent mitotic divisions. This

lineage-based scaling relationship is maintained even if directed transport is reduced,

ring canal size is altered, or in egg chambers with twice as many germline cells.

Analysis of lines that produce larger or smaller mature eggs reveals different strategies

could be used to alter final egg size.

<u>Introduction</u>

A fundamental question in biology is how size is regulated. There is an extensive

literature describing the relationship between body size and the size of organs or

appendages (Gayon, 2000; Huxley and Teissier, 1936), but we are just beginning to

understand how size is regulated at the cellular level. Cells vary in size from one

organism to another and from one cell type to another, but an essential feature of all

cells is that they must regulate the size and number of their subcellular structures. Many

studies have described the scaling of various organelles with cell size, including the

nucleus, the mitotic spindle, mitochondria, the ER, and others (Marshall, 2020), but very

little is known about how subcellular scaling is established and maintained within

developing tissues, where cells are growing, dividing, and rearranging.

The developing fruit fly egg chamber is a powerful model system to study scaling

of subcellular structures. The egg chamber is the functional unit of oogenesis which will

ultimately produce the mature egg; within the egg chamber, the developing oocyte and

its supporting nurse cells are surrounded by a layer of epithelial cells. The oocyte is

connected to the nurse cells through intercellular bridges, called ring canals, which

provide a channel for the delivery of essential materials from the nurse cells to the

oocyte. From stages 2-14, the egg chamber increases in volume by approximately

1,000-fold (Fig. 1A; Ali-Murthy et al., 2021); therefore, the egg chamber provides us with

the opportunity to learn how the size of subcellular structures is coordinated during

tissue growth.

One of the most well-established examples of subcellular scaling is that of the

nucleus. Cells typically maintain a constant ratio of nuclear to cytoplasmic volume,

which appears to be actively maintained in healthy cells. When cells divide, or if cell size

is artificially manipulated, the nucleus will adjust its size to re-establish the proper ratio

(Balachandra et al., 2022; Cantwell and Nurse, 2019). Within the developing egg

chamber, the growth of the nurse cells, their nuclei, and nucleoli is similar, which leads

to isometric scaling of these structures (Diegmiller et al., 2021; Imran Alsous et al.,

2017). Similar isometric scaling of nuclear, nucleolar, and cell size within the developing

C. elegans intestine has also been observed (Uppaluri et al., 2016). Nurse cell nuclear

growth is primarily driven by multiple rounds of endoreplication, and increased ploidy

correlates with increased cell size (Diegmiller et al., 2021; Doherty et al., 2021; Orr-

Weaver, 2015; Øvrebø and Edgar, 2018). Although all nurse cells and their nuclei grow

during oogenesis, it has recently been demonstrated that nurse cell size is not uniform

within the germline cell cluster; instead, there is a size hierarchy based on distance from

the oocyte. The four posterior nurse cells directly connected to the oocyte are the

largest, whereas the anterior most nurse cell that is four ring canals away from the

oocyte is the smallest (Diegmiller et al., 2021; Imran Alsous et al., 2017). Therefore,

within this system, we have the opportunity to monitor subcellular scaling in a growing

tissue with a cell size gradient.

The germline ring canals are essential structures whose size must be tightly

regulated to produce a viable mature egg. The germline cell cluster is formed through

four rounds of mitosis; at the end of each division, the midbody is modeled to form the

ring canal (Price et al., 2023). These mitotic divisions are synchronous across the

cluster, which gives rise to a stereotypical pattern of connections (Fig. 1E). After

completion of the four mitotic divisions, the ring canals begin to expand (Ong and Tan,

2010), ultimately reaching a final diameter of nearly 10 µm. Although it is known that

ring canal growth is required for fertility, the scaling of ring canal size during oogenesis

has not been determined.

Here, we characterize the scaling of two types of ring canals: the four ring canals

that connect the posterior most nurse cells to the oocyte and the anterior most ring

canal. We hypothesized that ring canal size and growth would correlate with that of the

associated nurse cells; if that were true, we would have expected to see an anterior-

posterior gradient in ring canal size and scaling. Instead, our data supported an

alternative hypothesis that ring canal scaling is most closely linked to lineage. The

largest, "first born" ring canals expand more slowly compared to smaller, later born ring

canals. This scaling pattern is maintained even if polarized transport is reduced, ring

canal size is altered, or in egg chambers with twice as many germline cells. Finally,

analysis of a set of artificially selected "big egg" and "small egg" lines revealed that the

lineage dependent pattern of scaling is maintained, but that subtle changes to ring canal
scaling could provide a mechanism to regulate final egg size.
95
<u>Materials and Methods</u>
Fly Stocks and Maintenance: Flies were maintained on a cornmeal molasses diet at
18-25°C. The "small egg" (9.31.4, 8.26.2, 8.11.1, 7.17.4, and 9.2.2) and "big egg"
(1.16.1, 2.15.4, 2.49.3, 1.40.2, and 3.34.1) *D. melanogaster* lines were obtained from C.
Miles (Augustana University)(Jha et al., 2015; Miles et al., 2011), the *otu-GAL4,UAS-*
Dcr-2 (recombinant of BDSC #58424 and BDSC #24648) was provided by J. Merkle
(University of Evansville), and the *UASp-ovhts:GFP* line was provided by L. Cooley
(Gerdes et al., 2020). The following lines were obtained from the Bloomington
Drosophila stock center: w^{1118} (BDSC# 3605), *UAS-dhc64C-RNAi* (BDSC #36583),
matαTub-GAL4 (BDSC #7063), maternal triple driver MTD-GAL4 (BDSC #31777), *nos-*
GAL4 (BDSC #32563), and *UAS-CG34200/tom6-RNAi* (BDSC #44463).
107
Immunofluorescence and Imaging: Female flies were maintained with ground yeast
in the presence of males for ~44-60 hours prior to dissection. Ovaries were dissected in
Schneider's media, fixed in 4% formaldehyde, and stained with an HtsRC antibody
(1:20, hts RC DSHB), phalloidin (1:500; either TRITC or FITC conjugated; ECM
Biosciences), and DAPI (1:500). Anti-mouse secondary antibodies were used at a 1:200
dilution (Jackson Immunolab). Tissue was imaged using a Nikon Ti2-E Inverted
microscope equipped with a Yokogawa CSU-X1 Spinning Disk and Hamamatsu ORCA
Fusion camera with a 20x PLAN APO/0.75 NA objective. Although we tried to be
consistent in scaling confocal images in the figures, there were differences in
fluorescence intensity within z-stacks, so scaling was adjusted to best highlight the
structures of interest in each image.
119
Mature Egg Collection and Volume Measurements: Male and female flies were
incubated with ground yeast for approximately 24 hours at 25°C prior to transferring
them to cages with apple juice plates and wet yeast. Mature eggs were collected and
imaged using a Zeiss LP-520 and ProgRes MF camera (Jenoptik) using ProgRes

Capture Pro Software (Jenoptik). Length and width measurements were made using Fiji

(Schindelin et al., 2012), and volumes were calculated using the formula: Volume =

$1/6*\pi*Width^2*Length$

Ring canal and Nuclear Measurements: Nuclear and ring canal measurements were

made using Fiji (Schindelin et al., 2012) with either the freehand drawing tool or the line

tool respectively. Measurements were performed in the plane that contained the

maximum cross section or diameter within the DAPI or HtsRC channel. For Fig. 3, egg

chambers with low levels of ovhts::GFP expression were excluded from the analysis.

Egg chamber volumes were calculated using the formula: Volume =

$1/6*\pi*Width^2*Length$

Statistical Analysis: Graphing and statistical analyses were performed using Prism

10.1.0 (GraphPad Software, LLC). Data are typically shown as individual data points or

as the mean +/- s.d. or the 95% confidence interval as indicated in legends. For linear

regression analysis, ANCOVA was used to determine whether there were significant

differences in the slopes or y-intercepts. One-way ANOVA with Tukey's multiple

comparisons test was used to compare mature egg size (Fig. 5A). The sample size for

each experiment (n) is indicated in the figure legends.

Results

Ring canal scaling is related to lineage

It has recently been shown that there is a size hierarchy within the germline cell

cluster; the largest nurse cell nuclei are located at the posterior, whereas the smallest

nucleus is within the anterior most nurse cell (Diegmiller et al., 2021; Imran Alsous et

al., 2017). Recent work in this system has demonstrated that nurse cell nuclear growth

is tightly coupled to cell growth (Diegmiller et al., 2021). Therefore, we can use nuclear

size as a proxy for nurse cell size.

We first wanted to determine whether there were differences in the growth rate of

nurse cell nuclei in different regions of the germline cell cluster. We focused our

analysis on two sets of nurse cells – the four posterior nurse cells that are directly

connected to the oocyte and the anterior most nurse cell (Fig. 1E). This should allow us

to compare the relative growth rates of the largest and smallest nurse cells. We found

that as the egg chamber increases in volume, the anterior and posterior nurse cell

nuclei also increase in size, but at slightly different rates (Fig. 1A-B). If we plot these

measurements on a log-log scale, then the slope of the regression line provides us with

the scaling exponent, or the way that the size of one structure is related to the size of

another structure. Although the intercept does not represent a biologically relevant

measurement (since you cannot measure the size of one structure when the size of the

other structure is 0), significant differences in the y-intercepts allow us to compare the

relative size of structures between different conditions or genetic backgrounds. Because

we are plotting the log of egg chamber volume and the log of nuclear area, isometric

nuclear scaling would be reflected by a slope of 2/3 or 0.666; a smaller slope would

indicate hypoallometric scaling, and a larger slope would indicate hyperallometric

scaling. We found that the posterior nuclei scale nearly isometrically with egg chamber

volume (Fig. 1B; scaling exponent of 0.652, 95% confidence interval is 0.627-0.678),

which is consistent with what has been previously shown (Diegmiller et al., 2021). The

nucleus in the anterior most nurse cell scales slightly hypoallometrically (Fig. 1B; scaling

exponent of 0.626, 95% confidence interval of 0.593-0.659), but this difference is not

significant ($p > 0.05$). The y-intercepts of these two lines are significantly different (Fig.

1B; $p < 0.0001$), which is consistent with the smaller size of the anterior cells (and their

nuclei) that has been previously reported (Diegmiller et al., 2021; Imran Alsous et al., 2017).

This suggests that in addition to absolute differences in nuclear size, there could be modest

differences in nurse cell (and nuclear) growth rates based on position within the germline

cell cluster.

If we hypothesize that ring canal growth correlates with nurse cell size and

growth, then we would predict that the posterior ring canals should be larger and

expand faster than the anterior ring canal. We measured the diameter of the ring canals

that connect the same nurse cells to either the oocyte (for the posterior ring canals) or

to another nurse cell (for the anterior ring canal). There was no difference in the scaling

exponent between these two ring canal types (Fig. 1C; 95% confidence interval of

0.218-0.241 for the posterior ring canals and 0.209-0.250 for the anterior ring canals),

but there was again a significant difference in the y-intercepts (Fig. 1C; p<0.0001),

which reflects the smaller size of the anterior ring canals. At first glance, these data

suggest ring canal size varies along the anterior-posterior axis, which correlates with

nuclear size, but that ring canal scaling is similar across the germline cell cluster.

However, what was obvious from our analysis was that within an egg chamber,

the size of the four posterior ring canals varied significantly (Fig. 1A,C). This variation in

size could be explained by the emergence of one ring canal at the end of each of the

four mitotic divisions that form the germline cyst. It has been shown that the ring canals

formed during the earlier mitotic divisions are larger than those that are formed during

the later divisions (Ong and Tan, 2010), at least within the germarium. Therefore, we

hypothesized that lineage could be the source of the posterior ring canal size

differences. If we separate out the posterior ring canals based on their mitotic division of

origin (Fig. 1E; M1 for the first mitotic division, M2 for the second mitotic division, etc.),

we found that the "first born" M1 ring canals were nearly always the largest, whereas

the "last born" M4 ring canals were typically the smallest (Fig. 1A,D). In addition to these

size differences, the posterior ring canals had different scaling exponents; the M1 ring

canals had a significantly lower scaling exponent compared to the M2, M3, and M4 ring

canals (M1 ring canals: 0.202, 95% confidence interval 0.188-0.216; M4 ring canals:

0.252, 95% confidence interval 0.234-0.271). Interestingly, the scaling exponent for the

anterior ring canals, which are also formed during the fourth mitotic division, was not

significantly different from that of the posterior M4 ring canals (Fig. 1D). This

observation suggests that ring canal size and growth are not linked to nurse cell size,

since the anterior nurse cells are the smallest and grow more slowly than the posterior

nurse cells. Instead, our data support the hypothesis that ring canals scale differently

based on their lineage. These differences could reflect a need for the initially smaller M4

ring canals to grow more quickly to try to "catch up" to the size of the larger M1 ring

canals.

Reducing directed transport reduces ring canal size and leads to a modest increase in

scaling

Ring canals serve a specific purpose within the egg chamber by providing a

channel for the directed transport of material from the nurse cells into the oocyte. Based

on the observed size differences, we hypothesized that the larger M1 ring canals may

allow more material to flow through them than the smaller M4 ring canals; this difference

in flow could provide a mechanism to drive faster growth of the M4 ring canals. If the

amount of directed transport through the ring canal inversely correlates with ring canal

growth, then reducing transport should increase growth. It has been shown that the

amount of biased transport of materials through the germline ring canals correlates with

their position along the anterior-posterior axis (Lu et al., 2021). Movement of materials

through the anterior ring canals is bidirectional, whereas the movement through the

posterior ring canals is significantly biased toward the oocyte. Depletion of either dynein

heavy chain (Dhc64C) or the linker protein Short stop (Shot) significantly reduces

biased transport at the posterior, in turn reducing oocyte size (Lu et al., 2021).

Therefore, if ring canal size scaling is sensitive to transport through the posterior ring

canals, then depletion of Dhc64C should alter the scaling exponents.

We first confirmed that depletion of Dhc64C led to a significant reduction in

oocyte size. To perform the depletion, we used *otu-GAL4,UAS-Dcr-2* which bypasses

the requirement for dynein during earlier stages of oogenesis (Lu et al., 2022; McGrail

and Hays, 1997). This phenotype was most obvious during mid-oogenesis (stage ~5-8),

so we focused our analysis on these egg chambers (Fig. 2A-B). We performed the

same analysis of ring canal diameter in the control and *dhc64C-RNAi* egg chambers.

We found that depletion of Dhc64C led to a significant decrease in ring canal diameter

(Fig. 2C,S1A; y-intercepts for both ring canal types were significantly different from

controls), and there was a modest, but not significant increase in ring canal scaling (Fig.

2C,S1A). A similar reduction in ring canal size was observed in Shot-depleted egg

chambers (Lu et al., 2021). If we separated the ring canals based on lineage, we again

found that depletion of Dhc64C led to a significant decrease in ring canal size and a

consistent, but not significant increase in the scaling exponent for all ring canal types

(Fig. 2D, S1B significant difference in y-intercepts for control/*dhc64C-RNAi* comparisons

for M2, M3, M4 and the anterior ring canals, $p < 0.05$). Despite these changes, the

pattern of lineage-based size scaling was maintained.

Increasing ring canal size does not affect the lineage-based patterns, but it does lead to
modest differences in scaling

      If ring canal size inversely correlates with scaling such that larger ring canals expand more slowly than smaller ring canals, then increasing ring canal diameter should reduce the scaling exponent. To test this hypothesis, we over-expressed a GFP-tagged form of ovhts, which is cleaved to produce HtsRC::GFP (Gerdes et al., 2020). As was previously reported (Gerdes et al., 2020), we observed a consistent increase in size for all ring canal types (Fig. 3A-F; y-intercepts were significantly different for M1, M2, M4, and anterior RCs, $p<0.0001$), but the lineage-based pattern of scaling exponents was still observed (Fig. 3B-F). Although the scaling exponents were consistently lower in egg chambers over-expressing HtsRC::GFP, these differences were only significant for the M3 ring canals (Fig. 3D, $p=0.008$). This suggests that increasing ring canal size is not sufficient to significantly reduce scaling for all ring canal types.

      Although the lineage-based scaling pattern in the HtsRC::GFP over-expressing egg chambers were similar to the control, our impression was that the differences in posterior ring canal size were not as large. Therefore, we calculated the ratio of the M1 ring canal diameter to the M4 ring canal diameter. In control egg chambers, this ratio is initially higher, but as the egg chamber increases in volume, the ratio decreases, approaching, but never reaching 1.0 (Fig. 3G). For the HtsRC::GFP over-expressing egg chambers, we still observed the same negative correlation, but the ratios were significantly lower than in controls (Fig. 3G; y-intercepts were significantly different, $p<0.0001$). This suggests that even though over-expression of HtsRC::GFP increases the size of all ring canals, the "last born" ring canals still grow faster and are sometimes even able to catch up to the size of the "first born" ring canals.

Lineage-based ring canal scaling is maintained in egg chambers with twice as many
germline cells

      Thus far, we have described previously unappreciated trends in the size scaling relationships of the germline ring canals based on lineage. Although size may contribute

to the growth differences, lineage seems to be the most consistent predictor of scaling.

To further test this relationship, we turned to a condition in which egg chambers contain

twice as many germline cells. As part of another project, it was found that when a

shRNA construct was used to induce a depletion of CG34200, or Tom6, using *nanos-*

GAL4, many egg chambers contained 31 nurse cells and ring canals, with a single

oocyte connected to five posterior nurse cells (Fig. 4A). This phenotype could occur if

the germline cells go through an additional (fifth) mitotic division; alternatively, if there is

a delay in abscission following the germline stem cell division, this could give rise to a 2-

cell cystoblast, which then goes through four mitotic divisions to produce the 32-cell cyst

(Mathieu et al., 2013; Matias et al., 2015). Additional work to further characterize

the Tom6 depletion phenotype is in progress (manuscript in preparation), but we could

use this condition to determine whether ring canal scaling changes in egg chambers

that contain additional germline cells and ring canals.

When Tom6 was depleted from the germline, we saw similar trends for ring canal

scaling. The ring canal scaling exponent increased from M1 to M5 (Fig. 4B-D). Although

the ring canals in *tom6-RNAi* egg chambers were typically larger, some ring canal types

had a significantly larger scaling exponent (Fig. 4D) compared to controls. Therefore,

although size could impact scaling, our data again support the model that lineage is the

strongest predictor of ring canal scaling.

Scaling differences were observed across a collection of "big" and "small" egg lines

When considering the importance of size scaling within the germline of the

developing egg chamber, we wondered whether it would be maintained in egg

chambers that develop into eggs of different sizes or whether differences in ring canal

size and/or scaling could contribute to differences in final egg volume. We chose to

monitor scaling within a collection of *D. melanogaster* lines that were artificially selected

to produce either "big" or "small" eggs (Jha et al., 2015; Miles et al., 2011). We first

confirmed that these lines still produce significantly larger or smaller mature eggs; all

the "big egg" lines produced significantly larger mature eggs than all the "small egg"

lines (Fig. 5A; $p < 0.0001$ ANOVA with Tukey's post-hoc). From this collection, we

performed further analysis on two of the "big egg" lines (1.40.2 and 3.34.1) and two of

the "small egg" lines (7.17.4 and 9.31.4).

In this set of four lines, we went on to monitor ring canal size. We hypothesized

that if lineage determines ring canal scaling, then we should see the same patterns

across the "big egg" and "small egg" lines, where the M1 ring canals have a lower

scaling exponent than the M4 ring canals. When comparing ring canal size and scaling

between the "big egg" and "small egg" lines, we considered different possibilities. For

example, if the M1 ring canals are larger in the "big egg" lines than the M1 ring canals in

the "small egg" lines, then the scaling could be similar between those lines, which would

support the model that lineage, and not ring canal size, most reliably determines

scaling. If ring canal size does impact growth, then in this scenario, the M1 ring canals

in the "big egg" lines might have a lower scaling exponent than the M1 ring canals in the

"small egg" lines. Alternatively, ring canal size may be similar across the "big egg" and

"small egg" lines, but differences in scaling could provide a mechanism to alter final egg

size.

After completing ring canal measurements, we found that the lineage-based

pattern of scaling exponents was consistent with what we had observed previously.

Within each line, the scaling exponents typically increased from M1 to M4 (Fig. 5B-F),

and there was no significant difference in the scaling exponent for the M4 ring canal

compared to the anterior ring canal (Fig. 5B). We were surprised to see that there was

not a consistent relationship between ring canal size and final egg volume. Although the

"big egg" lines did tend to contain larger ring canals, that was not always the case.

However, the line that produces the smallest mature eggs, 9.31.4, did consistently have

smaller ring canals than the other lines (Fig. 5A,C-F; y-intercept for 9.31.4 was

significantly different than "big egg" lines for all ring canal types; $p<0.0001$); this

difference was most obvious when comparing the size of the M4 ring canals (Fig. 5F).

Although the ring canals were smaller in the 9.31.4 line, the scaling exponents were not

significantly different than those of the two "big egg" lines (Fig. 5B-F). This suggests that

reducing ring canal size without altering scaling could provide a mechanism to reduce

final egg size.

The other "small egg" line, 7.17.4, showed a different behavior. This line

consistently had scaling exponents for the posterior ring canals that were significantly

smaller than the "big egg" lines (Fig. 5B-F); however, the anterior ring canal scaling was

not different (Fig. S2A). Unlike the other "small egg" line, the size of the posterior ring

canals in 7.17.4 was more similar to the size of those in the "big egg" lines, but the

lower scaling exponent suggests that they may be expanding at a slower rate. This line

also had a consistently lower nuclear scaling exponent for all nurse cell types (Fig.

S2B). This suggests that coordinated changes in ring canal and nurse cell (nuclear)

scaling could contribute to changes in final egg size. Additional work will be necessary

to determine whether there are other size and/or scaling differences across this

collection of "big egg" and "small egg" lines, which could provide insight into different

developmental strategies that could be used to alter final egg size.

Discussion

We still have much to learn about how cells within developing tissues coordinate

their size and the size and number of their subcellular structures, but here, we have

characterized size scaling of the germline ring canals within the developing egg

chamber. Our data support the model that birth order inversely correlates with ring canal

scaling; the larger, "first born" ring canals have a lower scaling exponent than the

smaller ring canals derived from later mitotic divisions (Fig. 1-4). Although some data

support a modest effect of directed transport (Fig. 2) or ring canal size (Fig. 3), lineage

consistently provided the strongest predictor of ring canal scaling. Analysis of artificially

selected lines that produce eggs of different sizes (Fig. 5) suggests that within *D.*

melanogaster, there could be multiple strategies used to alter final egg size.

Ring Canal Scaling

Ring canals are known to undergo significant growth during oogenesis, but to our

knowledge, this is the first time that their scaling has been described. It was not

surprising that the diameter of the ring canals would increase more slowly than other

structures within the egg chamber, such as the nurse cell nuclei or nucleoli (Diegmiller

et al., 2021). Although they are a common feature of developing germline cells across

the animal kingdom, most intercellular bridges are much smaller (typically less than 2

µm in diameter) than the germline ring canals within the egg chamber (Haglund et al.,

2011). A recent study reported that when they are formed, the ratio of intercellular

bridge diameter to cell diameter was ~5-15% (Singh et al., 2022), which suggests that

there might be a physical limit to the size of a stable intercellular bridge. One way to

overcome this limitation might be to alter ring canal structure. Most other intercellular

bridges contain a single "rim," but the germline ring canals within the egg chamber,

which undergo a nearly 20-fold increase in diameter, contain two (Haglund et al., 2011).

Perhaps this more complex structure enables more growth than other intercellular

bridges or stabilizes these larger channels.

Ring canals initially differ in size due to differences in the degree of contractile

ring constriction at the end of each mitosis, producing smaller ring canals with each

round of division (Ong 2010). It has been shown that once the mitotic divisions are

complete, the M1 ring canal is the first to initiate the growth phase (Ong 2010), and we

assumed that growth would be similar for all ring canal types. Instead, our data support

a lineage-based effect on ring canal scaling, with the ring canals from earlier mitotic

divisions increasing in size more slowly than the ring canals from later divisions (Fig. 1-

5). The scaling differences could reflect nurse cell size differences; if that were the case,

then the posterior M4 ring canal would scale differently than the anterior ring canal,

since the anterior cells (and their nuclei) are always smaller than the posterior cells (Fig.

1) (Diegmiller et al., 2021; Imran Alsous et al., 2017). The observation that the scaling

exponents for the M4 posterior ring canal and the anterior ring canal were not

significantly different (Fig. 1) argues against that model.

Another possibility is that there could be structural differences in the ring canals

that are formed during the different mitotic divisions. Early work suggests that the

germline cells do not double in size between each round of mitosis (Koch and King,

1966), so as the cluster develops, the dividing cells are smaller and smaller. In the *C.*

elegans embryo, the contractile ring contains structural memory of cell size; the

contractile rings in larger cells divide faster than those in smaller cells. This structural

memory has been proposed to be due to the incorporation of different numbers of

contractile units into rings of different sizes (Carvalho et al., 2009). Because the ring

canal develops from the remodeled midbody, and the midbodies become smaller with

each germline cell division (Price et al., 2023), it is possible that the structure of the

nascent intercellular bridge may retain some memory of the size of the dividing cell,

which may then impact its subsequent rate of growth.

Additional structural differences could also explain lineage-based growth

differences. For example, if the M1 contractile ring constricts less than contractile rings

formed in subsequent divisions, perhaps this larger intercellular bridge must be

reinforced by the recruitment of more actin filaments or stabilized by more cell-cell

junctions, which may impact growth. Extensive cell-cell contacts have been observed in

the developing germline cell cysts (Loyer et al., 2015; Riparbelli et al., 2022; Tilney et

al., 1996), but a comparison of their distribution within different regions of the cluster

has not been performed.

If ring canal size correlates with oocyte-directed transport, then lineage-based

ring canal scaling could be used to ultimately achieve similar rates of transport through

all ring canals. If this were the case, then we would expect to see ring canals of different

lineages approach a similar final size; the initially smaller M4 ring canals grow faster to

catch up to the size of the larger M1 ring canals. When we compared the ratio of the M1

diameter to the M4 diameter, we found that this ratio did approach, but never reached

1.0. When the ring canals were larger in the HtsRC::GFP over-expressing egg

chambers, this ratio was significantly lower (Fig. 3G), but we still observed a lineage-

based effect on scaling. It is possible that the ring canals do not adjust their growth rate

based on the size of other ring canals, or it could be that because our over-expression

begins around stage 2-3 of oogenesis, lineage-based structural differences have

already been established. If there is an upper limit or a stage-specific optimal size for

transport of materials into the oocyte, then perhaps the larger ring canals hit that upper

limit or reach the optimal size sooner, which could explain the reduced scaling

exponents when HtsRC is over-expressed (Fig. 3). Additional work will be necessary to

further explore the relative rate of transport through different types of ring canals and

whether that changes as the ring canals increase in size.

One final explanation we considered is that the growth of a posterior ring canal

could depend on how many other ring canals exist within that germline cell. For

example, the M1 ring canal that we were monitoring is connecting the oocyte to a nurse

cell that contains three additional ring canals (M2, M3, and M4 types), whereas the

posterior M4 ring canal is the only intercellular bridge for that nurse cell (Fig. 1E).

Therefore, the differences in scaling could be due to the presence or absence of

additional ring canals within that cell. One argument against that model is that in Tom6-

depleted egg chambers that contain twice as many germline cells, each of the posterior

nurse cells had an additional ring canal, but the scaling exponents were not lower than

what we observed in the controls (Fig. 4). Therefore, additional work is necessary to

determine the molecular mechanisms underlying these scaling differences.

Although lineage does appear to be the strongest predictor of ring canal scaling,

we cannot rule out a contribution of ring canal size to scaling. In egg chambers depleted

of Dhc64C, there was a reduction in ring canal size and increase in scaling (Fig. 2), and

in egg chambers over-expressing HtsRC::GFP, there was a consistent increase in ring

canal size and reduction in scaling (Fig. 3). However, the *tom6-RNAi* egg chambers

typically contained larger ring canals, but some ring canal types showed higher scaling

exponents (Fig. 4). Therefore, analysis of additional conditions will be necessary to

determine the impact of ring canal size on scaling.

The lineage-based pattern of ring canal scaling is observed in lines that produce eggs of

different sizes

Final egg size is likely determined by many genes that impact the size or growth

of multiple structures during oogenesis. One possibility is that when producing eggs of

different sizes, the size of all structures is similarly altered. For example, when

generating larger mature eggs, flies produce larger egg chambers that contain larger

ring canals that expand at a specific rate based on their lineage. For three of the lines

analyzed (1.40.2, 3.34.1, and 9.31.4), we found that there were no significant

differences in the lineage-based ring canal scaling exponents. In these egg chambers,

the line that produced the smallest mature eggs (9.31.4) tended to have the smallest

ring canals relative to egg chamber volume (Fig. 5C-F), which was especially obvious

for the M4 ring canals (Fig. 5F). Analysis of these lines, therefore, is consistent with this

developmental strategy.

Another approach could be to increase or decrease ring canal scaling, which
could then affect the transfer of materials into the oocyte, ultimately changing final egg
size. Whole genome sequencing of the entire collection of artificially selected "big egg"
and "small egg" lines identified a set of candidate genes that may contribute to egg size
evolution (Jha et al., 2015); that list included many proteins that localize to the germline
ring canals or have been implicated in ring canal size regulation, including
Cheerio/Filamin, Flapwing, Mbs, and Short stop (Jha et al., 2015). Cheerio/Filamin
localizes to germline ring canals and is required for formation of the actin-rich inner rim
(Robinson et al., 1997). Flapwing and Mbs have been shown to work together to inhibit
myosin activity during incomplete cytokinesis, and mutants form smaller ring canals
(Ong et al., 2010; Yamamoto et al., 2013). As previously discussed, Short stop is a
linker protein that promotes directed transport into the oocyte (Lu et al., 2021). One of
the small egg lines (7.17.4) had a lower scaling exponent for all posterior ring canals
(Fig. 5), which suggests that this could be a mechanism to alter final egg size.

Many studies have demonstrated the impact of environmental conditions on egg
size, but the molecular mechanisms that control this life history trait are still not known.
Some insight has come from a recent study that found that honey bees could reversibly
alter egg size in response to the perceived colony size. The larger eggs that were
produced in response to a smaller perceived colony size expressed higher levels of the
GTPase, Rho1, and other related cytoskeletal proteins; depletion of Rho1 reduced
honey bee final egg size (Han et al., 2022). In *Drosophila,* Rho family members have
been implicated in regulating germline actin structures, including the ring canals
(Murphy and Montell, 1996). Additional studies will be necessary to confirm the
connection between ring canal scaling and final egg size and to further explore
pathways that may be involved in establishing or adjusting ring canal scaling. Our study
focused on ring canal growth prior to nurse cell dumping, so it remains to be determined
how this stage of oogenesis proceeds in the "big egg" and "small egg" lines and whether
there could be differences in the duration of the dumping phase and/or the degree of
completion of material transfer between the different lines. In addition, because our
study looked at a narrow range of final egg sizes, future work could expand to include
analysis of oogenesis in insects that produce much larger and much smaller eggs.

Figure 1: Ring canal size and nuclear size vary within the egg chamber. (A) Single plane confocal images of stage 5 (I), early stage 9 (II), and stage 10 (III) control egg chambers (w^{1118}/+;;matαTub-GAL4) stained with an HtsRC antibody, phalloidin, and DAPI. Panels below show anterior, M4, and M1 ring canals from each of the indicated egg chambers (which were located in different z-planes than the one shown in the larger panels); ring canal identity is indicated by the colored bars, which match those used in (E). Measurements of these ring canals and associated nuclei are included below; measurements were made in the z-plane where each structure was in focus. Scale bar in larger panel is 50 μm; scalebars in smaller panels are 5 μm. (B-D) log-log plots of egg chamber (EC) volume, nuclear area, or ring canal (RC) diameter from stage 5-10b. All four posterior nuclei and ring canals (M1, M2, M3, and M4) were measured. Error bars in B,C are s.d. n=61 egg chambers. 95% confidence intervals are shown. (E) Cartoon showing the stereotypical connections that result from the four rounds of mitotic divisions. Fill colors correspond to those used in (D), and the thickness of the outlines and connecting lines correspond to distance from the oocyte (thicker lines are closer to the oocyte, and thinner lines are further from the oocyte).

Figure 2: Reducing directed transport and oocyte growth has a modest effect on ring canal size and scaling. (A) Single plane confocal images of control and *dhc64C-RNAi* egg chambers (*otu-GAL4,UAS-Dcr-2*) stained with an HtsRC antibody, phalloidin, and DAPI. Scale bar is 50 μm. (B,C) log-log plots of egg chamber (EC) volume, oocyte area, or ring canal (RC) diameter from stage 5-10b. 95% confidence intervals are shown. Error bars in C are s.d. (D) Scaling exponent for ring canals separated by lineage. Error bars are 95% confidence intervals. n=49 egg chambers for control and 28 egg chambers for *dhc64C-RNAi*.

Figure 3: Over-expression of HtsRC::GFP increases ring canal size without affecting the lineage-based scaling. (A) Single plane or maximum intensity projections of confocal images of control and HtsRC::GFP over-expressing (OE) egg chambers (*matαTub-GAL4*) stained with an HtsRC antibody, phalloidin, and DAPI.

Scale bar is 100 μm. (B-F) log-log plots of egg chamber (EC) volume and ring canal

(RC) diameter and (G) the ratio of M1/M4 ring canal diameter from stage 5-10b. 95%

confidence intervals are shown. n=50 egg chambers for control and 54 egg chambers

for HtsRC::GFP OE.

Figure 4: Depletion of Tom6 increases the number of germline cells but does not

alter the pattern of ring canal scaling. (A) Single plane confocal images of control

and *tom6-RNAi* egg chambers (*nos-GAL4*) stained with an HtsRC antibody, phalloidin,

and DAPI. Maximum intensity projections of the *tom6-RNAi* egg chambers are included

to show the additional nurse cells and ring canals in these egg chambers. Scale bar is

50 μm. (B,C) log-log plots of egg chamber (EC) volume and ring canal (RC) diameter

from stage 5-10b. n=74 egg chambers for control and 64 egg chambers for *tom6-RNAi*.

95% confidence intervals are shown. (D) Scaling exponent for ring canals separated by

lineage. Error bars indicate 95% confidence interval. Asterisks indicate significant

difference between control and *tom6-RNAi* (ANCOVA, $p < 0.05$).

Figure 5: Differences in posterior ring canal size and scaling in "big egg" and

"small egg" *D. melanogaster* **lines.** (A) Box and whiskers plot (10-90th percentile) of

mature egg volume of "big egg" and "small egg" *D. melanogaster* lines. One-way

ANOVA with Tukey's multiple comparisons test indicates mature eggs from all "small

egg" lines (9.31.4, 8.26.2, 8.11.1, 7.17.4, and 9.2.2) are significantly smaller than all "big

egg" lines (1.16.1, 2.15.4, 2.49.3, 1.40.2, and 3.34.1) ($p < 0.0001$). n= 20-77 eggs per

line. (B) Scaling exponent for ring canals separated by lineage. Error bars indicate 95%

confidence interval. There were significant differences in the scaling exponent across

the four lines for M1, M2, and M4 (ANCOVA, $p < 0.05$). (C-F) log-log plots of egg

chamber (EC) volume and ring canal (RC) diameter in two "big egg" (1.40.2 and 3.34.1)

and two "small egg" (7.17.4 and 9.31.4) lines separated by lineage. 95% confidence

intervals are shown. n=29 egg chambers for 1.40.2, 46 egg chambers for 3.34.1, 44 egg

chambers for 7.17.4, and 43 egg chambers for 9.31.4.

<u>Acknowledgements:</u>

We would like to thank Seth Donoghue and Andrew Stoehr for helpful suggestions on
the manuscript and Cecelia Miles for the big and small egg lines. This work was
supported by the National Institutes of Health (NIH R15HD084243 to L.L. and
R15GM099054 to W.G.). The confocal microscope was purchased with support from a
Major Research Instrumentation (MRI) award from the National Science Foundation
(#2116348). Stocks obtained from the Bloomington Drosophila Stock Center (NIH
P40OD018537) were used in this study. The following antibody was obtained from the
Developmental Studies Hybridoma Bank, created by the NICHD of the NIH and
maintained at the University of Iowa, Department of Biology, Iowa City, IA 52242: hts
RC antibody developed by L. Cooley.

References:

Ali-Murthy, Z., Fetter, R. D., Wang, W., Yang, B., Royer, L. A. and Kornberg, T. B.
(2021). Elimination of nurse cell nuclei that shuttle into oocytes during oogenesis.
J. Cell Biol. **220**, e202012101.

Balachandra, S., Sarkar, S. and Amodeo, A. A. (2022). The Nuclear-to-Cytoplasmic
Ratio: Coupling DNA Content to Cell Size, Cell Cycle, and Biosynthetic Capacity.
Annu. Rev. Genet. **56**, 165–185.

Cantwell, H. and Nurse, P. (2019). Unravelling nuclear size control. *Curr. Genet.* **65**,
1281–1285.

Carvalho, A., Desai, A. and Oegema, K. (2009). Structural Memory in the Contractile
Ring Makes the Duration of Cytokinesis Independent of Cell Size. *Cell* **137**, 926–
937.

Diegmiller, R., Doherty, C. A., Stern, T., Alsous, J. I. and Shvartsman, S. Y. (2021).
Size scaling in collective cell growth. *Development* dev.199663.

Doherty, C. A., Diegmiller, R., Kapasiawala, M., Gavis, E. R. and Shvartsman, S. Y.
(2021). Coupled oscillators coordinate collective germline growth. *Dev. Cell* **56**,
860-870.e8.

Gayon, J. (2000). History of the Concept of Allometry1. *Am. Zool.* **40**, 748–758.

Gerdes, J. A., Mannix, K. M., Hudson, A. M. and Cooley, L. (2020). HtsRC-Mediated
Accumulation of F-Actin Regulates Ring Canal Size During Drosophila
melanogaster Oogenesis. *Genetics* **216**, 717–734.

Haglund, K., Nezis, I. P. and Stenmark, H. (2011). Structure and functions of stable
intercellular bridges formed by incomplete cytokinesis during development.
Commun. Integr. Biol. **4**, 1–9.

Han, B., Wei, Q., Amiri, E., Hu, H., Meng, L., Strand, M. K., Tarpy, D. R., Xu, S., Li,
J. and Rueppell, O. (2022). The molecular basis of socially induced egg-size
plasticity in honey bees. *eLife* **11**, e80499.

Huxley, J. S. and Teissier, G. (1936). Terminology of Relative Growth. *Nature* **137**,
780–781.

Imran Alsous, J., Villoutreix, P., Berezhkovskii, A. M. and Shvartsman, S. Y.
(2017). Collective Growth in a Small Cell Network. *Curr. Biol.* **27**, 2670-2676.e4.

Januschke, J., Gervais, L., Dass, S., Kaltschmidt, J. A., Lopez-Schier, H.,
Johnston, D. S., Brand, A. H., Roth, S. and Guichet, A. (2002). Polar
Transport in the Drosophila Oocyte Requires Dynein and Kinesin I Cooperation.
Curr. Biol. **12**, 1971–1981.

Jha, A. R., Miles, C. M., Lippert, N. R., Brown, C. D., White, K. P. and Kreitman, M.
(2015). Whole-Genome Resequencing of Experimental Populations Reveals
Polygenic Basis of Egg-Size Variation in *Drosophila melanogaster. Mol. Biol.*
Evol. **32**, 2616–2632.

Koch, E. A. and King, R. C. (1966). The origin and early differentiation of the egg
chamber of Drosophila melanogaster. *J. Morphol.* **119**, 283–303.

Loyer, N., Kolotuev, I., Pinot, M. and Le Borgne, R. (2015). *Drosophila* E-cadherin is
required for the maintenance of ring canals anchoring to mechanically withstand
tissue growth. *Proc. Natl. Acad. Sci.* **112**, 12717–12722.

Lu, W., Lakonishok, M. and Gelfand, V. I. (2021). Gatekeeper function for Short stop
at the ring canals of the Drosophila ovary. *Curr. Biol.* **31**, 3207-3220.e4.

Lu, W., Lakonishok, M., Serpinskaya, A. S. and Gelfand, V. I. (2022). A novel
mechanism of bulk cytoplasmic transport by cortical dynein in Drosophila ovary.
eLife **11**, e75538.

Marshall, W. F. (2020). Scaling of subcellular structures. *Annu. Rev. Cell Dev. Biol.* **36**,
219–236.

Mathieu, J., Cauvin, C., Moch, C., Radford, S. J., Sampaio, P., Perdigoto, C. N.,
Schweisguth, F., Bardin, A. J., Sunkel, C. E., McKim, K., et al. (2013). Aurora

B and Cyclin B Have Opposite Effects on the Timing of Cytokinesis Abscission in *Drosophila* Germ Cells and in Vertebrate Somatic Cells. *Dev. Cell* **26**, 250–265.

Matias, N. R., Mathieu, J. and Huynh, J.-R. (2015). Abscission Is Regulated by the ESCRT-III Protein Shrub in Drosophila Germline Stem Cells. *PLOS Genet.* **11**, e1004653.

McGrail, M. and Hays, T. S. (1997). The microtubule motor cytoplasmic dynein is required for spindle orientation during germline cell divisions and oocyte differentiation in Drosophila. *Development* **124**, 2409–2419.

Miles, C. M., Lott, S. E., Hendriks, C. L. L., Ludwig, M. Z., Manu, null, Williams, C. L. and Kreitman, M. (2011). Artificial selection on egg size perturbs early pattern formation in Drosophila melanogaster. *Evol. Int. J. Org. Evol.* **65**, 33–42.

Murphy, A. M. and Montell, D. J. (1996). Cell type-specific roles for Cdc42, Rac, and RhoL in Drosophila oogenesis. *J. Cell Biol.* **133**, 617–630.

Ong, S. and Tan, C. (2010). Germline cyst formation and incomplete cytokinesis during Drosophila melanogaster oogenesis. *Dev. Biol.* **337**, 84–98.

Ong, S., Foote, C. and Tan, C. (2010). Mutations of DMYPT cause over constriction of contractile rings and ring canals during Drosophila germline cyst formation. *Dev. Biol.* **346**, 161–169.

Orr-Weaver, T. L. (2015). When bigger is better: the role of polyploidy in organogenesis. *Trends Genet. TIG* **31**, 307–315.

Øvrebø, J. I. and Edgar, B. A. (2018). Polyploidy in tissue homeostasis and regeneration. *Dev. Camb. Engl.* **145**, dev156034.

Price, K. L., Tharakan, D. M. and Cooley, L. (2023). Evolutionarily conserved midbody remodeling precedes ring canal formation during gametogenesis. *Dev. Cell* **58**, 474-488.e5.

Riparbelli, M. G., Persico, V. and Callaini, G. (2022). Cell-to-Cell Interactions during Early Drosophila Oogenesis: An Ultrastructural Analysis. *Cells* **11**, 2658.

Robinson, D. N., Smith-Leiker, T. A., Sokol, N. S., Hudson, A. M. and Cooley, L. (1997). Formation of the Drosophila Ovarian Ring Canal Inner Rim Depends on Cheerio. *Genetics* **145**, 1063–1072.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji - an Open Source platform for biological image analysis. *Nat. Methods* **9**, 10.1038/nmeth.2019.

Singh, J., Imran Alsous, J., Garikipati, K. and Shvartsman, S. Y. (2022). Mechanics
of stabilized intercellular bridges. *Biophys. J.* **121**, 3162–3171.

Tilney, L. G., Tilney, M. S. and Guild, G. M. (1996). Formation of actin filament
bundles in the ring canals of developing Drosophila follicles. *J. Cell Biol.* **133**, 61–
74.

Uppaluri, S., Weber, S. C. and Brangwynne, C. P. (2016). Hierarchical Size Scaling
during Multicellular Growth and Development. *Cell Rep.* **17**, 345–352.

Yamamoto, S., Bayat, V., Bellen, H. J. and Tan, C. (2013). Protein Phosphatase 1ß
Limits Ring Canal Constriction during Drosophila Germline Cyst Formation.
PLOS ONE **8**, e70502.

Figure 5