A suite of selective pressures supports the maintenance of alleles of a *Drosophila* immune

peptide

Running Title: Antagonistic roles of *Diptericin* alleles

Authors: Sarah R. Mullinax[1], Andrea M. Darby[2], Anjali Gupta[3], Patrick Chan[1], Brittny R.
Smith[1], and Robert L. Unckless[1*]

[1]Department of Molecular Biosciences, University of Kansas, Lawrence, KS, USA

[2]Department of Entomology, Cornell University, Ithaca, NY, USA

[3]Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS, USA

[*]Corresponding author

Email: unckless@ku.edu

Keywords: balancing selection, innate immunity, antimicrobial peptide, diptericin

Abstract

The innate immune system provides hosts with a crucial first line of defense against pathogens.

While immune genes are often among the fastest evolving genes in the genome, in *Drosophila*,

antimicrobial peptides (AMPs) are notable exceptions. Instead, AMPs may be under balancing

selection, such that over evolutionary timescales multiple alleles are maintained in populations.

In this study, we focus on the *Drosophila* antimicrobial peptide Diptericin A, which has a

segregating amino acid polymorphism associated with differential survival after infection with

the Gram-negative bacteria *Providencia rettgeri*. Diptericin A also helps control opportunistic

gut infections by common Drosophila gut microbes, especially those of *Lactobacillus plantarum*.

In addition to genotypic effects on gut immunity, we also see strong sex-specific effects that are

most prominent in flies without functional *diptericin A*. To further characterize differences in

microbiomes between different *diptericin* genotypes, we used 16S metagenomics to look at the

microbiome composition. We used both lab reared and wild caught flies for our sequencing and

looked at overall composition as well as the differential abundance of individual bacterial

families. Overall, we find flies that are homozygous for one allele of *diptericin A* are better

equipped to survive a systemic infection from *P. rettgeri*, but in general have a shorter lifespans

after being fed common gut commensals. Our results suggest a possible mechanism for the

maintenance of genetic variation of *diptericin A* through the complex interactions of sex,

systemic immunity, and the maintenance of the gut microbiome.

Introduction

An effective immune response is essential for organisms to protect themselves from pathogens. However, an excessive immune response causes damage to self either directly (autoimmunity, cytokine storms) or through dysbiosis *via* disruption of the composition of beneficial microflora [1–4]. The dynamic suite of microbes that hosts face necessitates initiating a robust systemic immune response, avoiding self-harm, and maintaining a beneficial microbiome. The challenge of maintaining a balanced immune response is exacerbated because the threat of microorganisms is contextual; thus, the immune system must distinguish harmful versus beneficial microbes in each context [5,6]. For example, an ingested microbe may be harmless or even beneficial as a food source in the digestive tract, but, when in circulation, that same microbe may prove harmful to the organism [7]. One consequence of balancing a robust innate immune system is that it may lead to the maintenance of genetic polymorphism in genes that encode for proteins intimately involved in the immune response [8,9]. Such patterns of maintained polymorphisms contrast the standard coevolutionary arms race model where hosts and pathogen continually adapt to each other leading to rapid evolutionary change in immune genes [10–12]. Though the coevolutionary arms race model appears apt in many cases, the case for the adaptive maintenance of alleles, balancing selection in the broad sense, on genes involved in the immune defense stems from a more nuanced view of the delicate interplay of systemic immunity, life history traits and the beneficial microbiome [13,14].

Balancing selection is the general term for the adaptive maintenance of multiple alleles in a population. In contrast to the evolutionary arms race model, natural variation in many immune genes is maintained by balancing selection [15–18]. The inference of balancing selection from

molecular population genetic data is difficult, especially when genes are small and in areas of

high recombination. Nonetheless, several examples of balancing selection exist, many of which

involve small effector proteins of the innate immune system [18–21].

Balancing selection on immune genes is likely to involve allelic benefits that are only

conditionally beneficial. One way that an allele could be conditionally beneficial is if there is

specificity between pathogen and allele [22] such that *allele A* better protects against *pathogen 1*

and *allele B* better protects against *pathogen 2*: *pathogen specificity hypothesis*. Another way that

an allele can be conditionally beneficial in the context of immune defense is if resistance alleles

are costly in the absence of infection. In this case, *allele A* better protects against both *pathogen*

1 and *pathogen 2*, but in the absence of infection *allele A* is costly for its host [23]. This cost

could be energetic or through autoimmune-like damage: *autoimmune hypothesis*. In reality, there

is likely a continuum between these alternative hypotheses.

Invertebrates lack an adaptive immune system, so the delicate balance between systemic and gut

immunity is achieved only through adjustments to the innate immune response [24]. It is

therefore important to understand how invertebrates optimize their systemic immune response

with as little detriment to their beneficial gut microbiota as possible.

Though first characterized for its role in systemic immunity, the (IMD) pathway is also the main

NF-κB immune pathway in the gut and contributes to the maintenance of the microbiome and

protection from gut infections [25–27]. To date, there is little research into how individual

AMPs help maintain a healthy microbiome composition in the fly, let alone how allelic variation

may affect microbiome composition [28,29]. The standard gut microbiome of lab reared *D.*

melanogaster consists of two prominent bacterial genera: *Lactobacillus* and *Acetobacter* [27].

Both genera are easily cultured in the lab, and the *Drosophila* gut can easily be manipulated,

making it a good model to study the effects of immune gene variation on microbiome

composition and how microbiome composition can, in turn, influence host fitness.

Antimicrobial peptides (AMPs) are a critical part of the innate immune system that act as broad-

spectrum antimicrobials, combating bacteria, fungi, and viruses. In the coevolutionary arms race

model, AMPs are on the front lines: directly interacting with microbes. AMPs have also been

demonstrated to have critical roles in other physiological functions, for example, dysregulation

of AMPs has been connected to diseases such as atopic dermatitis and Alzheimer's disease

[2,30,31]. AMPs also contribute to the aging process where the immune system becomes more

active as organisms ages to compensate for a decline in its effectiveness. However, this leads to

cytotoxicity that in turn shortens lifespan [3].

In *Drosophila,* AMPs play a crucial in both manage systemic infections and in the maintenance

of the gut microbiota [32]. AMPs, however, appear to evolve more slowly than most immune

gene families [33–36]. One explanation for this perceived lack of adaptive evolution is that

Drosophila AMPs genetic variation is maintained adaptively through balancing selection [15,37–

39], where different selective pressures favor different alleles. Naturally occurring allelic

variation in AMP loci are sometimes associated with variability in pathogen resistance [19,40–

42]. *Diptericin A* (hereon referred to as *diptericin* or *Dpt)* is one of the canonical effector genes

of the *Drosophila* Immune deficiency (IMD) pathway, and is generally associated with defense

against Gram-negative bacterial infection [43,44].

In both *D. melanogaster* and *D. simulans*, an amino acid polymorphism at the 69th

residue of the mature 83-residue Diptericin peptide segregates in most populations surveyed

[19]. In both species, this polymorphism is due to a point mutation that changes the ancestral

serine (S) allele to arginine (R), but the two species use different codons for arginine. Unckless *et*

al. [19] found a significant difference in survival based on *diptericin* genotype after systemic

infection with the Gram-negative bacteria, *Providencia rettgeri,* a natural pathogen of *D.*

melanogaster. The study used inbred fly lines and found homozygous serine flies have a better

survival rate 5 days post infection (~60%) than homozygous arginine flies (~20%) or flies with a

premature stop codon in *diptericin* (0%). Later work by Hanson *et al.* took a more general

approach to AMP specificity and found that *Diptericin* plays a disproportionate role in response

to infection with *P. rettgeri [45].* What remains unclear, however is why this presumably

deleterious arginine allele persist in populations of two different species. We hypothesize that it

either protects against a different suite of pathogens or it is beneficial in the absence of infection

through some life history tradeoff.

This study aimed to test two hypotheses about the maintenance of genetic variation in *Diptericin.*

First, different *Diptericin* alleles might protect against different pathogens – we refer to this as

the *pathogen specificity hypothesis*. It would be supported if the arginine allele were to be

associated with higher survival than the serine allele in some infections, since the serine allele is

already clearly more protective against *P. rettgeri* [19,42]. In contrast, flies with the serine allele

might be generally better at surviving infection, but this may have a cost to the organism. We

refer to this as the *autoimmune hypothesis,* where a protective immune allele is associated with a

tradeoff that is costly in the absence of infection. We test whether there are tradeoffs between

systemic immunity and other life history traits – particularly traits related to the gut microbiome

utilizing genetically controlled lines for *Diptericin* alleles generated by CRISPR/Cas9-editing.

Flies were systemically infected with a panel of six different bacteria and homozygous serine

flies had a better 5-day survival with each bacterium. We then used axenic and gnotobiotic flies

to look at the lifespan of flies since AMP overexpression and microbial proliferation are

commonly observed in aging flies [46–49]. *Lactobacillus plantarum* is harmful to female flies

with non-functional diptericin, while homozygous arginine flies poly-associated with *L.*

plantarum and *A. tropicalis* had a longer lifespan than homozygous serine flies. In this way we

found evidence for a tradeoff between the ability to fight systemic infection and the ability to

control opportunistic gut infections.

Results

A single amino acid change drastically influences survival after infection.

Unckless et al. [19] found that in inbred lines from the *Drosophila Genetic Reference Panel*

(DGRP), flies homozygous for serine as position 69 of the mature Diptericin peptide survive

systemic infection with *P. rettgeri* much better than those homozygous for the arginine peptide at

the same position. This experiment showed only an association between immune defense and the

serine/arginine polymorphism since these inbred lines were on several different genetic

backgrounds. To control genetic background, we used CRISPR/Cas9 genome editing to create

both an arginine allele (single nucleotide change, dpt^{S69R}) and multiple null alleles (1 or 3 base

pair deletions, *Δdpt* flies), as well as control *dpt^{S69}* (serine at position 69 of the mature peptide) in

diptericin (Fig S1A, Table S1). The phenotype for our CRISPR/Cas9 edited flies showed striking

similarity to the inbred lines. In systemic infection challenges with *P. rettgeri, dpt^{S69}* flies are

better protected from infection than *dpt^{S69R}* flies (p= 5.42e-08) and *Δdpt* flies (p=6.46e-09, Fig

1A). Remarkably, for inbred lines

and CRISPR/Cas9 edited lines,

survival for five days post infection

with *P. rettgeri* for the serine allele

lines is 50-60%, with the arginine

allele is 10-20%, and less than 5%

for null alleles.

We next challenged the

CRISPR/Cas9 edited flies with

systemic infection using multiple

other Gram-positive and Gram-

negative bacteria to determine

whether the arginine allele (*dpt^{S69R}*)

protects against some infections

better than the serine allele. Such a

finding would support the

hypothesis that allelic variation is

Genotype — *dpt^{S69}* — *dpt^{S69R}* — *Δdpt* — *imd$^-$*

Figure 1: Five-day survival post systemic infection.
Survival after systemic immunity challenge with the following bacteria A) *P. rettgeri,* B) *Staphylococcus succinus*, C) *Enterococcus faecalis*, D) *Lysinibacillus fusiformis* Strain Juneja, E) *Lactococcus lactis* F) *Serratia marcescens*. Each graph represents the combined results of 3 different infection dates of at least 20 males of each genotype for each date (at least 60 total). Significance is relative to dpt^{S69} (red line) using Cox proportional-hazards regression model. *P<0.05, **P<0.01, ***P<0.001.

maintained by different alleles providing specific protection against different microbes.

However, the differential response to systemic infection with *P. rettgeri* at an OD_{600}=0.1, as

described above, remains the largest difference in immune response between *Dpt* genotypes.

Dpt^{S69} flies survived better than *dpt^{S69R}* flies for all systemic infections tested (Fig 1, Table S2).

In the Gram-positive bacterial infections (*Enterococcus faecalis*, *S. succinus*, *L. fusiformis,* and

L. lactis), *dpt^{S69R}* flies had lower survival than that of *Δdpt* or *imd^-* (null allele for the Imd gene)

flies (Fig 1B-E). The only systemic infection where *dpt* genotype did not seem to matter was *S.*

marcescens, where all flies survive infection well except for *imd^-*, which die very quickly (Fig

1F).

We also tested if males and females showed differences in systemic immunity based on genotype

for *P. rettgeri, E. faecalis* and *L. plantarum* infections (Fig S2, Table S3). Generally, the *Dpt*

genotypes have qualitatively similar survival patterns between males and females when infected

with *P. rettgeri.* Females showed lower survival compared to males after systemic infection with

E. faecalis (p=0.0003). Higher male survival post infection was observed previously for both *E.*

faecalis and *P. rettgeri* with differences ascribed to the Toll pathway [50].

Overall, flies with the *dpt^{S69}* allele are better equipped to survive a systemic bacterial infection

than *dpt^{S69R}* flies, though this is most pronounced for *P. rettgeri.* Of the 6 bacteria tested, there

was no case where *dpt^{S69R}* flies survived the infection better than *dpt^{S69}* flies. These results do not

support the hypothesis that alleles are maintained to better combat different pathogens.

Diptericin genotype affects lifespan of mono- and poly- associated gnotobiotically reared flies.

Although our survey of systemic infections was not exhaustive, we did not find any instances where dpt^{S69R} flies were better able to fight infection, so we turned our attention to the role of *Dpt* in gut microbiome maintenance and immunity. The gut microbiome influences several life history traits in *Drosophila* and other organisms [51–54]. To dissect how *diptericin* genotype influences microbiome maintenance, we manipulated the microbiota in CRISPR/Cas9 flies, and measured longevity and bacterial load. We began with the longevity of axenically reared flies, since it represents a baseline survival without the presence of microbes. Flies with functional copies of *diptericin* (dpt^{S69} or dpt^{S69R}) did not differ in lifespan for either sex ($p_{genotype}=0.3762$), and *Δdpt* lines show a similar lifespan. However,

Figure 2: Axenic and gnotobiotic lifespan.
A) Survival plots for axenic flies and gnotobiotic flies mono-associated with L. plantarum or poly-associated with L. plantarum and A. tropicalis. Each curve represents 60 flies from 3 replicates of 20 flies. Significance is shown in comparison to SS (red line) using the model Lifespan~(Genotype*Sex)/Vial+Block. B) Mean lifespan was plotted against each sex and separated by bacterial association. Axenic flies have no bacterial association. Each point for each sex in a combination of 3 separate trials of 20 flies for a total of 60 flies. These interaction plots only have mean lifespan and thus are useful for seeing all the data at a glance. Error bars are mean plus or minus the standard error of the mean. *P<0.05, **P<0.01, ***P<0.001.

there was a difference in overall lifespan between the sexes among the CRISPR genome edited

lines (p_{sex}=1.04e-6). In dpt^{S69} or Δdpt female flies have a longer lifespan than male flies, as

observed generally for *D. melanogaster* previously [55]. In contrast, *imd*⁻ male flies have a much

longer lifespan than any of the other lines tested (Fig 2A – axenic row, Table S3).

We next tested the influence of Diptericin genotype on mono-associations with standard

constituents of the *Drosophila* gut microbiome. We found multiple sex effects when axenically

reared flies were fed *L. plantarum, L. brevis,* or *A. tropicalis* (Figs 2A and 2B, S3). Flies that

were fed *L. plantarum* show the most striking differences. Male flies continued to show similar

lifespans to each other, but female *Δdpt* and female *imd*⁻ flies both succumbed quickly post-

feeding, indicating functional *Diptericin* is important for gut immunity against opportunistic *L.*

plantarum in females.

We also fed axenic flies *P. rettgeri,* which does not readily cause gut infection, to axenically

reared flies. Recall that after systemic infection dpt^{S69} flies are more resistant to *P. rettgeri* than

dpt^{S69R} flies (Fig 1A). After monoassociation, however, we see no significant difference between

the two genotypes with *P. rettgeri* in either sex (p=0.612, Fig 2B). There is also no significant

difference between the *Δdpt* and lines with functional Diptericin (Fig 2B, S3). Surprisingly,

however, the null allele for *Imd* again shows a significant effect on survival in a sex-specific

manner: null *Imd* females die much earlier and null *Imd* males survive much longer when fed *P.*

rettgeri.

Some of the sex-specific differences in survival after monoassociation with bacteria may be

driven by intrinsic differences in feeding rates between the sexes (and potentially genotypes as

well) [56]. To determine whether the differences in male and female survival and load were due

in part to differences in feeding rates, we performed a feeding rate assay with blue dye mixed

with media (Luria-Bertani broth) or *P. rettgeri* (OD_{600}=15.0). We noted that females did eat

more in a single hour of feeding, but that there was also an effect of *diptericin* genotype (with

null flies eating less, Figure S4). Thus, it is possible that differences in longevity after

monoassociation are due to different rates of exposure to those bacteria because of different

feeding rates.

We next looked at how poly-associations with bacteria affected lifespan. First, we fed flies a 1:1

mixture of *L. plantarum* and *A. tropicalis*, two common gut microbes found in lab-reared and

wild caught flies [57,58]. We found that female *Δdpt* and *imd⁻* flies had a much shorter lifespan

than flies with functional Diptericin (Fig 2A and 2B). This is the same pattern observed in mono-

association with *L. plantarum*. However, we observe that *diptericin* genotype influences survival

when poly-associated with *L. plantarum* and *A. tropicalis*. Dpt^{S69R} female flies live longer than

dpt^{S69} female flies (Fig 2A and 2B; p=0.00782). This is the opposite of systemic infections,

where dpt^{S69} flies always survived better than dpt^{S69R} flies, and may indicate a tradeoff between

defense against systemic and gut immunity.

In poly-association with 1:1 *L. plantarum* and *P. rettgeri,* we observe many of the same patterns

as in the poly-association with *L. plantarum* and *A. tropicalis*. Again, female *Δdpt* and *imd⁻* flies

have a shorter lifespan than female dpt^{S69} and dpt^{S69R} flies (Fig 2B), and *dpt* genotype is

associated with lifespan. *Dpt^{S69R}* females have a longer lifespan than *dpt^{S69}* flies (p=0.000198,

Fig 2B).

Given the genotypic and sex effects on survival after oral association with *L.* plantarum, we also

looked at sex differences after systemically infecting conventionally reared flies with *L.*

plantarum. We saw no difference in survival between the sexes, as observed for systemic

infections with *E. faecalis* or *P. rettgeri* and saw no difference between *Δdpt* lines and lines with

functional Diptericin, as observed for axenic flies mono-associated with *L. plantarum* (Fig S2).

This could indicate that Diptericin plays different roles for systemic and gut immunity in relation

to *L. plantarum* in each sex. Further, *dpt^{S69}* male flies survived better than *dpt^{S69R}* male flies

(p=0.00438), in line with observations from other systemic infections in males (Fig 1).

Overall, we found a role of both specific *diptericin* genotype (serine vs. arginine) and the

presence of functional copies of *diptericin* for survival after introducing common gut microbes in

controlled conditions. Most striking was the sexually dimorphic role of both *Dpt* and *Imd-,* with

females being much more sensitive to genotype than males.

Gnotobiotic fly bacterial load

Given the differences in survival among sexes and genotype for different gnotobiotic

associations with bacteria, we determined whether bacterial load after associations also were

different among sexes and genotypes. To assess the influence of genotype on the immune

response of aging flies, we studied how well common gut bacteria colonized the gut over 20

days, representing ~20-33% of *D. melanogaster's* normal lab lifespan [59,60]. We generated

gnotobiotic flies by feeding specific bacteria for

two days and observed the bacterial load 2-, 10-

and 20-days post feeding.

When specifically looking at mono-association

with *L. plantarum* we observe that bacterial load

differences between genotypes occur within the

first 2 days post feeding but begin to disappear by

day 20 ($p_{genotype(day2)}$=2.6e-5, $p_{genotype(day20)}$=0.709,

Fig 3A, S5). Note that in the first two days, the

flies were raised on microbe-contaminated media,

but after 2 days were moved onto sterile food and

then transferred to new sterile food every 3 days.

This corroborates what we saw in the longevity

data in females. Within the first 15 days a large

proportion of *Δdpt* female flies died, and we

observed a higher bacterial load in these flies,

especially on day 2 post feeding (Fig 2). By day

20, differences in bacterial load in females

disappeared. Note that there is inherent sampling

bias, as only the flies able to survive until day 20

are sampled at that time point. In the case of *imd*

Figure 3: Bacterial load interaction plots for day 2, 10, and 20 post association with various microbes.
Two-day old flies were placed on food seeded with a bacterial suspension of each bacteria at an OD600 of 15. Each point for each sex is from 3 separate trials of 2 samples of 3 flies each for a total of 6 samples per point. A) monoassociations where each plot represents a different condition, B) polyassociations where each row plots both the *P. rettgeri* and *L. plantarum* load from the same flies but were plated on LB and MRS respectively. Error bars are mean plus or minus the standard error of the mean.

flies, no females survived until day 20, hence there is no data for imd^- flies on day 20 (Fig 3A,

S5).

P. rettgeri is the only canonical systemic pathogen we included in our gnotobiotic study, and the

patterns based on genotype were different between systemic infection and the gnotobiotic oral

infection data. When flies were fed *P. rettgeri,* we saw a range of bacterial loads, from zero

colonies to bacterial load levels on par with common gut bacteria. We also observed a genotype

effect between dpt^{S69} and dpt^{S69R} on day 10- and 20- post feeding in males ($p_{genotype(day10)}$=0.0044,

$p_{genotype(day20)}$=0.0004, model only included dpt^{S69} and dpt^{S69R}, Table S5). In both instances,

bacterial load in dpt^{S69R} flies is higher than in dpt^{S69} flies. This may indicate that dpt^{S69} flies are

better equipped to deal with both systemic and oral infection from *P. rettgeri*. Whether this is

due to a greater ability of dpt^{S69} flies to withstand the effects of infection (tolerance) by *P.*

rettgeri by remains a question. It is also important to note that *P. rettgeri* establishes poorly in

the gut of wildtype flies (personal observation), which may explain the noisy results for mono

associations after oral infection.

A range of bacterial loads were also observed when flies were poly-associated with *L. plantarum*

and *P. rettgeri* (Fig 3B, S6). There were no statistically significant differences between dpt^{S69}

and dpt^{S69R} bacterial for this poly association (Table S6). In fact, there were no significant

differences between any of the *P. rettgeri* bacterial loads, however *L. plantarum* does show

differences on day 2-post feeding (p=7.97e-6). These differences are mainly between flies with

non-functional *diptericin* (*imd-* and *Δdpt*) and flies with functional *diptericin*. Both *Δdpt* and

imd- flies had higher bacterial load than dpt^{S69} and dpt^{S69R} flies on day 2 post feeding which may

be an indication of the reason both these lines quickly succumb to feeding with *L. plantarum* at

least in the context of the poly-association with *P. rettgeri*.

We observed larger differences in bacterial load shortly after feeding with bacteria and those

differences became less by day 20-post feeding. We did not see any differences in *P. rettgeri*

load when mono-associated or when part of a poly-association, indicating the flies respond

differently to the same pathogen when introduced systemically or orally.

Evidence for life history tradeoffs mediated by *diptericin* genotype

Proteins involved in a robust immune response may have pleiotropic effects on other traits either

because of a direct interaction with that trait or because of inherent costs of immune defense

(self-damage, energy expenditure, etc.) [61–63]. We examined three life history traits

(desiccation stress survival, starvation stress survival, and uninfected longevity) to determine

whether *Dpt* genotype had such pleiotropic effects.

The ability to survive desiccation stress is an important life history trait for wild *Drosophila*

survival [64,65]. When subjecting our male CRISPR/Cas9 flies to desiccation, we observe

conventionally reared flies succumb faster to desiccation stress than axenically reared flies (Fig

4A, $p_{genotype} < 2e-16$). In conventionally reared males, dpt^{S69} flies have similar desiccation

resistance to dpt^{S69R} flies (p=0.917). However, dpt^{S69} flies survive desiccation stress better than

dpt^{S69R} flies when reared axenically ($p_{genotype}$=0.04126). We also compared *Drosophila* OreR and

W1118 (both homozygous for the serine allele of Diptericin) to the dpt^{S69} line (Fig S7, Table S7).

Unsurprisingly, despite the same *diptericin* genotype, all 3 lines show dramatically different

desiccation resistance. Therefore, *diptericin* genotype plays limited, if any role in variation in

desiccation resistance, as all 3 wildtype lines were derived from different genetic backgrounds.

We next looked at the effect

of genetic variation in

diptericin on male ability to

survive under starvation

stress. As with desiccation

resistance, we found

axenically reared flies

survive starvation stress

better than conventionally

reared flies (Fig 4B, S8,

$p_{treatment} < 2e-16$). However,

unlike the desiccation stress

conditions, both

conventionally reared and

axenically reared dpt^{S69R}

flies survive starvation

stress longer than dpt^{S69}

Figure 4: *Dpt* genotype is associated with variation in some life history traits

A) Conventionally and axenically reared male flies desiccation resistance (N>90 for each line apart from *Δdpt* line). There is no significance between dpt^{S69} (red line) and other genotypes. B) Conventionally and axenically reared male flies starvation resistance (N>90 for each line apart from *Δdpt* line). C) Female and male lifespan for conventionally reared flies. N>93 for each line and sex. Significance is shown in relation to dpt^{S69} (red line). Note that axenic longevity is shown in Figure 2. *P<0.05, **P<0.01, ***P<0.001.

flies (conventional: $p_{genotype} < 2e-16$, axenic: $p_{genotype} < 2e-16$). In conventionally reared flies

there is no difference in survival between dpt^{S69} flies and *Δdpt* flies (Fig 4B, Table S8). However,

in axenically reared flies, *Δdpt* flies have an intermediate survival phenotype between dpt^{S69} and

dpt^{S69R} flies. This may suggest an interaction between functional *diptericin* and the microbiome

that influences starvation.

Finally, we looked at overall longevity of female and male conventionally reared flies (in the

absence of any infection or other significant selection pressure). Female flies have a longer

lifespan than males (Fig 4C, S9, p_{sex}= 5.80e-13) regardless of *dpt* genotype. In males, *dpt^{S69}* flies

had a significantly longer lifespan than male *dpt^{S69R}* flies (mean of 60.3 days for *dpt^{S69}* and 54.1

for *dpt^{S69R}*, p= 0.0072), but not in female flies (mean of 61.9 days for *dpt^{S69}* and 59.0 for *dpt^{S69R}*,

p= 0.6434). However, in axenically reared flies only female *dpt^{S69}* flies have a longer lifespan

than male *dpt^{S69R}* flies. Interestingly, the female flies with the longest lifespan have non-

functional Diptericin (Fig 4C, *Δdpt* line and *imd⁻* line, Table S9). Males lacking functional

Diptericin show the same effect, but to a lesser extent. These results provide evidence that, in the

presence of a standard gut microbiota, both *Dpt* and a functional Imd pathway may decrease

longevity. This is consistent with others who found downregulation of NF-κB pathways and

AMPs increased lifespan in *Drosophila* [63,66], but the fact that flies with *Dpt* null alleles alone

are sufficient to increase lifespan is noteworthy.

Diptericin's influence on gut microbial diversity

To determine whether *Diptericin* genotype influences the composition of the bacterial

community in the gut, we sequenced amplicons of 16s ribosomal rRNA in conventionally reared

lab flies under two different rearing conditions: flies reared in standard *Drosophila* vials, and the

progeny of the cross between *dpt^{S69}* and *dpt^{S69R}* flies reared in cages for more than 2 generations.

The vials may capture the long-term impact of genotype on microbiota but makes ruling out

stochastic changes in communities difficult. The cages ensure identical microbiota during

development, but do not allow gradual effects of genotype on microbiota to accumulate. First,

we found flies that were co-reared in cages had similar microbiomes, with no discernable

differences by the alpha diversity metric, Shannon diversity (p=0.5239, Fig 5A). On the other

hand, the microbiomes of flies reared in vials was distinctly different overall compared to the

microbiomes of flies reared in the cages (p<0.001, Fig 5B). However, the differences between

genotypes were still minimal in the vial-reared flies and vial may be a large factor in differences

between lines. There were 2 dpt^{S69R} lines used and even though these are the same genotype and

genetic background there was a significant difference between the 2 lines (Fig 5B). This

difference is likely due

to the within-vial drift

in gut microbiome

communities [67],

which we attempted to

control for using flies

of different genotypes

reared in the same cage

(Fig 5A).

The lab is a controlled

environment that

allows for excellent

control of variables but

Figure 5 The influence of *Dpt* genotype on the gut microbiota
A) Shannon diversity of flies co-reared in cages. B) Shannon diversity of
individual genotypes reared in vials. C) Bray-Curtis dissimilarity of wild caught
flies. D) Differential abundance of Morganellaceae in wild caught flies. Different
letters show significance based on post-hoc Tukey test, P<0.05.

does not replicate the conditions found in nature. Thus, we looked at the microbiomes of wild

caught flies collected from the decaying fruit of an apple orchard. We found flies with

homozygous for dpt^{S69R}, homozygous for dpt^{S69}, and heterozygous genotypes at our collection

location (Table S10). Out of the 955 *D. melanogaster* flies successfully genotyped, only 20 flies

were homozygous for dpt^{S69R} were identified. All 20 homozygous dpt^{S69R} flies were profiled

along with 36 each of dpt^{S69} homozygous flies and heterozygous flies for a total of 92 flies

profiled (amplicon sequences of both *Dpt* and 16s rRNA are in Table S2).

We found that *dpt* genotype does not affect the overall composition of the microbiome of wild

caught flies based on Shannon diversity (alpha diversity) and Bray-Curtis dissimilarity (beta

diversity) (Fig 5C). However, when looking at differential abundance of individual bacterial

families there are differences based on genotype. We tested specific association between

diptericin genotype and normalized counts from 9 microbial families. Most intriguing is the

difference in abundance of *Morganellaceae* family reads since, *P. rettgeri* belongs to this family.

Heterozygous and homozygous serine flies have a slightly higher abundance than homozygous

arginine flies (Fig 5D), but these differences were not significant.

Overall, our results for associations between *Dpt* genotype and gut microbiome in both lab and

wild flies are weak. Such associations may require much larger sample sizes if the effects are

small – particularly given the noisy phenotype in the wild. Alternatively, the influence of *Dpt*

genotype on gut microbiome diversity may be specific to developmental life stages or microbes

that are relatively rare in these populations.

Discussion

The adaptive maintenance of multiple alleles of a single gene has been posited for nearly a century [68–72] and evidence for its pervasiveness continues to grow [73–78]. However, there are relatively few cases where we understand the adaptive benefits of both allele (sickle-cell anemia and malaria resistance being the best studied example [79]). In this study, we generated CRISPR/Cas9 genome-edited flies of different *diptericin* genotypes to investigate mechanisms of maintenance of allelic variation in AMPs. This allowed us to specifically study a single amino acid change on an otherwise genetically-controlled background. Overall, we found evidence that dpt^{S69} flies survive systemic infection better, while dpt^{S69R} flies survive some opportunistic gut infections better. Thus, more robust defense against systemic infection appears to interfere with maintaining a balanced gut microbiota – this tradeoff may result in natural populations maintaining both alleles. Importantly, we note that while our results are consistent with the adaptive maintenance of alleles in natural populations, these lab assays do not prove the adaptive value of different alleles in the field.

Systemic infections with *P. rettgeri* in CRISPR/Cas9 genome-edited lines confirmed the amino acid polymorphism in Diptericin as the basis for the difference in immune response of flies with different *diptericin* genotypes previously hypothesized *via* an association study [19]. Surprisingly, dpt^{S69} flies had a higher survival five days post infection than dpt^{S69R} flies in *all* systemic infections tested. This may mean either that the serine allele provides improved defense against all the tested systemic infections, or that alterations to other aspects of fly physiology (notably the gut microbiota) could predispose arginine flies to poorer survival. It is also possible that some of *diptericin's* influence on survival is mediated by changes in the microbiome. We

did not perform systemic infections on axenically-reared flies, but if *Dpt* genotype influences the

microbiome, which in turn influences survival after infection with other pathogens, *Dpt* should

not influence survival after systemic infection if flies are reared axenically.

Our study also reveals an important role of Diptericin in preventing opportunistic gut infections

by common gut microbes. This is especially evident when looking at lifespan after associating

axenic flies with *L. plantarum* in both mono- and poly- association contexts. In both sexes,

dpt^{S69R} flies have a longer lifespan than *dpt^{S69}* flies, and this effect is enhanced in poly-

association with *L. plantarum* and *A. tropicalis*. Other studies have posited a role for individual

AMPs in the maintenance of gut microbes in *Drosophila* [29], but this appears to be the first

example linking a functional copy of a single AMP to survival differences.

One of our most striking and unexpected results was the distinct sex differences in longevity

after association with common gut bacteria. Furthermore, fly genotype plays a significant role in

these differences. After association with *L. plantarum*, *imd-* females had the shortest mean

survival, while *imd-* males had the longest mean survival. The effect for *Δdpt* was similar:

females had much shorter average survival than the other two *dpt* alleles, but the three alleles

showed equivalent longevity in males. While the *imd-* sexual dimorphism is relatively consistent

across experiments with different microbes, the *Δdpt* effect seems to be more limited to

associations involving *L. plantarum*. This result not only highlights the importance of using both

sexes in microbiome research, but also confirms that functional Diptericin is important for

Drosophila melanogaster gut health. There is a growing body of literature on the effects of

sexual dimorphisms in immune response, but it is lacking for sexual dimorphisms in gut

immunity and our findings only emphasize the need to fill this gap (*Drosophila* immunity sexual

dimorphisms reviewed in Belmonte et al. 2020; also see [80]). Our feeding assay provides some

evidence that the differences between male and female survival after exposure to microbes is due

the amount of food (and therefore the number of microbes) consumed, but these differences

should be examined in much more detail. More generally, the sex by genotype effect suggests a

likely explanation for the maintenance of genetic variation: different fitness effects of alleles in

the different sexes [81]. Note that while our study focuses on the serine/arginine polymorphism,

at least 6 null alleles of *dpt* segregate in natural populations of *D. melanogaster* and the

frequency of those nulls shows clinal variation in both North American and Africa [82], thus the

nulls are likely maintained selectively too.

Finally, we note that we have framed our argument in terms of balancing selection in the broad

sense because we are agnostic as to the population dynamics upon which these different selective

pressures might act. For example, it may be that some populations are disproportionately

infected by *P. rettgeri* or other microbes for which the serine allele is favored, while other

populations are relatively pathogen-free and the arginine allele is favored. This scenario is partly

supported by the evidence of clinal patterns in allele frequency in Africa and (to a lesser extent)

North America [83]. Similarly, these selective pressures could vary temporally. On the other

hand, these pressures could act all in one population where serine is favored in infected

individuals and the arginine allele is favored in uninfected individuals.

Herein we highlight 3 main roles of Diptericin in *Drosophila*. 1) Diptericin genotype influences

systemic immune defense; 2) in certain conditions Diptericin genotype influences gut immunity;

and 3) Diptericin has sex-specific effects in the gut. These results highlight the need for

individuals and populations to modulate the immune system to balance systemic and gut

immunity, and how the different needs of females and males complicate this balance. The

dramatic differences in survival of males and females in response to oral infection by common

gut bacteria underlines the importance of looking at both sexes when examining the maintenance

of genetic diversity, as balancing selection may be caused by sexual dimorphism. Overall, our

results suggest that a complex interaction between sex, environmental context (starvation,

pathogen exposure – both systemic and oral), and genotype may contribute to the long-term

maintenance of immune alleles.

Materials and Methods

Drosophila Lines and Rearing

Conventionally reared flies were maintained in a 23°C incubator with a 12h light:12h dark

schedule on a cornmeal-molasses-yeast diet (64.3g/L cornmeal, 79.7mL/L molasses, 35.9g/L

yeast, 8g/L agar, 15.4mL of food acid mix (50mL Phosphoric Acid + 418mL Propionic Acid +

532mL deionized water) and 1g/L Tegosept. We used CRISPR/Cas9 genome editing to modify

the *diptericin A* gene. Briefly, DNA coding for guide RNA (gRNA) was inserted into the pUS-

BbsI plasmid (Table S1). A single stranded donor DNA (120bp) containing the desired edit (to

change from serine to arginine) and a silent mutated PAM site was synthesized by IDT

(Coralville, IA, USA). The plasmid and ssODN were then injected into Bloomington stock

#55821, which expresses Cas9 driven by the *vasa* promoter, by Genetivision, Inc. (Houston, TX,

USA). Individual flies developed from the injected embryos were collected and crossed with a

modified version of Bloomington stock # 7198 (a line with *CyO*/*Kruppel* balanced on the 2nd

chromosome, and *serrate/Dichaete* balanced on the 3rd chromosome). Our version, 7198^{A4}, was

provided by Stuart Macdonald and has the DSPR [84] A4 line's X chromosome instead of the

w[*] from the original 7198. The F1s were collected and again individually crossed with 7198^{A4}

for F2 crosses yielding individuals with a homozygous 2nd chromosome representing one of the

chromosomes carried by the original injected embryo. The Dpt gene was sequenced from the F2

cross progeny to determine whether edits occurred. This yielded several classes of alleles

including homozygous serine *dpt* (wildtype, *dpt*^{S69}), homozygous arginine *dpt* (*dpt*^{S69R}), and *dpt*

null (*Δdpt,* refers to lines with either 1 or 3 base pair deletion) (Fig S1A). Balancers were

removed, and lines were moved into the same genetic background through a series of crosses as

shown in Fig S1B. An *imd-* line was used as an IMD pathway negative control but note that this

line was from a completely different genetic background from the rest of the lines.

Axenic Fly Preparation

Microbe-free (axenic) lines were generated by first washing embryos in a 10% bleach solution to

dissolve the chorion for 2 minutes. The embryos were then washed in 70% ethanol for 30

seconds and water for another 30 seconds, then transferred to autoclaved molasses food (see

above). Some embryos from each treatment were placed onto De Man, Rogosa and Sharpe

(MRS) agar plates and incubated at 30°C for 48 hours to check that they did not contain viable

microbes. Axenic lines were continuously checked for the presence of contaminating microbes

(every 3-4 generations) by homogenizing flies and plating the homogenate on MRS agar.

Axenically and gnotobiotically (see below) reared flies were maintained in an incubator that was

isolated from conventionally reared flies. The incubator was kept at 23°C with a 12h light:12h

dark schedule. Axenic and gnotobiotic flies were kept on the same molasses diet that had been

autoclaved before dispensing into autoclaved vials. Axenic and gnotobiotic flies were only

handled inside a sterile hood (Baker SG 400, The Baker Company Inc., Sanford, ME).

Bacterial Strains

The following bacteria were used for systemic infection assays: *Providencia rettgeri* [85],

Providencia burhodegraneria Strain B [85], *Enterococcus faecalis* [85]), *Serratia marcescens*

[40], *Lysinibacillus fusiformis* Strain Juneja [86], and *Staphylococcus succinus* (isolated from

wild *Drosophila,* Unckless lab). Bacteria were grown from glycerol stocks on LB plates at 37°C

overnight.

The following bacteria were used in gnotobiotic experiments: *L. plantarum, L. brevis, A.*

tropicalis. All these strains were isolated from plating conventionally reared flies on MRS agar

in the Unckless lab at the University of Kansas. Individual colonies from plated fly homogenate

were grown overnight in MRS for DNA isolation. Bacterial species were identified using Sanger

sequencing with the 16S rRNA region primers 27F (AGAGTTTGATCCTGGCTCAG) and

1492R (CGGTTACCTTGTTACGACTT). We also utilized the same *P. rettgeri* as described for

systemic infection.

Survival Assays

For systemic infections, individual colonies of bacteria were picked and grown in 2mL LB broth

shaking overnight at 37°C. Bacterial suspensions were then diluted or concentrated to OD_{600}=0.1

for *P. rettgeri,* OD_{600}=10 for *S. succinus,* OD_{600}=1.5 for *E. faecalis*, OD_{600}=3.5 for *L. fusiformis*,

$OD_{600}=1.0$ for *L. lactis*, and $OD_{600}=4.0$ for *S. marcescens*. *L. plantarum* was grown in 5mL

MRS at 30°C overnight and was concentrated to $OD_{600}=10$ for systemic infections. To induce

systemic infection, 5-9 day old, conventionally reared flies were pricked in the thorax with a

needle dipped in a bacterial suspension [87]. Infections were done in triplicate with at least 20

flies for each replicate per line for a total of 60 flies per genotype per condition. Flies were

incubated at 23°C with a 12h light:12h dark schedule and survival was tracked daily for 5 days

post infection.

Gnotobiotic Longevity

Axenically reared flies were collected within 24 hours of eclosion. Flies were then kept on sterile

food for 2 days before sorting for longevity experiments. To begin longevity experiments, flies

were separating into groups of 10 individuals of each sex and put onto sterile food seeded with

50uL of bacterial suspension at an OD600 of 15 +/- 1. For each replicate, we used 2 vials of 10

flies each per sex per line for a total of 20 flies for each sex per genotype for a total of 60 flies

across all replicates. Flies were allowed to feed in the inoculated vials for 3 days before being

transferred to uninoculated sterile food vials. Flies were flipped to new sterile media every 4-5

days for the remainder of the experiment. Surviving flies were counted every 1-3 days until all

flies were dead.

Gnotobiotic Bacterial Load

To determine whether microbes became established in the gut, we homogenized flies during and

after the exposure and plated the homogenate. We measured bacterial load by inoculating flies in

the same manner as gnotobiotic longevity. Flies were separated into groups of 5 females and 5

males per vial. For 2-day experiments flies were kept on the seeded food for the entire

experiment. For 10– and 20- day experiments, flies were allowed to feed on the seeded food for

3 days before being transferred to sterile food. Flies continued to be transferred to new sterile

media every 3-4 days until day 10 or 20 post feeding. After the experimental (2, 10 or 20 day)

time period, flies were surface sterilized by washing in 70% ethanol followed by molecular grade

water. Flies were separated by sex and three individuals were homogenized together in 300uL of

sterile 1x PBS and the homogenate was plated on the appropriate media using a Whitley WASP

Touch® spiral plater (Don Whitley Scientific, UK). When there were not 3 flies still alive then

all remaining flies of a sex were used and squished in 100uL of 1x PBS per fly collected instead

of 300uL to keep all samples the same concentration per fly. Counts were adjusted accordingly.

Feeding Rate Assays

To measure the amount of food and bacteria consumed by males and females of different

genotypes, we made food containing blue dye by adding 11.2g FCF blue dye (Erioglaucine

disodium salt) per liter of food. We used newly eclosed flies (1-2 days post eclosion / 14 days

post oviposition) and separated sexes (in sterile conditions) and keep flies at a density of 10

flies/vial in fresh food vials. We held these flies in incubators for one day, so they would recover

stress of anesthesia during sexing. To introduce bacteria (or control media) into the food, we

pipetted 50uL of suspension or LB at an OD600 of 15 (+/- 1) into each vial and allowed the

suspension to absorbed into the food for 30-40 mins by keeping the vials open inside a sterile

hood. We next added the experimental flies and allowed them to feed for one hour. After 1 hour,

we anesthetized the flies on ice, then rinsed in ethanol and sterile water. Flies were homogenized

in 300µl of 1X PBS with a glass bead (maximum speed for 4 minutes). The homogenate was

centrifuged at 14000 RPM for 4 minutes and 200μl of the supernatant was used to measure

absorbance at 630nm. Absorbance differences were analyzed using the natural log of absorbance

as the response variable with genotype, sex, genotype by sex interaction, and block as

independent variables. Due to the blocking structure of the experiment, each treatment (no media

control, LB control, *P. rettgeri*) was analyzed separately.

Desiccation

Desiccation survival assays were performed on dpt^{S69} and dpt^{S69R} adult males 4-7 days post-

eclosion in conventionally and axenically reared flies. Ten males were placed into an empty vial

and closed off with a cotton flug. Each genotype had 5 vials for a total of 50 flies per line per

rearing condition in each replicate. The flugs were topped with silica gel (Fisher, #S684) and

sealed with parafilm to prevent any moisture from entering the vials. Vials were kept at 24°C on

12 hour day/night cycles. Survival was measured by counting flies hourly until the entire

population died. This was repeated once more with 50 flies per line per rearing condition in each

trial for a total of 100 flies with the exception for the *Δdpt* line which in total only had 30 flies

per condition in total across trials.

Starvation

Starvation survival assays were performed on dpt^{S69} and dpt^{S69R} adult males 4-7 days after

eclosion for axenically and conventionally reared flies. Ten males were placed into a vial with

autoclaved starvation media (1% agar). The 1% agar was used to starve flies of nutrition but not

desiccate them. Vials were kept at 24°C on 12-hour day/night cycles. Survival was measured by

counting surviving flies at three 8-hour intervals (8 am, 4 pm, and 12 am) until all flies died. This

was repeated twice more for a total of 3 trials with at least 40 flies per line per rearing condition

in each trial for a total of at least 120 flies.

16S Sequencing

Flies were reared in the lab in two distinct ways for 16S rRNA sequencing of conventionally

reared flies. First, flies were taken from vials of individual genotypes. This allows for any

moderate effects of fly genotype to equilibrate over time. Second, flies were taken from cages

that were started with heterozygous *diptericin* flies. These cages were started with the F1

progeny from crosses of dpt^{S69} and dpt^{S69R} flies and allowed to continue for 3 discrete

generations before flies were collected for 16S rRNA sequencing. This ensures that the

genotypes are exposed to the same microbes and any differences in microbiome are due to

genetic differences manifest in that generation.

We isolated DNA from individual flies using Gentra PureGene Tissue DNA Isolation Kit

(Qiagen #158388, Qiagen, Germantown, MD) following manufacturer's instructions. DNA

pellets were rehydrated in 10 μl DNA hydration solution. Flies from cage populations were

genotyped by using the dpt_cw primer pair (Table S1) followed by a restriction enzyme digest

using enzymes that cut DNA sequences for arginine (BccI, NEB # R0704S) or serine (AluI, NEB

R0137S).

DNA concentration was measured with Qubit fluorimeter (Invitrogen). Ten flies from each

inbred CRISPR genome edited line (5 females and 5 males from dpt^{S69}, dpt^{S69R}, dpt^{S69R-2} (a

second homozygous arginine line from CRISPR/Cas9 genome editing), and Δdpt (1 base pair

deletion) and ten flies of each genotype from the cage population (5 females and 5 males of SS,

SR, RR genotypes) were each brought to a DNA concentration of 5 ng/µl. Libraries were

prepped in accordance with the 16S Metagenomic Sequencing Library Preparation protocol from

Illumina for the 16S V3/V4 region. Sequencing was performed on the Illumina MiSeq platform

using v3 300bp paired end reads. Library preparation and sequencing was performed at the

University of Kansas Genome Sequencing Core (Lawrence, KS, USA).

Wild flies were collected from decaying apples and pears in an apple orchard in Kansas City, KS

(3341 N 139[th] St, Kansas City, KS 66109). Flies were immediately transported back to the lab

and sorted by species on CO_2 and frozen at -20°C. DNA was extracted from individual flies

using Gentra PureGene Cell & Tissue DNA Isolation Kit (Qiagen #158388). The samples were

tested for species (*D. simulans* vs. *D. melanogaster*)*, Wolbachia* status, and Dpt genotype using

primers listed in Table S1. Collections are summarized in Table S10. Libraries were prepped in

the same manner as the conventionally reared flies.

16S Bioinformatic Analysis

Demultiplexed reads were processed with QIIME2, v 2019.10 [88]. Primers were removed from

5' ends with Cutadapt using default parameters [89]. Reads were de-noised and trimmed for

quality with Divisive Amplicon Denoising Algorithm (DADA2) within the QIIME2

bioinformatics pipeline [90]. Forward and reverse reads were truncated at 280 bp and 245 bp

respectively. The remaining ASV table was exported from QIIME2 for further processing in R

[91]. Taxonomy was assigned to the ASV table using SILVA 16S rRNA gene reference

database, v.138 [92–94]. Reads assigned to Genus level *Wolbachia* (a *Drosophila* endosymbiont)

and Kingdom level Eukaryota were removed from further analysis. We also removed reads not

observed at least 3 times in at least 10 samples. Then, conventionally reared flies were rarefied to

17066 reads, while wild flies were rarefied to 10771 reads.

All statistical analysis of 16S data was performed in R (4.1.2) using the Phyloseq [95] package

and ggplot2 [96] package for visualization. The CRISPR and cage population data were analyzed

separately. For each population, alpha diversity was estimated using Shannon diversity in

Phyloseq using the estimate_richness() function. Bray-Curtis dissimilarity was calculated to look

at overall patterns of microbiome composition in Phyloseq using the ordinate() function.

Significance was determined at $\alpha=0.05$. Fixed effects models were fit with the package lme4 [97]

with the fixed effects genotype + sex + genotype*sex. When necessary, P-values were adjusted

for multiple comparisons using FDR correction method.

Statistical Analysis

R (version 4.1.2) was used to run statistical analyses. Survival data were plotted using the R

package survminer [98]. The analysis was performed using the Cox proportional-hazards

regression model in R [99]. For longevity, significance was determined using the model:

Lifespan~(Genotype*Sex)/Vial+Block. For gnotobiotic bacterial load, significance was

determined using the model: Colonies.mL~(Genotype)*Sex+Block.

Competing Interests

The authors state no competing interests.

Acknowledgments

We thank the Cider Hill Family Orchard for allowing us to collect wild flies on their property; S.

Macdonald lab for providing the modified 7198 and A4 lines; Kistie Brunsell for help with DNA

extractions of wild flies; and Elizabeth Everman for assistance with the feeding assay. The

University of Kansas Genome Sequencing Center Core (supported by NIH CMADP COBRE

P20-GM103638) performed 16s sequencing. SRM was supported by a University of Kansas Self

Graduate Fellowship, the work was supported by NIH R01-AI139154 to RLU.

References

1.   Miyauchi E, Shimokawa C, Steimle A, Desai MS, Ohno H. The impact of the gut microbiome
on extra-intestinal autoimmune diseases. Nat Rev Immunol. 2023;23: 9–23.
doi:10.1038/s41577-022-00727-y

2.   Wang M, Peng I-F, Li S, Hu X. Dysregulation of antimicrobial peptide expression
distinguishes Alzheimer's disease from normal aging. Aging. 2020;12: 690–706.
doi:10.18632/aging.102650

3.   Badinloo M, Nguyen E, Suh W, Alzahrani F, Castellanos J, Klichko VI, et al. Over-expression
of antimicrobial peptides contributes to aging through cytotoxic effects in Drosophila
tissues. Arch Insect Biochem Physiol. 2018;98: e21464. doi:10.1002/arch.21464

4.   Fajgenbaum DC, June CH. Cytokine Storm. N Engl J Med. 2020;383: 2255–2273.
doi:10.1056/NEJMra2026131

5.   Thaiss CA, Zmora N, Levy M, Elinav E. The microbiome and innate immunity. Nature.
2016;535: 65–74. doi:10.1038/nature18847

6.   Shi N, Li N, Duan X, Niu H. Interaction between the gut microbiome and mucosal immune
system. Mil Med Res. 2017;4: 14. doi:10.1186/s40779-017-0122-9

7.   Brown SP, Cornforth DM, Mideo N. Evolution of virulence in opportunistic pathogens:
generalism, plasticity, and control. Trends Microbiol. 2012;20: 336–342.
doi:10.1016/j.tim.2012.04.005

8.   Bagnicka E, Strzałkowska N, Jóźwik A, Krzyżewski J, Horbańczuk J, Zwierzchowski L.
Expression and polymorphism of defensins in farm animals. Acta Biochim Pol. 2010;57:
487–497.

9.   Pierce KR, Eichler T, Mosquera Vasquez C, Schwaderer AL, Simoni A, Creacy S, et al.
Ribonuclease 7 polymorphism rs1263872 reduces antimicrobial activity and associates
with pediatric urinary tract infections. J Clin Invest. 2021;131: e149807.
doi:10.1172/JCI149807

10.   Anderson JP, Gleason CA, Foley RC, Thrall PH, Burdon JB, Singh KB, et al. Plants versus
pathogens: an evolutionary arms race. Funct Plant Biol. 2010;37: 499–512.
doi:10.1071/FP09304

11.   Carrillo-Bustamante P, Keşmir C, de Boer RJ. A Coevolutionary Arms Race between Hosts
and Viruses Drives Polymorphism and Polygenicity of NK Cell Receptors. Mol Biol Evol.
2015;32: 2149–2160. doi:10.1093/molbev/msv096

12.   Marques JT, Carthew RW. A call to arms: coevolution of animal viruses and host innate
immune responses. Trends Genet. 2007;23: 359–364. doi:10.1016/j.tig.2007.04.004

13.   Liston A, Humblet-Baron S, Duffy D, Goris A. Human immune diversity: from evolution to
modernity. Nat Immunol. 2021;22: 1479–1489. doi:10.1038/s41590-021-01058-1

14.   Koenig D, Hagmann J, Li R, Bemm F, Slotte T, Neuffer B, et al. Long-term balancing
selection drives evolution of immunity genes in Capsella. Przeworski M, Baldwin IT, Gao Z,
editors. eLife. 2019;8: e43606. doi:10.7554/eLife.43606

15.   Chapman JR, Hill T, Unckless RL. Balancing Selection Drives the Maintenance of Genetic
Variation in Drosophila Antimicrobial Peptides. Genome Biol Evol. 2019;11: 2691–2701.
doi:10.1093/gbe/evz191

16.   Aguilar A, Roemer G, Debenham S, Binns M, Garcelon D, Wayne RK. High MHC diversity
maintained by balancing selection in an otherwise genetically monomorphic mammal.
Proc Natl Acad Sci. 2004;101: 3490–3494. doi:10.1073/pnas.0306582101

17.   Ferrer-Admetlla A, Bosch E, Sikora M, Marquès-Bonet T, Ramírez-Soriano A, Muntasell A,
et al. Balancing Selection Is the Main Force Shaping the Evolution of Innate Immunity
Genes1. J Immunol. 2008;181: 1315–1322. doi:10.4049/jimmunol.181.2.1315

18.   Tennessen JA, Blouin MS. Balancing Selection at a Frog Antimicrobial Peptide Locus:
Fluctuating Immune Effector Alleles? Mol Biol Evol. 2008;25: 2669–2680.
doi:10.1093/molbev/msn208

19.   Unckless RL, Howick VM, Lazzaro BP. Convergent balancing selection on an antimicrobial
peptide in Drosophila. Curr Biol CB. 2016;26: 257–262. doi:10.1016/j.cub.2015.11.063

20.   Chapman JR, Hellgren O, Helin AS, Kraus RHS, Cromie RL, Waldenström J. The Evolution of
Innate Immune Genes: Purifying and Balancing Selection on β-Defensins in Waterfowl. Mol
Biol Evol. 2016;33: 3075–3087. doi:10.1093/molbev/msw167

21.   Padhi A, Verghese B. Molecular diversity and evolution of myticin-C antimicrobial peptide
variants in the Mediterranean mussel, Mytilus galloprovincialis. Peptides. 2008;29: 1094–
1101. doi:10.1016/j.peptides.2008.03.007

22.   Sommer S. The importance of immune gene variability (MHC) in evolutionary ecology and
conservation. Front Zool. 2005;2: 16. doi:10.1186/1742-9994-2-16

23.   Williams TN, Mwangi TW, Wambua S, Alexander ND, Kortok M, Snow RW, et al. Sickle cell
trait and the risk of Plasmodium falciparum malaria and other childhood diseases. J Infect
Dis. 2005;192: 178–186. doi:10.1086/430744

24.   Ryu J-H, Ha E-M, Lee W-J. Innate immunity and gut–microbe mutualism in Drosophila. Dev
Comp Immunol. 2010;34: 369–376. doi:10.1016/j.dci.2009.11.010

25. Lhocine N, Ribeiro PS, Buchon N, Wepf A, Wilson R, Tenev T, et al. PIMS Modulates
Immune Tolerance by Negatively Regulating Drosophila Innate Immune Signaling. Cell Host
Microbe. 2008;4: 147–158. doi:10.1016/j.chom.2008.07.004

26. Bosco-Drayon V, Poidevin M, Boneca IG, Narbonne-Reveau K, Royet J, Charroux B.
Peptidoglycan Sensing by the Receptor PGRP-LE in the Drosophila Gut Induces Immune
Responses to Infectious Bacteria and Tolerance to Microbiota. Cell Host Microbe. 2012;12:
153–165. doi:10.1016/j.chom.2012.06.002

27. Broderick NA, Buchon N, Lemaitre B. Microbiota-Induced Changes in Drosophila
melanogaster Host Gene Expression and Gut Morphology. mBio. 2014;5.
doi:10.1128/mBio.01117-14

28. Hanson MA, Lemaitre B. New insights on Drosophila antimicrobial peptide function in host
defense and beyond. Curr Opin Immunol. 2020;62: 22–30. doi:10.1016/j.coi.2019.11.008

29. Marra A, Hanson MA, Kondo S, Erkosar B, Lemaitre B. Drosophila Antimicrobial Peptides
and Lysozymes Regulate Gut Microbiota Composition and Abundance. mBio. 2021;12:
e0082421. doi:10.1128/mBio.00824-21

30. Ong PY, Ohtake T, Brandt C, Strickland I, Boguniewicz M, Ganz T, et al. Endogenous
antimicrobial peptides and skin infections in atopic dermatitis. N Engl J Med. 2002;347:
1151–1160. doi:10.1056/NEJMoa021481

31. Rieg S, Steffen H, Seeber S, Humeny A, Kalbacher H, Dietz K, et al. Deficiency of dermcidin-
derived antimicrobial peptides in sweat of patients with atopic dermatitis correlates with
an impaired innate defense of human skin in vivo. J Immunol Baltim Md 1950. 2005;174:
8003–8010. doi:10.4049/jimmunol.174.12.8003

32. Buchon N, Silverman N, Cherry S. Immunity in Drosophila melanogaster — from microbial
recognition to whole-organism physiology. Nat Rev Immunol. 2014;14: 796–810.
doi:10.1038/nri3763

33. Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, et al. Evolution of genes
and genomes on the Drosophila phylogeny. Nature. 2007;450: 203–218.
doi:10.1038/nature06341

34. Jiggins FM, Kim KW. A screen for immunity genes evolving under positive selection in
Drosophila. J Evol Biol. 2007;20: 965–970. doi:10.1111/j.1420-9101.2007.01305.x

35. Sackton TB, Lazzaro BP, Schlenke TA, Evans JD, Hultmark D, Clark AG. Dynamic evolution of
the innate immune system in Drosophila. Nat Genet. 2007;39: 1461–1468.
doi:10.1038/ng.2007.60

36. Obbard DJ, Welch JJ, Kim K-W, Jiggins FM. Quantifying Adaptive Evolution in the
Drosophila Immune System. PLoS Genet. 2009;5. doi:10.1371/journal.pgen.1000698

37.   Comeron JM. Background Selection as Baseline for Nucleotide Variation across the
834   Drosophila Genome. PLOS Genet. 2014;10: e1004434. doi:10.1371/journal.pgen.1004434

835   38.   Unckless RL, Lazzaro BP. The potential for adaptive maintenance of diversity in insect
836   antimicrobial peptides. Philos Trans R Soc B Biol Sci. 2016;371. doi:10.1098/rstb.2015.0291

837   39.   Perlmutter JI, Chapman JR, Wilkinson MC, Nevarez-Saenz I, Unckless RL. A single amino
838   acid polymorphism in natural Metchnikowin alleles of Drosophila results in systemic
839   immunity and life history tradeoffs. PLOS Genet. 2024;20: e1011155.
840   doi:10.1371/journal.pgen.1011155

841   40.   Lazzaro BP, Sceurman BK, Clark AG. Genetic Basis of Natural Variation in D. melanogaster
842   Antibacterial Immunity. Science. 2004;303: 1873–1876. doi:10.1126/science.1092447

843   41.   Smith BR, Patch KB, Gupta A, Knoles EM, Unckless RL. The genetic basis of variation in
844   immune defense against Lysinibacillus fusiformis infection in Drosophila melanogaster.
845   PLOS Pathog. 2023;19: e1010934. doi:10.1371/journal.ppat.1010934

846   42.   Hanson MA, Grollmus L, Lemaitre B. Ecology-relevant bacteria drive the evolution of host
847   antimicrobial peptides in Drosophila. Science. 381: eadg5725.
848   doi:10.1126/science.adg5725

849   43.   Tanji T, Hu X, Weber ANR, Ip YT. Toll and IMD Pathways Synergistically Activate an Innate
850   Immune Response in Drosophila melanogaster. Mol Cell Biol. 2007;27: 4578–4588.
851   doi:10.1128/MCB.01814-06

852   44.   Broderick NA. Friend, foe or food? Recognition and the role of antimicrobial peptides in
853   gut immunity and Drosophila–microbe interactions. Philos Trans R Soc B Biol Sci.
854   2016;371. doi:10.1098/rstb.2015.0295

855   45.   Hanson MA, Dostálová A, Ceroni C, Poidevin M, Kondo S, Lemaitre B. Synergy and
856   remarkable specificity of antimicrobial peptides in vivo using a systematic knockout
857   approach. MacPherson AJ, Garrett WS, Hornef M, Hooper LV, editors. eLife. 2019;8:
858   e44341. doi:10.7554/eLife.44341

859   46.   Arias-Rojas A, Iatsenko I. The Role of Microbiota in Drosophila melanogaster Aging. Front
860   Aging. 2022;3. Available: https://www.frontiersin.org/articles/10.3389/fragi.2022.909509

861   47.   Badinloo M, Nguyen E, Suh W, Alzahrani F, Castellanos J, Klichko VI, et al. Overexpression
862   of antimicrobial peptides contributes to aging through cytotoxic effects in Drosophila
863   tissues. Arch Insect Biochem Physiol. 2018;98: e21464. doi:10.1002/arch.21464

864   48.   Biswajit Shit, Arun Prakash, Saubhik Sarkar, Pedro F. Vale, Imroze Khan. Ageing leads to
865   nonspecific antimicrobial peptide responses in Drosophila melanogaster.
866   bioRxiv. 2022; 2022.06.25.497570. doi:10.1101/2022.06.25.497570

867   49.   Hanson MA, Lemaitre B. Antimicrobial peptides do not directly contribute to aging in
868   Drosophila, but improve lifespan by preventing dysbiosis. Dis Model Mech. 2023;16:
869   dmm049965. doi:10.1242/dmm.049965

870   50.   Duneau DF, Kondolf HC, Im JH, Ortiz GA, Chow C, Fox MA, et al. The Toll pathway underlies
871   host sexual dimorphism in resistance to both Gram-negative and Gram-positive bacteria in
872   mated Drosophila. BMC Biol. 2017;15: 124. doi:10.1186/s12915-017-0466-3

873   51.   Ding R, Goh W-R, Wu R, Yue X, Luo X, Khine WWT, et al. Revisit gut microbiota and its
874   impact on human health and disease. J Food Drug Anal. 2019;27: 623–631.
875   doi:10.1016/j.jfda.2018.12.012

876   52.   Ludington WB, Ja WW. Drosophila as a model for the gut microbiome. PLOS Pathog.
877   2020;16: e1008398. doi:10.1371/journal.ppat.1008398

878   53.   Erkosar B, Leulier F. Transient adult microbiota, gut homeostasis and longevity: Novel
879   insights from the Drosophila model. FEBS Lett. 2014;588: 4250–4257.
880   doi:10.1016/j.febslet.2014.06.041

881   54.   Macke E, Tasiemski A, Massol F, Callens M, Decaestecker E. Life history and eco-
882   evolutionary dynamics in light of the gut microbiota. Oikos. 2017;126: 508–531.
883   doi:10.1111/oik.03900

884   55.   Lints FA, Bourgois M, Delalieux A, Stoll J, Lints CV. Does the Female Life Span Exceed That
885   of the Male: A Study in Drosophila melanogaster. Gerontology. 2009;29: 336–352.
886   doi:10.1159/000213136

887   56.   Wu Q, Yu G, Park SJ, Gao Y, Ja WW, Yang M. Excreta Quantification (EX-Q) for Longitudinal
888   Measurements of Food Intake in Drosophila. iScience. 2020;23: 100776.
889   doi:10.1016/j.isci.2019.100776

890   57.   Wong CNA, Ng P, Douglas AE. Low-diversity bacterial community in the gut of the fruitfly
891   Drosophila melanogaster. Environ Microbiol. 2011;13: 1889–1900. doi:10.1111/j.1462-
892   2920.2011.02511.x

893   58.   Bost A, Martinson VG, Franzenburg S, Adair KL, Albasi A, Wells MT, et al. Functional
894   variation in the gut microbiome of wild Drosophila populations. Mol Ecol. 2018;27: 2834–
895   2845. doi:10.1111/mec.14728

896   59.   Giannakou ME, Partridge L. Role of insulin-like signalling in Drosophila lifespan. Trends
897   Biochem Sci. 2007;32: 180–188. doi:10.1016/j.tibs.2007.02.007

898   60.   Biteau B, Karpac J, Hwangbo D, Jasper H. Regulation of Drosophila lifespan by JNK
899   signaling. Exp Gerontol. 2011;46: 349–354. doi:10.1016/j.exger.2010.11.003

61.   Viney ME, Riley EM, Buchanan KL. Optimal immune responses: immunocompetence
revisited. Trends Ecol Evol. 2005;20: 665–669. doi:10.1016/j.tree.2005.10.003

62.   Sadd BM, Siva-Jothy MT. Self-harm caused by an insect's innate immunity. Proc R Soc B
Biol Sci. 2006;273: 2571–2574. doi:10.1098/rspb.2006.3574

63.   Lin Y-R, Parikh H, Park Y. Stress resistance and lifespan enhanced by downregulation of
antimicrobial peptide genes in the Imd pathway. Aging. 2018;10: 622–631.
doi:10.18632/aging.101417

64.   Gibbs AG. Water balance in desert Drosophila: lessons from non-charismatic microfauna.
Comp Biochem Physiol A Mol Integr Physiol. 2002;133: 781–789. doi:10.1016/S1095-
6433(02)00208-8

65.   Wang R-A. Life history tradeoffs of pathogens and the treatment principle of
antibiogenesis. Genes Dis. 2017;4: 154–158. doi:10.1016/j.gendis.2017.07.004

66.   Khor S, Cai D. Control of lifespan and survival by Drosophila NF-κB signaling through
neuroendocrine cells and neuroblasts. Aging. 2020;12: 24604–24622.
doi:10.18632/aging.104196

67.   Wong AC-N, Chaston JM, Douglas AE. The inconstant gut microbiota of Drosophila species
revealed by 16S rRNA gene analysis. ISME J. 2013;7: 1922–1932.
doi:10.1038/ismej.2013.86

68.   Fisher RA. The genetical theory of natural selection. Oxford, England: Clarendon Press;
1930. pp. xiv, 272. doi:10.5962/bhl.title.27468

69.   Ford EB, De Beer G, Huxley JS. Polymorphism and taxonomy. [London]: Oxford University
Press; 1940.

70.   Dobzhansky T. GENETICS OF NATURAL POPULATIONS. XIX. ORIGIN OF HETEROSIS
THROUGH NATURAL SELECTION IN POPULATIONS OF DROSOPHILA PSEUDOOBSCURA.
Genetics. 1950;35: 288–302. doi:10.1093/genetics/35.3.288

71.   Cain AJ, Sheppard PM. NATURAL SELECTION IN CEPAEA. Genetics. 1954;39: 89–116.
doi:10.1093/genetics/39.1.89

72.   Dobzhansky T. Genetics of the Evolutionary Process. Columbia University Press; 1970.

73.   Hedrick PW, Thomson G. EVIDENCE FOR BALANCING SELECTION AT HLA. Genetics.
1983;104: 449–456. doi:10.1093/genetics/104.3.449

74.   Subramaniam B, Rausher MD. Balancing Selection on a Floral Polymorphism. Evolution.
2000;54: 691–695. doi:10.1111/j.0014-3820.2000.tb00070.x

75.   Koskella B, Lively CM. Evidence for Negative Frequency-Dependent Selection During
933   Experimental Coevolution of a Freshwater Snail and a Sterilizing Trematode. Evolution.
934   2009;63: 2213–2221. doi:10.1111/j.1558-5646.2009.00711.x

935   76.   Leffler EM, Gao Z, Pfeifer S, Ségurel L, Auton A, Venn O, et al. Multiple Instances of Ancient
936   Balancing Selection Shared Between Humans and Chimpanzees. Science. 2013;339: 1578–
937   1582. doi:10.1126/science.1234070

938   77.   Delph LF, Kelly JK. On the importance of balancing selection in plants. New Phytol.
939   2014;201: 45–56. doi:10.1111/nph.12441

940   78.   Lee D, Zdraljevic S, Stevens L, Wang Y, Tanny RE, Crombie TA, et al. Balancing selection
941   maintains hyper-divergent haplotypes in Caenorhabditis elegans. Nat Ecol Evol. 2021;5:
942   794–807. doi:10.1038/s41559-021-01435-x

943   79.   Piel FB, Patil AP, Howes RE, Nyangiri OA, Gething PW, Williams TN, et al. Global
944   distribution of the sickle cell gene and geographical confirmation of the malaria
945   hypothesis. Nat Commun. 2010;1: 104. doi:10.1038/ncomms1104

946   80.   Shianiou G, Teloni S, Apidianakis Y. Intestinal Immune Deficiency and Juvenile Hormone
947   Signaling Mediate a Metabolic Trade-off in Adult Drosophila Females. Metabolites.
948   2023;13: 340. doi:10.3390/metabo13030340

949   81.   Connallon T, Clark AG. Balancing Selection in Species with Separate Sexes: Insights from
950   Fisher's Geometric Model. Genetics. 2014;197: 991–1006.
951   doi:10.1534/genetics.114.165605

952   82.   Hanson MA, Lemaitre B, Unckless RL. Dynamic Evolution of Antimicrobial Peptides
953   Underscores Trade-Offs Between Immunity and Ecological Fitness. Front Immunol.
954   2019;10. Available: https://www.frontiersin.org/articles/10.3389/fimmu.2019.02620

955   83.   Hanson MA, Lemaitre B, Unckless RL. Dynamic Evolution of Antimicrobial Peptides
956   Underscores Trade-Offs Between Immunity and Ecological Fitness. Front Immunol.
957   2019;10. Available: https://www.frontiersin.org/articles/10.3389/fimmu.2019.02620

958   84.   King EG, Merkes CM, McNeil CL, Hoofer SR, Sen S, Broman KW, et al. Genetic dissection of
959   a model complex trait using the Drosophila Synthetic Population Resource. Genome Res.
960   2012;22: 1558–1566. doi:10.1101/gr.134031.111

961   85.   Juneja P, Lazzaro BP. Providencia sneebia sp. nov. and Providencia burhodogranariea sp.
962   nov., isolated from wild Drosophila melanogaster. Int J Syst Evol Microbiol. 2009;59: 1108–
963   1111. doi:10.1099/ijs.0.000117-0

964   86.   Smith BR, Unckless RL. Draft Genome Sequence of Lysinibacillus fusiformis Strain Juneja, a
965   Laboratory-Derived Pathogen of Drosophila melanogaster. Genome Announc. 2018;6:
966   e01571-17. doi:10.1128/genomeA.01571-17

87. Khalil S, Jacobson E, Chambers MC, Lazzaro BP. Systemic Bacterial Infection and Immune
Defense Phenotypes in Drosophila Melanogaster. JoVE J Vis Exp. 2015; e52613.
doi:10.3791/52613

88. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible,
interactive, scalable and extensible microbiome data science using QIIME 2. Nat
Biotechnol. 2019;37: 852–857. doi:10.1038/s41587-019-0209-9

89. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal. 2011;17: 10–12. doi:10.14806/ej.17.1.200

90. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13: 581–
583. doi:10.1038/nmeth.3869

91. R Core Development Team. R: A language and environment for statistical computing.
2021.

92. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA
gene database project: improved data processing and web-based tools. Nucleic Acids Res.
2013;41: D590–D596. doi:10.1093/nar/gks1219

93. Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, et al. The SILVA and "All-
species Living Tree Project (LTP)" taxonomic frameworks. Nucleic Acids Res. 2014;42:
D643–D648. doi:10.1093/nar/gkt1209

94. Glöckner FO, Yilmaz P, Quast C, Gerken J, Beccati A, Ciuprina A, et al. 25 years of serving
the community with ribosomal RNA gene reference databases and tools. J Biotechnol.
2017;261: 169–176. doi:10.1016/j.jbiotec.2017.06.1198

95. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. PLOS ONE. 2013;8: e61217.
doi:10.1371/journal.pone.0061217

96. Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al. Welcome to the
Tidyverse. J Open Source Softw. 2019;4: 1686. doi:10.21105/joss.01686

97. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. J
Stat Softw. 2015;67: 1–48. doi:10.18637/jss.v067.i01

98. Kassambara A, Kosinski M, Biecek P, Fabian S. survminer: Drawing Survival Curves using
"ggplot2." 2021. Available: https://CRAN.R-project.org/package=survminer

99. Therneau TM, until 2009) TL (original S->R port and R maintainer, Elizabeth A, Cynthia C.
survival: Survival Analysis. 2022. Available: https://CRAN.R-project.org/package=survival

1000

1001

1002 **Supplemental tables**

1003 **Table S1. Oligonucleotide sequences used in this study.**
1004

Primer name	Primer sequence	Use	Reference
wsp-81F	TGGTCCAATAAGTGATGAAGAAAC	*Wolbachia* status	Braig et al. 1998
Wsp-691R	AAAAATTAAACGCTACTCCA	*Wolbachia* status	Braig et al. 1998
Actin79B-F	CGTCTACCAGTCCATCATGA	Species identification	
Actin79B-R	AGACGGAGTACTTGCGCTC	Species identification	
Dpt_cw_F	GCCTATAAAAGAGCATCGAA	*Diptericin*	Clark and Wang, 1997
Dpt_cw_R	GCTAGACTCGGATACCAATCGAG	*Diptericin*	Clark and Wang, 1997
Dpt_gRNA18_S	CTTCGATGGCAACTCAGAACCGACG	gRNA	This work
Dpt_gRNA18_A	AAACGCTCGGTTCTGAGTTCGGATC	gRNA	This work
ssODN	CGCCACGAGATTGGACTGAATGGAGG ATATGGACAGCACTTGGGAGGACCAT ACGGCAACTCAGAACCGAGATGGAA AGTGGGAAGCACCTACACCTACAGA TTTCCGAATTTCTAAGCT	ssODN	This work

1005
1006
1007
1008

Table S2. Cox proportional hazard pairwise significance p-values for systemic infection. P-values less than 0.05 are underlined.

P. rettgeri	imd-	Δdpt	dptS69R
Δdpt	4.07E-02	-	-
dptS69R	4.54E-01	0.26329	-
dptS69	1.20E-04	1.10E-09	1.30E-06

S. succinus	imd-	Δdpt	dptS69R
Δdpt	0.4994	-	-
dptS69R	0.0672	0.0023	-
dptS69	0.4994	0.9828	0.012

E. faecalis	imd-	Δdpt	dptS69R
Δdpt	0.0264	-	-
dptS69R	0.0264	1.00E-06	-
dptS69	0.2969	0.3405	2.10E-03

L. fusiformis	imd-	Δdpt	dptS69R
Δdpt	0.435	-	-
dptS69R	0.048	0.22	-
dptS69	0.435	0.927	0.19

L. lactis	imd-	Δdpt	dptS69R
Δdpt	0.7515	-	-
dptS69R	0.0474	0.0474	-
dptS69	0.1966	0.0858	0.0015

S. marcescens	imd-	Δdpt	dptS69R
Δdpt	<2e-16	-	-
dptS69R	<2e-16	0.1151	-
dptS69	<2e-16	0.054	0.0042

1014 **Table S3. Cox proportion hazard ratio pairwise p-values for systemic infections with males and females.** P-
1015 values less than 0.05 are underlined.
1016

P. rettgeri							
	F-*imd-*	M-*imd-*	F-*Δdpt*	M-*Δdpt*	F- *dpt^{S69R}*	M- *dpt^{S69R}*	F- *dpt^{S69}*
M-*imd-*	0.2560	-	-	-	-	-	-
F-*Δdpt*	0.1592	0.8278	-	-	-	-	-
M-*Δdpt*	0.2356	0.9763	0.8278	-	-	-	-
F- *dpt^{S69R}*	0.9763	0.2981	0.1933	0.2560	-	-	-
M-*d dpt^{S69R}*	<u>0.0144</u>	0.1124	0.1510	0.1124	<u>0.0152</u>	-	-
F- *dpt^{S69}*	<u>5.60E-05</u>	<u>0.0014</u>	<u>0.0014</u>	<u>0.0008</u>	<u>8.80E-05</u>	0.3598	-
M- *dpt^{S69}*	<u>3.70E-06</u>	<u>9.60E-05</u>	<u>9.60E-05</u>	<u>7.70E-05</u>	<u>6.50E-06</u>	0.1639	0.5982

E. faecalis							
	F-*imd-*	M-*imd-*	F-*Δdpt*	M-*Δdpt*	F- *dpt^{S69R}*	M- *dpt^{S69R}*	F- *dpt^{S69}*
M-*imd-*	<u>0.0304</u>	-	-	-	-	-	-
F-*Δdpt*	<u>0.0036</u>	0.8954	-	-	-	-	-
M-*Δdpt*	<u>0.0012</u>	0.7414	0.7414	-	-	-	-
F- *dpt^{S69R}*	0.7414	0.1433	<u>0.0326</u>	<u>0.0158</u>	-	-	-
M- *dpt^{S69R}*	<u>0.0318</u>	0.9686	0.8893	0.7414	0.1566	-	-
F- *dpt^{S69}*	<u>0.0326</u>	0.9686	0.8779	0.7414	0.1701	0.9686	-
M- *dpt^{S69}*	<u>0.0114</u>	0.753	0.8664	0.9686	0.0511	0.7414	0.7414

L. plantarum							
	F-*imd-*	M-*imd-*	F-*Δdpt*	M-*Δdpt*	F- *dpt^{S69R}*	M- *dpt^{S69R}*	F- *dpt^{S69}*
M-*imd-*	<u><2e-16</u>	-	-	-	-	-	-
F-*Δdpt*	<u>3.80E-08</u>	<u><2e-16</u>	-	-	-	-	-
M-*Δdpt*	<u><2e-16</u>	<u>8.10E-09</u>	<u><2e-16</u>	-	-	-	-
F- *dpt^{S69}*	<u><2e-16</u>	<u>3.20E-09</u>	<u>7.60E-15</u>	0.734	-	-	-
M- *dpt^{S69R}*	<u><2e-16</u>	<u>1.80E-12</u>	<u>8.30E-15</u>	0.372	0.197	-	-
F- *dpt^{S69}*	<u><2e-16</u>	<u>6.00E-15</u>	<u>1.00E-10</u>	0.08	0.056	0.548	-
M- *dpt^{S69}*	<u><2e-16</u>	<u>5.40E-07</u>	<u><2e-16</u>	0.364	0.662	0.056	<u>0.013</u>

1017
1018

Table S4. P-values from Tukey HSD for axenic and gnotobiotic longevity. P-values less than 0.05 underlined.

Axenic							
	F-*imd-*	F-*Δdpt*	F-*dpt*^{S69R}	F-*dpt*^{S69}	M-*imd-*	M-*Δdpt*	M-*dpt*^{S69R}
F-*Δdpt*	1	-	-	-	-	-	-
F-*dpt*^{S69R}	<u>6.00E-04</u>	<u><2e-16</u>	-	-	-	-	-
F-*dpt*^{S69}	0.9447	0.8051	0.063	-	-	-	-
M-*imd-*	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	-	-	-
M-*Δdpt*	<u>1.00E-04</u>	<u><2e-16</u>	1	<u>0.0492</u>	<u><2e-16</u>	-	-
M-*dpt*^{S69R}	0.1713	0.0519	0.7467	0.8822	<u><2e-16</u>	0.8133	-
M-*dpt*^{S69}	<u>1.00E-04</u>	<u><2e-16</u>	1	<u>0.0218</u>	<u><2e-16</u>	0.9947	0.5114

L. plantarum							
	F-*imd-*	F-*Δdpt*	F-*dpt*^{S69R}	F-*dpt*^{S69}	M-*imd-*	M-*Δdpt*	M-*dpt*^{S69R}
F-*Δdpt*	<u>0.0064</u>	-	-	-	-	-	-
F-*dpt*^{S69R}	<u><2e-16</u>	<u><2e-16</u>	-	-	-	-	-
F-*dpt*^{S69}	<u><2e-16</u>	<u><2e-16</u>	0.185	-	-	-	-
M-*imd-*	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	-	-	-
M-*Δdpt*	<u><2e-16</u>	<u><2e-16</u>	0.9992	0.2479	<u><2e-16</u>	-	-
M-*dpt*^{S69R}	<u><2e-16</u>	<u><2e-16</u>	1	0.2568	<u><2e-16</u>	0.9999	-
M-*dpt*^{S69}	<u><2e-16</u>	<u><2e-16</u>	0.9531	<u>0.0056</u>	<u><2e-16</u>	0.5593	0.9198

P. rettgeri							
	F-*imd-*	F-*Δdpt*	F-*dpt*^{S69R}	F-*dpt*^{S69}	M-*imd-*	M-*Δdpt*	M-*dpt*^{S69R}
F-*Δdpt*	<u><2e-16</u>	-	-	-	-	-	-
F-*dpt*^{S69R}	<u>1.00E-04</u>	0.1564	-	-	-	-	-
F-*dpt*^{S69}	<u><2e-16</u>	0.5911	0.9977	-	-	-	-
M-*imd-*	<u><2e-16</u>	<u>1.12E-02</u>	<u><2e-16</u>	<u>1.00E-04</u>	-	-	-
M-*Δdpt*	<u>2.34E-02</u>	<u><2e-16</u>	4.81E-01	9.95E-02	<u><2e-16</u>	-	-
M-*dpt*^{S69R}	<u>0.0013</u>	<u>3.63E-02</u>	0.9997	0.9434	<u><2e-16</u>	0.8429	-
M-*dpt*^{S69}	0.0938	<u>1.00E-04</u>	0.6415	0.2208	<u><2e-16</u>	1	0.9079

L. brevis							
	F-*imd-*	F-*Δdpt*	F-*dpt*^{S69R}	F-*dpt*^{S69}	M-*imd-*	M-*Δdpt*	M-*dpt*^{S69R}
F-*Δdpt*	<u><2e-16</u>	-	-	-	-	-	-
F-*dpt*^{S69R}	<u><2e-16</u>	<u>0.0348</u>	-	-	-	-	-
F-*dpt*^{S69}	<u>4.00E-04</u>	<u>9.00E-04</u>	0.9874	-	-	-	-
M-*imd-*	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	-	-	-
M-*Δdpt*	<u>1.75E-02</u>	<u><2e-16</u>	1.18E-01	6.87E-01	<u><2e-16</u>	-	-
M-*dpt*^{S69R}	<u>2.06E-02</u>	<u><2e-16</u>	0.5139	9.63E-01	<u><2e-16</u>	0.9998	-
M-*dpt*^{S69}	4.02E-01	<u><2e-16</u>	<u>0.0333</u>	2.97E-01	<u><2e-16</u>	0.9803	0.9241

A. tropicalis							
	F-*imd-*	F-*Δdpt*	F-*dpt*^{S69R}	F-*dpt*^{S69}	M-*imd-*	M-*Δdpt*	M-*dpt*^{S69R}
F-*Δdpt*	<u><2e-16</u>	-	-	-	-	-	-
F-*dpt*^{S69R}	0.2941	<u><2e-16</u>	-	-	-	-	-
F-*dpt*^{S69}	<u><2e-16</u>	0.9003	<u><2e-16</u>	-	-	-	-
M-*imd-*	<u>0.0016</u>	<u><2e-16</u>	0.6689	<u>0.0255</u>	-	-	-
M-*Δdpt*	<u>0.0017</u>	<u><2e-16</u>	0.8677	<u>4.00E-04</u>	0.9987	-	-
M-*dpt*^{S69R}	0.9522	<u><2e-16</u>	0.9472	<u><2e-16</u>	0.0876	0.1404	-
M-*dptS69*	0.1403	<u><2e-16</u>	1	<u>2.00E-04</u>	0.8812	0.9821	0.8103

L. plantarum + P. rettgeri							
	F-*imd-*	F-*Δdpt*	F-*dpt*^{S69R}	F-*dpt*^{S69}	M-*imd-*	M-*Δdpt*	M-*dpt*^{S69R}
F-*Δdpt*	0.1291	-	-	-	-	-	-
F-*dpt*^{S69R}	<u><2e-16</u>	<u><2e-16</u>	-	-	-	-	-
F-*dpt*^{S69}	<u><2e-16</u>	<u><2e-16</u>	0.3736	-	-	-	-
M-*imd-*	<u><2e-16</u>	<u><2e-16</u>	<u>9.00E-04</u>	<u><2e-16</u>	-	-	-
M-*Δdpt*	<u><2e-16</u>	<u><2e-16</u>	<u>0.0176</u>	0.9872	<u><2e-16</u>	-	-
M-*dpt*^{S69R}	<u><2e-16</u>	<u><2e-16</u>	0.998	0.807	<u>1.00E-04</u>	0.1564	-
M-*dpt*^{S69}	<u><2e-16</u>	<u><2e-16</u>	0.1446	0.9998	<u><2e-16</u>	1	0.5051

L. plantarum + A. tropicalis							
	F-*imd-*	F-*Δdpt*	F-*dpt*^{S69R}	F-*dpt*^{S69}	M-*imd-*	M-*Δdpt*	M-*dpt*^{S69R}
F-*Δdpt*	<u>0.008</u>	-	-	-	-	-	-
F-*dpt*^{S69R}	<u><2e-16</u>	<u><2e-16</u>	-	-	-	-	-
F-*dpt*^{S69}	<u><2e-16</u>	<u><2e-16</u>	0.0632	-	-	-	-
M-*imd-*	<u><2e-16</u>	<u><2e-16</u>	0.9998	0.1957	-	-	-
M-*Δdpt*	<u><2e-16</u>	<u><2e-16</u>	0.9998	0.058	1	-	-
M-*dpt*^{S69R}	<u><2e-16</u>	<u><2e-16</u>	0.8864	0.7167	0.9893	0.9573	-
M-*dpt*^{S69}	<u><2e-16</u>	<u><2e-16</u>	0.6101	0.944	0.8792	0.7135	0.9997

Table S5. P-values from Tukey HSD for bacteria load. P-values less than 0.05 are underlined.

	F-*imd-*	F-*Δdpt*	F-*dptS69R*	F-*dptS69*	M-*imd-*	M-*Δdpt*	M-*dptS69R*
Day 2 L. plantarum							
F-*Δdpt*	0.8875	NA	NA	NA	NA	NA	NA
F-*dptS69R*	0.9595	0.9999	NA	NA	NA	NA	NA
F-*dptS69*	0.4774	0.7296	0.6348	NA	NA	NA	NA
M-*imd-*	9.66E-01	1.00E+00	1	0.5932	NA	NA	NA
M-*Δdpt*	0.2644	0.1289	1.53E-01	0.9983	0.1308	NA	NA
M-*dptS69R*	0.8632	1	9.99E-01	0.924	0.9982	0.4745	NA
M-*dptS69*	0.8237	0.9999	9.96E-01	0.9628	0.9928	0.5926	1
Day 10 L. plantarum							
F-*Δdpt*	0.8875	NA	NA	NA	NA	NA	NA
F-*dptS69R*	0.9595	0.9999	NA	NA	NA	NA	NA
F-*dptS69*	0.4774	0.7296	0.6348	NA	NA	NA	NA
M-*imd-*	9.66E-01	1.00E+00	1	0.5932	NA	NA	NA
M-*Δdpt*	0.2644	0.1289	1.53E-01	0.9983	0.1308	NA	NA
M-*dptS69R*	0.8632	1	9.99E-01	0.924	0.9982	0.4745	NA
M-*dptS69*	0.8237	0.9999	9.96E-01	0.9628	0.9928	0.5926	1
Day 20 L. plantarum (no imd females)							
F-*Δdpt*	NA	NA	NA	NA	NA	NA	NA
F-*dptS69R*	NA	0.9978	NA	NA	NA	NA	NA
F-*dptS69*	NA	1	0.9943	NA	NA	NA	NA
M-*imd-*	NA	0.9998	1	0.9991	NA	NA	NA
M-*Δdpt*	NA	9.35E-02	<u>0.0314</u>	0.2254	0.0553	NA	NA
M-*dptS69R*	NA	1.00E+00	0.9807	1	0.9952	0.3207	NA
M-*dptS69*	NA	3.40E-01	0.1438	0.5173	0.2122	1	0.6305
Day 2 P. rettgeri							
F-*Δdpt*	<u>0.0406</u>	NA	NA	NA	NA	NA	NA
F-*dptS69R*	0.5572	0.9645	NA	NA	NA	NA	NA
F-*dptS69*	<u>0.0234</u>	0.9952	0.7777	NA	NA	NA	NA
M-*imd-*	0.5805	0.9565	1	0.7576	NA	NA	NA
M-*Δdpt*	<u>2.10E-03</u>	0.912	0.4482	1	0.4228	NA	NA
M-*dptS69R*	<u>1.00E-03</u>	0.5392	0.1887	0.9679	0.1757	0.9842	NA
M-*dptS69*	<u>4.00E-04</u>	0.3479	0.1066	0.8962	0.0984	0.9237	1
Day 10 P. rettgeri							
F-*Δdpt*	0.9999	NA	NA	NA	NA	NA	NA
F-*dptS69R*	0.1574	0.1425	NA	NA	NA	NA	NA
F-*dptS69*	0.9844	0.998	0.5859	NA	NA	NA	NA
M-*imd-*	0.0707	0.0541	0.9999	0.351	NA	NA	NA
M-*Δdpt*	0.9334	0.9767	0.57	1	0.3138	NA	NA
M-*dptS69R*	<u>0.0086</u>	<u>0.0042</u>	0.9253	0.0653	0.9903	<u>0.0432</u>	NA
M-*dptS69*	0.904	0.958	0.8201	0.9999	0.5938	1	0.1563
Day 20 P. rettgeri							
F-*Δdpt*	1	NA	NA	NA	NA	NA	NA
F-*dptS69R*	<u>0.0241</u>	<u>3.00E-04</u>	NA	NA	NA	NA	NA
F-*dptS69*	0.7316	0.4202	0.1779	NA	NA	NA	NA
M-*imd-*	0.3846	0.0835	0.5982	0.9932	NA	NA	NA
M-*Δdpt*	1	1	<u>3.00E-04</u>	0.4122	0.081	NA	NA
M-*dptS69R*	0.0838	<u>0.0032</u>	0.996	0.5558	0.9479	<u>0.003</u>	NA
M-*dptS69*	0.9304	0.8325	<u>0.0445</u>	0.9984	0.8519	0.826	0.2147

Table S5 (continued)

	F-*imd-*	F-*Δdpt*	F-*dptS69R*	F-*dptS69*	M-*imd-*	M-*Δdpt*	M-*dptS69R*
Day 2 L. brevis							
F-*Δdpt*	0.0027	NA	NA	NA	NA	NA	NA
F-*dptS69R*	2.09E-02	1	NA	NA	NA	NA	NA
F-*dptS69*	0.0855	0.9932	0.9993	NA	NA	NA	NA
M-*imd-*	3.00E-04	0.8295	0.8593	0.5336	NA	NA	NA
M-*Δdpt*	<02e-16	6.19E-01	0.7332	0.3352	1	NA	NA
M-*dptS69R*	<02e-16	0.0511	0.1029	0.026	0.8176	0.7015	NA
M-*dptS69*	1.00E-04	0.5769	0.6552	0.3136	1	0.9999	0.9518
Day 10 L. brevis							
F-*Δdpt*	0.9892	NA	NA	NA	NA	NA	NA
F-*dptS69R*	1	0.9836	NA	NA	NA	NA	NA
F-*dptS69*	0.6678	0.9554	0.6632	NA	NA	NA	NA
M-*imd-*	0.9941	0.6931	0.9989	0.2425	NA	NA	NA
M-*Δdpt*	8.49E-02	0.2951	0.1117	0.9918	0.0087	NA	NA
M-*dptS69R*	0.9987	1	0.9971	0.9634	0.8893	0.4734	NA
M-*dptS69*	0.0801	0.2515	0.095	0.9423	0.0119	0.9995	0.3653
Day 20 L. brevis							
F-*Δdpt*	0.9999	NA	NA	NA	NA	NA	NA
F-*dptS69R*	0.9998	0.9808	NA	NA	NA	NA	NA
F-*dptS69*	1	0.9997	0.9999	NA	NA	NA	NA
M-*imd-*	0.0045	2.00E-04	0.0175	0.0056	NA	NA	NA
M-*Δdpt*	0.9469	0.9854	0.7085	0.9241	0	NA	NA
M-*dptS69R*	0.9981	1	0.9528	0.9962	6.00E-04	0.9999	NA
M-*dptS69*	0.954	0.9876	0.7694	0.9363	1.00E-04	1	0.9998
Day 2 A. tropicalis							
F-*Δdpt*	0.9999	NA	NA	NA	NA	NA	NA
F-*dptS69R*	0.9981	0.9373	NA	NA	NA	NA	NA
F-*dptS69*	0.9993	0.9626	1	NA	NA	NA	NA
M-*imd-*	2.06E-01	0.0304	0.5635	0.5008	NA	NA	NA
M-*Δdpt*	0.5631	0.1075	0.9417	0.9084	0.9672	NA	NA
M-*dptS69R*	0.1343	0.0157	0.4295	3.72E-01	1	0.9048	NA
M-*dptS69*	0.9994	0.9634	1	1.00E+00	0.4983	0.9069	0.3698
Day 10 A. tropicalis							
F-*Δdpt*	0.5974	NA	NA	NA	NA	NA	NA
F-*dptS69R*	1	0.2639	NA	NA	NA	NA	NA
F-*dptS69*	0.812	1	0.5422	NA	NA	NA	NA
M-*imd-*	0.9993	0.1249	1	0.3359	NA	NA	NA
M-*Δdpt*	0.004	0.0796	2.00E-04	0.1109	1.00E-04	NA	NA
M-*dptS69R*	0.9998	0.7771	9.94E-01	0.9394	0.9524	0.0035	NA
M-*dptS69*	<2e-16	2.00E-04	<2e-16	5.00E-04	<2e-16	0.2145	<2e-16
Day 20 A. tropicalis							
F-*Δdpt*	0.9956	NA	NA	NA	NA	NA	NA
F-*dptS69R*	1	0.9988	NA	NA	NA	NA	NA
F-*dptS69*	1	0.998	1	NA	NA	NA	NA
M-*imd-*	0.9999	0.6771	0.9677	0.9743	NA	NA	NA
M-*Δdpt*	0.1447	3.77E-02	0.0201	1.77E-02	8.00E-04	NA	NA
M-*dptS69R*	1	9.99E-01	1	1	0.9677	0.0201	NA
M-*dptS69*	4.90E-03	1.00E-04	1.00E-04	1.00E-04	<2e-16	0.3458	1.00E-04

1023
1024

Table S6. P-values from Tukey HSD for bacterial load from poly-association with *L. plantarum* and *P. rettgeri*. P-values less than 0.05 are underlined.

	F-*imd-*	F-*Δdpt*	F-*dptS69R*	F-*dptS69*	M-*imd-*	M-*Δdpt*	M-*dptS69R*
Day 2 *L. plantarum*							
F-*Δdpt*	0.2226	NA	NA	NA	NA	NA	NA
F-*dptS69R*	0.0101	0.6004	NA	NA	NA	NA	NA
F-*dptS69*	<2e-16	2.00E-04	0.1471	NA	NA	NA	NA
M-*imd-*	4.21E-01	1	0.7283	0.0017	NA	NA	NA
M-*Δdpt*	0.0031	4.83E-01	1	0.0344	0.6898	NA	NA
M-*dptS69R*	<2e-16	6.00E-04	0.2442	1	0.0038	0.0725	NA
M-*dptS69*	<2e-16	4.00E-04	0.2095	1	0.0029	0.0577	1
Day 10 *L. plantarum*							
F-*Δdpt*	0.2226	NA	NA	NA	NA	NA	NA
F-*dptS69R*	0.0101	0.6004	NA	NA	NA	NA	NA
F-*dptS69*	<2e-16	2.00E-04	0.1471	NA	NA	NA	NA
M-*imd-*	4.21E-01	1	0.7283	0.0017	NA	NA	NA
M-*Δdpt*	0.0031	4.83E-01	1	0.0344	0.6898	NA	NA
M-*dptS69R*	<2e-16	6.00E-04	0.2442	1	0.0038	0.0725	NA
M-*dptS69*	<2e-16	4.00E-04	0.2095	1	0.0029	0.0577	1
Day 20 *L. plantarum* (no imd females)							
F-*Δdpt*	NA	NA	NA	NA	NA	NA	NA
F-*dptS69R*	NA	0.831	NA	NA	NA	NA	NA
F-*dptS69*	NA	0.9883	0.9977	NA	NA	NA	NA
M-*imd-*	NA	0.5334	0.9998	0.9429	NA	NA	NA
M-*Δdpt*	NA	0.9999	0.484	0.857	0.1772	NA	NA
M-*dptS69R*	NA	0.9202	1	0.9999	0.9943	0.6188	NA
M-*dptS69*	NA	0.9999	0.5746	0.8987	0.2615	1	0.7064
Day 2 *P. rettgeri*							
F-*Δdpt*	0.0784	NA	NA	NA	NA	NA	NA
F-*dptS69R*	0.1199	1	NA	NA	NA	NA	NA
F-*dptS69*	0.434	0.9992	0.9966	NA	NA	NA	NA
M-*imd-*	0.8069	0.912	0.8957	0.9988	NA	NA	NA
M-*Δdpt*	0.0063	0.9457	0.9972	0.8097	0.3815	NA	NA
M-*dptS69R*	0.131	1	1	0.9977	0.9101	0.9957	NA
M-*dptS69*	0.0348	0.9902	0.9996	0.9254	0.6186	1	0.9993
Day 10 *P. rettgeri*							
F-*Δdpt*	1	NA	NA	NA	NA	NA	NA
F-*dptS69R*	0.993	0.9967	NA	NA	NA	NA	NA
F-*dptS69*	0.9835	0.9901	1	NA	NA	NA	NA
M-*imd-*	0.998	0.9994	1	1	NA	NA	NA
M-*Δdpt*	1	0.9997	0.9485	0.9085	0.9773	NA	NA
M-*dptS69R*	1	1	0.9998	0.9992	1	0.9981	NA
M-*dptS69*	0.9911	0.9954	1	1	1	0.9393	0.9998
Day 20 *P. rettgeri* (no imd females)							
F-*Δdpt*	NA	NA	NA	NA	NA	NA	NA
F-*dptS69R*	NA	0.9927	NA	NA	NA	NA	NA
F-*dptS69*	NA	0.9971	0.8027	NA	NA	NA	NA
M-*imd-*	NA	0.9999	0.911	1	NA	NA	NA
M-*Δdpt*	NA	9.64E-01	1	0.5715	0.7503	NA	NA
M-*dptS69R*	NA	1	0.9896	0.9966	0.9998	0.9478	NA
M-*dptS69*	NA	0.9991	1	0.8929	0.9652	0.9998	0.9986

1028

1029 **Table S7. Desiccation resistance Tukey HSD p-values.** P-values less than 0.05 are underlined. "A" refers to flies
1030 reared axenically, "C" refers to flies reared on a conventional diet.

1031

	A *dptS69*	A *Δdpt*	A *dptS69R*	A *imd-*	A OreR	A w1118	C *dptS69*	C *Δdpt*	C *dptS69R*	C *imd-*	C OreR
A *Δdpt*	0.9999	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
A *dptS69R*	0.3541	0.9978	NA	NA	NA	NA	NA	NA	NA	NA	NA
A *imd-*	0.9195	0.8332	<u>0.003</u>	NA	NA	NA	NA	NA	NA	NA	NA
A OreR	1	1	0.5898	0.8561	NA	NA	NA	NA	NA	NA	NA
A w1118	<u><2e-16</u>	<u>7.00E-04</u>	<u><2e-16</u>	<u>0.007</u>	<u><2e-16</u>	NA	NA	NA	NA	NA	NA
C *dptS69*	<u><2e-16</u>	0.1208	0.1473	<u><2e-16</u>	<u>1.00E-04</u>	<u><2e-16</u>	NA	NA	NA	NA	NA
C *Δdpt*	<u><2e-16</u>	<u>5.00E-04</u>	<u>2.00E-04</u>	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	0.3672	NA	NA	NA	NA
C *dptS69R*	<u><2e-16</u>	<u>0.0193</u>	<u>0.0086</u>	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	0.9995	0.8062	NA	NA	NA
C *imd-*	<u>0.0057</u>	0.7281	0.963	<u><2e-16</u>	<u>0.0242</u>	<u><2e-16</u>	0.934	<u>0.013</u>	0.3953	NA	NA
C OreR	<u>0.0024</u>	0.5452	0.8351	<u><2e-16</u>	<u>0.0105</u>	<u><2e-16</u>	0.9979	0.0653	0.8078	1	NA
C w1118	0.9996	1	0.8729	0.3974	1	<u><2e-16</u>	<u>3.00E-04</u>	<u><2e-16</u>	<u><2e-16</u>	0.0781	<u>0.0352</u>

1032

1033 **Table S8. Starvation resistance Tukey HSD p-values.** P-values less than 0.05 are underlined. "A" refers to flies
1034 reared axenically, "C" refers to flies reared on a conventional diet.
1035

	A *dptS69*	A *Δdpt*	A *dptS69R*	A *imd-*	A *OreR*	A *w1118*	C *dptS69*	C *Δdpt*	C *dptS69R*	C *imd-*	C *OreR*
A *Δdpt*	0.0295	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
A *dptS69R*	<2e-16	0.0022	NA	NA	NA	NA	NA	NA	NA	NA	NA
A *imd-*	<2e-16	<2e-16	0.9288	NA	NA	NA	NA	NA	NA	NA	NA
A OreR	0.0043	1	8.00E-04	<2e-16	NA	NA	NA	NA	NA	NA	NA
A W1118	<2e-16	<2e-16	<2e-16	0.0038	<2e-16	NA	NA	NA	NA	NA	NA
C *dptS69*	5.00E-04	<2e-16	<2e-16	<2e-16	<2e-16	<2e-16	NA	NA	NA	NA	NA
C *Δdpt*	0.0055	<2e-16	<2e-16	<2e-16	<2e-16	<2e-16	1	NA	NA	NA	NA
C *dptS69R*	<2e-16	0.0614	0.998	0.3139	0.0407	<2e-16	<2e-16	<2e-16	NA	NA	NA
C *imd-*	2.00E-04	0.9988	0.0306	<2e-16	0.9995	<2e-16	<2e-16	<2e-16	0.3851	NA	NA
C OreR	0.7443	0.9473	<2e-16	<2e-16	0.828	<2e-16	<2e-16	<2e-16	1.00E-04	0.325	NA
C *w1118*	<2e-16	<2e-16	1.00E-04	0.0374	0	1	<2e-16	<2e-16	<2e-16	<2e-16	<2e-16

1036
1037
1038

1039

1040 **Table S9. Conventional longevity Cox proportional hazards regression p-values.** P-values less than 0.05 are
1041 underlined. "F" and "M" refer to males and females, respectively; "het" refers to heterozygotes for serine/arginine.
1042

	F het	F imd-	F Δdpt	F dptS69R	F dptS69	M het	M imd-	M Δdpt	M dptS69R
F imd-	1	NA	NA	NA	NA	NA	NA	NA	NA
F Δdpt	1	1	NA	NA	NA	NA	NA	NA	NA
F dptS69R	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	NA	NA	NA	NA	NA	NA
F dptS69	<u><2e-16</u>	<u>3.00E-04</u>	<u><2e-16</u>	0.6785	NA	NA	NA	NA	NA
M het	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	0.9797	<u>0.0443</u>	NA	NA	NA	NA
M imd-	0.1346	0.3182	<u>0.0498</u>	<u>0.0025</u>	0.5097	<u>0</u>	NA	NA	NA
M Δdpt	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	0.948	0.9977	0.1163	<u>0.0321</u>	NA	NA
M dptS69R	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	0.0622	<u><2e-16</u>	0.4311	<u><2e-16</u>	<u><2e-16</u>	NA
M dptS69	<u><2e-16</u>	<u><2e-16</u>	<u><2e-16</u>	0.9984	0.9882	0.5841	<u>0.0492</u>	1	<u>0.0043</u>

1043

Table S10. Genotyped wild caught *D. melanogaster*.

Wolbachia/Sex	SS	SR	RR
Wolbachia+ female	203	64	8
Wolbachia+ male	369	77	9
Wolbachia- female	68	21	2
Wolbachia- male	102	31	1
Total	742	193	20

Supplemental Figures

Figure S1. CRISPR/Cas9 genome editing of the *dpt* locus
A) Alignment of CRISPR/Cas9 genome edited lines. CRISPR/Cas9 genome edited lines aligned to consensus sequence. Genotype/line name is indicated on the left-hand side of the figure. B) Crossing schematic for CRISPR clean up. A modified Bloomington stock # 7198 was used for CRISPR injections. The line contained *CyO*/*Kruppel* balanced on the 2nd chromosome, and *serrate*/*Dichaete* balanced on the 3rd chromosome. Our version, 7198[A4], was provided by Stuart Macdonald and has the DSPR (King et al. 2012) A4 line's X chromosome instead of the *w*[*] from the original 7198.

Figure S2. Male vs. female survival after systemic infection

Systemic infections were repeated with females and males. A) *P. rettgeri* and B) *E. faecalis* are normal systemic infection bacteria in *Drosophila*. C) *L. plantarum* was used to test differences in systemic based on sex based on the results from gnotobiotic longevity.

Figure S3. Gnotobiotic longevity
Survival plots of all bacteria. Dpt null lines have been separated into the two lines used, 1 base pair deletion and 3 bp deletion.

Figure S4. Feeding Rates

Female flies eat more food (as measured by blue dye ingestion) than males, but genotype also influences feeding. Each genotype has more than 100 individuals per treatment per sex. Control refers to no added media, LB Media is the addition of sterile MB, and *P. rettgeri* is the addition of *P. rettgeri* at a concentration of OD_{600} = 15 (±1). Absorbance was measured at 630nm.

Figure S5. Bacterial load at day 2-, 10-, and 20-days post feeding
All bacteria were plated on MRS media except for *P. rettgeri* that was plated on LB media to grow colonies. Significance can be found in Table 2.S6. Individual points represent a pool of 3 flies. There is no *imd*-female bar for *L. plantarum* on day 20 because all female flies had died at this timepoint.

Figure S6. Bacterial load of *L. plantarum* and *P. rettgeri* poly-association
The flies were co-feed with *L. plantarum* and *P. rettgeri* and plated on MRS and LB respectively to select for the different bacteria. There is no *imd-* female bar on day 20 because all flies had died at this timepoint.

Figure S7. Desiccation stress survival with additional wildtype lines (OreR and W1118)

OreR and W1118 are standard *Drosophila melanogaster* lab stocks. They are both homozygous serine but have different resistance to desiccation stress than *dpt^S69^*.

Figure S8. Starvation with additional wildtype lines (OreR and w1118)
OreR and W1118 are standard *Drosophila melanogaster* lab stocks. They are both homozygous serine but have different resistance to starvation stress than *dpt^S69^*.

Figure S9. Longevity with separated *Δdpt* lines and heterozygotes

Two *Δdpt* lines were used. One with a 1 base pair deletion and the other with a 3 base pair deletion. These lines were combined for ease of viewing in Fig 4C. Heterozygotes were made with reciprocal crosses