Cis-regulatory modes of *Ultrabithorax* inactivation in butterfly forewings

Amruta Tendolkar[1], Anyi Mazo-Vargas[1], Luca Livraghi[1], Joseph J. Hanly[1,2], Kelsey C. Van Horne[1], Lawrence E. Gilbert[3], Arnaud Martin[1,*]

[1]: Department of Biological Sciences, The George Washington University, Washington, DC, United States
[2]: Smithsonian Tropical Research Institute, Panama
[3]: Department of Integrative Biology, University of Texas – Austin, Austin, TX, United States

*: Corresponding author:
Arnaud Martin, Department of Biological Sciences, The George Washington University, 800 22nd Street NW, Washington, DC 20052. Email: arnaud@gwu.edu

ABSTRACT

Hox gene clusters encode transcription factors that drive regional specialization during animal development: *e.g.* the Hox factor Ubx is expressed in the insect metathoracic (T3) wing appendages and differentiates them from T2 mesothoracic identities. *Hox* transcriptional regulation requires silencing activities that prevent spurious activation and regulatory crosstalks in the wrong tissues, but this has seldom been studied in insects other than *Drosophila*, which shows a derived *Hox* dislocation into two genomic clusters that disjoined *Antennapedia* (*Antp*) and *Ultrabithorax* (*Ubx*). Here we investigated how *Ubx* is restricted to the hindwing in butterflies, amidst a contiguous *Hox* cluster. By analysing Hi-C and ATAC-seq data in the butterfly *Junonia coenia*, we show that a Topologically Associated Domain (TAD) maintains a hindwing-enriched profile of chromatin opening around *Ubx*. This TAD is bordered by a Boundary Element (BE) that separates it from a region of joined wing activity around the *Antp* locus. CRISPR mutational perturbation of this BE releases ectopic *Ubx* expression in forewings, inducing homeotic clones with hindwing identities. Further mutational interrogation of two non-coding RNA encoding regions and one putative *cis*-regulatory module within the *Ubx* TAD cause rare homeotic transformations in both directions, indicating the presence of both activating and repressing chromatin features. We also describe a series of spontaneous forewing homeotic phenotypes obtained in *Heliconius* butterflies, and discuss their possible mutational basis. By leveraging the extensive wing specialization found in butterflies, our initial exploration of *Ubx* regulation demonstrates the existence of silencing and insulating sequences that prevent its spurious expression in forewings.

INTRODUCTION

Hox genes are key specifiers of antero-posterior regional identity in animals, and thus require robust regulatory mechanisms that confine their expression to well-delimited sections of the body. Their genomic arrangement into *Hox* gene clusters has provided a rich template for the study of gene regulation, with mechanisms including chromatin silencing and opening, 3D conformational changes, and non-coding RNAs (Mallo and Alonso 2013). However, this rich body of work has been almost exclusively performed in mice and fruit flies. In order to decipher how diverse body plans and morphologies evolved, we must start assessing the mechanisms of Hox gene regulation in a wider range of organisms.

The *Ultrabithorax* (*Ubx*) gene encodes a Hox family transcription factor involved in the specification of segment identities in arthropods (Hughes and Kaufman 2002; Heffer and Pick 2013). In insects, the conserved expression of *Ubx* in the metathoracic (T3) segment is required for their differentiation from Ubx-free tissues in the mesothorax (T2) , and has been a key factor for the specialization of metathoracic serial appendages including T3 legs (Mahfooz et al. 2007; Refki et al. 2014; Tomoyasu 2017; Feng et al. 2022; Buffry et al. 2023) and hindwings or their derivatives (Tomoyasu 2017; Loker et al. 2021). The mechanisms of *Ubx* segment-specific expression have been intensively studied in *D. melanogaster* (Mallo and Alonso 2013; Hajirnis and Mishra 2021), where *Hox* genes are separated into two genomic loci, the Antennapedia (ANT-C, *Antp*) and Bithorax clusters (BX-C). In short, the BX-C complex that includes *Ubx*, *abdominal-A* (*abd-A*), and *Abdominal-B* (*Abd-B*) is compartmentalized into nine chromosomal domains that determine the parasegmental expression boundaries of these three genes (Maeda and Karch 2015). Each boundary is primarily enforced by insulators that separate Topologically Associating Domains (TADs) of open-chromatin, while also allowing interactions of enhancers with their target promoters (Postika et al. 2018; Srinivasan and Mishra 2020). The BX-C locus also includes non-coding RNAs, some of which are processed into miRNAs known to repress *abd-A* and *Abd-B* (Garaulet and Lai 2015). *Fub-1/bxd* long non-coding RNAs (lncRNAs) situated 5' of *Ubx* are thought to participate in *Ubx* regulation in the PS5 (posterior T3 to anterior A1) parasegment (Ibragimov et al. 2022). An intronic lncRNA dubbed *lncRNA:PS4* is expressed in the PS4 parasegment (posterior T2 - anterior T3), and appears to stabilize *Ubx* in this region in mutant contexts (Hermann et al. 2022). Little is known about how *Hox* genes are regulated outside of flies, where they co-localize into a single *Hox* cluster, and where *Antp* and *Ubx* thus occur in contiguous positions (Gaunt 2022; Mulhair and Holland 2022). A few *Hox*-related miRNAs are evolutionarily conserved across the locus in arthropods (Pace et al. 2016), and an early study in *Tribolium* characterized the embryonic expression of a *Hox* cluster non-coding transcripts (Shippy et al. 2008).

These knowledge gaps lead us to consider the use of butterflies and moths (Lepidoptera) as alternative model systems for the study of *Ubx* function and regulation. Lepidopteran forewings and hindwings are functionally and morphologically differentiated, and CRISPR mosaic knock-outs (mKOs) showed that *Ubx* is necessary for the specification of hindwing color patterns, shape, and venation (Tendolkar et al. 2021). In three species of nymphalid butterflies (*Heliconius erato*, *Junonia coenia*, and *Bicyclus anynana*), CRISPR-mediated loss-of-function of *Ubx* induces regional-specific homeotic transformations of hindwing patterns into their forewing counterpart (Matsuoka and Monteiro 2018; Tendolkar et al. 2021), reminiscent of homeotic aberrations that are sporadically observed in butterfly wings (Sibatani 1983; Nijhout and Rountree 1995). The ectopic activation of *Ubx* into the pupal forewing results in the gain of hindwing features, suggesting Ubx is sufficient to drive T3-like identity when

expressed in T2 (Lewis et al. 1999; Tong et al. 2014). Besides its roles in adult wing differentiation, *Ubx*
is also known to repress thoracic leg identity in transient embryonic appendages of the first abdominal
segment, called pleuropods (Zheng et al. 1999; Masumoto et al. 2009; Tong et al. 2017; Tendolkar et al.
2021; Matsuoka et al. 2022).
The general organization of Hox gene clusters has been well described in Lepidoptera, but their
regulation has been seldom studied. Lepidopteran genomes have accumulated divergent *Hox3* copies,
named *Shox* genes, that are required during early embryonic development but do not appear to play
homeotic functions (Ferguson et al. 2014; Livraghi 2017; Mulhair et al. 2022). An lncRNA and two
miRNAs were identified in the intergenic region between *abd-A* and *Abd-B* in the silkworm (Wang et al.
2017, 2019). In butterfly wings, the regulation of *Ubx* shows strong patterns of segment-specific
regulation at two levels. First, the *Ubx* transcript has been consistently identified as the most
differentially expressed mRNA between the two wing sets (Hanly et al. 2019; Wang et al. 2022). Second,
comparison of ATAC-seq signals reveal that forewing vs. hindwing have identical open-chromatin
profiles during development across the genome, except at the *Ubx* gene itself (Lewis and Reed 2018;
van der Burg et al. 2019). Thus, the ability of the *Ubx* locus to be robustly activated in hindwings and
repressed in forewings is likely driving most subsequent differences between these tissues. In this
study, we provide an initial assessment of the regulation of the *Ubx* locus during butterfly wing
development. To do this, we leverage genomic resources and CRISPR mutagenesis with a focus on two
laboratory species belonging to the Nymphalinae sub-family, *J. coenia* and *Vanessa cardui* (Livraghi et
al. 2017; Martin et al. 2020; van der Burg et al. 2020; Mazo-Vargas et al. 2022). We identify putative
regulatory regions likely involved in the repression and activation of *Ubx* expression, and discuss the
potential mechanisms restricting it to hindwings. Finally, we describe a collection of spontaneous wing
homeotic mutants in *Heliconius spp.* and elaborate on the categories of mutations that could underlie
these phenotypes by misregulating *Ubx*.

RESULTS

Gene expression at the *Ubx* locus during wing development

We provide annotations of the *Ubx* genomic region in four Nymphalinae butterflies (**Fig. 1A**). These feature existing genomic resources for our model species *J. coenia* and *V. cardui* (van der Burg et al. 2020; Lohse et al. 2021b; Zhang et al. 2021), as well as for *Aglais* (*Nymphalis*) *io* (Lohse et al. 2021a). The publicly available annotations for these three species include evidence from developmental transcriptomes, and we added to this set a manual annotation of the *Ubx* locus from the oak leaf butterfly *Kallima inachus*, for which forewing *vs*. hindwing transcriptomes have been sequenced across a replicated developmental time series (Yang et al. 2020; Wang et al. 2022).

All Nymphalinae show a similar organization of the region spanning *Ubx*. Interestingly, the first intron of *Ubx* encodes a long non-coding RNA in opposite orientation to *Ubx*, that we dub here *lncRNA:Ubx-IT1* (abbr. *Ubx-IT1*), based on the recommended nomenclature (Seal et al. 2022). Orthologous versions of *Ubx-IT1* are detected in most NCBI RefSeq genome annotations throughout Lepidoptera (*e.g.* the ncRNA *NCBI:XR_960726* in *Plutella xylostella*), implying it is a conserved feature of the *Ubx* locus in this insect order. Finally, both annotations from *V. cardui, A. io,* and *J. coenia* show a long intergenic non-coding transcript starting in antisense orientation about 10-15 kb 5' of *Ubx*, that we dub here *lincRNA:Ubx-AS5'* (abbr. *Ubx-AS5'*). This transcript was neither detected in *K. inachus* transcriptomes nor in RNA datasets outside of the Nymphalinae sub-family, and could be specific to this lineage.

Next we reanalyzed the *K. inachus* wing transcriptomes (Wang et al. 2022), and profiled the expression of *Ubx* region transcripts during wing development (**Fig. 1B**). As expected from previous studies (Hanly et al. 2019; Paul et al. 2021; Merabet and Carnesecchi 2022; Wang et al. 2022), *Ubx* showed a strong expression bias in hindwings, spanning the larval imaginal disks to the intermediate pupal stage. Of note, Ubx is confined to the peripodial membranes of insect T2 wing appendages (Weatherbee et al. 1998, 1999; Prasad et al. 2016), which may explain residual detection in some of the forewing samples here. *Ubx-IT1* was significantly enriched in hindwings compared to forewings, albeit at ~100-fold lower count levels than *Ubx* in the same samples. The *Hox* gene *Antp* showed a minor forewing enrichment, confirming that while *Ubx* expression is robustly repressed in T2 forewing tissues, *Antp* expression is permitted in both T2 and T3 appendages (Matsuoka and Monteiro 2021, 2022; Paul et al. 2021). Expression of *abd-A* was undetected in most wing samples.

131
Figure 1. Annotation of the *Ubx* genomic interval in four butterflies of the Nymphalinae sub-family. (A) Genomic
intervals spanning *Antp*, *Ubx*, and *abd-A*, featuring published transcript annotations from NCBI Reference Genomes of *V. cardui* and *A. io*, and
manual re-annotations of the *J. coenia* and *K. inachus* genomes using published RNAseq dataset (see Methods). Exons are shown with coding
(thick) and non-coding (thin) sections. No *lincRNA:Ubx-AS5'* transcripts were detected in *K. inachus*. **(B)** Expression profiling of transcripts of
the *Ubx* region in *K. inachus*, based on a reanalysis of published wing RNA-seq transcriptomes (Wang et al. 2022). Expression levels are plotted
as DESeq2 normalized counts plots. Pairwise Wald tests adjusted for multiple test correction each assess differential expression between
forewings and hindwings. ns : non-significant ; * : $p < 0.05$; ** : $p < 0.01$; *** : $p < 0.001$.

Chromatin 3D conformation reveals a Boundary Element between *Antp* and *Ubx*

Genome-wide Hi-C sequencing can be used to generate heatmaps that capture the conformation of 3D chromatin in tissues, and has been used extensively to study *Drosophila Hox* cluster organisation into TADs that prevent regulatory crosstalk between adjacent genes (Ibragimov et al. 2022; Moniot-Perron et al. 2023). Here we used Hi-C to assess the 3D chromatin architecture of the *Hox* cluster interval in the butterfly *J. coenia*, using existing datasets that were generated from fifth instar larval forewings (van der Burg et al. 2020; Mazo-Vargas et al. 2022). In larval forewings, the *Hox* chromatin conformation landscape consists of three well-delimited TADs, the first one spanning *proboscipedia* (*pb*) to *Sex comb reduced* (*Scr*), the second one around *Antp*, and the third one *Ubx*, *abd-A, and Abd-B* (**Figs. 2 and 3A**). A Boundary Element (BE), was robustly called (see Methods) at the region separating the *Antp* and *Ubx* TADs, situated in the *Ubx* last intron. Because TAD boundary prediction has a coarse resolution, we arbitrarily define the candidate BE region as a 15-kb interval centered in the *Ubx* last intron, and dub it *Antp-Ubx_BE*. A binding motif analysis identified 4 CTCF binding sites in a 1-kb interval within *Antp-Ubx_BE*, two of which were found in a tightly linked, convergent orientation (**Fig. S1**), which is consistent with TAD insulating role in mediating chromatin loop-extrusion (Guo et al. 2015). This concordance between Hi-C profiling and CTCF motif prediction thus indicates that *Antp-Ubx_BE* region functions as an insulator between regulatory domains of *Antp* and *Ubx*.

Differential forewing vs. hindwing chromatin-opening across the *Antp-Ubx* interval

In flies, the *Ubx/abd-A* section is organized into regulatory domains that display differential activities across the antero-posterior axis, following what has been called the open-for-business model (Maeda and Karch 2015; Gaunt 2022). Here we tested if this pattern extends to butterfly species with a contiguous *Hox* cluster. To do this we used ATAC-seq datasets from *J. coenia* forewing (T2), hindwing (T3), and whole-head tissues sampled across fifth instar larval and early pupal stages, similarly to previous studies (van der Burg et al. 2020; Mazo-Vargas et al. 2022; Van Belleghem et al. 2023). These data reveal that chromatin opening along the *Antp/Ubx/abd-A* interval is partitioned into several regions showing a transition of T2 to T3 activity (**Fig. 2B**). From the anterior to posterior *Hox* colinear order (*i.e.* from *Antp* towards *abd-A*), chromatin-opening forms a block of forewing-enriched activity close to *Antp* and its 5' region, to a block of activity in both forewings and hindwings that stops at the *Antp-Ubx_BE*. This region is consistent with the fact that *Antp* is expressed in both wing pairs (**Fig. 1B**). From *Antp-Ubx_BE,* the interval including *Ubx* and a large upstream region is strongly enriched for hindwing opening, consistently with previous studies that found it to be the only genomic region showing this pattern (Lewis and Reed 2018; van der Burg et al. 2019). Last, the region surrounding *abd-A* is devoid of differntial open-chromatin activity between forewings and hindwings, in accordance with the exclusion of its expression from thoracic segments (Warren et al. 1994; Tong et al. 2014)

Figure 2. A region of hindwing-specific chromatin-opening is bordered by a TAD BE in the last intron of *Ubx*.
(A) Hi-C contact heatmap in fifth instar forewings of *J. coenia* and TAD separation scores around *Ubx*. A TAD boundary element (*Antp-Ubx_BE*)
is inferred in the last intron of *Ubx* (vertical dotted line). **(B)** Differential ATAC-seq profiles, re-analyzed from a previous dataset (Mazo-Vargas
et al. 2022). Top : open-chromatin profiles of forewings (FW, magenta), and hindwings (HW, green), respectively subtracted from larval head
signal (purple, negative when wing signals at background-level). Bottom : subtractive ATAC-seq profile (HW-FW) revealing hindwing-enriched
chromatin in the *Ubx* locus. *Antp-Ubx_BE* is in the vicinity of an isolated region of forewing-enriched opening (blue arrowhead). **(C)** *PhastCons*
genomic alignment scores, with overall alignability suggesting minimal structural variation across this interval in Lepidoptera and Trichoptera.

Figure 3. Hindwing-enriched chromatin-opening around *Ubx*, and the *Antp-Ubx_BE* boundary, are both maintained in mid-pupal
hindwings. (A) Hi-C heatmap in *J. coenia* fifth instar larval forewings, and subtractive ATAC-seq profiles at this stage (hindwing-forewing), as
expanded from Fig. 2 across the *Hox* cluster. **(B)** Hi-C heatmap in *J. coenia* mid-pupal hindwings, and subtractive ATAC-seq profiles at this
stage (forewing-hindwing). Inferred TAD boundaries are shown as vertical dotted lines. Blue arrowhead : position of the *Antp-Ubx_BE* sgRNA.

Comparison of 3D conformation and open-chromatin profiles between larval forewings and mid-pupal hindwings

The Hi-C dataset analyzed above was prepared from larval forewings, and forewings do not express Ubx (**Fig. 1B**). Next, we repeated our analysis on a Hi-C dataset generated in pupal hindwings instead (van der Burg et al. 2020), *i.e.* in a later-stage tissue expressing *Ubx*. We found two main differences in this contact landscape compared to the larval forewing (**Fig. 3**). First, the TAD spanning from *proboscipedia* (*pb*) to *fushi-tarazu* (*ftz*) faded in intensity, while in contrast, the TAD around *Antp* remained strongly organized. Second, *Ubx* lost its physical association to the *abd-A* and *Abd-B* domains, and gained a TAD boundary situated in the *Ubx-AS5'* intron. It is difficult to disentangle effects from staging (larval vs. pupal) and tissues (forewing vs. hindwing) in this comparison. Specifically, these differences we observed may be due to chromatin remodeling between stages, as somewhat expected during metamorphosis (Gutierrez-Perez et al. 2019). Alternatively, it is also possible hindwing development requires insulate *Ubx* from more posterior enhancers. These issues will require further investigation, for instance using profiling of histone marks, with pairwise forewing-hindwing comparison at single stages. Nonetheless the later hindwing sample showed a maintenance of *Antp-Ubx* separation. First, while *Ubx* formed a smaller TAD spanning its coding exons 1-2, this region conserved a domain of hindwing-enriched open-chromatin. Second, boundary prediction called two possible, tightly linked TAD limits in the *Antp-Ubx_BE* region, showing that the last intron of *Ubx* still acts as an insulating region. In conclusion, our preliminary comparison of *Hox* 3D conformation indicates that the *Antp-Ubx_BE* is relatively stable across two stages and wing serial homologs.

Mutagenic perturbation of *Antp-Ubx_BE* results in forewing homeosis

Next, we reasoned that the forewing-enriched ATAC-seq peak present in the inferred boundary interval (**Fig. 4A**) might mediate the binding of insulator proteins (Savitsky et al. 2016; Stadler et al. 2017), or act as a transcriptional silencer (Segert et al. 2021). Several genomic features support the former hypothesis. First, the only forewing-enriched ATAC-seq peak across a 150-kb region (spanning the *Ubx* gene and the *Antp-Ubx* intergenic region), coincides with the midpoint between the two tentative *hicFindTADs* boundary predictions inferred from HiC data (**Fig. 2B**). Second, during motif scans conducted across that 150-kb region we found 8 predicted binding-sites for the *Drosophila* CCCTC-Binding Factor (CTCF) clustered in a 5-kb region around the differentially accessible region, and none elsewhere in the last Ubx intron (**Fig. 4A**), suggesting the forewing-enriched ATAC-seq peak may function as a transcriptional insulator (Gambetta and Furlong 2018; Postika et al. 2018; Kyrchanova et al. 2020; Kaushal et al. 2022). Last, the two candidate CTCF binding motifs that are within the forewing-enriched ATAC-seq peak are also conserved across Lepidoptera and Trichoptera (**Fig. S1**), two lineages that diverged around 300 Mya (Kawahara et al. 2019; Thomas et al. 2020).

To test this hypothesis, we used CRISPR targeted mutagenesis to perturb *Antp-Ubx_BE* and assess its functionality, and designed a single sgRNA in a conserved sequence within the forewing-enriched ATAC-seq (**Fig. S1**). Remarkably, CRISPR mutagenesis of the *Antp-Ubx_BE* target induced G_0 mutants with homeotic transformations of their forewings into hindwings (**Figs. 4B-C and S2**), including identity shifts in patterns, venation, and wing shape. It is important to note that none of the resulting crispants showed hindwing effects. Thus, we can reasonably attribute forewing homeotic phenotypes to indel mutations restricted to the intronic region, without disruption of the *Ubx* transcript, as this would result in hindwing phenotypes (Matsuoka and Monteiro 2021; Tendolkar et al. 2021).

228
Figure 4. CRISPR perturbation of *Antp-Ubx_BE* results in FW→HW homeoses. **(A)** *Antp-Ubx_BE* sgRNA targeting (cyan triangle) of a FW-
enriched ATAC-peak (magenta) within the *Ubx* last intron. **(B-C)** Two examples of *J. coenia Antp-Ubx_BE* crispants showing mosaic FW→HW
homeoses, shown in dorsal views. CL-WT : contralateral, horizontally flipped images of forewings from the same individuals. WT HW : wild type
hindwings from the same individual and mutant forewing side. Both individuals show disruption of their Radial veins (R$_1$-R$_5$ area). The specimen
shown in C displays a partial, ectopic eyespot (asterisk). **(D-E)** Immunofluorescent detection of the UbdA epitope (green) in fifth instar wings
disks of *Antp-Ubx_BE* crispants, revealing ectopic antigenicity in forewings. WT forewings of similar stage, and HW from the same crispant
individuals, are shown for comparison as insets. Green autofluorescence was observed in tracheal tissues.

Homeotic clones are first visible in *Antp-Ubx_BE* crispants at the pupal stage, with streaks of
thinner cuticle, sometimes associated with local necrosis or with suture defects in the ventral midline, in
particular where leg and wing pouches adjoin (**Fig. S3**). Color pattern homeotic clones were salient at
the adult stage, with for example, clonal losses of the forewing specific white-band, and partial
acquisitions of the large M_1-M_2 hindwing eyespot. In one specimen, an ectopic, partial M_1-M_2 hindwing
eyespot appeared in the R_5-M_1 region, suggesting a perturbation of the eyespot induction process in
this wing. Nymphalid forewings have five radial veins (R_{1-5}), which provide sturdiness for flight (Wootton
1993), while hindwings have only two Radial veins. Forewing homeotic mutants showed mosaic venation
defects in the Radial vein area (**Fig. 4B**). Finally, higher expressivity mutant forewings were smaller and
rounder, reminiscent of hindwing shape.
Next, we dissected fifth instar larval wing disks from developing *Antp-Ubx_BE* crispants, and
monitored the expression of Ubd-A (Ubx and Abd-A epitopes), normally restricted to the hindwing and
only present in the forewing peripodial membrane (Weatherbee et al. 1999). Crispants showed forewing
clones with strong ectopic expression of Ubd-A (**Figs. 4D-E and S4**). This result supports the inference
that *Antp-Ubx_BE* forewing homeoses are due to the de-repression of *Ubx* in this tissue.
251
Mutational interrogation of lncRNA-encoding regions at the *Ubx* locus
We used CRISPR mutagenesis to test the function of the two lncRNA-encoding loci at the *Ubx* locus.
Mutagenesis of the *Ubx-IT1* first exon in *J. coenia*, and of the *Ubx-T1* promoter in *V. cardui*, both
resulted in crispants with small homeotic phenotypes in forewings and hindwings (**Figs. 5 and S5**). This
result contrasts with *Ubx* exon mKO experiments, which only generate hindwing phenotypes (Tendolkar
et al. 2021). Given the scarcity of *Ubx-IT1* crispants obtained (11 out of 236 adults), and the small size of
the homeotic clones within them, we infer the occasional phenotypes may be due to rare alleles. Thus,
rather than evidence of functionality of the *Ubx-IT1* transcript, the homeotic phenotypes may rather
reflect the effects of regulatory perturbation on *Ubx* itself, with some random mutations in this intronic
region resulting in hindwing *Ubx* loss-of-function, and some others triggering derepression in forewings.
Likewise, next we mutagenized the first exon of *Ubx-AS5'*, located upstream of the *Ubx* promoter, and
obtained twelve hindwing mutants and a single forewing mutant (**Fig. 6 and S6**). As with *Ubx-IT1*
CRISPR experiments, these results may be explained by regulatory disruption of *Ubx* transcription, with
a higher ratio of hindwing phenotypes compared to forewings linked to the proximity of the *Ubx*
promoter. Overall, we conclude that the mutational interrogation at these loci can result in dual loss
(hindwing) and gain (forewing) of *Ubx* function effects. Deciphering whether or when these effects
affected *Ubx* expression via local *cis*-regulatory modules, impairment of lncRNA transcripts, or larger
indels overlapping with *Ubx* exons, will require further study (see Discussion).

Table 1. CRISPR mutational interrogation experiments at putative *Ubx* regulatory regions

Species	sgRNA(s)	Inj. Embryos Ninj	L1 larvae Nhat	Pupae or L5 larvae	Adults Nadu	Crispants Nmut	Inj. time h AEL	Cas9:sgRNA ng/µL	Hatching Rate Nhat/Ninj	Crispant Rate Nmut/Ninj
J. coenia	*Antp-Ubx_BE*	59	50	50	44	6	2.5-3.5	500 : 250	84.7%	10.2%
		118	40	40	31	6	1.75-2.75	250 : 125	33.9%	5.1%
		89	44	*44*	*39 **	17	2.25-3.5	500 : 250	49.4%	19.1%
	Total	**266**	**90**	**134**	**115**	**29**			**33.8%**	**10.9%**
V. cardui	*IT1_sgRNA1*	204	67	50	50	2	1-3	250 : 125	32.8%	1.0%
		108	49	31	31	3	2-3	125 : 62.5	45.4%	2.8%
		145	60	39	39	2	2.25-3.5	500 : 250	41.4%	1.4%
	Total	**457**	**176**	**120**	**120**	**7**			**38.5%**	**1.5%**
J. coenia	*IT1_sgRNA2*	59	40	7	6	0	0.5-2.5	500 : 250	67.8%	0.0%
		124	112	112	110	4	2.25-4.75	500 : 250	90.3%	3.2%
	Total	**183**	**152**	**119**	**116**	**4**			**83.1%**	**2.2%**
V. cardui	*AS5_sgRNA1*	334	183	57	52	5	2-3	250 : 125	54.8%	1.5%
		122	87	2	2	0	2-4	500 : 250	71.3%	0.0%
	Total	**456**	**270**	**59**	**54**	**5**			**59.2%**	**1.1%**
J. coenia	*AS5_sgRNA1*	**309**	**181**	**181**	**181**	**8**	2-4.5	500 : 250	**58.6%**	**2.6%**
J. coenia	*Ubx11a2+3 +c5+6*	317	18	-	-	2	1-3	500 : 75 ea.	5.7%	0.6%
		203	35	0	0	0	1.5-3.5	500 : 75 ea.	17.2%	0.0%
	Total	**520**	**53**	**-**	**-**	**2**			**10.2%**	**0.4%**
V. cardui	*Ubx11a2+c5*	50	5	3	3	2	4-4.5	500 : 500	10.0%	4.0%
		151	29	6	5	2	2-2.75	500 :125:125	19.2%	1.3%
		361	18	13	16	6	0.5-2	500 :125:125	5.0%	1.7%
	Total	**562**	**52**	**22**	**24**	**10**			**9.3%**	**1.8%**
V. cardui	*Ubx11c5*	168	99	27	26	3	3.75-4.75	250 : 125	58.9%	1.8%
		62	22	9	9	2	0.5-0.75	500 : 250	35.5%	3.2%
		131	93	8	8	3	1.5-3	500 : 250	71.0%	2.3%
		114	63	20	20	6	3.5-4.5	500 : 250	55.3%	5.3%
	Total	**475**	**277**	**64**	**63**	**14**			**58.3%**	**2.9%**
V. cardui	*Ubx11b9*	32	18	6	5	1	1.25-2.25	500 : 250	56.3%	3.1%
		63	49	9	6	1	3.5-4.5	500 : 250	77.8%	1.6%
	Total	**95**	**67**	**15**	**11**	**2**			**70.5%**	**2.1%**
J. coenia	*Ubx11b9*	41	13	13	13	3	2.5-4	125 : 62.5	31.7%	7.3%
		48	21	14	14	1	2-3	250 : 125	43.8%	2.1%
	Total	**89**	**34**	**27**	**27**	**4**			**38.2%**	**4.5%**

* : upper estimate, includes 16 fifth instar larvae that were dissected for immunostainings, of which 7 were mutants (FW UbdA+), and 3 dissected mutant pupae.

Figure 5. Rare, dual homeoses obtained from CRISPR mutagenesis of the *lncRNA_Ubx-IT1* 5' region. (A) Genomic context of the sgRNA targets (here shown in *J. coenia*), in the promoter and first exon of the non-coding *Ubx-IT1* transcript. **(B-C)** Dorsal and ventral views of a *J. coenia* crispant displaying dual homeoses, *i.e.* with both FW→HW (presumably due to *Ubx* gain-of-expression), and HW→FW clones (akin to *Ubx* null mutations). Insets on the right describe forewing mutant clones (*IT1 mKO*), in apposition to CL-WT (contralateral forewings from the same individual), and WT HW (wild type hindwings from the same individual and mutant forewing side). **(D)** Examples of dual homeoses obtained when targeting orthologous sites in *V. cardui*.

278 **Figure 6. Homeoses obtained from CRISPR mutagenesis of the lncRNA *Ubx-AS5'* first exon**. **(A)** CRISPR sgRNA targets (here shown in
279 *J. coenia*), in the first exon of the non-coding *Ubx-AS5'* transcript. **(B)** A single *J. coenia* crispant showed a FW→HW transformation. Insets on
280 the right describe forewing mutant clones (*AS5' mKO*), in apposition to CL-WT (contralateral forewings from the same individual), and WT HW
281 (wild-type hindwings from the same individual and mutant forewing side). **(C-D)** Examples of HW→FW homeoses obtained in J. coenia or when
282 targeting orthologous sites in *V. cardui*. Scale bars: 500 μm.

Dual effects of mutagenesis in a putative *Ubx cis*-regulatory module

In an attempt to probe for *Ubx* hindwing-specific regulatory sequences, we focused on a ~ 25kb region
in the first intron of *Ubx* that displays an ATAC-seq signature of hindwing enrichment in open-
chromatin relative to forewings, hereafter dubbed *CRM11* (**Fig. 7A**). We sub-divided this differentially
accessible region into four peaks (*11a, b, c* and *d*). Targeting the ATAC-seq peaks with multiple sgRNAs
spanning sub-domains *11a* and *11c* (*UbxCRE11a2c5* in *V. cardui*, *11a2a3c5c6* in *J. coenia*), or with a

single target in *11c* (*UbxCRE11c5* in *V. cardui*) yielded dual homeoses : FW→HW and HW→FW (**Figs. 7B-**
D and S7). Hindwing effects were reminiscent of *Ubx* protein coding knockouts (Tendolkar et al. 2021),
indicating that these crispant alleles with a hindwing phenotype produce *Ubx* loss-of-function effects.
Individuals with hindwing clones 2.75 times more common than individuals with forewings in this
dataset. Similarly to the lncRNA loci perturbation experiments, dual homeoses may indicate the
presence of hindwing activators and forewing repressors in the *CRM11* region, with various CRISPR
alleles producing a spectrum of indels and effects (see Discussion). It is noteworthy that while single-
target experiments showed little lethality (55% hatching rate), dual or quadruple injection mixes
resulted in low hatching rates of injected embryos (~ 10%). Multiple targeting thus appears to induce
high-rates of embryonic lethality, possibly due to chromosomal damage (Cullot et al. 2019; Zuccaro et
al. 2020). Dual targeting with *a2+c5* also yielded partial HW→FW homeoses in *V. cardui* under the form
of ectopic white eyespot foci phenotypes (**Fig. 7E**), as occasionally observed in *Ubx* null crispants
(Tendolkar et al. 2021), seemingly due to hypomorphic or heterozygous allelic states.
Next, we focused on a single target shared between both *V. cardui* and *J. coenia* in the *11b* sub-
domain. A whole genome alignment between 23 lepidopteran species and 2 trichopteran species
indicated that region *11b* is deeply conserved, suggesting important functional constraints on its
sequence (**Fig. S8A-B**). Mutagenesis of *11b* yielded a relatively high hatching rate (mean = 51.8 %),
indicating a rare occurrence of the deleterious mutational effects observed in multiple targeting (see
above). Four *J. coenia* crispants and two *V. cardui* crispants were obtained, all exclusively showing
hindwing phenotype devoid of forewing effects. HW→FW homeoses included a variety of phenotypes all
seen in *Ubx* CDS mutants (Tendolkar et al. 2021), including transformations of the orange Discalis
elements and the white band in *J. coenia*, and partial shifts in eyespot identity (**Fig. S8C**). Together the
consistency in direction of transformations and the deep conservation of the *11b* region suggests it may
encode an enhancer necessary for the transcriptional activation of *Ubx* in hindwings.

314
Figure 7. CRISPR perturbation of *Ubx CRM11* generates occasional dual homeotic phenotypes. (A) Overview of ATAC-seq differential
chromatin accessibility profiles (hindwing - head tissues, green ; forewing - head tissue, magenta) across the *Ubx* first exon. Several regions
show differential opening between wings, one of which (*CRM11*), was targeted here for CRISPR perturbation (sites *a2* and *c5* indicate sgRNA
targets). **(B)** Dual homeosis phenotypes obtained in *V. cardui* following dual-targeting of *UbxCRE11a2c5*, including homeoses in color patterns
and scale morphology. **(D)** Additional example of a *V. cardui UbxCRE11a2c5* crispant with a forewing phenotype (gain of hindwing hair patches,
arrowheads). **(E)** Example of mild hindwing homeoses showing a white eyespot focus on the dorsal and ventral sides. These effects were

previously shown to occur in coding *Ubx* CRISPR knock-out experiments (Tendolkar et al, 2021). Contralateral (CL) WT wings are shown for
comparison with mutant wings (B-E). Colored dashed lines: wing veins. Scale bars: 500 µm.

A sample of spontaneous homeotic mutants in *Heliconius* butterflies

Homeotic shifts between forewings and hindwings can occur naturally in Lepidoptera, and have been

documented as pattern aberrations in museum specimens (Sibatani 1980, 1983). As a complement to

CRISPR-induced homeoses, we document here a rich sample of forewing/hindwing homeotic mutants

in the genus *Heliconius*, systematically collected by L. E. Gilbert between 1987 and 2016 in captive

stocks at UT Austin, as well as in the wild. Across these 15 spontaneous mutants, 12 show HW➔FW

clones (**Fig. S9**), against 3 specimens with FW➔HW effects (**Fig.8**). Mutant clones in this dataset were

always posterior to the M_2 vein. Only 2 mosaic phenotypes were found on a dorsal side, with the 13

others appearing ventrally. These homeotic mosaics show pattern shifts with complete fore/hindwing

conversions of scale types, as seen for instance in the loss of gray wing coupling scales on posterior

ventral forewings (**Fig. 8A-B**), or conversely, in their acquisition in posterior hindwings (arrowheads in

Fig. S9B-D, H). Homeoses also include noticeable local changes in wing shape, particularly in

hindwings (asterisks in **Fig. S9**). Taken together, these effects are akin to CRISPR-induced

perturbations at the *Ubx* locus. We speculate that fore/hindwing homeotic aberrations, found in nature

and captive stocks, result from mutations at the *Ubx* locus itself.

338
339 **Figure 8. Mosaic forewing homeoses in *Heliconius* butterfly spontaneous mutants.** Wild-type and mutant sides from the same
340 individuals are shown in each panel, with one side digitally flipped to match left-to-right orientation. **A.** *Heliconius melpomene rosina*, ventral
341 view. Wild-caught in the Osa Peninsula (Costa Rica), October 1989. **B.** *Heliconius cydno galanthus*, ventral view (magnified inset in **B'**). Stock
342 culture from Organisation for Tropical Studies station, La Selva (Costa Rica), June 1990 **C.** *Heliconius himera*, dorsal view (magnified inset in
343 **C'**). Stock Culture in the butterfly farm Heliconius Butterfly Works in Mindo (Ecuador), March 2008.

DISCUSSION

Figure 9. Summary of wing homeosis phenotypes obtained from mutational interrogation. (A) CRISPR targets at non-coding regions across the *Ubx* region, here visualized in *J. coenia*. **(B)** Summary of injection and adult phenotype data obtained across CRISPR experiments. FW/HW crispants : total number of individuals with forewing or hindwing homeotic clones, regardless of the injected species. Individuals with dual homeosis are counted in both categories. N_{mut}/N_{inj} : number of crispants obtained (N_{mut}), over the number of injected embryos for each species. Bold: experiments with consistent effects in only one segment. See **Table 1** for details.

An intronic region with ATAC-seq hindwing-enrichment regulates *Ubx*

All CRISPR targets yielded homeotic phenotypes (**Fig. 9**), with two kinds of interference with *Ubx* expression – forewing gain-of-function effects, and hindwing loss-of-function effects – and indicating the presence of regulatory sequences (broadly defined), that repress or enhance *Ubx* expression in this region. It is crucial here to highlight the limitations of the method, in order to derive proper insights about the functionality of the regulatory regions we tested. In essence, butterfly CRISPR experiments generate random mutations by non-homologous end joining repair, that are usually deletions (Connahs et al. 2019; Mazo-Vargas et al. 2022; Van Belleghem et al. 2023), and they require genotyping in a second (G_1) generation to be properly matched to a phenotype (genotyping G_0 mosaic wings is limited, because adult wings lost scale building cells that underlie a given phenotype). Previous data from other organisms suggests that Cas9 nuclease targeting can generate larger than expected mutations that evade common genotyping techniques (Shin et al. 2017; Adikusuma et al. 2018; Kosicki et al. 2018; Cullot et al. 2019; Owens et al. 2019). Even under the assumption that such mutations are relatively rare in butterfly embryos, the fact we injected >100 embryos in each experiment makes their occurrence likely (**Fig. 9**).

When targeting hindwing-enriched ATAC-seq peaks within the first intron of *Ubx* – from *CRM11* to the hindwing-enriched open-chromatin peak that coincides with the first exon of *Ubx-IT1* – we obtained a mixture of hindwing and forewing phenotypes. Given the potential heterogeneity of allele sizes in these experiments, it is difficult to conclude robustly about the function of individual targets. Nonetheless, the phenotypic data and in particular the obtention of dual homeoses suggest we disrupted sequences that are necessary to *Ubx* activation in hindwings, as well as to its repression in forewings. Bifunctional *cis*-regulatory elements that can switch between enhancer and silencer roles are prevalent in *Drosophila* (Gisselbrecht et al. 2020; Segert et al. 2021; Pang et al. 2022). The *CRM11* and *IT1* targets adjoin or overlap with open-chromatin signals in both wing sets (**Figs. 5A and 7A**), providing circumstantial evidence that these regions might serve as bifunctional elements. Similar observations were recently made in mutational interrogation experiments of the butterfly *WntA* patterning gene (Mazo-Vargas et al. 2022). Alternatively, silencers and enhancers may be tightly linked and interact in close proximity to shape gene expression (Méndez-González et al. 2023), implying in our

case that forewing and hindwing phenotypes are mediated by alleles spanning adjacent but distinct
elements. A formal test of these mechanisms would require germline transmission and genotyping of
these alleles, which was unsuccessful in our attempts at crossing *Ubx cis*-regulatory crispants.
In contrast with the dual effects obtained when targeting *CRM11a+c* (**Fig. 9**), *CRM11b*
perturbation resulted in hindwing-limited effects, and may suggest that an *Ubx* enhancer was
consistently compromised in this specific dataset. The high lethality and small size of mutant wing
streaks suggest that only individuals with sparse, small mutant mitotic clones can survive to the adult
stage. If this is true, *CRM11* may contain pleiotropic enhancers that are vital for normal *Ubx* function at
earlier stages, but expression-reporter studies will be required to test this.
390
Parsing lncRNA-encoding regions – correlation or cause?
LncRNAs are emerging as important regulators of gene expression and developmental processes
(Zhang et al. 2019; Statello et al. 2021). *IT1* targeting generated a majority of forewing phenotypes,
suggesting perturbation of *Ubx* repression in the T2 segment. However, *IT1* showed low expression in
forewing RNAseq datasets from *K. inachus,* and higher expression in the hindwing (**Fig. 1B**), a pattern
inconsistent with a repressive role of the antisense *IT1* transcript on *Ubx* expression. It is generally
challenging to disentangle the effects of transcription of a non-coding element from the potential
effects of adjacent enhancers (Natoli and Andrau 2012; Pease et al. 2013). Therefore, an alternative
explanation would be that the phenotypes are confounded by the overlap and proximity to open-
chromatin regions, which may play *cis*-regulatory roles on *Ubx* via DNA-protein interactions, rather than
via the lncRNA. If this is the case, it is possible that the targeted *Ubx-IT1* site, which yielded homeoses
in both directions and bears both forewing and hindwing open-chromatin, is a bifunctional *cis*-
regulatory element that can shift regulatory activities between these tissues (Gisselbrecht et al. 2020).
Targeted mutagenesis of the *Ubx-AS5'* first exon mainly generated hindwing phenotypes, although with
a relatively low-efficiency. Because this target is about 10 kb away from the *Ubx* promoter itself, it is
plausible that the observed phenotypes were due to large deletions reaching the promoter region of
Ubx. Because mutational interrogation alone cannot discern if phenotypic effects arose from regulatory
failure at the chromatin or transcript level, determining if *AS5'* and *IT1* are functional lncRNAs will need
further examination.
410
A TAD boundary element likely acts as an insulator preventing *Ubx* forewing expression
Tight maintenance of TAD boundaries at the *Hox* locus is crucial for accurate segment identity and is
facilitated by insulator proteins (Stadler et al. 2017; Gambetta and Furlong 2018; Ramírez et al. 2018).
The *Antp-Ubx_BE* element we targeted is in a good position to block interactions between *Antp* and *Ubx*
(**Figs. 2-3**). Consistent with this idea, the last intron of *Ubx* contains 8 CTCF binding motifs that are all
clustered within 5-kb around the forewing-enriched ATAC peak, including two sites at highly conserved
positions that are only 100-bp away from the CRISPR target (**Fig. S1**). CTCF sites prevent cross-talk
between regulatory domains in the fly BX-C complex, and result in *Hox* misexpression when deleted
(Postika et al. 2018; Kyrchanova et al. 2020; Kaushal et al. 2022; Kahn et al. 2023). Thus, the density of
CTCF sites in this region may be indicative of a *bona fide* insulator active in forewings.
CRISPR mutagenesis of *Antp-Ubx_BE* generated FW➔HW homeoses associated with a gain of
UbdA antigenicity in forewings, with no effects in the other direction, in stark contrast with other targets
(**Fig. 9B**). This suggests a possible de-insulation of the TAD boundary in the crispant clones, resulting in
a TAD fusion or in a long-range interaction between a T2-specific enhancer and *Ubx* promoter. Similar

de-insulating effects of deletion alleles have been described at the *Notch* locus in *Drosophila* (Arzate-
Mejía et al. 2020), in digit-patterning mutants in mice and humans (Lupiáñez et al. 2015; Anania et al.
2022), or at murine and fly *Hox* loci depleted of CTCF-mediated regulatory blocking (Narendra et al.
2015; Gambetta and Furlong 2018; Kyrchanova et al. 2020). It will be interesting to profile the binding
of insulator proteins and transcriptional repressors across the butterfly *Hox* TAD landscape to shed
more light onto the mechanisms of *Ubx* insulation, using *in vivo* assays (Bowman et al. 2014), or *in silico*
predictions that take advantage of updated binding matrices for insect insulator proteins (Mitra et al.
2018). Of note, our CRISPR target is adjacent to an hindwing-enriched peak that also presented CTCF
binding sites (**Fig. 4A**). Following a similar logic, this site could be a candidate insulator specific to
Ubx-expressing tissues like the hindwing, a hypothesis that will require further testing.
435
436
Making sense of spontaneous wing homeotic mutants
In this article, we documented a large sample of spontaneous homeotic mutants obtained in *Heliconius*
spp. All homeotic clones were limited to the wing posterior compartments (*ie*. posterior to the M$_2$ vein),
possibly because of parasegmental, compartment-specific regulatory domains that played historic roles
in the study of *Drosophila* BX-C regulation (Maeda and Karch 2015). Sibatani documented in
Lepidoptera that "*the mosaics of F/H homeosis tend to occur most frequently in the posterior half of*
the wing, the boundary of the anterior and posterior halves occurring somewhere in space M$_1$-M$_2$"
(Sibatani 1983). Our collection of spontaneous *Heliconius* mutants only displayed clones in posterior
regions, consistently with this trend. However, our CRISPR perturbation assays of *J. coenia* and *V. cardui*
cis-regulatory regions also generated anterior clones, with all targets. Deciphering how butterfly *Ubx*
regulation is compartmentized between parasegmental or wing antero-posterior domains will require
additional investigation. Most *Heliconius* homeoses were in the hindwings (*ie*. putative *Ubx* loss-of-
expression clones), and among these, all but one were ventral (**Fig. S9**). Three mutants showed forewing
homeoses (*ie*. putative *Ubx* gain-of-expression clones), two of them ventral and one of them dorsal (**Fig.**
8). The systematic reviews of wing homeosis in Lepidoptera found that forewing homeoses are almost
as common as hindwing ones (Sibatani 1980, 1983). Our mutational interrogation assays, while coarse
in nature, revealed the existence of activating and repressing *cis*-regulatory sequences at the *Ubx* locus
itself. Spontaneous FW🔄HW homeoses observed in butterflies and moths may thus result from somatic
mutations or transposition events at this locus.
456
457 **MATERIALS AND METHODS**
458
Genome annotations and transcriptomic analysis
Nymphalid genome sequences of the *Hox* cluster and their annotations were extracted from the NCBI
Assembly and Lepbase online repositories (Challis et al. 2016; Kitts et al. 2016) as follows : *V. cardui*
from NCBI *ilVanCard2.1* and LepBase *Vc_v1* ; *A.* (*Nymphalis*) *io* from NCBI *ilAglIoxx1.1*; *J. coenia* from
Lepbase *Jc_v2*; *P xylostella* from NCBI *ilPluXylo3.1*. The *Ubx* regions from *ilVanCard2.2, Vc_v1, and Jc_v2*
were manually re-annotated using wing transcriptome data on the NCBI SRA (BioProjects
PRJNA661999, PRJNA293289, PRJNA237755, PRJNA385867, and *PRJNA498283*) The genome
sequence of *K. inachus* was obtained from the Dryad repository (Yang et al. 2020). Differential gene
expression analysis across the *K. inachus Ubx* locus were carried out using wing transcriptome data
available on the NCBI SRA (BioProject *PRJNA698433*), following a manual re-annotation of a

preliminary gene models provided by Peiwen Yang and Wei Zhang (Wang et al. 2022). All transcripts
analyses were performed using the *STAR* intron-aware aligner and *DEseq2* expression analysis package
as previously described (Love et al. 2014; Dobin and Gingeras 2016; Hanly et al. 2019, 2022). Expression
levels were calculated as genome-wide normalized counts and pairwise Wald tests were performed to
assess differential expression between forewings and hindwings. Multiple test adjustment was
performed using Benjamini and Hochberg correction.
475

Hi-C and ATAC-seq analyses
Hi-C data from *J. coenia* fifth instar larval forewings and 48-72 h APF pupa hindwings are available at
the NCBI SRA BioProject *PRJNA641138* (van der Burg et al. 2020). Triplicated ATAC-seq datasets for
larval and pupal wing and head tissues of *J. coenia* and *V. cardui* (van der Burg et al. 2019, 2020; Mazo-
Vargas et al. 2022) are available on the NCBI SRA BioProjects *PRJNA497878*, *PRJNA695303*, and
PRJNA559165. All the ATAC-seq and Hi-C data were re-analysed on *J. coenia* and *V. cardui Ubx*
genomic regions as previously described (Mazo-Vargas et al. 2022). Briefly, matrices of interactions
were constructed by mapping paired reads against the *Junonia coenia* genome (Mazo-Vargas et al.,
2022) using *hicBuildMatrix* (Ramírez et al. 2018). Matrices from larvae and pupae were normalized
using *hicNormalize* and corrected with the Knight-Ruiz matrix balancing algorithm.The definitions of
topologically associating domains (TADs) can be influenced by various factors such as the choice of
software, parameters, sequencing depth, and the presence of experimental noise. To ensure reliability, it
is recommended to compare TAD calls with independent datasets, such as histone marks or known
factors associated with TAD boundaries. In the absence of these specific datasets, we employed a
different combination of parameters in the *hicFindTADs* tool and compared the resulting TAD calls. HiC
matrices at 10 kb and 20 kb bin resolutions were utilized, and TAD insulation scores were evaluated
using a false-discovery rate correction for multiple testing, with *p-value* thresholds of 0.01 and 0.005.
Consistent TAD boundary calls with negative TAD separation scores were selected to define domain
limits at 10 kb and 20 kb matrix resolutions.
495

496　**CTCF motif binding predictions**
497　The program *fimo* was used to scan for the *J. coenia* candidate TAD boundary region
498　(HiC_scaffold_12:6430000-6444000) for canonical CTCF binding sites, using the positional weight
499　matrix MA0205.1 deposited in the JASPAR database (Holohan et al. 2007; Cuellar-Partida et al. 2012;
500　Castro-Mondragon et al. 2022).
501

Genomic conservation analyses
We generated whole-genome alignments of 25 Lepidoptera and 2 Trichoptera reference species from
NCBI Assembly using *ProgressiveCactus* (Armstrong et al. 2020), and *HAL tools* (Hickey et al. 2013) for
converting the resulting HAL file to the MAF format. We provided a species topology tree of 23
Lepidoptera species to *PhyloFit* (Hubisz et al. 2011) to fit a multiple sequence alignment on the
reference *J. coenia Ubx* locus, using *HKY85* as the substitution model. Using *PhastCons* (Siepel et al.
2005)*,* we then generated conservation score plots using standard parameters (target-coverage = 0.45;
expected-length = 12; rho = 0.1) stored in BED and WIG file formats.
510

Butterfly rearing and CRISPR microinjections
J. coenia and *V. cardui* colonies were maintained at 25°C and 60-70% relative humidity in a growth
chamber with a 14:10 light:dark photoperiod. Larval rearing on artificial diets, egg collection, and

microinjections followed previously described methods (Martin et al. 2020; Tendolkar et al. 2021).
Cas9:sgRNA heteroduplexes were prepared as previously described (Martin et al., 2020). Frozen
aliquots of Cas9-2xNLS (2.5 μL ; 1,000 ng/μL) and sgRNA (2.5 μL ; 500 ng/μL) were mixed in 2:1 and
4:1:1 mass ratios for single and dual target injections, respectively. CRISPR sgRNA targets are listed in
Table S1.
519
Antibody stainings
Fifth instar wing disks were dissected in ice cold Phosphate Buffer Saline (PBS), fixed for 15-20 min at
room temperature in methanol-free formaldehyde diluted to 4% in PBS / 2mM EGTA (egtazic acid),
washed in PBS with 0.1% Triton X-100 (PT), stored in PT with 0.5% Bovine Serum albumin (PT-BSA),
incubated overnight at 4°C in PT-BSA with a 1:5 dilution of the FP6.87 antibody serum (mouse
monoclonal, Developmental Studies Hybridoma Bank), and washed in PT. A 1:250 dilution of anti-Mouse
IgG antibody coupled to AlexaFluor488 or Rabbit AlexaFluor555 was made in PT-BSA and spun down
at 14,000 rcf to remove aggregates, and incubated with wings for 2 h at room temperature, before
additional washes, incubation in 50% glycerol-PBS with DAPI (4′,6-diamidino-2-phenylindole) nuclear
stain, and incubation and mounting in 60% glycerol-PBS with 2mM of EDTA
(Ethylenediaminetetraacetic acid).
531
Imaging
Full-mount photographs of *J. coenia* and *V. cardui* were taken on a Nikon D5300 digital camera mounted
with an AF-S VR MicroNikkor 105mm f/2.8G lens, with magnified views taken using a Keyence VHX-5000
digital microscope mounted with VH-Z00T and VH-Z100T lenses. Immunofluorescent stainings were
imaged on an Olympus BX53 epifluorescent microscope mounted with UPLFLN 4x/0.13 and 10X/0.3
objectives.
538

ACKNOWLEDGEMENTS
We thank Ling Sheng Loh and the undergraduate researchers from the Martin Lab for assistance with
micro-injections and insect rearing, Rachel Canalichio and the GWU Harlan Greenhouse personnel for
growing host plants, Patricia Hernandez for sharing microscopes, and Alex Wild for assistance with
Heliconius microphotographs at UT Austin. We wish to acknowledge James Lewis and Bob Reed for
stimulating insights on open-chromatin biology and the *Hox* locus, as well as for generating Hi-C
libraries published in previous publications that we re-analyzed here. This work was supported by the
NSF awards IOS-1656553 and IOS-2110534 to AM, the Wilbur V. Harlan Research Fellowship to AT, the
NSF Postdoctoral Research Fellowship in Biology to AMV, and the Smithsonian Institution Biodiversity
Genomics Fellowship to JJH.

REFERENCES

Adikusuma F, Piltz S, Corbett MA, et al (2018) Large deletions induced by Cas9 cleavage. Nature 560:E8–E9

Anania C, Acemel RD, Jedamzick J, et al (2022) In vivo dissection of a clustered-CTCF domain boundary reveals developmental principles of regulatory insulation. Nature Genetics 54:1026–1036

Armstrong J, Hickey G, Diekhans M, et al (2020) Progressive Cactus is a multiple-genome aligner for the thousand-genome era. Nature 587:246–251. https://doi.org/10.1038/s41586-020-2871-y

Arzate-Mejía RG, Josué Cerecedo-Castillo A, Guerrero G, et al (2020) In situ dissection of domain boundaries affect genome topology and gene transcription in Drosophila. Nature communications 11:894

Bowman SK, Deaton AM, Domingues H, et al (2014) H3K27 modifications define segmental regulatory domains in the Drosophila bithorax complex. Elife 3:e02833

Buffry AD, Kittelmann S, McGregor AP (2023) Characterisation of the role and regulation of Ultrabithorax in sculpting fine-scale leg morphology. Frontiers in Cell and Developmental Biology 11:

Castro-Mondragon JA, Riudavets-Puig R, Raulusevicuite I, et al (2022) JASPAR 2022: the 9th release of the open-access database of transcription factor binding profiles. Nucleic acids research 50:D165–D173

Challis RJ, Kumar S, Dasmahapatra KKK, et al (2016) Lepbase: the Lepidopteran genome database. bioRxiv 056994

Connahs H, Tlili S, van Creij J, et al (2019) Activation of butterfly eyespots by Distal-less is consistent with a reaction-diffusion process. Development 146:dev169367

Cuellar-Partida G, Buske FA, McLeay RC, et al (2012) Epigenetic priors for identifying active transcription factor binding sites. Bioinformatics 28:56–62

Cullot G, Boutin J, Toutain J, et al (2019) CRISPR-Cas9 genome editing induces megabase-scale chromosomal truncations. Nature communications 10:1136

Dobin A, Gingeras TR (2016) Optimizing RNA-Seq mapping with STAR. In: Data mining techniques for the life sciences. Springer, pp 245–262

Feng S, Rastogi C, Loker R, et al (2022) Transcription factor paralogs orchestrate alternative gene regulatory networks by context-dependent cooperation with multiple cofactors. Nature Communications 13:1–19

Ferguson L, Marletaz F, Carter J-M, et al (2014) Ancient expansion of the Hox cluster in Lepidoptera generated four homeobox genes implicated in extra-embryonic tissue formation. PLoS Genet 10:e1004698

Gambetta MC, Furlong EE (2018) The insulator protein CTCF is required for correct Hox gene expression, but not for embryonic development in Drosophila. Genetics 210:129–136

Garaulet DL, Lai EC (2015) Hox miRNA regulation within the Drosophila Bithorax complex: Patterning behavior. Mechanisms of development 138:151–159

Gaunt SJ (2022) Seeking sense in the Hox gene cluster. Journal of Developmental Biology 10:48

Gisselbrecht SS, Palagi A, Kurland JV, et al (2020) Transcriptional silencers in drosophila serve a dual role as transcriptional enhancers in alternate cellular contexts. Molecular cell 77:324–337

Guo Y, Xu Q, Canzio D, et al (2015) CRISPR inversion of CTCF sites alters genome topology and enhancer/promoter function. Cell 162:900–910

Gutierrez-Perez I, Rowley MJ, Lyu X, et al (2019) Ecdysone-Induced 3D chromatin reorganization involves active enhancers bound by pipsqueak and polycomb. Cell reports 28:2715–2727

Hajirnis N, Mishra RK (2021) Homeotic Genes: Clustering, Modularity, and Diversity. Frontiers in Cell and Developmental Biology 9:

Hanly JJ, Livraghi L, Heryanto C, et al (2022) A large deletion at the cortex locus eliminates butterfly wing patterning. G3 12:jkac021

Hanly JJ, Wallbank RW, McMillan WO, Jiggins CD (2019) Conservation and flexibility in the gene regulatory landscape of heliconiine butterfly wings. Evodevo 10:1–14

Heffer A, Pick L (2013) Conservation and variation in Hox genes: how insect models pioneered the evo-devo field. Annu Rev Entomol 58:161–179

Hermann A, Kosman D, McGinnis W, Tour E (2022) The expression of Drosophila melanogaster Hox gene Ultrabithorax is not overtly regulated by the intronic long noncoding RNA lncRNA:PS4 in a wild-type genetic background. G3 Genes|Genomes|Genetics 12:jkab374. https://doi.org/10.1093/g3journal/jkab374

Hickey G, Paten B, Earl D, et al (2013) HAL: a hierarchical format for storing and analyzing multiple genome alignments. Bioinformatics 29:1341–1342. https://doi.org/10.1093/bioinformatics/btt128

Holohan EE, Kwong C, Adryan B, et al (2007) CTCF genomic binding sites in Drosophila and the organisation of the bithorax complex. PLoS genetics 3:e112

Hubisz MJ, Pollard KS, Siepel A (2011) PHAST and RPHAST: phylogenetic analysis with space/time models. Brief Bioinform 12:41–51. https://doi.org/10.1093/bib/bbq072

Hughes CL, Kaufman TC (2002) Hox genes and the evolution of the arthropod body plan. Evolution & development 4:459–499

Ibragimov A, Bing XY, Shidlovskii Y, et al (2022) The insulating activity of the Drosophila BX-C chromatin boundary Fub-1 is parasegmentally regulated by lncRNA read-through. bioRxiv

Kahn TG, Savitsky M, Kuong C, et al (2023) Topological screen identifies hundreds of Cp190-and CTCF-dependent Drosophila chromatin insulator elements. Science Advances 9:eade0090

Kaushal A, Dorier J, Wang B, et al (2022) Essential role of Cp190 in physical and regulatory boundary formation. Science Advances 8:eabl8834

Kawahara AY, Plotkin D, Espeland M, et al (2019) Phylogenomics reveals the evolutionary timing and pattern of butterflies and moths. Proceedings of the National Academy of Sciences 116:22657–22663

Kitts PA, Church DM, Thibaud-Nissen F, et al (2016) Assembly: a resource for assembled genomes at NCBI. Nucleic acids research 44:D73–D80

Kosicki M, Tomberg K, Bradley A (2018) Repair of double-strand breaks induced by CRISPR–Cas9 leads to large deletions and complex rearrangements. Nature biotechnology 36:765–771

Kyrchanova O, Maksimenko O, Ibragimov A, et al (2020) The insulator functions of the Drosophila polydactyl C2H2 zinc finger protein CTCF: Necessity versus sufficiency. Science advances 6:eaaz3152

Lewis DL, DeCamillis MA, Brunetti CR, et al (1999) Ectopic gene expression and homeotic transformations in arthropods using recombinant Sindbis viruses. Current Biology 9:1279–1287. https://doi.org/10.1016/S0960-9822(00)80049-4

Lewis JJ, Reed RD (2018) Genome-wide regulatory adaptation shapes population-level genomic landscapes in Heliconius. Molecular biology and evolution

Livraghi L (2017) Hox3 duplication and divergence in the Lepidoptera. PhD Thesis, Oxford Brookes University

Livraghi L, Martin A, Gibbs M, et al (2017) CRISPR/Cas9 as the Key to Unlocking the Secrets of Butterfly Wing Pattern Development and Its Evolution. Advances in Insect Physiology

Lohse K, Mackintosh A, Vila R, Consortium DT of L (2021a) The genome sequence of the European peacock butterfly, Aglais io (Linnaeus, 1758). Wellcome Open Research 6:258

Lohse K, Wright C, Talavera G, et al (2021b) The genome sequence of the painted lady, Vanessa cardui Linnaeus 1758. Wellcome Open Research 6:324

Loker R, Sanner JE, Mann RS (2021) Cell-type-specific Hox regulatory strategies orchestrate tissue identity. Current Biology 31:4246–4255

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome biology 15:1–21

Lupiáñez DG, Kraft K, Heinrich V, et al (2015) Disruptions of topological chromatin domains cause pathogenic rewiring of gene-enhancer interactions. Cell 161:1012–1025

Maeda RK, Karch F (2015) The open for business model of the bithorax complex in Drosophila. Chromosoma 124:293–307

Mahfooz N, Turchyn N, Mihajlovic M, et al (2007) Ubx regulates differential enlargement and diversification of insect hind legs. PLoS One 2:e866

Mallo M, Alonso CR (2013) The regulation of Hox gene expression during animal development. Development 140:3951–3963

Martin A, Wolcott NS, O'Connell LA (2020) Bringing immersive science to undergraduate laboratory courses using CRISPR gene knockouts in frogs and butterflies. Journal of Experimental Biology 223:

Masumoto M, Yaginuma T, Niimi T (2009) Functional analysis of Ultrabithorax in the silkworm, Bombyx mori, using RNAi. Development genes and evolution 219:437–444

Matsuoka Y, Monteiro A (2018) Melanin pathway genes regulate color and morphology of butterfly wing scales. Cell reports 24:56–65

Matsuoka Y, Monteiro A (2021) Hox genes are essential for the development of eyespots in Bicyclus anynana butterflies. Genetics 217:iyaa005

Matsuoka Y, Monteiro A (2022) Ultrabithorax modifies a regulatory network of genes essential for butterfly eyespot development in a wing sector-specific manner. Development 149:dev200781

Matsuoka Y, Murugesan SN, Prakash A, Monteiro A (2022) Lepidopteran prolegs are novel traits, not leg homologs. bioRxiv

Mazo-Vargas A, Langmüller AM, Wilder A, et al (2022) Deep cis-regulatory homology of the butterfly wing pattern ground plan. Science 378:304–308

Méndez-González ID, Williams TM, Rebeiz M (2023) Changes in locus wide repression underlie the evolution of Drosophila abdominal pigmentation. PLoS genetics 19:e1010722

Merabet S, Carnesecchi J (2022) Hox dosage and morphological diversification during development and evolution. In: Seminars in Cell & Developmental Biology. Elsevier

Mitra S, Biswas A, Narlikar L (2018) DIVERSITY in binding, regulation, and evolution revealed from high-throughput ChIP. PLoS Computational Biology 14:e1006090

Moniot-Perron L, Moindrot B, Manceau L, et al (2023) The Drosophila Fab-7 boundary modulates Abd-B gene

activity by guiding an inversion of collinear chromatin organization and alternate promoter use. Cell Reports 42:111967

Mulhair P, Crowley L, Boyes DH, et al (2022) Diversity, duplication, and genomic organization of homeobox genes in Lepidoptera. Genome Research gr-277118

Mulhair PO, Holland PW (2022) Evolution of the insect Hox gene cluster: Comparative analysis across 243 species. In: Seminars in Cell & Developmental Biology. Elsevier

Narendra V, Rocha PP, An D, et al (2015) CTCF establishes discrete functional chromatin domains at the Hox clusters during differentiation. Science 347:1017–1021

Natoli G, Andrau J-C (2012) Noncoding transcription at enhancers: general principles and functional models. Annual review of genetics 46:1–19

Nijhout HF, Rountree DB (1995) Pattern induction across a homeotic boundary in the wings of Precis coenia (Hbn.) (Lepidoptera: Nymphalidae). International Journal of Insect Morphology and Embryology 24:243–251

Owens DD, Caulder A, Frontera V, et al (2019) Microhomologies are prevalent at Cas9-induced larger deletions. Nucleic acids research 47:7402–7417

Pace RM, Grbić M, Nagy LM (2016) Composition and genomic organization of arthropod Hox clusters. EvoDevo 7:1–11

Pang B, van Weerd JH, Hamoen FL, Snyder MP (2022) Identification of non-coding silencer elements and their regulation of gene expression. Nature Reviews Molecular Cell Biology 1–13

Paul R, Giraud G, Domsch K, et al (2021) Hox dosage contributes to flight appendage morphology in Drosophila. Nature Communications 12:2892

Pease B, Borges AC, Bender W (2013) Noncoding RNAs of the Ultrabithorax domain of the Drosophila bithorax complex. Genetics 195:1253–1264

Postika N, Metzler M, Affolter M, et al (2018) Boundaries mediate long-distance interactions between enhancers and promoters in the Drosophila Bithorax complex. PLoS genetics 14:e1007702

Prasad N, Tarikere S, Khanale D, et al (2016) A comparative genomic analysis of targets of Hox protein Ultrabithorax amongst distant insect species. Scientific reports 6:27885

Ramírez F, Bhardwaj V, Arrigoni L, et al (2018) High-resolution TADs reveal DNA sequences underlying genome organization in flies. Nature communications 9:189

Refki PN, Armisén D, Crumière AJJ, et al (2014) Emergence of tissue sensitivity to Hox protein levels underlies the evolution of an adaptive morphological trait. Developmental biology 392:441–453

Savitsky M, Kim M, Kravchuk O, Schwartz YB (2016) Distinct roles of chromatin insulator proteins in control of the Drosophila bithorax complex. Genetics 202:601–617

Seal RL, Tweedie S, Bruford EA (2022) A standardised nomenclature for long non-coding RNAs. IUBMB life

Segert JA, Gisselbrecht SS, Bulyk ML (2021) Transcriptional silencers: Driving gene expression with the brakes on. Trends in Genetics 37:514–527

Shin HY, Wang C, Lee HK, et al (2017) CRISPR/Cas9 targeting events cause complex deletions and insertions at 17 sites in the mouse genome. Nature communications 8:15464

Shippy TD, Ronshaugen M, Cande J, et al (2008) Analysis of the Tribolium homeotic complex: insights into mechanisms constraining insect Hox clusters. Development genes and evolution 218:127–139

Sibatani A (1980) Wing homoeosis in Lepidoptera: a survey. Developmental Biology 79:1–18

Sibatani A (1983) A compilation of data on wing homoeosis in Lepidoptera. J Res Lepidop 22:1–46

Siepel A, Bejerano G, Pedersen JS, et al (2005) Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. Genome Res 15:1034–1050. https://doi.org/10.1101/gr.3715005

Srinivasan A, Mishra RK (2020) Lessons on gene regulation learnt from the Drosophila melanogaster bithorax complex. International Journal of Developmental Biology 64:151–158

Stadler MR, Haines JE, Eisen MB (2017) Convergence of topological domain boundaries, insulators, and polytene interbands revealed by high-resolution mapping of chromatin contacts in the early Drosophila melanogaster embryo. Elife 6:e29550

Statello L, Guo C-J, Chen L-L, Huarte M (2021) Gene regulation by long non-coding RNAs and its biological functions. Nature reviews Molecular cell biology 22:96–118

Tendolkar A, Pomerantz AF, Heryanto C, et al (2021) Ultrabithorax Is a Micromanager of Hindwing Identity in Butterflies and Moths. Front Ecol Evol 9:. https://doi.org/10.3389/fevo.2021.643661

Thomas JA, Frandsen PB, Prendini E, et al (2020) A multigene phylogeny and timeline for Trichoptera (Insecta). Systematic Entomology 45:670–686

Tomoyasu Y (2017) Ultrabithorax and the evolution of insect forewing/hindwing differentiation. Current opinion in insect science 19:8–15

Tong X, Hrycaj S, Podlaha O, et al (2014) Over-expression of Ultrabithorax alters embryonic body plan and wing patterns in the butterfly Bicyclus anynana. Developmental biology 394:357–366

Tong XL, Fu MY, Chen P, et al (2017) Ultrabithorax and abdominal-A specify the abdominal appendage in a
dosage-dependent manner in silkworm, Bombyx mori. Heredity 118:578
Van Belleghem SM, Ruggieri AA, Concha C, et al (2023) High level of novelty under the hood of convergent
evolution. Science 379:1043–1049
van der Burg KR, Lewis JJ, Brack BJ, et al (2020) Genomic architecture of a genetically assimilated seasonal
color pattern. Science 370:721–725
van der Burg KR, Lewis JJ, Martin A, et al (2019) Contrasting roles of transcription factors Spineless and EcR in
the highly dynamic chromatin landscape of butterfly wing metamorphosis. Cell reports 27:1027–1038
Wang H, Hu H, Xiang Z, et al (2019) Identification and characterization of a new long noncoding RNA iab-1 in the
Hox cluster of silkworm, Bombyx mori identification of iab-1. Journal of Cellular Biochemistry 120:17283–
17292. https://doi.org/10.1002/jcb.28990
Wang H, Tong X, Liu M, et al (2017) Fine mapping of a degenerated abdominal legs mutant (Edl) in silkworm,
Bombyx mori. Plos one 12:e0169224
Wang S, Teng D, Li X, et al (2022) The evolution and diversification of oakleaf butterflies. Cell 185:3138–3152
Warren RW, Nagy L, Selegue J, et al (1994) Evolution of homeotic gene regulation and function in flies and
butterflies. Nature 372:458
Weatherbee SD, Frederik Nijhout H, Grunert LW, et al (1999) Ultrabithorax function in butterfly wings and the
evolution of insect wing patterns. Current Biology 9:109–115. https://doi.org/10.1016/S0960-
9822(99)80064-5
Weatherbee SD, Halder G, Kim J, et al (1998) Ultrabithorax regulates genes at several levels of the wing-
patterning hierarchy to shape the development of the Drosophila haltere. Genes & Development 12:1474–
1482. https://doi.org/10.1101/gad.12.10.1474
Wootton RJ (1993) Leading edge section and asymmetric twisting in the wings of flying butterflies (Insecta,
Papilionoidea)
Yang J, Wan W, Xie M, et al (2020) Chromosome-level reference genome assembly and gene editing of the
dead-leaf butterfly Kallima inachus. Molecular Ecology Resources 20:1080–1092
Zhang L, Steward RA, Wheat CW, Reed RD (2021) High-quality genome assembly and comprehensive
transcriptome of the painted lady butterfly Vanessa cardui. Genome biology and evolution 13:evab145
Zhang X, Wang W, Zhu W, et al (2019) Mechanisms and functions of long non-coding RNAs at multiple regulatory
levels. International journal of molecular sciences 20:5573
Zheng Z, Khoo A, Fambrough Jr D, et al (1999) Homeotic gene expression in the wild-type and a homeotic
mutant of the moth Manduca sexta. Development genes and evolution 209:460–472
Zuccaro MV, Xu J, Mitchell C, et al (2020) Allele-specific chromosome removal after Cas9 cleavage in human
embryos. Cell 183:1650–1664

SUPPLEMENTARY INFORMATION

Species	sgRNA name	Target Sequence (5' to 3') PAM sequence not shown
J. coenia	*Antp-Ubx_BE*	CTCGAATATGGAGATATCGG
	UbxCRE11a3	ACGGACCTCCGCTTTCCTGG
	UbxCRE11c6	AACTGGTGCAGTGCCTTGTA
J. coenia + V. cardui	*UbxCRE11a2*	CTACTCTGTTCGGACATTCG
	UbxCRE11c5	GCTGCCGCGAGTCTGAATCG
	UbxCRE11b9	TTCATGTATGAACCATGACG
	UbxIT1#1	CCTTCGCATAAGTTCGGATAGG
	Bxd1	TATCGGTCGTTCGTCACACA
V. cardui	*UbxIT1#2*	CTCGGCTATGTGTCGAGGGC

762 **Table S1. List of sgRNAs used in CRISPR experiments.**

766
Figure S1. Prediction of two conserved CTCF binding sites at *Antp-Ubx_BE*. (**A**) Sequence-level view of a
180-bp genomic interval including the *Antp-Ubx_BE* sgRNA (turquoise) in *J. coenia*, overlapping with an ATAC-seq
peak of forewing-enriched chromatin opening (red). The CRISPR target is about 100 bp away from two predicted
binding sites for the *Drosophila* CTCF insulator protein. (**B**) High-level of nucleotide conservation at thesgRNA site
and CTCF motifs across Lepidoptera and Trichoptera representative genomes, indicative of functional constraints
on these sequences.

773
774 **Figure S2. CRISPR perturbation of the *Antp-Ubx* boundary element results in FW-to-HW homeosis.** (**A**)
775 Example of an *Antp-Ubx_BE* crispant with a unilateral phenotype on the right forewing. (**B**) Additional examples of
776 forewing homeoses in *Antp-Ubx_BE* crispant. Wing sets (forewing mKO mutants and corresponding contralateral
777 WT) are shown with one of the wings horizontally flipped to show the mutant wings in left-to-right orientation..
778 Cyan arrows : small mutant clones. Cyan asterisks : large mutant clones.
779
780

Figure S3. Pupal defects following FW→HW homeosis in *Antp-Ubx_BE* crispants. (**A-B**) Contralateral (CL) and, forewing mosaic knockout (mKO) mutants following CRISPR targeting of *Antp-Ubx_BE* in *J. coenia*. The two pupae show suture defects in the midline appendages (arrows). (**A'-B'**) Magnified views of the crispant forewings, showing defective cuticle (arrowheads). (**C-C'**) Crispant adult butterfly emerged from the pupa in panel B. White arrowheads in C' highlight the match between dorsal forewing clones and the pupal forewing cuticle defects shown in B'. Scale bars : 1 mm.

788
789 **Figure S4. Additional examples of ectopic UbdA and FW→HW homeosis in *Antp-Ubx_BE* crispant larval**
790 **forewings.** (**A-F**) Each panel shows forewings with ectopic detection of UbdA (FP6.87 monoclonal antibody,
791 green), dissected at the fifth instar stage. Panels D and F are wing sets from individual crispants. Panels E and C
792 are mutant contralateral wings of the mutant forewings shown in Figs. 4D and E, respectively.
793
794

Ubx_IT1

796
797 **Figure S5. Additional mutant phenotypes from CRISPR-mediated interrogation of *lncRNA_Ubx-IT1* 5'**
798 **region in *J. coenia* (top) and *V. cardui* (bottom).** Cyan arrows : mutant clones.
799

Ubx_AS5'

800
801 **Figure S6. Additional mutant phenotypes from CRISPR-mediated interrogation of the *lncRNA_Ubx-A*S5'**
802 **region in *J. coenia* and *V. cardui*.** Cyan arrows : mutant clones. Cyan arrowheads : white eyespot foci.

803
804
805

806
807 **Figure S7. Additional mutant phenotypes from CRISPR-mediated interrogation of *CRM11* in *J. coenia* and**
808 ***V. cardui* show bidirectional homeoses and non-homeotic eyespot changes.** Cyan arrows : mutant clones.
809 Cyan arrowheads : white eyespot foci.

810
811 **Figure S8. CRISPR perturbation of the conserved *Ubx_CRE11b* results in HW➞FW homeoses**. (**A-B**) The
812 *UbxCRE11b9* sgRNA targets a hindwing-enriched ATAC peak with strong conservation across genomes from 23
813 Lepidoptera and 2 Trichoptera species (gray : PhastCons scores). Colored bars denote variation from the *J. coenia*
814 reference (**C**) *Jc_UbxCRE_11b9* crispant butterflies exclusively showed HW➞FW transformed clones (cyan arrows in
815 both *J. coenia* and *V. cardui*).
816
817
818
819

Figure S9 (previous page, continued above). Hindwing homeoses in *Heliconius* butterfly spontaneous mutants

from pure stocks, hybrid cultures and wild-caught individuals from the L.E. Gilbert collection (UT Austin). White

arrowheads: homeotic clones including the acquisition of ventral forewing coupling scales. Asterisks : local

deformation of hindwings relative to wild-type. All hindwing homeoses are ventral except in panel L. **A.** *Heliconius*

cydno galanthus x H. melpomene rosina (Costa Rica), cross J31, August 1987. **B.** *Heliconius cydno gustavi*,

captive stock from Saladito (Colombia), September 1991. **C.** *Heliconius melpomene madeira* (Brazil) *x Heliconius*

melpomene plesseni (Ecuador), September 2012. **D.** *H. m. rosina* (Costa Rica) *x Heliconius melpomene madeira*

(Brazil) *x H. cydno galanthus* (Costa Rica) mixed population, December 2015. **E.** *H. m. rosina,* captive stock from

Osa Peninsula (Costa Rica), September 1991. **F.** *Heliconius hewitsoni,* captive stock from Osa Peninsula (Costa

Rica), July 2005. **G.** *Heliconius cydno cydnides*, captive stock from natural hybrid zone in Dagua Pass (Colombia),

May 1989. **H.** *H. m. rosina* (Costa Rica) *x H. m. madeira* (Brazil) *x H. c. galanthus* (Costa Rica) mixed population,

June 2016. **I.** *H. c. galanthus x H. m. rosina* crossed three times, and back to *H. c. galanthus,* August 2014. **J.**

Heliconius melpomene malleti (Ecuador) x *H. m. plesseni* (Ecuador) hybrid stock, 2010. **K.** *H. m. rosina* captive

stock, Costa Rica. **L.** *H. m. rosina* captive stock, Osa Peninsula (Costa Rica), March 1987, in dorsal view.

836