Plasticity of circadian and circatidal rhythms in activity and

transcriptomic dynamics in a freshwater snail

Takumi Yokomizo[1] and Yuma Takahashi[1]*

[1]Graduate School of Science, Chiba University, Chiba 263-8522, Japan

*Corresponding author: Yuma Takahashi

Address: 1-33, Yayoi, Inage, Chiba, Chiba, Japan

Email: takahashi.yum@gmail.com

Abstract

Organisms have diverse biological clocks synchronized with environmental cycles depending on their habitats. Anticipation of tidal changes has driven the evolution of circatidal rhythms in some marine species. In a freshwater snail, *Semisulcospira reiniana*, individuals in nontidal areas exhibit circadian rhythms, whereas those in tidal areas exhibit both circadian and circatidal rhythms. We investigated whether the circatidal rhythms are genetically determined or induced by environmental cycles. After exposure to a simulated tidal cycle, both tidal and nontidal populations exhibited circatidal rhythms in locomotive activity. Transcriptome analysis revealed that genes with circatidal oscillation increased due to entrainment to the tidal cycle in both populations and dominant rhythmicity was consistent with the environmental cycle. These results suggest plasticity in the endogenous rhythm in both populations. However, circatidal oscillating genes were more abundant in the tidal population than in the nontidal population, suggesting that a greater number of genes are associated with circatidal clocks in the tidal population compared to the nontidal population. Our findings suggest that the plasticity of biological rhythms and genetic changes in the molecular network between clocks and downstream genes may have contributed to the adaptation to the tidal environment in *S. reiniana*.

Keywords: endogenous rhythm, biological clock, phenotypic plasticity, tidal environment, transcriptome

Introduction

Organisms have faced the challenge of adapting to periodic variations in environments since ancient times. The rotation and inertial force of Earth, along with the gravitational forces of the sun and moon, generate environmental variations across multiple timescales, including day–night, tidal, lunar, and seasonal cycles. To cope with these fluctuations, organisms possess self-sustained, temperature-compensated timekeeping systems capable of entraining to environmental cues [1–3]. The circadian rhythm, observed in a wide range of organisms, ranging from bacteria [4] to mammals [5–7], allows for adaptation to day–night cycles. Biological clocks enable organisms to anticipate environmental changes and regulate physiological processes with appropriate timing. The coordination between environmental cycles and biological clock parameters, such as phase and period, plays a vital role in enhancing fitness and facilitating adaptation to rhythmic environments [8–10].

Synchronizing biological rhythms with environmental cycles has adaptive significance and can result in variations in endogenous rhythms depending on the circumstances. For instance, studies on the model fungus *Neurospora discreta* have shown that local adaptation to different rhythmic environments, regulated by the circadian clock, leads to habitat-specific endogenous rhythms and influences reproductive success [11]. Variations in endogenous rhythms can arise not only from local adaptation but also from plasticity driven by social roles. Honeybees, for example, exhibit social environment-dependent plasticity in their circadian rhythm within a single population [12,13]. These findings suggest that genetic and/or nongenetic changes in biological rhythms contribute to enhancing individual performance.

Organisms residing in marine or intertidal habitats encounter substantial and

complex environmental changes, including fluctuations in water level, salinity,

temperature, light, and food availability, driven by daily and tidal cycles. The circatidal

rhythm, an approximately 12-h biological rhythm synchronized with tides, is observed

in diverse taxa [14–17]. Owing to the critical importance of anticipating tidal changes

for marine species, the selective pressure exerted by the tidal cycle can drive the

adaptive evolution of circatidal clocks. The relationship between circadian and

circatidal rhythms, characterized by different periodicities and zeitgebers, remains

controversial. For example, studies on the mangrove cricket suggest that the circatidal

timekeeping system operates independently of circadian clocks [18,19]. This

dissociation in the circadian and circatidal clocks of mangrove crickets may be

attributed to the adaptive evolution of a novel timekeeping system that emerged in their

mangrove habitat. Conversely, investigations of various marine species have

demonstrated plasticity in rhythm expression within individuals or populations under

different environmental conditions [20–22]. Notably, certain circadian clock genes in

the Pacific oyster *Crassostrea gigas* exhibit oscillation at tidal frequencies under tidal

conditions [23,24]. In the amphipod crustacean *Parhyale hawaiensis*, the core circadian

clock gene *Bmal1* is suggested to be essential for circatidal rhythms [25]. These studies

give rise to the hypothesis that the circadian and circatidal rhythms may be generated by

a single biological timekeeping system, with circatidal clocks sharing some molecular

components with circadian clocks. The expression of circatidal rhythms and successful

adaptation to tidal environments likely involve a combination of plasticity and genetic

changes.

Rivers present complex rhythmic environments due to diurnal and tidal cycles.

Although a river follows a 24-h light–dark (LD) cycle throughout its course,

downstream areas experience an environmental cycle with a period of 12.4 hours due to

the tidal cycle. To investigate variations in endogenous rhythm across habitats with

distinct environmental cycles, we focused on the freshwater snail *Semisulcospira*

reiniana, which inhabits both nontidal and tidal areas of rivers. Our previous study

revealed that snails in nontidal areas exhibit circadian rhythms, whereas those in tidal

areas exhibit both circadian and circatidal rhythms [26]. However, it remains unclear

whether the differential rhythmicity arises from evolutionary differentiation or plasticity

in response to environmental cycles. In the present study, we evaluated genetic and

nongenetic changes in the endogenous rhythm of snails living in habitats characterized

by heterogeneous environmental cycles. We exposed snails to a simulated tidal

environment in the laboratory and examined the activity and transcriptome rhythms of

individuals from both tidal and nontidal populations.

Materials and Methods

Species and sampling sites

$\qquad$ *Semisulcospira reiniana*, a common freshwater snail species found in Japan,

was collected from nontidal (3–5 m above sea level; 35° 15′ 15″ N, 136° 41′ 09″ E) and

tidal areas (<1 m above sea level; 35° 08′ 49″ N, 136° 40′ 37″ E) of the Kiso River in

June and November 2021. The two populations were approximately 20 km apart along

the river. The collected snails were acclimated in the laboratory under 23°C and

12–12-h LD conditions in freshwater for at least one month.

Entrainment to the simulated tidal cycle

After acclimation, snails from the tidal and nontidal populations were divided

into control and treatment groups. The treatment group individuals were exposed to a

simulated tidal cycle by raising and lowering the water level in a water tank ($60 \times 30 \times$

45 cm; Fig. S1). This tidal simulation was conducted twice in the laboratory under 23°C

and 12–12-h LD (12L12D) conditions. Containers ($\varphi 7 \times 9$ cm) with water vents were

placed 32 cm from the bottom of the tank. Four snails were placed in each container.

For the high tide condition, the water level was filled to a depth of 39 cm, submerging

the snails completely. To simulate the low tide condition, the water level was dropped to

a depth of 24 cm, exposing the snails completely to air. Water supply and draining were

controlled by timers. In the first simulation, water supply switched to drainage at 03:00

and 15:00, and drainage switched to supply at 09:00 and 21:00. In the second simulation,

water supply switched to drainage at 05:00 and 17:00, and drainage switched to supply

at 11:00 and 23:00. The entrainment treatment lasted for four weeks. Snails from the

first tidal simulation were used for behavioural observations, whereas those from the

second simulation were used for gene expression analysis. Individuals in the control

group were kept in freshwater without water level oscillation under 12L12D conditions.

Behavioural observation

To examine the effect of 12-h water level oscillation on the endogenous activity

rhythm of *S. reiniana*, individuals from the control and treatment groups in both

nontidal and tidal populations were observed under constant laboratory conditions.

Eight snails per population were individually placed in containers (13×11.7 cm, 4.7 cm

in height) filled with freshwater. The snails were observed under constant darkness

(DD) at a temperature of 23°C in the laboratory for 96 and 84 h in the control and

treatment groups, respectively. A camera malfunction resulted in a shorter observation

period for the treatment group compared with the control group. Water circulation was

maintained using a pump to keep the water clean. Images were captured every 30 s

using a model 400-CAM061 camera (Sanwa Direct, Japan) and were used to create

time-lapse movies. UMATracker tracking software [27] was used to track the positions

of snails. After correcting major tracking errors, the coordinates of the snails were

averaged every three frames to minimize the influence of small deviations. The total

locomotion distance travelled in 1 h was calculated using the snail trajectories.

RNA sampling

To investigate the effect of a 12-h water level oscillation on the endogenous

gene expression rhythm of *S. reiniana*, time-course RNA sequencing (RNA-seq) was

conducted using individuals from the control and treatment groups in both nontidal and

tidal populations. Snails were preserved in containers without water under the DD

condition at a constant temperature (23°C) for tissue collection. Starting at subjective

high tide, three individuals per population were dissected every 3.1 h for a total of 49.6

h (17 sampling time points) in both control and treatment groups. The epidermis of each

individual was preserved in 750 μL of RNA*later* Stabilization Solution (Invitrogen).

The samples were stored at 4°C for approximately 3 h and then stored at −80°C until

total RNA extraction.

RNA extraction and sequencing

Total RNA was extracted using the Maxwell 16 LEV Plant RNA Kit with the

Maxwell 16 Research Instrument (Promega, USA) following the manufacturer's

instructions. Electrophoresis was performed on a 1% agarose gel to assess RNA

degradation. RNA concentrations were estimated using the Qubit 2.0 fluorometer

(Invitrogen), and RNA purity was assessed using the NanoDrop Lite Spectrophotometer

(Thermo Scientific). Total RNA from three individuals in the same population at each

sampling time point was pooled in equal amounts for pooled RNA-seq. A cDNA library

was constructed using the TruSeq RNA Sample Prep Kit, and paired-end (150 bp)

pooled RNA-seq was performed on the Illumina NovaSeq6000 platform. Adaptor

sequences and low-quality reads were removed using Trimmomatic (version 0.38) [28],

and FastQC (version 0.11.8; http://www.bioinformatics.babraham.ac.uk/projects/fastqc/)

was used for quality control. The remaining high-quality reads from all samples were

used for *de novo* assembly via Trinity software (version 2.9.1) [29]. To estimate gene

expression levels, reads from each sample were mapped to the reference transcripts

using RSEM software (version 1.3.0) [30] to obtain the transcripts per million (TPM)

for each gene. The reference transcripts were used to create supertranscripts, and a

BLAST search was performed against all protein sequences of *C. gigas* [31] to identify

homologs for each gene with the best hit and an e-value <0.0001. Genes showing

circatidal oscillation were analyzed based on the identified homologs.

Statistical analyses

The activity rhythm of individuals in the control and treatment groups was

examined using Lomb–Scargle periodogram analysis via ActogramJ, a software

package based on ImageJ for chronobiological data analysis and visualization [32]. The

mean locomotor distance of surviving individuals was used for rhythm detection.

Owing to the limited movement of individuals in the treatment group at the end of the

observation, activity data from 72 h after the onset were used for the periodogram

analysis. For transcriptome analysis, genes with low expression (average TPM ≤ 1) or

small oscillation amplitude (peak/trough ≤ 1.3) were excluded. Probabilistic principal

component analysis (PCA) was performed to summarize the gene expression pattern of

each sample. After removing an outlier sample, Tukey's honestly significant difference

(HSD) test was conducted to identify differences in expression patterns between

populations or entrainment treatments. Gene expression rhythmicity was analyzed using

RAIN [33], a nonparametric algorithm for identifying rhythmic components in large

biological datasets. Circatidal (12.4 ± 3.1 h) and circadian (24.8 ± 3.1 h) oscillating

genes ($p < 0.01$) were identified. A differential rhythmicity score (S_{DR}) was calculated

for each gene to assess the differential rhythmicity of gene expression between the

control and treatment groups. Based on periodicity analysis performed via RAIN, genes

with a P-value of circatidal periodicity <1 were used for the calculation of S_{DR}, which

was defined as follows:

$$S_{DR} = \frac{Z_P + Z_R}{\sqrt{2}},$$

where Z_P and Z_R represent the Z-scores for changes in periodicity and

amplitude between groups, respectively [34]. Changes in periodicity were calculated as

log (P-value of the control group) – log (P-value of the treatment group), whereas

changes in amplitude were calculated as $\log_2$ (amplitude of the treatment group /

amplitude of the control group). The amplitude of each gene expression was defined as

the difference between the maximum and minimum TPM values. The P-value for S_{DR}

was computed using a Gaussian distribution fitted to the empirical distribution, and it

was adjusted using the Benjamini & Hochberg procedure for multiple testing. To assess

the relative significance of circadian and circatidal rhythms in the transcriptome, the

log-scaled ratio of the P-value for circadian periodicity to that for circatidal periodicity

was calculated for each gene. Functional enrichment analysis of biological processes in

circatidal oscillating genes or differential rhythmic genes based on S_{DR} was performed

using Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway

over-representation analysis and KEGG gene set enrichment analysis (KEGG-GSEA)

conducted through clusterProfiler [35]. Circatidal oscillating genes identified via RAIN

in the treatment group were used as the test set for KEGG pathway over-representation

analysis. The gene list for GSEA was generated based on S_{DR}. All statistical analyses

were conducted using R version 4.2.2.

Results

Activity rhythm

Snails exhibited rhythmic activity patterns under DD conditions in the

laboratory, with a gradual decrease in activity observed in the treatment group (Figs. 1

and 2). In the control group, individuals from the nontidal and tidal populations

exhibited a circadian activity rhythm with periods of 21.6 h and 22.5 h, respectively,

under laboratory DD conditions (Figs. 1a, b). The tidal population did not exhibit the

circatidal rhythm, indicating synchronization with the laboratory 12L12D condition. In

contrast, the treatment group exhibited a circatidal activity rhythm with periods of 12.4

h and 11.9 h in the nontidal and tidal populations, respectively, under laboratory DD

conditions (Figs. 2a, b). This indicates entrainment to the simulated tidal cycle, with the

activity peak preceding the expected high tide peak.

Transcriptome rhythm

In total, we obtained 1,578,392 contigs with a mean length of 513 bp. Among

the 954 metazoan core gene orthologs, we identified 948 (99.3%). We obtained 908,817

supertranscripts containing 58,952 contigs annotated against all protein sequences of *C.*

gigas. In the tidal population, 13,929 and 14,508 genes in the control group and

treatment group, respectively, met the criteria of average TPM value >0 and oscillation

amplitude (peak/trough) ≥1.3. In the nontidal population, the corresponding numbers

were 13,448 and 13,997 genes in the control group and treatment group, respectively.

Based on probabilistic PCA, one sample in the control group of the nontidal population

showed a greatly different expression pattern compared with the other samples (Fig. S2)

and was excluded from subsequent analysis. Probabilistic PCA of the remaining

samples indicated significant differences in expression patterns between populations but

not between the control and treatment groups (Fig. 3a; Table 1). In the nontidal

population, we identified 190 (1.4%) and 292 (2.1%) circatidal (12.4 ± 3.1 h) oscillating

genes in the control group and treatment group, respectively (Figs. 3b–d). In the tidal

population, the corresponding numbers were 271 (1.9%) and 340 (2.3%) circatidal

oscillating genes in the control group and treatment group, respectively (Figs. 3b–d).

The proportion of genes oscillating in the circatidal period increased in both populations

with entrainment to the simulated tidal cycle. However, the abundance of oscillating

genes was higher in the tidal population compared with the nontidal population,

regardless of the control or treatment group. We also detected several oscillating genes

with a circadian period. In the nontidal population, 343 (2.6%) and 328 (2.3%)

oscillating genes were found in the control and treatment groups, respectively (Figs.

3b–d). In the tidal population, the corresponding numbers were 420 (3.0%) and 301

(2.1%) in the control and treatment groups, respectively (Figs. 3b–d). Unlike circatidal

oscillating genes, the proportion of genes oscillating in the circadian period decreased in

both populations with entrainment to the simulated tidal cycle. No circadian clock genes

showed circadian or circatidal rhythmicity in the control and treatment groups in both

populations (Figs. S3 and S4).

251   Differential rhythmicity analysis revealed changes in transcriptome rhythmicity

between the control and treatment groups (Fig. S5). Genes in the first quadrant

exhibited clearer periodicity and larger amplitude in the treatment group compared with

the control group, indicating increased circatidal rhythmicity. In the nontidal population,

13 genes showed significantly increased circatidal rhythmicity in the treatment group

(FDR < 0.05; Table S1). In the tidal population, 16 genes showed significantly increased

circatidal rhythmicity in the treatment group (FDR < 0.05; Table S2). Although most

transcripts did not show significant differential rhythmicity in expression patterns in the

tidal and nontidal populations, the log-scaled ratio of the P-value for circadian

periodicity to that for circatidal periodicity changed with entrainment to the tidal cycle

(Fig. 4). In both populations, the log-scaled P-value ratio was negative in the control

groups, indicating dominance of the circadian rhythm over the circatidal rhythm under

LD conditions without a tidal cycle. However, the ratio was positive in the treatment

groups, indicating dominance of the circatidal rhythm over the circadian rhythm after

entrainment to the tidal cycle. KEGG pathway over-representation analysis revealed

that circatidal oscillating genes in the treatment group of the tidal population were

enriched for some pathways, including "Aminoacyl-tRNA biosynthesis" (KO:

crg00970) and "Histidine metabolism" (KO: crg00340) (Fig. 5a). KEGG-GSEA based

on S_{DR} revealed enrichment of the biological process "Ribosome" (KO: crg03010) for

genes with increased circatidal rhythmicity in the tidal population (Fig. 5b). Based on

both KEGG pathway over-representation analysis and KEGG-GSEA, no enriched

pathways were detected in the nontidal population.

Discussion

The coordination of biological processes with environmental cycles is

regulated by biological clocks, such as the circadian clock, which enhances fitness in

organisms in rhythmic environments [36]. Marine and intertidal habitats, characterized

by the tidal cycle, experience complex and drastic environmental variations, unlike

terrestrial and inland water habitats. The freshwater snail *S. reiniana*, found in tidal

areas, exhibits the circatidal rhythm, which is believed to be an adaptation to the tidal

environment [26]. In the present study, we investigated whether the differential

rhythmicity in *S. reiniana* between tidal and nontidal populations is a result of

evolutionary differentiation or plastic expression of rhythms depending on dominant

environmental cycles. We demonstrated that even individuals in the nontidal population

can exhibit the circatidal rhythm through entrainment to the tidal cycle. However, our

results regarding the transcriptome rhythm suggest that more genes or pathways are

influenced by the tidal cycle in the tidal population compared with the nontidal

population.

We first examined the influence of the simulated tidal cycle on the activity

rhythm under constant conditions in each population. The free-running period of

individuals in the control group under DD conditions was shorter than 24 h, regardless

of the population. This short circadian period aligns with Aschoff's rule, which predicts

that diurnal animals have a free-running period longer than 24 h and nocturnal animals,

such as *S. reiniana* [37], have a free-running period shorter than 24 h [38]. Although

individuals in the control group exhibited a circadian activity rhythm, the periodicity

was not distinct. The lack of additional circadian zeitgebers, such as daily temperature

cycles, under the 24-h light condition in the laboratory may have contributed to noise or

ultradian rhythms. The activity rhythms of individuals were markedly altered by

exposure to the simulated tidal cycle. Individuals in the treatment group exhibited a

circatidal activity rhythm regardless of the population, indicating that the circatidal

rhythm was plastically expressed in response to the tidal cycle, even in individuals from

the nontidal population. Our results also suggest that water level fluctuations serve as a

zeitgeber for the circatidal rhythm in *S. reiniana*. The peak of the activity rhythm

slightly differed from that of snails under a natural tidal condition. Snails captured from

tidal areas exhibit a rhythmic activity pattern with a peak that coincides with high tide

[26]. However, individuals in the treatment group in the present study exhibited peaks

preceding the expected high tide. This discrepancy may be attributed to the earlier

arrival of high tide compared with the programmed switching time, as the water supply

rate was adjusted to reach high tide in slightly under 6 h. Periodogram analysis also

detected peaks around 19 h in the activity rhythms of individuals in the treatment group.

The origin of this periodicity, which could result from a shortened circadian rhythm or a

combination of circadian and circatidal rhythms, remains unidentified.

Based on probabilistic PCA, we observed distinguishable gene expression

patterns between populations rather than between the control and treatment groups,

indicating minimal differentiation in transcriptome expression between populations.

Focusing on rhythmic transcripts, we found an increase in the number of circatidal

oscillating genes in both populations following entrainment to the tidal cycle,

accompanied by a decrease in the number of circadian oscillating genes. These results

suggest that the circatidal rhythm is plastically expressed in both populations, and the

reduction in circadian oscillating genes supports the transition from the circadian to the

circatidal rhythm under a single biological timekeeping system. Additionally, we

identified a greater number of circatidal oscillating genes in the tidal population

compared with the nontidal population, implying that adaptation to the tidal

environment may contribute to the increase in circatidal clock–controlled genes in

individuals from the tidal population. We also detected hundreds of circatidal oscillating

genes in the control group of nontidal populations. The expression rhythm with a period

of 12.4 h in these genes might arise from the harmonics of circadian clocks, as observed

in mammals [39–41]. Although we identified several circadian clock genes, none of

these genes exhibited circadian rhythmicity in their expression. This finding is

consistent with previous studies that reported the absence of circadian rhythmic

expression in clock genes in several intertidal organisms [22,42,43]. The interference of

circatidal rhythms with the rhythmic expression of circadian clock genes and weak

peripheral clocks in the epidermis or asynchronization between tissues in the samples

could potentially explain the lack of rhythmic expression in these genes.

To assess the extent to which the tidal cycle influenced circatidal rhythmicity in

the transcriptome, we calculated the S_{DR} for each gene in both populations. The S_{DR}

distribution in the tidal population was similar to that in the nontidal population,

indicating comparable changes in circatidal rhythmicity. The number of genes showing

a significant increase in circatidal rhythmicity did not differ substantially between

populations. These results suggest that the degree of changes in circatidal rhythmicity in

the transcriptome is similar in individuals exposed to the tidal cycle, regardless of

population. Although the majority of genes did not exhibit significant changes in the

strength of circatidal rhythmicity between the control and treatment groups, we

identified 13 and 16 genes with significantly increased rhythmicity in the treatment

group of the nontidal and tidal populations, respectively. These genes are likely

entrained by the simulated tidal cycle, leading to enhanced circatidal rhythmicity.

Additionally, we calculated the ratio of P-values for circadian periodicity to circatidal

periodicity for each gene. Comparing the ratios between the control and treatment

groups revealed that the dominant rhythmicity of the transcriptome aligns with the

environmental cycle.

We conducted KEGG over-representation analysis and KEGG-GSEA to

identify enriched biological processes among the circatidal oscillating genes and

differential rhythmic genes. In the tidal population, we detected three pathways,

whereas no pathways were identified in the nontidal population. Considering the larger

number of circatidal oscillating genes in the tidal population, it is likely that individuals

in the tidal population possess a higher abundance of circatidal clock–controlled genes

and biological processes regulated by circatidal clocks compared with individuals in the

nontidal population. The enrichment of the "Aminoacyl-tRNA biosynthesis" (KO:

crg00970) and "Ribosome" (KO: crg03010) pathways in the circatidal oscillating genes

of the treatment group in the tidal population suggests the involvement of circatidal

rhythms in biological processes related to translation, as aminoacyl-tRNA is responsible

for delivering amino acids for mRNA-guided protein synthesis at the ribosome.

Furthermore, the detection of the "Histidine metabolism" (KO: crg00340) pathway

suggests a potential association between proteins containing histidine and the response

to the tidal cycle.

Our findings on the activity rhythm demonstrate the plasticity of the

endogenous rhythm in individuals from both tidal and nontidal populations. The genetic

basis of the circatidal rhythm may be conserved in the genome, with the tidal zeitgeber

triggering the expression of the circatidal rhythm in individuals from the nontidal

population. The plasticity of the endogenous rhythm likely played a role in the

adaptation of *S. reiniana* to tidal areas in rivers. We have previously revealed minimal

genetic differentiation between the two populations [26], supporting the contribution of

plasticity to range expansion. By investigating other rivers where *S. reiniana* is limited

to nontidal areas, we can further explore the relationship between the plasticity of the

endogenous rhythm and the establishment in tidal areas. Although our findings suggest

the existence of a genetic basis for the circatidal rhythm in individuals from the nontidal

population, it remains unclear whether they possess independent circatidal clocks

alongside circadian clocks or a single timekeeping system governing both circadian and

circatidal rhythms. The latter hypothesis is gaining support, particularly in studies on

biological clocks in marine molluscs [22,24,44]. Further investigations are needed to

elucidate the genetic mechanisms underlying circatidal clocks in freshwater snails.

However, the absence or decrease of circadian rhythm in activity and gene expression

after entrainment to the tidal cycle supports the notion that circadian and circatidal

rhythms are generated by changes in the periodicity of a single biological clock in

response to environmental cycles. Unravelling the molecular relationship between

circadian and circatidal clocks would contribute to testing the debated hypotheses on the

mechanism of circatidal oscillation generation [45–47].

Our study indicates that individuals in the nontidal population also possess the

circatidal clock and express the circatidal rhythm when exposed to the tidal cycle.

However, transcriptome analysis suggests that individuals in the tidal population exhibit

increased regulation through the circatidal clock. Overall, the plasticity of the

endogenous rhythm and subsequent genetic changes in a molecular network involving a

small number of genes may contribute to adaptation to the tidal environment.

Investigating the parallel evolution of biological clocks using individuals from

independent rivers would provide a comprehensive understanding of the mechanisms

underlying adaptation to tidal environments, including both genetic and nongenetic

changes in biological clocks.

399 Acknowledgements

400   This work was supported by research grants JSPS KAKENHI Grant Numbers

401   17H03729 and 20K06822, Fujiwara Natural History Public Interest Incorporated

402   Foundation and Research Institute of Marine Invertebrates to Y.T., and Grant-in-Aid for

403   JSPS Fellows Grant Number 21J20682 to T.Y. Computations were partially performed

404   on the NIG supercomputer at ROIS National Institute of Genetics.

References

References

1. Dunlap JC. 1999 Molecular bases for circadian clocks. *Cell* **96**, 271–290.
(doi:10.1016/S0092-8674(00)80566-8)

2. Bell-Pedersen D, Cassone VM, Earnest DJ, Golden SS, Hardin PE, Thomas TL,
Zoran MJ. 2005 Circadian rhythms from multiple oscillators: lessons from
diverse organisms. *Nat Rev Genet* **6**, 544–556. (doi:10.1038/nrg1633)

3. Yerushalmi S, Green RM. 2009 Evidence for the adaptive significance of
circadian rhythms. *Ecol Lett* **12**, 970–981.
(doi:10.1111/j.1461-0248.2009.01343.x)

4. Kondo T, Strayert CA, Kulkarnit RD, Taylor W, Ishiura M, Goldent SS, Hirschie
C, Ii J. 1993 Circadian rhythms in prokaryotes: Luciferase as a reporter of
circadian gene expression in cyanobacteria. *Proc Natl Acad Sci U S A* **90**,
5672–5676.

5. Takahashi JS. 1995 Molecular neurobiology and genetics of circadian rhythms in
mammals. *Annu Rev Neurosci* **18**, 531–553.
(doi:10.1146/annurev.ne.18.030195.002531)

6. King DP, Takahashi JS. 2000 Molecular genetics of circadian rhythms in
mammals. *Annu Rev Neurosci* **23**, 713–742.

7. Reppert SM, Weaver DR. 2001 Molecular analysis of mammalian circadian
rhythms. *Annual Review of Physiology* **63**, 647–676.
(doi:10.1146/ANNUREV.PHYSIOL.63.1.647)

8. Yerushalmi S, Yakir E, Green RM. 2011 Circadian clocks and adaptation in
Arabidopsis. *Mol Ecol* **20**, 1155–1165. (doi:10.1111/j.1365-294X.2010.04962.x)

9. Rubin MJ, Brock MT, Baker RL, Wilcox S, Anderson K, Davis SJ, Weinig C.
2018 Circadian rhythms are associated with shoot architecture in natural settings.
New Phytologist **219**, 246–258. (doi:10.1111/nph.15162)

10. Rubin MJ *et al.* 2017 Circadian rhythms vary over the growing season and
correlate with fitness components. *Mol Ecol* **26**, 5528–5540.
(doi:10.1111/mec.14287)

11. Koritala BSC *et al.* 2020 Habitat-specific clock variation and its consequence on
reproductive fitness. *J Biol Rhythms* **35**, 134–144.

12. Bloch G, Robinson GE. 2001 Reversal of honeybee behavioural rhythms. *Nature*
410, 1048–1048. (doi:10.1038/35074183)

13.   Shemesh Y, Cohen M, Bloch G. 2007 Natural plasticity in circadian rhythms is

mediated by reorganization in the molecular clockwork in honeybees. *The FASEB*

Journal **21**, 2304–2311. (doi:10.1096/FJ.06-8032COM)

14.   Tessmar-Raible K, Raible F, Arboleda E. 2011 Another place, another timer:

marine species and the rhythms of life. *BioEssays* **33**, 165–172.

(doi:10.1002/bies.201000096)

15.   Barnwell FH. 1966 Daily and tidal patterns of activity in individual fiddler crab

(Genus *Uca*) from the Woods Hole region. *Biol Bull* **130**, 1–17.

16.   Chabot CC, Ramberg-Pihl NC, Watson WH. 2016 Circalunidian clocks control

tidal rhythms of locomotion in the American horseshoe crab, *Limulus polyphemus*.

Mar Freshw Behav Physiol **49**, 75–91. (doi:10.1080/10236244.2015.1127679)

17.   Satoh A, Yoshioka E, Numata H. 2008 Circatidal activity rhythm in the mangrove

cricket *Apteronemobius asahinai*. *Biol Lett* **4**, 233–236.

(doi:10.1098/rsbl.2008.0036)

18.   Takekata H, Matsuura Y, Goto SG, Satoh A, Numata H. 2012 RNAi of the

circadian clock gene *period* disrupts the circadian rhythm but not the circatidal

rhythm in the mangrove cricket. *Biol Lett* **8**, 488–491.

(doi:10.1098/rsbl.2012.0079)

19.   Takekata H, Numata H, Shiga S, Goto SG. 2014 Silencing the circadian clock

gene *Clock* using RNAi reveals dissociation of the circatidal clock from the

circadian clock in the mangrove cricket. *J Insect Physiol* **68**, 16–22.

(doi:10.1016/j.jinsphys.2014.06.012)

20.   O'Neill JS, Lee KD, Zhang L, Feeney K, Webster SG, Blades MJ, Kyriacou CP,

Hastings MH, Wilcockson DC. 2015 Metabolic molecular markers of the tidal

clock in the marine crustacean *Eurydice pulchra*. *Current Biology* **25**,

R326–R327. (doi:10.1016/J.CUB.2015.02.052)

21.   Mat AM, Massabuau JC, Ciret P, Tran D. 2014 Looking for the clock mechanism

responsible for circatidal behavior in the oyster *Crassostrea gigas*. *Mar Biol* **161**,

89–99. (doi:10.1007/s00227-013-2317-2)

22.   Schnytzer Y, Simon-Blecher N, Li J, Ben-Asher HW, Salmon-Divon M, Achituv

Y, Hughes ME, Levy O. 2018 Tidal and diel orchestration of behaviour and gene

expression in an intertidal mollusc. *Sci Rep* **8**. (doi:10.1038/s41598-018-23167-y)

23.   Mat AM, Perrigault M, Massabuau JC, Tran D. 2016 Role and expression of *cry1*

in the adductor muscle of the oyster *Crassostrea gigas* during daily and tidal valve

activity rhythms. *Chronobiol Int* **33**, 949–963.

(doi:10.1080/07420528.2016.1181645)

24. Tran D, Perrigault M, Ciret P, Payton L. 2020 Bivalve mollusc circadian clock

genes can run at tidal frequency. *Proceedings of the Royal Society B: Biological*

Sciences **287**. (doi:10.1098/rspb.2019.2440)

25. Kwiatkowski ER, Schnytzer Y, Rosenthal JJC, Emery P. 2023 Behavioral

circatidal rhythms require *Bmal1* in *Parhyale hawaiensis*. *Current Biology* **33**,

1867–1882. (doi:10.1016/J.CUB.2023.03.015)

26. Yokomizo T, Takahashi Y. 2022 Endogenous rhythm variation and adaptation to

the tidal environment in the freshwater snail, *Semisulcospira reiniana*. *Front Ecol*

Evol **10**, 1218. (doi:10.3389/FEVO.2022.1078234)

27. Yamanaka O, Takeuchi R. 2018 UMATracker: An intuitive image-based tracking

platform. *Journal of Experimental Biology* **221**, 1–5. (doi:10.1242/jeb.182469)

28. Bolger AM, Lohse M, Usadel B. 2014 Trimmomatic: A flexible trimmer for

Illumina sequence data. *Bioinformatics* **30**, 2114–2120.

(doi:10.1093/bioinformatics/btu170)

29. Grabherr MG *et al.* 2011 Full-length transcriptome assembly from RNA-Seq data

without a reference genome. *Nat Biotechnol* **29**, 644–652. (doi:10.1038/nbt.1883)

30. Li B, Dewey CN. 2014 RSEM: accurate transcript quantification from RNA-seq

data with or without a reference genome. *BMC Bioinformatics* **12**.

(doi:10.1201/b16589)

31. Peñaloza C, Gutierrez AP, Eöry L, Wang S, Guo X, Archibald AL, Bean TP,

Houston RD. 2021 A chromosome-level genome assembly for the Pacific oyster

Crassostrea gigas. *Gigascience* **10**, 1–9.

(doi:10.1093/GIGASCIENCE/GIAB020)

32. Schmid B, Helfrich-Förster C, Yoshii T. 2011 A new ImageJ plug-in 'actogramJ'

for chronobiological analyses. *J Biol Rhythms* **26**, 464–467.

(doi:10.1177/0748730411414264)

33. Thaben PF, Westermark PO. 2014 Detecting rhythms in time series with rain. *J*

Biol Rhythms **29**, 391–400. (doi:10.1177/0748730414553029)

34. Kuintzle RC, Chow ES, Westby TN, Gvakharia BO, Giebultowicz JM, Hendrix

DA. 2017 Circadian deep sequencing reveals stress-response genes that adopt

robust rhythmic expression during aging. *Nat Commun* **8**.

(doi:10.1038/ncomms14529)

35.   Yu G, Wang LG, Han Y, He QY. 2012 ClusterProfiler: An R package for
comparing biological themes among gene clusters. *OMICS* **16**, 284–287.
(doi:org/10.1089/omi.2011.0118)

36.   Sharma VK. 2003 Adaptive significance of circadian clocks. *Chronobiol Int* **20**,
901–919. (doi:10.1081/CBI-120026099)

37.   Urabe M. 1998 Diel change of activity and movement on natural river beds in
Semisulcospira reiniana. *The Japanese journal of malacology* **57**, 17–27.
(doi:10.18941/venusjjm.57.1_17)

38.   Beaulé C. 2008 Aschoff's Rules. *Encyclopedia of Neuroscience* , 190–193.
(doi:10.1007/978-3-540-29678-2_383)

39.   Hughes ME, DiTacchio L, Hayes KR, Vollmers C, Pulivarthy S, Baggs JE, Panda
S, Hogenesch JB. 2009 Harmonics of circadian gene transcription in mammals.
PLoS Genet **5**. (doi:10.1371/journal.pgen.1000442)

40.   Zhu B. 2020 Decoding the function and regulation of the mammalian 12-h clock.
J Mol Cell Biol **12**, 752–758. (doi:10.1093/jmcb/mjaa021)

41.   Zhu B, Zhang Q, Pan Y, Mace EM, York B, Antoulas AC, Dacso CC, O'Malley
BW. 2017 A cell-autonomous mammalian 12 hr clock coordinates metabolic and
stress rhythms. *Cell Metab* **25**, 1305–1319. (doi:10.1016/j.cmet.2017.05.004)

42.   Zhang L, Hastings MH, Green EW, Tauber E, Sladek M, Webster SG, Kyriacou
CP, Wilcockson DC. 2013 Dissociation of circadian and circatidal timekeeping in
the marine crustacean *Eurydice pulchra*. *Current Biology* **23**, 1863–1873.
(doi:10.1016/j.cub.2013.08.038)

43.   Satoh A, Terai Y. 2019 Circatidal gene expression in the mangrove cricket
Apteronemobius asahinai. *Sci Rep* **9**. (doi:10.1038/s41598-019-40197-2)

44.   Mat AM *et al.* 2020 Biological rhythms in the deep-sea hydrothermal mussel
Bathymodiolus azoricus. *Nat Commun* **11**. (doi:10.1038/s41467-020-17284-4)

45.   Hastings MH, Naylor E. 1980 Ontogeny of an endogenous rhythm in *Eurydice
pulchra*. *J Exp Mar Biol Ecol* **46**, 137–145. (doi:10.1016/0022-0981(80)90027-1)

46.   Palmer JD. 1995 Review of the dual-clock control of tidal rhythms and the
hypothesis that the same clock governs both circatidal and circadian rhythms.
Chronobiol Int **12**, 299–310.

47.   Enright JT. 1976 Plasticity in an isopod's clockworks: shaking shapes form and
affects phase and frequency. *Journal of Comparative Physiology A* **107**, 13–37.
(doi:10.1007/BF00663916)

Data Accessibility

545   All raw transcriptome read data were deposited in the DDBJ Sequenced Read Archive

546   under accession numbers SAMD00632165–SAMD00632232. The activity and

547   transcriptome data are available in the Figshare (doi: 10.6084/m9.figshare.23672424

548   and doi: 10.6084/m9.figshare.23672691, respectively).

Author Contributions

551   T.Y. and Y.T. designed the experiments. T.Y. performed the experiments and conducted

552   the data analysis. T.Y. drafted the manuscript. T.Y. and Y.T. edited the manuscript.

554 **Table**

555
556 **Table 1.** Tukey's HSD test of differences in gene expression patterns based on PC1.

	Mean difference	P-value
Nontidal vs. Tidal	37.67	4.7×10^{-5}
Control vs. Treatment	−14.80	0.09
Nontidal Control vs. Nontidal Treatment	−10.74	0.82
Nontidal Control vs. Tidal Control	41.51	0.0067
Nontidal Control vs. Tidal Treatment	22.77	0.26
Nontidal Treatment vs. Tidal Control	−52.25	3.2×10^{-4}
Nontidal Treatment vs. Tidal Treatment	33.51	0.036
Tidal Control vs. Tidal Treatment	−18.74	0.41

Figure legends

Figure 1. Activity patterns of individuals in the nontidal and tidal populations of the control group under laboratory DD conditions. (a) Locomotor distance of the nontidal and tidal populations over 96 h. Error bars represent the standard error of the mean (SEM). Subjective day and night are indicated by grey and black bars above the patterns, respectively. (b) Lomb–Scargle periodogram analysis showing peaks in the circadian range.

Figure 2. Activity patterns of individuals in the nontidal and tidal populations of the treatment group under laboratory DD conditions. (a) Locomotor distance of the nontidal and tidal populations over 84 h. Error bars represent the SEM. Subjective day and night are indicated by grey and black bars above the patterns, respectively. The expected tide level in the laboratory is shown as a grey dotted line. (b) Lomb–Scargle periodogram analysis showing peaks in the circatidal range.

Figure 3. Gene expression patterns and rhythms of individuals in the nontidal and tidal populations under laboratory DD conditions. Control samples were kept under LD conditions without the tidal cycle. Treatment samples were exposed to the simulated tidal cycle for four weeks. (a) Probabilistic PCA of the expression of all genes after filtering in the nontidal and tidal populations. (b) Heatmaps of standardized expression patterns of rhythmic transcripts with circatidal periods in the nontidal and tidal individuals under DD conditions detected via RAIN. The dotted wavy lines above the heatmaps of treatment samples represent the simulated tide level. (c) Venn diagrams detailing the number of circatidal and circadian oscillating genes detected in the nontidal and tidal populations. (d) Bar plot showing the number of circatidal and circadian oscillating genes detected in the nontidal and tidal populations.

Figure 4. The dominant rhythmicity of the transcriptome in the control and treatment groups of the nontidal and tidal populations evaluated by log-scaled ratio of the P-value of circadian periodicity to that of circatidal periodicity in the transcriptome. The dashed line represents the same P-values for circadian and circatidal periodicity. Error bars represent the SEM.

Figure 5. KEGG pathway analysis of the tidal population. (a) KEGG pathway over-representation analysis of the tidal population. Gene ratio refers to the ratio of the number of differentially expressed genes (DEGs) to the total number of genes in a specific pathway. The size of the circles represents the count of DEGs. Circles are coloured based on the adjusted P-value. (b) KEGG-GSEA of the tidal population. Gene ratio refers to the ratio of the number of core enriched genes to the total number of genes in a specific pathway. The size of the circles represents the count of the core enriched genes. No pathways involved in decreased circatidal rhythmicity were detected.