Striatal dopamine release reflects a multi-factorial prediction error

Kauê Machado Costa[1*], Nishika Raheja[1], Jash Mirani[1], Courtney Sercander[1], and Geoffrey Schoenbaum[1*]

[1] National Institute on Drug Abuse Intramural Research Program, National Institutes of Health, Baltimore, MD, 21224, USA

* Corresponding authors: kaue.m.costa@gmail.com and geoffrey.schoenbaum@nih.gov

Abstract

Dopamine is classically thought to drive learning based on errors in the prediction of rewards and punishments. However, animals also learn to predict cues with no intrinsic value, and it is unclear if such latent learning also relies on dopaminergic prediction errors. Here, we tested this by recording dopamine release in the nucleus accumbens and dorsomedial striatum while rats executed a sensory preconditioning task that incorporates both types of learning. We found that dopamine release in both regions correlated with errors in predicting value-neutral cues during latent learning and with errors in predicting reward during reward-based conditioning. Moreover, dopamine in the nucleus accumbens reflected inferred value in the probe test, supported by orbitofrontal cortex activity. Our findings suggest that dopamine signals a multi-factorial prediction error, capable of supporting model-based learning.

Introduction

When navigating the world, animals learn both to predict rewards and punishments (value-based learning) and the relational structure of different elements of the environment, even if they are seemingly irrelevant (value-neutral or latent learning) (*1*). The first form of learning can be explained by a "model-free" framework, where learning is based on cached values, while the second form requires internal representations of the external world, or a "model-based" framework. The neuromodulator dopamine has been strongly implicated in value-based learning, as across several species and tasks dopamine activity reflects the difference between experienced and expected value, or reward prediction errors (RPEs) (*2–5*). This has led to the powerful and influential proposal that dopamine supports something akin to a model-free temporal difference reinforcement learning (TDRL) process in the brain (*2*).

However, recent work has questioned this interpretation. Dopamine signals have been found to correlate with, and influence learning about, variables that are not strictly value, including movement probability, interval timing, and sensory stimulus properties (*6–9*). Dopamine neuron activity also signals errors in predictions derived from value that must be inferred from task models (*10, 11*). Some of this functional diversity has been related to differences in projection-specific subpopulations (*12, 13*). For example, dopamine release in dorsal striatal regions, like the dorsomedial striatum (DMS), has been shown to correlate more with value-orthogonal variables than dopamine release in the nucleus accumbens core (NAcc) (*14–16*). However, this functional diversity is not sufficient to explain all of the data, since even within the NAcc, dopamine release can correlate with variables that are orthogonal to value, like novelty and sensory salience (*17, 18*), and individual RPE-encoding dopamine neurons in the midbrain also signal errors in the prediction of task-irrelevant sensory properties of rewards, or sensory prediction-errors (SPEs) (*7, 19*). Finally, optogenetic manipulations that causally affect RPE-based learning also impact learning about the sensory features of rewards (*20–22*).

To conciliate these findings with classical work, several new formal hypotheses on dopamine function have been proposed. These posit that dopamine neuron activity keeps tracks RPEs based on separate predictive "threads", "bases", or "channels",

resulting in RPE-signals that are dissociable according to their defining sensory or task-
based properties (*23–25*). These models are essentially modifications of the current
model-free framework to allow more complexity, diversity, or specificity in the RPE
signal, however they remain tied to learning about motivationally significant elements.
Therefore, while these updated models can explain new findings on dopaminergic
errors in learning about specific reward properties and the cues and actions associated
with them, they do not predict any role for such error signaling in learning to predict
events in the environment that are not related to reward or punishment.

Yet, such latent learning is common, and there is evidence that it can be influenced by
brief activation and inhibition of midbrain dopamine neurons (*22*). While these results
can be potentially explained by a permissive role for phasic or even tonic dopamine in
value-neutral learning (*26, 27*), it is also possible that they are evidence that dopamine
operates as an error signaling mechanism that is *general* or *multifactorial*, which
represents predictive errors across multiple dimensions, one of which is value (*7, 17,*
19, 28–31).

Here, we used optophysiological dopamine recordings and a classical learning theory
task (*32*) to assess whether dopamine provides an error-like signal during value-neutral
or latent learning. We recorded striatal dopamine signals while rats executed a 3-phase
sensory preconditioning task (SPC), which incorporates value-neutral, explicit value-
based, and inferred value-based prediction errors in its structure (*10, 33*). We also took
advantage of the known role of the lateral orbitofrontal cortex (lOFC) in supporting
model-based inference in this task and inactivated this region during the critical probe
session while recording dopamine signals (*33*). We found that dopamine release in both
NAcc and DMS correlated with sensory prediction errors during the formation of the
valueless cue-cue associations and with classical RPEs during conditioning. In the
probe session, lOFC inactivation had a selective effect on NAcc dopamine release
consistent with disruption of an inference process. These findings suggest that striatal
dopamine signals reflect errors across different types of predictions, including when
value is not at issue**.**

Results

We transfected male rats with dLight1.2 and implanted them with optic fiber cannulas in the NAcc and DMS to allow simultaneous multi-site optophysiological recordings of dopamine release dynamics (Figure 1A and S1A) (*34–37*). To allow for better comparisons across rats and sessions, z-scored dopamine-dependent fluorescence was normalized to the maximum signal observed in each session (see Methods). We also transfected the same rats with either hM4d (inhibitory DREADD receptor; n=8; OFCi group) or mCherry (control; n=6; CTRL group) in the lOFC to allow us to chemogenetically inhibit activity in this region (Figure 1B) (*38, 39*). After at least 4 weeks for recovery and viral expression, rats were food restricted, shaped to retrieve food-pellets from the food port, and then subjected to SPC training (Figure 1C). Dopamine release was recorded during all sessions, and JHU37160 dihydrochloride (JH60; i.p. 0.2 mg/kg), a high-potency DREADD agonist (*40*), was injected prior to the probe test to inactivate the principal neurons in the lOFC of the OFCi rats, as validated previously (*38, 39*). Against this behavioral backdrop, we analyzed dopamine release dynamics in each recording site to address:

1) whether dopamine release in the probed locations reflected error-driven associative learning in the absence of overt value or reward,

2) if release in these same sites exhibited classic RPE signals during conditioning, and

3) how dopamine release was structured during the probe test, when the preconditioned cues were inferred to have value, a property known to depend on processing in lOFC.

Behavioral results and the role of lOFC in inference-based behavior

First, we consider the behavioral results. During the preconditioning phase, food port responding during the presentation of each cue was low and rats did not distinguish between them (Figure 1D; 2-way ANOVA for cue and group effects: $P > 0.113$ and $F < 2.131$ for all comparisons). During conditioning, responding progressively and selectively increased to B, the cue paired with food delivery (Figure 1E; 3-way ANOVA

for effects of session progression, cues, and group; session effect: $P=0.014^*$,

$F_{(5,60)}=3.152$; cue effect: $P=0.002^{**}$, $F_{(1,12)}=14.88$; interaction of cue and session effects:

session effect: $P=0.008^{**}$, $F_{(5,60)}=3.497$; all other effects: $P>0.742$, $F<0.327$).

Subsequently, in the probe test, all rats responded more to B, the food-paired cue, than

to the non-reinforced cue, D, indicating that they had correctly learned to attribute higher

value to the food-paired cue (Figure 1F; Wilcoxon test, $P=0.031^*$ for CTRL and

$P=0.008^{**}$ for OFCi). Furthermore, rats in the CTRL group responded more during A

than C ($P=0.031^*$), indicating that they had learnt the A-B association in the pre-

conditioning phase and were able to use it in the probe test to infer that A (and not C)

might also lead to reward (as it predicted B). Rats in the OFCi group, on the other hand,

had similar responding to cues A and C ($P=0.313$), despite showing normal B versus D

discrimination, indicating a selective disruption of inference-guided behavior, replicating

previous results obtained with other methods of lOFC inactivation (*33, 41*).

Figure 1. Histological verification and behavioral results. A: Location of fiber tips in the striatum for all recorded rats and striatal locations (top panels) and representative examples (bottom panels). Insert in the middle bottom panel contains a representative demonstration of the expression of dLight1.2 in relation to fiber tip location. **B:** Spread of DREADD construct expression (mCherry reporter) in lOFC. Left panels contains representative histological microphotographs of a CTRL and an OFCi rat. Right panels contain schematic reconstructions of the viral spread across the lOFC in both groups.

Lighter shading represents area of maximal spread, while darker shading reflects the

area of minimal spread. **C:** Schematic of the SPC behavioral task. **D:** Behavioral

responding, as measured by total time spent in the food port during cue presentation,

for all rats in the preconditioning phase. Due an absence of group effects (CTRL vs

OFCi) in the preconditioning and conditioning phases, these data were pooled together

in this figure for clarity (see Figure S2 for results split according to groups). **E:**

Behavioral responding in the conditioning phase. Note the progressively increased

responding to the reinforced cue B in relation to the nonreinforced cue D. **F:** Behavioral

responding during the probe test, where all rats were injected with the DREADD agonist

JH60. **G:** Difference index (difference divided by the sum) between responding to cues

B and D and cues A and C, for the CTRL and OFCi groups. A positive difference index

indicates a higher responding to B or A, while a negative one indicates higher

responding to C or D, in each of the comparisons. Difference index values were tested

with one-sample Wilcoxon tests against a hypothetical 0 (equal responding to both

cues). This is essentially another way of representing the data from the graphs in F and

it confirmed a significant higher responding for B versus D in both groups (P=0.031* for

CTRL and P=0.008** for OFCi) and a similar higher responding for A only in the CTRL

group (P=0.031*) and not in the OFCi group (P=0.148). Note from F and G that rats in

the CTRL group, where OFC activity was not affected, respond more to cue A, which in

preconditioning predicted the subsequently reinforced cue B, than to cue C, which was

paired with the subsequently unreinforced cue D. In the OFCi group, where OFC activity

was chemogenetically inhibited, there was no difference in responding to cues A and C.

Also notably, rats in both groups responded significantly more to B than to D,

irrespective of OFC inactivation. Data are represented as mean ± SEM. *P<0.05;

**P<0.01.

Dopamine release in the NAcc and DMS exhibit error-like correlates during pre-conditioning

We next asked how dopamine release responded during the pre-conditioning phase, when the rats were exposed to arbitrary pairings of neutral cues, in the absence of any value, overt reward, or even response requirement. Focusing on release around cue presentation, and starting with responses in the NAcc, we observed clear dopamine peaks at the onset of each cue (A,B,C, and D). Initially, these peaks were similar (Figure 2A and 2B) and each diminished with repeated exposure (2-way ANOVA for effect of trial progression and cue predictability; trial effect: $P<0.0001^{***}$, $F_{(2,54)} = 22.76$), a pattern consistent with signaling of surprise or novelty (*42–44*) and which has been associated with learning that underlies latent inhibition (*18, 45*).

However, concurrently with this expected effect, we also found that peak responses to the predicted events, B and D, diminished faster and to a greater extent than those to the unpredicted stimuli, A and C (Figure 2B and C; cue predictability effect: $P=0.0004^{**}$, $F_{(1,27)} = 16.48$). This additional effect indicates that the predictive structure influences the dopamine responses to each cue, essentially providing an error-like correlate during value-less sensory-sensory learning – a SPE riding on top of novelty-related changes (Figure 2).

We observed a qualitatively similar response pattern in the DMS, where dopamine peak responses were larger at the first presentation of each cue, reduced over repeated exposures, and were lower for predicted events than for unpredicted events (Figure 2D and 2E; trial effect: $P=0.023^*$, $F_{(2,54)} = 4.039$; cue predictability effect: $P=0.004^{**}$, $F_{(1,27)} = 9.577$). However, the changes in peak responses over time were less pronounced and seemed to occur at a slower progression rate than in the NAcc, indicating that perhaps the changes observed in DMS responses were following those in the NAcc, consistent with previous evidence suggesting an ascending ventromedial to dorsolateral hierarchy of learning-induced plasticity in midbrain-striatal circuits (*46, 47*).

Figure 2. Dopamine signals in NAcc and DMS reflect general prediction errors during sensory preconditioning.

A: Heatmap representation of average NAcc dopamine responses during all 12 preconditioning trials, recorded with dLight1.2 (N=14). Note the larger dopamine responses during the first trials to both the unpredicted and predicted cues, and the progressive diminishment of the responses with subsequent trials, which is more pronounced for the responses to predicted stimuli. **B:**.Average traces of NAcc dopamine responses to predicted and unpredicted cues during the start (first two trials), middle (trials 6 and 7), and end (trials 11 and 12) of the preconditioning phase. **C:** Peak NAcc dopamine responses measured over the course of the preconditioning training. Note the similar responses to both types of cues in the beginning of training, and the significant difference between responses to predicted and unpredicted cues. **D-E:** similar to A-C, but for DMS dopamine signals, simultaneously recorded with NAcc signals. We highlight that in the DMS, the progressive reduction of dopamine responses to both types of cues was not as strong as in the NAcc. Data are represented as mean ± SEM. *$P<0.05$; **$P<0.01$.

To confirm the importance of the predictive structure in determining the changes in

dopamine release that we observed, we trained another group of rats (n=5), this time

transfected with GRAB-DA2m and implanted with an optic fiber cannula only in NAcc

and without viral injection of DREADDs into lOFC. These rats underwent the same pre-

conditioning training as above (Figure 3A-C), except that we added a session at the end

in which we switched the order in which the cues were presented (Figure 3D). In other

words, in this added session, the rats initially received presentations of A→B and C→D,

as they had the prior two sessions, followed by presentations of B→A and D→C. If the

changes in dopamine release were due to the predictive structure of the cues, then the

release should be immediately restored to D and B when they appear unpredicted by

another cue, and it should slowly diminish to A and C, as the new predictive structure is

learnt. Consistent with this idea, we first found that VMS dopamine signals in this new

group showed similar SPE signature as in the previous dLight1.2-transfected rats during

the first 2 sessions - responses to predicted cues B and D were lower than responses to

the unpredicted cues A and C - confirming that this finding is robust across experimental

groups and dopamine sensors (Figure 3A-C; trial effect: P=0.0002***, $F_{(2,18)}$ = 14.86;

cue predictability effect: P=0.008**, $F_{(1,9)}$ = 11.43). Then, when the order of cue

presentation was reversed in the second half of the added preconditioning session, the

responses to B and D, now unpredicted, were restored, and responses to cues A and C,

now predicted, were slightly diminished (Figure 3D-F; 2-way ANOVA for effect of cue

predictability – unpredicted versus predicted - and cue identity – B/D versus A/C;

predictability effect: P=0.0228*, $F_{(1,9)}$ = 7.512; cue effect: P=0.3102, $F_{(1,9)}$ = 1.156;). This

pattern strongly suggests that VMS dopamine release reflects the predictive structure of

cue presentation in exactly the same way that the classic RPE response reflects the

order of presentation of a cue and an event with overt value – an SPE for lack of a

better term.

Figure 3. Encoding of sensory-sensory prediction errors by dopamine signals in NAcc depends on the predictive relationship between cues.

A: Heatmap representation of the average NAcc dopamine signals during all trials of preconditioning training in a separate cohort of rats (N=5), recorded with GRAB-DA2m. Note that the response pattern is similar to those recorded with dLight1.2. **B:** Average traces of dopamine responses during the start, middle, and end of preconditioning, defined the same way as in Figure 2. **C:** Peak dopamine responses during preconditioning training in this cohort of rats. Note the similar pattern to the cohort in Figure 2. **D:** Heatmap representation of dopamine signals in the preconditioning switch session. In the first three trials the structure of the task was the same as in the previous training sessions, with A preceding B and C preceding D. In the last three trials, that

order was switched, with B not coming before A and D before C, with no external

signaling to the rats. Note that responses to B and D, which were almost entirely

suppressed in trial 3, were immediately restored in the trial 4, when these cues

presented unexpectedly, and responses to cues A and C, which were unexpected, start

to become slightly more suppressed once they become predicted by cues B and D. **E:**

Average dopamine responses to each type of cue before and after the switch. **F:** Peak

dopamine responses to cues A/C and B/D when they were both predicted (first three

trials for B/D and last three trials for A/C) and unpredicted (first three trials for A/C and

last three trials for B/D). For both sets of cues, regardless of their identity or exposure

history, NAcc dopamine responses were larger when they were unpredicted than when

they were predicted. Data are represented as mean ± SEM.

Dopamine signals in NAcc reflect classical reward prediction errors during conditioning

Consistent with numerous prior studies, we found that peaks in dopamine release in the

NAcc strongly signaled RPEs during the conditioning phase of the task (Figure 2)(*14,*

16). Peaks in release were initially high in response to pellet delivery and then

decreased over time, while peaks in response to onset of the pellet-paired cue, B,

increased (Figure 4A). This pattern could be resolved on a trial-by-trial (Figure 4A) or

session-by-session basis (Figure 4B), with the difference between peak responses to

the pellet delivery (US) and the cue (CS) becoming progressively lower with training

(Figure 4C; 2-way ANOVA for effects of session progression and cue – B versus D;

session effect: $P<0.0001$***, $F_{(5,65)} = 14.48$; cue effect: $P<0.0001$***, $F_{(1,13)} = 153.2$;

interaction effect: $P<0.0001$***, $F_{(5,65)} = 9.809$)(*10*). Importantly, responses to the non-

reinforced cue, D, did not change significantly over time. These results confirm that

dopamine signals in NAcc - recorded in the current rats, fiber locations, and using this

task and methods - exhibited classic RPE correlates during conditioning.

Furthermore, we found that dopamine peaks in the DMS also seemed to reflect RPEs,

albeit again with a lower update rate as compared to the NAcc, and with less of a

difference in relation to the peaks to the CS- (Figure 4D-F; session effect: $P=0.492$

$F_{(5,65)} = 0.892$; cue effect: $P<0.0001$***, $F_{(1,13)} = 36.26$; interaction effect: $P=0.0221$*,

$F_{(5,65)} = 2.842$). This parallels our findings in the preconditioning phase, where changes

to signals in DMS seem to lag changes to signals in NAcc.

Figure 4. Dopamine release in VMS and DMS reflects classic, value-based, "reward" prediction errors in the conditioning.

A: Heatmap representation of average NAcc dopamine signals in all trials of the conditioning phase of the cohort recorded with dLight1.2 (N=14). Note the progressive reduction of responses to US presentation and concurrent increase is responses to cue B, the reinforced CS+, with less changes in the responses to cue D, the unreinforced CS-. **B:** Average traces of dopamine signals for each session (6 trials per condition per session) of conditioning. **C:** Difference between the peak NAcc dopamine responses to

277   US and CS presentation, for both cues B and D. A progressive decrease in this

278   difference is key indicator of an RPE response (*10*). **D-F:** Same as A-C, but for DMS

279   dopamine signals. Note that DMS dopamine also shows signs of RPE encoding, but

280   much less prominently as NAcc. Data are represented as mean ± SEM. *$P<0.05$;

281   **$P<0.01$.

Dopamine release in the NAcc and DLS responses to cues with inferred value are
dependent on lOFC activity

Finally, we arrive at the probe phase of the main SPC experiment. Up until now we have collapsed the two groups because there was no effect of group on any of the previous comparisons (Figures S2 and S3; 3-way ANOVA for effects of session/trial progression, cues, and group; all group main or interaction effects: $P>0.1$, $F<0.5$). In the NAcc, as expected, responses to the reinforced cue B were significantly higher than to the unreinforced cue D in both the CTRL and OFCi groups (Figure 5A-B; paired t-tests; CTRL: P= 0.002**; OFCi: P=0.007**). In the DMS, there was no significant difference between responses to B and D cues in either group (Figure 5A-B; paired t-tests; CTRL: P= 0.087; OFCi: P=0.466), which is likely reflective of the diminished RPE encoding observed in DMS and the lower number of trials in the probe session.

Surprisingly, in the CTRL group dopamine responses to cues A and C were similar during the probe phase (Figure 5). This was true in both areas (paired t-test, A vs C in CTRL group; NAcc: P=0.326; DMS: P=0.801). That measured dopamine release did not differ for the two cues was unexpected, particularly in light of the differential behavior of these rats to the two cues in the probe test (Figure 1) as well as prior unit recording data showing that dopamine neuron spiking is higher to cue A than to cue C (*10*). These results are consistent with prior demonstrations that striatal dopamine release and dopamine neuron firing are not always correlated, and that certain behavioral variables can be observed in one measure but not the other (*48*).

Nevertheless, even if the release appears similar, the hypothesized circuit function predicts that release should reflect different information for the two different cues; specifically, while release to cue C might reflect non-specific generalization and perhaps elevated salience due to the long period since it was last seen, release to cue A should reflect, at least in part, the specific predictive relationship between A and B and between B and reward. We know from prior work in this setting that this is the case for the behavior to A, as it is sensitive to devaluation of the predicted reward (*49*). Critically, behavior based on this information is selectively sensitive to inactivation of lOFC (*33*). This predicts that inactivation of lOFC should selectively affect dopamine release to A .

312   Consistent with this prediction, we found that OFC inactivation did reduce the relative

313   peak responses to cue A in NAcc (Figure 5A-B), with responses to all other cues being

314   relatively similar (paired t-tests; A vs C in OFCi group; NAcc: P=0.044*) (unpaired t-

315   tests; A response in CTRL vs OFCi; NAcc: P=0.043*; all other comparisons of CTRL vs

316   OFCi showed P>0.05 for both NAcc and DMS).

Figure 5. Dopamine signals in NAcc and DLS reveal a contribution of lOFC on learning-dependent dopamine signals during probe. A: Average traces of NAcc

dopamine responses to all cues in the probe test, for both CTRL (N=6) and OFCi (N=8)

rats. All recordings and behavior in this session were performed after the i.p. injection of

0.2 mg/kg of JH60, which inhibited OFC activity in the OFCi group. Note that the only

difference between responses in the two groups is a blunted dopamine peak to A in the

OFCi group. **B:** Peak NAcc dopamine responses to all cues in the probe session,

confirming the significant difference only in responses to A between CTRL and OFCi

rats. **C-D:** same as in A-B, but for DMS dopamine responses. In the DMS, there was no

significant difference between responses to any of the cues between the CTRL and

OFCi rats. Data are represented as mean ± SEM. *$P<0.05$.

Discussion

Here we recorded dopamine in a three-phase SPC task and found that release in both

NAcc and DMS reflected errors in predicting neutral cues during the initial

preconditioning phase of training. These signals were dissociable from sensory salience

and detected at sites that also showed classic RPE signals during subsequent

conditioning. Thus, the results replicate previous findings linking striatal dopamine

release to salience and signaling of value-based prediction errors (*2*, *14*, *17*, *18*, *43*),

while also showing a novel, value-independent error-like response during learning about

neutral cue pairs. The significance of the latter finding is bolstered by previous work

using optogenetics showing that dopamine activity is both necessary and sufficient for

driving sensory-sensory learning in a similar preconditioning task (*22*). The causal

manipulations in this previous work could not differentiate between an error-driven

signal and a more general permissive role of dopamine for value-neutral learning, thus

the current data are critical in ruling out these possibilities in favor of a prediction error-

like mechanism.

Importantly, showing error signals in this setting goes beyond prior work by us and

others showing error signals during learning of specific features of otherwise valuable

events (i.e. rewards) (*7*, *11*, *19*, *20*, *50*), which can be explained with adjustments to

TDRL algorithms, such as the addition of dissociable "threads", "bases", or "channels"

for keeping track of different components of rewarding events (*23–25*). These models

cannot easily explain why neutral cues evoke error signals. Instead, the response

patterns observed here to neutral cues are much better explained by a model-based

process, or at least a hybrid model incorporating something like successor

representations (*28*, *29*, *31*, *39*).

These results, coupled with much prior work (*2*, *7*, *11*, *16*, *19*, *20*, *50*), suggest that

dopamine activity is shaped by multiple factors or dimensions that characterize events,

of which value is but one component (*28*, *29*, *31*, *39*). This broader interpretation holds

better explanatory power than several alternatives. For example, one potential

challenge to our conclusion that dopamine release reflects more than value is the

expansion of the concept of "value" itself. For example, novel sensory cues may be

assigned intrinsic value, given that they could potentially convey information about the

external world (*51*, *52*), or perhaps previous experiences, in the home cage or during

shaping, led the rats to acquire the general belief that sounds are broadly predictive of

rewards prior to preconditioning (*53*). These explanations seem unlikely to us, since

preconditioned cues trained in this manner do not support conditioned reinforcement

(*54*). Further, if these cues were intrinsically valuable, like a primary reward, then the

dopamine responses to the unpredicted cues A and C should have increased over time

during preconditioning, just like responses to B during conditioning, but they did not.

Instead, the observed pattern, in which dopamine responses to predicted cues were

inhibited faster, seems more compatible with the formation of a predictive association

between two cues without intrinsic value.

A related set of ideas is that dopamine responses track stimulus salience (*17*, *30*, *43*).

This component of the signal differs from a value prediction error because it factors in

attributes of the cues themselves and the relational structure of task elements, such as

novelty, stimulus intensity, and unexpectedness. Its presence should increase

associability or learning about a cue, both at that time and subsequently. Our findings

here and in prior causal work are generally compatible with this interpretation, as

predicted cues would indeed, by definition, be less surprising than unpredicted cues.

However, contrary to predictions for salience, there was no impact of activating

dopamine neurons on *subsequent* learning in our prior work (*22*).

Indeed, novelty, sensory salience, and signaled sensory and value prediction errors,

could all be conceptualized as prediction errors that operate on different variables and

across different hierarchical structures of time scale and information content (*7*, *31*).

Similar frameworks have been proposed to shape perception and general cognition,

and the idea that the brain itself is fundamentally dedicated to processing, and

minimizing, prediction errors is currently an influential and powerful hypothesis (*55*).

The involvement of dopamine in model-based learning is also supported by the effects

of lOFC inactivation in the probe test, where it selectively disrupted behavioral

responding and dopamine release in NAcc to cue A, the cue endowed with inferred

value by virtue of the prior training. This result is consistent with the impact of muscimol

inactivation of lOFC on responding in the probe test in our prior work (*33*), and with the

widespread observation that the lOFC is strongly associated with inferential or model-

based behavior across a variety of tasks, modalities, and species (*56*).

Dopamine recordings in the probe test did show some unexpected features. In the

control group, dopamine release was similar both to cue A, the preconditioned cue

predictive of reward, and cue C, the control cue. If the only mechanism at work were

one based on inference and if the only information represented by the error signal were

value, then the signal should preferentially have shown up to cue A. In this regard, the

results contrast with the results of a previous single unit recording study (*10*), in which

the firing of putative dopamine neurons in the ventral midbrain differed between cues A

and C in the probe test of a similar task.

However there may be a number of non-mutually exclusive explanations for this finding.

For starters, the optophysiological methods used here may be not sensitive enough to

pick up what may be a very subtle difference in the dopamine responses to cues A and

C in the control group, or to put it another way, other components of the dopamine

response (i.e., local features of the spatiotemporal patterns of release, differential

release from individual axon terminals, etc.) may better reflect inferred prediction errors

than bulk responses to cue onset. It is also possible that in the unit recording study (*10*)

the sampled neuronal population may have contained a disproportionate number of

neurons that projected to other striatal regions not sampled in our current work, where

different variables may be represented. However, the methods and analyses applied

here, which fit the standard of the field, were successful in detecting prediction errors in

preconditioning and conditioning, including classically defined RPEs, as well as some

differences imposed by lOFC inactivation. Therefore, this result seems unlikely to be

due to these artifactual causes.

More likely, release may follow a different time course than spiking activity, with different

sensitivities to aspects of our design. Such speculation is consistent with recent work

showing that dopamine neuron firing and release in target regions can be dissociated,

likely due to the sculpting effects of local striatal mechanisms (*48*, *57–59*). Similar

release in controls to the two preconditioned cues might reflect a greater sensitivity of

these release dynamics to commonalities across the cues that are unique to our task.

For instance, in preconditioning, all the cues are initially presented without any reward,

and this similarity may be the foremost determinant of bulk dopamine release signals in

the probe, even if other processes are in play. This might be quite different for other

tasks designed to study inferential behavior, like reinforcer devaluation, where no

common unrewarded latent state can be inferred during the initial learning phase (*11*).

Within this framework, one might predict that it would require a selective manipulation of

areas that process inferential information – such as the lOFC - to reveal a potential

difference in dopamine responses that depend on such processes, which is indeed what

we found.

In conclusion, our findings support the proposal that dopamine signaling in the striatum

mimics a *multi-factorial* prediction error term that comprises different variables and

whose information content varies across different regions. These prediction errors in the

NAcc and DMS reflect the relational structure of value-neutral sensory-sensory

associations, as well as cached value-based associations and novelty. We also present

evidence that accumbal dopamine responses to cues with inferred value are

differentially sensitive to lOFC disruption, suggesting that dopamine signals that appear

similar may be dependent on different underlying mechanisms according to their task

relevance.

References

1. E. C. Tolman, Cognitive maps in rats and men. *Psychol. Rev.* **55**, 189–208 (1948).

2. W. Schultz, P. Dayan, P. R. Montague, A Neural Substrate of Prediction and Reward. *Science (80-.).* **275**, 1593–1599 (1997).

3. K. M. Costa, G. Schoenbaum, Dopamine. *Curr. Biol.* **32**, R817–R824 (2022).

4. E. E. Steinberg, R. Keiflin, J. R. Boivin, I. B. Witten, K. Deisseroth, P. H. Janak, A causal link between prediction errors, dopamine neurons and learning. *Nat Neurosci.* **16**, 966–973 (2013).

5. C. Y. Chang, G. R. Esber, Y. Marrero-Garcia, H.-J. Yau, A. Bonci, G. Schoenbaum, Brief optogenetic inhibition of dopamine neurons mimics endogenous negative reward prediction errors. *Nat. Neurosci.* **19**, 1–8 (2015).

6. A. E. Hamilos, G. Spedicato, Y. Hong, F. Sun, Y. Li, J. A. Assad, Slowly evolving dopaminergic activity modulates the moment-to-moment probability of reward-related self-timed movements. *Elife.* **10** (2021), doi:10.7554/ELIFE.62583.

7. Y. K. Takahashi, H. M. Batchelor, B. Liu, A. Khanna, M. Morales, G. Schoenbaum, Dopamine Neurons Respond to Errors in the Prediction of Sensory Features of Expected Rewards. *Neuron.* **95**, 1395-1405.e3 (2017).

8. S. Soares, B. V. Atallah, J. J. Paton, Midbrain dopamine neurons control judgment of time. *Science (80-.).* **354**, 1273–1277 (2016).

9. L. S. Gonzalez, A. A. Fisher, S. P. D'Souza, E. M. Cotella, R. A. Lang, J. E. Robinson, Ventral striatum dopamine release encodes unique properties of visual stimuli in mice. *Elife.* **12** (2023), doi:10.7554/ELIFE.85064.

10. B. F. Sadacca, J. L. Jones, G. Schoenbaum, Midbrain dopamine neurons compute inferred and cached value prediction errors in a common framework. *Elife.* **5**, e13665 (2016).

11. G. K. Papageorgiou, M. Baudonnat, F. Cucca, M. E. Walton, Mesolimbic Dopamine Encodes Prediction Errors in a State-Dependent Manner. *Cell Rep.* **15**, 221–228 (2016).

12. J. Roeper, Dissecting the diversity of midbrain dopamine neurons. *Trends Neurosci.* **36**, 336–342 (2013).

467   13.   N. Farassat, K. M. Costa, S. Stojanovic, S. Albert, L. Kovacheva, J. Shin, R. Egger, M.
468   Somayaji, S. Duvarci, G. Schneider, J. Roeper, In vivo functional diversity of midbrain
469   dopamine neurons within identified axonal projections. *Elife*. **8** (2019),
470   doi:10.7554/eLife.48408.

471   14.   W. Menegas, B. M. Babayan, N. Uchida, M. Watabe-Uchida, Opposite initialization to
472   novel cues in dopamine signaling in ventral and posterior striatum in mice. *Elife*. **6**, 751–
473   767 (2017).

474   15.   N. F. Parker, C. M. Cameron, J. P. Taliaferro, J. Lee, J. Y. Choi, T. J. Davidson, N. D.
475   Daw, I. B. Witten, Reward and choice encoding in terminals of midbrain dopamine
476   neurons depends on striatal target. *Nat. Neurosci.* **19**, 845–854 (2016).

477   16.   J. J. Day, M. F. Roitman, R. M. Wightman, R. M. Carelli, Associative learning mediates
478   dynamic shifts in dopamine signaling in the nucleus accumbens. *Nat. Neurosci.* **10**,
479   1020–1028 (2007).

480   17.   M. G. Kutlu, J. E. Zachry, P. R. Melugin, S. A. Cajigas, M. F. Chevee, S. J. Kelly, B.
481   Kutlu, L. Tian, C. A. Siciliano, E. S. Calipari, Dopamine release in the nucleus accumbens
482   core signals perceived saliency. *Curr. Biol.* **31**, 4748-4761.e8 (2021).

483   18.   M. G. Kutlu, J. E. Zachry, P. R. Melugin, J. Tat, S. Cajigas, A. U. Isiktas, D. D. Patel, C.
484   A. Siciliano, G. Schoenbaum, M. J. Sharpe, E. S. Calipari, Dopamine signaling in the
485   nucleus accumbens core mediates latent inhibition. *Nat. Neurosci.* **25**, 1071–1081 (2022).

486   19.   T. A. Stalnaker, J. D. Howard, Y. K. Takahashi, S. J. Gershman, T. Kahnt, G.
487   Schoenbaum, Dopamine neuron ensembles signal the content of sensory prediction
488   errors. *Elife*. **8** (2019), doi:10.7554/eLife.49315.

489   20.   R. Keiflin, H. J. Pribut, N. B. Shah, P. H. Janak, Ventral Tegmental Dopamine Neurons
490   Participate in Reward Identity Predictions. *Curr. Biol.* **29**, 93-103.e3 (2019).

491   21.   C. Y. Chang, M. Gardner, M. G. Di Tillio, G. Schoenbaum, Optogenetic Blockade of
492   Dopamine Transients Prevents Learning Induced by Changes in Reward Features. *Curr.*
493   *Biol.* **27**, 3480-3486.e3 (2017).

494   22.   M. J. Sharpe, C. Y. Chang, M. A. Liu, H. M. Batchelor, L. E. Mueller, J. L. Jones, Y. Niv,
495   G. Schoenbaum, Dopamine transients are sufficient and necessary for acquisition of
496   model-based associations. *Nat. Neurosci.* **20**, 735–742 (2017).

497   23.   Y. K. Takahashi, T. A. Stalnaker, L. E. Mueller, S. K. Harootonian, A. J. Langdon, G.
498   Schoenbaum, Dopaminergic prediction errors in the ventral tegmental area reflect a
499   multithreaded predictive model. *Nat. Neurosci. 2023 265*. **26**, 830–839 (2023).

500   24.   B. Millidge, Y. Song, A. Lak, M. E. Walton, R. Bogacz, *bioRxiv*,
501   doi:10.1101/2023.05.09.540067.

502   25.   R. S. Lee, B. Engelhard, I. B. Witten, N. D. Daw, *bioRxiv*,
503   doi:10.1101/2022.02.28.482379.

504   26.   Y. Wang, O. Toyoshima, J. Kunimatsu, H. Yamada, M. Matsumoto, Tonic firing mode of
505   midbrain dopamine neurons continuously tracks reward values changing moment-by-
506   moment. *Elife*. **10** (2021), doi:10.7554/ELIFE.63166.

507   27.   W. Schultz, Recent advances in understanding the role of phasic dopamine activity.
508   *F1000Research 2019 81680*. **8**, 1680 (2019).

509   28.   M. P. H. Gardner, G. Schoenbaum, S. J. Gershman, Rethinking dopamine as generalized
510   prediction error. *Proc. R. Soc. B Biol. Sci.* **285**, 20181645 (2018).

511   29.   A. J. Langdon, M. J. Sharpe, G. Schoenbaum, Y. Niv, Model-based predictions for
512   dopamine. *Curr. Opin. Neurobiol.* **49**, 1–7 (2018).

513   30.   K. C. Berridge, The debate over dopamine's role in reward: the case for incentive
514   salience. *Psychopharmacology (Berl)*. **191**, 391–431 (2007).

515   31.   B. Lau, T. Monteiro, J. J. Paton, The many worlds hypothesis of dopamine prediction
516   error: implications of a parallel circuit architecture in the basal ganglia. *Curr. Opin.*
517   *Neurobiol.* **46**, 241–247 (2017).

518   32.   W. J. Brogden, Sensory pre-conditioning. *J. Exp. Psychol.* **25**, 323–332 (1939).

519   33.   J. L. Jones, G. R. Esber, M. A. McDannald, A. J. Gruber, A. Hernandez, A. Mirenzi, G.
520   Schoenbaum, Orbitofrontal cortex supports behavior and learning using inferred but not
521   cached values. *Science (80-.)*. **338**, 953–956 (2012).

522   34.   E. Martianova, S. Aronson, C. D. Proulx, Multi-Fiber Photometry to Record Neural Activity
523   in Freely-Moving Animals. *JoVE (Journal Vis. Exp.* **2019**, e60278 (2019).

524   35.   T. Patriarchi, J. R. Cho, K. Merten, M. W. Howe, A. Marley, W. H. Xiong, R. W. Folk, G. J.
525   Broussard, R. Liang, M. J. Jang, H. Zhong, D. Dombeck, M. von Zastrow, A. Nimmerjahn,

V. Gradinaru, J. T. Williams, L. Tian, Ultrafast neuronal imaging of dopamine dynamics with designed genetically encoded sensors. *Science (80-.).* **360** (2018), doi:10.1126/science.aat4422.

36. G. Lopes, N. Bonacchi, J. Frazão, J. P. Neto, B. V. Atallah, S. Soares, L. Moreira, S. Matias, P. M. Itskov, P. A. Correia, R. E. Medina, L. Calcaterra, E. Dreosti, J. J. Paton, A. R. Kampff, Bonsai: an event-based framework for processing and controlling data streams. *Front. Neuroinform.* **9** (2015), doi:10.3389/FNINF.2015.00007.

37. F. Sun, J. Zhou, B. Dai, T. Qian, J. Zeng, X. Li, Y. Zhuo, Y. Zhang, Y. Wang, C. Qian, K. Tan, J. Feng, H. Dong, D. Lin, G. Cui, Y. Li, Next-generation GRAB sensors for monitoring dopaminergic activity in vivo. *Nat. Methods 2020 1711.* **17**, 1156–1166 (2020).

38. K. M. Costa, A. Sengupta, G. Schoenbaum, The orbitofrontal cortex is necessary for learning to ignore. *Curr. Biol.* **31**, 2652-2657.e3 (2021).

39. K. M. Costa, R. Scholz, K. Lloyd, P. Moreno-Castilla, M. P. H. Gardner, P. Dayan, G. Schoenbaum, The role of the lateral orbitofrontal cortex in creating cognitive maps. *Nat. Neurosci. 2022 261.* **26**, 107–115 (2022).

40. J. Bonaventura, M. A. G. Eldridge, F. Hu, J. L. Gomez, M. Sanchez-Soto, A. M. Abramyan, S. Lam, M. A. Boehm, C. Ruiz, M. R. Farrell, A. Moreno, I. M. Galal Faress, N. Andersen, J. Y. Lin, R. Moaddel, P. J. Morris, L. Shi, D. R. Sibley, S. V. Mahler, S. Nabavi, M. G. Pomper, A. Bonci, A. G. Horti, B. J. Richmond, M. Michaelides, High-potency ligands for DREADD imaging and activation in rodents and monkeys. *Nat. Commun.* **10**, 1–12 (2019).

41. M. P. H. Gardner, J. S. Conroy, M. H. Shaham, C. V. Styer, G. Schoenbaum, Lateral Orbitofrontal Inactivation Dissociates Devaluation-Sensitive Behavior and Economic Choice. *Neuron.* **96**, 1192-1203.e4 (2017).

42. S. Kakade, P. Dayan, Dopamine: Generalization and bonuses. *Neural Networks.* **15**, 549–559 (2002).

43. W. Schultz, Dopamine reward prediction-error signalling: a two-component response. *Nat. Rev. Neurosci.* **17**, 183–195 (2016).

44. N. Bunzeck, E. Düzel, Absolute Coding of Stimulus Novelty in the Human Substantia Nigra/VTA. *Neuron.* **51**, 369–379 (2006).

45.   J. Morrens, Ç. Aydin, A. Janse van Rensburg, J. Esquivelzeta Rabell, S. Haesler, Cue-
Evoked Dopamine Promotes Conditioned Responding during Learning. *Neuron*. **106**,
142-153.e7 (2020).

46.   S. N. Haber, J. L. Fudge, N. R. McFarland, Striatonigrostriatal pathways in primates form
an ascending spiral from the shell to the dorsolateral striatum. *J. Neurosci.* **20**, 2369–82
(2000).

47.   I. Willuhn, L. M. Burgeno, B. J. Everitt, P. E. M. Phillips, Hierarchical recruitment of phasic
dopamine signaling in the striatum during the progression of cocaine use. *Proc. Natl.*
Acad. Sci. U. S. A. **109**, 20703–8 (2012).

48.   A. Mohebi, J. R. Pettibone, A. A. Hamid, J. M. T. Wong, L. T. Vinson, T. Patriarchi, L.
Tian, R. T. Kennedy, J. D. Berke, Dissociable dopamine dynamics for learning and
motivation. *Nat. 2019 5707759*. **570**, 65–70 (2019).

49.   E. E. Hart, M. J. Sharpe, M. P. Gardner, G. Schoenbaum, Responding to preconditioned
cues is devaluation sensitive and requires orbitofrontal cortex during cue-cue learning.
Elife. **9** (2020), doi:10.7554/eLife.59998.

50.   B. Engelhard, J. Finkelstein, J. Cox, W. Fleming, H. J. Jang, S. Ornelas, S. A. Koay, S. Y.
Thiberge, N. D. Daw, D. W. Tank, I. B. Witten, Specialized coding of sensory, motor and
cognitive variables in VTA dopamine neurons. *Nat. 2019 5707762*. **570**, 509–513 (2019).

51.   C. Kidd, B. Y. Hayden, The Psychology and Neuroscience of Curiosity. *Neuron*. **88**, 449–
460 (2015).

52.   J. Gottlieb, P. Y. Oudeyer, M. Lopes, A. Baranes, Information-seeking, curiosity, and
attention: computational and neural mechanisms. *Trends Cogn. Sci.* **17**, 585–593 (2013).

53.   S. Kobayashi, W. Schultz, Reward contexts extend dopamine signals to unrewarded
stimuli. *Curr. Biol.* **24**, 56–62 (2014).

54.   M. J. Sharpe, H. M. Batchelor, G. Schoenbaum, Preconditioned cues have no value.
Elife. **6** (2017), doi:10.7554/ELIFE.28362.

55.   K. Friston, The free-energy principle: a unified brain theory? *Nat. Rev. Neurosci. 2010*
112. **11**, 127–138 (2010).

56.   R. C. Wilson, Y. K. Takahashi, G. Schoenbaum, Y. Niv, Orbitofrontal cortex as a cognitive

map of task space. *Neuron*. **81**, 267–279 (2014).

57.   C. Liu, X. Cai, A. Ritzau-Jost, P. F. Kramer, Y. Li, Z. M. Khaliq, S. Hallermann, P. S.
587   Kaeser, An action potential initiation mechanism in distal axons for the control of
588   dopamine release. *Science (80-.)*. **375**, 1378–1385 (2022).

589   58.   P. F. Kramer, S. G. Brill-Weil, A. C. Cummins, R. Zhang, G. A. Camacho-Hernandez, A.
590   H. Newman, M. A. G. Eldridge, B. B. Averbeck, Z. M. Khaliq, Synaptic-like axo-axonal
591   transmission from striatal cholinergic interneurons onto dopaminergic fibers. *Neuron*. **110**,
592   2949-2960.e4 (2022).

593   59.   H. Zhang, D. Sulzer, Glutamate Spillover in the Striatum Depresses Dopaminergic
594   Transmission by Activating Group I Metabotropic Glutamate Receptors. *J. Neurosci.* **23**,
595   10585–10592 (2003).

Acknowledgements

We thank Marios Panayi, Angela Langdon, Matthew Gardner, and Evan Hart and for critical discussions. This work was supported by the Intramural Research Program at the National Institute on Drug Abuse. KMC received a post-doctoral fellowship from the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG; MA 8509/1-1).

Author contributions

KMC and GS designed experiments, interpreted results, and wrote the manuscript. KMC performed the experiments with technical assistance from NR, JM, and CS. KMC analyzed the data and wrote the first draft of the manuscript. GS supervised the research.

Competing interests

The authors declare no competing interests. The opinions expressed in this article are the authors' own and do not reflect the view of the NIH/DHHS.

Materials and correspondence

Additional information on materials and protocols are available upon request to Geoffrey Schoenbaum (geoffrey.schoenbaum@nih.gov).

Methods

Experimental Model and Subject Details

Experiments were performed on a total of 31 male Long-Evans rats (>3 months of age

at the start of the experiment, Charles River Laboratories) housed on a 12 hr light/dark

cycle at 25 °C. Rats were food restricted to ~85% of their original weight for the duration

of the experiments and were tested at the NIDA-IRP in accordance with NIH guidelines

determined by the Animal Care and Use Committee. All rats had *ad libitum* access to

water during the experiment and were fed 16-20 g of food per day, including rat chow

and pellets consumed during the behavioral task. Behavior was performed during the

light phase of the light/dark schedule. Prior to surgery and each experiment, rats were

handled by the experimenter, as previously described (*38, 39*).

Surgical procedures

For the main preconditioning experiment, rats (N=24) were anesthetized with 1-2%

isoflurane and prepared for aseptic surgery. They received unilateral infusions of AAV5-

hSyn-dLight1.2 into the NAcc (AP +1.7 mm, ML + or -1.7 mm, and DV -6.3 and -6.2 mm

from the brain surface), DMS (AP -0.4 mm, ML + or -2.6 mm, and DV -3.7 and -3.6 mm

from the brain surface) and dorsolateral striatum (DLS, AP +0.2 mm, ML + or -3.8 mm,

and DV -3.7 and -3.6 mm from the brain surface). Recordings from DLS were poorly

correlated with our task events, and seemingly correlated more with individual

movements; as that was not the focus of our study, these signals were not further

included in our analysis. A separate group of rats received unilateral infusions of AAV9-

hSyn-GRABDA2m (N=5) or AAV9-hSyn- GRAB-DA-mut (N=2) only in the NAcc, and

no additional viral infusions. A total 0.7 µL of each dopamine sensor-transfecting virus

was delivered in each site at 0.1 µL/min via an infusion pump. Optic fiber cannulas (200

µm diameter; Neurophotometrics, CA) were implanted in each site in the location of the

second (most dorsal) viral infusion.

Rats for the main preconditioning experiment, which were transfected with dLight1.2

and implanted in different areas of the striatum, also received in the same surgery

infusions of either AAV8-CaMKIIa-hM4d-mCherry (an inhibitory, Gi-coupled, designer

receptor exclusively activated by designer drugs (DREADD); OFCi group, N=12) or

AAV8-hSyn-mCherry (CTRL group, N=12), bilaterally into the lOFC (AP −3.0 mm, ML ±

3.2 mm, and DV -4.4 and -4.5 mm from the brain surface) (*38*). A total 0.5 µL of each

DREADD virus was delivered in each site at 0.1 µL/min via an infusion pump. All viruses

were obtained from Adgene. Exposed fiber ferrules and a protective black 3D-printed

headcap were secured to the skull with dental cement. After surgery, rats were given

Cephalexin (15 mg/kg po qd) for two weeks to prevent any infection. One rat in the

OFCi group had health complications after surgery and was removed from the study.

Fiber photometry

Dopamine-dependent fluorescence signals were recorded using custom ordered multi-

pronged fiber optic patch cables (200 µm diameter, 0.37 NA, Doric Lenses, Canada)

that were attached to the optic fiber ferrules on the skull of the rats with brass sleeves

(Thorlabs, NJ). Up to three fibers were connected at a time in each rat for recordings,

and they were shielded and secured with a custom 3D-printed headcap-swivel shielding

system that allowed for the relatively free movement of the rats without the use of optic

commutators and prevented the spillover of light during recordings.

Recordings were conducted using an FP3002 system (Neurophotometrics, CA), by

providing both 470 (active signal) and 415 nm (isosbestic reference) excitation light

through the patch cord in interleaved LED pulses at 100 Hz (50 Hz acquisition rate for

each channel). The light was reflected through a dichroic mirror and onto a 20×

Olympus objective. Excitation power was measured at ~150-200 µW at the tip of the

patch cord. Emitted fluorescent light was captured thorough the same cords, separated

with an image splitting filter, and captured via a high quantum efficiency CMOS camera.

Signals were acquired and synchronized with behavioral events using Bonsai (*36*). To

avoid photobleaching, recordings were triggered so that the system was only active

during behavioral trials (see next section).

Signals were processed using custom scripts in MATLAB (MathWorks, MA). Raw

fluorescence signals from the 470 (active) and 415 (reference) channels were first

filtered with a fourth-order median filter. Then the reference channel data was fitted to

the active signal using a second order polynomial fit, and the fitted data was subtracted

from the active channel, removing dopamine-independent variations in fluorescence.

The resulting signal was z-scored for each trial, using the 10 seconds before each trial

onset as a baseline. The resulting z-scores were normalized for each rat and session,

were the maximum observed peak was set to 100% and the mean was set at 0%.

For quality control, a group of rats (n=2) were infused with GRAB-DA-mut, a mutated

version of the GRAB-DA sensor that does not show dopamine-dependent fluorescence

and tested in similar tasks as the other experimental groups. All recordings in this group

showed no fluorescence peaks around any behavior event, confirming that the signals

we observed in the other groups were indeed due to dopamine release (Figure S1 and

S4).

Behavioral apparatus and general procedures

Rats were trained and tested at least eight weeks after the surgeries. All experiments

were conducted in standard behavioral boxes (12" x 10" x 12," Coulbourn Instruments,

PA), which were individually housed in light- and sound-attenuating boxes (Jim Garmon,

JHU Psychology Machine Shop). Each box was equipped with a food cup (recessed

magazine, a pellet dispenser, two wall speakers (one for white noise, and the other for

tones and sirens), a clicker, a house light, and a panel light. Head entries into the food

cup was measured based on breaks of an infra-red beam. For conditioning, 45 mg

sucrose pellets (5TUT, TestDiet, MO) were used as reinforcers. Intertrial intervals (ITIs)

varied around ~6 min on average and all cues lasted 10 seconds. Each rat experienced

one session per day, and the order of presentation of each cue was counterbalanced

across rats. Behavioral responses were quantified as the percentage of time that each

rat spent in the food cup during the last 5 seconds of each CS.

A computer equipped with GS3 software (Coulbourn Instruments, PA) controlled the

equipment. One Arduino microcontroller in each box recorded all events from the

Coulbourn equipment using a strobe based logic and relayed it as a serial input to the

Bonsai photometry interface. One master Arduino relayed trial initiation and end from

the Coulbourn interface to the FP3002 photometry hardware, triggering photometry

recordings only during trials and turning them off during the ITIs.

Sensory preconditioning task

Rats were first lightly shaped to retrieve sucrose pellets from the food cup. They were

then submitted to two sensory preconditioning sessions; these used a total of four

different auditory stimuli, drawn from stock equipment available from Coulbourn, which

included tone, siren, clicker, and white noise. Assignment of these stimuli to task cues

was counterbalanced across rats. In each pre-conditioning session, rats received six

blocked presentations of A-B pairs (A presented unexpectedly, followed by the

immediate presentation of B) and C-D pairs (similar arrangement as the A-B pairs), with

no direct reinforcement being present.

After pre-conditioning, rats in the main experiment were conditioned for six sessions,

where they received randomized presentations of B immediately followed by delivery of

two sucrose pellets, and D without any reinforcement.

In the final probe session, rats experience three reminder trials each where B was

presented followed by reward, and D was presented with no outcome, immediately

followed by blocked presentations, six each, of A and C, with no outcome. Prior to this

probe, each rat received an i.p. injection of JH60 (0.2 mg/kg, dissolved in 0.9% NaCl)

and was left in their home cage for at least 20 minutes before the start of the session, to

allow for the DREADD agonist to effectively inhibit transfected lOFC neurons in the

OFCi group (*38, 40*). Some rats in both groups did not show proper discriminative

conditioning in both the conditioning and probe sessions. To control for this, only rats

that responded twice as much to cue B than to cue D in the probe reminder trials were

included in the final study (N=6 for CTRL and N=8 for OFCi).

Two subsets of rats, infused only with GRAB-DA2m or GRAB-DA-mut in the VMS and

implanted with an optic fiber cannula in that same site, were subjected to the same

aforementioned preconditioning protocol for two sessions, after which they underwent a

third preconditioning session where the in first three trials of each pairing the cues were

presented in the normal order (A→B and C→D), whilst in the last three trials of each

pairing the cue order was reversed (B→A and D→C).

Histological procedures

After completion of the experiment, rats were perfused with chilled phosphate buffer

saline (PBS) followed by 4% paraformaldehyde in PBS. The brains were then immersed

in 18% sucrose in PBS for at least 24 hours and frozen. The brains were sliced at 40

μm and stained with DAPI (Vectashield-DAPI, Vector Lab, Burlingame, CA), and

processed for immunohistochemical detection of green fluorescent protein (Figures 1A

and S4B and C). For immunohistochemistry, the brain slices were first blocked in 10%

goat serum made in 0.1% Triton X-100/1× PBS and then incubated in anti-GFP

antibodies (1/1000, RT, overnight, Mouse anti-GFP, 632381, Takara Bio USA, WI)

followed by Alexa-488 secondary antibodies (1/200 , RT, 2h, Donkey anti-mouse Alexa-

488, 715-546-150, Jackson, Immunoresearch, PA). Fluorescent microscopy images of

the slides were acquired with a BZ-X800 Keyence microscope (Figures 1A and B, and

S4B and C).

Statistical analyses

Statistical analyses were performed in GraphPad Prism (GraphPad Software, San

Diego, CA). Error bars in figures denote the standard error of the mean. Effects of

experimental variables on behavioral and dopamine signal measures were examined

with t-tests if the data was normally distributed or Wilcoxon tests otherwise, as well as

repeated-measures 2-way and 3-way ANOVAs combined with Sidak's or Tukey's post-

hoc tests, respectively. Statistical significance threshold for all tests was set at $P<0.05$.

Supplementary Information

Figure S1. Acquisition and validation of dopamine photometry recordings. A: Cartoon schematic of the photometry approach for multi-site recording of dopamine signals in freely moving rats. Rats were implanted with multiple optic fiber cannulas and transfected with AAVs that led to the expression of dopamine sensors in target striatal regions, signals were collected using a Neurophotometrics FP3002 system, and reference (405 nm; ligand-insensitive fluorescence) and active (465) channels are separated using time-division multiplexing. Figure inspired by Martianova et al (*34*) and created with Biorender.com. **B:** Validation of active dopamine sensors (dLight1.2 and GRAB-DA2m) by comparison with a mutated sensor that does not exhibit dopamine-dependent changes in fluorescence (*37*). The panels show data from representative rats, with bold black trace representing mean responses overlayed on gray traces representing 16 individual trials centered around unexpected pellet deliveries in the operant box. Note the clear dopamine responses in the rats transfected with the active sensors, but a complete absence of response in the rats transfected with the mutated control fluorophore.

Figure S2. Behavioral responses during preconditioning and conditioning divided by groups. A: Percentage of time in port during preconditioning for both CTRL and OFCi rats. **B:** Same metric, but in conditioning. As mentioned in main text, there was no difference in behavior performance between groups in these phases.

Figure S3. Preconditioning and conditioning dopamine recordings divided by groups. A: Peak dopamine responses to predicted and unpredicted cues during preconditioning for both CTRL and OFCi rats. **B:** Difference in peak dopamine responses to the US and CS – a measure of RPE representation – during conditioning sessions. As mentioned in main text, there was no difference in quantified dopamine responses between groups in these phases.

Figure S4. Histological validation of rats transfected with GRAB-DA2m and

GRAB-DA-mut. A: Schematic representation of fiber tip locations in the brains of the

four rats transfected with GRAB-DA2m and used for the preconditioning switch

experiment, and for the two rats transfected with GRAB-DA-mut, used as a control for

our photometry recordings. **B:** Representative histology of a rat transfected with GRAB-

DA2m. C: Same as B, but for a rat transfected with GRAB-DA-mut.