1   **Plasticity drives geographic variation and trait coordination in blue oak drought**
2   **physiology**
3

4   Leander D.L. Anderegg[1], Robert P. Skelton[2], Jessica Diaz[3], Prahlad Papper[3], David D.
5   Ackerly[3,4], Todd E. Dawson[3,4]
6
1 – UC Santa Barbara, Ecology, Evolution & Marine Biology
Address: Ecology, Evolution & Marine Biology
University of California, Santa Barbara
Santa Barbara, CA 93106-9620
landeregg@ucsb.edu
541.790.1096
2 – South African Environmental Observation Network and School of Animal, Plant and
Environmental Sciences, University of Witwatersrand, Johannesburg, South Africa
3 – UC Berkeley, Integrative Biology
4 – UC Berkeley, Environmental Science, Policy & Management

17
18
19   Abstract
20

Two decades of widespread drought-induced forest mortality events on every forested continent
have raised the specter of future unpredictable, rapid ecosystem changes in 21[st] century forests.
Yet our ability to predict drought stress, much less drought-induced mortality across the
landscape remains limited. This uncertainty stems at least in part from an incomplete
understanding of within-species variation in hydraulic physiology, which reflects the interaction
of genetic differentiation among populations (ecotypic variation) and phenotypic plasticity in
response to growth environment. We examined among-population genetic differentiation in a
number of morphological and hydraulic traits in California blue oak (*Quercus douglasii*) using a
30 year old common garden. We then compared this genetic trait differentiation and trait-trait
integration to wild phenotypes in the field from the original source populations.
We found remarkably limited among-population genetic differentiation in all traits in the
common garden, but considerable site-to-site variation in the field. However, it was difficult to
explain trait variation in the field using site climate variables, suggesting that gridded climate
data does not capture the drivers of plasticity in drought physiology in this species. Trait-trait
relationships were also considerably stronger in the field than in the garden, particularly links
between leaf morphology, leaf hydraulic efficiency and stem hydraulic efficiency. Indeed, while
twelve of 45 potential trait-trait relationships showed significant wild phenotypic correlations,
only four relationships showed both genetic and phenotypic correlations, and five relationships
showed significantly different genetic and phenotypic correlations. Collectively, our results
demonstrate limited ecotypic variation in drought-related physiology but considerable
geographic variation in physiology and phenotypic integration in the wild, both driven largely by
plasticity.

43
44
45
46

Introduction:

In the 21st century, trees living in a hotter, more variable and often drier world will need to acclimate or adapt to avoid local extirpation, or shift their geographic ranges through multi-generational migration. Recent drought- or heat-induced forest mortality events highlight the vulnerability of even highly drought-adapted forests to climate change (Allen *et al.* 2010; Brodribb *et al.* 2020; Hammond *et al.* 2022). Mass tree die-offs also illustrate the potential for extremely rapid ecosystem changes in a warming world, which have profound ecological consequences and impacts on human society (Anderegg *et al.* 2012). Unfortunately, despite multiple decades of concerted effort to understand the causes of drought-induced forest mortality, we still struggle to predict when and where trees will die off during drought (Trugman *et al.* 2021).

Part of the large uncertainty about drought vulnerability is a poor understanding of physiological variation within species (Trugman *et al.* 2021). Within-species variation can be substantial (Martinez-Vilalta *et al.* 2009; Mclean *et al.* 2014; Anderegg *et al.* 2021), can decrease climate vulnerability in some populations (Laforest-Lapointe *et al.* 2014; Garcia-Forner *et al.* 2016), and can differ between co-occurring species (Anderegg & HilleRisLambers 2016). Drought resistance is the complex result of numerous plant traits, all of which potentially vary across populations within a species. A key drought tolerance trait, namely xylem resistance to embolism, often shows limited among-population variation in conifers (Lamy *et al.* 2013) and angiosperms (Skelton *et al.* 2019). But morphological characteristics, for example high leaf mass per area (LMA) and high leaf dry matter content (LDMC) may indirectly allow plants to maintain function under increasingly negative water potentials during drought (Lamont *et al.* 2002; Poorter *et al.* 2009; Bartlett *et al.* 2012). Meanwhile, shifting allocation to different tissues, for example decreasing leaf area to sapwood area ratio ($A_L:A_S$), and increasing tissue-specific hydraulic efficiency can help plants avoid drought stress by minimizing the water potential drop required to maintain transpirational flux (Gleason *et al.* 2013; Mencuccini *et al.* 2019). Across species, coordinated physiological strategies or trait syndromes lead to correlated variation across many of these traits (Reich *et al.* 2003; Chave *et al.* 2009; Mencuccini *et al.* 2015; Sanchez Martinez *et al.* 2020). However, within individual species, these broad evolutionary trait-trait correlations often break down (Anderegg *et al.* 2018; Messier *et al.* 2018; Rosas *et al.* 2019). Consequently, both the total amount of within-species variation in drought resistance-related traits and the key physiologic axes that drive spatial variation in drought resistance remain poorly understood.

Possibly more important than the total amount of within-species variation are the drivers of this variation. Intraspecific variation can arise from two distinct processes with drastically different implications for near-term climate responses. First, spatio-temporal variation in environmental selection on heritable fitness-related phenotypes can lead to genetic differentiation among populations and local adaptation of functional traits (traits that mediate plant performance in a given environment) (Alberto *et al.* 2013). Substantial geographic variation in drought physiology due to local adaptation would imply that certain populations or genotypes (and not others) are drought resistant and that gene-flow (assisted or natural) is necessary for other populations to manifest improved drought resistance (Hoffmann *et al.* 2021). Local adaptation is widespread in plants, particularly species with large population sizes (Leimu & Fischer 2008; Savolainen *et al.* 2013). Indeed, provenance trials or common gardens (where multiple populations of a species are planted in a common environment) almost always detect genetic differentiation among populations in trees (Alberto *et al.* 2013; Ramírez-Valiente *et al.* 2022)

and landscape genetic/genomic approaches often reveal evidence for local adaptation
(Savolainen *et al.* 2013; Capblancq *et al.* 2020; Gugger *et al.* 2021; Dauphin *et al.* 2022).
However, 'home site advantage' (a hallmark of local adaptation) has proven rarer than expected
(see meta-analyses Leimu & Fischer 2008; Oduor *et al.* 2016), though it is harder to test for, as it
requires reciprocal transplant experiments and lifetime fitness estimates.
Alternatively, adaptive trait plasticity, or the ability of each genotype to manifest a broad
range of phenotypes depending on environmental cues, can generate geographic variation in
water stress-related traits independent of genetic variation. If plasticity is not itself locally
adapted (i.e., all populations or genotypes have the same amount of plasticity), all populations of
a species could manifest drought resistant traits given the right environmental cues. As sessile
organisms that experience a range of environments over their lifetime and even in different parts
of the same organism (e.g., light availability in different parts of the canopy), plants show
marked plasticity in many traits (Poorter *et al.* 2009; Palacio-López *et al.* 2015; Keenan &
Niinemets 2016). Indeed, substantial plastic responses to drought and cold stress have been
observed in multiple tree species (Gimeno *et al.* 2009; Gárate-Escamilla *et al.* 2019). However,
the prevalence of active adaptive plasticity (as opposed to passive plasticity that results from
reduced growth or other stresses under environmental limitation) remains the subject of long
running debate (van Kleunen & Fischer 2005; Palacio-López *et al.* 2015).
Blue oak (*Quercus douglasii* Hook & Arn) is a culturally and ecologically important tree
species and the dominant species in many of the oak savannas in California, USA. Blue oak
experienced substantial and spatially widespread mortality during a major drought from 2012-
2016, particularly in the southern portion of its geographic range (Brown *et al.* 2018;
McLaughlin *et al.* 2020). This mortality highlights the potential vulnerability of blue oak in a
changing climate, necessitating an improved understanding of blue oak drought tolerance in
space and time to support proactive oak management. Blue oak grows across a wide latitude and
huge range of water availability, and thus likely exhibits substantial within-species phenotypic
variation. While blue oak exhibits minimal within-species variation in its xylem vulnerability to
embolism (P50, or the water potential at which xylem suffers 50% embolism, a key drought
tolerance metric, Skelton *et al.* 2019), drought avoidance traits such as hydraulic efficiency and
allocation to transpiring leaf area as well as additional traits related to drought tolerance, such as
leaf robustness (specific leaf area or SLA, leaf dry matter content or LDMC) are expected to
vary considerably across blue oak populations.
We quantified the extent and drivers of geographic variation in drought-related traits
within mature blue oak trees. We asked 1) how much do drought-related traits vary across the
landscape, 2) how is drought resistance coordinated across different traits and tissues, and 3)
does spatial variation and trait coordination arise from local adaptation or plasticity? We sought
to disentangle the relative roles of local adaptation versus plasticity by measuring drought-related
traits in a common garden experiment compared to the traits of the source populations in the
wild. In the garden, trait differences between populations are indicative of between-population
genetic differentiation/local adaptation (assuming limited maternal effects). Meanwhile,
differences between the traits measured in the common garden and those measured in the wild
source populations are due to plasticity.
135
Methods:
Blue oak (*Quercus douglasii* Hook & Arn.) is a deciduous species, which is endemic to
California but widespread and abundant throughout the state in low elevation woodlands around
the California Central Valley (Fig. S1). We measured morphological and hydraulic traits of

leaves and stems from seven populations of blue oak in the wild and in a ~30 year old,
reproductively mature common garden. These populations were selected to span the geographic
range and range of moisture availability of *Q. douglasii* (Fig. S1, Table S1), using only
populations whose original acorn source (for the common garden) had been accurately relocated,
and whose source populations were accessible based on current land tenure. Climate data for the
source populations, both 1951-1980 climate normal and 2018 water year meteorology, were
extracted from the California state-wide Basin Characterization Model (Flint *et al.* 2013).
147
Common garden sampling
To investigate the amount of among-population genetic trait differentiation, we sampled an
existing provenance trial/common garden planted at the Hopland Research and Extension Center
(CA, USA). This trial was planted in 1992 with acorns collected from 26 *Q. douglasii*
populations across California and planted in a randomized block design (J. McBride, pers.
comm.; see also McBride *et al.* 1997). We subsampled seven populations, capturing as much of
Q. douglasii's aridity range as possible. For each of the seven populations, we sampled five to
six individuals for hydraulic traits, leaf traits and branch wood density. Between April and mid-
June of 2018, we collected >1 m long branches from the sunlit, south-facing portion of the
canopy in the early morning (before 9am local time), relaxed the xylem by repeatedly recutting
the stem underwater, and then placed the branches in large plastic bags for immediate transport
back to the lab. One to two individuals from each of the seven sampled populations were
collected from the garden at each sampling point (once or twice weekly) to control for temporal
variation in hydraulic traits. Three small terminal twigs and subtending leaves were collected at
the same time for the measurement of morphological traits. Tree diameter at 50cm and total tree
height was also surveyed for every common garden tree in the winter of 2017 and used to
calculate the total stem volume of the largest stem (assuming the stem was a cylinder) as a metric
of total growth rate since planting.
166
167   *Wild population sampling*
168   We collected branches for trait measurement from six to eight mature trees from the acorn source
169   populations used in the garden. Large branches (typically >1.5m long and >4cm basal diameter)
170   from the south-facing canopy were collected in the same manner as in the common garden. Tree
171   cores were also collected from the sampled trees for calculation of radial growth rates.
172
Morphological Traits
We selected three terminal branches from the south-facing sun-exposed canopy for
morphological measurements (Table 1), cut at the prior year bud scar and including all current
year stem and leaf tissue. Branches were rehydrated for >12 hrs using the 'partial rehydration'
method (Pérez-Harguindeguy *et al.* 2013). We then measured specific leaf area (SLA, cm^2 wet
leaf area per g^{-1} leaf dry mass), leaf dry matter content (LDMC, g leaf dry mass per g^{-1} leaf wet
mass), median leaf size (cm^2 fresh leaf area), leaf dry mass to stem dry mass ratio ($M_L:M_S$), and
leaf area to sapwood area ratio ($A_L:A_S$, cm^2 leaf area per mm^2 stem area underneath bark) on
these current year terminal twigs using calipers, balances, and ImageJ image analysis software.
We calculated mean leaf size as the average area per leaf for each terminal branch. We also
measured wood density (WD, g dry mass per cm^3 wet volume) on one to five branch disks 2-5cm
in diameter cut from the basal end of branches collected for hydraulic sampling. Bark was
removed from branch disks and wet volume was measured via water displacement on a balance.

For all analyses using individual or population average traits, we also included leaf and
stem P50 values (the xylem tension causing 50% embolism) from Skelton *et al.* (2019), which
were measured on branches collected at the same time and using the same methods as branches
collected for hydraulic efficiency measurements reported here. However, P50 measurements had
few or no replicates per individual and were thus not suitable for branch-level analyses.
191
Hydraulic traits
Due to extremely long vessel lengths in oaks (often >1 m in branches of *Q. douglasii*, based on
pressurized branches recut at the basal end until air bubbles were seen at the distal end; RPS and
LDLA *pers obs*), classic hydraulic methods using stem segments were impossible without
considerable open vessel artifacts. Instead, we performed hydraulic measurements on terminal
branches of 1-2 mm basal diameter using the vacuum chamber method (Kolb *et al.* 1996). We
targeted current year growth when possible (similar to branch sampling for morphology), but
included up to three years of growth when the current year was <2 cm long and stem diameter
was not sufficiently large to fit in hydraulic tubing. For each terminal branch, all subtending
leaves were cut under water at the petiole with a razor blade, and the entire stem was inserted
into a vacuum chamber. Flow from a scale, through the branch and out the cut petioles was
induced by subjecting the stem to a ~60 kPa vacuum. Nevertheless, any stem that was suspected
of having an open vessel (i.e. any stem that had a high apparent conductance) was checked for
open vessels by attaching to pressurized nitrogen for ~10 minutes and checking under water for
air bubbles from the stem or petioles. The leaf area of all subtending leaves was measured via
flatbed scanner and ImageJ (Schneider *et al.* 2012), basal stem diameter at 4 radii averaged, and
the length of the stem and all branches measured with digital calipers.
Raw stem conductance was standardized two different ways. First, raw conductance was
standardized per unit leaf area, here termed k_{stem}, indicating leaf area-specific stem conductance
(i.e. not standardized by path length, so including the effects of differential stem growth in long
versus short stems, Table 1). Raw conductance was also divided by stem cross-sectional area and
multiplied by total stem length (including length of branches if stem was branched) to produce
sapwood area specific conductivity, or K_s.
Leaf hydraulic conductance was also measured on terminal branches (cut from the same
>1 m long branch as stem conductance) using the 'rehydration kinetics' (RK) method (Brodribb
& Holbrook 2003). Because leaves often had very small and irregular petioles, we measured
multiple leaves attached to a terminal twig. Leaf conductance was an order of magnitude lower
than stem conductance, meaning the effect of the stem in the conductance observed via the RK
measurements was negligible. Total leaf area for each sample was calculated via flatbed scanner
and ImageJ for each sample, and leaf-area specific leaf conductance, k_{leaf} was calculated. Only
measurements from leaves with an initial water potential < −0.2 MPa and > −2 MPa were
analyzed to avoid large measurement errors in samples with small pressure gradients or potential
embolism.
225
Growth rate:
We calculated growth rates for the common garden individuals based on a winter of 2017survey
of tree diameter at 50 cm and stem height. We calculated total stem volume growth since
planting, assuming the stem was a cylinder with the diameter of the basal diameter and a height
of tree height, using only the largest stem of multi-stemmed individuals. Growth rates were
calculated for the wild trees from tree cores, which were collected via increment borer at 1.3 m

height, mounted, sanded, and 5 years worth of rings identified using a dissecting microscope.
The length of the 2013-2018 5-year growth period was measured via digital calipers and radial
growth was converted to Basal Area Increment (BAI) based on tree DBH. Tree BAI is a function
of tree size (Figure S2), so BAI was standardized for tree size by calculating the percent of size-
specific maximum Basal Area Increment (BAI) over the 2013-2018 period. Size-specific
maximum BAI (the fastest rate a tree of a given DBH was observed to grow) was calculated for
each tree's DBH based on the 90th quantile regression of BAI versus tree DBH for all study trees
(n=32) plus tree core data from 29 trees spanning five additional sites across California to
expand the range of sampled tree sizes (total DBH range 7.8cm-104cm, Figure S2). The
observed BAI was then divided by the maximum BAI for each tree to produce '% max BAI'.
Tree cores were collected in Oct of 2018, but could not be collected from one site due to loss of
site access (n=6 sites rather than 7).
244
Variance decomposition analysis:
We performed variances decompositions, separately in the garden and the wild, on all branch-
level trait measurements to quantify trait variation within canopy (among branches), among
individuals within populations, and among populations. We fit linear mixed-effects models
separately for the garden or the wild traits, with a fixed intercept term, a random intercept for
population and a random intercept for individual nested within population. We employed the
lmer() function from the *lme4* (v 1.1-28) and *lmerTest* (v 3.1-3) packages. We then extracted the
random effect variance parameters, and calculated the proportion of total trait variance that was
attributed to among-population differences (population random effect), individual differences
within populations (individual random effect) or variation within individual tree canopies
(residual variance). We also calculated the coefficient of variation (CV = trait mean / trait
standard deviation) for each trait in the garden and in the wild. Finally, to compare the total trait
variation in the garden versus in the wild while accounting for unequal sample sizes, we
bootstrapped variance estimates, randomly sampling with replacement each dataset 1000 times
with sample number set to the minimum sample size of the two datasets (wild versus garden) for
each trait. We then compared the median bootstrapped trait variance for the wild versus garden.
261
Among-population trait differentiation and climatic variation
We tested for significant among-population trait differences by averaging trait values per
individual and then performing one-way ANOVAs separately in the garden and in the wild, and
calculated the omega-squared (ω^2) as an estimate of the proportion of variance explained by
among-population differences. Replicate was never a statistically significant factor in the garden,
and was excluded from the final ANOVAs (fitted using the aov() function in the *stats* package, v
4.1.2). We then used AICc (Akiake's Information Criterion corrected for small sample sizes,
AICc() function from the *MuMIn* package, v 1.43.17) to select the best single climatic predictors
of among-population differences for each trait. First, we determined whether a random intercept
for population was required to account for non-independence among individuals in a population
by fitting an 'over the top' fixed effects structure (Zuur *et al.* 2009), including mean annual
precipitation, mean annual potential evapotranspiration and mean minimum temperature) and
using likelihood ratio test to determine whether the population random effect was needed (using
the gls() and lme() functions from the *nlme* package, version 3.1-155). We then used AICc to
select the best climate predictor for each trait using linear models (lm in the *stats* package) or
linear mixed models with a population random intercept (lmer in the *lme4* package) and assessed

significance of the best predictor using t-tests, using Satterthwaite's degrees of freedom from the
lmerTest package (Kuznetsova *et al.* 2016). For the common garden, potential predictors
included only 30 year climate normals of mean annual precipitation (PPT_{30yr}), mean annual
potential evapotranspiration (PET_{30ry}), mean annual actual evapotranspiration (AET_{30yr}), mean
annual climatic water deficit ($CWD_{30yr}=PET_{30yr}-AET_{30yr}$), and mean annual minimum
temperature ($Tmin_{30yr}$). For wild traits, we also included growth year (2017-2018) weather
information as well: wet season (Nov-May) minimum temperature ($Tmin_{gy}$), maximum
temperature ($Tmax_{gy}$), precipitation (PPT_{gy}), potential evapotranspiration (PET_{gy}), actual
evapotranspiration (AET_{gy}) and climatic water deficit (CWD_{gy}), as well as the anomaly of the
2018 water year from the site 30 year normal (PPT_{anom}, PET_{anom}, AET_{anom}, CWD_{anom}). We
visually examined quantile-quantile plots and other patterns in the residuals to identify outliers
and ensure model assumptions were met.
290
Trait-trait coordination
To examine the magnitude of phenotypic (genetic plus plastic variation in the wild) and genetic
(variation in the garden) trait correlations, we calculated Pearson's correlation coefficients for all
pairs of individual-averaged traits in the wild and in the common garden. We visualized the
correlation structure using the corrplot() function in the *corrplot* package (v 0.92). We then used
a similar procedure to the bootstrapped variance comparisons to test for significant differences
between genetic and phenotypic trait correlations, randomly sampling individuals with
replacement from the entire wild and entire common garden datasets to produce two populations
of equal size to the smallest of the garden or wild complete pairwise observations, resampling
1000 times per trait combination. We then calculated the difference between the trait-trait
correlations in each bootstrapped pair of garden and wild datasets and determined the two-tailed
probability that the distribution of differences did not include 0 (e.g., alpha < 0.05 if either the
2.5[th] percentile of differences was greater than zero or the 97.5[th] percentile of differences is less
than zero, alpha < 0.1 if the 5[th] and 95[th] percentile range did not include zero, and alpha < 0.2 if
the 10[th] to 90[th] percentile range did not include zero).
306
307 All code and data for analyses and figure generation can be found at
308 https://github.com/leanderegg/BlueOakGarden.git, and all data are also available on Dryad at
309 [insert Dryad link upon acceptance].
310
311

Results

Genetic variation among populations was low in all morphological, allocation and
hydraulic traits in the common garden (Fig 1a). However, all traits except for stem sapwood
area-specific hydraulic conductivity (K_s) and leaf and stem vulnerability to embolism ($P50_{leaf}$,
$P50_{stem}$) showed substantial among-population variation in the wild. Among-population variation
in the wild was consistently 25%-30% of total trait variation in most traits, with SLA and LDMC
showing even larger among-population variation (46% and 40%, respectively, Fig 1b).
Meanwhile, branch-to-branch variation within tree canopies consistently made up the majority of
trait variation in the common garden, constituting >50% of total variation in all traits except WD
(42% of total) and over 75% of total variation in k_{leaf}, k_{stem}, and K_s (Fig. 1a). In the wild, within-
tree variation was typically less than 50% of total variation, except in $M_L:M_S$ and the three
hydraulic traits (Fig 1b).

324
325
326
When traits were averaged to the individual, 40-60% of the total tree-to-tree variation in
almost all traits was between populations in the wild (Fig 2a) and population differences were
significant in 8 of 11 traits (all but K_s, P50$_{leaf}$, and P50$_{stem}$) plus growth (ANOVAs p <0.05, Table
S2). Meanwhile, typically <30% of tree-to-tree variation was between populations in the garden,
and population differences were only significant in 2 of 10 traits (SLA and leaf size) plus growth
in the garden (Table S2). Leaf size was the one notable trait that showed significant and similar
among-population differentiation in both the wild and garden.
334
However, despite the consistent among-population variation in wild traits, climate
predictors (historical climate normals, meteorology of the water year of sampling, or the
sampling year anomaly from historical climate) very rarely explained differences among
populations (Fig. 2b). Based on AICc of linear or linear mixed models (when there was support
for a site random intercept, see Table S2), univariate climate predictors were only statistically
significant for four wild traits (Fig 2b) with WD decreasing with AET$_{anom}$, P50$_{stem}$ decreasing
(growing more negative) with AET$_{30yr}$, and M_L:M_S decreasing and k_{stem} increasing with Tmin$_{gy}$
(Figs S3,S4). (Fig 2b, Table S2). Meanwhile, in the garden three traits and growth were
significantly associated with source population climate (Table S2). In the garden, increasing
PPT$_{30yr}$ was significantly associated with decreasing wood density and increasing leaf size, while
increasing PET$_{30yr}$ was significantly associated with increasing SLA and decreasing growth (Fig
S3,S4).
While the among-population variance in the garden was always a small fraction of the
among-site variation in the wild, this did not generally result in decreased total trait variance the
garden (Figure 3). Bootstrapped estimates of the variance in individual averaged traits (i.e.,
among individual variance) generally showed larger median variance estimates in wild traits than
the common garden, but ratio of the garden variance to wild variance was only significantly less
than one (alpha <0.05) for SLA, LDMC, and WD. Meanwhile, M_L:M_S and k_{leaf} had similar
variances in the wild and the garden despite significant among-population differentiation on the
landscape and not in the garden, and A_L:A_S and k_{stem} had only marginally higher wild variances.
K_s, P50$_{[stem]}$ and P50$_{[leaf]}$ had similar variances in the wild compared to the garden and no
significant among-population differentiation in either location (Table S2), while growth could
not be compared because of differing metrics (% of max basal area increment vs 30 yr stem
volume). Ultimately, population-average trait values were entirely uncorrelated in the garden and
in the wild except for M_L:M_S (Figure S2), even for the traits with similar amounts of total
variation in the two settings.
361
362
Trait coordination
Trait-trait coordination was much stronger in the wild than in the common garden,
suggesting phenotypic correlations are not generally driven by genetic correlations or constraints
but rather by coordinated plasticity. Only six of 55 possible trait-trait correlations were
significant in both the garden and the wild (Fig. 4), almost all of which are unsurprising (e.g.,
A_L:A_S ~ M_L:M_S, SLA ~ LDMC) or mathematically related (K_s ~ k_{stem} or K_s~A_L:A_S because K_s
was calculated as k_{stem} * A_L:A_S * branch length). The main notable correlations consistent across

the garden and the wild trees were positive relationships between average leaf size and leaf vs
stem allocation on both an area basis ($A_L:A_S$) and a mass basis ($M_L:M_S$). Moreover, average leaf
size was much more strongly associated with variation in $A_L:A_S$ than number of leaves per
branch in both the garden and the wild (Fig. S3), suggesting that leaf expansion may be a key
regulator of leaf-to-stem allocation.
Beyond these correlations, only three additional trait correlations were significant in the
garden, while 10 additional trait correlations were significant in the wild (Fig 4). The change in
trait-trait correlations from garden to wild was statistically significant ($\alpha < 0.05$) in four trait pairs
and marginally significant ($\alpha < 0.1$) in two more pairs (based on 1000 bootstrapped correlation
comparisons, Fig 5). The most striking change in trait coordination occurred amongst k_{leaf}, k_{stem}
and $A_L:A_S$. Whereas there was only weak coordination between k_{stem}, k_{leaf} and $A_L:A_S$ in the
garden, there was a strong positive correlation between leaf area-specific branch and leaf
conductance in the wild (significantly different in the wild vs garden, $p < 0.05$, Fig 5) as well as
strong negative relationships between both leaf and stem conductance and $A_L:A_S$. This plastic
increase in hydraulic integration in wild trees appears to be driven by leaf size, which is strongly
negatively correlated with k_{leaf} and k_{stem} in the wild (change in correlation is statistically
significant for k_{stem}). Across the landscape, trees with smaller leaves had lower $A_L:A_S$, but
consequently greater k_{leaf} and k_{stem}.
393
394
Trait-Growth relationships
In the garden, traits were generally not strongly predictive of population average growth
(30 year height). Contrary to expectations, LDMC was positively and k_{stem} negatively correlated
with growth rates. Leaf size was the only trait correlated with growth in the expected direction,
with populations with larger leaves having higher average growth rates (Fig. S4). In the wild, no
traits were correlated with radial growth rates (calculated as the percent of maximum size-
specific Basal Area Increment), with the exception of K_s (Fig. S5), though growth was only
measured in six of the seven populations due to lost access at one site after initial trait sampling.
K_s was negatively correlated with growth, contrary to expectations.
404
Discussion
We found very little among-population genetic differentiation but consistent among-
population variation across the landscape in morphological and hydraulic traits of a widespread
oak species. This landscape-scale plasticity could not be easily explained using gridded climate
or annual meteorological data, but drove trait-trait coordination – particularly of leaf and stem
hydraulic traits – that was otherwise lacking in the common garden.
411
Traits in the common garden: little local adaptation
In a relatively mesic common garden (978 mm of mean annual precipitation, greater than
all but two of the sampled populations, Table S1), we found very little evidence for genetically
based trait variation among populations. Populations showed significantly different growth rates

(height attained at 30 years) in the garden, but only SLA and leaf size showed significant among-
population variation (Figure 2). Among individual trees in the common garden, there was
remarkably little trait coordination to suggest strong genetic correlations among traits, even
though many of these traits are correlated across species (Mencuccini *et al.* 2019; Sanchez
Martinez *et al.* 2020). Moreover, this lack of coordination was not necessarily an artifact of
limited total trait variation in the common garden, as only three of 11 possible traits showed
significantly smaller variance in the garden than in the wild (Fig. 3). Thus, the trait variation
manifested in the garden was not merely the wild trait variance less among-site plasticity, and
neither the lack of trait coordination (Fig 4) nor the lack of correlation between wild and garden
trait values (Fig. S5) was likely driven by restricted garden trait variance.
At the same time, the only traits that showed expected relationships with growth rates in
the common garden were leaf to stem allocation traits ($M_L:M_S$ and potentially $A_L:A_S$), with
leafier populations showing faster growth, and leaf size, with larger leaved populations showing
faster growth. We conclude that changes to allometry were the primary drivers of functional
differences among populations in the common garden in line with classic predictions about the
drivers of growth rate variation (Lambers & Poorter 1992), and that this variation in allocation
was linked in part to genetically determined variation in leaf size (Fig. S3). Meanwhile, leaf dry
matter content (LDMC) and leaf-specific stem conductance (k_s) were significantly correlated
with growth rates, but in a direction contrary to expectations. For LDMC, we interpret this as a
consequence rather than a cause of growth rate variation (faster growing trees had accumulated
more photosynthate in their leaves by the sampling period and thus had higher LDMC), and for
k_s, rather than high conductivity promoting growth (Hajek *et al.* 2014) slower growth in high k_s
populations appears to be a consequence of high hydraulic efficiency in the two most arid
populations that grew slowly in the garden.
440
Traits in the wild
In contrast to the common garden results, all traits except K_s and $P50_{stem, leaf}$ showed
substantial among-population trait differentiation across *Quercus douglasii*'s geographic range.
While all traits exhibited large branch-to-branch variation within a canopy (in both the garden
and the wild), among-population trait differences were as large or larger than differences among
trees in the wild (Fig 1b, 2a). Thus, it appears that spatial variation in drought-related physiology
is substantial in blue oak, and is primarily driven by trait plasticity.
Surprisingly, this site-to-site trait variability was not strongly related to available site
climate data. Neither 30 year site climate normals nor meteorological data from the water year of
sample collection, nor the climate anomaly of the sampled year compared to climate normals
predicted among-site trait variation. For the four traits that were significantly predicted by site
climate, winter minimum temperatures ($Tmin_{gy}$) was the best predictor for two traits ($M_L:M_S$ and
k_{stem}) and a water balance metrics (some actual evapotranspiration metric, AET_{anom} and AET_{30yr})
was only a significant predictor for two traits (WD and k_{leaf}, respectively). We hypothesize that
this discrepancy is the result of *Quercus douglasii*'s deep rooting strategy, which combined with
the substantial geologic and edaphic variation among the sampled sites to largely decouple the
actual water availability (the 'weather underground', sensu (McLaughlin *et al.* 2020)) from
above-ground climate.
Strikingly, the plasticity-driven trait variation across California resulted in strong trait-
trait coordination on the landscape, particularly among hydraulic efficiency and allocation traits.
Leaf and branch morphology and P50 were largely uncoupled from other traits in both the wild

and the common garden (Fig 4). However, k_{leaf} and k_{stem} significantly changed their correlation
strength from uncorrelated in the garden to positively correlated in the wild, and both traits
showed a strong correlation with both leaf size and $A_L:A_S$ in the wild. Collectively, this resulted
in strong integration across a leaf and stem 'hydraulic module', wherein trees with small leaves
had small $A_L:A_S$ ratios and high leaf area-specific leaf and stem hydraulic conductance. We
found no evidence for a tradeoff between safety and efficiency in either leaves or stems (Fig S6).
We also found very little evidence for trait-growth relationships in the wild, despite much
larger among-population trait differentiation. The only significant relationship was a decrease in
growth with increasing K_s, which may be consistent with increasing drought avoidance in the
driest populations (Fig S5). This lack of wild trait-growth relationship is perhaps not surprising,
given that both traits and environmental conditions and resources differ among populations,
making it difficult to disentangle cause versus effect.
474
Leaf area-driven hydraulic coordination
In theory, the coordination or integration of plant traits across tissues should optimize
whole-plant function (Reich 2014). However, perhaps partly as a result of traits rarely being
measured simultaneously for multiple tissues, this coordination is not universally observed
among tissues (Vleminckx *et al.* 2021) or types of traits (Sanchez Martinez *et al.* 2020), and the
actual mechanisms of this coordination (e.g., coordinated evolution, developmental constraints,
etc.) remain elusive. The leaf and stem hydraulic integration that we found across wild
populations suggests that, within individual species, plastically determined leaf size, presumably
coordinated by leaf expansion, may be a key mechanism coordinating hydraulic efficiency across
tissues.
We found little evidence for genetic correlations among hydraulic traits in a common
environment (Fig 4a), suggesting that hydraulic variation within species was not due to genetic
linkage or coordinated microevolution. However, strong hydraulic integration between the leaf
area-specific hydraulic conductance of leaves and stems across wild populations indicates
coordinated trait plasticity. This coordination seems to be mediated by allocation to leaf area
($A_L:A_S$), which was largely driven by the average size of leaves (rather than leaf number, Figure
S3). Average leaf size had a larger coefficient of variation than stem diameter (0.41 versus 0.31,
respectively), and had a stronger negative correlation with both k_{leaf} ($\rho = -0.43$) and k_{stem} ($\rho=$
-0.54) than $A_L:A_S$ itself did with either trait ($\rho= -0.39$ and $\rho= -0.30$ for k_{leaf} and k_{stem}
respectively). Further supporting the importance of plasticity, average leaf size was uncorrelated
with either k_{leaf} or k_{stem} in the garden, and the increase in the correlation from garden to wild was
strong for both traits and statistically significant for k_{stem} ($p < 0.05$). Meanwhile, the more
commonly measured sapwood-specific branch conductivity (K_s), which is standardized for
branch xylem conductive area and path length, was highly variable but not differentiated among
populations (Fig 1) and unrelated to either leaf size or k_{leaf} in the wild. This may partly be due to
the difficulty of measuring path length for our K_s estimates, because in order to avoid open
vessels in this long-vesseled oak species we had to measure the conductance of whole terminal
branches cut at the petioles (rather than branch segments) and estimate the path length from the
total branch length, even though leaf attachment to the branch occurred along much of its length.
Thus, k_{stem} (conductance of the whole branch per unit leaf area) was likely a more accurate
measurement of hydraulic efficiency than K_s (conductivity per sapwood area for a 1cm stem
length) in this study.

Ultimately, it appears that whole-branch conductance of terminal branches and leaf
conductance were coordinated in the wild by plastic changes in leaf size, presumably regulated
by environmental cues governing spring leaf expansion, rather than changes in the xylem
efficiency itself. K_s, while highly variable, did not differ among populations either in the garden
or the wild (Fig 1) and therefore could not drive this pattern. A developmentally fixed hydraulic
architecture (in both leaves and stems) that then gets allocated to supply a variable amount of
leaf area based on leaf expansion presents a parsimonious mechanism for regulating whole-tissue
hydraulic resistance, as regulation of transpirational area may be much easier to regulate in
response to environmental cues through turgor driven leaf expansion than anatomical changes to
xylem efficiency. Moreover, the negative relationship between leaf size and k_{leaf}/k_{stem}
documented here contrasts with the lack of relationship between k_{leaf} and leaf size among species
and suggests a developmental constraint rather than adaptive mechanism.
519
Management implications
Our results from mature trees would indicate that genetically informed management or
restoration of blue oaks may not require strong consideration of the aridity of the source
materials, as mature tree drought resilience appears to be determined by canalized thresholds of
xylem vulnerability to drought (Skelton *et al.* 2019) and largely plastic adjustments to drought
avoidance-related traits (e.g. k_{branch} and k_{leaf}). While we could not predict trait variation across the
landscape based on site climate (Fig 2b), the substantial among-population variation in leaf-to-
stem allocation and hydraulic efficiency are presumably adaptive and therefore constitute plastic
adjustments to drought exposure by increasing plant hydraulic efficiency. This decreases the
minimum leaf water potential required to attain a given rate of transpiration (Whitehead & Jarvis
1981; Mencuccini *et al.* 2019; Trugman *et al.* 2019) and is quite consistent with within-species
adjustments to promote drought avoidance in numerous other systems (Mencuccini & Grace
1995; Martinez-Vilalta *et al.* 2009; Anderegg & HilleRisLambers 2016; Rosas *et al.* 2019;
Anderegg *et al.* 2021).
Our findings of limited local adaptation are also consistent with the relatively small
amount of local adaptation found in blue oak growth rates (Fig 1) and phenology (Papper &
Ackerly 2021), and the relatively high rates of gene flow in blue oak (Papper 2021). While other
California oak species have been found to show strong genetic structure across their range (Sork
et al. 2013; Gugger *et al.* 2021), evidence suggests that blue oak population differentiation across
its broad geographic distribution is more limited. Given the strong potential for plastic
adjustments to plant hydraulics documented here, this might suggest that most populations can
(and may) manifest more drought avoidant phenotypes in a warmer and drier future climate,
regardless of their climate history.
Two important caveats to our finding of limited drought-related local adaptation are 1)
that we only studied reproductively mature trees and 2) that we only studied traits in a relatively
mesic garden. Local adaptation at the earliest life stages may be most relevant for restoration,
and limited local adaptation in adult phenotypes does not guarantee limited local adaptation in
juvenile phenotypes. Moreover, we were only able to perform trait measurements in a relatively
wet common garden. If the expression of genetic differences depends on the environment – e.g.,
if there are strong GxE interactions for drought-related phenotypes – then our results could differ
entirely in a drier common garden location. It remains possible that genotypes from drier
populations have larger potential for plasticity (as documented in a *Eucalyptus* species, Mclean
et al. 2014), which could imply that the same source populations planted in a dry location could

show substantial among-population variation due to the dry-adapted populations manifesting
more plasticity than wet-adapted populations.
555
556

Conclusion
In a study of range-wide trait variation in blue oak (*Quercus douglasii*), we found little
evidence for local adaptation in any of numerous morphological or hydraulic traits but
substantial realized trait varíation across the landscape. Our results suggest that this landscape-
scale trait variation involved plastic trait adjustments, and that this plasticity drove among-tissue
trait coordination. Thus, plastic adjustments, likely during the process of leaf development and
expansion, drive the coordinated manifestation of drought avoidant phenotypes with high
hydraulic efficiency in leaves and stems and increased carbon investment in stems relative to
leaves. However, these phenotypes could not be predicted by site climate and were poorly linked
to growth rates, indicating that our understanding of blue oak drought exposure and integrated
performance remains limited.
568
569

Acknowledgements
We acknowledge the Traditional Custodians and Owners of California, and recognize
their continuing connection to land upon which this research was conducted. We particularly
recognize the Chumash, Sierra Miwok, Konkow, Pomo, Northern Wintu, on whose traditional
territory our labs and study sites sit. We thank the Hopland Research Extension Center for access
to the common garden and keeping the trees alive for multiple decades and the Ranger Station at
Sonora for access to trees on their property. We thank CALFIRE for saving the common garden
trees from the Mendocino Complex Fire. We also acknowledge funding from the National
Science Foundations (NSF 1457400 to DDA & TED; NSF DBI-1711243, 2003205, and 2216855
to LDLA) and the National Oceanographic and Atmospheric Administration (Climate and
Global Change Fellowship to LDLA).

589 **Literature Cited**
590

Alberto, F.J., Aitken, S.N., Alía, R., González-Martínez, S.C., Hänninen, H., Kremer, A., *et al.*
(2013). Potential for evolutionary responses to climate change - evidence from tree
populations. *Global Change Biology*, 19, 1645–1661.
Allen, C.D., Macalady, A.K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., *et al.*
(2010). A global overview of drought and heat-induced tree mortality reveals emerging
climate change risks for forests. *Forest Ecology and Management*, 259, 660–684.

597 Anderegg, L.D. & HilleRisLambers, J. (2016). Drought stress limits the geographic ranges of
598   two tree species via different physiological mechanisms. *Global Change Biology*, 22,
599   1029–1045.
600 Anderegg, L.D.L., Berner, L.T., Badgley, G., Sethi, M.L., Law, B.E. & Hillerislambers, J.
601   (2018). Within-species patterns challenge our understanding of the leaf economics
602   spectrum. *Ecol. Lett.*, 21, 734–744.
603 Anderegg, L.D.L., Loy, X., Markham, I.P., Elmer, C.M., Hovenden, M.J., Hillerislambers, J., *et*
604   *al.* (2021). Aridity drives coordinated trait shifts but not decreased trait variance across
605   the geographic range of eight Australian trees. *New Phytologist*, 15, 343.
606 Anderegg, W.R.L., Kane, J.M. & Anderegg, L.D.L. (2012). Consequences of widespread tree
607   mortality triggered by drought and temperature stress. *Nature Climate change*, 1–7.
608 Bartlett, M.K., Scoffoni, C., Ardy, R., Zhang, Y., Sun, S., Cao, K., *et al.* (2012). Rapid
609   determination of comparative drought tolerance traits: using an osmometer to predict
610   turgor loss point. *Methods in Ecology and Evolution*, 3, 880–888.
611 Brodribb, T.J. & Holbrook, N.M. (2003). Stomatal Closure during Leaf Dehydration, Correlation
612   with Other Leaf Physiological Traits. *PLANT PHYSIOLOGY*.
613 Brodribb, T.J., Powers, J., Cochard, H. & Choat, B. (2020). Hanging by a thread? Forests and
614   drought. *Science*, 368, 261–266.
615 Brown, B.J., McLaughlin, B.C., Blakey, R.V. & Morueta-Holme, N. (2018). Future vulnerability
616   mapping based on response to extreme climate events: Dieback thresholds in an endemic
617   California oak. *Diversity Distrib.*, 24, 1186–1198.
618 Capblancq, T., Fitzpatrick, M.C., Bay, R.A., Exposito-Alonso, M. & Keller, S.R. (2020).
619   Genomic Prediction of (Mal)Adaptation Across Current and Future Climatic Landscapes.
620   *Annual Review of Ecology, Evolution, and Systematics*, 51, 245–269.
621 Chave, J., Coomes, D., Jansen, S., Lewis, S.L., Swenson, N.G. & Zanne, A.E. (2009). Towards a
622   worldwide wood economics spectrum. *Ecology Letters*, 12, 351–366.
623 Dauphin, B., Rellstab, C., Wüest, R.O., Karger, D.N., Holderegger, R., Gugerli, F., *et al.* (2022).
624   Re-thinking the environment in landscape genomics. *Trends in Ecology & Evolution*.
625 Flint, L.E., Flint, A.L., Thorne, J.H. & Boynton, R. (2013). Fine-scale hydrologic modeling for
626   regional landscape applications: the California Basin Characterization Model
627   development and performance. *Ecological Processes*, 2, 25.
628 Gárate-Escamilla, H., Hampe, A., Vizcaíno-Palomar, N., Robson, T.M. & Benito Garzón, M.
629   (2019). Range-wide variation in local adaptation and phenotypic plasticity of fitness-
630   related traits in Fagus sylvatica and their implications under climate change. *Global*
631   *Ecology and Biogeography*, 28, 1336–1350.
632 Garcia-Forner, N., Sala, A., Biel, C., Savé, R. & Martinez-Vilalta, J. (2016). Individual traits as
633   determinants of time to death under extreme drought in Pinus sylvestrisL. *Tree*
634   *physiology*, 36, 1196–1209.
635 Gimeno, T.E., Pías, B., Lemos-Filho, J.P. & Valladares, F. (2009). Plasticity and stress tolerance
636   override local adaptation in the responses of Mediterranean holm oak seedlings to
637   drought and cold. *Tree physiology*, 29, 87–98.
638 Gleason, S.M., Butler, D.W., Waryszak, P. & Graham, E.L.E. (2013). Shifts in Leaf and Stem
639   Hydraulic Traits across Aridity Gradients in Eastern Australia. *International Journal of*
640   *Plant Sciences*, 174, 1292–1301.

Gugger, P.F., Fitz-Gibbon, S.T., Albarrán-Lara, A., Wright, J.W. & Sork, V.L. (2021).
Landscape genomics of Quercus lobata reveals genes involved in local climate adaptation
at multiple spatial scales. *Molecular Ecology*, 30, 406–423.
Hajek, P., Leuschner, C., Hertel, D., Delzon, S. & Schuldt, B. (2014). Trade-offs between xylem
hydraulic properties, wood anatomy and yield in Populus. *Tree Physiology*, 34, 744–756.
Hammond, W.M., Williams, A.P., Abatzoglou, J.T., Adams, H.D., Klein, T., López, R., *et al.*
(2022). Global field observations of tree die-off reveal hotter-drought fingerprint for
Earth's forests. *Nat Commun*, 13, 1761.
Hoffmann, A.A., Miller, A.D. & Weeks, A.R. (2021). Genetic mixing for population
management: From genetic rescue to provenancing. *Evolutionary Applications*, 14, 634–
652.
Keenan, T.F. & Niinemets, U. (2016). Global leaf trait estimates biased due to plasticity in the
shade. *Nature Plants*, 1–6.
van Kleunen, M. & Fischer, M. (2005). Constraints on the evolution of adaptive phenotypic
plasticity in plants. *New Phytologist*, 166, 49–60.
Kolb, K.J., Sperry, J.S. & Lamont, B.B. (1996). A method for measuring xylem hydraulic
conductance and embolism in entire root and shoot systems. *Journal of Experimental ….*
Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. (2016). lmerTest Package: Tests in
linear mixed effects models. R package version 2.0-30. *Journal of Statistical Software*,
82, 1–12.
Laforest-Lapointe, I., Martinez-Vilalta, J. & Retana, J. (2014). Intraspecific variability in
functional traits matters: case study of Scots pine. *Oecologia*, 175, 1337–1348.
Lambers, H. & Poorter, H. (1992). Inherent Variation in Growth Rate Between Higher Plants: A
Search for Physiological Causes and Ecological Consequences. Elsevier, pp. 283–362.
Lamont, B.B., Groom, P.K. & Cowling, R.M. (2002). High leaf mass per area of related species
assemblages may reflect low rainfall and carbon isotope discrimination rather than low
phosphorus and nitrogen concentrations. *Functional Ecology*, 16, 403–412.
Lamy, J.-B., Delzon, S., Bouche, P.S., Alía, R., Vendramin, G.G., Cochard, H., *et al.* (2013).
Limited genetic variability and phenotypic plasticity detected for cavitation resistance in
a Mediterranean pine. *New Phytologist*.
Leimu, R. & Fischer, M. (2008). A Meta-Analysis of Local Adaptation in Plants. *PLOS ONE*, 3,
e4010.
Martinez-Vilalta, J., Cochard, H., Mencuccini, M., Sterck, F., Herrero, A., Korhonen, J.F.J., *et*
al. (2009). Hydraulic adjustment of Scots pine across Europe. *New Phytologist*, 184,
353–364.
McBride, J., Norberg, E., Bertenshaw, J.L., Kloss, S. & Mossadegh, A. (1997). Genetic
Variation in Shoot Growth, Phenology, and Mineral Accumulation of Northern and
Central Sierra Nevada Foothill Populations of Blue Oak 1. *USDA Forest Service General*
Technical Reports, 160.
McLaughlin, B.C., Blakey, R., Weitz, A.P., Feng, X., Brown, B.J., Ackerly, D.D., *et al.* (2020).
Weather underground: Subsurface hydrologic processes mediate tree vulnerability to
extreme climatic drought. *Global Change Biology*, 26, 3091–3107.
Mclean, E.H., Prober, S.M., Stock, W.D., Steane, D.A., Potts, B.M., Vaillancourt, R.E., *et al.*
(2014). Plasticity of functional traits varies clinally along a rainfall gradient in Eucalyptus
tricarpa. *Plant, Cell & Environment*, 37, 1440–1451.

Mencuccini, M. & Grace, J. (1995). Climate influences the leaf area/sapwood area ratio in Scots pine. *Tree physiology*, 15, 1–10.

Mencuccini, M., Minunno, F., SALMON, Y., Martinez-Vilalta, J. & Hölttä, T. (2015). Coordination of physiological traits involved in drought-induced mortality of woody plants. *New Phytologist*, n/a-n/a.

Mencuccini, M., Rosas, T., Rowland, L., Choat, B., Cornelissen, H., Jansen, S., *et al.* (2019). Leaf economics and plant hydraulics drive leaf : wood area ratios. *New Phytologist*, 224, 1544–1556.

Messier, J., Violle, C., Enquist, B.J., Lechowicz, M.J. & McGill, B.J. (2018). Similarities and differences in intrapopulation trait correlations of co-occurring tree species: consistent water-use relationships amid widely different correlation patterns. *American Journal of Botany*, 105, 1477–1490.

Oduor, A.M.O., Leimu, R. & van Kleunen, M. (2016). Invasive plant species are locally adapted just as frequently and at least as strongly as native plant species. *Journal of Ecology*, 104, 957–968.

Palacio-López, K., Beckage, B., Scheiner, S. & Molofsky, J. (2015). The ubiquity of phenotypic plasticity in plants: a synthesis. *Ecol Evol*, 5, 3389–3400.

Papper, P.D. (2021). Three scales of gene flow in California White Oaks. PhD. UC Berkeley.

Papper, P.D. & Ackerly, D.D. (2021). Partitioning genetic and environmental components of phenological variation in Quercus douglasii (Fagaceae). *madr*, 68, 425–433.

Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., *et al.* (2013). New handbook for standardised measurement of plant functional traits worldwide. *Aust. J. Bot.*, 61, 167.

Poorter, H., Niinemets, U., Poorter, L., Wright, I.J. & Villar, R. (2009). Causes and consequences of variation in leaf mass per area (LMA): a meta-analysis. *New Phytologist*, 182, 565–588.

Ramírez-Valiente, J.A., Santos del Blanco, L., Alía, R., Robledo-Arnuncio, J.J. & Climent, J. (2022). Adaptation of Mediterranean forest species to climate: Lessons from common garden experiments. *Journal of Ecology*, 110, 1022–1042.

Reich, P.B. (2014). The world-wide "fast–slow" plant economics spectrum: a traits manifesto. *Journal of Ecology*, 102, 275–301.

Reich, P.B., Wright, I.J., Cavender Bares, J., Craine, J.M., Oleksyn, J., WESTOBY, M., *et al.* (2003). The Evolution of Plant Functional Variation: Traits, Spectra, and Strategies. *Int. J Plant Sci.*, 164, S143–S164.

Rosas, T., Mencuccini, M., Barba, J., Cochard, H., Saura-Mas, S. & Martinez-Vilalta, J. (2019). Adjustments and coordination of hydraulic, leaf and stem traits along a water availability gradient. *New Phytologist*, 223, 632–646.

Sanchez Martinez, P., Martinez-Vilalta, J., Dexter, K.G., Segovia, R.A. & Mencuccini, M. (2020). Adaptation and coordinated evolution of plant hydraulic traits. *Ecol. Lett.*, 23, 1599–1610.

Savolainen, O., Lascoux, M. & Merilä, J. (2013). Ecological genomics of local adaptation. *Nat Rev Genet*, 14, 807–820.

Schneider, C.A., Rasband, W.S. & Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat Meth*, 9, 671–675.

Skelton, R.P., Anderegg, L.D.L., Papper, P., Reich, E., Dawson, T.E., Kling, M., *et al.* (2019).
No local adaptation in leaf or stem xylem vulnerability to embolism, but consistent
vulnerability segmentation in a North American oak. *New Phytologist*, 223, 1296–1306.
Sork, V.L., Aitken, S.N., Dyer, R.J., Eckert, A.J., Legendre, P. & Neale, D.B. (2013). Putting the
landscape into the genomics of trees: approaches for understanding local adaptation and
population responses to changing climate. *Tree Genetics & Genomes*, 9, 901–911.
Trugman, A.T., Anderegg, L.D.L., Anderegg, W.R.L., Das, A.J. & Stephenson, N.L. (2021).
Why is Tree Drought Mortality so Hard to Predict? *Trends in Ecology & Evolution*,
S0169534721000355.
Trugman, A.T., Anderegg, L.D.L., Sperry, J.S., Wang, Y., Venturas, M. & Anderegg, W.R.L.
(2019). Leveraging plant hydraulics to yield predictive and dynamic plant leaf allocation
in vegetation models with climate change. *Global Change Biology*, 25, 4008–4021.
Vleminckx, J., Fortunel, C., Valverde-Barrantes, O., Timothy Paine, C.E., Engel, J., Petronelli,
P., *et al.* (2021). Resolving whole-plant economics from leaf, stem and root traits of 1467
Amazonian tree species. *Oikos*, 130, 1193–1208.
Whitehead, D. & Jarvis, P.G. (1981). Coniferous forests and plantations. In: (ed. Kozlowski,
T.T.). Academic Press, New York, NY USA.
Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A. & Smith, G.M. (2009). *Mixed Effects*
Models and Extensions in Ecology with R. Springer, New York, NY.

756 **Table 1:** Trait definitions and abbreviations as employed by this study.

Trait	Definition
SLA	Specific leaf area ($cm^2\,g^{-1}$)
LDMC	Leaf dry matter content (g dry mass per g fresh mass)
WD	Wood density
$M_L:M_S$	Terminal branch leaf dry mass to stem dry mass ratio (g/g)
$A_L:A_S$	Terminal branch leaf area to sapwood area ratio ($cm^2\,mm^{-2}$)
Leaf size	Average area of a single leaf based on all leaves on a terminal branch (cm^2)
k_{leaf}	Leaf conductance per unit leaf area ($mmol\,m^{-2}\,s^{-1}\,MPa^{-1}$)
k_{stem}	Leaf area-specific stem conductance of full current year terminal branch ($mmol\,cm^{-2}\,s^{-1}\,kPa^{-1}$)
K_s	Sapwood area-specific stem conductivity per branch length (k[stem] * Al:As * length, $mmol\,cm\,s^{-1}\,kPa^{-1}$)
$P50_{leaf,\,stem}$	Leaf and stem P50 measured with optical technique from Skelton et al. 2019
Growth	**Garden**: stem volume attained at 30 yrs, based on basal diameter and height of largest stem assuming cylindrical stem. **Wild**: percent of size-specific maximum basal area increment (based on rangewide 90[th] percentile BAI at a given DBH)

Figure 1: *Variance decomposition of all branch-level measurements in the common garden (a) and wild source populations (b). Grey bars beside the colored bars show the magnitude of the trait coefficient of variation. $P50_{stem}$, $P50_{leaf}$ and Growth do not have replicates within tree, so only contain between tree and between population variance components.*

763
764
Figure 2: *(a) % of the total tree-to-tree variation attributable to population in the garden versus*
attributable to sites in the wild. Traits with significant between-site variation (ANOVA p <0.05)
in the wild are shown as filled points, while traits with significant between population variation
in the garden are shown with crosses. (b) The among-population and among-site variation that
could be explained with climate variables. Y-axis shows R^2 of best climate predictor based on
AICc, filled points indicate predictors that are statistically significant for wild traits, open points
show traits with significant among-population differentiation but non-significant climate
predictors, and crosses with bold text show garden traits with significant climate predictors
(Table S2). Black line shows 1:1 relationship.

779
Figure 3: *Ratio of trait variance in the common garden compared to the wild (values <1*
indicating more trait variance in the wild). Bars show the ratio of total trait variance (median of
1000 bootstrapped variance estimates, lines show $5^{th} - 95^{th}$ percentile of ratio), gray bars
indicate significantly lower variance in the garden. Points show the ratio of the between-
population variance component alone (Ks and $P50_{leaf}$ had 0 estimated between-population
variance in the wild and thus no points).
786

787
788 **Figure 4:** *Tree level trait-trait correlations in the garden (a) and in the wild (b). Ellipses and*
789 *color show direction and strength of the correlation, points show statistically significant*
790 *relationships, and grey boxes show relationships that are significant in both the garden and the*
791 *wild.*
792

793
Figure 5: *Multiple trait correlations were significantly different in the wild versus the garden*
(based on 1000 bootstrapped comparisons of correlations), particularly k[stem] and k[leaf]
(significant correlation between leaf and stem hydraulic efficiency in whole terminal branch
tissues in the field but not the garden) and in relationships relating to leaf size (again suggesting
something interesting is going on with the drivers of plasticity in leaf expansion/development
that leads to phenotypic integration in the wild).
800