Light perception by the root is essential for gibberellin-induced primary-root

elongation in tomato

Uria Ramon[1], Amit Adiri[1], Hadar Cheriker[1], Yogev Burko[2], Ido Nir[2], and David Weiss[1]

[1]Institute of Plant Sciences and Genetics in Agriculture, The Hebrew University of

Jerusalem, P.O. Box 12, Rehovot 76100, Israel.

[2]The Institute of Plant Sciences, ARO, Volcani Center, HaMaccabbim Road 68, POB

15159, Rishon LeZion 7505101, Israel.

ORCIDs

Uria Ramon: 0000-0002-1157-1410

Amit Adiri: 0009-0002-4730-3386

Yogev Burko: 0000-0002-3607-6280

Ido Nir: 0000-0002-6223-6700

David Weiss: 0000-0002-3253-8441

Author for correspondence: David Weiss, Email- david.weiss@mail.huji.ac.il ; Tel-+972-

8-9489436

Word count:

Introduction: 790

Materials and Methods: 1404

Results: 2078

Discussion: 1046

Total: 5318

Number of figures: 7 (in color 6)

Number of supporting figures: 13

Number of supporting tables: 1

Number of datasets: 1

Summary

- The promoting effect of gibberellin (GA) on primary-root elongation was demonstrated in several species, including Arabidopsis (*Arabidopsis thaliana*). However, in some other species its effect is less clear.

- We have used the tomato (*Solanum lycopersicum*) GA-deficient mutants *gib-1* and *ga 20-oxidase* (*ga20ox1*) to study the role of GA in primary-root elongation and examined the involvement of light in this response.

- Application of GA to *gib-1* following germination, promoted primary-root elongation, but later, when roots were deeper in the dark, GA had no effect. RNA-seq analysis revealed a typical GA transcriptional response in dark-grown roots, but the output toward cell-expansion was blocked. When dark-grown roots were illuminated deep in the ground, the hormone promoted their elongation. We demonstrated that direct red light to the root is essential for GA-induced elongation. A similar light dependent GA-elongation response was found in wheat (*Triticum aestivum*) roots, but not in Arabidopsis.

- We suggest that in tomato GA promotes root elongation following germination when roots are exposed to low light intensity underground, and this contributes to seedling establishment. As root grow deeper into the soil, the insensitivity to GA may be important for continues root growth under fluctuating water availability, as water deficiency suppresses GA accumulation.

Introduction

The plant hormone gibberellin (GA) regulates numerous developmental processes, including organ elongation (Yamaguchi, 2008; Hedden & Sponsel, 2015). Bioactive GAs bind to the GIBBERELLIN-INSENSITIVE DWARF1 (GID1) receptor, leading to the interaction of GID1s with the master growth suppressor, the DELLA protein (Locascio *et al*., 2013). This facilitates the degradation of DELLA by the 26S proteasome, a process mediated by the F-box SLEEPY1 (SLY1). DELLA degradation leads to transcriptional reprogramming and activation of GA responses, including elongation (Hauvermale *et al*., 2012).

The role of GA in primary root elongation has been studied mainly in Arabidopsis (*Arabidopsis thaliana*) and its central role was demonstrated (Ubeda- Tomas *et al*., 2008, 2009; Achard *et al*., 2009; Rizza *et al*., 2017). It was shown that GA regulates both cell division in the root meristem (Shtin *et al*., 2022), and cell elongation in the root elongation zone (Ubeda-Tomas *et al*., 2008). The Arabidopsis GA deficient mutant *ga1-3* has shorter primary root, and normal elongation can be restored by GA application or loss of DELLA activity (Fu & Harberd, 2003). Using the GA perception sensor nlsGPS1, Rizza *et al*. (2017) showed that active endogenous GA accumulates in the root elongation zone. Ubeda-Tomas *et al*. (2008) showed that inhibition of GA signaling specifically in the endodermis of Arabidopsis roots is sufficient to disrupt root elongation, indicating that the endodermis is the key site for GA action in the regulation of root elongation. This conclusion was supported by Shani *et al*., (2013) that found accumulation of exogenous bioactive tagged-GAs in the endodermis of the root elongation zone. While root elongation of *ga1-3* was induced by low concentration of GA, this treatment had no effect on shoot growth (Arizumii *et al*., 2008), suggesting that roots are very sensitive to GA. Moreover, it was shown that very low concentrations of GA are required to saturate root elongation response (Tanimoto & Hirano, 2013).

The importance of GA in the regulation of root elongation in other plant species is less clear, with numerous conflicting reports (Torrey, 1976; Feldman, 1984; Phinney, 1985;

Tanimoto, 2005; Tanimoto & Hirano, 2013; Ramon *et al.*, 2020). While Whaley and
Kephart (1957) show that GA application to maize (*Zea mays*) promotes root elongation,
Svensson (1972) reported that the hormone has no effect. In Medicago (*Medicago*
truncatula), GA treatment suppressed, and the GA-biosynthesis inhibitor paclobutrazol
(PAC) increased primary root elongation (Fonouni-Farde *et al.*, 2019), suggesting that
GA inhibits Medicago root elongation. Gou *et al.* (2010) showed in *Populus* trees that
reduced GA activity promotes lateral root elongations and has no effect on primary-root
length. Butcher & Street (1960) found that elevating GA concentrations in tomato
(*Solanum lycopersicum*), progressively promote root elongation, whereas Tognoni *et al.*
(1967), showed that application of GA to tomato plants inhibits root elongation in a
concentration-dependent manner.

In a previous study we showed that PAC inhibited primary root elongation in tomato
seedlings (Ramon *et al.*, 2020). Application of low GA concentrations to the PAC-treated
plants slightly promoted root elongation, but higher concentrations, inhibited elongation,
exhibiting a bell-shaped curve of response. In contrast, in the shoots, GA strongly
promoted elongation and the effect was saturated only at very high concentrations and
exhibited a typical saturated-type curve of response. We also found that application of
GA to the wild-type (WT) M82 tomato plants, slightly inhibited primary root elongation,
suggesting that in WT tomato roots, GA levels are saturated. Genetic evidence, support
a role for GA in tomato primary-root elongation; the extreme tomato dwarf mutant *gid1^{TRI}*,
which lack the activity of all GA receptors (Illouz-Eliaze *et al.*, 2019) and the strong GA
deficient mutant *gib-1* (Koornneef *et al.*, 1990; Bensen & Zeevaart, 1990) have shorter
primary roots (Barlow *et al.*, 1991; Ramon *et al.*, 2020).

Most studies on GA and root elongation in Arabidopsis were performed with light-grown
seedlings on agar plates (Arizumii *et al.*, 2008; Ubeda-Tomas *et al.*, 2008; Shani *et al.*,
2013; Rizza *et al.*, 2017). Roots however, normally grow in the soil, in the dark and escape
from light by negative phototropism (Zhang *et al.*, 2013). Using the D-root system, Silva-
Navas *et al.* (2015) showed that roots respond differently to many stimuli under light or
dark conditions. Here we have studied how GA affect tomato root elongation, using the
gib-1 and *ga20ox1* mutants. We found that direct red light to the root is essential for GA-

induced elongation. The results suggest that penetration of light to the upper layers of the
soil drives the effect of GA on primary root elongation, but later, when the roots grow
dipper into the soil, to the dark, GA has no more effect. This phenomenon is probably
conserved in some, but not all plant species, as similar light dependent GA-induced root
elongation was found in wheat, but not in Arabidopsis.

Materials and Methods

Plant materials, seed germination, growth conditions and hormone treatments

Tomato cv. M82 (sp/sp), *gib-1* (Koornneef *et al.*, 1990; Bensen & Zeevaart, 1990) and
ga20 oxidase1 (*ga20ox1*, Shohat *et al.*, 2021) in M82 background were used this study.
Plants/seedlings were grown in vermiculite in a growth room set to a photoperiod of 12
h/12 h, day/night, light intensity of 150µmol m^{-2} s^{-1} and 25°C. In other experiments, plants
were grown in a glasshouse under natural day-length conditions, a light intensity of 700–
1000 µmol m^{-2} s^{-1} and 18–30°C. The tomato seeds were sterilized with ethanol 70% for
2 minutes, and then with sodium hypochlorite 3% for 10 minutes. The seeds washed
thoroughly by sterilized double distillated water (DDW). For germination of the strong GA
deficient mutant *gib-1*, embryos were rescued as described by Illouz-Eliaz et al., (2019)
and for germination of *ga 20-oxidase1* (*ga20ox1*, Shohat *et al.*, 2021) mutant seeds were
scarified.

Arabidopsis Columbia (Col-0) and *ga1-3* mutant in Col-0 background were also used in
this study. The seeds were sterilized with 6% sodium hypochlorite and 4% hydrochloric
acid for 2 h and then placed in the dark at 4°C for 48 h. Arabidopsis plants were grown in
a growth room, under a controlled temperature (22°C) and long photoperiod (16-h light/8-
h dark) provided by cool-white lamps (150 µmol m^{-2} s^{-1}) in pots with vermiculite. *ga1-3*
seeds were soaked with 1µM GA$_3$ (Sigma-Aldrich, St. Louis, USA) shake in the dark for
7 days and then washed thoroughly with sterilized DDW several times to remove all GA$_3$
residues.

Wheat cv. 'Abu Fashi' (GA-responsive line, Amram *et al.*, 2015) seeds were germinated
in rolled papers in flasks filled with water (Watt et al., 2013). The seeds were germinated

in the dark at 16°C. Following germination, the rolled papers were transferred to new

flasks filled with water (mock), 5 mg/L PAC or PAC and 10µM GA$_3$. Half of the papers for

each treatment were covered with aluminum foil to block the light (Supporting information

Fig. S1). The papers were placed in a growth room with a photoperiod of 16h/8h,

light/dark, a light intensity of 150µmol m^{-2} s^{-1} provided by cool-white lamps, and a

temperature of 22°C.

GA$_3$, and PAC) were applied by spraying, drenching or incorporated into the growth

medium.

Root elongation assay in Petri dishes (agar plates)

The tomato seeds/embryos were grown in Petri dishes filled with 4.4 g/L Murashige &

Skoog (MS) medium (Sigma-Aldrich), 8 g/L plant agar (Duchefa Biochemie, RV Haarlem,

The Netherlands), pH adjusted to 5.7 with NaOH. Arabidopsis seeds were placed on 1\2

MS plates (2.2 g/L) containing 5 g/L sucrose. The plates were placed vertically under

continuous white light (100µmol m^{-2} s^{-1}, provided by cool-white lamps). For dark condition

the plates were covered with aluminum foil after germination. In some experiments,

following germination, the location of the root tip was marked on the plate and after 5-7

days the plates were photographed (Canon PowerShot SX60 HS, Melville, NY, USA) and

primary root length, from the mark to the root tip, was measured using ImageJ software

(http://rsb.info.nih.gov/ij).

D-root and D-shoot growth system

To grow seedlings on agar plates (see above) and keep the root in the dark and the shoot

in the light, we have used a modified Dark-Root (D-root) growth system (Silva-Navas et

al., 2015; Supporting information Fig. S2).   We used black plastic comb that was fitted

into the plates separating the shoot and the root system and covered the lower part (root

part) of the plate with aluminum foil. To keep the shoot in the dark while exposing the

root to the light we used Dark-shoot (D-shoot) growth system. We used the same system

as for D-root, but covered the upper part (shoot part) of the plate with aluminum foil. The

plates were placed vertically under white light (100µmol m^{-2} s^{-1}, provided by cool-white

lamps).

Light treatments

The tomato *ga20ox1* mutant seeds were scarified and then placed on agar D-shoot plates and transferred into a growth chamber (Percival, Perry, Iowa, USA), at 25°C, under the specified constant light conditions: blue light (19µmol m^{-2} s^{-1}), red light (8µmol m^{-2} s^{-1}), and far-red light (17µmol m^{-2} s^{-1}). Light for all experiments was provided by light-emitting diodes (LED30-HL1). The wavelengths and light intensity were measured by LI-180 Spectrometer (LI-COR Biosciences, Lincoln, NE, USA) (Supporting information methods S3).

Molecular Cloning/Constructs and Plant Transformation.

To generate p35S:GID1a construct, GID1a (Illouz-Eliaz *et al*., 2019) cDNA was inserted into the pART7 vector downstream to the 35S promoter into the KpnI and XbaI sites. The construct was subcloned into the pART27 binary vector by NotI site and introduced into *Agrobacterium tumefaciens* strain GV3101 by electroporation. The construct was transferred into M82 cotyledons using transformation and regeneration methods described before (McCormick, 1991). Kanamycin-resistant to plants were grown and at least seven independent transgenic lines were selected and self-pollinated to generate homozygous transgenic lines.

Library Constructions and RNAseq analysis

Total RNA was extracted from root tips and leaves using a RNeasy Plant Mini Kit (Qiagen). RNA-seq libraries were prepared at the Crown Genomics institute of the Nancy and Stephen Grand Israel National Center for Personalized Medicine, Weizmann Institute of Science, Israel. Libraries were prepared using the INCPM-mRNA-seq protocol. Briefly, the polyA fraction (mRNA) was purified from 500ng of total input RNA followed by fragmentation and the generation of double-stranded cDNA. After Agencourt Ampure XP beads cleanup (Beckman Coulter), end repair, a base addition, adapter ligation, and PCR amplification steps were performed. Libraries were quantified by Qubit (Thermo Fisher Scientific, Waltham, MA 15 USA) and TapeStation (Agilent, Santa Clara, CA, USA). Sequencing was done on a NextSeq instrument (Illumina, San Diego, CA, USA) using a

HO 75 cycles kit, allocating approximately 18 million reads per sample (single read sequencing).

Sequence data analysis

Stretches of Poly-A/T and Illumina adapters were trimmed from the reads using cutadapt (Martin *et al.*, 2011); resulting reads shorter than 30bp were discarded. Reads were mapped to the Solanum lycopersicum reference genome; release SL3.00 using STAR *(*Dobin *et al.,* 2013*)* (with End To End option and out Filter Mismatch Nover Lmax set to 0.04). Annotation file was downloaded from SolGenomics, ITAG release 3.2. Expression levels for each gene were quantified using htseq-count (Andres et al., 2014), using the gtf file above (using union mode). Differentially expression analysis was performed using DESeq2 (Love *et al.*, 2014) with the betaPrior, cooksCutoff and independent Filtering parameters set to False. Raw P values were adjusted for multiple testing using the procedure of Benjamini and Hochberg (Bejamini & Hochberg, 1995). The RNA-Seq data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus (Edgar *et al.*, 2002):

https://dataview.ncbi.nlm.nih.gov/object/PRJNA1002623?reviewer=m2cghdn5fcve5npf4 qpqd8es63.

RNA Extraction, cDNA Synthesis and RT-qPCR Analysis

Total RNA was extracted with a RNeasy Plant Mini Kit (Qiagen, Hilden, Germany). For synthesis of cDNA, SuperScript II reverse transcriptase (18064014; Invitrogen, Waltham, MA, USA) and 3 mg of total RNA were used according to the manufacturer's instructions. RT-qPCR analysis was performed using an Absolute Blue qPCR SYBR Green ROX Mix kit (AB-4162/B; Thermo Fisher Scientific, Waltham, MA, USA). Reactions were performed using a Rotor-Gene 6000 Cycler (Corbett Research, Sydney, Australia). A standard curve was obtained using dilutions of the cDNA sample. The expression was quantified using the Rotor-Gene's software (Corbett Research). Three independent technical repeats were performed for each sample. Relative expression was calculated by dividing the

231 expression level of the examined gene by that of *SlACTIN*. The target-gene-to-*ACTIN*
232 ratio was then averaged.

Confocal Imaging and cell length measurement

234 Root tips were stained with Propidium iodide (PI) solution ($0.01\mu g\,ml^{-1}$). Laser beam of
235 552 nm were used for excitation of PI. The samples were placed on a glass slide with a
236 drop of water and sealed with coverslip. Fluorescence images were acquired using Leica
237 SP8 confocal microscope (Boston, MA, USA). The cells lengths were measured by
238 ImageJ software (http://rsb.info.nih.gov/ij).

Statistical analysis

All assays were conducted with three or more biological replicates and analyzed using
JMP software (SAS Institute, Cary, NC, USA). Means comparison was conducted using
analysis of variance (ANOVA) with post-hoc Tukey–Kramer honest significant difference
(HSD) test (for multiple comparisons), Student's t-test (for one comparison) and Dunnett's
test (for comparison against control) ($P < 0.05$). For assays without normal distribution
Mann-Whitney -Wilcoxon test was conducted. The experiments were performed three
times.

Accession numbers

248 Sequence data from this article can be found in the Sol Genomics Network
249 (https://solgenomics.net/) under the following accession numbers: ACTIN,
250 Solyc11g005330; GA20ox1, Solyc03g006880; GA20ox3, Solyc11g072310; GID1a,
251 Solyc01g098390.

Results

We previously found that GA levels in WT tomato roots are saturated (Ramon et al.,
2020), therefore, we have used the strong tomato GA-deficient mutant *gib-1* (Bensen &
Zeevaart, 1990) to study the effect of GA on primary-root elongation. *gib-1* exhibited very
low percentage of germination even after seed scarification, thus, we had to rescue *gib-*
1 embryos. *gib-1* and WT (M82) embryos were grown on vermiculite in pots and after

producing four expanded leaves, the length of the primary roots was measured. Similar

to *gid1^{TRI}* (Ramon *et al.*, 2020), primary roots of *gib-1* were shorter than those of WT (Fig.

1a). Microscopic analysis revealed that epidermal cells of *gib-1* primary roots are shorter

than those of WT roots (Supporting information Fig. S4). We then examined the effect of

GA application on *gib-1* primary-root elongation. Two weeks-old *gib-1* seedlings, grown

in vermiculite, were treated twice a week for one month with 10μM GA_3 and then the

length of leaves, stems and primary roots was measured. While the hormone strongly

promoted shoot growth, it had no effect on primary root elongation (Fig. 1b,c). To exclude

the possibility that the limiting factor for root elongation in *gib-1* is shortage of assimilates

(Stafen *et al.*, 2022), due to the very small leaf area and not a direct effect of GA, we

measured the elongation rate of the primary roots of *gib-1* and WT immediately after

embryo rescue, on agar plates. In this early developmental stage, differences in foliage

area between WT and *gib-1* seedlings are neglected (Supporting information Fig. S5).

We found that elongation rate of *gib-1* roots was ca. three times lower than that of WT

(Fig. 1d), suggesting a direct effect of GA on root elongation.

Since long-term treatment with GA had no effect on *gib-1* root elongation, we

hypothesized that GA is not active in the roots deeper in the growth medium. We

therefore, examined GA transcriptional responses in the roots. To this end, we grew WT

and *gib-1* for two weeks on vermiculite, and then treated *gib-1* plants with 10μM GA_3. A

previous study in Arabidopsis showed that after a short period from the GA treatment (1–

3 h), growth-related genes were only slightly affected by the hormone, but after 24 h,

these genes exhibited clear upregulation (Park et al., 2017). We therefore collected, for

RNAseq analysis, shoots and primary root tips, including the elongation zone, 24 h after

the GA treatment. We first compared non-treated *gib-1* and WT shoot and roots. In shoots

we found ca. 3000 differentially expressed genes (DEGs) between the two genotypes

and in roots approximately 1900 DEG (Fig. 2a, Supporting information Fig. S6a,

Supporting information Dataset S1). *gib-1* roots exhibited clear transcriptional responses

to GA application (Fig. 2b, Supporting information Fig. S6b, Supporting information

Dataset S1), including a typical negative feedback responses of *GA20ox1* and *GA20ox3*,

whose expression was strongly inhibited both in *gib-1* shoots and roots by GA application

(Fig. 2c,). We also analyzed the expression of cell-expansion genes (Park et al., 2017).

While many of these genes were upregulated by the GA treatment in *gib-1* shoots, in the

roots they were hardly affected (Fig. 2d). We previously speculated that in tomato, root

elongation is less responsive to GA than the shoot due to lower expression of the

dominant GA receptor *GID1a* (Ramon et al., 2020), whose activity is responsible for the

strong stem-elongation response to GA (Illouz-Eliaz *et al.*, 2019). *GID1a* exhibited similar

expression in shoots and roots (Fig. 2e). The other two receptors, *GID1b1* and *GID1b2*,

exhibited much higher expression in the roots. The F-box *SLY1* showed similar

expression in shoot and roots and the tomato *DELLA*, *PROCERA,* exhibited higher

expression in the shoot (Fig. 2e). These suggest that all canonical GA signaling

components are normally expressed in the root. Moreover, overexpression of *GID1a* in

M82 roots slightly inhibited root elongation and eliminated completely the effect of GA

(Supporting information Fig. S7), probably due to super-optimal GA activity in the root

(Ramon *et al.*, 2020). Taken together, these results suggest that GA signaling is active in

the root, but the output toward cell elongation is blocked.

Many studies that showed clear effect of GA on primary root elongation were performed

on agar plates and the roots were exposed light (Ubeda- Tomas *et al.*, 2008, 2009; Achard

et al., 2009; Rizza *et al.*, 2017; Qin *et al.*, 2022). We therefore tested if GA could promote

elongation of *gib-1* primary root on agar plates in the light. Rescued *gib-1* embryos were

grown on agar plates with or without GA_3 and the plates were placed vertically in the light

or under dark conditions for five days. Both in light and dark, GA induced hypocotyl

elongation (Fig. 3a and b). However, the hormone promoted the elongation of *gib-1*

primary roots only in the light (Fig. 3a and c). We also examined microscopically the length

of the primary-root epidermal cell above the elongation zone and found that GA increased

cell length in the light but had no effect on cell elongation in the dark (Supporting

information Fig. S8).

If roots are insensitive to GA in the dark, then under dark conditions the length of M82

roots should be similar to those of *gib-1*. We grew M82 and *gib-1* on agar plates under

light or dark conditions. M82 hypocotyls were much longer than those of *gib-1* under light

or dark conditions. However, while M82 primary roots were much longer than those of

gib-1 under light conditions, under dark conditions their length was similar (Fig. 3c,

Supporting information Fig. S9). To test if direct light to the root is required for GA-induced
elongation, we used the D-root agar plate system (Silva-Navas *et al.*, 2015; Supporting
information Fig. S2). This system allows growing seedling on agar plates with the shoots
exposed to light while the roots grow in the dark, similar to soil-growing seedlings. While
GA promoted primary root elongation in the light, it had a very mild effect on root
elongation in the D-root system (Fig. 3d). It should be noted that the D-root system does
not block completely the light to the root, but just limiting it (Silva-Navas *et al.*, 2015). We
further examined if a weaker GA-deficient mutant shows similar light-dependent GA-
induced primary-root elongation. To this end, we have used the CRISPR-derived mutant
ga20ox1 (Shohat et al., 2021). Primary roots of this mutant are shorter than those of the
WT M82 (Supporting information Fig. S10). Scarified *ga20ox1* seeds were placed on agar
plates in the light, or covered for the dark treatment, with or without 1 µM GA_3. GA
promoted primary root elongation in *ga20ox1* seedlings only under light conditions (Fig.
3e).

We hypothesized that GA induces root elongation right after germination, when light is
able to penetrate the upper layer of the soil (Tester & Morris, 1987; Mandoli *et al.*, 1990),
but later, deeper in the soil, in a complete darkness, the roots cannot respond to GA. To
test this hypothesis, we applied 10µM GA_3 immediately after placing the rescued *gib-1*
embryos on the vermiculite, or 10 days later, when the roots were deeper (approximately
2 cm long) in the vermiculite. While GA_3 application immediately after sowing promoted
primary root elongation, at the later stage the hormone had no effect (Fig. 4a). Moreover,
a single application of GA (10µM GA_3) right after germination completely restored *gib-1*
root elongation and their primary-root length, three weeks later, was similar to that of the
WT (Fig. 4b). To exclude the possibility that only very young roots can respond to GA, but
later it becomes insensitive to the hormone, we examined root elongation of *gib-1*
seedlings grown on agar plates in the light with or without GA or on plates first without
GA and only when primary roots were approximately 2.5 cm long, GA_3 was added. GA
promoted elongation when applied at early or later stages (Fig. 4c), ruling out the
possibility that GA can induce elongation only at early stages of primary-root
development, and further support the hypothesis that perception of light by the root is
essential for GA-induced elongation.

We next tested if the effect of GA on root elongation can be induced deep in the soil if we

expose the root tip to light. To this end, we removed the bottoms of the pots, and the pots

containing vermiculite were placed on LED lighting strips, exposing root tips to light from

the bottom (Supporting information Fig. S11). *gib-1* embryos were grown on the

vermiculite and two weeks later, when roots were approximately 2.5 cm long, the light

was turned on and GA$_3$ was applied. A week later, we measured the length of the primary

roots of all treatments. GA application to LED-lighted *gib-1* roots promoted their

elongation, but without bottom lighting it had no effect (Fig 4d), suggesting again that

direct light to the root is required for GA to promote their elongation.

We hypothesized that light drives GA-induced primary root elongation in the top layer of

the soil. Thus, GA-induced elongation should be activated by low light intensity. We

examined the effect of light intensity on GA-induced hypocotyl and primary-root

elongation. Rescued *gib-1* embryos were placed on agar plates with or without GA$_3$ in

dark or under 0.1, 1, 10 or 100µmol m^{-2} s^{-1} of white light (WL). The strongest effect of GA$_3$

on hypocotyls elongation was found in the dark and it was slightly decreased with the

increasing light intensity (Fig 5a and b). On the other hand, GA had no effect on primary-

root elongation in the dark, but a clear effect was found already at the lowest light intensity

(0.1 µmol m^{-2} s^{-1}) (Fig 5a and c).

To study the active light spectrum required for GA-induced root elongation, we first tested

in *ga20ox1* the effect of direct WL light to the root on agar plate. For this we have used

D-shoot plates (shoot in dark and root in light, Supporting information Fig. S2) and D-root

plates (shoot in light and root in dark), with or without 1µM GA$_3$. GA promoted primary

root elongation only when the roots were exposed to light (D-shoot) but not when the root

grew in the dark in the D-root plates (Fig. 6a). We then used the D-shoot system to study

the active light spectrum. Scarified *ga20ox1* seeds on D-shoot agar plates with or without

1µM GA$_3$, were vertically placed under blue, red or far-red light (see methods for light

spectrum). GA promoted primary root elongation only under red light (Fig. 6b), suggesting

that red-light perception by the root is essential for GA-induced primary root elongation.

There is an extensive body of information regarding the promoting effect of GA on

primary-root elongation in Arabidopsis. However, most of these studies have been

performed using light-grown roots (Arizumii *et al.*, 2008; Ubeda-Tomas *et al.*, 2008; Shani
et al., 2013; Rizza *et al.*, 2017). We therefore examined if light has similar effect on GA-
induced primary-root elongation in Arabidopsis. WT Col and the GA deficient *ga1-3*
seedlings were grown in vermiculite and after three weeks, when *ga1-3* primary roots
were approximately 2.5 cm long, the mutant was treated with GA (10µM GA_3). Different
from tomato, GA promoted the elongation of *ga1-3* primary roots deep in the soil
(Supporting information Fig. S12a). Moreover, GA promoted the elongation of *ga1-3*
primary roots also on agar plates under light and dark (full dark, D-root and D-shoot)
conditions (Supporting information Fig. S12b).

We also examined the possible interaction between light and GA in the regulation of root
elongation in wheat. To this end, the germination paper methods (Watt et al., 2013) was
used. Rolled germination papers containing wheat seeds were placed first in Erlenmeyer
flasks filled with water. After germination, the rolled papers were transferred to new flasks
filled with water (mock), 5 mg/l PAC or PAC and 10µM GA_3. Half of the papers for each
treatment were covered with aluminum foil to block the light (Supporting information Fig.
S1). The coleoptiles emerged from the paper and further developed in the light in all
treatment. After 5 days the length of the primary root was measured. When roots were
exposed to light, PAC treatment inhibited the elongation of the primary root and GA
restored elongation (Fig. 7). When roots developed in the dark, the primary roots were
longer than those developed in the light, but PAC and GA had no effect on their
elongation. Taken together, these results suggest that in wheat, similar to tomato, GA-
induced primary-root elongation depends on direct light to the root.

404 Discussion
Rapid elongation of the primary root, following germination, is essential for seedling
establishment; it anchorage the seedling and allows water and nutrient uptake. Various
factors regulate this rapid, directional elongation, including environmental factors, such
as light and gravity that direct the growth downward into the soil (negative phototropism
and gravitropism) and growth hormones. Auxin plays a central role in root growth, but
other hormones, including brassinosteroids and GA are also known to promote primary

root elongation (Garay-Arroyo *et al.*, 2012). While the role of auxin has been studied

intensively and is conserved in all tested plant species, the role of GA in different species

is less clear (Castro-Camba *et al.*, 2022). There are conflicting reports on the effect of GA

on root elongation and while numerous studies showed clear promoting effect, some

other showed no effect or even inhibitory effect (Tognoni *et al.* 1967; Fonouni-Farde *et*

al., 2019). The discrepancy between the reports could be due to application of GA to WT

plants with saturated GA levels in the root, in some studies, or application of the hormone

to GA deficient mutants, in other studies. It can also be due to application of low GA

concentrations that promote elongation in some studies and application of super-optimal

GA concentrations that inhibit root elongation, in other studies (Ramon *et al.*, 2021). The

results of this study show that it can also be due to the experimental setup; growing the

plants on plates in the light, or in the soil with roots growing in the dark. Silva-Navas *et*

al. (2015) showed that roots behave differently under light and dark conditions. Our

results suggest that in tomato and wheat the effect of GA on root elongation depends on

light. Thus, the effect of the hormone would be different if it is examined on agar plates in

the light or in the soil, in the dark.

The interaction between light and GA was demonstrated in germinating seeds, elongating

hypocotyls and stems and developing roots. According to these studies, light affects GA

metabolism, transport and responses (Arana *et al.*, 2014; Matsuo *et al.*, 2019; Lyu *et al.*,

2021; Li *et al.*, 2020; van Gelderen *et al.*, 2023). In soybean (*Glycine max*), blue light, via

cryptochrome (CRY), suppresses stem elongation by the induction of *GA 2-oxidase*

(*GA2ox*) transcription, leading to GA deactivation (Lyu *et al.*, 2021). Low red/far-red ratio,

mediated by phytochrome B (PhyB), affects GA accumulation and shoot elongation in

Pinus tabuliformis seedlings (Li *et al.*, 2020). It also affects GA responsiveness in

Arabidopsis seeds (Arana *et al.*, 2014). Increased far-red light to Arabidopsis shoots

promotes GA accumulation and its translocation to the root, where it inhibits lateral root

growth (van Gelderen *et al.*, 2023). Our results suggest that continuous red, but not blue

or far-red light, promotes GA-induced primary root elongation. This implies that activation

of PhyB is required for this GA response (Quail, 2002). Our RNAseq data show that both

PhyB1 and *PhyB2* genes are expressed in tomato roots (Supporting information Fig.

S13).

How red light and PhyB affect GA elongation responses in tomato roots is unknow. In dark-grown roots, all GA signaling components were normally expressed and GA induced or suppressed the expression of numerous genes and pathways. However, cell expansion genes were not affected by the hormone (Fig. 2). These results suggest that light affects GA-induced root-elongation response downstream of the GA signaling pathway. DELLA regulates down-stream pathways by the interactions with numerous transcription factors (TF) and their inhibition (Locascio et al., 2013). Degradation of DELLA by GA in the root probably releases these TFs, leading to transcriptional activation of many pathways independently of light. These may include the regulation of root architecture (Castro-Camba *et al.*, 2022), secondary xylem development, lignification (Wang et al., 2017) and endodermal suberin formation (Binenbaum *et al.*, 2023). Several DELLA-interacting TFs are involved in GA regulation of cell elongation (de Lucas *et al.*, 2008; Feng *et al.*, 2008; Gallego-Bartolome *et al.*, 2012; Chaiwanon & Wang, 2015). It is possible that in tomato roots the activity of these TFs depends on the activation of PhyB by red light.

Roots normally grow in the dark and escape from light by negative phototropism (Zhang *et al.*, 2013). However, in most cases, shortly after germination, in the upper layer of the soil, the young roots are exposed to low light intensity (Tester & Morris, 1987; Mandoli *et al.*, 1990). We suggest that in tomato, at this early stage the light signal interacts with the GA response pathway to accelerate root elongation. Rapid elongation at this early stage of root development contributes to seedling establishment. At later stages however, when the root elongation zone was deeper in the soil, in the dark, GA had no more effect on root elongation. It is possible that the insensitivity of root elongation to GA deeper in the soil, contributes to root growth under changing environment. Under moderate drought conditions, bioactive GA levels in tomato decrease due to the inhibition of GA biosynthesis and increased GA deactivation (Shohat *et al.*, 2021). The reduced GA levels suppress shoot growth to reduce water loss. At the same time, under the same dry conditions, root elongation is maintained (xerotromism), enabling roots to grow deeper and find new sources of water (Sharp *et al.*, 2004; Miao *et al.*, 2021). If root elongation was dependent on GA, this differential growth response between shoot and root, under moderate drought conditions, would not be possible.

Our results show that light is essential for GA-induced root elongation in tomato and
wheat, but not in Arabidopsis. If the GA insensitivity deep in the ground in the dark was
evolved to allow efficient xerotropism, it can explain the differences between these
species. Wild Arabidopsis is a winter annual with a short life cycle, originating from well-
watered habitats of the western Eurasia (Kramer, 2015), and normally not exposed to
prolonged drought. On the other hand, wild tomato originates from the arid land of the
Andean region of South America, and wheat, from the middle east, and both experience
periods of drought in their longer life cycle (Chetelat *et al.*, 2009; Chetelat *et al.*, 2019;
Harlan & Zohary, 1966). Thus, changing root to shoot ratio under water limiting conditions
may be essential for wild tomato and wheat survival, but not for wild Arabidopsis.

Acknowledgements

This research was supported by the Israel Science Foundation (617/20) to DW. We thank
Dr. Natanella Illouz-Eliaz for her valuable suggestions. We also thank Ziva Amsellem for
technical assistance.

Author Contributions: UR, IN, YB and DW designed the research; UR, HC, and AA
performed experiments; UR, AA and DW analyzed data; UR, IN, YB and DW wrote the
manuscript.

Data Availability

All data can be found in the manuscript and in the supporting information.

References

Achard P, Gusti A, Cheminant S, Alioua M, Dhondt S, Coppens F, Beemster GTS,
Genschik P. **2009**. Gibberellin signaling controls cell proliferation rate in Arabidopsis.
Current Biology **19**: 1188–1193

Amram A, Fadida-Myers A, Golan G, Nashef K, Ben-David R, Peleg Z. 2015. Effect of GA-sensitivity on wheat early vigor and yield components under deep sowing. *Frontiers in Plant Science* **6**:487

Arana MV, Sánchez-Lamas M, Strasser B, Ibarra SE, Cerdán PD, Botto JF, Sánchez RA. 2014. Functional diversity of phytochrome family in the control of light and gibberellin-mediated germination in Arabidopsis. *Plant Cell & Environment* **37**: 2014–2023.

Ariizumi T, Murase K, Sun T, Steber CM. **2008**. Proteolysis-Independent Downregulation of DELLA repression in Arabidopsis by the gibberellin receptor GIBBERELLIN INSENSITIVE DWARF1. *The Plant Cell* **20**: 2447–2459.

Barlow PW, Brain P, Parker JS. 1991. Cellular growth in roots of a gibberellin-deficient mutant of tomato (*Lycopersicon escolentum* Mill.) and its wild-type. *Journal of Expiramental Botany* **42**: 339-351.

Bensen RJ, Zeevaart JAD. 1990. Comparison of *ent*-kaurene synthetase A and B activities in cell-free extracts from young tomato fruits of wild-type and *gib-1*, *gib-2*, and *gib-3* tomato plants. *Journal of Plant Growth Regulation* **9**: 237–242.

Binenbaum J, Wulff N, Camut L, Kiradjiev K, Anfang M, Tal I, Vasuki H, Zhang Y, Sakvarelidze-Achard L, Davière JM, et al. 2023. Gibberellin and abscisic acid transporters facilitate endodermal suberin formation in Arabidopsis. *Nature Plants* **9**: 785–802

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society* **57**: 289-300.

Butcher DN, Street HE. **1960**. Effects of kinetin on the growth of excised tomato roots. *Physiologia Plantarum* **13**: 46–55.

Castro-Camba R, Sánchez C, Vidal N, Vielba JM. 2022. Plant development and crop yield: the role of gibberellins. *Plants* **11**(19), 2650

Chaiwanon J, Wang Z-Y. 2015. Spatiotemporal brassinosteroid signaling and antagonism with auxin pattern stem cell dynamics in Arabidopsis roots. *Current Biology* **25**: 1031-1042

Chetelat RT, Pertuzé RA, Faúndez L, Graham EB, Jones CM. 2009. Distribution, ecology and reproductive biology of wild tomatoes and related nightshades from the Atacama Desert region of northern Chile. *Euphytica* **167**:77–93.

Chetelat RT, Qin X, Tan M, Burkart-Waco D, Moritama Y, Huo X, Wills T, Pertuzé, R. 2019. Introgression lines of *Solanum sitiens*, a wild nightshade of the Atacama Desert, in the genome of cultivated tomato. *The Plant Journal* **100**: 836-850

de Lucas M, Daviére JM, Rodríguez-Falcón M, Pontin M, Iglesias-Pedraz JM, Lorrain S, Fankhauser C, Blázquez MA, Titarenko E, Prat S. 2008. A molecular framework for light and gibberellin control of cell elongation. *Nature* **451**: 480–484.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15-21.

Edgar R, Domrechev M, Lash AE. 2002. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. Nucleic Acids Research **30**: 207-210.

Feldman LJ. **1984**. Regulation of root development. *Annual review of plant physiology* **35**: 223–242.

Feng S, Martinez C, Gusmaroli G, Wang Y, Zhou J, Wang F, et al. 2008. Coordinated regulation of *Arabidopsis thaliana* development by light and gibberellins. *Nature* **451**: 475–479.

Fonouni-Farde C, Miassod A, Laffont C, Morin H, Bendahmane A, Diet A, Frugier F. **2019**. Gibberellins negatively regulate the development of *Medicago truncatula* root system. *Scientific Reports* **9**: 1–9.

Fu X, Harberd NP. **2003**. Auxin promotes Arabidopsis root growth by modulating gibberellin response. *Nature* **421**: 740–743.

Gallego-Bartolome J, Minguet, EG, Grau-Enguix F, Abbas M, Locascio A, Thomas SG, Alabadi D, Blazquez MA. 2012. Molecular mechanism for the interaction between gibberellin and brassinosteroid signaling pathways in Arabidopsis. *Proceedings of the National Academy of Sciences* **109**: 13446–13451.

Garay-Arroyo A, De La Paz Sánchez M, Garcıá-Ponce B, Azpeitia E, Álvarez-Buylla ER. 2012. Hormone symphony during root growth and development. *Developmental Dynamics* **241**:1867–1885

van Gelderen K, , van der Velde K, Kang C-k, Hollander J, Petropoulos O, Akyüz T, Pierik R. 2023. Gibberellin transport affects (lateral) root growth through HY5 during Far-Red light enrichment. bioRxiv doi: https://doi.org/10.1101/2023.04.21.537844

Gou J, Strauss SH, Tsai CJ, Fang K, Chen Y, Jiang X, Busov VB. 2010. Gibberellins regulate lateral root formation in populous through interactions with auxin and other hormones. *The Plant Cell* **22**: 623–639

Harlan JR, Zohary D. 1966. Distribution of wild wheats and barley. *Science*. **153**:1074–1080.

Hauvermale AL, Ariizumi T, Steber CM. 2012. Gibberellin signaling: a theme and
variations on DELLA repression. *Plant Physiology* **160**: 83-92.

Hedden P, Sponsel V. 2015. A Century of Gibberellin Research. *Journal of Plant Growth*
Regulation **34**: 740–760

Illouz-Eliaz N, Ramon U, Shohat H, Blum S, Livne S, Mendelson D, Weiss D. **2019**.
Multiple gibberellin receptors contribute to phenotypic stability under changing
environments. *The Plant Cell* **31**: 1506–1519.

Koornneef M, Bosma TDG, Hanhart CJ, Veen H Van Der, Zeevaart JAD. **1990**. The
isolation and characterization of gibberellin-deficient mutants in tomato. *Theoretical and*
Applied Genetics **80**: 852–857.

581
582 **Kramer U. 2015**. Planting molecular functions in an ecological context with
583 Arabidopsis thaliana. *eLife* **4**:e06100.
584

Li W, Liu SW, Ma JJ, Liu HM, Han FX, Li Y, Niu SH. 2020. Gibberellin signaling is
required for far-red light-induced shoot elongation in *Pinus tabuliformis* seedlings. *Plant*
Physiology **182**: 658–668.

Locascio A, Blázquez MA, Alabadí D. 2013. Genomic analysis of DELLA protein
activity. *Plant & Cell Physiology* **54**: 1229–1237.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biology* **15**:1-21.

Lyu X, Cheng Q, Qin C, Li Y, Xu X, Ji R, Mu R, Li H, Zhao T, Liu J, et al. 2021.
GmCRY1s modulate gibberellin metabolism to regulate soybean shade avoidance in
response to reduced blue light. *Molecular Plant* **14**: 298–314.

Mandoli DF, Ford GA, Waldron LJ, Nemson JA, Briggs WR. 1990. Some spectral
properties of several soil types: implications for photomorphogenesis. *Plant Cell &*
Environment **13**: 287–294.

Martin M. **2011.** Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet Journal* **17**: 1-10.

Matsuo S, Nanya K, Imanishi S, Honda I, Goto E. 2019. Effects of blue and red lights
on gibberellin metabolism in tomato seed. *The Horticulture Journal* **88**: 76–82

McCormick S. 1991. Transformation of tomato with *Agrobacterium tumefaciens*. *Plant Tissue Culture Manual* **B6**: 1-9.

Miao R, Yuan W, Wang Y, Garcia-Maquilon I, Dang X, Li Y, Zhang J, Zhu Y, Rodriguez PL, Xu W. 2021. Low ABA concentration promotes root growth and hydrotropism through relief of ABA INSENSITIVE 1-mediated inhibition of plasma membrane H+ATPase 2. *Science Advances* **7**: eabd4113

Park J, Oh D-H, Dassanayake M, Nguyen KT, Ogas J, Choi G, Sun T-p. 2017 Gibberellin signaling requires chromatin remodeler PICKLE to promote vegetative growth and phase transitions. *Plant Physiology* **173**: 1463–1474.

Phinney BO. 1985. Gibberellin A_1 dwarfism and shoot elongation in higher plants. *Biologia Plantarium* **27**: 172–179.

Qin H, Pandey BK, Li Y, Huang G, Wang J, Quan R, Zhou J, Zhou Y, Miao Y, Zhang et al. 2022. Orchestration of ethylene and gibberellin signals determines primary root elongation in rice. *The Plant Cell* **34**:1273-1288.

Quail PH. 2002. Phytochrome photosensory signalling networks. *Nature Reviews Molecular Cell Biology* **3**: 85–93.

Ramon U, Weiss D, Illouz-Eliaz N. 2020. Underground gibberellin activity: differential gibberellin response in tomato shoots and roots. *New Phytologist* **229**: 1196–200.

Rizza A, Walia A, Lanquar V, Frommer WB, Jones AM. 2017. In vivo gibberellin gradients visualized in rapidly elongating tissues. *Nature Plants* **3**: 803–813.

Svensson SB. 1972. A comparative study of the changes in root growth, induced by coumarin, auxin, ethylene, kinetin and gibberellic acid. *Physiologia Plantarum* **26**: 115–135.

Shani E, Weinstain R, Zhang Y, Castillejo C, Kaiserli E, Chory J, Tsien RY, Estelle M. 2013. Gibberellins accumulate in the elongating endodermal cells of Arabidopsis root. *Proceedings of the National Academy of Sciences of the United States of America* **110**: 4834–4839.

Sharp RE, Poroyko V, Hejlek LG, Spollen WG, Springer GK, Bohnert HJ, Nguyen HT. 2004. Root growth maintenance during water defcits: physiology to functional genomics. *Journal of Experimental Botany* **55**: 2343–2351.

Shohat H, Ceriker H, Vasuki H, Illouz-Eliaz N, Blum S, Amsellem Z, Tarkowska D, Aharoni A, Eshed Y, Weiss D. 2021. Inhibition of gibberellin accumulation by water deficiency promotes fast and long-term 'drought avoidance' responses in tomato. *New Phytologist* **232**:1985-98.

Shtin M, Dello Ioio R, Del Bianco M. 2022. It's time for a change: the role of gibberellin in root meristem development. *Frontiers in Plant Science* **13**:882517.

Silva-Navas J, Moreno-Risueno MA, Manzano C, Pallero-Baena M, Navarro-Neila
S, Téllez-Robledo B, Garcia-Mina JM, Baigorri R, Gallego FJ, del Pozo JC. 2015. D-
Root: a system for cultivating plants with the roots in darkness or under different light
conditions. *The Plant Journal* **84**: 244–255.

Stafen CF, Kleine-Vehn J, Maraschin FS. 2022. Signaling events for photomorphogenic
root development. *Trends in Plant Science* **27**: 1266-1282.

Tanimoto E. **2005**. Regulation of root growth by plant hormones-roles for auxin and
gibberellin. *Critical Reviews in Plant Sciences* **24**: 249–265.

Tanimoto E, Hirano K. 2013. Role of gibberellins in root growth. In: Eshel A, Beeckman
T. Fourth Edition, Plant Roots: The Hidden Half. CRC Press **13**: 1-14

Tester M, Morris C. 1987. The penetration of light through soil. *Plant Cell & Environment*
10: 281-286

Tognoni F, Halevy AH, Wittwer SH. **1967**. Growth of bean and tomato plants as affected
by root absorbed growth substances and atmospheric carbon dioxide. *Planta* **72**: 43–52.

Torrey JG. **1976**. Root hormones and plant growth. *Annual Review of Plant Physiology*
27: 435–459.

Ubeda-Tomás S, Swarup R, Coates J, Swarup K, Laplaze L, Beemster GTS, Hedden
P, Bhalerao R, Bennett MJ. **2008**. Root growth in Arabidopsis requires
gibberellin/DELLA signalling in the endodermis. *Nature Cell Biology* **10**: 625–628.

Wang GL, Que F, Xu ZS, Wang F, Xiong AS. 2017. Exogenous gibberellin enhances
secondary xylem development and lignification in carrot taproot. *Protoplasma* **254**: 839–
848.

Watt M, Moosavi S, Cunningham SC, Kirkegaard JA, Rebetzke GJ, Richards RA.
2013. A rapid, controlled-environment seedling root screen for wheat correlates well with
rooting depths at vegetative, but not reproductive, stages at two field sites. *Annals of*
Botany **112**: 447–455.

Whaley WG, Kephart J. **1957**. Effect of gibberellic acid on growth of maize roots. *Science*
125: 234.

Yamaguchi S. 2008. Gibberellin Metabolism and its Regulation. *Annual Review of Plant*
Biology **59**: 225-251.

Zhang KX, Xu HH, Yuan TT, Zhang L, Lu YT. 2013. Blue-light-induced PIN3 polarization
for root negative phototropic response in Arabidopsis. *The Plant Journal* **76**: 308–321

Figure Legends

Fig. 1. GA application to *gib-1* had no effect on primary-root elongation. (a). WT M82 and *gib-1* plants. (b and c) Two weeks-old *gib-1* seedlings, grown in vermiculite, were treated twice a week for one month with 10µM GA$_3$. (b) representative GA-treated and non-treated plants. The average length of the primary roots is presented (n=10 ±SE). (c) *gib-1* leaf (third leaf from the top), stem and primary root length of GA treated or non-treated plants. Data are graphically presented as whisker and box plots (n=10 ±SE). White triangles show the average values (d) Embryos, rescued from *gib-1* and WT seeds, were placed on agar plates and the length of the primary-root was measured each day for six days (n=8 ±SE). Bars in a = 1 cm in b = 2.5 cm.

Fig 2. Transcriptional response to GA in tomato root and shoot. M82 and *gib-1* seedlings were grown for two weeks on vermiculite, and then *gib-1* plants were treated with 10 µM GA$_3$ and 24 h later, shoots and primary root tips were taken (three biological replications) for RNAseq analysis. (a and b) Violine plot of differentially expressed genes (DEG) between WT M82 and *gib-1* shoot and root (a) and *gib-1* with or without GA (b). (c) Normalized read counts of *GA20ox1* and *GA20ox3* in *gib-1* roots and shoots with or without GA treatment. Values are the average of three biological repeats ±SE. (d) Heatmap of GA regulated cell expansion genes (Park et al., 2017) in *gib-1* shoots and roots. Coloring of the genes is according to the color bar (Log2 fold change). XTH-xyloglucan endotransglucosylase. (e) Heatmap showing normalized read counts of GA-signaling genes in WT M82 shoots and roots. Values are the average of three biological repeats. Coloring of the genes is according to the color bar (normalized read counts).

Fig. 3. Light is required for the promoting effect of GA on primary-root elongation. (a, b and c) Seedlings (*gib-1*) were grown on agar plates with or without 1µM GA$_3$ and the plates were placed vertically in the light or under dark conditions and after 5 days hypocotyl and primary-root length was measured. (a) Representative seedlings (bars = 1 cm). (b) Hypocotyl length. (c) Primary root length. Data in a and b are graphically presented as whisker and box plots (n=16) ± SE. White triangles show the average values. (d) Representative M82 and *gib-1* seedlings grown on agar plates with or without 1µM GA$_3$ for 5 days in light or dark (bars = 1 cm). (e) Seedlings (*gib-1*) were grown on agar plates with or without 1µM GA$_3$ and the plates were placed vertically in the light or under dark or in the light but using the D-root plates. After 5 days the length of the primary roots was measured. Values are mean of 16 seedlings ±SE. (f). *ga20ox1* seedlings were grown on agar plates with or without 1µM GA$_3$ and the plates were placed vertically in the light or under dark conditions and after 5 days primary-root length was measured. Values are mean of 15 seedlings ±SE. Stars in b, c, e and f represent significant differences between respective treatments by Student's t test (*P < 0.05; **P<0.01; ***P<0.001). ns-not significant.

Fig. 4. GA induced primary root elongation right after germination but not deeper in the growth medium. (a) Rescued *gib-1* embryos were placed on vermiculite and treated with 10μM GA$_3$ immediately, or 10 days later. One week after the GA treatment (for each treatment), the length of the primary roots was measured. Data are graphically presented as whisker and box plots (n= 16 ± SE). (b) Rescued M82 and *gib-1* embryos were placed on vermiculite and 10μM GA$_3$ was immediately added to *gib-1* and two weeks later the length of the primary roots was measured. Values are mean of 13 seedlings ±SE. (c) Rescued *gib-1* embryos were placed on agar plates without or with 1μM GA$_3$ *gib-1* or first without GA and only when primary roots were approximately 2.5 cm long, 1μM GA$_3$ was added. The seedlings were grown in the light for 6 days and then the length of the primary roots was measured. Data are graphically presented as whisker and box plots (n= 12 ± SE). (d) *gib-1* embryos were put on the vermiculite in bottomless pots that were placed (or not) on LED lighting strips and two weeks later, when roots were approximately 2.5 cm long, the light was turned on and part of the plants were treated with 10μM GA$_3$. A week later, the length of the primary roots was measured for all treatments. Values are mean of 10 seedlings ±SE. Stars represent significant differences between respective treatments or lines by Student's t test (*P < 0.05; **P<0.01; ***P<0.001; ****P<0.0001). ns- not significant. DAG- days after germination. White triangles in a and c show the average values.

Fig. 5. Low light intensity is sufficient for GA to promote primary root elongation. Rescued *gib-1* embryos were placed on agar plates with or without GA$_3$ in dark or under 0.1, 1, 10 or 100 μmol m^{-2} s^{-1} of white light. After 10 days hypocotyl and primary root length was measured. (a) Representative seedling. Hypocotyls (b) and primary root (c) length. Values are mean of 9 seedlings ±SE. Bars = 1 cm.

Fig. 6. Direct red light to the root is essential for GA-induced elongation in *ga20ox1* mutant. (a) *ga20ox1* seedlings were grown on D-root or D-shoot agar plates with or without 1μM GA$_3$ and under white light and after 5 days primary-root length was measured. (c) *ga20ox1* seedlings were grown on agar plates with or without 1μM GA$_3$ under blue, red or far-red light (for light spectrum see Supporting information Fig. S3) for 5 days and then primary-root length was measured. Values in a and b are mean of 16 seedlings ±SE. Stars represent significant differences between respective treatments by Student's t test (P<0.01). ns- not significant.

Fig. 7. Light is essential for gibberellin-induced primary-root elongation in wheat. Wheat seeds were placed between two wet germination papers and the papers were rolled and placed in Erlenmeyer flasks filled with water. Half of the papers for each treatment were

covered with aluminum foil to block the light. After germination, the rolled papers were transferred to new flasks filled with water (mock) or 5mg/l PAC or PAC and 10μM GA$_3$. After 5 days the length of the primary root was measured. (a) Representative wheat seedlings 5 days after the beginning of the hormonal treatments. (b) Primary root length. Data are graphically presented as whisker and box plots (n= 20 ± SE).. Stars in represent significant differences between respective treatments or lines by Student's t test (**$P<0.01$; ****$P<0.0001$). ns- not significant. White triangles in b show the average values. Bar = 1cm

Fig. 1. GA application to *gib-1* had no effect on primary-root elongation. (a). WT M82 and
gib-1 plants. (b and c) Two weeks-old *gib-1* seedlings, grown in vermiculite, were treated
twice a week for one month with 10µM GA₃. (b) representative GA-treated and non-treated
plants. The average length of the primary roots is presented (n=10 ±SE). (c) *gib-1* leaf
(third leaf from the top), stem and primary root length of GA treated or non-treated plants.

Data are graphically presented as whisker and box plots (n=10 ±SE). White triangles
show the average values (d) Embryos, rescued from *gib-1* and WT seeds, were placed
on agar plates and the length of the primary-root was measured each day for six days
(n=8 ±SE). Bars in a = 1 cm in b = 2.5 cm.

Fig 2. Transcriptional response to GA in tomato root and shoot. M82 and *gib-1* seedlings were grown for two weeks on vermiculite, and then *gib-1* plants were treated with 10 μM GA₃ and 24 h later, shoots and primary root tips were taken (three biological replications) for RNAseq analysis. (a and b) Violine plot of differentially expressed genes (DEG) between WT M82 and *gib-1* shoot and root (a) and *gib-1* with or without GA (b). (c) Normalized read counts of *GA20ox1* and *GA20ox3* in *gib-1* roots and shoots with or without GA treatment. Values are the average of three biological repeats ±SE. (d) Heatmap of GA regulated cell expansion genes (Park et al., 2017) in *gib-1* shoots and roots. Coloring of the genes is according to the color bar (Log2 fold change). XTH-xyloglucan endotransglucosylase. (e) Heatmap showing normalized read counts of GA-signaling genes in WT M82 shoots and roots. Values are the average of three biological repeats. Coloring of the genes is according to the color bar (normalized read counts).

Fig. 3. Light is required for the promoting effect of GA on primary-root elongation. (a, b
and c) Seedlings (*gib-1*) were grown on agar plates with or without 1µM GA₃ and the
plates were placed vertically in the light or under dark conditions and after 5 days
hypocotyl and primary-root length was measured. (a) Representative seedlings (bars = 1
cm). (b) Hypocotyl length. (c) Primary root length. Data in a and b are graphically
presented as whisker and box plots (n=16) ± SE. White triangles show the average
values. (d) Representative M82 and *gib-1* seedlings grown on agar plates with or without
1µM GA₃ for 5 days in light or dark (bars = 1 cm). (e) Seedlings (*gib-1*) were grown on
agar plates with or without 1µM GA₃ and the plates were placed vertically in the light or
under dark or in the light but using the D-root plates. After 5 days the length of the primary

roots was measured. Values are mean of 16 seedlings ±SE. (f). *ga20ox1* seedlings were
grown on agar plates with or without 1μM GA_3 and the plates were placed vertically in the
light or under dark conditions and after 5 days primary-root length was measured. Values
are mean of 15 seedlings ±SE. Stars in b, c, e and f represent significant differences
between respective treatments by Student's t test (*$P < 0.05$; **$P<0.01$; ***$P<0.001$). ns-
not significant.

Fig. 4. GA induced primary root elongation right after germination but not deeper in the growth medium. (a) Rescued *gib-1* embryos were placed on vermiculite and treated with 10µM GA_3 immediately, or 10 days later. One week after the GA treatment (for each treatment), the length of the primary roots was measured. Data are graphically presented as whisker and box plots (n= 16 ± SE). (b) Rescued M82 and *gib-1* embryos were placed on vermiculite and 10µM GA_3 was immediately added to *gib-1* and two weeks later the length of the primary roots was measured. Values are mean of 13 seedlings ±SE. (c) Rescued *gib-1* embryos were placed on agar plates without or with 1µM GA_3 *gib-1* or first without GA and only when primary roots were approximately 2.5 cm long, 1µM GA_3 was added. The seedlings were grown in the light for 6 days and then the length of the primary roots was measured. Data are graphically presented as whisker and box plots (n= 12 ± SE). (d) *gib-1* embryos were put on the vermiculite in bottomless pots that were placed (or not) on LED lighting strips and two weeks later, when roots were approximately 2.5 cm long, the light was turned on and part of the plants were treated with 10µM GA_3. A week later, the length of the primary roots was measured for all treatments. Values are mean of 10 seedlings ±SE. Stars represent significant differences between respective treatments or lines by Student's t test (*P < 0.05; **P<0.01; ***P<0.001; ****P<0.0001). ns- not significant. DAG- days after germination. White triangles in a and c show the average values.

.

(a)

(b) **(c)**

Fig. 5. Low light intensity is sufficient for GA to promote primary root elongation. Rescued *gib-1* embryos were placed on agar plates with or without GA_3 in dark or under 0.1, 1, 10 or 100 μmol m^{-2} s^{-1} of white light. After 10 days hypocotyl and primary root length was measured. (a) Representative seedling. Hypocotyls (b) and primary root (c) length. Values are mean of 9 seedlings ±SE. Bars = 1 cm.

Fig. 6. Direct red light to the root is essential for GA-induced elongation in *ga20ox1*
mutant. (a) *ga20ox1* seedlings were grown on D-root or D-shoot agar plates with or
without 1µM GA₃ and under white light and after 5 days primary-root length was
measured. (c) *ga20ox1* seedlings were grown on agar plates with or without 1µM GA₃
under blue, red or far-red light (for light spectrum see Supporting information Fig. S3) for
5 days and then primary-root length was measured. Values in a and b are mean of 16
seedlings ±SE. Stars represent significant differences between respective treatments by
Student's t test (P<0.01). ns- not significant.

(a)

(b)

Fig. 7. Light is essential for gibberellin-induced primary-root elongation in wheat. Wheat
seeds were placed between two wet germination papers and the papers were rolled and
placed in Erlenmeyer flasks filled with water. Half of the papers for each treatment were
covered with aluminum foil to block the light. After germination, the rolled papers were
transferred to new flasks filled with water (mock) or 5mg/l PAC or PAC and 10µM GA$_3$.
After 5 days the length of the primary root was measured. (a) Representative wheat
seedlings 5 days after the beginning of the hormonal treatments. (b) Primary root length.
Data are graphically presented as whisker and box plots (n= 20 ± SE). Stars in represent
significant differences between respective treatments or lines by Student's t test
(**P<0.01; ****P<0.0001). ns- not significant. White triangles in b show the average
values. Bar = 1cm

New Phytologist **Supporting Information**

Article title: Light perception by the root is essential for gibberellin-induced primary-root elongation in tomato

Authors: Uria Ramon, Amit Adiri, Hadar Cheriker, Yogev Borko, Ido Nir and David Weiss

Article acceptance date: Click here to enter a date.

The following Supporting Information is available for this article:

Fig. S1. The paper germination system for wheat root elongation assay (light and dark).

Fig. S2. D-root and D-shoot growth systems.

Fig. S3. The light spectrum of the different light treatment.

Fig. S4. Root epidermal cell length of WT M82 and *gib-1*.

Fig. S5. WT M82 and *gib-1* seedlings

Fig. S6. Differentially expressed genes (DEG).

Fig. S7. Overexpression of GID1a. root elongation.

Fig. S8. GA increases root cell length in the light but not in the dark.

Fig. S9. Primary root length of WT M82 and *gib-1* under light and dark conditions.

Fig. S10. Primary-root length of M82 and *ga20ox1.*

Fig. S11. The bottom lighting growth system.

Fig. S12. GA promoted primary root elongation in Arabidopsis under light and dark conditions.

Fig. S13. Normalized reads counts of the tomato photoreceptors in shoot and root.

Table S1 Primers used in this study

Dataset S1. Differentially expressed genes (DEG) between M82 and *gib-1* shoots and roots and between *gib-1* shoot and roots, treated or not treated with GA (see separate file).

Fig. S1. The paper germination system for wheat-root elongation assay.
Seeds were placed between two rolled wet paper, in the outer circle. For dark
conditions (to the roots), the rolled papers were covered with aluminium foil.

Fig. S2. D-root and D-shoot growth systems.

Fig. S3. The light spectrum of the different light treatment. White light (white line), blue light (blue line), red light (red line) and far red light (dark red line).

Fig. S4. Cell length of WT M82 and *gib-1* primary roots. Microscopic analysis of epidermal cells of WT M82 and *gib-1* primary roots.

Fig. S5. WT M82 and *gib-1* seedlings. Bar = 1 cm

a

b

Fig. S6. Differentially expressed genes (DEG). (a) Venn diagram represents DEG between WT M82 and *gib-1* shoots and roots. (b) Venn diagram represents DEG between *gib-1* untreated and GA treated shoots and roots

Fig. S7. Overexpression of GID1a. (a) GID1a expression in WT M82 and transgenic 35S:GID1a roots. Values are mean of three biological replicates ± SE. (b) Primary root length in WT M82 and transgenic plants overexpressing GID1a treated with 5mg/l paclobutrazol (PAC) or PAC with 10μM GA_3. Data are graphically presented as whisker and box plots (n= 12 ± SE). White triangles show the average values. Stars represent significant differences between respective treatments or lines by Student's t test (*$P < 0.05$; **$P<0.01$).

Fig. S8. GA increases root cell length in the light but not in the dark. *gib-1* seedlings were grown on agar plates with or without 1μM GA$_3$. The plates were placed vertically in the light or under dark conditions for 5 days. The length of the primary-root epidermal cell above the elongation zone was examined microscopically. Average cell length is presented (n=30 ± SE). Bar = 100 μm

Fig. S9. Primary root length of WT M82 and *gib-1* under light and dark conditions. WT M82 and *gib-1* seedlings were grown on agar plates. The plates were placed vertically in the light or under dark conditions for 7 days and then the length of the primary roots was measured. Values are mean of 12 seedlings ± SE. Stars represent significant differences between respective treatments by Student's t test (P<0.01). ns- not significant.

Fig. S10. *ga20ox1* has short roots. M82 and *ga20ox1* seedlings were grown in vermiculite for 10 days and then primary-root length was measured. The average length is presented (n=7 ±SE). Star represents significant difference by Student's t test (P<0.05).

Fig. S11. The bottom-lighting growth system. Left- LED lighting strips fixed to the growing tray. Middle- Try with LED lighting strips and pots without their bottom. Right- Try with LED lighting and pots without their bottom, filled with vermiculite and *gib-1* seedlings.

Fig. S12. GA promoted primary root elongation in Arabidopsis under light and dark conditions. (a) WT Col and ga1-3 seedlings were grown in vermiculite and after three weeks, when *ga1-3* primary roots were approximately 2.5 cm long, the mutant was treated with 10μM GA$_3$. Values are mean of 25 seedlings ± SE. (b) Primary root length of ga1-3 grown for one week on agar plates containing 0.5% sucrose, with or without 1μM GA$_3$, under light or dark conditions or in the D-root or D-shoot growth systems in the light. Data are graphically presented as whisker and box plots (n= 15 ± SE). Stars represent significant differences between respective treatments by Student's t test (**P<0.01, ***P<0.005, ****P<0.001). White triangles show the average values.

Fig. S13. Normalized reads count of the different photoreceptors (with theie accession numbers) in tomato M82 shoot and root. The data were taken from the RNAseq analysis (see Material and Methods).

Tabl2 S1. Primers used in this study:

Gene	Used for	Sequence (5'-3')
ACTIN	RT-qPCR	Forward- GTCCTCTTCCAGCCATCCAT Reverse- ACCACTGAGCACAATGTTACCG
GID1a	RT-qPCR	Forward- ATGGCAAGAAATAATGAAGCTGT Reverse- CGAAGCTACTTATCTCATCCATG