The Lifespan Trajectories of Brain Activities Related to Cognitive Control

Zhenghan Li[1,2,3,4], Isaac T. Petersen[5], Lingxiao Wang[6,7,3], Joaquim Radua[8,9,10], Guo-
chun Yang[5,11,4*], Xun Liu[4,12]

[1] Institute of Brain Science and Department of Physiology, School of Basic Medical
Sciences, Hangzhou Normal University, Hangzhou, China

[2] Zhejiang Philosophy and Social Science Laboratory for Research in Early Develop-
ment and Childcare, Hangzhou Normal University, Hangzhou, China

[3] Zhejiang Key Laboratory for Research in Assessment of Cognitive Impairments,
Hangzhou, China

[4] Key Laboratory of Behavioral Science, Institute of Psychology, Chinese Academy of
Sciences, Beijing, China

[5] Department of Psychological and Brain Sciences, University of Iowa, Iowa City, IA,
USA

[6] Centre for Cognition and Brain disorders, The Affiliated Hospital of Hangzhou Nor-
mal University, Hangzhou, Zhejiang Province, China

[7] Institute of Psychological Science, Hangzhou Normal University, Hangzhou,
Zhejiang Province, China

[8] Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), CIBERSAM,
University of Barcelona, Barcelona, Spain

[9] Institute of Psychiatry, Psychology and Neuroscience, King's College London, Lon-
don, United Kingdom

[10] Department of Clinical Neuroscience, Karolinska Institute, Stockholm, Sweden

[11] Cognitive Control Collaborative, University of Iowa, Iowa City, IA, USA

[12] Department of Psychology, University of Chinese Academy of Sciences, Beijing,
China

*Correspondence: guochun-yang@uiowa.edu

Abstract

Cognitive control plays a pivotal role in guiding human goal-directed behavior, and revealing its lifespan trajectory is crucial for optimizing cognitive functioning at different ages, especially for stages of rapid development and decline. While existing studies have shed light on the inverted U-shaped trajectory of cognitive control function both behaviorally and anatomically, little is known about the corresponding changes in functional brain activation with age. To bridge this gap, we conducted a comprehensive meta-analysis of 129 neuroimaging studies using conflict tasks, encompassing 3,388 participants whose age spanned from 5 to 85 years old. We applied the seed-based d mapping (SDM), generalized additive model (GAM) and model comparison approaches to investigate age-related changes of brain activity, chart the lifespan trajectories and pinpoint peaks of cognitive control brain activity. The present study have three major findings: 1) The inverted U-shaped lifespan trajectory is the predominant pattern; 2) Cognitive control related brain regions exhibit heterogeneous lifespan trajectories: the frontoparietal control network (such as the inferior frontal gyrus and inferior parietal lobule) follows inverted U-shaped trajectories, peaking between 24 and 41 years, while the dorsal attention network (such as the frontal eye field and superior parietal lobule) demonstrates flatter trajectories with age; 3) Both the youth and the elderly show weaker brain activities and greater left laterality than young adults. These results collectively reveal the lifespan trajectories of cognitive control, highlighting heterogeneous fluctuations in brain networks with age.

Keywords

Cognitive control; Lifespan; Brain activity; Neuroimaging meta-analysis; Heterogeneity

Introduction

The cognitive abilities of human beings dynamically change throughout the entire lifespan, experiencing rapid development in the early stages, and gradual decline in the later period of life. As one of the most fundamental cognitive functions, cognitive control is deeply engaged in various domains of high-level capabilities that human greatly outperform other species, such as decision making, planning and problem solving[1]. Cognitive control refers to the cognitive processes that enable individuals to manage and regulate their attention, thoughts, and actions, which plays a vital role in goal-directed behavior, allowing us to focus on the target and ignore distractors[2,3]. For instance, cognitive control enables us to concentrate on reading in a library despite the presence of people chatting nearby.

Cognitive control provides fundamental support for normal human behaviors. The young adults typically maintain an optimal mature level of cognitive control[4]. On the other hand, the youth (including children and adolescents) and elderly individuals may struggle with behavioral problems because of their suboptimal cognitive control system[5]. For instance, school-age children may suffer from attentional deficits that are related to a noticeable deficit of control system[6]. The adolescents are characterized by more risk-taking or even susceptive to addiction, which is related to the underdeveloped cognitive control system[7,8] or insufficient top-down regulation on the limbic emotional system[4,9]. Elders may have difficulty focusing on target information as a result of decay

in cognitive control[10] or a less efficient engagement of neural resources[11]. While the

state-of-art progress of cognitive/behavioral changes has been well-documented and

shaped how the diagnosis of developmental/ageing related disorders[12], the change of

related neural system has been under-investigated. Understanding how the neural un-

derpinning of cognitive control changes over the lifespan can yield valuable insights

into the developmental and aging mechanisms of the human brain. This knowledge may

lead to the identification of potential biomarkers for age-related disorders before any

behavioral symptoms become apparent[13], and may assist in customizing cognitive train-

ing strategies based on brain activities[14].

Inverted U-shaped Lifespan Trajectory of Cognitive Control: Behavioral and

Anatomical Evidence

Researchers generally believe that the cognitive control ability follows an inverted

U-shaped trajectory across the lifespan[5,15,16]. Specifically, infants initially possess lim-

ited cognitive control that gradually develops as the brain anatomy matures. These func-

tions continue to progress, reaching their peaks during young adulthood and maintain-

ing relative stability until middle age, after which they begin to decline in elderly age[17].

This inverted U-shaped trajectory has been generally supported by the behavioral evi-

dence. The Eriksen Flanker task (a task requiring participants to respond to central stim-

uli while ignoring flanking distractions), with its validity of detecting cognitive control

across the lifespan[18], has been widely utilized, and results suggest a clear U-shape tra-

jectory of conflict cost (measured by worsened behavioral performance, e.g., reaction

time, in incongruent compared to congruent conditions) with age[19-21]. Similar results

have been observed in other conflict tasks, such as color-word Stroop (a task requiring

participants to name the ink color of a word that is incongruent with the word's semantic

meaning)[19]. However, it is notable that a recent massive cross-sectional study[21] found

a weaker decline in performance on the Stroop and Simon (a task requiring participants

to respond based on one attribute of a stimulus while ignoring its conflicting spatial

location) tasks compared to the Flanker task, highlighting the possible variations in the

lifespan trajectory across different conflict tasks.

Recent large-cohort studies have also found that gray and white matter volumes

across all brain regions exhibit overall inverted U-shaped trajectories with age, with the

gray matter volume peaking at early adolescence and the white matter volume peaking

at young adulthood[22,23]. Gray matter volume exhibits a rapid increase during the early

childhood, reflecting the growth and branching of dendrites and synaptogenesis; how-

ever, following its peak at early adolescent, there is a gradual volume thinning and

density reduction due to synaptic pruning, where unused neural connections are elimi-

nated, making brain networks more efficient[24]. White matter, on the other hand, under-

goes continuous increase throughout adolescence until young adulthood due to mye-

lination, a process that can enhance the efficiency of neural transmission[25]. During late

adulthood, normal ageing yields protracted decline of brain structure, with the volumes

of both gray matter and white matter reduced[16]. These potentially provide the anatom-

ical foundations for the lifespan trajectory of cognitive control, among other func-

tions[5,26]. Consistently, the gray matter volumes of the frontal and parietal regions, which

are essential in cognitive control tasks[27], have also been found to increase during early

childhood and atrophy in early elderly age[28].

However, it remains largely unknown how brain activities related to cognitive con-

trol change over the lifespan.

Age Matters in Brain Activities Related to Cognitive Control

Previous research has primarily focused on brain activities in either youths or elderly

adults, rather than examining changes across the entire lifespan. With conflict para-

digms, children and adolescents are often found to have lower brain activity than young

adults in frontoparietal regions (refs.[29-34], but see Bunge et al.[35]). For example, a study

utilizing Flanker task revealed that children aged 8−12 years had reduced activation of

dorsolateral prefrontal regions compared to young adults, suggesting an immature cog-

nitive control system[33]. However, brain activity differences in elderly adults as com-

pared to young adults during cognitive control tasks have been less consistently re-

ported[16]. Some studies have found that elderly adults have lower neural activity in fron-

toparietal regions than young adults[36-38], possibly because elderly adults may be unable

to engage in an equal level of control-related activity due to functional decline. On the

other hand, other studies have found that elderly adults may exhibit greater brain activ-

ity in frontoparietal regions than young adults[39,40], possibly because they have recruited

additional brain regions to compensate for their decreased efficiency in utilizing control

resources. Adding to the debate, it has been proposed that the cognitive control function

in elderly adults might not decline at all[41,42]. These conflicting findings underscore the

complexity of understanding age-related differences in cognitive control.

Few studies have directly tested the change of brain activities related to cognitive

control across the lifespan. One existing study observed a positive association between

the activation of the bilateral prefrontal cortex and different age groups[43]. Given the

relatively small sample size (N = 30), the reliability of these findings is somewhat lim-

ited. As a result, it is difficult to draw a clear conclusion about how brain activities

related to cognitive control change across the entire age range.

In addition to understanding how the activations of individual brain regions change

with age, previous research has also suggested that the overall distribution of brain ac-

tivity changes with age. Two notable phenomena have emerged as significant findings

in the field of aging research: compensatory recruitment of additional brain regions and

a decrease in hemispheric asymmetry in brain activities. The concept of compensatory

recruitment posits that, while aging may lead to reduced functionality in specific brain

areas, older adults can mitigate this decline by engaging alternative neural circuits or

regions. This compensatory mechanism serves as a form of neural adaptation to the

challenges of aging, allowing for the preservation of cognitive function despite age-

related physiological changes[11]. Similarly, some developmental studies have also re-

ported compensatory brain recruitment in young children see review[44]. In parallel, pre-

vious studies have reported a decrease in hemispheric asymmetry in older populations.

This phenomenon, often referred to as the hemispheric asymmetry reduction in older

adults (HAROLD) model, proposes that older adults exhibit a reduced laterality of brain

activation compared to younger individuals[45]. This phenomenon might reflect a form

of neural compensation, where both hemispheres are utilized to perform tasks that once

predominantly engaged only one hemisphere[46].

The Hierarchical Trajectories across Different Brain Regions

Neurodevelopment follows a hierarchical trajectory, characterized by distinct rates of

development and decline across various brain networks. Research has revealed a clear

developmental hierarchy, where higher-order cortical regions tend to undergo more

complex lifespan trajectories compared with the lower-order subcortical regions[47]. Ad-

ditionally, it has been observed that higher-order associative regions typically reach the

peak later than the lower-order primary areas[48-50]. Notably, the frontal region is among

the last regions to reach their peaks and one of the earliest to show declination[5,15,51].

Importantly, the brain regions associated with cognitive control have been pro-

posed to exhibit a hierarchical organization. At the apex of this hierarchy is the fron-

toparietal control network (FPCN), which is crucial for supporting multiple higher-or-

der functions[52,53]. During conflict resolution, FPCN is closely associated with central

executive control processes, which involve maintaining goals and modulating sensory

inputs in a top-down manner[54]. The dorsal attention network (DAN), responsible for

directing selective attention allocation[55], is generally thought to be modulated by the

FPCN[53,54]. Another network pertinent to conflict processing is the cingulo-opercular

network (CON), which is primarily involved in detecting salient stimuli[56] and monitor-

ing conflicts[27]. Research suggests that the CON and DAN occupy lower positions in

the hierarchy compared to the FPCN[57], with some evidence indicating that the CON

may be situated at a higher level than the DAN[58]. Previous investigations into the mor-

phological and connectivity-based characteristics of these networks have demonstrated

that they follow distinct lifespan trajectories[59,60]. However, the specific lifespan trajec-

tories of engagement of these networks during cognitive control tasks remain to be fully

elucidated.

The Necessity of Conducting Meta-analyses

One direct way to test the trajectory of brain activation related to cognitive control is to

conduct a large cohort of neuroimaging study with participants covering a wide age

range. To the best of our knowledge, such studies have not yet been conducted. An

alternative approach is to utilize meta-analyses to combine the results of existing studies

targeting different age groups. Compared to the large cohort studies, meta-analyses are

more readily available and resource-saving. In addition, meta-analyses show several

advantages over individual studies. By combining results from various studies, meta-

analyses can increase the statistical power and generalizability of the findings and thus

result in more precise conclusions. Meta-analyses can also reduce the heterogeneity and

bias that may exist in individual studies because of different methods, populations, or

confounding variables[61]. Importantly, meta-analyses can reveal patterns or trends that

may not be immediately evident in individual studies, such as nonlinear effects or in-

teractions[61]. There have been several neuroimaging meta-analyses that attempted to ex-

amine the changes of cognitive control brain activity with age[37,62,63], but a few limita-

tions prevent them from appropriately answering questions about the lifespan trajectory

of cognitive control functions. These limitations include not covering the full range of

age across the lifespan[37,62], or having too few studies in some age ranges (e.g., elderly

adults)[63]. In addition, these studies have primarily focused on the spatial convergence

and/or diversity of reported coordinates across different age ranges, which provides

limited insights into lifespan trajectory of activity strength. Although a few studies have

attempted parametric meta-regression to examine the age-related differences in cogni-

tive control[62,64], they have been constrained by utilizing linear models that may over-

look non-linear trajectory patterns, such as the inverted U-shaped trend.

The Present Study

The goal of this study is to reveal the lifespan trajectory of brain activities responsible

for cognitive control. Instead of conducting a comprehensive meta-analysis encompass-

ing various aspects of cognitive control such as working memory, cognitive flexibility,

and conflict processing, we choose to focus on conflict processing as an indicator of

cognitive control for several reasons. First, conflict processing reflects the ability to

maintain a goal in mind while avoiding distractions, which is a fundamental aspect of

cognitive control[2]. Second, the mechanisms underlying conflict processing in young

adults are relatively well-known, with the frontoparietal and cingulo-opercular net-

works repeatedly reported as the most involved brain regions[27,65], providing a baseline

reference for our study. Third, conflict tasks with neuroimaging data have been widely

applied to both younger and elderly groups, making a systematic meta-analysis feasible.

Lastly, different conflict tasks share components on key aspects of cognitive control,

such as conflict monitoring[66] and inhibitory control[67]. Consequently, different conflict

tasks can be measured using a clearly defined congruency effect (i.e., the contrast be-

tween incongruent and congruent/neutral conditions), which enables us to conduct ef-

fect-size based meta-analyses. Including other sub-processes of cognitive control may

introduce heterogeneity and make the effect sizes incomparable.

Based on previous research, we hypothesized that brain activities related to cog-

nitive control increase rapidly from childhood to adolescence, reach a peak in young

adulthood, and then gradually decline with increasing age, following an inverted U-

shaped trajectory. Additionally, we hypothesized that different brain networks may

show some heterogeneity in their lifespan trajectories.

Results

Sample Description

A total of 3,611 articles were identified including 3,484 articles searched from the database, 111 articles adopted from a previous study[27], and 16 articles manually searched from the references of crucial articles. After excluding duplicates and applying exclusion criteria, 119 articles including 129 studies with $N = 3,388$ participants and 1,579 brain activation foci, were included in this meta-analytic study (supplementary Table S1). Of the studies included, 125 were published in peer-reviewed journals, and 4 were master's theses. All studies were published or completed between 1994 and 2022. The average age of participants ranged from 8 to 74 years, with the individual age ranged from 5 to 85 years. Included studies are written in English (115 articles) and Chinese (4 articles). All included studies reported the task type used, including 73 studies utilizing Stroop-like task (57%), 26 studies utilizing Simon task (20%), 25 studies utilizing Flanker task (19%), 2 studies utilizing a combination of Simon and Flanker tasks (2%), 1 study utilizing a combination of Simon and Stroop tasks (1%), and 2 studies utilizing multi-source interference task (2%). Besides, the contrasts conducted to reveal brain activations were also reported, with 98 studies (76%) resulting from the contrast of Incongruent trials > Congruent trials, 25 studies (19%) resulting from the contrast of Incongruent trials > Neutral trials, and 6 studies (5%) resulting from the union contrast

of Incongruent trials > Congruent trials and Incongruent trials > Neutral trials. Regard-

ing the handedness of participants in the included studies, 89 studies (69%) included

right-handed participants only, 5 studies (4%) included both left and right-handed par-

ticipants, while 35 studies (27%) did not report this information. Furthermore, 76 stud-

ies (59%) included only correct response trials, 2 studies (2%) included both correct

and incorrect response trials, while 51 studies (39%) did not report this information. To

eliminate the influence of these confounding factors, we included them as covariates in

the modeling analyses. A detailed description of these features for each study is availa-

ble in the Supplemental Table S1.

Regions Related to Cognitive Control Identified by both ALE and SDM

To enhance the replicability and robustness of our finding and enable direct comparison

with a previous study[27], we conducted the mean analysis across all studies with the ALE

and SDM approaches separately. First, we performed the single dataset analysis with

the GingerALE software to explore the brain regions consistently reported in all the

included studies. Results showed significant activation in the frontoparietal regions,

including the left dorsolateral prefrontal cortex, bilateral frontal eye field, right inferior

frontal gyrus and bilateral superior parietal lobule; the cingulo-opercular regions, in-

cluding the pre-supplementary motor area and bilateral insula; and other regions, in-

cluding left inferior temporal gyri (Figure 1A and supplementary Table S2). Second,

we calculated the average brain activation based on the effect sizes reported in all stud-

ies using the SDM-PSI software. Similar to the ALE results, we found significant acti-

vation in the frontoparietal regions (left inferior parietal lobule, right inferior frontal

gyrus, and right middle frontal gyrus), the cingulo-opercular regions (left anterior cin-

gulate cortex), and other regions including bilateral inferior temporal gyrus, right cau-

date nucleus, right cerebellum, and left anterior thalamic projections (Figure 1B and

supplementary Table S3). Despite their methodological distinction, that is, the SDM

approach additionally incorporates the statistical values of brain activation compared

to the ALE, these two approaches yielded similar results (Figure 1C), both replicating

a previous study including only the young adult studies[27]. The count of voxels revealed

that the overlapped area (3,496 voxels) accounted for 96.3% of the regions from the

ALE analysis (3,635 voxels), and 15.0% of the regions from the SDM analysis (23,240

voxels). This result also suggested that SDM analysis showed a greater sensitivity than

the ALE analysis under the same threshold (voxel-wise FWE corrected $p < 0.001$, clus-

ter ≥ 10 voxels).

Figure 1. Overview of significant clusters across all studies regardless of age in the ALE meta-analysis
(A), the SDM meta-analysis (B), and their overlap (C). ALE = activation likelihood estimation; SDM =
seed-based *d* mapping.

Trajectories of Cognitive Control Regions Identified in the Mean Analysis

Having identified nine brain regions in the mean analysis using SDM-PSI, we pro-

ceeded to examine how the activation levels of these regions change with age. To this

end, we extracted brain activity data from all studies for each identified region. Before

performing the meta-regression, we verified the effectiveness of using mean age as a

predictor with a simulation approach (see Supplementary Note S1). We then subjected

the extracted data to separate GAM analyses, factoring in the confounding covariates.

We did not conduct simplified model fitting (e.g., quadratic model) as we did not have

pre-defined hypotheses about the trajectory shapes for these regions. The GAM anal-

yses revealed significant age-related changes in the activation levels of 5 out of the 9

regions, ps < .05, false discovery rate (FDR) corrected (Figure 2, panels A, B, C, E and

G). The other four regions showed no significant age-related changes. Notably, none of

the clusters showed significant publication bias based on Egger's test (ps > 0.86), and

they all showed low between-study heterogeneity (τs < 0.13, Qs < 12.11, I^2s < 25%).

This indicates that the observed results are not likely influenced by biased reporting or

substantial variability in the included studies.

To test whether the visually inverted U-shaped trajectories reach significance, we

applied the two-line test approach proposed by Simonsohn[68] (see supplementary Note

S2). Results suggest that all five regions (panels A, B, C, E, and G in Figure 2) involve

an increase in activity from childhood to young adulthood (approximately up to 30

years of age). Three of them (panels C, E and G) showed consistent decrease in the later

stages of age, and the other two (panels A and B) showed no significant decrease (sup-

plementary Note S2).

314
Figure 2. Lifespan trajectories within regions identified in the mean analysis. Panels A-I correspond to
left anterior cingulate cortex, left inferior parietal lobule, right inferior frontal gyrus, right inferior tem-
poral gyrus, left inferior temporal gyrus, right middle frontal gyrus, right caudate nucleus, right cerebel-
lum and left anterior thalamic projections, respectively. Scattered plots are the effect sizes as a function
of age, with curves fitted by GAM. The sizes of the scattered dots show the square root of model weights
(1/variance) for each study. Shaded areas around the curves represent standard errors. Dashed lines indi-
cate peak ages. Panels A and B do not show the peak age due to an insignificant decrease at the later
stage (see supplementary Note S2).

A notable limitation of the model fitting using regions identified from the mean

analysis is the considerable sizes of some regions, with the maximum region (i.e., the

pre-supplementary motor area) encompassing over 9,000 voxels. This vastness means

that the average trajectory of voxels within these regions could obscure the distinct

trajectories specific to different subregions, which are crucial in understanding the hi-

erarchical patterns of brain development and decline. Recognizing this issue, our next

step was to delve deeper and examine the brain trajectory with a more granular ap-

proach.

Detecting Different Trajectories of Brain Activities

The present study aimed to investigate the lifespan trajectories of cognitive control

brain activities. To explore various possibilities, we conducted several contrast analyses

(Figure 7, also see Methods for details). These analyses did not reveal any regions that

exhibited significantly higher or lower brain activity in the youth compared to others

(the combination of young and older adults) (Table 1). Similarly, we did not observe

any regions with higher or lower brain activity in older adults compared to others (the

combination of the youth and young adults) (Table 1). These results excluded the pos-

sibilities of increase/decrease-then-stable and stable-then-increase/decrease trajectories

(Figure 7, panels D, F, G, and I). In addition, we failed to observe any region showing

lower activity in young adults than others (the combination of the youth and older

adults), and thereby ruled out the possibility of the upright U-shaped trajectory (Table

1, Figure 7E).

Table 1. Brain areas activated in the contrast of one age group versus others (voxel-
wise FWE-corrected, *p* < 0.001, with minimum cluster size ≥ 10 voxels) with the SDM-
PSI.

Order	# Voxels	*Z*	*p*	L/R	MNI coordinate			Anatomical location	BA
					x	y	z		
Young adults > Others									
1	1165	4.713	< 0.001	R	52	16	4	inferior frontal gyrus	48
2	607	4.895	< 0.001	L	−46	16	26	inferior frontal gyrus	48
3	569	4.289	< 0.001	R	54	−48	42	inferior parietal lobule	40
4	215	3.405	< 0.001	L	−8	0	58	supplementary motor area	6
5	202	3.830	< 0.001	L	−36	−54	42	inferior parietal lobule	40
6	106	3.245	< 0.001	R	10	4	6	caudate nucleus	/
7	76	3.231	< 0.001	L	−38	24	0	insula	47
8	33	3.119	< 0.001	L	−36	12	−6	insula	48
9	10	2.794	< 0.001	R	4	−16	44	anterior cingulate cortex	23
Others > Young adults									
None									
The youth > Others									
None									
Others > The youth									
None									
The elderly > Others									
None									
Others > The elderly									
None									

Note. The orders in the table correspond to the numbers of each plot in Figure 3. ALE
= activation likelihood estimation; MNI = Montreal Neurological Institute; BA = Brod-
mann area; L = left; R = right.

However, we identified greater activity in young adults compared to others in the

frontoparietal regions, including bilateral inferior frontal gyrus and bilateral inferior

parietal lobule; the cingulo-opercular regions, including left supplementary motor area,

left insula, and right anterior cingulate cortex; and a subcortical region right caudate

nucleus (Figure 3 and Table 1). Notably, none of the clusters showed significant publi-

cation bias based on Egger's test ($ps > 0.79$), and they all showed low between-study

heterogeneity ($\tau s < 0.17$, $Qs < 12.21$, $I^2s < 25\%$). This indicates that the observed results

are not likely influenced by biased reporting or substantial variability in the included

studies.

Fitting the Lifespan Trajectories with the Generalized Additive Model (GAM)

Results of the GAM applied to the regions identified from the contrast between young

adults and others are shown in Figure 3 and supplementary Table S4. For each region,

the GAM could fit the data significantly with a smooth curve, $Fs > 4.7$, $ps < 0.002$,

FDR corrected, with degrees of freedom varying from 3.2 to 7.0. Peak ages of the in-

verted U-shaped trajectories were between 24.2 and 38.5 years.

Model Simplification of the Lifespan Trajectories

Considering the GAM may overfit the data, we fitted the results with simpler models

commonly adopted in lifespan developmental literature[69-71], including the quadratic,

cubic, square root and quadratic logarithmic models, and estimated the goodness of fit

by comparing their Akaike information criterion (AIC) (see section "Model Simplifi-

cation and Model Comparison" in Methods). Weights of each model based on AIC

values for each region are presented in supplementary Table S5. Results showed that

the quadratic model provided the best fit for capturing the age-related changes in left

inferior temporal gyrus, while the square root model demonstrated the best goodness of

fit for all the other regions. We also calculated the peak age for each region based on

the optimal model, and results showed that the peak ages ranged from 33.4 to 40.6 years.

In addition, we found the peak ages obtained from the above optimal model (i.e., square

root or quadratic models) and the GAM are consistent, $r = 0.69$, $p < 0.05$. The detailed

results of the optimal model fitting are shown in Figure 3 and supplementary Table S6.

Figure 3. Brain regions showing inverted U-shaped trajectory patterns. Scattered plots are the effect size
as a function of age, with curves fitted by GAM (blue color) and the best simplified model (red color).
Shaded areas around the curves represent standard errors. Dashed vertical lines show peak ages estimated
from GAM (blue) and simplified model (red). The sizes of the scattered dots show the square root of
model weights (1/variance) for each study. The numbers at the up-left corner of each plot correspond to
the orders in Table 1. 1: right inferior frontal gyrus, 2: left inferior frontal gyrus, 3: right inferior parietal
lobule, 4: left supplementary motor area, 5: left inferior parietal lobule, 6: right caudate nucleus, 7: left
insula, 8: left insula, 9: right anterior cingulate cortex.

Fitting the Whole Brain with Square Root and Linear Models

Based on model comparisons, we found that the square root model provided the best

goodness of fit for the age-related change of the brain activities. To supplement the

inverted U-shaped results from the contrast analysis, the square root function was then

submitted to whole-brain meta-regression analyses in SDM-PSI.

By fitting the activation over the whole brain, we found 7 significant brain regions,

including the bilateral inferior frontal gyrus, right superior longitudinal fasciculus, left

inferior parietal lobule, right insula, right caudate nucleus, and left anterior thalamic

projections (Figure 4A and supplementary Table S7). These results further supported

the existence of the inverted U-shaped regions we initially identified.

In addition, we explored the whole-brain trajectories with a linear meta-regression.

However, no significant regions were observed (Figure 4B), even under a more tolerant

threshold of uncorrected $p < 0.01$.

Figure 4. Significant clusters (voxel-wise FWE-corrected, $p < 0.001$, voxels ≥ 10) showing square root
pattern (A) and linear pattern (B) with age in the model fitting.

Dissociated Brain Networks with Distinct Lifespan Trajectories

We note that the inverted U-shaped regions constitute only part of the cognitive control-related regions (Figure 2), and the remaining regions appear to show a relatively flat trajectory. To further elucidate the spatial distribution of brain regions following these different trajectory patterns, we used the results from the mean SDM analysis (see section "Regions Related to Cognitive Control Identified by both ALE and SDM") as the mask and replotted the results of the contrast analysis between young adults and others with two different thresholds, and then compared them with the Yeo's 7-network atlas[72] (Figure 5). Results showed that the distribution of inverted U-shaped regions was more consistent with the frontoparietal control network, and the distribution of non-U-shaped regions was more closely related to the dorsal attention network (Figure 5B). The count of voxels revealed that a larger portion of the inverted U-shaped regions overlapped with the frontoparietal control network (2,538 voxels) than with the dorsal attention network (932 voxels), while the overlap with the cingulo-opercular network was in between (1,867 voxels). Conversely, a larger portion of the non-U-shaped regions overlapped with the dorsal attention network (3,009 voxels) than with the frontoparietal control network (2,447 voxels), while the overlap with the cingulo-opercular network was lower (1,594 voxels). In addition, the visualization of spatial distribution patterns revealed a dissociation between the middle frontal gyrus and its adjacent rostral and caudal areas (Figure 5A).

Figure 5. Dissociated brain regions based on their trajectory patterns. A) The regions following inverted U-shaped trajectories (red color) and non-U-shaped trajectories (blue color). B) The axial view of the same results. The border lines display the frontoparietal control network (black), dorsal attention network (white), and their boundary (gray) from Yeo's 7-network atlas[72]. Cingulo-opercular network was not plotted due to its less clear dissociation among the two maps. The two scatter plots show two example regions showing the non-U-shaped trajectory.

To further investigate the potential functional difference between the two sets of brain regions, we decoded the related terms with the Neurosynth decoder[64] (see section "Neurosynth Decoding Analysis" in Methods). Results showed that the inverted U-shaped regions were related to the term "cognitive control" with a relatively high correlation ($r = 0.187$), but were less related to "attentional" ($r = 0.100$) and "monitoring" ($r = 0.072$). Note the key word "monitoring" refers to the major function of the cingulo-opercular network[27]. On the other hand, the non-U-shaped regions were related to the term "attentional" ($r = 0.240$) but were less related to "monitoring" ($r = 0.090$) and "control" ($r = 0.062$) (supplementary Table S8). This result suggests that the inverted and non-U-shaped regions may be associated with cognitive control and attention, respectively, which is consistent with the frontoparietal control network and dorsal attention network as identified in the atlas overlapping analysis.

To provide additional insight into the non-U-shaped trajectory, we specifically

chose two coordinates for analysis (Figure 5B), one ([34, 4, 52]) representing a peak

region from the average brain activity analysis using SDM-PSI (supplementary Table

S3), and the other ([22, −63, 42]) representing a region displaying a null effect from the

contrast analysis with p between 0.49 and 0.51, indicating a trajectory that does not

align with either an inverted or upright U-shaped pattern. The GAM analysis showed

that neither coordinate could be adequately fitted by a smoothed curve, with ps > 0.27.

These results elucidate how the non-U-shaped regions exhibit relatively stable brain

activities across the lifespan compared to regions showing inverted U-shaped patterns.

Lifespan Trajectory of the Laterality

We also tested how the laterality of the brain activity changes with age (see section

"Laterality Analysis" in Methods). We first modeled the laterality trajectory using the

GAM. Results showed a significant model fitting, $F(4.3, 5.3) = 3.38$, $p = 0.013$. More-

over, we fitted the data with the four simplified models (i.e., the quadratic, cubic, square

root, and quadratic logarithmic models). The results showed that a quadratic logarith-

mic function provided the best goodness of fit, with the $\beta_{\log(age)} = -2.48$ and $\beta_{(\log(age))^2} =$

0.36, both ps < 0.001. The two-line test suggests the hypothetical peaks from both mod-

els did not reach significance (ps > .22), however. A visually upright U-shaped trajec-

tory indicated the youth and elderly adults tended to be more left lateralized across the

whole brain. A further comparison of the relative levels of laterality across different

age groups revealed that both the youth and the elderly groups exhibited greater left

lateralization than the young adult group, *p*s < .05, one-tailed. This left lateralized pat-

tern could be illustrated by the brain map estimated with voxel-wise laterality calcula-

tion (Figure 6B). In addition, the prefrontal regions appeared to be the major area driv-

ing the U-shaped trajectory, with the youth and elderly adults showing more left lat-

eralized regions than the young adult group.

A previous study[27] showed that the semantic conflicts (e.g., Stroop) are left-lat-

eralized and the response conflicts (e.g., Simon) are right lateralized. However, inclu-

sion of the conflict type as nuisance covariables in our model fitting analysis could

essentially rule out the possibility that the laterality trajectory results were driven by

different conflict types.

Figure 6. The laterality as a function of age. A) The trajectory fitted with a quadratic logarithmic model
(red) and the GAM (blue). Higher values mean more left-lateralized and lower values mean more right-
lateralized. B) Visualization of the laterality for each group. Regions in the left hemisphere show the left
laterality, and vice versa.

Discussion

The objective of this meta-analysis was to investigate the age-related changes in brain

regions associated with cognitive control across the lifespan. Using data from 129 stud-

ies that encompass participants aged 8 to 74 years on average, we examined how the

activities of cognitive control regions vary with age. Our analyses yielded three primary

findings: 1) Among different possible trajectories, only the inverted U-shaped trajecto-

ries were reliably observed across the whole brain; 2) The cognitive control related

brain regions exhibit heterogeneous lifespan trajectories: the frontoparietal control net-

work (such as the inferior frontal gyrus and inferior parietal lobule) follows inverted U-

shaped trajectories, peaking between 24 and 41 years, while the dorsal attention net-

work (such as the frontal eye field and superior parietal lobule) demonstrates flatter

trajectories with age; 3) The youth and the elderly demonstrate weaker brain activities

and a relatively greater extent of left laterality compared to the young adults. These

results provide strong evidence for the existence of cognitive control regions exhibiting

inverted U-shaped trajectory, and also show the heterogenous lifespan trajectories in

different brain regions.

The Inverted U-shaped Trajectory of Brain Activity Related to Cognitive Control

The main result is that a wide range of cognitive control regions follow inverted U-

shaped lifespan trajectories, but no regions showed decrease-then-stable (Figure 7D),

501   upright U-shaped (Figure 7E), stable-then-increase (Figure 7F), increase-then-stable

502   (Figure 7G), stable-then-decrease (Figure 7I), or linear trajectories (Figure 7, panels A,

503   and C).

504
Figure 7. The lifespan trajectories explored in our study. Panels A-C show linear decrease, flat, and linear
increase patterns, respectively, and were modelled with the linear function. Panels E and H show the
upright and inverted U-shapes, respectively, and were tested with the contrast between young adults and
others, as well as with the quadratic function. Panels D, F, G, and I show combinations of a stable period
and an increase/decrease period across the lifespan, and were tested with the contrast between the youth
and others, or between the older adults and others. Panels J, K and L show the variants of inverted U-
shaped trajectories, which capture the possibly early peak feature. They were tested with square root,
quadratic logarithmic, and cubic functions, respectively. See Methods for detailed models.

The greater activation in the frontoparietal control network among young adults

compared to the youth and the elderly supports the notion that cognitive control abilities

may not be fully developed in children and may decline in the elderly. This finding is

consistent with the idea that the cognitive control system is most effective in young

adulthood, suggesting a possible correlation between the higher functional activations

in the brain and the superior performance of young adults on cognitive control tasks[17].

The inverted U-shaped trajectories of cognitive control regions, instead of an increase-

then-stable trajectory, are also consistent with the idea that cognitive control is an im-

portant aspect of fluid intelligence, which is known to peak in early to mid-adulthood

and decline with age[5]. Hence, our findings may reflect the general rise-and-fall patterns

in the higher-order brain functions.

The inverted U-shaped trajectory of brain activation might be associated with the

development of brain structure and functional changes. First, it may reflect the well-

documented structural changes that occur in these regions across the lifespan, which

include synaptic pruning, myelination, cortical thinning, and white matter matura-

tion[22,23]. For example, the density of dopamine receptors increases during adolescence

and young adulthood and subsequently declines with age[73]. These changes can affect

the efficiency and connectivity of neural circuits within the frontoparietal control net-

work[15,49]. Second, the inverted U-shaped trajectory may also arise from functional

changes resulting from the modulation of neurotransmitters, hormones, and environ-

mental factors[74]. Understanding change patterns of brain structure and function is crit-

ical for developing interventions and treatments aimed at improving cognitive control

abilities across the lifespan.

The present results further revealed that the inverted U-shaped lifespan trajectories

of cognitive control regions are not uniform. The GAM fitting results (Figure 3, sup-

plementary Table S4) showed that the subregions exhibited varying degrees of modu-

lation effects by age. Moreover, the model simplification demonstrated different under-

lying trajectory curves, with most regions showing a skewed shape that could best be

fitted with a square root model, except that one region showed a symmetric quadratic

shape. The quadratic lifespan trajectory has been well-documented in previous stud-

ies[22,47,69,71], while the application of the square root model has been relatively rare[75].

The square root model can better capture the early peak in the trajectory, which aligns

with the hypothesis. We also identified different peak ages for those regions, ranging

from 24 to 41 years, suggesting that cognitive control regions may not develop at the

same rate. Hence, the developmental peaks of cognitive control occur at distinct time

points across various brain regions.

Hierarchical development trajectories in different brain networks

Previous research has indicated that the attentional orientation function is preserved

during aging[41,76]. Consistently, we found that the dorsal attention network regions un-

derlying the attentional orientation showed a relatively flat pattern, in contrast to the

inverted U-shaped trajectory in frontoparietal control network regions. The dissociation

between these two sets of regions may reflect hierarchical age effects on brain function.

The brain regions are organized in a functional hierarchy, with the frontoparietal

control network to be at the highest level. It acts as a connectivity hub, receiving inputs

from and providing feedback to other systems, including the dorsal attention network[77].

Within the cingulo-opercular network, we did not observe a clear dissociation of tra-

jectory patterns, which may correspond with its intermediate position in the hierarchy

between the frontoparietal control and dorsal attention networks[58]. During conflict tasks,

the three sets of regions are coordinated in a hierarchical manner. The frontoparietal

control network maintains task goals and resolves conflicts, the cingulo-opercular net-

work monitors the existence of conflict, and the dorsal attention network directs atten-

tion towards task-relevant stimuli[78]. Even within the prefrontal cortex itself, a hierar-

chical organization exists, with middle frontal areas occupying the peak position[79]. This

is in line with our finding that the middle frontal cortex is dissociated from rostral and

caudal frontal regions.

Furthermore, different brain regions exhibit different age effects. Higher-order re-

gions typically have more complex lifespan trajectories[47] and reach peaks during later

periods[49,50]. Specifically, prefrontal control regions are among the last to mature and

one of the earliest to decline[5,15,51]. Therefore, the different lifespan trajectory patterns

among frontoparietal control, cingulo-opercular and dorsal attention regions likely re-

flect their different positions within the functional hierarchy and the specific patterns

of age-related changes.

No Evidence for the Compensatory and Asymmetry Reduction Theories

Critically, there was no region showing higher activity in the elderly compared to the

young adults, but we observed several regions showing the opposite (see supplementary

Note S3 and Table S9). This finding is inconsistent with the compensatory theory[11],

which proposes that the elderly recruit additional brain regions to compensate for age-

related cognitive decline. Moreover, we did not observe a strong laterality in young

adults, diverging from the findings from previous studies[46]. In contrast, we found that

elderly adults showed a tendency of left laterality, which diverges from the hemispheric

asymmetry reduction in older adults (HAROLD) theory[45]. The HAROLD theory sug-

gests that older adults typically exhibit less lateralization, either as a compensatory re-

sponse to functional deficits or as a reflection of neural dedifferentiation. Stronger pat-

terns of left laterality were also identified in childhood, primarily driven by the prefron-

tal region (Figure 6). The left laterality at both ends of the age spectrum reflects differ-

ential age effects on the cerebral hemispheres. The left hemisphere is known to undergo

earlier development[80] and later decline[81] compared to the right hemisphere. Conse-

quently, a stronger imbalance in activity favoring the left hemisphere may occur in both

childhood and elderly adulthood. Notably, it is likely that previous studies showing

stronger laterality in young adults may have been biased by too small sample sizes and

the use of non-quantitative methods for calculation of laterality[82,83]. Moreover, since

both left and right lateralized results were reported in the literature on laterality[81], it is

reasonable to observe a low laterality for the young adults in the current meta-analysis.

Implications for Other Cognitive Control Subdomains

A more comprehensive understanding of the lifespan trajectory of cognitive control

relies on studying different subdomains of cognitive control, such as conflict processing,

working memory, and cognitive flexibility[67]. Previous studies have suggested that dif-

ferent subdomains of cognitive control exhibit both similar and separable neural mech-

anisms[84]. Moreover, these subdomains might follow different lifespan trajectories[19].

Findings from our study may contribute valuable insights into the potential differences

in brain activity across different components of cognitive control during development.

Consistent with the inverted U-shaped trajectory, we confirmed that the young

adults exhibit higher levels of cognitive control brain activities compared to the youth

during conflict processing but not otherwise (supplementary Figure S3, Table S9). This

observation is in line with previous studies[30-34]. To the best of our knowledge, previous

research using conflict tasks has rarely reported findings that contradict this direction

except [35]. However, more controversial patterns have emerged in studies focusing on

other subdomains of cognitive control, such as working memory[85-87] and response in-

hibition[88-90]. For instance, with visual-spatial working memory task, Scherf, Sweeney

[85] found an age-related increase in the recruitment of brain regions in the lateral pre-

frontal cortex, whereas Geier, Garver [87] reported both age-related increase and decrease

in different prefrontal regions. These studies employed similar paradigms but different

contrast methods (i.e., delay vs. non-delay and long-delay vs. short-delay), which ex-

emplifies the many diversities within the working memory domain[91]. Similarly, in the

subdomain of response inhibition, studies have shown discrepancies in brain engage-

ment with different designs and different measurements[92,93]. On the contrary, conflict

processing provides a more consistent framework for measuring brain activities across

age groups. Given these differences, it is prudent to be cautious in generalizing our

current findings to the lifespan trajectories of other subdomains of cognitive control,

although an inverted U-shaped trajectory could be a good candidate hypothesis to begin

with. Future research is necessary to develop more precise cross-age comparisons

within these other cognitive control subdomains.

Methodology Implications

By incorporating all the studies, our results demonstrate that the SDM has relatively

higher sensitivity than that of the ALE in identifying involved brain regions, consistent

with the conclusion from a previous study[94]. Moreover, the SDM also has the added

advantage of fully utilizing existing effect size data and coordinates[95], allowing us to

compare the relative activity strength among various age groups, such as the contrast

between young adults and elderly groups. More importantly, this approach allows for

meta-regressions to examine parametric relationships between brain region activity and

age, providing insights into the lifespan trajectories of cognitive control regions. Our

simulation analysis also confirms the efficacy of using mean variables as predictors for

such regression. Future studies could implement the meta-regression approach to better

link cognitive-behavioral measurements with brain functions. However, it is important

to acknowledge that the coordinate based SDM analysis is an approximation of the

actual statistical images. As such, image-based approaches are preferred whenever fea-

sible[96-98].

Limitation of Results

One caveat to consider in this study is the non-uniform distribution of age among the

included studies. Specifically, most of the young adult studies (80 out of 84) had aver-

age ages lower than 45 years, resulting in a noticeable gap in the age range of 45 to 60

years. While these studies may have recruited an adequate number of individuals within

this age range, the focus on the mean age may limit our study to capture this variation.

Consequently, it is important to acknowledge that the observed age distribution could

potentially impact the results of the regression analysis. We hope that future research

could allocate more attention to the middle-aged period, considering the significant

cognitive and neural changes that may occur during this stage, such as the onset of

cognitive decline[18,20,99]. Besides, it is crucial to avoid the occurrence of ecological fal-

lacy[100] when interpreting the results of meta-regression analyses. The ecological fallacy

arises when associations observed at the group level are erroneously assumed to apply

to individuals within those groups. Therefore, while our analysis reveals significant as-

sociations between brain activities and age across various studies, it does not provide

direct insights into the specific age effects at the individual level. Future research in-

corporating individual-level investigations (e.g., longitudinal follow-up studies) is cru-

cial to obtaining a more comprehensive understanding of these relationships.

Conclusions

Our meta-analysis adopted advanced meta-regression approaches to chart the lifespan

trajectories of cognitive control brain activities. We observed inverted U-shaped chang-

ing patterns in regions aligned with the frontoparietal control network, with the peaks

occurring between 24 and 41 years. In contrast, the dorsal attention network presented

a relatively flat trajectory. This dissociation may reflect the hierarchy of brain develop-

ment in different regions. No other trajectory patterns were observed, highlighting the

predominance of the inverted U-shaped pattern in the lifespan trajectory of cognitive

control. Furthermore, we found older adults showed lower brain activity strength and a

more asymmetric brain distribution than young adults, in contrast to the hypotheses

from the compensatory and asymmetric reduction theories of aging. In sum, these re-

sults demonstrate the multifaceted nature of age-related changes in cognitive control

brain function.

Methods

Literature Preparation

Literature Search

We report how we determined our sample size, all data exclusions (if any), all manipulations, and all measures in the study. We first searched both English and Chinese articles on the youth and the elderly from PUBMED, Web of Science and CNKI (China National Knowledge Infrastructure) till 2022. The following search terms were applied in titles, abstracts, table of contents, indexing, and key concepts: *("Stroop" OR "Flanker" OR "Simon" OR "SNARC" OR "Navon" OR "interference" OR "cognitive conflict") AND ("fMRI" OR "functional resonance imaging" OR "functional imaging" OR "neuroimaging" OR "PET") AND ("children" OR "kids" OR "adolescents" OR "teenagers" OR "underage" OR "aged" OR "old" OR "older" OR "elder" OR "elderly" OR "senior" OR "development" OR "developmental" OR "aging" OR "life span").* The above process yielded 3,484 articles. Besides, 111 studies on the young adult from a previous meta-analysis study[27] were adopted, which had applied very similar recruitment criteria. Moreover, we screened 16 articles citing or being cited by the crucial literature. After removing duplicates, the literature search identified 2,930 articles.

 Exclusion Criteria

We excluded any articles that met one or more of the following predefined exclusion

criteria: 1) not in English or Chinese; 2) not including healthy human participants; 3)

case study; 4) not empirical study; 5) not functional resonance imaging (fMRI) or pos-

itron emission tomography (PET) study; 6) not whole-brain results (i.e., not have cov-

ered the whole gray matter); 7) not in Talairach or Montreal Neurological Institute

(MNI) space; 8) not reflecting the congruency effect (i.e., contrasts between incongru-

ent and congruent or between incongruent and neutral conditions); 9) not reporting ex-

act mean age of participants.

A total of 119 articles were identified as eligible for inclusion in our meta-analyses.

Supplementary Figure S1 shows the preferred reporting items for systematic reviews

and meta-analyses (PRISMA)[101] flow chart for the literature screening process. The 119

articles included 129 studies (individual contrasts reported in the articles) with 3,388

participants and 1,579 activation foci reported. None of the experiments share the same

group of participants.

Coding Procedure

A coding manual was formulated to record pertinent study information, including au-

thors, publication dates, experimental tasks, contrasts, and sample demographics (such

as the average age and sample size). To ensure coding accuracy, two authors inde-

pendently coded all studies, with discrepancies resolved through discussion or refer-

ence to the original studies. In instances where studies lacked essential information,

such as peak coordinates for relevant contrasts, participant age averages, or data for

specific age groups, efforts were made to contact the authors via e-mail to obtain the

relevant data. In addition, both coordinates and effect sizes (i.e., Hedge's *g*) were ex-

tracted from each study. Further, Talairach space coordinates were transformed to MNI

coordinates using the Lancaster transform[102].

Meta-Analytic Procedure

Activation Likelihood Estimation (ALE)

In order to obtain a comprehensive understanding of cognitive control-related brain ac-

tivity across all age groups and to replicate a prior study[27], we initially conducted a

single dataset meta-analysis using BrainMap GingerALE software[103] (version 3.0.2,

http://www.brainmap.org). This meta-analytical approach, known as activation likeli-

hood estimation, utilizes the spatial convergence of brain activity across multiple stud-

ies to determine the probability of activation in specific regions. Foci from individual

studies were transformed into a standardized coordinate space and modeled as Gaussian

probability values that accounted for variability in the number of participants in each

study. In situations where foci overlapped across studies, multiple Gaussians were as-

sociated with a single focus, and ALE selected the Gaussian with maximum probability

for each focus[103]. Subsequently, ALE score maps were generated by comparing these

modeled Gaussian distributions with a null distribution that simulated random brain

effects. The null distribution was generated using the same sample size and number of

foci groups as the experimental dataset for 1,000 times[104]. ALE scores were then used

to calculate p-values, which were based on the proportion of values higher than a certain

threshold in the null distribution. This resulted in a statistical ALE map that differenti-

ated true brain effects from random effects. A cluster-defining threshold of $p < 0.001$

and a minimum cluster size of 10 voxels (80 mm^3) were utilized to compute ALE maps,

which matches the threshold applied in the seed-based d mapping (SDM) approach (see

below).

Seed-based Effect Size Mapping

SDM is an alternative approach to statistically synthesize results from multiple fMRI

experiments[95]. Similar to ALE, SDM employs a coordinate-based random-effects ap-

proach to amalgamate peak coordinate information into a standard space across several

experiments. However, while ALE solely considers the binary feature (i.e., active ver-

sus inactive) of peak coordinates, SDM takes into account the quantitative effect size

(can be positive or negative) connected to each peak and reconstructs the initial para-

metric maps of individual experiments before amalgamating them into a meta-analytic

map[105]. Therefore, the use of a distinct algorithm in SDM from ALE allows us to scru-

tinize the robustness and replicability of the outcomes obtained via ALE. More im-

portantly, SDM enables the inclusion of covariates in the meta-regression analyses to

reflect the changes in brain function across the lifespan.

We conducted three types of analyses using the SDM approach. Firstly, we esti-

mated the mean activation across all age groups and compared the results with ALE's

single dataset meta-analysis results. Secondly, we conducted contrasts between two

groups of studies (e.g., between young adults and a combination of the youth and el-

derly groups) to identify brain regions that showed different levels of activity across

age. This analysis method served to investigate the hypothesized lifespan trajectories,

such as the inverted U-shaped pattern by elucidating neural activity variations linked to

age. Thirdly, we defined specific models (e.g., linear and square root models) to fit the

whole brain to validate brain regions adhering to the hypothetical lifespan changing

patterns. This type of analysis aimed to explore various lifespan trajectories, recogniz-

ing that different brain regions might follow distinct model functions.

These analyses were conducted using the software of SDM with permutation of

subject images (SDM-PSI) (version 6.22, https://www.sdmproject.com). Effect size

maps were built for the 129 individual experiments. This was accomplished by (a) con-

verting the statistical value of each peak coordinate into an estimate of effect size

(Hedge's g) using standard formulas[106] and (b) convolving these peaks with a fully an-

isotropic unnormalized Gaussian kernel ($\alpha = 1$, FWHM = 20 mm) within the bounda-

ries of a gray matter template (voxel size = $2\times2\times2$ mm^3). Imputation (50 times) was

conducted for each study separately to obtain an reliable estimate of brain activation

maps[105]. Besides, the individual effect size maps were combined using a random-effect

general linear model. To assess the statistical significance of activations in the resulting

meta-analytic effect size map, 1,000 random permutations of activation peaks within

the gray matter template were compared. Finally, the meta-analytic maps were

thresholded using a voxel-wise family-wise error (FWE) corrected threshold of $p <$

0.001 and a cluster-wise extent threshold of 10 voxels[107].

Mean Analyses Across all Studies

This analysis aimed to characterize the activation distributions of cognitive control-

related brain regions across all studies, which was conducted utilizing the "Mean" func-

tion in SDM-PSI software. In order to verify the reliability of the SDM analysis results,

we compared the results with the single dataset meta-analysis using ALE. Results from

this analysis were further used as regions of interest (ROIs) in the subsequent model

fitting analyses (see the section "Generalized Additive Model (GAM) Fitting").

Contrast Analyses

To test our hypothesis that cognitive control related brain activities follow an inverted U-shaped trajectory with age, we categorized each study based on the mean age of participants into youth (< 18 years), young adults (18−59 years), and elderly (>= 60 years) groups. The age boundaries were determined to minimize age distribution overlap. We utilized SDM-PSI to perform a contrast analysis between the group of young adults and the combination of other groups in order to examine whether there are brain regions that exhibit an inverted U-shaped lifespan trajectory. This was achieved by assigning studies from the young adult group as 1 and all other studies as −1. This analysis yielded two results, one showing higher activity in young adults than the youth and elderly groups, and the other showing the opposite. Like the mean analysis, results from this analysis were used as ROIs in the subsequent model fitting analyses (see sections "Generalized Additive Model (GAM) Fitting" and "Model Simplification and Model Comparison"). The possibility of publication bias for resultant clusters was examined using Egger's test[108], in which any result showing $p < 0.05$ was regarded as having significant publication bias. Heterogeneity was evaluated using the I^2 index, which quantifies the proportion of total variability attributable to heterogeneity between studies. A value less than 25% indicates low heterogeneity among the included studies[109].

In addition, to explore other possible trajectories, such as the increase-and-stable pattern[5], we conducted contrast analyses between the youth and the combined group of

young and old adults, as well as between old adults and the combined group of youth

and young adults. Furthermore, to address the controversies in previous studies, we

conducted contrast analyses between old and young adult groups, and between the

youth and young adult groups, respectively.

Meta-regression Analyses

To better describe the possible lifespan trajectories of the whole brain, we carried out

meta-regression analyses with the age and/or its derivatives as regressors. Two regres-

sions were conducted across the whole-brain, including a linear regression (with only

age as the regressor) and a square root regression (with age and its square root as sepa-

rate regressors). The linear regression aimed to test regions with increasing/decreasing

activity with age, and the square root regression aimed to test regions with the inverted

U-shaped trajectories based on the model fitting results (see below).

Data Extraction

Masks were generated for each ROI derived from the mean and contrast analyses in

SDM-PSI as described above. Subsequently, we extracted the effect sizes for each mask.

Fifty values were obtained from the SDM iterations and subsequently averaged for each

study in each region. The iterated variances were also averaged in a similar way. Addi-

tionally, we removed the outliers (beyond 3 standard deviations from the mean) in the

following model fitting analyses.

Generalized Additive Model (GAM) Fitting

To precisely estimate the inverted U-shaped trajectories, we adopted the GAM to fit the

curves. The GAM allows for flexible, nonparametric smoothing of predictor varia-

bles[110], and has been widely used to depict the lifespan trajectories[23,111]. We imple-

mented GAMs using the "mgcv" package[110] in R. For each ROI, we fitted a GAM with

the following formula:

$g \sim s(age) + covariates,$

where g is the effect size (dependent variable), s(age) represents a smoothing spline of

age (predictor variable), and covariates represent the dummy-coded categorical covari-

ate regressors. These regressors correspond to four aspects of the included studies: (1)

the presence of various conflict types (e.g., Stroop or Simon), (2) the mixed subject

samples based on handedness (e.g., right handed only or both handed), (3) the different

contrasts in reporting congruency effects (e.g, incongruent – congruent or incongruent

– neutral), and (4) different trial types regarding whether they excluded error trials. We

incorporated these covariates to control for their potential confounding effects related

to age, which could otherwise influence our results. We also adjusted the estimate with

a weight parameter, which was the reciprocal of variance. We used penalized regression

splines, with the amount of smoothing determined automatically based on generalized

cross validation.

For each ROI, we quantified the peak age by choosing the highest prediction of a

fine-grained age scale (1,000 points from 8 to 74 years old). We also calculated the

estimated degree of freedom (EDF) for the smooth curve by summing up the degree of

freedom for each penalized term (i.e., s(age).1 to s(age).9).

Model Simplification and Model Comparison

While the GAM analysis may yield good fitting results on the data, it is important to

acknowledge its potential limitations. One concern is that it can fit the data with high

degree of freedoms (up to 7.0, supplementary Table S4), which makes it susceptible to

over-fitting and harder to generalize. Another issue is its poor interpretability. Therefore,

we next sought to fit the data with simpler models.

To this end, we used the "metafor" package in R to fit these effect sizes with the

age and its derivatives as predictors. Specifically, we tested four non-linear models (see

below formulas and Figure 7). Quadratic and cubic models were included based on

previous studies[69,70]; quadratic logarithmic and square-root models were included to

capture the possible skewed trajectory, which would reflect the asymmetric trajectory

of development and decline of cognitive control[5]. Each model was fitted to each ROI

separately. In addition, we included the same covariates as the GAM analysis in each

regression model. We calculated the Akaike information criterion (AIC) to evaluate the

goodness of fit for each model.

1) Quadratic model

$g \sim \text{age} + \text{age}^2 + \text{covariates}$

2) Cubic model

$g \sim \text{age} + \text{age}^2 + \text{age}^3 + \text{covariates}$

3) Quadratic logarithmic model

$g \sim \log(\text{age}) + (\log(\text{age}))^2 + \text{covariates}$

4) Square root model

$g \sim \text{age} + \text{sqrt}(\text{age}) + \text{covariates}$

Neurosynth Decoding Analysis

To investigate the potential functional difference between the two sets of brain regions

reported in section "Dissociated Brain Networks with Distinct Lifespan Trajectories",

we generated two binary maps from the contrast analysis between young adults and the

combination of other groups, and then submitted them to the Neurosynth decoding sys-

tem[64]. As the non-U-shaped map is defined as the converse of the inverted U-shaped

map, they were obtained by applying thresholds of $0.1 < p < 0.9$ and $p < 0.05$, respec-

tively. These lenient threshold boundaries were used to minimize the impact of sparsity

on the decoding results. The two-boundary threshold was used in generating the non-

U-shaped map, as the original statistical map from the SDM-PSI analysis was one-

tailed, which means the threshold of $p > 0.9$ indicates the upright U-shaped trend instead

of a flat trajectory. Additionally, to focus on the brain regions specifically related to

cognitive control, the two maps were masked by using results from the mean SDM

analysis (see section "Regions Related to Cognitive Control Identified by both ALE

and SDM" in Results). In the decoding results, we deleted terms that were related to

brain regions (e.g., "frontal"), not functional specific (e.g., "task"), and duplicated (e.g.,

"attention" was removed if there was already "attentional").

Laterality Analysis

We calculated the laterality based on the effect size of reported brain coordinates from

each study. We computed the sum of effect sizes across coordinates for the left and right

hemisphere, respectively, yielding one global effect size each (i.e., g_L and g_R). Then,

we calculated the index of brain laterality with the following equation[112]:

$$\text{laterality} = \frac{g_L - g_R}{g_L + g_R} \ ,$$

which was then submitted to the GAM and simplified models (i.e., the quadratic, cubic,

square root, and quadratic logarithmic models). In addition, to directly assess the dif-

ferences in laterality levels across different age groups, we implemented two linear

meta-regression models: one comparing the youth with young adults, and the other

comparing the elderly with young adults. In each model, we introduced a variable that

encoded the specific groups of studies involved, designating the youth and elderly

groups with a code of 1, and the young adult group with a code of -1. Once again, we

included covariates in the above-mentioned analyses to control for potential confound-

ing factors, such as variations in task types.

To illustrate the contribution of different brain regions to the age-related change

of laterality, we calculated the laterality for each voxel[113]. We first used the SDM-PSI

to conduct a mean analysis for each age group (i.e., the youth, young adult and elderly),

yielding three z-maps. Then the laterality was computed with the above equation for

each voxel from the left hemisphere. The opposite values were calculated for the right

hemisphere. To avoid the bias due to asymmetric brain hemispheres, we removed

voxels without corresponding mirror coordinates. The results were visualized confining

to the brain regions estimated from the grand mean analyses (see section "Seed-based

Effect Size Mapping" in Results).

Transparency and Openness

We adhered to the MARS guidelines for meta-analytic reporting[114]. All meta-analytic

data and analysis codes are available at https://anonymous.4open.science/r/Cogni-

tive_Control_Developmental_Trajectory-66E5. Data were analyzed using R, version

4.3.0 (R Core Team, 2018), and the packages mgcv, version (1.8.42)[115] and metafor,

version 3.0-2[116]. This review project was not preregistered. It should be noted that ethics

committee approval was not obtained for this study because it involved the analysis of

existing data rather than the collection of new data from human participants. However,

all datasets used in the analysis were obtained from studies that had undergone ethical

review and approval.

References

1. Cohen JD. Cognitive control: Core constructs and current considerations. In:
The Wiley handbook of cognitive control (ed Egner T) (2017).

2. Miller EK, Cohen JD. An integrative theory of prefrontal cortex function. *Annu*
Rev Neurosci **24**, 167-202 (2001).

3. Li Z, Goschl F, Yang G. Dissociated Neural Mechanisms of Target and
Distractor Processing Facilitated by Expectations. *J Neurosci* **40**, 1997-1999
(2020).

4. Constantinidis C, Luna B. Neural Substrates of Inhibitory Control Maturation
in Adolescence. *Trends Neurosci* **42**, 604-616 (2019).

5. Craik FI, Bialystok E. Cognition through the lifespan: mechanisms of change.
Trends Cogn Sci **10**, 131-138 (2006).

6. Gupta R, Kar BR. Development of attentional processes in ADHD and normal
children. *Prog Brain Res* **176**, 259-276 (2009).

7. Casey BJ, Getz S, Galvan A. The adolescent brain. *Dev Rev* **28**, 62-77 (2008).

8. Diamond A. Executive Functions. *Annu Rev Psychol* **64**, 135-168 (2013).

9. Casey BJ, Heller AS, Gee DG, Cohen AO. Development of the emotional brain.
Neurosci Lett **693**, 29-34 (2019).

10. Zanto TP, Gazzaley A. Attention and ageing. In: *The Oxford handbook of*
attention.). Oxford University Press (2014).

11. Reuter-Lorenz PA, Cappell KA. Neurocognitive Aging and the Compensation
Hypothesis. *Current Directions in Psychological Science* **17**, 177-182 (2008).

12. Association AP. Diagnostic and statistical manual of mental disorders : DSM-
5.). 5th edn. American Psychiatric Association (2013).

13. Ing A*, et al.* Identification of neurobehavioural symptom groups based on
shared brain mechanisms. *Nat Hum Behav* **3**, 1306-1318 (2019).

14. Adnan A, Chen AJW, Novakovic-Agopian T, D'Esposito M, Turner GR. Brain
Changes Following Executive Control Training in Older Adults. *Neurorehabil*
Neural Repair **31**, 910-922 (2017).

15. Casey BJ, Tottenham N, Liston C, Durston S. Imaging the developing brain:
what have we learned about cognitive development? *Trends Cogn Sci* **9**, 104-
110 (2005).

16. Grady C. The cognitive neuroscience of ageing. *Nat Rev Neurosci* **13**, 491-505
(2012).

17. De Luca CR, Leventer RJ. Developmental trajectories of executive functions
across the lifespan. In: *Executive functions and the frontal lobes*). Psychology
Press (2010).

18. Weintraub S*, et al.* Cognition assessment using the NIH Toolbox. *Neurology* **80**,
S54-64 (2013).

19. Ferguson HJ, Brunsdon VEA, Bradford EEF. The developmental trajectories of
executive function from adolescence to old age. *Sci Rep* **11**, 1382 (2021).

20. Li SC, Hammerer D, Muller V, Hommel B, Lindenberger U. Lifespan
development of stimulus-response conflict cost: similarities and differences
between maturation and senescence. *Psychol Res* **73**, 777-785 (2009).

21. Erb CD, Germine L, Hartshorne JK. Cognitive control across the lifespan:
Congruency effects reveal divergent developmental trajectories. *J Exp Psychol*
Gen **152**, 3285-3291 (2023).

22. Brouwer RM*, et al.* Genetic variants associated with longitudinal changes in
brain structure across the lifespan. *Nat Neurosci* **25**, 421-432 (2022).

23. Bethlehem RAI*, et al.* Brain charts for the human lifespan. *Nature* **604**, 525-533
(2022).

24. Luna B. Developmental Changes in Cognitive Control through Adolescence. In:
Advances in Child Development and Behavior (ed Bauer P). JAI (2009).

25. Giedd JN, Rapoport JL. Structural MRI of pediatric brain development: what
have we learned and where are we going? *Neuron* **67**, 728-734 (2010).

26. Hammerer D, Muller V, Li SC. Performance monitoring across the lifespan:
still maturing post-conflict regulation in children and declining task-set
monitoring in older adults. *Neurosci Biobehav Rev* **46 Pt 1**, 105-123 (2014).

27. Li Q, Yang G, Li Z, Qi Y, Cole MW, Liu X. Conflict detection and resolution
rely on a combination of common and distinct cognitive control networks.
Neurosci Biobehav Rev **83**, 123-131 (2017).

28. Giedd JN*, et al.* Brain development during childhood and adolescence: a
longitudinal MRI study. *Nat Neurosci* **2**, 861-863 (1999).

29. Crone EA, Dahl RE. Understanding adolescence as a period of social-affective
engagement and goal flexibility. *Nat Rev Neurosci* **13**, 636-650 (2012).

30. Andrews-Hanna JR, Mackiewicz Seghete KL, Claus ED, Burgess GC, Ruzic L,
Banich MT. Cognitive control in adolescence: neural underpinnings and
relation to self-report behaviors. *PLoS One* **6**, e21598 (2011).

31. Veroude K, Jolles J, Croiset G, Krabbendam L. Changes in neural mechanisms
of cognitive control during the transition from late adolescence to young
adulthood. *Dev Cogn Neurosci* **5**, 63-70 (2013).

32. Wilk HA, Morton JB. Developmental changes in patterns of brain activity
associated with moment-to-moment adjustments in control. *Neuroimage* **63**,
475-484 (2012).

33. Konrad K*, et al.* Development of attentional networks: an fMRI study with
children and adults. *Neuroimage* **28**, 429-439 (2005).

34. Marsh R*, et al.* A developmental fMRI study of self-regulatory control. *Hum*
Brain Mapp **27**, 848-863 (2006).

35. Bunge SA, Dudukovic NM, Thomason ME, Vaidya CJ, Gabrieli JD. Immature
frontal lobe contributions to cognitive control in children: evidence from fMRI.
Neuron **33**, 301-311 (2002).

36. Milham MP*, et al.* Attentional control in the aging brain: insights from an fMRI
study of the stroop task. *Brain Cogn* **49**, 277-296 (2002).

37. Yaple ZA, Stevens WD, Arsalidou M. Meta-analyses of the n-back working
memory task: fMRI evidence of age-related changes in prefrontal cortex
involvement across the adult lifespan. *Neuroimage* **196**, 16-31 (2019).

38. Prakash RS, Erickson KI, Colcombe SJ, Kim JS, Voss MW, Kramer AF. Age-
related differences in the involvement of the prefrontal cortex in attentional
control. *Brain Cogn* **71**, 328-335 (2009).

39. Fernandez NB, Hars M, Trombetti A, Vuilleumier P. Age-related changes in
attention control and their relationship with gait performance in older adults
with high risk of falls. *Neuroimage* **189**, 551-559 (2019).

40. Sebastian A*, et al.* Differential effects of age on subcomponents of response
inhibition. *Neurobiology of aging* **34**, 2183-2193 (2013).

41. Verhaeghen P, Cerella J. Aging, executive control, and attention: a review of
meta-analyses. *Neurosci Biobehav Rev* **26**, 849-857 (2002).

42. Verhaeghen P. *The elements of cognitive aging: Meta-analyses of age-related
differences in processing speed and their consequences*. OUP USA (2014).

43. Wood G, Ischebeck A, Koppelstaetter F, Gotwald T, Kaufmann L.
Developmental trajectories of magnitude processing and interference control:
an FMRI study. *Cereb Cortex* **19**, 2755-2765 (2009).

44. Crone EA, Steinbeis N. Neural Perspectives on Cognitive Control Development
during Childhood and Adolescence. *Trends Cogn Sci* **21**, 205-215 (2017).

45. Cabeza R. Hemispheric asymmetry reduction in older adults: the HAROLD
model. *Psychol Aging* **17**, 85-100 (2002).

46. Park DC, Reuter-Lorenz P. The adaptive brain: aging and neurocognitive
scaffolding. *Annu Rev Psychol* **60**, 173-196 (2009).

47. Shaw P*, et al.* Neurodevelopmental trajectories of the human cerebral cortex. *J
Neurosci* **28**, 3586-3594 (2008).

48. Teissier A, Pierani A. Wiring of higher-order cortical areas: Spatiotemporal
development of cortical hierarchy. *Semin Cell Dev Biol* **118**, 35-49 (2021).

49. Gogtay N*, et al.* Dynamic mapping of human cortical development during
childhood through early adulthood. *Proc Natl Acad Sci U S A* **101**, 8174-8179
(2004).

50. Sydnor VJ*, et al.* Intrinsic activity development unfolds along a sensorimotor-
association cortical axis in youth. *Nat Neurosci* **26**, 638-649 (2023).

51. Raz N*, et al.* Selective aging of the human cerebral cortex observed in vivo:
differential vulnerability of the prefrontal gray matter. *Cereb Cortex* **7**, 268-282
(1997).

52.   Duncan J. The multiple-demand (MD) system of the primate brain: mental
programs for intelligent behaviour. *Trends Cogn Sci* **14**, 172-179 (2010).

53.   Dixon ML*, et al*. Heterogeneity within the frontoparietal control network and
its relationship to the default and dorsal attention networks. *Proc Natl Acad Sci*
U S A **115**, E1598-E1607 (2018).

54.   Spreng RN, Sepulcre J, Turner GR, Stevens WD, Schacter DL. Intrinsic
architecture underlying the relations among the default, dorsal attention, and
frontoparietal control networks of the human brain. *J Cogn Neurosci* **25**, 74-86
(2013).

55.   Corbetta M, Shulman GL. Control of goal-directed and stimulus-driven
attention in the brain. *Nat Rev Neurosci* **3**, 201-215 (2002).

56.   Seeley WW*, et al*. Dissociable intrinsic connectivity networks for salience
processing and executive control. *J Neurosci* **27**, 2349-2356 (2007).

57.   Margulies DS*, et al*. Situating the default-mode network along a principal
gradient of macroscale cortical organization. *Proc Natl Acad Sci U S A* **113**,
12574-12579 (2016).

58.   Zhou Y, Friston KJ, Zeidman P, Chen J, Li S, Razi A. The Hierarchical
Organization of the Default, Dorsal Attention and Salience Networks in
Adolescents and Young Adults. *Cereb Cortex* **28**, 726-737 (2018).

59.   DuPre E, Spreng RN. Structural covariance networks across the life span, from
6 to 94 years of age. *Netw Neurosci* **1**, 302-323 (2017).

60.   Betzel RF, Byrge L, He Y, Goni J, Zuo XN, Sporns O. Changes in structural
and functional connectivity among resting-state networks across the human
lifespan. *Neuroimage* **102 Pt 2**, 345-357 (2014).

61.   Hart H, Radua J, Nakao T, Mataix-Cols D, Rubia K. Meta-analysis of functional
magnetic resonance imaging studies of inhibition and attention in attention-
deficit/hyperactivity disorder: exploring task-specific, stimulant medication,
and age effects. *JAMA psychiatry* **70**, 185-198 (2013).

62.   Zhang Z, Peng P, Eickhoff SB, Lin X, Zhang D, Wang Y. Neural substrates of
the executive function construct, age-related changes, and task materials in
adolescents and adults: ALE meta-analyses of 408 fMRI studies. *Dev Sci* **24**,
e13111 (2021).

63.   Long J, Song X, Wang Y, Wang C, Huang R, Zhang R. Distinct neural
activation patterns of age in subcomponents of inhibitory control: A fMRI meta-
analysis. *Front Aging Neurosci* **14**, 938789 (2022).

64.   Yarkoni T, Poldrack RA, Nichols TE, Van Essen DC, Wager TD. Large-scale
automated synthesis of human functional neuroimaging data. *Nat Methods* **8**,
665-670 (2011).

65.   Nee DE, Wager TD, Jonides J. Interference resolution: insights from a meta-
analysis of neuroimaging tasks. *Cogn Affect Behav Neurosci* **7**, 1-17 (2007).

66. Botvinick MM, Cohen JD, Carter CS. Conflict monitoring and anterior
1078    cingulate cortex: an update. *Trends Cogn Sci* **8**, 539-546 (2004).

1079 67. Diamond A. Executive functions. *Annu Rev Psychol* **64**, 135-168 (2013).

1080 68. Simonsohn U. Two Lines: A Valid Alternative to the Invalid Testing of U-
1081    Shaped Relationships With Quadratic Regressions. *Advances in Methods and*
1082    *Practices in Psychological Science* **1**, 538-555 (2018).

1083 69. Zuo XN*, et al.* Growing together and growing apart: regional and sex
1084    differences in the lifespan developmental trajectories of functional homotopy. *J*
1085    *Neurosci* **30**, 15034-15043 (2010).

1086 70. Coupe P, Catheline G, Lanuza E, Manjon JV, Alzheimer's Disease
1087    Neuroimaging I. Towards a unified analysis of brain maturation and aging
1088    across the entire lifespan: A MRI analysis. *Hum Brain Mapp* **38**, 5501-5518
1089    (2017).

1090 71. Amlien IK, Sneve MH, Vidal-Pineiro D, Walhovd KB, Fjell AM. The Lifespan
1091    Trajectory of the Encoding-Retrieval Flip: A Multimodal Examination of
1092    Medial Parietal Cortex Contributions to Episodic Memory. *J Neurosci* **38**, 8666-
1093    8679 (2018).

1094 72. Yeo BT*, et al.* The organization of the human cerebral cortex estimated by
1095    intrinsic functional connectivity. *J Neurophysiol* **106**, 1125-1165 (2011).

1096 73. Brenhouse HC, Andersen SL. Developmental trajectories during adolescence in
1097    males and females: a cross-species understanding of underlying brain changes.
1098    *Neurosci Biobehav Rev* **35**, 1687-1703 (2011).

1099 74. Luna B, Marek S, Larsen B, Tervo-Clemmens B, Chahal R. An integrative
1100    model of the maturation of cognitive control. *Annu Rev Neurosci* **38**, 151-170
1101    (2015).

1102 75. Grimm KJ, Ram N, Estabrook R. *Growth Modeling: Structural Equation and*
1103    *Multilevel Modeling Approaches*. Guilford Publications (2016).

1104 76. Verissimo J, Verhaeghen P, Goldman N, Weinstein M, Ullman MT. Evidence
1105    that ageing yields improvements as well as declines across attention and
1106    executive functions. *Nat Hum Behav* **6**, 97-110 (2022).

1107 77. Cole MW, Reynolds JR, Power JD, Repovs G, Anticevic A, Braver TS. Multi-
1108    task connectivity reveals flexible hubs for adaptive task control. *Nat Neurosci*
1109    **16**, 1348-1355 (2013).

1110 78. Petersen SE, Posner MI. The attention system of the human brain: 20 years after.
1111    *Annu Rev Neurosci* **35**, 73-89 (2012).

1112 79. Badre D, Nee DE. Frontal Cortex and the Hierarchical Control of Behavior.
1113    *Trends Cogn Sci* **22**, 170-188 (2018).

1114 80. Thatcher RW, Walker RA, Giudice S. Human cerebral hemispheres develop at
1115    different rates and ages. *Science* **236**, 1110-1113 (1987).

1116   81.   Dolcos F, Rice HJ, Cabeza R. Hemispheric asymmetry and aging: right
1117   hemisphere decline or asymmetry reduction. *Neurosci Biobehav Rev* **26**, 819-
1118   825 (2002).

1119   82.   Cabeza R*, et al.* Age-related differences in neural activity during memory
1120   encoding and retrieval: a positron emission tomography study. *J Neurosci* **17**,
1121   391-400 (1997).

1122   83.   Grady CL, Bernstein LJ, Beig S, Siegenthaler AL. The effects of encoding task
1123   on age-related differences in the functional neuroanatomy of face memory.
1124   *Psychol Aging* **17**, 7-23 (2002).

1125   84.   He L*, et al.* Unity and diversity of neural representation in executive functions.
1126   *J Exp Psychol Gen* **150**, 2193-2207 (2021).

1127   85.   Scherf KS, Sweeney JA, Luna B. Brain basis of developmental change in
1128   visuospatial working memory. *J Cogn Neurosci* **18**, 1045-1058 (2006).

1129   86.   Olesen PJ, Macoveanu J, Tegner J, Klingberg T. Brain activity related to
1130   working memory and distraction in children and adults. *Cereb Cortex* **17**, 1047-
1131   1054 (2007).

1132   87.   Geier CF, Garver K, Terwilliger R, Luna B. Development of working memory
1133   maintenance. *J Neurophysiol* **101**, 84-99 (2009).

1134   88.   Booth JR*, et al.* Neural development of selective attention and response
1135   inhibition. *Neuroimage* **20**, 737-751 (2003).

1136   89.   Durston S*, et al.* A shift from diffuse to focal cortical activity with development.
1137   *Dev Sci* **9**, 1-8 (2006).

1138   90.   Rubia K*, et al.* Progressive increase of frontostriatal brain activation from
1139   childhood to adulthood during event-related tasks of cognitive control. *Hum*
1140   *Brain Mapp* **27**, 973-993 (2006).

1141   91.   Rottschy C*, et al.* Modelling neural correlates of working memory: a
1142   coordinate-based meta-analysis. *Neuroimage* **60**, 830-846 (2012).

1143   92.   Criaud M, Boulinguez P. Have we been asking the right questions when
1144   assessing response inhibition in go/no-go tasks with fMRI? A meta-analysis and
1145   critical review. *Neurosci Biobehav Rev* **37**, 11-23 (2013).

1146   93.   Simmonds DJ, Pekar JJ, Mostofsky SH. Meta-analysis of Go/No-go tasks
1147   demonstrating that fMRI activation associated with response inhibition is task-
1148   dependent. *Neuropsychologia* **46**, 224-232 (2008).

1149   94.   Radua J*, et al.* A new meta-analytic method for neuroimaging studies that
1150   combines reported peak coordinates and statistical parametric maps. *Eur*
1151   *Psychiatry* **27**, 605-611 (2012).

1152   95.   Albajes-Eizagirre A, Solanes A, Vieta E, Radua J. Voxel-based meta-analysis
1153   via permutation of subject images (PSI): Theory and implementation for SDM.
1154   *Neuroimage* **186**, 174-184 (2019).

96.   Salimi-Khorshidi G, Smith SM, Keltner JR, Wager TD, Nichols TE. Meta-
analysis of neuroimaging data: a comparison of image-based and coordinate-
based pooling of studies. *Neuroimage* **45**, 810-823 (2009).

97.   Muller VI*, et al.* Ten simple rules for neuroimaging meta-analysis. *Neurosci*
Biobehav Rev **84**, 151-161 (2018).

98.   Luijten M, Schellekens AF, Kuhn S, Machielse MW, Sescousse G. Disruption
of Reward Processing in Addiction : An Image-Based Meta-analysis of
Functional Magnetic Resonance Imaging Studies. *JAMA Psychiatry* **74**, 387-
398 (2017).

99.   Sowell ER, Thompson PM, Toga AW. Mapping changes in the human cortex
throughout the span of life. *Neuroscientist* **10**, 372-392 (2004).

100.   Thompson SG, Higgins JP. How should meta-regression analyses be
undertaken and interpreted? *Stat Med* **21**, 1559-1573 (2002).

101.   Page MJ*, et al.* The PRISMA 2020 statement: an updated guideline for reporting
systematic reviews. **88**, 105906 (2021).

102.   Lancaster JL*, et al.* Bias between MNI and talairach coordinates analyzed using
the ICBM-152 brain template. *Human Brain Mapping* **28**, 1194-1205 (2007).

103.   Turkeltaub PE, Eickhoff SB, Laird AR, Fox M, Wiener M, Fox P. Minimizing
within-experiment and within-group effects in activation likelihood estimation
meta-analyses. *Human Brain Mapping* **33**, 1-13 (2012).

104.   Eickhoff SB, Bzdok D, Laird AR, Kurth F, Fox PT. Activation likelihood
estimation meta-analysis revisited. *NeuroImage* **59**, 2349-2361 (2012).

105.   Radua J, Mataix-Cols D. Meta-analytic methods for neuroimaging data
explained. *Biol Mood Anxiety Disord* **2**, 6 (2012).

106.   Hedges LV. Distribution theory for Glass's estimator of effect size and related
estimators. *Journal of Educational Statistics* **6**, 107-128 (1981).

107.   Radua J*, et al.* A new meta-analytic method for neuroimaging studies that
combines reported peak coordinates and statistical parametric maps. *Eur*
Psychiat **27**, 605-611 (2012).

108.   Egger M, Smith GD, Phillips AN. Meta-analysis: principles and procedures.
BMJ (Clinical research ed) **315**, 1533-1537 (1997).

109.   Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in
meta-analyses. *BMJ (Clinical research ed)* **327**, 557-560 (2003).

110.   Wood SN. mgcv: GAMs and generalized ridge regression for R. *J R news* **1**, 20-
25 (2001).

111.   Zuo XN, He Y, Betzel RF, Colcombe S, Sporns O, Milham MP. Human
Connectomics across the Life Span. *Trends Cogn Sci* **21**, 32-45 (2017).

112.   Seghier ML. Laterality index in functional MRI: methodological issues. *Magn*
Reson Imaging **26**, 594-601 (2008).

113.   Hoffman P, Morcom AM. Age-related changes in the neural networks
1195   supporting semantic cognition: A meta-analysis of 47 functional neuroimaging
1196   studies. *Neurosci Biobehav Rev* **84**, 134-150 (2018).
1197   114.   Appelbaum M, Cooper H, Kline RB, Mayo-Wilson E, Nezu AM, Rao SM.
1198   Journal Article Reporting Standards for Quantitative Research in Psychology:
1199   The APA Publications and Communications Board Task Force Report. *Am*
1200   *Psychol* **73**, 3-25 (2018).
1201   115.   Wood SN. Fast Stable Restricted Maximum Likelihood and Marginal
1202   Likelihood Estimation of Semiparametric Generalized Linear Models. *Journal*
1203   *of the Royal Statistical Society Series B: Statistical Methodology* **73**, 3-36
1204   (2011).
1205   116.   Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package. *J*
1206   *Stat Softw* **36**, 1-48 (2010).
1207
1208

Acknowledgement

We thank Fergus I.M. Craik, Beatriz Luna, and Haiyan Wu for valuable suggestions on this manuscript. We also thank Jing Yang and Yifan Zhang for the data check of extracted coordinates. Z.L. discloses support for the research of this work from Scientific Research Fund of Zhejiang Provincial Education Department (Y202249966), Starting Research Fund from Hangzhou Normal University (2021QDL079), and STI 2030-Major Projects (2021ZD0201705). I.T.P discloses support for the research of this work from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) (HD098235). G.Y. disclose support for the research of this work from Jiefeng Jiang, and China Postdoctoral Science Foundation (2019M650884). X.L. discloses support for the research of this work from the Ministry of Science and Technology of the People's Republic of China (2021ZD0200505).

Author Contributions

Conceptualization: Z.L. and G.Y.; Methodology: Z.L. and G.Y.; Formal analysis: Z.L. and G.Y.; Data curation: Z.L. and G.Y.; Writing original draft: Z.L. and G.Y.; Writing, review, and editing: Z.L., I.T.P., J.R., L.W., X.L. and G.Y.; Funding: Z.L., I.T.P., X.L. and G.Y; Supervision: G.Y.

Competing Interests

The authors declare no competing interests.

Inclusion & Ethics

All collaborators of this study fulfilled the criteria for authorship required by Nature Portfolio journals have been included as authors, as their participation was essential for

implementation of the study and improvement of the manuscript. Roles and responsi-

bilities were agreed among collaborators ahead of the research. This work does not

include findings that are locally relevant. As a meta-analysis, this study does not involve

direct interactions with human participants or the collection of new data. Each of the

original studies included in this meta-analysis obtained ethical approval from their re-

spective institutions, and we confirm that we complied with all applicable ethical guide-

lines and regulations.

1238